

Fig. 4. Effect of expression of HSP70 on production of TGF- β 1 and expression of pro-inflammatory cytokines. Transgenic mice overexpressing HSP70 (HSP70 tg) and wild-type (WT) mice were treated with (BLM) or without (vehi) bleomycin (5 mg/kg) once-only at day 0 and cells in BALF were collected at day 3 (A and B). Cells were incubated for 24 h and the level of TGF- β 1 in the culture medium was determined by ELISA (A). Total RNA was extracted and subjected to real-time RT-PCR using a specific primer set for each gene. Values were normalized to the *gapdh* gene and expressed relative to the control sample (B). Values shown are mean \pm S.E.M. ($n = 3-6$). ** or *** $P < 0.01$.

shown in Fig. 5E, either treatment with TGF- β 1 or transfection with siRNA for HSP70 induced morphological change (from a cobble-stone-like epithelial monolayer to dispersed spindle-shaped mesenchymal cells with reduced cell-cell contact), one of characteristic features of induction of EMT.

A number of transcription factors (such as Slug) are involved in the induction of EMT [30,31]. We found that the transfection of cells with siRNA for HSP70 enhanced the expression of *slug* mRNA in the presence and absence of TGF- β 1 (Fig. 5A). The results in Fig. 5 suggest that the expression of HSP70 suppresses the bleomycin-induced EMT of A549 cells by suppressing the expression of Slug.

We also examined the effect of HSP70 expression on the TGF- β 1-dependent activation of lung fibroblasts *in vitro*. Treatment of HFL-1 cells (human embryonic lung fibroblasts) with TGF- β 1-induced the expression of *col1a1*, *α -sma* and *hsp47* mRNAs and collagen I protein (Fig. 6A and B), suggesting that TGF- β 1 activated fibroblasts to myofibroblasts. The transfection of cells with siRNA for HSP70 did not affect the TGF- β 1-dependent alteration in expression of these genes (Fig. 6A and B), suggesting that the expression of HSP70 did not affect the TGF- β 1-dependent activation of fibroblasts.

3.3. Effect of GGA on bleomycin-induced pulmonary fibrosis

We subsequently examined the effect of GGA on bleomycin-induced pulmonary damage, inflammatory response, fibrosis and

dysfunction. Oral administration of GGA suppressed the bleomycin-induced increase in the number of inflammatory cells present in BALF (Fig. 7A), and decreased pulmonary damage (Fig. 7B) and epithelial apoptosis (Fig. 7C and D) 3 days after the administration of bleomycin. Immunohistochemical analysis with an antibody against HSP70 revealed that treatment with bleomycin slightly induced the pulmonary expression of HSP70, and the simultaneous administration of GGA enhanced this expression (Fig. 7E). The extent of bleomycin-induced pulmonary fibrosis 14 days after the administration of bleomycin was also reduced by the administration of GGA; GGA suppressed both the bleomycin-induced collagen deposition and the increase in pulmonary hydroxyproline content (Fig. 7F and G).

We also examined the effect of GGA on bleomycin-induced alterations to lung mechanics. As shown in Fig. 7H, the bleomycin-induced decrease in total respiratory system compliance and the increase in total respiratory system elastance and tissue elastance were significantly suppressed by the administration of GGA, suggesting that GGA exerts a protective effect against bleomycin-induced lung dysfunction. Taking these findings with the results shown in Figs. 2 and 3, it is likely that GGA suppresses the negative effects of bleomycin and improves lung function by inducing HSP70 expression.

In order to test this idea, we examined the effect of quercetin, an inhibitor for HSP70, on the ameliorative effect of GGA for bleomycin-induced pulmonary fibrosis and dysfunction. As shown in Fig. 7F–H, the ameliorative effect of GGA on bleomycin-induced pulmonary fibrosis and dysfunction was not observed in the presence of simultaneous administration of quercetin, suggesting that GGA suppresses bleomycin-induced pulmonary fibrosis and dysfunction by inducing HSP70 expression.

4. Discussion

An ameliorative effect of HSP70 due to its cytoprotective, anti-inflammatory and molecular chaperone (quality control of proteins) properties has been reported for animal models of various diseases. For example, using transgenic mice overexpressing HSP70, we have reported that HSP70 protects against irritant-produced lesions in the stomach and small intestine, inflammatory bowel disease-related experimental colitis and ultraviolet (UV)-induced skin damage [23,32–35]. The potential therapeutic applicability of HSP70 for use in other diseases, such as neurodegenerative diseases, ischemia-reperfusion damage and diabetes has also been suggested [36–38]. Interestingly, GGA, an anti-ulcer drug and HSP-inducer has been reported to suppress not only gastric lesions but also lesions of the small intestine, inflammatory bowel disease-related experimental colitis and neurodegenerative diseases [32,34,37,39]. On the other hand, the role of HSP70 in IPF has not been fully evaluated. It was recently reported that oral administration of GGA suppresses bleomycin-induced pulmonary fibrosis [40]. However, because GGA induces expression of HSPs other than HSP70, and has various pharmacological activities other than induction of HSPs (such as an increase in blood flow, stimulation of surface mucus production and direct protection of cell membranes [41–43]), it remains unclear whether GGA achieves its anti-fibrotic activity through up-regulation of expression of HSP70. In this study, we used transgenic mice overexpressing HSP70 to identify the role of HSP70 in bleomycin-induced pulmonary fibrosis and found that the transgenic mice showed a phenotypic resistance to this treatment. This is the first genetic evidence of the protective role of HSP70 against IPF-related fibrosis. The results presented here also suggest that GGA achieves its anti-fibrotic activity via the up-regulation of HSP70 expression.




Fig. 5. Effect of siRNA for HSP70 on TGF-β1-induced EMT-like phenotypes. A549 cells were transfected with 1.2 μg of siRNA for HSP70 (siHSP70) or non-silencing (ns) siRNA and incubated for 24 h. Cells were then incubated with the indicated concentration of TGF-β1 for 48 h. Expression of each gene was examined by real-time RT-PCR as described in the legend of Fig. 4 (A). Expression of each protein was examined by immunoblotting as described in the legend of Fig. 1 (B). Immunostaining with an antibody against E-cadherin (C) or α-SMA (D) was done as described in Section 2. Cell morphology was examined by phase-contrast microscopic observation (E). Values shown are mean ± S.E.M. (n = 3). * P < 0.05; ** or *** P < 0.01.




Fig. 6. Effect of siRNA for HSP70 on the TGF-β1-dependent activation of fibroblasts. HFL-I cells were transfected with 1.2 μg of siRNA for HSP70 (siHSP70) or non-silencing (ns) siRNA and incubated for 24 h. Cells were then incubated with the indicated concentrations of TGF-β1 for 24 h. Expression of each gene was examined by real-time RT-PCR as described in the legend of Fig. 4 (A). The levels of collagen I and HSP70 in cell lysate and collagen I in medium were examined by immunoblotting as described in the legend of Fig. 1 (B). Values shown are mean ± S.E.M. (n = 3). *P < 0.05; ** or *** P < 0.01.

Given that an increase in alveolar epithelial cell apoptosis has been observed in human IPF [44], the apoptotic process is believed to play an important role in the development of IPF. We here showed that bleomycin-induced pulmonary damage and epithelial cell apoptosis were suppressed in transgenic mice overexpressing HSP70. It is well known that HSP70 has anti-apoptotic effects through various mechanisms such as binding to apoptotic protease activating factor (Apaf)-1 to prevent the activation of caspase-9, suppression of the apoptotic pathway downstream of caspase-3 activation, and suppression of apoptosis-inducing factor-induced chromatin condensation [45–48]. We recently reported that HSP70 inhibits the activation of bcl-2-associated X protein (BAX), which is important for apoptosis-related mitochondrial dysfunction [32]. We consider that these mechanisms are involved in the HSP70-dependent suppression of bleomycin-induced lung epithelial cell apoptosis, because the activation of BAX and of caspases were reported to play an important role in bleomycin-induced apoptosis in lung epithelial cells [49,50].

In addition to this anti-apoptotic (cytoprotective) effect of HSP70, its anti-inflammatory effect was recently revealed and is thought to be important for its protective role against various diseases. HSP70 suppresses the activation of nuclear factor kappa B (NF-κB; an inflammation-inducing transcription factor) through various mechanisms such as suppression of inflammatory stimuli-induced degradation of IκB-α (an inhibitor of NF-κB) [18,51]. We previously reported that inflammatory responses (such as expression of pro-inflammatory cytokines and infiltration of leucocytes) in the stomach, colon and skin were suppressed in transgenic mice overexpressing HSP70 and that this suppression is mediated by the inhibition of NF-κB

[23,32,34,35]. In this study, we showed that a bleomycin-induced increase in leucocytes in BALF and the pulmonary expression of pro-inflammatory cytokines were suppressed in transgenic mice overexpressing HSP70. We also showed that production of TGF-β1 was suppressed in bleomycin-treated transgenic mice overexpressing HSP70 compared to corresponding wild-type mice. This is the first demonstration of the inhibitory effect of HSP70 on the production of TGF-β1. It is known that TNF-α induces the expression of TGF-β1 [52] suggesting that the inhibitory effect of HSP70 on NF-κB and the resulting decrease in the level of TNF-α are responsible for this inhibitory effect of HSP70 on TGF-β1 production. Since TGF-β1 plays a major role in bleomycin-induced pulmonary fibrosis through various mechanisms such as activation of fibroblasts and stimulation of EMT of epithelial cells, the inhibitory effect of HSP70 on the production of TGF-β1 should be responsible for its protective effect against bleomycin-induced pulmonary fibrosis.

We also examined the effect of HSP70 expression on TGF-β1-dependent cellular responses involved in pulmonary fibrosis. We found that the TGF-β1-dependent induction of EMT-like phenotypes in lung epithelial cells (up-regulation of expression of markers of myofibroblasts and down-regulation of expression of markers of epithelial cells) and up-regulation of the expression of Slug (a transcription factor inducing EMT) were stimulated by the suppression of HSP70 expression. On the other hand, the TGF-β1-dependent activation of fibroblasts was not affected by the suppression of HSP70 expression. These results suggest that the expression of HSP70 suppresses the bleomycin-induced increase in lung myofibroblast number via suppression of the TGF-β1-dependent EMT of epithelial cells rather than via the activation of fibroblasts. It was recently reported that expression of HSP70 in cultured rat kidney proximal tubular epithelial cells inhibited TGF-β1-induced EMT, although the mechanism governing this inhibition is unknown [53]. Thus, it seems that expression of HSP70 generally suppresses EMT of epithelial cells. It is possible that the inhibitory effect of extracellular HSP70 on mitogen-activated protein kinases (MAPKs) [54] is involved in the inhibitory effect of HSP70 on the TGF-β1-induced EMT of epithelial cells, because MAPKs are involved in TGF-β1-dependent signal transduction pathways [55]. Furthermore, it is also possible that HSP70 achieves its inhibitory effect on EMT via the inhibition of NF-κB, because it was recently reported that NF-κB stimulates EMT via a mechanism that is independent on TGF-β1 [56].

While bleomycin-induced pulmonary fibrosis has been used as an animal model of IPF, this model does however has some limitations, such as the spontaneous resolution of fibrosis, which is rare in human IPF [57]. Furthermore, although assessment tools used in bleomycin-induced pulmonary fibrosis are primarily based on histology and quantitative collagen analysis, clinical management of IPF relies on lung function analysis. Therefore, in this study, we used a computer-controlled small-animal ventilator to monitor the bleomycin-induced decrease in compliance and increase in elastance, which are known to be associated with human IPF [58]. We found that these bleomycin-induced alterations in lung mechanics were ameliorated in transgenic mice overexpressing HSP70, suggesting that an increased expression of HSP70 could potentially improve the pulmonary dysfunction associated with human IPF.

As described above, it was recently reported that administration of GGA ameliorates bleomycin-induced pulmonary fibrosis [40], a result that we confirmed in this study. We also found that the administration of GGA improved lung function in the presence of bleomycin treatment. As described in Section 1, current agents for the treatment of IPF have not been found to improve the




Fig. 7. Effect of oral administration of GGA on bleomycin-induced pulmonary damage, inflammatory response, fibrosis and dysfunction. C57/BL6 mice were orally administered with GGA (200 mg/kg) once per day for 3 days (A–E) or 14 days (F–H). Quercetin (200 mg/kg) was orally administered into mice 30 min before each administration of GGA (F–H). Mice were treated with (BLM) or without (vehi) bleomycin (5 mg/kg) once-only at day 0 and cells in BALF (A), pulmonary damage (B–D), pulmonary fibrosis (F and G) and lung mechanics (H) were assessed as described in the legends of Figs. 2 and 3. Immunohistochemical analysis with an antibody against HSP70 was performed as described in Section 2 (E). Values shown are mean ± S.E.M. (n = 3–12). *, * or ⁵ < 0.05; **, ** or ⁵⁵P < 0.01.

prognosis [1,3,4,59]. We consider that HSP70-inducers such as GGA could be beneficial for the prophylaxis of IPF. The development of new molecules as candidate drugs to treat this disease must pass through the clinical trials process and may encounter

unanticipated side effects. Thus, based on the results of this study, we propose that clinical studies should be performed to prove the effectiveness of GGA for treating IPF given that the safety of GGA has already been shown clinically.




Fig. 7. (Continued).

Acknowledgements

This work was supported by Grants-in-Aid for Scientific Research from the Ministry of Health, Labour, and Welfare of Japan, as well as the Japan Science and Technology Agency and Grants-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology, Japan.

References

- American Thoracic Society. Idiopathic pulmonary fibrosis: diagnosis and treatment. International consensus statement. American Thoracic Society (ATS), and the European Respiratory Society (ERS). *Am J Respir Crit Care Med* 2000;161:646–64.
- Bjoraker JA, Ryu JH, Edwin MK, Myers JL, Tazelaar HD, Schroeder DR, et al. Prognostic significance of histopathologic subsets in idiopathic pulmonary fibrosis. *Am J Respir Crit Care Med* 1998;157:199–203.
- American Thoracic Society/European Respiratory Society International Multi-disciplinary Consensus Classification of the Idiopathic Interstitial Pneumonias. This joint statement of the American Thoracic Society (ATS), and the European Respiratory Society (ERS) was adopted by the ATS board of directors, June 2001 and by the ERS Executive Committee, June 2001. *Am J Respir Crit Care Med* 2002;165:277–304.
- Walter N, Collard HR, King Jr TE. Current perspectives on the treatment of idiopathic pulmonary fibrosis. *Proc Am Thorac Soc* 2006;3:330–8.
- Moore BB, Hogaboam CM. Murine models of pulmonary fibrosis. *Am J Physiol Lung Cell Mol Physiol* 2008;294:L152–60.
- Kinnula VL, Myllarniemi M. Oxidant-antioxidant imbalance as a potential contributor to the progression of human pulmonary fibrosis. *Antioxid Redox Signal* 2008;10:727–38.
- Sheppard D. Transforming growth factor beta: a central modulator of pulmonary and airway inflammation and fibrosis. *Proc Am Thorac Soc* 2006;3:413–7.
- Kisseleva T, Brenner DA. Fibrogenesis of parenchymal organs. *Proc Am Thorac Soc* 2008;5:338–42.
- Willis BC, Borok Z. TGF-beta-induced EMT: mechanisms and implications for fibrotic lung disease. *Am J Physiol Lung Cell Mol Physiol* 2007;293:L525–34.
- Bartram U, Speer CP. The role of transforming growth factor beta in lung development and disease. *Chest* 2004;125:754–65.
- Kasai H, Allen JT, Mason RM, Kamimura T, Zhang Z. TGF-beta1 induces human alveolar epithelial to mesenchymal cell transition (EMT). *Respir Res* 2005;6:56.
- Kim KK, Kugler MC, Wolters PJ, Robillard L, Galvez MG, Brumwell AN, et al. Alveolar epithelial cell mesenchymal transition develops in vivo during pulmonary fibrosis and is regulated by the extracellular matrix. *Proc Natl Acad Sci USA* 2006;103:13180–5.
- Kim Y, Kugler MC, Wei Y, Kim KK, Li X, Brumwell AN, et al. Integrin alpha3-beta1-dependent beta-catenin phosphorylation links epithelial Smad signaling to cell contacts. *J Cell Biol* 2009;184:309–22.
- Wu Z, Yang L, Cai L, Zhang M, Cheng X, Yang X, et al. Detection of epithelial to mesenchymal transition in airways of a bleomycin induced pulmonary fibrosis model derived from an alpha-smooth muscle actin-Cre transgenic mouse. *Respir Res* 2007;8:1.

- [15] Kim KK, Wei Y, Szekeres C, Kugler MC, Wolters PJ, Hill ML, et al. Epithelial cell alpha3beta1 integrin links beta-catenin and Smad signaling to promote myofibroblast formation and pulmonary fibrosis. *J Clin Invest* 2009;119:213–24.
- [16] Mathew A, Morimoto RI. Role of the heat-shock response in the life and death of proteins. *Ann N Y Acad Sci* 1998;851:99–111.
- [17] Hirakawa T, Rokutan K, Nikawa T, Kishi K. Geranylgeranylacetone induces heat shock proteins in cultured guinea pig gastric mucosal cells and rat gastric mucosa. *Gastroenterology* 1996;111:345–57.
- [18] Tang D, Kang R, Xiao W, Wang H, Calderwood SK, Xiao X. The anti-inflammatory effects of heat shock protein 72 involve inhibition of high-mobility-group box 1 release and proinflammatory function in macrophages. *J Immunol* 2007;179:1236–44.
- [19] Weiss YG, Maloyan A, Tazelaar J, Raj N, Deutschman CS. Adenoviral transfer of HSP-70 into pulmonary epithelium ameliorates experimental acute respiratory distress syndrome. *J Clin Invest* 2002;110:801–6.
- [20] Koh Y, Lim CM, Kim MJ, Shim TS, Lee SD, Kim WS, et al. Heat shock response decreases endotoxin-induced acute lung injury in rats. *Respirology* 1999;4:325–30.
- [21] Hagiwara S, Iwasaka H, Matsumoto S, Noguchi T, Yoshioka H. Association between heat stress protein 70 induction and decreased pulmonary fibrosis in an animal model of acute lung injury. *Lung* 2007;185:287–93.
- [22] Singleton KD, Wischmeyer PE. Effects of HSP70.1/3 gene knockout on acute respiratory distress syndrome and the inflammatory response following sepsis. *Am J Physiol Lung Cell Mol Physiol* 2006;290:L956–61.
- [23] Tanaka K, Namba T, Arai Y, Fujimoto M, Adachi H, Sobue G, et al. Genetic evidence for a protective role for heat shock factor 1 and heat shock protein 70 against colitis. *J Biol Chem* 2007;282:23240–52.
- [24] Woessner Jr JF. The determination of hydroxyproline in tissue and protein samples containing small proportions of this imino acid. *Arch Biochem Biophys* 1961;93:440–7.
- [25] Mima S, Tsutsumi S, Ushijima H, Takeda M, Fukuda I, Yokomizo K, et al. Induction of claudin-4 by nonsteroidal anti-inflammatory drugs and its contribution to their chemopreventive effect. *Cancer Res* 2005;65:1868–76.
- [26] Lovgren AK, Jania LA, Hartney JM, Parsons KK, Audoly LP, Fitzgerald GA, et al. COX-2-derived prostacyclin protects against bleomycin-induced pulmonary fibrosis. *Am J Physiol Lung Cell Mol Physiol* 2006;291:L144–56.
- [27] Plumier JC, Ross BM, Currie RW, Angelidis CE, Kazlaris H, Kollias G, et al. Transgenic mice expressing the human heat shock protein 70 have improved post-ischemic myocardial recovery. *J Clin Invest* 1995;95:1854–60.
- [28] Ishii H, Mukae H, Kakugawa T, Iwashita T, Kaida H, Fujii T, et al. Increased expression of collagen-binding heat shock protein 47 in murine bleomycin-induced pneumopathy. *Am J Physiol Lung Cell Mol Physiol* 2003;285:L957–63.
- [29] Inoue K, Takano H, Yanagisawa R, Sakurai M, Abe S, Yoshino S, et al. Effects of nanoparticles on lung physiology in the presence or absence of antigen. *Int J Immunopathol Pharmacol* 2007;20:737–44.
- [30] Peinado H, Olmeda D, Cano A. Snail, Zeb and bHLH factors in tumour progression: an alliance against the epithelial phenotype? *Nat Rev Cancer* 2007;7:415–28.
- [31] Thiery JP, Sleeman JP. Complex networks orchestrate epithelial–mesenchymal transitions. *Nat Rev Mol Cell Biol* 2006;7:131–42.
- [32] Suemasu S, Tanaka K, Namba T, Ishihara T, Katsu T, Fujimoto M, et al. A role for HSP70 in protecting against indomethacin-induced gastric lesions. *J Biol Chem* 2009;284:19705–1.
- [33] Tanaka K, Tsutsumi S, Arai Y, Hoshino T, Suzuki K, Takaki E, et al. Genetic evidence for a protective role of heat shock factor 1 against irritant-induced gastric lesions. *Mol Pharmacol* 2007;71:985–93.
- [34] Asano T, Tanaka K, Yamakawa N, Adachi H, Sobue G, Goto H, et al. HSP70 confers protection against indomethacin-induced lesions of the small intestine. *J Pharmacol Exp Ther* 2009;330:458–67.
- [35] Matsuda M, Hoshino T, Yamashita Y, Tanaka KI, Maji D, Sato K, et al. Prevention of ultraviolet B radiation-induced epidermal damage by expression of heat shock protein 70. *J Biol Chem* 2009.
- [36] Gupta AA, Bomhoff GL, Swerdlow RH, Geiger PC. Heat treatment improves glucose tolerance and prevents skeletal muscle insulin resistance in rats fed a high-fat diet. *Diabetes* 2009;58:567–78.
- [37] Katsuno M, Sang C, Adachi H, Minamiyama M, Waza M, Tanaka F, et al. Pharmacological induction of heat-shock proteins alleviates polyglutamine-mediated motor neuron disease. *Proc Natl Acad Sci USA* 2005;102:16801–6.
- [38] Kuboki S, Schuster R, Blanchard J, Pritts TA, Wong HR, Lentsch AB. Role of heat shock protein 70 in hepatic ischemia-reperfusion injury in mice. *Am J Physiol Gastrointest Liver Physiol* 2007;292:G1141–9.
- [39] Ohkawara T, Nishihira J, Takeda H, Miyashita K, Kato K, Kato M, et al. Geranylgeranylacetone protects mice from dextran sulfate sodium-induced colitis. *Scand J Gastroenterol* 2005;40:1049–57.
- [40] Fujibayashi T, Hashimoto N, Jijiwa M, Hasegawa Y, Kojima T, Ishiguro N. Protective effect of geranylgeranylacetone, an inducer of heat shock protein 70, against drug-induced lung injury/fibrosis in an animal model. *BMC Pulm Med* 2009;9:45.
- [41] Terano A, Hiraishi H, Ota S, Sugimoto T. Geranylgeranylacetone, a novel anti-ulcer drug, stimulates mucus synthesis and secretion in rat gastric cultured cells. *Digestion* 1986;33:206–10.
- [42] Kunisaki C, Sugiyama M. Effect of teprenone on acute gastric mucosal lesions induced by cold-restraint stress. *Digestion* 1992;53:45–53.
- [43] Ushijima H, Tanaka K, Takeda M, Katsu T, Mima S, Mizushima T. Geranylgeranylacetone protects membranes against nonsteroidal anti-inflammatory drugs. *Mol Pharmacol* 2005;68:1156–61.
- [44] Thannickal VJ, Horowitz JC. Evolving concepts of apoptosis in idiopathic pulmonary fibrosis. *Proc Am Thorac Soc* 2006;3:350–6.
- [45] Saleh A, Srinivasula SM, Balkir L, Robbins PD, Alnemri ES. Negative regulation of the Apaf-1 apoptosome by Hsp70. *Nat Cell Biol* 2000;2:476–83.
- [46] Beere HM, Wolf BB, Cain K, Mosser DD, Mahboubi A, Kuwana T, et al. Heat-shock protein 70 inhibits apoptosis by preventing recruitment of procaspase-9 to the Apaf-1 apoptosome. *Nat Cell Biol* 2000;2:469–75.
- [47] Ravagnan L, Gurbuxani S, Susin SA, Maise C, Daugas E, Zamzami N, et al. Heat-shock protein 70 antagonizes apoptosis-inducing factor. *Nat Cell Biol* 2001;3:839–43.
- [48] Jaattela M, Wissing D, Kokholm K, Kallunki T, Egeblad M. Hsp70 exerts its anti-apoptotic function downstream of caspase-3-like proteases. *EMBO J* 1998;17:6124–34.
- [49] Li X, Zhang H, Soledad-Conrad V, Zhuang J, Uhal BD. Bleomycin-induced apoptosis of alveolar epithelial cells requires angiotensin synthesis de novo. *Am J Physiol Lung Cell Mol Physiol* 2003;284:L501–7.
- [50] Lee VY, Schroedl C, Brunelle JK, Buccellato LJ, Akinci OI, Kaneto H, et al. Bleomycin induces alveolar epithelial cell death through JNK-dependent activation of the mitochondrial death pathway. *Am J Physiol Lung Cell Mol Physiol* 2005;289:L521–8.
- [51] Weiss YG, Bromberg Z, Raj N, Raphael J, Goloubinoff P, Ben-Neriah Y, et al. Enhanced heat shock protein 70 expression alters proteasomal degradation of IkkappaB kinase in experimental acute respiratory distress syndrome. *Crit Care Med* 2007;35:2128–38.
- [52] Sullivan DE, Ferris M, Pociask D, Brody AR. The latent form of TGFbeta(1) is induced by TNFalpha through an ERK specific pathway and is activated by asbestos-derived reactive oxygen species in vitro and in vivo. *J Immunotoxicol* 2008;5:145–9.
- [53] Mao H, Li Z, Zhou Y, Zhuang S, An X, Zhang B, et al. HSP72 attenuates renal tubular cell apoptosis and interstitial fibrosis in obstructive nephropathy. *Am J Physiol Renal Physiol* 2008;295:F202–14.
- [54] Luo X, Zuo X, Zhou Y, Zhang B, Shi Y, Liu M, et al. Extracellular heat shock protein 70 inhibits tumour necrosis factor-alpha induced proinflammatory mediator production in fibroblast-like synoviocytes. *Arthritis Res Ther* 2008;10:R41.
- [55] He S, Liu X, Yang Y, Huang W, Xu S, Yang S, et al. Mechanisms of transforming growth factor beta/Smad signalling mediated by mitogen-activated protein kinase pathways in keloid fibroblasts. *Br J Dermatol* 2009.
- [56] Huber MA, Azoitei N, Baumann B, Grunert S, Sommer A, Pehamberger H, et al. NF-kappaB is essential for epithelial–mesenchymal transition and metastasis in a model of breast cancer progression. *J Clin Invest* 2004;114:569–81.
- [57] Moeller A, Ask K, Warburton D, Gaudie J, Kolb M. The bleomycin animal model: a useful tool to investigate treatment options for idiopathic pulmonary fibrosis? *Int J Biochem Cell Biol* 2008;40:362–82.
- [58] Ask K, Labiris R, Farkas L, Moeller A, Froese A, Farncombe T, et al. Comparison between conventional and “clinical” assessment of experimental lung fibrosis. *J Transl Med* 2008;6:16.
- [59] Luppi F, Cerri S, Beghe B, Fabbri LM, Richeldi L. Corticosteroid and immunomodulatory agents in idiopathic pulmonary fibrosis. *Respir Med* 2004;98:1035–44.

Suppression of Expression of Endoplasmic Reticulum Chaperones by *Helicobacter pylori* and Its Role in Exacerbation of Non-steroidal Anti-inflammatory Drug-induced Gastric Lesions^{*S}

Received for publication, May 26, 2010, and in revised form, September 21, 2010. Published, JBC Papers in Press, September 22, 2010, DOI 10.1074/jbc.M110.148882

Takushi Namba[‡], Tatsuya Hoshino[‡], Shintaro Suemasu[‡], Mika Takarada-Iemata[§], Osamu Hori[§], Naomi Nakagata[¶], Akinori Yanaka^{||}, and Tohru Mizushima^{*1}

From the [‡]Graduate School of Medical and Pharmaceutical Sciences, Kumamoto University, Kumamoto 862-0973, the [§]Department of Neuroanatomy, Kanazawa University Graduate School of Medical Science, Kanazawa 920-8640, the [¶]Center for Animal Resources and Development, Institute of Resource Development and Analysis, Kumamoto University, Kumamoto 860-0811, and the ^{||}Division of Clinical Pharmacology, Faculty of Pharmaceutical Sciences, Tokyo University of Science, Noda 278-8510, Japan

Both the use of non-steroidal anti-inflammatory drugs (NSAIDs), such as indomethacin, and infection with *Helicobacter pylori* are major causes of gastric ulcers. Although some clinical studies suggest that infection with *H. pylori* increases the risk of developing NSAID-induced gastric lesions, the molecular mechanism governing this effect is unknown. We recently found that in cultured gastric cells, expression of endoplasmic reticulum (ER) chaperones (such as 150-kDa oxygen-regulated protein (ORP150) and glucose-regulated protein 78 (GRP78)) is induced by NSAIDs and confers protection against NSAID-induced apoptosis, which is important in the development of NSAID-induced gastric lesions. In this study we have found that co-culture of gastric cells with *H. pylori* suppresses the expression of ER chaperones. This suppression was regulated at the level of transcription and accompanied by a reduction in the level of activating transcription factor 6 (ATF6), one of the transcription factors for ER chaperone genes. *In vivo*, inoculation of mice with *H. pylori* suppressed the expression of ER chaperones at gastric mucosa both with and without administration of indomethacin. Inoculation with *H. pylori* also stimulated formation of indomethacin-induced gastric lesions and mucosal cell death. In addition, we found that heterozygous ORP150-deficient mice are sensitive to the development of indomethacin-induced gastric lesions and mucosal cell death. The results of this study suggest that *H. pylori* exacerbates NSAID-induced gastric lesions through suppression of expression of ER chaperones, which stimulates NSAID-induced mucosal cell death.

The balance between aggressive and defensive factors determines whether gastric ulcers develop. The gastric mucosa is

challenged by a variety of aggressive factors, and of these, both non-steroidal anti-inflammatory drugs (NSAIDs)² and *Helicobacter pylori* are major causes of gastric lesions. Therefore, an important question is whether infection with *H. pylori* increases the risk of developing NSAID-induced gastric lesions (in other words, if eradication of *H. pylori* would reduce the risk of developing NSAID-induced gastric lesions). Recent clinical studies suggest that infection with *H. pylori* increases the risk of developing NSAID-induced gastric lesions (1–4); however, some studies have shown the opposite effect (5, 6). Animal models could be useful to address this issue. For example, some reports have demonstrated that NSAID-induced gastric lesions in mongolian gerbils are exacerbated by infection with *H. pylori* (7–9), although the molecular mechanism governing this exacerbation is unclear.

An inhibitory effect of NSAIDs on cyclooxygenase (COX) activity and the resulting decrease in the gastric level of prostaglandins (PGs), especially PGE₂, was believed to be the only explanation for the gastric side effects of NSAIDs because PGE₂ is a strong protective factor for gastric mucosa (10). However, the increased incidence of gastrointestinal lesions and the decrease in PG levels induced by NSAIDs are not always linked with each other. For example, it has been shown that higher doses of NSAIDs are required for producing gastric lesions than are required for inhibiting COX at the gastric mucosa (11), suggesting that there are additional mechanisms involved in the development of NSAID-induced gastric lesions. We have recently demonstrated that NSAIDs induce apoptosis in cultured gastric mucosal cells and at gastric mucosa in a manner independent of COX inhibition (12–16) and have suggested that both COX inhibition (measured as a decrease in the gastric PGE₂ level) and gastric mucosal apoptosis are required for the formation of NSAID-induced gastric lesions (16–18). There-

* This work was supported by grants-in-aid of Scientific Research from the Ministry of Health, Labour, and Welfare of Japan, grants-in-aid for Scientific Research from the Ministry of Education, Culture, Sports, Science, and Technology of Japan, and grants-in-aid of the Japan Science and Technology Agency.

^S The on-line version of this article (available at <http://www.jbc.org>) contains supplemental Figs. S1–S5.

¹ To whom correspondence should be addressed. Tel.: and Fax: 81-96-371-4323; E-mail: mizu@gpo.kumamoto-u.ac.jp.

² The abbreviations used are: NSAID, non-steroidal anti-inflammatory drug; ATF, activating transcription factor; CagA, cytotoxin-associated gene A; CHOP, C/EBP homologous transcription factor; ER, endoplasmic reticulum; ERSE, ER stress response element; GRP78, glucose-regulated protein 78; IRE1, protein-kinase and site-specific endoribonuclease; PERK, protein kinase R-like ER kinase; ORP150, 150-kDa oxygen-regulated protein; PG, prostaglandin; S1P, site-1 protease; S2P, site-2 protease; VacA, vacuolating cytotoxin A; XBP, X box binding protein.

fore, protection against gastric mucosal apoptosis is important for protecting gastric mucosa against the formation of NSAID-induced lesions. As for the molecular mechanism governing this apoptosis, we have proposed the following pathway. Permeabilization of cytoplasmic membranes by NSAIDs stimulates Ca^{2+} influx and increases intracellular Ca^{2+} levels, which in turn induces the endoplasmic reticulum (ER) stress response (12, 13, 19, 20). In the ER stress response, an apoptosis-inducing transcription factor, C/EBP homologous transcription factor (CHOP), is induced. We have previously shown that CHOP is essential for NSAID-induced apoptosis (13). CHOP induces expression of p53 up-regulated modulator of apoptosis and the resulting activation of Bax, mitochondrial dysfunction, and the activation of caspases and apoptosis (18, 21).

The ER stress response is induced by accumulation of unfolded proteins in the ER, a process involving three types of ER transmembrane proteins: protein kinase and site-specific endoribonuclease (IRE1), protein kinase R-like ER kinase (PERK), and activating transcription factor 6 (ATF6) (22–24). ER stressors phosphorylate PERK, which in turn phosphorylates eukaryotic initiation factor-2 α , leading to the activation of ATF4 expression (25). In the presence of ER stressors, p90-ATF6 (full-length ATF6) is translocated from the ER to the Golgi apparatus, where it is sequentially cleaved by site-1 protease (S1P) and site-2 protease (S2P) into p50-ATF6 (24). Activation of IRE1 causes frame switch splicing of X box-binding protein 1 (XBP-1), which produces the active (spliced) form of XBP-1 (26). All of ATF4, p50-ATF6, and XBP-1 specifically activate the transcription of ER stress response-related genes. ER stress response-related proteins contain not only CHOP but also ER chaperones (such as 150-kDa oxygen-regulated protein (ORP150) and glucose-regulated protein 78 (GRP78)) that confer protection against ER stressors by refolding unfolded proteins in the ER. We have recently reported that up-regulation of expression of GRP78 and ORP150 by NSAIDs protects gastric cells from NSAID-induced apoptosis *in vitro* (19, 20), suggesting that ER chaperones are defensive factors for gastric mucosa. Although it has been reported that expression of ER chaperones is induced with the development of gastric lesions by water-immersion stress (27), there is no direct (such as genetic) evidence supporting the notion that ER chaperones are defensive factors for gastric mucosa.

The identification of *H. pylori* in the human stomach has changed the diagnosis and treatment of gastric diseases because it is now clear that *H. pylori* plays an important role in various gastric diseases, such as chronic gastritis, peptic ulcers, and gastric cancers (28, 29). This idea is supported by clinical results that eradication of *H. pylori* significantly decreases the risk of these gastric diseases (2, 30, 31). Infection with *H. pylori* damages gastric mucosa through production of ammonia and cytotoxic proteins such as vacuolating cytotoxin A (VacA) and cytotoxin-associated gene A (CagA) and induction of host inflammatory responses (32–36). In addition to these mechanisms, it is also possible that *H. pylori* damages gastric mucosa through decreasing the expression of host defensive factors for gastric mucosa. It was recently reported that *H. pylori* inhibit the expression of heat shock proteins and mucin, both of which are major defensive factors for gastric mucosa (35, 37). How-

ever, the effect of *H. pylori* on other defensive factors, including ER chaperones, has not been tested. In this study we found that co-culture of gastric cells with *H. pylori in vitro* decreases the expression of ER chaperones, and we suggest that this suppression is mediated by the degradation of ATF6. We also show that inoculation of mice with *H. pylori* not only suppresses the expression of ER chaperones at gastric mucosa but also exacerbates NSAID-induced gastric lesions and mucosal cell death. We have also found that heterozygous ORP150-deficient mice are sensitive to the development of NSAID-induced gastric lesions and mucosal cell death. The results of this study suggest that *H. pylori* exacerbates NSAID-induced gastric lesions through suppression of expression of ER chaperones and the resulting stimulation of NSAID-induced mucosal cell death.

EXPERIMENTAL PROCEDURES

Chemicals and Animals—RPMI 1640 and *Helicobacter* selection agar were obtained from Nissui Pharmaceutical Co (Osaka, Japan). Aneropack Bikouki was from Mitsubishi Gas Chemical (Tokyo, Japan). Horse serum was from Invitrogen. Paraformaldehyde, epoxomycin, cycloheximide, pepstatin A, and fetal bovine serum (FBS) were from Sigma. E-64-d was from the Peptide Institute Inc. (Tokyo, Japan). Indomethacin was obtained from Wako Co. (Osaka, Japan). Brain heart infusion was from Difco. Antibodies against GRP78, ATF4, ATF6, actin, lamin, the N-terminal region of Bax (Bax N20), and CHOP were from Santa Cruz Biotechnology (Santa Cruz, CA). Antibodies against GRP94, GRP58, and protein disulfide isomerase were from StressGen (San Diego, CA). An antibody against cytochrome *c* was from BD Biosciences, and that against connexin 43 was from Invitrogen. An antibody against GFP was from Clontech (Mountain View CA), and that against ORP150 was from our laboratory stocks (38). Terminal deoxynucleotidyl-transferase was obtained from TOYOBO (Osaka, Japan). Biotin 14-ATP, Alexa Fluor 488 goat anti-rabbit (or anti-mice) immunoglobulin G, Alexa Fluor 594 goat anti-rat immunoglobulin G, Lipofectamine (TM2000), Dynabeads Protein G, and Alexa Fluor 488 conjugated with streptavidin were purchased from Invitrogen. Mounting medium for immunohistochemical analysis (VECTASHIELD) was from Vector Laboratories (Burlingame, CA). 4',6-diamidino-2-phenylindole dihydrochloride (DAPI) was from Dojindo (Kumamoto, Japan). The RNeasy kit and HiPerFect transfection reagents were obtained from Qiagen (Valencia, CA), the first-strand cDNA synthesis kit was from Takara (Kyoto, Japan), and the iQ SYBR Green Supermix was from Bio-Rad. The Dual Luciferase Assay System was from Promega (Southampton, UK). Mice heterozygous for a truncated/inactivated mutant form of ORP150 (ORP150^{+/-}) and their wild-type counterparts (ORP150^{+/+}) (6–8 weeks of age) were prepared as described previously (38, 39). The experiments and procedures described here were carried out in accordance with the Guide for the Care and Use of Laboratory Animals as adopted and promulgated by the National Institutes of Health (Bethesda, MD) and were approved by the Animal Care Committee of Kumamoto University.

***H. pylori* Inoculation and Gastric Damage Assay**—*H. pylori* inoculation was done as described previously (40) with some modifications. *H. pylori* strain ATCC43504 (CagA⁺ and

H. pylori and ER Chaperones

VacA⁺) (a gift from Dr. Oguma, Okayama University) was cultured in a brain heart infusion broth containing 10% horse serum at 37 °C under a microaerophilic atmosphere. *H. pylori* were also cultured on brain heart infusion agar supplemented with heat-inactivated 7% horse blood or on *Helicobacter* selection agar under a gas-pack jar with an Anaeropack Bikouki. Mice were orally inoculated with *H. pylori* at a dose of 2.0×10^8 *H. pylori*/animal/0.5 ml of PBS every second day for 6 days (total 3 times). *H. pylori* lysates were prepared by sonication of cells in PBS. The protein concentration of lysates was determined by the Bradford method (41).

The gastric ulcerogenic response was examined as described previously (18), with some modifications. *H. pylori*-inoculated or non-inoculated mice (1 day after the final inoculation of *H. pylori*) fasted for 6 h were orally administered with indomethacin (10 mg/kg, 10 ml 1% methylcellulose/kg). Twelve hours later, the animals were sacrificed, after which their stomachs were removed, and the areas of gastric mucosal lesions were measured by an observer unaware of the treatment they had received. Calculation of the scores involved measuring the area of all the lesions in square millimeters and summing the values to give an overall gastric lesion index.

Cell Culture—AGS and MKN45 are human adenocarcinoma gastric cell lines. Cells were cultured in RPMI1640 medium supplemented with 10% FBS, 100 units/ml penicillin, and 100 µg/ml streptomycin in a humidified atmosphere of 95% air with 5% CO₂ at 37 °C.

Transfections were carried out using Lipofectamine (TM2000) according to the manufacturer's instructions. Cells were used for experiments after a 24-h recovery period. Transfection efficiency was determined in parallel plates by transfection of cells with a pEGFP-N1 control vector. Transfection efficiencies were greater than 80% in all experiments. The plasmid pCMVshort-EGFP-ATF6α (a gift from Dr. Mori, Kyoto University) (42) was transfected into AGS cells, according to the manufacturer's protocols.

siRNA Targeting of Genes—The siRNAs for ATF6 and ATF4 and nonspecific siRNA were purchased from Qiagen. AGS cells were transfected with siRNA using HiPerFect transfection reagent according to the manufacturer's instructions.

Pulse-Chase Analysis—A pulse-chase experiment was carried out with 0.1 mCi/ml Expre^{35S} protein labeling mix (PerkinElmer Life Sciences) as described previously (43), with some modifications. Cells were labeled with [³⁵S]methionine and [³⁵S]cysteine in methionine- and cysteine-free RPMI1640 medium for 30 min. To chase labeled proteins, cells were washed with fresh complete medium three times and incubated in fresh complete medium with or without *H. pylori*. ATF6 was immunoprecipitated with its antibody and separated by SDS-polyacrylamide gel electrophoresis followed by autoradiography (Fuji BAS 2500 imaging analyzer).

Immunostaining—Plasmids (pCMVshort-EGFP-ATF6α) were transfected into AGS cells, and cells were co-cultured with *H. pylori* in the Lab-Tek II chamber slide system (Nalge Nunc International, Rochester, NY). Cells were fixed in 4% paraformaldehyde for 20 min, blocked with goat serum for 15 min, then incubated for 12 h with antibody against GRP94 in the presence of 2.5% bovine serum albumin before finally being incubated for

2 h with Alexa Fluor 594 goat anti-rat IgG. Samples were mounted with VECTASHIELD. Images were captured on a confocal laser-scanning fluorescence microscope (FLUOVIEW FV500-IX-UV, Olympus).

RT-PCR and Real-time RT-PCR Analyses—Real-time RT-PCR was performed as previously described with some modifications (44). Total RNA was extracted from gastric tissues or cultured cells using an RNeasy kit according to the manufacturer's protocol. Samples (2.5 µg RNA) were reverse-transcribed using a first-strand cDNA synthesis kit. Synthesized cDNA was used in real-time RT-PCR (Chromo 4 instrument; Bio-Rad) experiments using iQ SYBR GREEN Supermix and analyzed with Opticon Monitor Software. Specificity was confirmed by electrophoretic analysis of the reaction products and by the inclusion of template- or reverse transcriptase-free controls. The cycle conditions were 2 min at 50 °C followed by 10 min at 90 °C and finally 45 cycles of 95 °C for 30 s and 63 °C for 60 s. To normalize the amount of total RNA present in each reaction, actin cDNA was used as an internal standard.

Primers were designed using the Primer3 website and are listed as forward and reverse, respectively. Human primers were: *atf4* (5'-tcaaacctcatgggttctcc-3') and (5'-gtgtcatcaacgtggctcag-3'); *atf6*, (5'-ctccgagatcagcagaggaa-3') and (5'-aatgactcagggatggtgct-3'); *chop*, (5'-tgcctttctctcggact-3') and (5'-tgtgacctctgctggtctg-3'); *grp78*, (5'-tagcgtatggtgctgctg-3') and (5'-ttgtcaggggtcttacc-3'); *orp150*, (5'-gaagatgcagagcccatttc-3') and (5'-tctgctccaggacctctaa-3'); *xbp-1*, (5'-aacagagtagcagcgcagactg-3') and (5'-ggatcttaaacatagaggcttggtg-3'); *xbp-1* (*u*), (5'-agcactcagactacgtgac-3') and (5'-ccagaatgcccaacaggata-3'); *actin*, (5'-ggacttcgagcaagagatgg-3') and (5'-agcattgttggtcgctacg-3'). Mouse primers were: *atf6* (5'-catcaaaagctctcggtc-3') and (5'-gggtcgtctctgtggtggtt-3'); *grp78*, (5'-gcttcgataatcagccaac-3') and (5'-gcaggaggaattccagta-3'); *orp150*, (5'-cagactgaagaggcgaacc-3') and (5'-ttcctgttcaggctcagctc-3'); *chop*, (5'-acagaggtcacacgcacatc-3'), and (5'-gggactgacctctgttt-3'); *gapdh*, (5'-tgcctttctctcggact-3') and (5'-tgtgacctctgctggttctg-3').

For regular RT-PCR, we used an initial denaturation step of 94 °C for 1 min followed by 30 cycles of denaturation at 94 °C for 30 s, annealing at 60 °C for 20 s, and elongation at 72 °C for 1 min. A final elongation step at 72 °C for 10 min completed the RT-PCR. The amplified PCR products were separated by 3% agarose gel electrophoresis and then visualized with ethidium bromide.

Immunoblotting Analysis—Total protein and nuclear protein extracts were prepared as described previously (20). The protein concentration of each sample was determined by the Bradford method (41). Samples were applied to polyacrylamide SDS gels and subjected to electrophoresis, after which the proteins were immunoblotted with appropriate antibodies.

Histological, Immunohistochemical, and Terminal Deoxynucleotidyltransferase-mediated Biotinylated UTP Nick End Labeling (TUNEL) Analyses—Gastric tissue samples were fixed in 4% buffered paraformaldehyde and embedded in paraffin before being cut into 4-mm sections.

For histological examination (hematoxylin and eosin (H&E) staining), sections were stained first with Mayer's hematoxylin and then with 1% eosin alcohol solution. Samples were

mounted with Malinol and inspected with the aid of an Olympus BX51 microscope.

For immunohistochemical analysis, sections were blocked with 2.5% goat serum for 10 min, incubated for 12 h with antibody against ORP150 or GRP78 in the presence of 2.5% bovine serum albumin, and finally incubated for 2 h with Alexa Fluor 488 goat anti-rabbit immunoglobulin G in the presence of DAPI (5 mg/ml). Samples were mounted with VECTASHIELD and inspected using fluorescence microscopy (Olympus BX51).

For TUNEL assays, sections were incubated first with proteinase K (20 mg/ml) for 15 min at 37 °C, then with terminal deoxynucleotidyltransferase and biotin 14-ATP for 1 h at 37 °C and finally with Alexa Fluor 488 conjugated with streptavidin for 1 h. Samples were mounted with VECTASHIELD and inspected using fluorescence microscopy (Olympus BX51).

Luciferase Assay—The pGL-3/ERSE (ER stress response element) plasmid, which was constructed by inserting ERSE (5'-ccaatcagaagtggcagc-3') just upstream of the luciferase gene (45), was kindly provided by Dr. Gotoh (Kumamoto University). The pGL-3/*grp78*pro plasmid, which was constructed by inserting the human *grp78* promoter (from -304 to +7 region) into the same region (46), was generously provided by Dr. Mori (Kyoto University).

The luciferase assay was performed as described previously (47). Cells were transfected with 1 μg of one of the *Photinus pyralis* luciferase reporter plasmids (pGL-3 or its derivatives) and 0.125 μg of the internal standard plasmid bearing the *Renilla reniformis* luciferase reporter (pRL-SV40). *P. pyralis* luciferase activity in cell extracts was measured using the Dual Luciferase Assay System and then normalized for *R. reniformis* luciferase activity.

Statistical Analysis—Two-way analysis of variance followed by the Tukey test or the Student's *t* test for unpaired results was used to evaluate differences between more than three groups or between two groups, respectively. Differences were considered to be significant for values of *p* < 0.05.

RESULTS

Effect of *H. pylori* on Expression of ER Chaperones in Vitro—It was previously reported that co-culture of AGS cells with *H. pylori* at a bacteria:cell ratio of 200:1 causes partial induction of apoptosis (48, 49). In the current study we examined the expression of ER chaperones under these conditions and, using MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide) assay, determined cell viability to be 80% after a 24-h incubation of AGS cells with *H. pylori* (data not shown). As shown in Fig. 1, *A* and *B*, treatment of cells with *H. pylori* decreased the levels of ORP150 and GRP78. Similar results were observed in another gastric cell line (MKN45) (Fig. 1*C*). A slight reduction of levels of other ER chaperones (such as GRP94 and protein disulfide isomerase but not GRP58) was also observed in cells treated with *H. pylori* (Fig. 1, *A* and *B*). We also found that treatment of AGS cells with *H. pylori* cell lysates also decreased the levels of ORP150 and GRP78; however, the extent of these decreases was not as apparent as that observed with *H. pylori* (Fig. 1, *D* and *E*). Unlike the ER chaperones, the level of CHOP was increased by co-culture of cells with *H. pylori* (Fig. 1, *A* and *B*).

Real-time RT-PCR analysis revealed that the *H. pylori*-dependent down-regulation of expression of ORP150 and GRP78 and up-regulation of expression of CHOP was also observed at the level of mRNA (Fig. 2*A*). We also performed a luciferase reporter assay using a reporter plasmid where the promoter of the *grp78* gene was inserted upstream of the luciferase gene (pGL-3/*grp78*pro). As shown in Fig. 2*B*, co-culture of cells with *H. pylori* decreased the luciferase activity in cells with pGL-3/*grp78*pro but not in those with the control vector (pGL-3). These results suggest that co-culture of cells with *H. pylori* inhibits the transcription of ER chaperone genes.

Mechanism for Suppression of Expression of ER Chaperones by *H. pylori*—To understand the molecular mechanism governing suppression of expression of ER chaperones by *H. pylori*, we first examined the effect of *H. pylori* on the level of ER stress response-related transcription factors (ATF6, ATF4, and XBP-1). Co-culture of cells with *H. pylori* decreased the level of ATF6 protein (p90-ATF6) but not *atf6* mRNA (Figs. 1, *A* and *B*, and 2*C*). We also found that the level of p50-ATF6 in nuclear extracts decreased after co-culture of cells with *H. pylori* (Fig. 1, *F* and *G*). On the other hand, the levels of ATF4 protein and *atf4* mRNA did not alter after co-culture with *H. pylori* (Figs. 1, *A* and *B*, and 2*C*). Furthermore, although treatment of cells with indomethacin increased the level of the spliced form (active form) of *xbp-1* mRNA, treatment of cells with *H. pylori* did not result in a similar observation (Fig. 2*D*). We also found that treatment of cells with indomethacin or *H. pylori* decreases or increases, respectively, the level of un-spliced form (inactive form) of *xbp-1* mRNA by real-time RT-PCR analysis (Fig. 2*E*). These results suggest that ATF6, rather than ATF4 and XBP-1, is responsible for suppression of expression of ER chaperones by *H. pylori*.

To confirm this idea, we performed a luciferase assay using a reporter plasmid where the ATF6 binding consensus sequence, ERSE, was inserted upstream of the luciferase gene (pGL-3/ERSE). As shown in Fig. 2*B*, treatment of cells with *H. pylori* decreased the luciferase activity in cells with pGL-3/ERSE, suggesting that the transcriptional activity of ATF6 decreased after treatment of cells with *H. pylori*. This is consistent with the observation that the level of p50-ATF6 decreased in *H. pylori*-treated cells (Fig. 1, *F* and *G*).

For further confirmation of the idea that ATF6 plays an important role in *H. pylori*-dependent suppression of expression of ER chaperones, we examined the effect of siRNA for ATF6 on the expression of ER chaperones. As shown in Fig. 3, transfection of cells with siRNA for ATF6 suppressed expression of not only ATF6 but also ORP150 and GRP78, affecting both mRNA and protein levels. The transfection did not affect the expression of *chop* mRNA (Fig. 3*A*). On the other hand, transfection of cells with siRNA for ATF4 suppressed expression of ATF4 but not that of ORP150 and GRP78 (supplemental Fig. S1). The results in Fig. 3 suggest that the reduction in the level of ATF6 is partly involved in the *H. pylori*-dependent suppression of expression of ER chaperones but not for induction of expression of CHOP.

We also examined the effect of *H. pylori* cell lysates on levels of p90-ATF6 and p50-ATF6. As shown in Fig. 1, *D*–*G*, levels of p90-ATF6 and p50-ATF6 were slightly decreased by

H. pylori and ER Chaperones

600DPI
7/2




FIGURE 1. Down-regulation of expression of ER chaperones by *H. pylori*. AGS (A, B, and D-G) or MKN-45 (C) cells were co-cultured with *H. pylori* at a bacteria:cell ratio of 200:1 (A-C, F, and G) or with *H. pylori* lysates (D-G) for the indicated periods. Whole cell extracts (A, C, and D) or nuclear extracts (F) were analyzed by immunoblotting with an antibody against ORP150, GRP78, CHOP, ATF6, ATF4, GRP94, GRP58, protein disulfide isomerase (PDI), actin, or lamin. The intensity of each band in three independent experiments (one of them is shown in A, D, and F) was determined and expressed relative to the control (B, E, and G, respectively). Values are the mean \pm S.D. ($n = 3$). **, $p < 0.01$; *, $p < 0.05$.

H. pylori and ER Chaperones




FIGURE 2. Inhibitory effects of *H. pylori* on transcription of ER stress response-related genes. AGS cells were co-cultured with *H. pylori* at a bacteria:cell ratio of 200:1 for the indicated periods (A and C–E) or treated with indomethacin (IND) for 24 h (D and E). A, C, and E, the relative expression of each gene was monitored by real-time RT-PCR using a specific primer for each gene. Values normalized to the *actin* gene are expressed relative to the control sample. B, AGS cells were co-transfected with pRL-SV40 (internal control plasmid carrying the *R. reniformis* luciferase gene) and pGL-3 or its derivatives (pGL-3/*grp78pro* and pGL-3/ERSE) and cultured for 24 h. Cells were then co-cultured with or without *H. pylori* at a bacteria:cell ratio of 200:1 for 24 h, and *P. pyralis* luciferase activity was measured and normalized for *R. reniformis* luciferase activity. The 100% value of the *P. pyralis* luciferase activity is 5.4×10^6 , 7.4×10^5 , or 2.4×10^4 units for pGL-3/*grp78pro*, pGL-3/ERSE, or pGL-3, respectively. D, RT-PCR was performed with total RNAs and primer sets for detecting the un-spliced (*xbp-1(u)*) and spliced (*xbp-1(s)*) forms of *xbp-1* mRNA, which were separated by agarose gel electrophoresis. Values are the mean \pm S.D. ($n = 3$). **, $p < 0.01$; *, $p < 0.05$; n.s., not significant.

treatment of cells with *H. pylori* cell lysates, but the decrease occurred more slowly than that of ORP150 and GRP78.

As described above, *atf6* mRNA expression was not affected by *H. pylori* (Fig. 2C). Thus, either suppression of translation or post-translational modification of ATF6 (such as degradation by proteases) may be responsible for the observed reduction in the level of ATF6 after co-culture of cells with *H. pylori*. To address this issue, we first examined the effect of *H. pylori* on the level of ATF6 in cells pretreated with cycloheximide, an inhibitor of protein synthesis. As shown in Fig. 4A, the *H. pylori*-

dependent decrease in the level of ATF6 was observed even in cells pretreated with cycloheximide. We also found that the *H. pylori*-dependent decrease in the level of ATF6 was observed for GFP-ATF6, whose expression is regulated by the strong cytomegalovirus promoter (Fig. 4B). We also examined the effect of *H. pylori* on the stability of p90-ATF6 by the pulse-chase experiment. As shown in Fig. 4, C and D, the labeled p90-ATF6 disappeared more rapidly in the presence of *H. pylori* treatment than its absence. These results suggest that post-translational modification of ATF6, such as protein degradation, is responsible for the lower level of ATF6 observed after treatment of cells with *H. pylori*.

In addition to cleavage by S1P and S2P, it is known that ATF6 is continuously degraded by the proteasome-ubiquitin pathway (50, 51). Thus, using specific inhibitors, we examined the contribution of these systems to the *H. pylori*-dependent decrease in the level of ATF6. As shown in Fig. 4E, an inhibitor of the proteasome-ubiquitin system, epoxomycin, weakly suppressed the *H. pylori*-dependent decrease in the level of ATF6. On the other hand, an inhibitor of S1P, 4-(2-aminoethyl)-benzenesulfonyl fluoride (AEBSF), did not affect the level of ATF6 in the presence of *H. pylori* (Fig. 4F). Furthermore, we found that inhibitors of lysosomal proteases (pepstatin A (an inhibitor of aspartate proteases) and E-64-d (an inhibitor of cysteine protease)) also weakly suppressed the *H. pylori*-dependent decrease in the level of ATF6 (Fig. 4G). Interestingly, combination of epoxomycin and inhibitors of lysosomal proteases resulted in clear

suppression of the *H. pylori*-dependent decrease in the level of ATF6 (Fig. 4H). The results in Fig. 4 suggest that *H. pylori* decreases the level of ATF6 partly through modulation of its degradation by the proteasome-ubiquitin and lysosomal systems.

We also examined the effect of *H. pylori* on subcellular localization of ATF6 using GFP-ATF6. As shown in Fig. 4I, GFP-ATF6 co-localized with GRP94 (ER marker). Although the level of GFP-ATF6 was decreased, the localization of ATF6 was not clearly affected by treatment of cells with *H. pylori* (Fig. 4I).

H. pylori and ER Chaperones




FIGURE 3. Effect of siRNA for ATF6 on expression of ER chaperones. AGS cells were transfected with siRNA for ATF6 (siATF6) or nonspecific siRNA (ns) and were incubated for 48 h (A) or 72 h (B). The mRNA (A) and protein (B and C) expression was monitored and expressed as described in the legends of Figs. 1 and 2. Values are the mean \pm S.D. ($n = 3$). **, $p < 0.01$; *, $p < 0.05$; n.s., not significant.

Effect of H. pylori on the Gastric Ulcerogenic Response and Expression of ER Chaperones Induced by Indomethacin in Mice—To evaluate the *in vivo* relevance of our *in vitro* observation that *H. pylori* suppress the expression of ER chaperones, we first examined the effect of oral inoculation of *H. pylori* on the expression of ER chaperones at gastric mucosa. We monitored by real-time RT-PCR analysis the mRNA expression of ER chaperones and CHOP after administration of indomethacin and/or *H. pylori* in gastric tissues. Oral inoculation of *H. pylori* to mice suppressed the background (without indomethacin administration) expression of *orp150* and *grp78* mRNAs but not *chop* mRNA (Fig. 5A). Indomethacin administration up-regulated the expression of *orp150*, *grp78*, and *chop* mRNAs, whereas the expression of *orp150* and *grp78* mRNAs but not that of *chop* mRNA was significantly suppressed by prior administration of *H. pylori* (Fig. 5A). Immunohistochemical analyses also demonstrated that oral inoculation with *H. pylori* decreased the levels of ORP150 and GRP78 at gastric mucosa in both the presence and absence of indomethacin administration (Fig. 5B). We consider that the staining of ORP150 and GRP78 in Fig. 5B is specific, because no positive staining was observed without a primary antibody (supplemental Fig. S2). We also performed immunoblotting analysis, and suppression of the gastric expression of ORP150 and GRP78 by inoculation with *H. pylori* in both the presence and absence of indomethacin treatment was confirmed (Fig. 5, C and D). Indomethacin-induced expression of GRP78 and CHOP was also confirmed (Fig. 5, C and D). Furthermore, we found that the gastric level of p90-ATF6 was decreased by inoculation with *H. pylori* in both the presence and absence of indomethacin treatment (Fig. 5, C and D). We also found that there is a tendency that inoculation with *H. pylori* decreases the levels of ORP150 and GRP78 in the presence of indomethacin treatment in the small intestine but not other organs (supplemental Fig. S3A).

The observation that inoculation with *H. pylori* reduces the gastric expression of ER chaperones suggests that this inoculation stimulates protein aggregation in ER in gastric cells. To

address this idea, we examined the effect of *H. pylori* inoculation on the gastric level of connexin 43, which is known to be degraded by ER-associated degradation when it is aggregated in the ER (52, 53). As shown in supplemental Fig. S3, B and C, the gastric level of connexin 43 was decreased by *H. pylori* inoculation, suggesting that *H. pylori* stimulate aggregation of the protein in ER.

The effect of pre-administration of *H. pylori* on the development of gastric lesions after oral administration of indomethacin was examined. Administration of indomethacin produced gastric lesions, and this lesion production was significantly enhanced by pre-administration of *H. pylori* (Fig. 6A). Administration of *H. pylori* alone did not significantly produce gastric lesions under the conditions used (Fig. 6A). Histological analysis with H&E staining also supported the notion that indomethacin-produced gastric lesions are exacerbated by pre-administration of *H. pylori* (Fig. 6B).

As mentioned above, gastric mucosal cell death plays an important role in the formation of NSAID-induced gastric lesions. We, therefore, examined the effect of pre-administration of *H. pylori* on this process. The level of gastric mucosal cell death was determined by TUNEL assay. An increase in the number of TUNEL-positive cells was observed after indomethacin administration, and this increase was more apparent in mice pre-administered with *H. pylori* than in control mice (Fig. 6C). We also examined the effect of *H. pylori* on indomethacin-induced expression of ORP150, GRP78, and p90-ATF6 or apoptosis *in vitro*. Treatment of cells with *H. pylori* decreased the levels of these proteins and increased apoptotic cells in both the presence and absence of indomethacin treatment (supplemental Fig. S4, A and B). These results suggest that *H. pylori* exacerbate indomethacin-induced gastric lesion formation by stimulating indomethacin-induced gastric mucosal cell death.

The results in Figs. 5 and 6 suggest that *H. pylori* exacerbates indomethacin-induced gastric lesion formation through down-regulation of expression of ER chaperones. To test this idea using a genetic approach, the development of gastric lesions after oral administration of indomethacin was compared between heterozygous ORP150-deficient mice (ORP150^{-/+}) and wild-type mice (ORP150^{+/+}). Indomethacin-induced gastric lesions were significantly worse in heterozygous ORP150-deficient mice than in wild-type controls (Fig. 7A). ORP150 deficiency did not affect the background level of gastric lesions (Fig. 7A). Immunohistochemical analyses confirmed that the level of ORP150 in gastric mucosa was lower in heterozygous ORP150-deficient mice than wild-type mice in both the presence and absence of indomethacin administration (Fig. 7B). These results show that ORP150 plays an important role




FIGURE 4. Mechanism for the *H. pylori*-dependent decrease in the level of ATF6. A and E–H, AGS cells were pre-incubated with or without each drug for 1 h and further incubated with or without *H. pylori* at a bacteria:cell ratio of 200:1 for 24 h in the presence (E, G, and H) or absence (A and F) of the same concentration of each drug as in the preincubation step. B and I, AGS cells were transfected with pCMVshort-EGFP-ATF6 α (42) and co-cultured with or without *H. pylori* at a bacteria:cell ratio of 200:1 for 24 h (B) or indicated periods (I). A, B, and E–H, whole cell extracts were analyzed by immunoblotting (WB) with an antibody against GFP, ATF6, or actin. C, AGS cells were pulse-labeled for 30 min with [³⁵S]methionine and [³⁵S]cysteine and then chased with excess amounts of cold methionine and cysteine for the indicated periods in the absence or presence of *H. pylori* at a bacteria:cell ratio of 200:1. Labeled proteins were extracted, immunoprecipitated with antibody against ATF6, subjected to SDS-PAGE, and autoradiographed. D, the band intensity of p90-ATF6 was determined and expressed relative to the control. I, cells were fixed, stained with antibody against GRP94, and analyzed by confocal laser-scanning fluorescence microscope (magnification, 600 times). Values are the mean \pm S.D. ($n = 3$). **, $p < 0.01$; *, $p < 0.05$. AEBSF, 4-(2-aminoethyl)benzenesulfonyl fluoride.

in protecting gastric mucosa against indomethacin-induced lesions.

We also examined the indomethacin-induced gastric mucosal cell death in heterozygous ORP150-deficient mice. Indomethacin-induced gastric mucosal cell death was more apparent in heterozygous ORP150-deficient mice than in wild-type mice (Fig. 7C). These results suggest that ORP150 protects the gastric mucosa from indomethacin-induced cell death. Combining the results in Figs. 5–7, we consider that *H. pylori* exacerbates indomethacin-induced gastric lesion formation partly through down-regulation of ER chaperones and

the resulting stimulation of indomethacin-induced gastric mucosal cell death.

DISCUSSION

There have been contradictory reports about whether infection with *H. pylori* increases the risk of developing NSAID-induced gastric lesions (in other words, whether eradication of *H. pylori* reduces the risk of developing NSAID-induced gastric lesions) (1–6, 54). This may be due to differences in diagnostic criteria (endpoints), standards for patient recruitment, and populations used for these studies. The most we can conclude is

H. pylori and ER Chaperones




FIGURE 5. Effect of *H. pylori* on expression of ER chaperones at gastric mucosa. *H. pylori* were orally inoculated into mice (C57/BL6) at a dose of 2.0×10^8 *H. pylori*/animal every second day for 6 days (total 3 times). One day after the final inoculation, *H. pylori*-inoculated or control mice were orally administered 10 mg/kg of indomethacin (IND), and their stomachs were removed after 12 h. A, total RNA was extracted and subjected to real-time RT-PCR using a specific primer for each gene. Values normalized to the *gapdh* gene are expressed relative to the control sample. B, sections of gastric tissues were subjected to immunohistochemical analysis with an antibody against ORP150 or GRP78 and DAPI staining (magnification, 200 times). C and D, protein expression was monitored and expressed as described in the legend of Fig. 1. Values are given as the mean \pm S.E. **, $p < 0.01$; *, $p < 0.05$; n.s., not significant.

H. pylori and ER Chaperones




FIGURE 6. Effect of *H. pylori* on indomethacin-induced gastric lesions and mucosal cell death. Mice (C57/BL6) were administered *H. pylori* and indomethacin (IND), as described in the legend of Fig. 5. *A*, the stomach was scored for hemorrhagic damage. Values are the mean \pm S.E. * $p < 0.05$; n.s., not significant. *B*, sections of gastric tissues were subjected to H&E staining (magnification, 200 times). *C*, sections of gastric tissues were subjected to TUNEL assay and DAPI staining (magnification, 200 times). n.s., not significant.

that under certain conditions infection with *H. pylori* increases the risk of developing NSAID-induced gastric lesions. Thus, it is important to examine the effect of *H. pylori* on factors that affect the formation of NSAID-induced gastric lesions. In this study we have focused on ER chaperones and found that co-culture of gastric cells with *H. pylori* decreases the level of ER chaperones. This is the first observation that *H. pylori* affect the expression of ER stress response-related proteins. However, although we used here the transient infection model of *H. pylori*, the infection in humans is chronic. The *H. pylori* strain used in this study does not colonize mice (data not shown), and thus, studies in the future need to be done with strains that do colonize mice.

By using real-time RT-PCR and luciferase reporter assays, we have shown that the *H. pylori*-dependent decrease in the level of ER chaperones *in vitro* is regulated at the level of transcription. Of three ER stress response-related transcription factors (ATF6, ATF4, and XBP-1), only the level of ATF6 (but not the *atf6* mRNA) was decreased by co-culture of cells with *H. pylori*, suggesting that ATF6 is involved in the *H. pylori*-dependent suppression of transcription of ER chaperone genes. Because the *H. pylori*-dependent decrease in the level of ER chaperones was observed in cells whose protein synthesis was inhibited and *H. pylori* decreased the stability of p90-ATF6, post-translational modification (protein degradation) of ATF6 would be involved in this process. Analysis with each inhibitor suggested that the proteasome-ubiquitin system rather than degradation

by S1P is involved in this degradation of ATF6. The observation that the level of p50-ATF6 (the proteolytic product of S1P and S2P) did not increase after co-culture of cells with *H. pylori* further supports this notion. We also suggest that protein degradation in lysosomes is involved in this degradation of ATF6. It is known that VacA perturbs endocytic traffic at a late stage (55, 56), and as such it is possible that *H. pylori* affects the traffic of ATF6 to lysosomes and its degradation in lysosomes. Furthermore, because the suppression of *H. pylori*-dependent degradation of ATF6 by inhibitors of proteasomal and lysosomal proteases was weak, other proteases seem to be involved in this degradation.

We found that not only *H. pylori* themselves but also cell extracts of *H. pylori* suppress the expression of ER chaperones *in vitro*. However, the suppression of expression of ER chaperones by cell extracts of *H. pylori* was not as great as that induced by *H. pylori* themselves, and cell extracts of *H. pylori* did not affect the level of ATF6 so distinctly (Fig.

1, D–G). Furthermore, the decrease in level of ATF6 occurred more slowly than that of ORP150 and GRP78 in the presence of cell extracts of *H. pylori* (Fig. 1), suggesting that the decrease in levels of ORP150 and GRP78 observed with cell extracts of *H. pylori* is not due to the decrease in levels of ATF6. In other words, results suggest that the mechanism for the decrease in levels of ORP150 and GRP78 is different between *H. pylori* cells and cell extracts of *H. pylori*. We also compared the signal pathway for induction of apoptosis between *H. pylori* cells and cell extracts of *H. pylori*. As shown in supplemental Fig. S5, the decrease in Bax and increase in cytochrome *c* in cytosol fractions (an indicator for mitochondria-mediated apoptosis) and induction of expression of CHOP were not observed with apoptosis induced by cell extracts of *H. pylori* so apparently as that induced by *H. pylori* cells, suggesting that the signal pathway for induction of apoptosis is also different between *H. pylori* cells and cell extracts of *H. pylori*. Although siRNA for ATF6 suppressed the expression of ER chaperones, the extent of suppression was not as apparent as that seen with *H. pylori*. These results suggest that in addition to the mechanism described above (the ATF6-mediated mechanism), an ATF6-independent and as yet unknown mechanism that can be reproduced with cell extracts of *H. pylori* should also be mainly involved in the *H. pylori*-dependent suppression of expression of ER chaperones.

We have previously reported that suppression of expression of GRP78 and ORP150 by siRNA stimulated NSAID-induced

H. pylori and ER Chaperones




FIGURE 7. Indomethacin-induced gastric lesions and mucosal cell death in heterozygous ORP150-deficient mice. Heterozygous ORP150-deficient mice (ORP150^{+/-}) and wild-type mice (ORP150^{+/+}) were orally administered with 10 mg/kg of indomethacin (IND), and their stomachs were removed after 12 h. Gastric lesions (A), expression of ORP150 (B), and mucosal apoptosis (C) were assayed as described in the legends of Figs. 5 and 6. Values are mean \pm S.E. **, $p < 0.01$; *, $p < 0.05$; n.s., not significant. n.s., not significant.

apoptosis in cultured gastric cells (19, 20). We have also suggested that NSAID-induced apoptosis at gastric mucosa plays an important role in the formation of NSAID-induced gastric lesions (16–18). These results suggest that ER chaperones play a protective role against the development of NSAID-induced gastric lesions; however, there is no direct evidence supporting this notion. In this study we have shown that heterozygous ORP150-deficient mice display phenotypes sensitive to indomethacin-induced gastric mucosal cell death and gastric lesion formation. This is the first genetic evidence that an ER chaperone is protective against NSAID-induced gastric lesion formation. This result also suggests that inducers of ER chaperones may be therapeutically beneficial against NSAID-induced gastric lesions, as is the case for heat shock proteins inducers (18, 57).

The *in vitro* observation that expression of ER chaperones is suppressed by *H. pylori* suggests that *H. pylori* would suppress the expression of GRP78 and ORP150 at the gastric mucosa and stimulate NSAID-induced gastric mucosal cell death and lesion formation. In fact, we have shown that pre-inoculation of mice with *H. pylori* not only suppresses the expression of ER chaperones

ones but also stimulates NSAID-induced cell death and gastric lesion formation.

We also showed *in vitro* that co-culture of gastric cells with *H. pylori* up-regulates the expression of CHOP, suggesting that this up-regulation is involved in the *H. pylori*-dependent stimulation of NSAID-induced cell death. There are two possible mechanisms that could explain this up-regulation of CHOP. One is that *H. pylori* directly affects the expression of CHOP. However, because siRNA for ATF6 did not up-regulate the expression of CHOP, *H. pylori*-dependent degradation of ATF6 must not be involved. The other possibility is that this up-regulation is a result of the suppression of expression of ER chaperones, as we previously reported that suppression of expression of GRP78 and ORP150 by siRNA induces the expression of CHOP in the presence of NSAIDs (19, 20).

Although we suggest that the *H. pylori*-dependent exacerbation of indomethacin-induced gastric lesion formation is mediated by the suppression of expression of ER chaperones, various other mechanisms are likely to be involved in this exacerbation. For example, cytotoxic proteins produced by *H. pylori*, such as VacA and CagA, which

are known to induce apoptosis in gastric cells (32–36), may stimulate indomethacin-induced cell death, resulting in exacerbation of indomethacin-induced gastric lesions. CagA disrupts the epithelial apical junction complex (58), which may also be involved in *H. pylori*-dependent exacerbation of indomethacin-induced gastric lesions. We believe that this animal model for *H. pylori*-dependent exacerbation of indomethacin-induced gastric lesion formation will be useful in future studies for examining the relationship between *H. pylori* and NSAIDs and their involvement in the production of gastric lesions.

Acknowledgments—We thank Dr. Oguma (Okayama University) and Drs. Gotoh (Kumamoto University) and Mori (Kyoto University) for generously providing the *H. pylori* strain or plasmids, respectively.

REFERENCES

- Huang, J. Q., Sridhar, S., and Hunt, R. H. (2002) *Lancet* 359, 14–22
- Chan, F. K., Sung, J. J., Chung, S. C., To, K. F., Yung, M. Y., Leung, V. K., Lee, Y. T., Chan, C. S., Li, E. K., and Woo, J. (1997) *Lancet* 350, 975–979
- Murakami, K., Okimoto, T., Kodama, M., Tanahashi, J., Yasaka, S., Inoue, K., Uchida, M., Anan, J., Mizukami, K., Abe, T., Watada, M., and Fujioka,

- T. (2009) *J Gastroenterol* **44**, 40–43
4. Papatheodoridis, G. V., Sougioultzis, S., and Archimandritis, A. J. (2006) *Clin. Gastroenterol Hepatol.* **4**, 130–142
 5. Chan, F. K., Chung, S. C., Suen, B. Y., Lee, Y. T., Leung, W. K., Leung, V. K., Wu, J. C., Lau, J. Y., Hui, Y., Lai, M. S., Chan, H. L., and Sung, J. J. (2001) *N. Engl. J. Med.* **344**, 967–973
 6. de Leest, H. T., Steen, K. S., Lems, W. F., Bijlsma, J. W., van de Laar, M. A., Huisman, A. M., Vonkeman, H. E., Houben, H. H., Kadir, S. W., Kostense, P. J., van Tulder, M. W., Kuipers, E. J., Boers, M., and Dijkmans, B. A. (2007) *Helicobacter* **12**, 477–485
 7. Kanatani, K., Ebata, M., Murakami, M., and Okabe, S. (2004) *J. Physiol. Pharmacol.* **55**, 207–222
 8. Chang, C. C., Chen, S. H., Lien, G. S., Lou, H. Y., Hsieh, C. R., Fang, C. L., and Pan, S. (2005) *World J. Gastroenterol.* **11**, 104–108
 9. Yoshida, N., Sugimoto, N., Hirayama, F., Nakamura, Y., Ichikawa, H., Naito, Y., and Yoshikawa, T. (2002) *Gut* **50**, 594–598
 10. Lichtenberger, L. M. (2001) *Biochem. Pharmacol.* **61**, 631–637
 11. Ligumsky, M., Golanska, E. M., Hansen, D. G., and Kauffman, G. L., Jr. (1983) *Gastroenterology* **84**, 756–761
 12. Tanaka, K., Tomisato, W., Hoshino, T., Ishihara, T., Namba, T., Aburaya, M., Katsu, T., Suzuki, K., Tsutsumi, S., and Mizushima, T. (2005) *J. Biol. Chem.* **280**, 31059–31067
 13. Tsutsumi, S., Gotoh, T., Tomisato, W., Mima, S., Hoshino, T., Hwang, H. J., Takenaka, H., Tsuchiya, T., Mori, M., and Mizushima, T. (2004) *Cell Death Differ.* **11**, 1009–1016
 14. Tomisato, W., Tanaka, K., Katsu, T., Kakuta, H., Sasaki, K., Tsutsumi, S., Hoshino, T., Aburaya, M., Li, D., Tsuchiya, T., Suzuki, K., Yokomizo, K., and Mizushima, T. (2004) *Biochem. Biophys. Res. Commun.* **323**, 1032–1039
 15. Tomisato, W., Tsutsumi, S., Rokutan, K., Tsuchiya, T., and Mizushima, T. (2001) *Am. J. Physiol. Gastrointest. Liver Physiol.* **281**, G1092–G1100
 16. Aburaya, M., Tanaka, K., Hoshino, T., Tsutsumi, S., Suzuki, K., Makise, M., Akagi, R., and Mizushima, T. (2006) *J. Biol. Chem.* **281**, 33422–33432
 17. Tomisato, W., Tsutsumi, S., Hoshino, T., Hwang, H. J., Mio, M., Tsuchiya, T., and Mizushima, T. (2004) *Biochem. Pharmacol.* **67**, 575–585
 18. Suemasu, S., Tanaka, K., Namba, T., Ishihara, T., Katsu, T., Fujimoto, M., Adachi, H., Sobue, G., Takeuchi, K., Nakai, A., and Mizushima, T. (2009) *J. Biol. Chem.* **284**, 19705–19715
 19. Tsutsumi, S., Namba, T., Tanaka, K. I., Arai, Y., Ishihara, T., Aburaya, M., Mima, S., Hoshino, T., and Mizushima, T. (2006) *Oncogene* **25**, 1018–1029
 20. Namba, T., Hoshino, T., Tanaka, K., Tsutsumi, S., Ishihara, T., Mima, S., Suzuki, K., Ogawa, S., and Mizushima, T. (2007) *Mol. Pharmacol.* **71**, 860–870
 21. Ishihara, T., Hoshino, T., Namba, T., Tanaka, K., and Mizushima, T. (2007) *Biochem. Biophys. Res. Commun.* **356**, 711–717
 22. Kaufman, R. J. (2002) *J. Clin. Invest.* **110**, 1389–1398
 23. Ron, D. (2002) *J. Clin. Invest.* **110**, 1383–1388
 24. Yoshida, H., Okada, T., Haze, K., Yanagi, H., Yura, T., Negishi, M., and Mori, K. (2000) *Mol. Cell. Biol.* **20**, 6755–6767
 25. Harding, H. P., Novoa, I., Zhang, Y., Zeng, H., Wek, R., Schapira, M., and Ron, D. (2000) *Mol. Cell* **6**, 1099–1108
 26. Mori, K. (2003) *Traffic* **4**, 519–528
 27. Lou, L. X., Geng, B., Yu, F., Zhang, J., Pan, C. S., Chen, L., Qi, Y. F., Ke, Y., Wang, X., and Tang, C. S. (2006) *Life Sci.* **79**, 1856–1864
 28. Covacci, A., Telford, J. L., Del Giudice, G., Parsonnet, J., and Rappuoli, R. (1999) *Science* **284**, 1328–1333
 29. Suerbaum, S., and Michetti, P. (2002) *N. Engl. J. Med.* **347**, 1175–1186
 30. Hawkey, C. J., Tulassay, Z., Szczepanski, L., van Rensburg, C. J., Filipowicz-Sosnowska, A., Lanás, A., Wason, C. M., Peacock, R. A., and Gillon, K. R. (1998) *Lancet* **352**, 1016–1021
 31. Wu, C. Y., Kuo, K. N., Wu, M. S., Chen, Y. J., Wang, C. B., and Lin, J. T. (2009) *Gastroenterology* **137**, 1641–1648.e1–2
 32. Yamasaki, E., Wada, A., Kumatori, A., Nakagawa, I., Funao, J., Nakayama, M., Hisatsune, J., Kimura, M., Moss, J., and Hirayama, T. (2006) *J. Biol. Chem.* **281**, 11250–11259
 33. Boquet, P., Ricci, V., Galmiche, A., and Gauthier, N. C. (2003) *Trends Microbiol.* **11**, 410–413
 34. Mine, T., Endo, C., Kushima, R., Kushima, W., Kobayashi, I., Muraoka, H., Taki, R., and Fujita, T. (2000) *Aliment. Pharmacol. Ther.* **14**, 199–204
 35. Kim, H., Seo, J. H., and Kim, K. H. (2003) *Ann. N.Y. Acad. Sci.* **1010**, 90–94
 36. Shibata, W., Hirata, Y., Maeda, S., Ogura, K., Ohmae, T., Yanai, A., Mitsuno, Y., Yamaji, Y., Okamoto, M., Yoshida, H., Kawabe, T., and Omata, M. (2006) *J. Pathol.* **210**, 306–314
 37. Yeo, M., Park, H. K., Kim, D. K., Cho, S. W., Kim, Y. S., Cho, S. Y., Paik, Y. K., and Hahm, K. B. (2004) *Proteomics* **4**, 3335–3342
 38. Kitao, Y., Ozawa, K., Miyazaki, M., Tamatani, M., Kobayashi, T., Yanagi, H., Okabe, M., Ikawa, M., Yamashita, T., Stern, D. M., Hori, O., and Ogawa, S. (2001) *J. Clin. Invest.* **108**, 1439–1450
 39. Asahi, H., Koshida, K., Hori, O., Ogawa, S., and Namiki, M. (2002) *BJU Int.* **90**, 462–466
 40. Sawai, N., Kita, M., Kodama, T., Tanahashi, T., Yamaoka, Y., Tagawa, Y., Iwakura, Y., and Imanishi, J. (1999) *Infect. Immun.* **67**, 279–285
 41. Bradford, M. M. (1976) *Anal. Biochem.* **72**, 248–254
 42. Nadanaka, S., Yoshida, H., Kano, F., Murata, M., and Mori, K. (2004) *Mol. Biol. Cell* **15**, 2537–2548
 43. Haze, K., Yoshida, H., Yanagi, H., Yura, T., and Mori, K. (1999) *Mol. Biol. Cell* **10**, 3787–3799
 44. Namba, T., Tanaka, K., Ito, Y., Ishihara, T., Hoshino, T., Gotoh, T., Endo, M., Sato, K., and Mizushima, T. (2009) *Am. J. Pathol.* **174**, 1786–1798
 45. Gotoh, T., Oyamomari, S., Mori, K., and Mori, M. (2002) *J. Biol. Chem.* **277**, 12343–12350
 46. Yoshida, H., Haze, K., Yanagi, H., Yura, T., and Mori, K. (1998) *J. Biol. Chem.* **273**, 33741–33749
 47. Namba, T., Homan, T., Nishimura, T., Mima, S., Hoshino, T., and Mizushima, T. (2009) *J. Biol. Chem.* **284**, 4158–4167
 48. Shirin, H., Sordillo, E. M., Oh, S. H., Yamamoto, H., Delohery, T., Weinstein, I. B., and Moss, S. F. (1999) *Cancer Res.* **59**, 2277–2281
 49. Ashktorab, H., Neapolitano, M., Bomma, C., Allen, C., Ahmed, A., Dubois, A., Naab, T., and Smoot, D. T. (2002) *Microbes Infect.* **4**, 713–722
 50. Hong, M., Li, M., Mao, C., and Lee, A. S. (2004) *J. Cell. Biochem.* **92**, 723–732
 51. Ye, J., Rawson, R. B., Komuro, R., Chen, X., Davé, U. P., Prywes, R., Brown, M. S., and Goldstein, J. L. (2000) *Mol. Cell* **6**, 1355–1364
 52. Huang, T., Wan, Y., Zhu, Y., Fang, X., Hiramatsu, N., Hayakawa, K., Paton, A. W., Paton, J. C., Kitamura, M., and Yao, J. (2009) *J. Cell. Biochem.* **107**, 973–983
 53. VanSlyke, J. K., and Musil, L. S. (2002) *J. Cell Biol.* **157**, 381–394
 54. Sakamoto, C., Sugano, K., Ota, S., Sakaki, N., Takahashi, S., Yoshida, Y., Tsukui, T., Osawa, H., Sakurai, Y., Yoshino, J., Mizokami, Y., Mine, T., Arakawa, T., Kuwayama, H., Saigenji, K., Yakabi, K., Chiba, T., Shimosegawa, T., Sheehan, J. E., Perez-Gutthann, S., Yamaguchi, T., Kaufman, D. W., Sato, T., Kubota, K., and Terano, A. (2006) *Eur. J. Clin. Pharmacol.* **62**, 765–772
 55. Molinari, M., Galli, C., Norais, N., Telford, J. L., Rappuoli, R., Luzio, J. P., and Montecucco, C. (1997) *J. Biol. Chem.* **272**, 25339–25344
 56. Li, Y., Wandinger-Ness, A., Goldenring, J. R., and Cover, T. L. (2004) *Mol. Biol. Cell* **15**, 1946–1959
 57. Tomisato, W., Takahashi, N., Komoto, C., Rokutan, K., Tsuchiya, T., and Mizushima, T. (2000) *Dig. Dis. Sci.* **45**, 1674–1679
 58. Amieva, M. R., Vogelmann, R., Covacci, A., Tompkins, L. S., Nelson, W. J., and Falkow, S. (2003) *Science* **300**, 1430–1434

