

## D. 考察

ウイルスライブラリーに存在する株の大部分を継代することにより、MDCK 細胞に馴化することを明らかにした。このことは、ウイルスライブラリーを用いてインフルエンザワクチンを作製することが可能であることを示すものである。今後は、これらの株について試作ワクチンを作製し、そのワクチン効果を検討する予定である。

今回我々は、WV ワクチン単独での経鼻接種による交叉防御効果が S V ワクチンとアジュバントとの併用経鼻接種の場合とほとんど同じで、ワクチン株と同じ亜型の変異株に対して十分な交叉防御効果を示すことを見出した。さらに WV ワクチンの経鼻接種でのワクチン株と同じ株に対する防御効果は、同ワクチンの皮下接種の場合よりはるかに強いものであった。現在、臨床で用いることができる粘膜アジュバントは存在しないことから、WV ワクチンが臨床に使用出来得る交叉防御効果を誘導する経鼻接種用ワクチンであることを強く示唆した。

これまで WV ワクチンの交叉防御効果に抗ウイルス中和効果が関与するとの可能性について直接明らかにした報告はほとんどない。鼻洗浄液および肺洗浄液の量は 1ml と非常に多く、各分泌液が 300 倍以上も希釈され、その結果、中和抗体価が検出されないことも考えられる。そこで我々は各洗浄液を濃縮することにより抗ウイルス中和抗体価の測定を行い、その検出に成功した。そして、WV ワクチンの経鼻接種による交叉反応性の中和抗体価の大小が、交叉防御効果のそれと相関することを明らかにした。

以上の結果を踏まえ、今後はウイルスライブラリーを用いて作製した WV ワクチンを経鼻接種することにより、交叉防御効果が見られるウイルス株の範囲を検討し、より広い範囲で交叉防御効果を示すワクチン用ウイルス株の選択を行う計画である。

## E. 結論

1. インフルエンザライブラリーの株を用いて培養細胞で高い増殖能を有するワクチン用種ウイルスを得ることを確認した

2. WV ワクチン単独での経鼻接種は、S V ワクチンとアジュバントとの併用経鼻接種と同様の十分な交叉防御効果を示すことを明らかにした。

## F. 研究発表

### 1. 論文発表

1. Kayamuro H, Yoshioka Y, Abe Y, Arita S, Katayama K, Nomura T, Yoshikawa T, Kubota-Koketsu R, Ikuta K, Okamoto S, Mori Y, Kunisawa J, Kiyono H, Itoh N, Nagano K, Kamada H, Tsunoda S, Tsutsumi Y. 2010. Systematic characterization of mucosal and systemic immune responses elicited by interleukins as mucosal vaccine adjuvant against influenza virus. *J. Virol.* 84:12703-12712.

2. Somboonthum P, Koshizuka T, Okamoto S, Matsuura M, Gomi Y, Takahashi M, Yamanishi K, Mori Y. 2010. Rapid and efficient introduction of a foreign gene into bacterial artificial chromosome-cloned varicella vaccine by Tn7-mediated site-specific transposition. *Virology*, 402:215-221.

### 2. 学会発表

1. 岡本 成史、松岡 須美子、山田 博司、五味 康行、奥野 良信、明石 満、森 康子、山西 弘一、インフルエンザ不活化全粒子ワクチン経鼻接種による交叉防御免疫応答の検討、第 14 回日本ワクチン学会学術集会、東京都千代田区、2010 年 12 月 11-12 日。

2. Okamoto S, Matsuoka S, Yamada H, Gomi Y, Okuno Y, Akagi T, Akashi M, Mori Y, Yamanishi K, Protective immune response by intranasal immunization with split-virion influenza A vaccine with adjuvant and whole-virion influenza A vaccine, Cell Symposia: Influenza - Translating basic insights, Washington DC, USA, December 2-4, 2010.

**H. 知的財産権の出願・登録状況**  
なし。

## 研究成果の刊行に関する一覧表

### 書籍

著者氏名	論文タイトル名	書籍全体の編集者名	書籍名	出版社名	出版地	出版年	ページ

### 雑誌

発表者氏名	論文タイトル名	発表誌名	巻号	ページ	出版年
Matsuura M, Takemoto M, <u>Yamanishi K</u> , Mori Y.	Human herpesvirus 6 major immediate early promoter has strong activity in T cells and is useful for heterologous gene expression.	Viol J.	8	9	2011
Tang H, Kawabata A, Yoshida M, Oyaizu H, Maeki T, <u>Yamanishi K</u> , Mori Y.	Human herpesvirus 6 encoded glycoprotein Q1 gene is essential for virus growth.	Virology	407(2)	360-7	2010
Koshizuka T, Ota M, <u>Yamanishi K</u> , Mori Y.	Characterization of varicella-zoster virus-encoded ORF0 gene--comparison of parental and vaccine strains.	Virology	405(2)	280-8	2010
Somboonthum P, Koshizuka T, Okamoto S, Matsuura M, Gomi Y, Takahashi M, <u>Yamanishi K</u> , Mori Y.	Rapid and efficient introduction of a foreign gene into bacterial artificial chromosome-cloned varicella vaccine by Tn7-mediated site-specific transposition.	Virology	402(1)	215-21	2010
Sadaoka T, Yanagi T, <u>Yamanishi K</u> , Mori Y.	Characterization of the varicella-zoster virus ORF50 gene, which encodes glycoprotein M.	J Virol.	84(7)	3488-502	2010
Isoda N, Tsuda Y, Sakoda Y, and <u>Kida H</u> .	Improvement of the H5N1 influenza virus vaccine strain to decrease the pathogenicity in chicken embryos.	Arch Virol	in press		2010
Feng F, Miura N, Isoda N, Sakoda Y, Okamatsu M, <u>Kida H</u> and Nishimura S.	Novel trivalent anti-influenza reagent.	Bioorg Med Chem Lett	20	3772-3776	2010

発表者氏名	論文タイトル名	発表誌名	巻号	ページ	出版年
Iwai A, Shiozaki T, Kawai T, Akira S, Kawaoka Y, Takada A, <u>Kida H</u> and Miyazaki T.	Influenza A virus polymerase inhibits type I interferon induction by binding to interferon {beta} promoter stimulator 1.	J Biol Chem	285		2010
Okamatsu M, Tanaka T, Yamamoto N, Sakoda Y, Sasaki T, Tsuda Y, Isoda N, Kokumai N, Takada A, Umemura T and <u>Kida H</u> .	Antigenic, genetic, and pathogenic characterization of H5N1 highly pathogenic avian influenza viruses isolated from dead whooper swans (Cygnus cygnus) found in northern Japan in 2008.	Virus Genes	41	351-357	2010
Sakoda Y, Sugar S, Batchluun D, Erdene-Ochir TO, Okamatsu M, Isoda N, Soda K, Takakuwa H, Tsuda Y, Yamamoto N, Kishida N, Matsuno K, Nakayama E, Kajihara M, Yokoyama A, Takada A, Sodnomdarjaa R and <u>Kida H</u> .	Characterization of H5N1 highly pathogenic avian influenza virus strains isolated from migratory waterfowl in Mongolia on the way back from the southern Asia to their northern territory.	Virology	406	88-94	2010
Tanaka T, Sunden Y, Sakoda Y, <u>Kida H</u> , Ochiai K and Umemura T.	Lipopolysaccharide treatment and inoculation of influenza A virus results in influenza virus-associated encephalopathy-like changes in neonatal mice.	J Neurovirol	16	125-132	2010
Miyake T, Soda K, Itoh Y, Sakoda Y, Ishigaki H, Nagata T, Ishida H, Nakayama M, Ozaki H, Tsuchiya H, Torii R, <u>Kida H</u> and Ogasawara K.	Amelioration of pneumonia with Streptococcus pneumoniae infection by inoculation with a vaccine against highly pathogenic avian influenza virus in a non-human primate mixed infection model.	J Med Primatol	39	58-70	2010
Itoh Y, Ozaki H, Ishigaki H, Sakoda Y, Nagata T, Soda K, Isoda N, Miyake T, Ishida H, Okamoto K, Nakayama M, Tsuchiya H, Torii R, <u>Kida H</u> and Ogasawara K.	Subcutaneous inoculation of a whole virus particle vaccine prepared from a non-pathogenic virus library induces protective immunity against H7N7 highly pathogenic avian influenza virus in cynomolgus macaques.	Vaccine	28	780-789	2010

発表者氏名	論文タイトル名	発表誌名	巻号	ページ	出版年
Igarashi M, Ito K, Yoshida R, Tomabechi D, <u>Kida H</u> and Takada A.	Predicting the antigenic structure of the pandemic (H1N1) 2009 influenza virus hemagglutinin.	PLoS One	5	e8553	2010
Ichinohe T, Ainai A, Ami Y, Nagata N, Iwata N, Kawaguchi A, Suzaki Y, Odagiri T, Tashiro M, Takahashi H, Strayer D, Carter W, Chiba J, Tamura S, Sata T, Kurata T, and <u>Hasegawa H.</u>	Intranasal administration of adjuvant-combined vaccine protects monkeys from challenge with the highly pathogenic influenza A H5N1 virus.	J Med Virol	82(10)	1754-61	2010
Kuroda M, Katano H, Nakajima N, Tobiume M, Ainai A, Sekizuka T, <u>Hasegawa H</u> , Tashiro M, Sasaki Y, Arakawa Y, Hata S, Watanabe M, Sata T.	Characterization of quasispecies of pandemic 2009 influenza A virus (A/H1N1/2009) by de novo sequencing using a next-generation DNA sequencer.	PLoS One.	5(4)	e10256	2010
Yamazaki T, Nagashima M, Ninomiya D, Arai Y, Teshima Y, Fujimoto A, Ainai A, <u>Hasegawa H</u> , Chiba J.	Passive Immune-Prophylaxis against Influenza Virus Infection by the Expression of Neutralizing Anti-Hemagglutinin Monoclonal Antibodies from Plasmids.	Jpn J Infect Dis.	64(1)	40-9	2011
Ainai A, Tashiro M, <u>Hasegawa H.</u>	Cross-protective immunity against influenza virus infections induced by intranasal vaccination together with a TLR3-mucosal adjuvant.	Hum Vaccin.	7 [Epub ahead of print]		2011
<u>Tetsuya Uchida</u>	Development of a cytotoxic T-lymphocyte-based, broadly protective influenza vaccine. Microbiol.	Immunol.	55	19-27	2011
Maiko Taneichi, Yuriko Tanaka, Terutaka Kakiuchi, <u>Tetsuya Uchida</u>	Liposome-coupled peptides induce long-lived memory CD8+ T cells without CD4+ T cells.	PLoS ONE	5(11)	e15091	2010
Yuriko Tanaka, Maiko Taneichi, Michiyuki Kasai, Terutaka Kakiuchi, <u>Tetsuya Uchida</u>	Liposome-coupled antigens are internalized by antigen-presenting cells via pinocytosis and cross-presented to CD8+ T cells.	PLoS ONE	5(12)	e15225	2010

発表者氏名	論文タイトル名	発表誌名	巻号	ページ	出版年
内田哲也	細胞性免疫誘導型リポソームワクチン	Bio Clinica	25	29-33	2010
内田哲也	今日の新型インフルエンザワクチンの問題点と将来展望	日本医師会雑誌	139	1491-1494	2010
Yoshida,T., Saito,A., Iwasaki,Y., Iijima,S., Kurosawa,T., Katakai,Y., Yasutomi,Y., Reimann,K.A., Hayakawa,T. and Akari,H.	Characterization of natural killer cells in tamarins: a technical basis for studies of innate immunity.	Frontiers Microbiol.	in press		2011
Xing, Li., Wang, J.C., Li, T-C., Yasutomi,Y., Lara,J., Khurdyakaov,Y., Schofield D., Emerson,S., Purcell,R., Takeda,N., Miyamura,T. and Holland,R.C.	Spatial configuration of hepatitis E virus antigenic domain.	J.Virol.	85	1117-1124	2011
Chono,H., Matsumoto,K., Tsuda,H., Saito,N., Lee,K., Kim,S., Shibata,H., Ageyama,N., Terao,K., Yasutomi,Y., Mineno J., Kim,S., Inoue,M. and Kato,I.	Acquisition of HIV-1 resistance in T lymphocytes using an ACA-specific E.coli mRNA interferase.	Human Gene Ther.	22	35-43	2011
Saito,A., Nomaguchi,M., Iijima,S., Lee,Y-J., Kono,K., Nakayama,E.E., Shioda,T., Yasutomi,Y., Adachi,A., Matano,T., Akari,H.	A novel monkey-tropic HIV-1 derivative encoding only minimal SIV sequences can replicate in cynomolgus monkeys.	Micorbes Infect.	13	58-64	2011
Naruse,T.K., Zhiyong,C., Yanagida,R., Yamashita,T., Saito,Y., Mori,K., Akari,H., Yasutomi,Y., Matano,T. and Kimura,A.	Diversity of MHC class I genes in Burmese-origine rhesus macaques.		62	601-611	2010
Okabayashi,S., Uchida,K., Nakayama,H., Ohno,C., Hanari,K., Goto,I. and Yasutomi,Y.	Periventricular Leucomalacia (PVL)-like lesions in two neonatal cynomolgus monkeys (macaca fascicularis).	J.Comp. Pathol.	Epub		2010
Yasuhiro Yasutomi.	Establishment of Specific Pathogen-Free Macaque Colonies in Tsukuba Primate Research Center of Japan for AIDS research.	Vaccine		B75-B77	2010

発表者氏名	論文タイトル名	発表誌名	巻号	ページ	出版年
Fujimoto,K., Takano,J., Narita,T., Hanari,K., Shimozawa,N., Sankai,T., Yoshida T., Terao,K., Kurata,T. and Yasutomi,Y.	Simian Retrovirus type D infection in a colony of cynomolgus monkeys.	Comp.Med.	60	51-53	2010
Cueno,M.E., Karamatsu,K., Yasutomi, Y., Laurena,A.C. and Okamoto.T.	Preferential expression and immunogenicity of HIV-1 Tat fusion protein expressed in tomato plant.	Transgenic Res.	19	889-895	2010
Kayamuro H, Yoshioka Y, Abe Y, Arita S, Katayama K, Nomura T, Yoshikawa T, Kubota-Koketsu R, Ikuta K, Okamoto S, Mori Y, Kunisawa J, Kiyono H, Itoh N, Nagano K, Kamada H, Tsunoda S, Tsutsumi Y.	Systematic characterization of mucosal and systemic immune responses elicited by interleukins as mucosal vaccine adjuvant against influenza virus.	J. Virol.	84	12703-12712	2010
Somboonthum P, Koshizuka T, Okamoto S, Matsuura M, Gomi Y, Takahashi M, Yamanishi K, Mori Y.	Rapid and efficient introduction of a foreign gene into bacterial artificial chromosome-cloned varicella vaccine by Tn7-mediated site-specific transposition.	Virology	402	215-221	2010

## 【参考資料】





RESEARCH

Open Access

# Human herpesvirus 6 major immediate early promoter has strong activity in T cells and is useful for heterologous gene expression

Masaaki Matsuura<sup>1,2</sup>, Masaya Takemoto<sup>1</sup>, Koichi Yamanishi<sup>1</sup>, Yasuko Mori<sup>1,3\*</sup>

## Abstract

**Background:** Human herpesvirus-6 (HHV-6) is a beta-herpesvirus. HHV-6 infects and replicates in T cells. The HHV-6-encoded major immediate early gene (MIE) is expressed at the immediate-early infection phase. Human cytomegalovirus major immediate early promoter (CMV MIEp) is commercially available for the expression of various heterologous genes. Here we identified the HHV-6 MIE promoter (MIEp) and compared its activity with that of CMV MIEp in various cell lines.

**Methods:** The HHV-6 MIEp and some HHV-6 MIEp variants were amplified by PCR from HHV-6B strain HST. These fragments and CMV MIEp were subcloned into the pGL-3 luciferase reporter plasmid and subjected to luciferase reporter assay. In addition, to investigate whether the HHV-6 MIEp could be used as the promoter for expression of foreign genes in a recombinant varicella-zoster virus, we inserted HHV-6 MIEp-DsRed expression cassette into the varicella-zoster virus genome.

**Results:** HHV-6 MIEp showed strong activity in T cells compared with CMV MIEp, and the presence of intron 1 of the MIE gene increased its activity. The NF- $\kappa$ B-binding site, which lies within the R3 repeat, was critical for this activity. Moreover, the HHV-6 MIEp drove heterologous gene expression in recombinant varicella-zoster virus-infected cells.

**Conclusions:** These data suggest that HHV-6 MIEp functions more strongly than CMV MIEp in various T-cell lines.

## Background

Human herpesvirus 6 (HHV-6) was first isolated in 1986 from the peripheral blood of patients with lymphoproliferative disorders and AIDS [1,2]. The virus was subsequently shown to be ubiquitous in healthy adults [3]. HHV-6 has been isolated from infants with exanthema subitum, a common childhood disease [4]. Later, HHV-6 isolates were classified into two variants, A and B (HHV-6A and HHV-6B), based on molecular and biological criteria [5-8]. HHV-6B causes exanthema subitum [4], while the pathogenesis of HHV-6A is still unknown. HHV-6 has the unique feature of being able to replicate and produce progeny in T cells [9,10]. The HHV-6 genome is a double-stranded DNA of approximately 160 kbp, consisting of a unique long region of 140 kbp

flanked by 10-kbp direct repeats, and there is 90% identity between the two variants [11].

HHV-6 belongs to the beta-herpesvirus subfamily, which includes human cytomegalovirus (HCMV) and human herpesvirus 7 (HHV-7) [12]. The beta-herpesviruses have extensive domains of similar genomic organization, with conserved herpesvirus gene blocks in the unique region of their genome [13]. HCMV's major immediate early (MIE) enhancer-containing promoter has been developed [14,15]; it is currently commercially available and is used to drive the expression of various genes. The MIE promoter controls the expression of two IE transcripts, designated IE1 (UL123) and IE2 (UL122) [16]. HHV-6 has positional homologs of UL123 and UL122; they are U89 and U86, which are designated IE1 and IE2, respectively [11,13,17,18]. The HHV-6 IE1 and IE2 transcripts are formed by alternative splicing [19,20]. Recently Takemoto et al. reported that the R3 region in the right end of HHV-6 is a strong enhancer

\* Correspondence: ymori@nibio.go.jp

<sup>1</sup>Laboratory of Virology, Division of Biomedical Research, National Institute of Biomedical Innovation, 7-6-8, Saito-Asagi, Ibaraki, Osaka 567-0085, Japan  
Full list of author information is available at the end of the article



of another HHV-6 immediate early gene, U95 [21]. R3 is positioned between U95 and U89; therefore, the region containing R3 is predicted to also contain promoter activity for the IE1 and IE2 genes. In other words, this location is predicted to be a positional homolog of the HCMV MIE promoter.


In this study, we identified the promoter region that regulates the HHV-6 MIE gene, and analyzed its activity. As expected, part of the R3 region was critical for the promoter activity. We also found that the first intron encoded by the IE1 gene enhanced HHV-6 MIE promoter (HHV-6 MIEp) activity, and that HHV-6 MIEp with the first intron had significantly stronger activity than the HCMV MIE promoter, especially in T-cell lines. The HHV-6 MIEp was able to express heterologous genes in a recombinant varicella-zoster virus, indicating that it could be useful for expressing various genes in a similar manner as the CMV MIE promoter.

## Results

### The HHV-6 major immediate-early promoter had stronger activity than the CMV promoter in T-cell lines


The 5' end of the mRNA encoded by the HHV-6 immediate early 1 (IE1) gene is located at base 139442 of the HHV-6 strain HST genome [11,22]. The 971-bp region upstream of the IE1 gene, including the R3 repeat, was suspected to include the HHV-6 major immediate-early promoter (HHV-6MIEp). The promoter region used in this study is illustrated in Figure 1A. First, to investigate the relative strength of the HHV-6 MIE promoter in various cell types, reporter gene assays were performed using the luciferase gene expression system. A plasmid containing the luciferase gene under the HHV-6MIEp was transfected into MRC-5, MeWo, U373, Molt-3, SupT1, and Jurkat cells. The pRL-TK plasmid, encoding Renilla luciferase under the transcriptional control of the herpes simplex virus thymidine kinase (HSV-TK) promoter, was co-transfected to normalize the transfection efficiency. The data show the fold-increase relative to the value of cells transfected with a blank plasmid, pGL3-basic (Promega). As shown in Figure 1B, the activity of the HHV-6 MIE promoter was weaker than that of the CMV promoter (CMV MIEp) in MRC-5, U373 and Mewo cells, while the activity was stronger than that of the CMV promoter in Molt-3, SupT1, and Jurkat cells.

The mRNAs encoded by the HHV-6 IE1 gene are produced by alternative splicing (Figure 1A). It is known that introns within some genes can elevate the protein expression level by either enhancing the promoter activity or stabilizing the mRNA [23]. In HCMV, the addition of intron A from the IE1 gene to the IE promoter/enhancer increases the promoter activity [24]. Therefore, we examined the role of the introns encoded by the




**Figure 1 Comparison of the activities of the HHV-6 MIE promoter and CMV promoter.** (A) The HHV-6 genome is a double-stranded DNA molecule of approximately 160 kbp, and is composed of a single long unique sequence (U) flanked by identical direct repeats (DR<sub>L</sub> and DR<sub>R</sub>). IE1 maps to open reading frames U90 and U89. The HHV-6 MIE promoter of 971 bp, located upstream of exon 1 of IE1, was amplified by PCR from the HHV-6B strain HST genome. Bases are numbered starting with the transcriptional start site for the IE1 gene. (B) The HHV-6 MIE promoter and CMV promoter were subcloned into the pGL3-basic plasmid, and the resultant plasmids were transfected into various cell lines (Jurkat, Molt-3, SupT1, MRC-5, MeWo, and U373 cells). At 24-hr post-transfection, the cells were harvested and subjected to the luciferase activity assay. The mean fold-activity relative to that of blank pGL3-basic plasmid-transfected cells and standard deviation for three independent experiments were plotted.

HHV-6 MIE genes in the HHV-6 MIE promoter activity. To examine this, several HHV-6 MIE promoter variants containing introns 1-4 were constructed (Figure 2A), and the activities were compared by performing the reporter assay in various cells. As shown in Figure 2B, in the presence of intron 1, the promoter activity was significantly upregulated in all the cells compared to the HHV-6 MIE promoter without intron 1. In contrast, the further addition of introns 1-2, 1-3, or 1-4 downregulated the promoter activity (Figure 2B). Therefore the HHV-6 MIE promoter containing intron 1 (HHV-6MIEp-in1), whose length is 1245-bp, was included in the remaining experiments.



**Figure 2** The activity of the HHV-6 MIE promoter including intron(s) of the IE gene. Schematic representation of the HHV-6 MIE promoter variants, including introns 1-4 of the IE1 gene, used for analysis of the introns. The numbers indicate the nucleotide position from the 5' end of mRNA encoded by the IE1 gene. The +1 indicates the 5' end. The name and size of each promoter used here are shown at the right. (B) Luciferase assays were performed after transfection in various cell lines (Molt-3, Jurkat, SupT-1, MRC-5, Mewo, and U373). The mean fold-activity of each HHV-6 MIE promoter variant relative to that of the blank pGL3-basic plasmid is shown by the horizontal bar. Standard deviation for three independent experiments is indicated. One asterisk indicates that the P value is < 0.05 and two asterisks indicate that the P value is < 0.01 in comparison with HHV-6MIEp-d2 (without intron 1), as determined by Student's unpaired two-tailed t-test.

Next, to determine the region that contributes to the promoter activity, various deletion mutants of both HHV6MIEp and HHV6MIEp-in1 were constructed (Figure 3A), and their activities were examined and compared by reporter assays in various cell lines. As shown in Figure 3B, the HHV-6MIEp-d3 promoter activity decreased compared to that of HHV-6MIEp-d2 (both with and without intron 1), showing that the region at nt positions from -381 to -552, which lies within R3, is important for the activity. In addition, the activities of HHV-6MIEp and HHV-6MIEp-in1 were significantly stronger than CMV MIEp activity in Jurkat, Molt-3, and SupT1 cells, suggesting that the HHV-6 MIE promoter has higher activity than the CMV promoter in certain cells, especially in T cells. This property of



**Figure 3** Comparison of promoter activity in deletion mutants of HHV-6MIEp and HHV-6MIEp-in1. (A) Schematic representation of 5'-deletion mutants of the HHV-6MIEp (black arrows) and of the HHV-6MIEp-in1, which is the HHV-6MIEp including intron 1 (white arrows). The name and size of each promoter are shown at the right (without intron 1) and left (with intron 1). Putative transcription factor-binding sites, predicted by TFSEARCH (<http://www.cbrc.jp/research/db/TFSEARCH.html>), are shown. (B) Luciferase assays were performed in various cell lines. The mean fold-activity relative to that of the blank pGL3-basic plasmid is indicated by the horizontal bars. The standard deviation for three independent experiments is indicated. The activities of CMV MIEp, deletion mutants of the HHV-6MIEp, and deletion mutants of the HHV-6MIEp-in1, are indicated by gray, black, and white bars, respectively. One asterisk indicates that the P value is < 0.05 and two asterisks indicate that the P value is < 0.01 in comparison with the CMV MIEp, as determined by Student's unpaired two-tailed t-test.


the HHV-6 MIE promoter might render it as a promising candidate for efficient protein expression in T cells. The region at nt positions -381 to -552, which lies within R3, is predicted to have an NF-κB-binding site and AP-1-binding site (Figure 3A). Takemoto et al. reported that the NF-κB-binding site in the R3 region plays an important role in U95 promoter activity [21]. We hypothesized that the NF-κB-binding site plays a major role in the HHV-6MIEp promoter activity as well. To investigate this, we constructed a promoter in which the NF-κB-binding site was deleted (HHV-6MIEpΔNF-

$\kappa$ Bin1) (Figure 4), and examined its activity in various cell lines. As shown in Figure 4, the NF- $\kappa$ B-binding site-deleted promoter HHV-6MIEp $\Delta$ NF- $\kappa$ Bin1 exhibited significantly decreased promoter activity in all cell lines, indicating that the NF- $\kappa$ B-binding site in the HHV-6MIEp plays an important role in its promoter activity.


**The HHV-6 MIE promoter could drive the expression of foreign gene in a recombinant varicella virus**

We recently constructed a recombinant varicella vaccine Oka strain (vOka) expressing the MuV (mumps virus) HN (hemagglutinin-neuraminidase) gene, as a possible candidate for a polyvalent vaccine for both varicella zoster virus (VZV) and MuV infections [25]. In that study, the CMV promoter was used to control the HN gene. Since the HHV-6 MIE promoter and CMV promoter showed similar activity in MRC-5 cells and MeWo cells, which are susceptible to VZV infection, we next examined whether the HHV-6 MIE promoter could control the expression of foreign genes in VZV.


To investigate this, we incorporated the HHV-6 MIE promoter, with the DsRed2 gene and BGH poly (A) signal sequence, into the vOka BAC genome by Tn7-mediated site-specific transposition (Figure 5). Since the full-length HHV-6 MIE promoter including intron 1 (HHV-6MIEpin1) had the strongest activity of all the promoter variants, we used it for this construct. The DsRed2 gene, which encodes a red fluorescent protein, was used as a reporter gene. The insertion of foreign gene cassette was confirmed by RFLP analysis using *Bam*HI and southern blot analysis. As shown in



**Figure 4 The NF- $\kappa$ B-binding site is critical for the promoter activity of 6MIEp.** (A) To investigate the importance of the NF- $\kappa$ B-binding site for the promoter activity of 6MIEp, a 5'-deletion mutant of the 6MIEp lacking the NF- $\kappa$ B-binding site (white letters in black box) was constructed. (B) The Luciferase assay was performed in various cell lines. The mean fold-activity relative to that of the blank pGL3-basic plasmid is indicated by the horizontal bars. The standard deviation for three independent experiments is indicated.



**Figure 5 Construction of the 6MIEpin1-DsRed-vOka genome.** The varicella vaccine Oka strain (vOka)-BAC genome (A) is about 125-kbp long and includes terminal repeats (TRL and TRS), a unique long (UL) domain, internal repeats (IRL and IRS), and a unique short domain (US). The LacZ $\alpha$ -mini-attTn7 sequence was inserted between ORF12 and ORF13 of the vOka-BAC genome by RecA-mediated recombination, generating vOka-BAC-Tn (B). The LacZ $\alpha$ -mini-attTn7 sequence in the vOka-BAC-Tn genome permitted site-specific insertion of the HHV-6MIEpin1-DsRed-BGH poly(A) signal sequence cassette (C) by Tn7-mediated transposition, resulting in the HHV-6MIEp-DsRed-vOka-BAC genome (D). Black arrowheads indicate the *Bam*HI sites. Horizontal bars indicate the region of the probe used for Southern blot analysis.



**Figure 6 Confirmation of the insertion of HHV-6MIEp-DsRed into the vOka-BAC genome by Southern blot.** (A) The HHV-6MIEpin1-DsRed-vOka-BAC DNA and the vOka-BAC DNA were digested with *Bam*HI, loaded onto a 0.5% agarose gel, and separated by electrophoresis. The DNA fragments were visualized with a UV transilluminator. Arrowheads indicate the band shift following transposition. Each DNA size is shown on the right side of the panel. (B) The blots were hybridized with ORF12, ORF13, DsRed, or HHV-6MIEp probes. Bands were detected by the Enhanced Chemiluminescence (ECL) Direct Nucleic Acid Labeling and Detection System. Lane M: size markers, lane 1: vOka-BAC DNA, lane 2: HHV-6MIEp-DsRed-vOka-BAC DNA.

Figure 6A, there was a shift in size from 7.8-kbp in the vOka-BAC DNA to 7.5-kbp in the HHV-6MIEpin1-DsRed-vOka-BAC DNA. Furthermore, in the Southern blot analysis, the probes for HHV-6MIEp and DsRed detected bands only in the HHV-6MIEpin1-DsRed-vOka-BAC genome (Figure 6B), indicating that the HHV-6MIEpin1-DsRed cassette had been inserted into the vOka genome.

To reconstitute infectious virus from the HHV-6MIEpin1-DsRed-vOka-BAC DNA, MRC-5 cells were transfected with the BAC DNA. Five days after the transfection, typical cytopathic effects (CPEs) were shown. Along with the CPEs, green fluorescence from green fluorescent protein (GFP), which gene was included in BAC sequence, and red fluorescence from DsRed2 were observed by fluorescence microscopy (Figure 7A); this indicated that the HHV-6MIEpin1-DsRed-vOka-BAC had been reconstituted as an infectious recombinant virus expressing DsRed under control of the HHV-6 MIE promoter.

The expression of the DsRed was confirmed by Western blotting analysis (Figure 7B). Recombinant


vOka-infected MRC-5 cell lysates were separated by SDS-PAGE and analyzed by Western blotting with an anti-DsRed mAb or anti-VZV gB Ab. The expression of gB, which is a late gene [26], was examined as a positive control of VZV infection. As shown in Figure 7B, the expression of gB was found in lysates from cells infected with either the control rvOka-BAC or HHV-6MIEpin1-DsRed-rvOka-BAC, while the anti-DsRed mAb specifically reacted with a 29-kDa band only in the HHV-6MIEpin1-DsRed-rvOka-BAC-infected cell lysates. These data indicated that the HHV-6 MIE promoter can be used to drive the expression of foreign genes in VZV-infected cells.

## Discussion

The HCMV major immediate early promoter (HCMV MIEp) has been established and used as a tool to drive gene expression by researchers worldwide. HHV-6 also belongs to the beta-herpesviruses, and has a positional homolog of the HCMV MIE gene. As described in the Introduction, HHV-6 replicates and produces progeny in T cells very well; we therefore speculated that the MIE promoter would have stronger promoter activity in T cells than in other cells. Here we identified the region of the HHV-6 major immediate early promoter (HHV-6 MIEp), described in Figure 1. The promoter activity of HHV-6 MIEp was stronger than that of HCMV MIEp in T-cell lines, but not in other adherent cell lines. This feature of HHV-6 MIEp activity is consistent with the fact that HHV-6 is T-cell tropic.

HHV-6 MIEp is predicted to have an NF- $\kappa$ B-binding site. The activity of a mutant HHV-6 MIEp, with the NF- $\kappa$ B-binding site deleted, was dramatically decreased, indicating that the NF- $\kappa$ B-binding site is critical for the promoter activity of HHV-6 MIEp. However, the HCMV MIEp activity was weak compared to that of HHV-6 MIEp in T-cell lines in our study, even though HCMV MIEp also has an NF- $\kappa$ B-binding site that plays a major role in its promoter activity [27,28]. Therefore, another binding site in addition to the NF- $\kappa$ B-binding site might contribute to the T-cell-specific promoter activity of HHV-6 MIEp, or another binding site in HCMV MIEp might have a repressive effect in T cells.

Although the AP-2 and PEA3 binding sites were not found in HHV-6 MIE promoter region by TSEARCH, R3 region has these binding sites [17,29]. However, in the study of U95 promoter, it has been reported that PEA3 binding sites in R3 region did not bind any proteins[21]. Therefore, PEA3 binding site might have no or low effect on the MIEp activity. The deletion promoter, HHV-6 MIEp-d1, lost two complete AP-2 binding sites and one AP-2 binding site with one nucleotide mutation, compared to full length promoter. Nevertheless, the activity of HHV-6 MIEp-d1 was similar to that



**Figure 7 The expression of heterologous gene under the HHV-6 promoter in recombinant VZV-infected cells.** (A) The HHV-6MIEpin1-DsRed-vOka-BAC DNA was transfected into MRC-5 cells. The infectious virus, reconstituted from the HHV-6MIEp-DsRed-vOka-BAC DNA, caused typical cytopathic effects along with green fluorescence and red fluorescence at 5 days post-transfection. (B) The HHV-6MIEpin1-DsRed-vOka-BAC-infected MRC-5 cells and vOka-BAC-infected MRC-5 cells were lysed in sample buffer, and subjected to Western blot analysis. Blots were reacted with an anti-DsRed mAb or anti-VZV gB Abs. The position and molecular mass in kDa of marker proteins are indicated at the left. Lane 1: vOka-BAC-infected MRC-5 cells, lane 2: HHV-6MIEpin1-DsRed-vOka-BAC-infected MRC-5 cells.

of HHV-6 MIEp. Therefore, the AP-2 binding sites might have low effect on the MIEp activity.

Adding the first intron (intron 1) of IE1 to HHV-6 MIEp enhanced the promoter activity significantly. When intron 1 was added, the activity of HHV-6 MIEp became markedly greater than that of HCMV in T cells. In adherent cell lines such as MRC-5 and MeWo cells, the activity of HHV-6 MIEp with intron 1 became similar to that of HCMV MIEp. Intron 1 of the IE1 region is predicted to have two CCAAT enhancer binding protein (C/EBP) binding sites and an OCT-1-binding site (Figure 3). The transcriptional regulators that bind to these sites might enhance the promoter activity of HHV-6 MIEp. Interestingly, the promoter construct that contained introns 1 and 2 was less active than the promoter containing only intron 1. Further investigation is needed to elucidate the mechanisms involving the intron regions.

We recently developed a recombinant VZV vaccine strain containing the mumps virus HN gene. In this study, we examined whether the HHV-6 MIEp containing intron 1 functioned as a heterologous expression promoter in the VZV vaccine strain. Indeed, in the recombinant VZV, HHV-6 MIEp functioned to drive the expression of the DsRed gene, which is a heterologous gene. These findings indicate that, like the commercially available HCMVp, HHV-6 MIEp is useful for expressing heterologous genes in a VZV vaccine strain.

## Conclusions

Our results show that HHV-6 MIE promoter functions more strongly than CMV MIEp in various T-cell lines. Moreover, the first intron of HHV-6 IE1 gene enhances the promoter activity of HHV-6 MIEp. In addition, the HHV-6 MIEp could drive heterologous gene expression in recombinant varicella-zoster virus-infected cells. These results suggest that HHV-6 MIEp can be used for driving gene expression.

## Methods

### Cells

MRC-5 cells, human lung fibroblasts, were cultured in modified minimum essential medium (MEM) supplemented with 10% fetal bovine serum (FBS). MeWo cells, a human melanoma cell line, and U373 cells, a human astrocytoma cell line, were cultured in Dulbecco's modified Eagle's medium supplemented with 8% FBS. Molt-3 cells, SupT1 cells, and Jurkat cells, which are lymphoblastic T-cell lines, were cultured in RPMI1640 medium supplemented with 8% FBS.

### Plasmids for the luciferase reporter assay

The HHV-6 major immediate-early promoter (HHV-6MIEp) sequence and its deletion mutants were

amplified by PCR from the HHV-6B strain HST [30]. The primer sequences are shown in Table 1. The 971-bp fragment located from -983 to -13 bp upstream of exon 1 of IE1, which was amplified using the primer pair 6MIEpF and 6MIEpR, was defined as 6MIEp. The 5' primers named 6MIEpF-732, 6MIEpF-552, 6MIEpF-531, 6MIEpF-381, 6MIEpF-214, 6MIEpF-165, and 6MIEpF-102 were used to generate a series of 5'-deletion mutants. The 3' primers named 6MIEpex2R, 6MIEpex3R, 6MIEpex4R, and 6MIEpex5R were used to amplify HHV-6MIEp including introns 1 to 4, respectively. These amplified fragments were digested and inserted into the pGL3-basic vector (Promega) at the *HindIII* and *XhoI* or *KpnI* site.

The CMV MIE promoter sequence was excised with *NruI* and *BamHI* from pcDNA3.1(+) (Invitrogen), and inserted into pGL3-basic (Promega) at the *SmaI* and *BglII* sites.

The pRL-TK plasmid (Promega), which contains the Renilla luciferase reporter gene under the HSV TK promoter, was used to normalize the transfection efficiency.

### Luciferase reporter assay

Adherent cells (MRC-5, MeWo, and U373) were plated on 24-well plates at a density of  $1 \times 10^5$  cells per well on the day before transfection, and were transfected with 1  $\mu$ g of reporter plasmid and 0.25  $\mu$ g of pRL-TK plasmid (Promega), using Lipofectamine 2000 (Invitrogen) according to the manufacturer's instructions. Samples containing  $4 \times 10^5$  suspended cells (Molt-3, Jurkat, or SupT1) were transfected with 1  $\mu$ g of reporter plasmid and 0.25  $\mu$ g of pRL-TK using Lipofectamine2000.

Firefly and Renilla luciferase activities were measured with the Dual-Luciferase Reporter Assay System (Promega) according to the manufacturer's protocol, using a luminometer (Berthold, TriStar LB941). Cells were lysed in 1  $\times$  lysis buffer (50  $\mu$ L/well) for 15 min at room temperature, and each cell lysate was added to a luminometer tube containing 100  $\mu$ L of assay reagent. The mixture was blended quickly by flicking, and placed in the luminometer for a 1-sec measurement. The transfection efficiency was normalized to the Renilla luciferase activity. The data (mean + SD) were collected from three independent transfections.

### Generation of a recombinant vOka-BAC genome containing HHV-6 MIE promoter

To generate the HHV-6MIEpin1-pFastBac plasmid, the gentamicin-resistance gene and the polyhedrin (PH)-promoter region of the pFastBac1 plasmid (Invitrogen) were replaced with 6MIEp including the intron 1 (HHV-6MIEpin1) sequence.

The DsRed fragment was amplified by PCR using the primer pair DsRed2-HindF and DsRed2-HindR, and

**Table 1 Primers**

Primer	Sequence*
6MIEpF	5'-tct <u>ctc gag</u> agt taa aga tca gcg ggt ac-3'
6MIEpF-732	5'-agt <u>cgg tac cgg</u> cga atg aga act cta aaa gct c-3'
6MIEpF-552	5'-agt <u>cgg tac</u> cta ctg tgg ttg ggg tct ttc cta c-3'
6MIEpF-531	5'-acc <u>ggg acc</u> tac cca ggc taa cga gaa cc-3'
6MIEpF-381	5'-agt <u>cgg tac cac</u> att cct gtt tca tga tgt gta gc-3'
6MIEpF-214	5'-agt <u>cgg tac ctc</u> ctg ttt ttg agt aag ata tga c-3'
6MIEpF-165	5'-agt <u>cgg tac cag</u> cta att tcc att cca tat ttg tc-3'
6MIEpF-102	5'-agt <u>cgg tac</u> cta cag cga ttg gct cct tca tcc tc-3'
6MIEpR	5'-agt <u>cct cga</u> gca ctg aac tgg ctg taa ctt ctg c-3'
6MIEpex2R	5'-tct <u>aag ctt</u> cag caa tcc aat aat tga tg-3'
6MIEpex3R	5'-cat <u>aag ctt</u> gca tac gtt cct cat tgg at-3'
6MIEpex4R	5'-cat <u>aag ctt</u> cca aag ttt tga att ctt ca-3'
6MIEpex5R	5'-cat <u>aag ctt</u> ttt gga tgc aag tgc caa cg-3'
DsRed2-HindF	5'-acc <u>aag ctt</u> tac cgg tcg cca cca tgg cct-3'
DsRed2-HindR	5'-acc <u>aag ctt</u> tta tct aga tcc ggt gga tcc-3'
ORF12TnFw	5'-tat <u>ctc gag</u> agg tac cgg tga ctt cag ag-3'
ORF12TnRv	5'-cga <u>gga tcc</u> aat caa cca atc aga cct-3'
ORF13TnFw	5'-gag <u>gat cgg</u> tac cca caa tat caa gtg gt-3'
ORF13TnRv	5'-gac <u>tcg agc</u> cta ttc gtg tca tct aga tgg-3'

\*:underlines indicate restriction enzyme sites.

*Hind*III sites were introduced at both the 5' and 3' ends. The pDsRed2-C1 plasmid (Clontech), in which the *Hind*III site had been eliminated, was used as the PCR template. Following amplification, the PCR products were inserted into the HHV-6MIEpin1-pFastBac plasmid at the *Hind*III site, generating the HHV-6MIEpin1-DsRed-pFastBac plasmid (Figure 5C). The BGH poly (A) signal sequence was derived from pFastBac plasmid.

The vOka-BAC was obtained using pHA-2 cloning vector (a kind gift from Dr. Ulrich Koszinowski[31]), as described previously[32]. The LacZ $\alpha$ -mini-attTn7 cassette was inserted into vOka-BAC (Figure 5A) to produce vOka-BAC-Tn (Figure 5B) using RecA-mediated recombination, essentially as described previously [32]. In brief, *E. coli* DH10B electrocompetent cells harboring circular vOka-BAC DNA were co-transformed with 1  $\mu$ g of the targeting vector, pKO5M-Tn (pKO5M is a kind gift from Dr. Kawaguchi[33]), which contain the LacZ $\alpha$ -mini-attTn7 region[33,34], and 3  $\mu$ g of pDF25 (Tet)- (a kind gift from Dr. J. Heath [35]) by electroporation, using a Gene Pulser II (Bio-Rad, Hercules, CA). The surviving co-integrant colonies, selected by their resistance to chloramphenicol and zeocin, and by a Lac + phenotype on an LB plate containing X-Gal and IPTG, were made electrocompetent and transformed with 1  $\mu$ g of pDF25(Tet). The *E. coli* DH10B

colonies containing the correct survival recombination were then selected by the following criteria: resistance to chloramphenicol, sensitivity to zeocin, and a Lac + phenotype on LB containing X-Gal and IPTG. The insertion of the LacZ $\alpha$ -mini-attTn7 sequence into the BAC genome was confirmed by PCR and Southern blotting (Data not shown).

The HHV-6MIEpin1-DsRed cassette was inserted into the vOka-BAC-Tn genome using Tn7-mediated site-specific transposition, essentially as described previously [34]. In brief, *E. coli* DH10B harboring the vOka-BAC-Tn genome was transformed with HHV-6MIEpin1-DsRed-pFastBac and pMON7124 (Invitrogen), a helper plasmid for transposition. The pMON7124 plasmid DNA was isolated from DH10Bac cells (Invitrogen). The transformed *E. coli* was cultured on LB containing X-gal and IPTG for blue/white selection. The white colonies were analyzed by PCR to verify the insertion of the DsRed expression cassette (data not shown). This completed the construction of the HHV-6MIEpin1-DsRed-vOka-BAC genome (Figure 5D).

#### Southern blot analysis

The HHV-6MIEpin1-DsRed-vOka-BAC DNA was extracted using a NucleoBond BAC 100 kit (Macherey-Nagel) following the manufacturer's instructions.

The BAC DNA was then digested with *Bam*HI, loaded onto a 0.5% agarose gel, and separated by electrophoresis at 20 V for 72 hrs. The DNA fragments were visualized with a UV transilluminator and then transferred onto a nylon membrane (Hybond-N+) (GE Healthcare Bio-sciences). The blots were hybridized with ORF12, ORF13, DsRed, or HHV-6MIEp probes labeled with horseradish peroxidase. These probes were amplified by PCR using the following primer pairs: ORF12TnFw/ORF12TnRv, ORF13TnFw/ORF13TnRv, DsRed-HindF/DsRed-HindR, and 6MIEpF-552/6MIEpex2R, respectively (the primer sequences are shown in Table 1). Bands were detected by the Enhanced Chemiluminescence (ECL) Direct Nucleic Acid Labeling and Detection System (GE Healthcare Bio-sciences) following the manufacturer's instructions.

#### Reconstitution of infectious virus from vOka-BAC DNA

Reconstitution of the recombinant virus, named HHV-6MIEpin1-DsRed-rvOka, was performed as described previously [32,36]. Briefly, MRC-5 cells were transfected with 1 µg of HHV-6MIEpin1-DsRed-vOka-BAC DNA by electroporation, using a Nucleofection unit (Amaxa Biosystems). The transfected cells were then cultured in MEM supplemented with 3% FBS for 3-5 days, and were observed under a microscope until a typical cytopathic effect with green and red fluorescence appeared.

#### Western blot analysis

The HHV-6MIEp-DsRed-vOka-BAC-infected MRC-5 cells were lysed in sample buffer [32 mM Tris-HCl (pH 6.8), 1.5% SDS, 5% glycerol, 2.5% 2-mercaptoethanol], separated by SDS-polyacrylamide gel electrophoresis (PAGE), and electrotransferred onto PVDF membranes (Bio-Rad Laboratories). A monoclonal antibody (mAb) against DsRed (Clontech) was purchased, and an anti-VZV gB monospecific antibody (Ab) was produced in our laboratory [26]. Blots were blocked with blocking buffer (PBS, 5% skim milk, 0.1% Tween-20) and reacted with the anti-DsRed mAb or anti-gB Ab in blocking buffer. The protein bands were developed with horseradish peroxidase-conjugated secondary antibodies (GE Healthcare) and ECL detection reagents (GE Healthcare Bio-Sciences), following the manufacturer's instructions.

#### Acknowledgements

We thank Dr. Ulrich Koszinowski (Max von Pettenkofer Institut für Virologie, Ludwig-Maximilians-Universität München, Germany) for providing the pHA-2 plasmid, Dr. John Heath (School of Biosciences, University of Birmingham, Birmingham, UK) for providing the pDF25(Tet) plasmid, Dr. Yasushi Kawaguchi (The Institute of Medical Science, The University of Tokyo, Japan) for providing the pKOSM plasmid. This study was supported in part by a grant in aid of Cluster, Ministry of Education, Culture, Sports, Science and Technology of Japan.

#### Author details

<sup>1</sup>Laboratory of Virology, Division of Biomedical Research, National Institute of Biomedical Innovation, 7-6-8, Saito-Asagi, Ibaraki, Osaka 567-0085, Japan. <sup>2</sup>Kanonji Institute, the Research Foundation for Microbial Diseases of Osaka University, 2-9-41, ahata-cho, Kanonji, Kagawa, 768-0061, Japan. <sup>3</sup>Division of Clinical Virology, Kobe University Graduate School of Medicine, 7-5-1, Kusunoki-cho, Chuo-ku, Kobe, 650-0017, Japan.

#### Authors' contributions

MM performed and analyzed the experiments, and drafted the manuscript. TM participated in the design of the study partly and performed the experiments partly. KY analyzed the study. YM participated in its design and coordination, analyzed the study, and drafted the manuscript. All authors read and approved the final manuscript.

#### Competing interests

The authors declare that they have no competing interests.

Received: 18 August 2010 Accepted: 11 January 2011

Published: 11 January 2011

#### References

1. Josephs SF, Salahuddin SZ, Ablashi DV, Schachter F, Wong-Staal F, Gallo RC: **Genomic analysis of the human B-lymphotropic virus (HBLV).** *Science* 1986, **234**:601-603.
2. Salahuddin SZ, Ablashi DV, Markham PD, Josephs SF, Sturzenegger S, Kaplan M, Halligan G, Biberfeld P, Wong-Staal F, Kramarsky B, et al: **Isolation of a new virus, HBLV, in patients with lymphoproliferative disorders.** *Science* 1986, **234**:596-601.
3. Linde A, Dahl H, Wahren B, Fridell E, Salahuddin Z, Biberfeld P: **IgG antibodies to human herpesvirus-6 in children and adults and in primary Epstein-Barr virus infections and cytomegalovirus infections [corrected].** *J Virol Methods* 1988, **21**:117-123.
4. Yamanishi K, Okuno T, Shiraki K, Takahashi M, Kondo T, Asano Y, Kurata T: **Identification of human herpesvirus-6 as a causal agent for exanthem subitum.** *Lancet* 1988, **1**:1065-1067.
5. Ablashi D, Agut H, Berneman Z, Campadelli-Fiume G, Carrigan D, Ceccerini-Nelli L, Chandran B, Chou S, Collandre H, Cone R, Dambaugh T, Dewhurst S, DiLuca D, Foa-Tomasi L, Fleckenstein B, Frenkel N, Gallo R, Gompels U, Hall C, Jones M, Lawrence G, Martin M, Montagnier L, Neipel F, Nicholas J, Pellett P, Razaque A, Torrelli G, Thomson B, Salahuddin S, Wyatt L, Yamanishi K: **Human herpesvirus-6 strain groups: a nomenclature.** *Arch Virol* 1993, **129**:363-366.
6. Ablashi DV, Balachandran N, Josephs SF, Hung CL, Krueger GR, Kramarsky B, Salahuddin SZ, Gallo RC: **Genomic polymorphism, growth properties, and immunologic variations in human herpesvirus-6 isolates.** *Virology* 1991, **184**:545-552.
7. Biberfeld P, Kramarsky B, Salahuddin SZ, Gallo RC: **Ultrastructural characterization of a new human B lymphotropic DNA virus (human herpesvirus 6) isolated from patients with lymphoproliferative disease.** *J Natl Cancer Inst* 1987, **79**:933-941.
8. Schirmer EC, Wyatt LS, Yamanishi K, Rodriguez WJ, Frenkel N: **Differentiation between two distinct classes of viruses now classified as human herpesvirus 6.** *Proc Natl Acad Sci USA* 1991, **88**:5922-5926.
9. Lusso P, Markham PD, Tschachler E, di Marzo Veronese F, Salahuddin SZ, Ablashi DV, Pahwa S, Krohn K, Gallo RC: **In vitro cellular tropism of human B-lymphotropic virus (human herpesvirus-6).** *J Exp Med* 1988, **167**:1659-1670.
10. Takahashi K, Sonoda S, Higashi K, Kondo T, Takahashi H, Takahashi M, Yamanishi K: **Predominant CD4 T-lymphocyte tropism of human herpesvirus 6-related virus.** *J Virol* 1989, **63**:3161-3163.
11. Isegawa Y, Mukai T, Nakano K, Kagawa M, Chen J, Mori Y, Sunagawa T, Kawanishi K, Sashihara J, Hata A, et al: **Comparison of the complete DNA sequences of human herpesvirus 6 variants A and B.** *J Virol* 1999, **73**:8053-8063.
12. Braun DK, Dominguez G, Pellett PE: **Human herpesvirus 6.** *Clin Microbiol Rev* 1997, **10**:521-567.
13. Gompels JA, Nicholas J, Lawrence G, Jones M, Thomson BJ, Martin ME, Efsthathiou S, Craxton M, Macaulay HA: **The DNA sequence of human herpesvirus-6: structure, coding content, and genome evolution.** *Virology* 1995, **209**:29-51.



14. Boshart M, Weber F, Jahn G, Dorsch-Hasler K, Fleckenstein B, Schaffner W: **A very strong enhancer is located upstream of an immediate early gene of human cytomegalovirus.** *Cell* 1985, **41**:521-530.
15. Thomsen DR, Stenberg RM, Goins WF, Stinski MF: **Promoter-regulatory region of the major immediate early gene of human cytomegalovirus.** *Proc Natl Acad Sci USA* 1984, **81**:659-663.
16. Malone CL, Vesole DH, Stinski MF: **Transactivation of a human cytomegalovirus early promoter by gene products from the immediate-early gene IE2 and augmentation by IE1: mutational analysis of the viral proteins.** *J Virol* 1990, **64**:1498-1506.
17. Dominguez G, Dambaugh TR, Stamey FR, Dewhurst S, Inoue N, Pellett PE: **Human herpesvirus 6B genome sequence: coding content and comparison with human herpesvirus 6A.** *J Virol* 1999, **73**:8040-8052.
18. Nicholas J: **Nucleotide sequence analysis of a 21-kbp region of the genome of human herpesvirus-6 containing homologues of human cytomegalovirus major immediate-early and replication genes.** *Virology* 1994, **204**:738-750.
19. Papanikolaou E, Kouvatzis V, Dimitriadis G, Inoue N, Arsenakis M: **Identification and characterization of the gene products of open reading frame U86/87 of human herpesvirus 6.** *Virus Res* 2002, **89**:89-101.
20. Schiewe U, Neipel F, Schreiner D, Fleckenstein B: **Structure and transcription of an immediate-early region in the human herpesvirus 6 genome.** *J Virol* 1994, **68**:2978-2985.
21. Takemoto M, Shimamoto T, Isegawa Y, Yamanishi K: **The R3 region, one of three major repetitive regions of human herpesvirus 6, is a strong enhancer of immediate-early gene U95.** *J Virol* 2001, **75**:10149-10160.
22. Kondo K, Shimada K, Sashihara J, Tanaka-Taya K, Yamanishi K: **Identification of human herpesvirus 6 latency-associated transcripts.** *J Virol* 2002, **76**:4145-4151.
23. Le Hir H, Nott A, Moore MJ: **How introns influence and enhance eukaryotic gene expression.** *Trends Biochem Sci* 2003, **28**:215-220.
24. Chapman BS, Thayer RM, Vincent KA, Haigwood NL: **Effect of intron A from human cytomegalovirus (Towne) immediate-early gene on heterologous expression in mammalian cells.** *Nucleic Acids Res* 1991, **19**:3979-3986.
25. Somboonthum P, Yoshii H, Okamoto S, Koike M, Gomi Y, Uchiyama Y, Takahashi M, Yamanishi K, Mori Y: **Generation of a recombinant Oka varicella vaccine expressing mumps virus hemagglutinin-neuraminidase protein as a polyvalent live vaccine.** *Vaccine* 2007, **25**:8741-8755.
26. Sadaoka T, Yoshii H, Imazawa T, Yamanishi K, Mori Y: **Deletion in open reading frame 49 of varicella-zoster virus reduces virus growth in human malignant melanoma cells but not in human embryonic fibroblasts.** *J Virol* 2007, **81**:12654-12665.
27. Cherrington JM, Mocarski ES: **Human cytomegalovirus ie1 transactivates the alpha promoter-enhancer via an 18-base-pair repeat element.** *J Virol* 1989, **63**:1435-1440.
28. Sambucetti LC, Cherrington JM, Wilkinson GW, Mocarski ES: **NF-kappa B activation of the cytomegalovirus enhancer is mediated by a viral transactivator and by T cell stimulation.** *EMBO J* 1989, **8**:4251-4258.
29. Martin ME, Nicholas J, Thomson BJ, Newman C, Honess RW: **Identification of a transactivating function mapping to the putative immediate-early locus of human herpesvirus 6.** *J Virol* 1991, **65**:5381-5390.
30. Takemoto M, Koike M, Mori Y, Yonemoto S, Sasamoto Y, Kondo K, Uchiyama Y, Yamanishi K: **Human herpesvirus 6 open reading frame U14 protein and cellular p53 interact with each other and are contained in the virion.** *J Virol* 2005, **79**:13037-13046.
31. Adler H, Messerle M, Wagner M, Koszinowski UH: **Cloning and mutagenesis of the murine gammaherpesvirus 68 genome as an infectious bacterial artificial chromosome.** *J Virol* 2000, **74**:6964-6974.
32. Yoshii H, Somboonthum P, Takahashi M, Yamanishi K, Mori Y: **Cloning of full length genome of varicella-zoster virus vaccine strain into a bacterial artificial chromosome and reconstitution of infectious virus.** *Vaccine* 2007, **25**:5006-5012.
33. Tanaka M, Kagawa H, Yamanashi Y, Sata T, Kawaguchi Y: **Construction of an excisable bacterial artificial chromosome containing a full-length infectious clone of herpes simplex virus type 1: viruses reconstituted from the clone exhibit wild-type properties in vitro and in vivo.** *J Virol* 2003, **77**:1382-1391.
34. Somboonthum P, Koshizuka T, Okamoto S, Matsuura M, Gomi Y, Takahashi M, Yamanishi K, Mori Y: **Rapid and efficient introduction of a foreign gene into bacterial artificial chromosome-cloned varicella vaccine by Tn7-mediated site-specific transposition.** *Virology* 2010, **402**:215-221.
35. Lalioti M, Heath J: **A new method for generating point mutations in bacterial artificial chromosomes by homologous recombination in Escherichia coli.** *Nucleic Acids Res* 2001, **29**:E14.
36. Nagaïke K, Mori Y, Gomi Y, Yoshii H, Takahashi M, Wagner M, Koszinowski U, Yamanishi K: **Cloning of the varicella-zoster virus genome as an infectious bacterial artificial chromosome in Escherichia coli.** *Vaccine* 2004, **22**:4069-4074.

doi:10.1186/1743-422X-8-9


Cite this article as: Matsuura et al.: Human herpesvirus 6 major immediate early promoter has strong activity in T cells and is useful for heterologous gene expression. *Virology Journal* 2011 **8**:9.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
www.biomedcentral.com/submit





## Human herpesvirus 6 encoded glycoprotein Q1 gene is essential for virus growth

Huamin Tang<sup>a</sup>, Akiko Kawabata<sup>a,b</sup>, Mayumi Yoshida<sup>b</sup>, Hiroko Oyaizu<sup>a,b</sup>, Takahiro Maeki<sup>a,b</sup>, Koichi Yamanishi<sup>c</sup>, Yasuko Mori<sup>a,b,\*</sup>

<sup>a</sup> Laboratory of Virology and Vaccinology, Division of Biomedical Research, National Institute of Biomedical Innovation, 7-6-8, Saito-Asagi, Ibaraki, Osaka 567-0085, Japan

<sup>b</sup> Division of Clinical Virology, Kobe University Graduate School of Medicine, Kobe, Japan

<sup>c</sup> National Institute of Biomedical Innovation, 7-6-8, Saito-Asagi, Ibaraki, Osaka 567-0085, Japan

### ARTICLE INFO

#### Article history:

Received 28 June 2010

Returned to author for revision 23 July 2010

Accepted 18 August 2010

Available online 21 September 2010

#### Keywords:

HHV-6

gQ1

BAC

Knockout gene

Ligand

### ABSTRACT

Human herpesvirus 6 (HHV-6) glycoprotein Q1 (gQ1), a unique gene in HHV-6, forms a complex with glycoproteinH (gH) and gL, which is the viral ligand for its cellular receptor, CD46. However, whether gQ1 is essential for virus growth is unknown, because a system is lacking for making gene knockouts for HHV-6. Recently, bacterial artificial chromosome (BAC) and *E. coli* mutagenesis techniques have been applied to herpesvirus investigation. Here we successfully inserted the HHV-6A genome into a BAC, and obtained reconstituted infectious virus from the HHV-6A-containing BAC DNA. Using this system, we generated a gQ1 mutant virus genome, which failed to yield reconstituted infectious virus, whereas its revertant virus could be produced, indicating that the HHV-6 gQ1 gene is essential for virus growth. Therefore, we successfully applied BAC and *E. coli* mutagenesis techniques to the study of HHV-6, and discovered that HHV-6 gQ1 is an essential gene for virus growth.

© 2010 Elsevier Inc. All rights reserved.

### Introduction

Human herpesvirus 6 (HHV-6) is a T-cell tropic virus, belonging to the betaherpesvirus subfamily and related to human herpesvirus 7 (HHV-7) and human cytomegalovirus (HCMV). It was first isolated from the peripheral blood lymphocytes of patients with lymphoproliferative disorders and AIDS (Salahuddin et al., 1986). The clinical isolates of HHV-6 can be categorized into two variants, HHV-6A and HHV-6B (Ablashi et al., 1991; Aubin et al., 1991; Campadelli-Fiume et al., 1993; Wyatt et al., 1990). Recently, HHV-6 was proposed to be an environmental factor contributing to diseases like multiple sclerosis, epilepsy, and others. However, the mechanisms underlying HHV-6's involvement in these diseases remain to be elucidated (Pietilainen et al., 2009; Theodore et al., 2008), in part because of a lack of biological tools for HHV-6.

Glycoproteins or their complexes on the surface of enveloped viruses play crucial roles in the viruses' infectivity. In HIV, a trimeric complex of gp120 forms a unique conformational structure that binds to the cellular HIV-1 receptor, CD4 (Amess et al., 1998). In several herpesviruses, unique glycoproteins have been reported that function by themselves or with other glycoproteins to promote virus entry. The binding of glycoprotein D (gD) of herpes simplex virus 1 (HSV-1), to its cellular receptor is a critical step for virus entry (Spear et al., 2000).

In Epstein–Barr virus (EBV), gp42 forms a complex with gH and gL. This complex binds to human leukocyte antigen (HLA) class II, and is required for the virus's entry into B cells (Li et al., 1997). The glycoprotein complex gH/gL/UL128-132 in HCMV is required for the virus' entry into epithelial and endothelial cells (Ryckman et al., 2008a), although its cellular partner remains to be identified.

The gQ proteins are unique to HHV-6 and HHV-7. Mori et al. (2003) first identified the gQ1 protein in HHV-6 and found that it forms a complex with gH and gL. They found that this complex binds to CD46, the cellular receptor for HHV-6 (Mori et al., 2004). Later, another gQ protein, gQ2 was identified, and found to be part of the gH/gL/gQ1 complex (Akkapaiboon et al., 2004; Mori, 2009). We speculated that gQ1 is essential for HHV-6 propagation, based on the functions of gD in HSV-1, gp42 in EBV, and UL128-132 in HCMV, but there were no direct experimental data to support this hypothesis. To test this hypothesis, new biologic research tools for studying HHV-6 were needed.

In recent years, a number of human herpesvirus genomes have been successfully cloned into F plasmids as BACs, including HSV-1 and HSV-2 (Meseda et al., 2004; Stavropoulos and Strathdee, 1998), varicella-zoster virus (VZV) (Nagaïke et al., 2004), EBV (Kanda et al., 2004), HCMV (Borst et al., 1999), and Kaposi's sarcoma associated herpesvirus (KSHV) (Zhou et al., 2002). This technique allows the stable maintenance of a viral genome as a BAC in *E. coli*, and the mutagenesis of viral genome in *E. coli* using the bacterial recombination machinery (Wagner et al., 2002). Reconstituted infectious viruses are generated by the transfection of the BAC plasmids into mammalian cells. The phenotypes of mutant viruses can help

\* Corresponding author. Division of Clinical Virology, Kobe University Graduate School of Medicine, 7-5-1, Kusunoki-cho, Chuo-ku, Kobe, 650-0017, Japan. Fax: +81 78 382 6879.

E-mail address: [ymori@med.kobe-u.ac.jp](mailto:ymori@med.kobe-u.ac.jp) (Y. Mori).

characterize the function(s) of their disrupted genes, and this information might lead to new treatments for controlling virus infections. In HHV-6, Borenstein and Frenkel successfully ligated a BAC construct with the HHV-6A genome and maintained it in *E. coli*; however, they could not reconstitute infectious virus using the BAC containing the HHV-6A genome in a T-cell line (Borenstein and Frenkel, 2009). Arbuckle et al. (2010) constructed an infectious HHV-6A virus harboring GFP and BAC sequences in the U53/U54 intergenic region, however, they did not take it through *E. coli* and back to human cells.

Here we cloned the HHV-6A strain U1102 genome as a BAC plasmid, and found that this HHV-6A BAC could be stably maintained in *E. coli*. Infectious virus was successfully reconstituted by transfecting the HHV-6A BAC into a T-cell line, followed by co-culturing with umbilical cord blood mononuclear cells (CBMCs). The expression of viral proteins was detected by Western blotting analysis, and the reconstituted virions were confirmed by electron microscopy (EM). We then generated in the BAC genome a deletion mutant of the gQ1 gene, and found that this mutation resulted in a failure to reconstitute the virus. The revertant could be reconstituted, suggesting that gQ1 was essential for HHV-6A propagation *in vitro*.

## Results

### Strategy of HHV-6A BAC construction and reconstitution of infectious virus

The plasmid pHA-2, containing the guanine phosphoribosyl transferase gene (*gpt*), *gfp*, *loxP*, and BAC sequences, was used in this study (Adler et al., 2000). The BAC construct was inserted into the HHV-6A genome between the U53 and U54 poly A sequences. The DNA fragments of U53 and U54 were amplified from the HHV-6A genome by PCR using the primers shown in Table 1. The PCR product was digested with *NotI* and *PacI* and ligated with pBluescript SK (–) (Stratagene) plasmid that had been digested with *NotI*. The resultant plasmid was then digested with *PacI* and ligated with the fragment containing the BAC sequence released from the pHA-2 digested with *PacI*. The construct was named pHA-U5354. After digestion with *NotI*, the plasmid was transfected into CBMCs by electroporation using a Nucleofection unit (Amaza).

On the next day, the CBMCs were co-cultured with HHV-6A (U1102 strain)-infected CBMCs. On the third day after the transfection, the recombination selection drugs mycophenolic acid and xanthine were added to the culture medium to a final concentration of 12.5 µg/ml and 110 µg/ml, respectively. The recombinant viruses were enriched by performing the cell-to-cell infection three times in the selection-drug medium, and then the circular viral DNAs were isolated from the infected CBMCs and transformed into *E. coli* DH10B

(Invitrogen) by electroporation using a Bio-Rad *E. coli* Pulser. The transformants were transferred to agar plates containing 17 µg/ml chloramphenicol. The recombination strategy is shown in Fig. 1.

The BACs from *E. coli* DH10B were isolated using NucleoBond AX purchased from Macherey-Nagel. HHV-6ABAC DNA was transfected into Jjhan cells by electroporation using a Nucleofection unit (Amaza). After electroporation, the cells were cultured in RPMI medium containing 8% FBS. On the second day after electroporation, the cells were co-cultured with CBMCs in RPMI medium containing 8% FBS. Green fluorescence from the infected CBMCs was easily confirmed after 4–5 days in co-culture. After three cell-to-cell infections (using CBMCs as the uninfected cells), the cells were harvested and prepared for Western blot analysis and EM.

### Successful reconstitution of infectious virus using HHV-6ABAC DNA

After the recombination between PHAU5354 and HHV-6A genome in CBMCs, the resultant circular DNAs were isolated and then transformed into DH10B competent cells. We cultured the positive colonies from the chloramphenicol plates and isolated the BACs as described above. These BAC DNAs were digested with *BamHI* and separated on agarose gels. As shown Fig. 2A, different digestion patterns of the BAC DNAs were observed, which might have reflected genetic variances within the U1102 strain before the insertion of the BAC into the HHV-6A genome. We then transfected the HHV-6ABAC DNAs into Jjhan cells, and enriched for the reconstituted virus by co-culturing the transfected Jjhan cells with CBMCs. Infectious virus could only be successfully reconstituted using BAC DNA from the G-1 strain (lane 1 in Fig. 2A). The proportion of GFP-positive cells increased dramatically when we performed the cell-to-cell infection three times (Fig. 2B).


### Confirmation of viral gene expression and infectious virions in HHV-6ABAC-transfected cells

Borenstein et al. successfully inserted a BAC sequence into the HHV-6A genome, but neither the viral protein nor HHV-6A virions could be identified in the HHV-6A BAC-transfected Sup T 1 cells. Viral protein expression could only be confirmed in the case of complementation by UV light-inactivated virus (Borenstein and Frenkel, 2009). We transfected Jjhan cells with HHV-6ABAC and then co-cultured them with CBMCs, which are commonly used for propagating HHV-6. After three rounds of cell-to-cell infection, the GFP expression from the BAC *gfp* gene was greatly increased (Fig. 2B).

We harvested these cells and prepared samples for Western blot analysis. The expression of U14, an early gene product of HHV-6 and gQ1, gH, and glycoprotein B (gB), which are late gene products of HHV-6, could be detected as in wild-type HHV-6-infected CBMCs

**Table 1**  
Primer sequence.

Primer name	Sequence
AgQ1Xho1F	5-accctcgagccaccatggcaaccgcaagactg-3
AgQ1693 Ecor1R	5-accgaattcacatagatgctcttccacttg-3
Agp105F922	5-tcttcaacctcatggactcag-3
AgQ1spstnotR	5-accctgcaggcggcccgacggagaatggcaaccgcaag-3
AgQ1de2F	5-acgaaggcttcagcccatagtttctactctgaaaggatgtagctctcagctctgtgaaaggatgacgacgataagtaggg-3
AgQ1de2R	5-gtttgtctcgcgcatacgaattttcaagaactgagacgaagactactctctcagatagaaccaaccaattaaccaattctgattag-3
ReF1	5-atgattacgaaggcttcag-3
ReR1	5-acaggatccatggcaaccgcaaggctgag-3
ReF2	5-acaggatccaggatgacgacgataagtag-3
ReR2	5-tgtttgtctcgcgcatacgaattttcaagaactgagacgaagatggcaaccgcaaggctgagcgtatgaaccgagatccaaccaattaaccaattctgattag-3
ReR3	5-tgtttgtctcgcgcatacgaattttc-3
U53Fnot1	5-accagcggccgctttggtaggtgattcctttg-3
U531650rpac1	5-accttaattaacactctgaattcttctataac-3
U541651fpac1	5-accttaattaataattctggacatggtgaaac-3
U54Rnot13650	5-accagcggccgacgacgatcttcaagctgtatc-3



**Fig. 1.** Strategy for cloning the HHV-6A BAC. (a) The HHV-6A genome consists of three major internal repeat elements (R1–R3), the origin of replication (*oriLyt*) and the direct repeat termini (*DR<sub>L</sub>* and *DR<sub>R</sub>*). (b) The recombinant plasmid, PHAU5354 contained an approximately 1.5-kbp sequence from U53, and 2.0-kbp sequence from U54 amplified from the HHV-6A genome by PCR, and the BAC vector sequence including the *gpt* gene, *gfp* gene, and *loxP* sites at both termini. The recombinant plasmid was digested with *NotI* and electroporated into CBMCs. These cells were then infected with HHV6A. Homologous recombination occurred between PHAU5354 and the HHV-6A genome. (c) The resultant HHV-6ABAC is depicted at the bottom of the figure. The circular recombinant viral genomes were then harvested and electroporated into DH10B. Single BAC clones were isolated.

(Fig. 3A). Interestingly, the 80-kDa gQ1 gene product (Akkapaiboon et al., 2004), which is incorporated into HHV-6A virions, was also detected. Next, we asked whether mature HHV-6 virions were also produced in these cells. We therefore observed these samples under EM. As shown in Fig. 3B, mature virions could be identified in both the cytoplasmic and extracellular spaces. This observation confirmed that we had successfully reconstituted the infectious virus of HHV-6A by transfecting HHV-6ABAC DNA into JJHAN cells and co-culturing them with CBMCs.

#### Generation of a gQ1 deletion mutant virus and its revertant virus

The main purposes of cloning intact viral genomes as BACs are to manipulate *E. coli* mutagenesis systems to disrupt the viral genes and to analyze the biocharacteristics of the mutant viruses. The gQ gene is unique to HHV-6. The gQ1 protein is reported to form a tetrameric

complex with gQ2, gH, and gL that acts as viral ligand for CD46, the cellular receptor for HHV-6A (Akkapaiboon et al., 2004). However, the specific function of gQ1 in HHV-6-infected cells, or even whether it is essential for HHV-6 propagation, has been unknown. In addition, even if gQ1 is non-essential, it is not known what, if any, phenotype results when the gQ1 gene is knocked out.

We therefore constructed a gQ1 mutant virus, and analyzed its function in HHV-6 infection. The N-terminal segment of gQ1 was deleted from the HHV-6A genome, and its revertant was also generated. (The detailed procedure can be found in Materials and methods, and is depicted in Fig. 4). The resultant BAC DNAs (HHV-6ABACΔgQ1 and HHV-6ABACΔgQ1re) and HHV-6ABAC DNA were examined by restriction digestion with *Bam*H1.

As shown in Fig. 5, the digestion of HHV-6ABACΔgQ1re (lane 3) with *Bam*H1 yielded the same pattern as HHV-6ABAC DNA (lane 1); however, a different digestion pattern was seen with HHV-