

本研究は、東北大学動物実験専門委員会の承認を得た上で、実験動物に対する十分な動物愛護上の配慮の下、実験を行っている。膵炎患者に対する膵Perfusion CTによる血流評価は東北大学倫理委員会の承認を得て行っており、患者に十分な説明を行い、同意を得て行っている。

#### C. 研究結果

衝撃波治療群と対照群のいずれの群においても、5ヵ月齢から死亡例が増加し、7ヵ月齢までに全例死亡した。血糖値、血中インスリン、リパーゼ、トリプシンの値に両群間で差を認めなかった。今後、組織学的な検討を行う。

膵臓に慢性的な炎症細胞浸潤と線維化が認められる、自己免疫性膵炎患者11名に対して膵Perfusion CTを行った。自己免疫性膵炎患者では、67%(6/9)の患者に膵外分泌障害を認め、27%(3/11)の患者に糖尿病を認めた。自己免疫性膵炎患者では、対照群と比較し有意にFv値が低下していた。

#### D. 考察

衝撃波治療を行わなかった対照群においても死亡率が高かったことから、頻回の麻酔が負荷になった可能性が考えられた。一方、血中膵酵素の増加を認めなかったことから、今回用いた低出力の衝撃波は、膵実質の破壊を起ささない安全な出力と考えられた。今後、組織学的な検討を行うとともに、実験プロトコルの再検討も行う。

膵Perfusion CTを用いた検討において、膵機能

低下に伴い、膵血流の低下が示唆された。今後、慢性膵炎発症ラットを用いた、膵機能と膵血流の評価、及び衝撃波治療による血流改善の評価を行う。

#### E. 結論

慢性膵炎に対する低出力体外衝撃波治療の有効性については、さらなる検討を要する。膵血流評価法の確立にもさらなる検討が必要である。

#### G. 研究発表

##### 1. 論文発表

① Hirota M, Tsuda M, Tsuji Y, et al. Perfusion computed tomography findings of autoimmune pancreatitis. *Pancreas* 2011, (in press).

##### 2. 学会発表

② 廣田衛久, 津田雅視, 下瀬川徹. 自己免疫性膵炎患者の膵Perfusion CT. 第1回膵Perfusion CT画像研究会 (2010年4月21日、新潟) .

③ Hirota M, Satoh K, Masamune A, et al. Pancreatic perfusion analysis of autoimmune pancreatitis. 第14回国際膵臓学会・第41回日本膵臓学会 (2010年7月11-13日、福岡) .

#### H. 知的財産権の出願・登録状況

なし

III. 研究成果の刊行に関する一覧表

発表者氏名	論文タイトル名	発表誌名	巻号	ページ	出版年
Kikuchi Y, Ito K, Ito Y, Shiroto T, Tsuburaya R, Aizawa K, Hao K, Fukumoto Y, Takahashi J, Takeda M, Nakayama M, Yasuda S, Kuriyama S, Tsuji I, Shimokawa H.	Double-blind and placebo-controlled study of the effectiveness and safety of extracorporeal cardiac shock wave therapy for severe angina pectoris.	Circ J.	74	589-591	2010
Ito Y, Ito K, Shiroto T, Tsuburaya R, Gao JY, Takeda M, Fukumoto Y, Yasuda S, Shimokawa H.	Cardiac shock wave therapy ameliorates left ventricular remodeling after myocardial ischemia reperfusion injury in pigs in vivo.	Coron Artery Dis.	21	304-311	2010
伊藤健太, 下川宏明	体外衝撃波を用いた非侵襲性血管新生治療	日内学雑誌	99	188-194	2010
Akamatsu D, Sato A, Goto H, Watanabe T, Hashimoto M, Shimizu T, Sugawara H, Sato H, Nakano Y, Miura T, Zukeran T, Serizawa F, Hamada Y, Tsuchida K, Tsuji I, Satomi S.	Nitroglycerin-mediated vasodilatation of the brachial artery may predict long-term cardiovascular events irrespective of the presence of atherosclerotic disease.	J Atheroscler Thromb.	17	1266-1274	2010
Serizawa F, Ito K, Matsubara M, Sato A, Shimokawa H, Satomi S.	Extracorporeal shock wave therapy induces therapeutic lymphangiogenesis in a rat model of secondary lymphoedema.	Eur J Vasc Endovasc Surg.	in press	in press	2011

#### IV. 研究成果の刊行物・別刷



## Double-Blind and Placebo-Controlled Study of the Effectiveness and Safety of Extracorporeal Cardiac Shock Wave Therapy for Severe Angina Pectoris

Yoku Kikuchi, MD; Kenta Ito, MD; Yoshitaka Ito, MD; Takashi Shiroto, MD; Ryuji Tsuburaya, MD; Kentaro Aizawa, MD; Kiyotaka Hao, MD; Yoshihiro Fukumoto, MD; Jun Takahashi, MD; Morihiko Takeda, MD; Masaharu Nakayama, MD; Satoshi Yasuda, MD; Shinichi Kuriyama, MD\*; Ichiro Tsuji, MD\*; Hiroaki Shimokawa, MD

**Background:** Low-energy shock wave (SW) therapy has improved myocardial ischemia in both a porcine model and in patients with severe angina pectoris.

**Methods and Results:** To further confirm the effectiveness and safety of SW therapy, 8 patients with severe angina pectoris were treated with SW therapy in a double-blind, placebo-controlled and cross-over manner. SW therapy, but not placebo, significantly improved chest pain symptoms and cardiac function without any complications or adverse effects.

**Conclusions:** Extracorporeal cardiac SW therapy is an effective, safe and non-invasive therapeutic option for severe angina pectoris. (*Circ J* 2010; **74**: 589–591)

**Key Words:** Angina pectoris; Angiogenesis; Myocardial ischemia; Shock wave

The number of patients with severe angina pectoris without indications for coronary artery bypass grafting (CABG) or percutaneous coronary intervention (PCI) is rapidly increasing worldwide and their prognosis still remains poor.<sup>1,2</sup> Thus, it is crucial to develop new therapeutic strategies for these patients. We have previously demonstrated that extracorporeal cardiac shock wave (SW) therapy with low-energy SW ( $\approx 10\%$  of the energy density used for urolithiasis) ameliorates myocardial ischemia and dysfunction in a porcine model of chronic myocardial ischemia *in vivo*.<sup>3,4</sup> We subsequently demonstrated in an open trial that our SW therapy effectively improved chest pain symptoms and exercise tolerance without any adverse effects in 9 patients with severe angina pectoris.<sup>3,5</sup> In the present study, to further confirm the effectiveness and safety of our SW therapy, we performed a double-blind placebo-controlled trial in patients with severe angina pectoris.

### Methods

We enrolled 8 consecutive patients with severe angina pectoris who already had undergone CABG or PCI, but who no longer had further indications for these therapies even though they still suffered from stable effort angina under

intensive medication (M/F, 5/3; age,  $70 \pm 3$  years) (Table).

The patients were treated with one series of placebo and the SW therapy in a double-blind and cross-over manner with an interval of 3 months. One series of therapy comprised 3 sessions per week. Throughout the study, the patient and the doctor in charge were not informed of the type of therapy. We performed the SW therapy (200 shots/spot at  $0.09 \text{ mJ}/\text{mm}^2$  for 40–60 spots per session; Modulith SLC, Storz Medical, Kreuzlingen, Switzerland) as described previously.<sup>3,5</sup> As placebo, the patients underwent the procedure of SW therapy but without irradiation. The patients were followed-up for 3 months after completion of the therapy. We evaluated symptoms using the Canadian Cardiovascular Society (CCS) class score, the patient's requirement for nitroglycerin,<sup>5</sup> exercise tolerance in a 6-min walk, and cardiopulmonary exercise test, and cardiac function assessed by MRI (Achieva 1.5 T, Philips, Eindhoven, Netherlands). The left ventricular ejection fraction (LVEF) was measured using contiguous short-axis slices obtained by cine MRI; end-diastolic and end-systolic endocardial traces were used to determine end-diastolic and end-systolic left ventricular (LV) volumes, respectively. We also evaluated the number of circulating progenitor cells in peripheral blood by FACS analysis 2 days before the 1<sup>st</sup> session and 1 h after the 3<sup>rd</sup> session in 7 of the 8 patients

Received December 25, 2009; revised manuscript received January 6, 2010; accepted January 11, 2010; released online February 4, 2010 Time for primary review: 11 days

Department of Cardiovascular Medicine, \*Department of Public Health and Forensic Medicine, Tohoku University Graduate School of Medicine, Sendai, Japan


Mailing address: Kenta Ito, MD, Department of Cardiovascular Medicine, Tohoku University Graduate School of Medicine, 1-1 Seiryomachi, Aoba-ku, Sendai 980-8574 Japan. E-mail: ito-kenta@cardio.med.tohoku.ac.jp

ISSN-1346-9843 doi:10.1253/circj.CJ-09-1028

All rights are reserved to the Japanese Circulation Society. For permissions, please e-mail: [cj@j-circ.or.jp](mailto:cj@j-circ.or.jp)

Patient no.	Age (years)	Gender	CAD	Previous treatment	OMI	ASO	HT	DM	HL
1	69	M	3VD	CABG, PCI	+	+	+	+	+
2	61	M	3VD	CABG, PCI	+	-	+	+	+
3	70	M	3VD	CABG	+	+	+	-	+
4	78	F	3VD	CABG, PCI	+	-	+	-	+
5	80	M	3VD	CABG, PCI	+	-	+	+	+
6	60	F	3VD	CABG, PCI	+	-	+	+	+
7	72	F	3VD	CABG, PCI	+	-	-	-	+
8	70	M	3VD	CABG, PCI	+	+	+	+	+

CAD, coronary artery disease; OMI, old myocardial infarction; ASO, arteriosclerosis obliterans; HT, hypertension; DM, diabetes mellitus; HL, hyperlipidemia; VD, vessel disease; CABG, coronary artery bypass grafting; PCI, percutaneous coronary intervention.



(technical problem in the remaining one patient).

The present study was approved by the Ethical Committees of Tohoku University in 2005, and informed consent was given by each patient.

Results are expressed as mean  $\pm$  SEM. Comparisons during the time course after SW therapy were made by repeated measure ANOVA followed by Bonferroni/Dunn post hoc test. All statistical analyses were performed using StatView (SAS Institute, Cary, NC, USA), and  $P < 0.05$  was considered to be statistically significant.

## Results

The SW therapy, but not placebo, significantly improved symptoms (CCS class score) and nitroglycerin use (Figure).

The SW therapy also significantly improved the 6-min walking distance and tended to improve both maximum exercise capacity and peak oxygen uptake (peak  $\dot{V}O_2$ ). LVEF and LV stroke volume evaluated by MRI were significantly improved only with the SW therapy, although LV end-diastolic volume and plasma brain natriuretic peptide level remained unchanged. The number of CD34<sup>+</sup>/KDR<sup>+</sup> and CD34<sup>+</sup>/KDR<sup>-</sup>/c-kit<sup>+</sup> cells in peripheral blood also remained unchanged with both therapies (data not shown). No procedural complications or adverse effects were noted during or after either therapy as in the previous studies.<sup>3-7</sup>

## Discussion

We have previously demonstrated that low-energy SW

therapy enhanced angiogenesis and improved myocardial ischemia in a pig model of chronic myocardial ischemia,<sup>3,4</sup> and that SW therapy improved the symptoms and myocardial perfusion in patients with severe angina pectoris in an open trial.<sup>15</sup> The present double-blind and placebo-controlled study further demonstrates that our extracorporeal cardiac SW therapy is an effective therapeutic option for severe angina pectoris, providing convincing evidence for its effectiveness and safety.

During the past 2 decades, regenerative therapies using genes, cytokines, and progenitor cells have been under investigation for ischemic cardiovascular diseases.<sup>8</sup> However, these therapies have not been consistently effective in humans, despite promising results in early preclinical studies.<sup>8,12</sup> A potential explanation for these inconsistent results is the complex crosstalk among multiple pathways, in which enhancement of only 1 factor among numerous angiogenic factors may not be enough to achieve clinical benefit. Furthermore, animal studies of cell therapy have revealed that the number of newly generated vascular cells is too low to induce any functional improvement, suggesting that the paracrine action of transplanted cells stimulates intrinsic angiogenic capacity.<sup>13</sup> In contrast, low-energy SW upregulates multiple angiogenic pathways (eg, VEGF, flt-1, SDF-1, and nitric oxide synthase).<sup>4,7,14</sup>

There are several limitations to the present study. First, the number of patients is small. Although more than 150 patients with severe angina pectoris were reviewed as potential candidates for this study, most of them were excluded due to insufficient medication, potential indications of CABG or PCI, and co-existence of malignant tumor. However, we were able to reconfirm the beneficial effects of SW therapy in the present double-blind and placebo-control study, as we had observed in a previous open study.<sup>5</sup> A future large-scale trial would validate the present results. Second, maximum exercise capacity and pVO<sub>2</sub> were not significantly improved while the symptoms and 6-min walking distance were significantly improved by the SW therapy. In 6 of the 8 patients, exercise was stopped because of leg pain or fatigue before reaching the anaerobic threshold. Thus, exercise tolerance might have been underestimated because of arteriosclerosis obliterans and/or physical deconditioning. Another parameter, such as arteriovenous difference in lactate concentration under overdrive pacing, might have been a better index of ischemia. Third, the number of circulating progenitor cells in peripheral blood was not increased in the present study. Thus, it remains to be examined whether our SW therapy promotes recruitment of bone marrow-derived cells by the ischemic myocardium of humans.<sup>14,15</sup> Fourth, the detailed molecular mechanisms of the beneficial effects of SW in humans remain to be clarified in future studies.<sup>3-7</sup>

In conclusion, the present double-blind, placebo-controlled study further confirmed the effectiveness and safety of our extracorporeal cardiac SW therapy for the treatment of severe

angina pectoris, although large-scale multi-center study is needed.

### Acknowledgments

We thank Dr EH Maringhaus, Storz Medical AG, Switzerland, for valuable comments. This study was supported in part by the grants-in-aid from the Japanese Ministry of Education, Culture, Sports, Science and Technology, Tokyo, and the Japanese Ministry of Health, Labor and Welfare, Tokyo, Japan.

### References

- Ma E, Iso H, Takahashi H, Yamagishi K, Tanigawa T. Age-period-cohort analysis of mortality due to ischemic heart disease in Japan, 1955 to 2000. *Circ J* 2008; **72**: 966–972.
- World Health Organization. Program Fact Sheet #310: 2004 Death Rates. Available at: <http://www.who.int/mediacentre/factsheets/fs310/en/index.html> (accessed 6 January, 2010).
- Ito K, Fukumoto Y, Shimokawa H. Extracorporeal shock wave therapy as a new and non-invasive angiogenic therapy. *Tohoku J Exp Med* 2009; **219**: 1–9.
- Nishida T, Shimokawa H, Oi K, Tatewaki H, Uwatuoku T, Abe K, et al. Extracorporeal cardiac shock wave therapy markedly ameliorates ischemia-induced myocardial dysfunction in pigs in vivo. *Circulation* 2004; **110**: 3055–3061.
- Fukumoto Y, Ito A, Uwatuoku T, Matoba T, Kishi T, Tanaka H, et al. Extracorporeal cardiac shock wave therapy ameliorates myocardial ischemia in patients with severe coronary artery disease. *Coron Artery Dis* 2006; **17**: 63–70.
- Uwatuoku T, Ito K, Abe K, Oi K, Hizume T, Sunagawa K, et al. Extracorporeal cardiac shock wave therapy improves left ventricular remodeling after acute myocardial infarction in pigs. *Coron Artery Dis* 2007; **18**: 397–404.
- Oi K, Fukumoto Y, Ito K, Uwatuoku T, Abe K, Hizume T, et al. Extracorporeal shock wave therapy ameliorates hindlimb ischemia in rabbits. *Tohoku J Exp Med* 2008; **214**: 151–158.
- Gupta R, Tongers J, Losordo DW. Human studies of angiogenic gene therapy. *Circ Res* 2009; **105**: 724–736.
- Epstein SE, Fuchs S, Zhou YF, Baffour R, Kornowski R. Therapeutic interventions for enhancing collateral development by administration of growth factors: Basic principles, early results and potential hazards. *Cardiovasc Res* 2001; **49**: 532–542.
- Dimmeler S, Zeith AM, Schneider MD. Unchain my heart: The scientific foundations of cardiac repair. *J Clin Invest* 2005; **115**: 572–583.
- Losordo DW, Schatz RA, White CJ, Udelson JE, Veereshwarayya V, Durgin M, et al. Intramyocardial transplantation of autologous CD34<sup>+</sup> stem cells for intractable angina: A phase I/IIa double-blind, randomized controlled trial. *Circulation* 2007; **115**: 3165–3172.
- van Ramshorst J, Bax JJ, Beeres SL, Dibbets-Schneider P, Roes SD, Stokkel MP, et al. Intramyocardial bone marrow cell injection for chronic myocardial ischemia: A randomized controlled trial. *JAMA* 2009; **301**: 1997–2004.
- Gnecchi M, Zhang Z, Ni A, Dzau VJ. Paracrine mechanisms in adult stem cell signaling and therapy. *Circ Res* 2008; **103**: 1204–1219.
- Aicher A, Heeschen C, Sasaki K, Urbich C, Zeith AM, Dimmeler S. Low-energy shock wave for enhancing recruitment of endothelial progenitor cells: A new modality to increase efficacy of cell therapy in chronic hind limb ischemia. *Circulation* 2006; **114**: 2823–2830.
- Yip HK, Chang LT, Sun CK, Youssef AA, Sheu JJ, Wang CJ. Shock wave therapy applied to rat bone marrow-derived mononuclear cells enhances formation of cells stained positive for CD31 and vascular endothelial growth factor. *Circ J* 2008; **72**: 150–156.

# Cardiac shock wave therapy ameliorates left ventricular remodeling after myocardial ischemia-reperfusion injury in pigs *in vivo*

Yoshitaka Ito, Kenta Ito, Takashi Shiroto, Ryuji Tsuburaya, Gao Jun Yi, Morihiro Takeda, Yoshihiro Fukumoto, Satoshi Yasuda and Hiroaki Shimokawa

**Objectives** Left ventricular (LV) remodeling after acute myocardial infarction (AMI) is associated with a poor prognosis and an impaired quality of life. We have shown earlier that low-energy extracorporeal cardiac shock wave (SW) therapy improves chronic myocardial ischemia in pigs and humans and also ameliorates LV remodeling in a pig model of AMI induced by permanent coronary ligation. However, in the current clinical setting, most of the patients with AMI receive reperfusion therapy. Thus, in this study we examined whether our SW therapy also ameliorates LV remodeling after myocardial ischemia-reperfusion (I/R) injury in pigs *in vivo*.

**Methods** Pigs were subjected to a 90-min ischemia and reperfusion using a balloon catheter and were randomly assigned to two groups with or without SW therapy to the ischemic border zone ( $0.09 \text{ mJ/mm}^2$ , 200 pulses/spot, 9 spots/animal, three times in the first week ( $n=15$  each).

**Results** Four weeks after I/R, compared with the control group, the SW group showed significantly ameliorated LV remodeling in terms of LV enlargement ( $131 \pm 9$  vs.  $100 \pm 7$  ml), reduced LV ejection fraction ( $28 \pm 2$  vs.  $36 \pm 3\%$ ), and elevated left ventricular

end-diastolic pressure ( $11 \pm 2$  vs.  $4 \pm 1$  mmHg) (all  $P<0.05$ ,  $n=8$  each). The SW group also showed significantly increased regional myocardial blood flow ( $-0.06 \pm 0.11$  vs.  $0.36 \pm 0.13 \text{ ml/min/g}$ ,  $P<0.05$ ), capillary density ( $1.233 \pm 31$  vs.  $1.560 \pm 60/\text{mm}^2$ ,  $P<0.001$ ), and endothelial nitric oxide synthase activity ( $0.24 \pm 0.03$  vs.  $0.41 \pm 0.05$ ,  $P<0.05$ ) in the ischemic border zone compared with the control group ( $n=7$  each).

**Conclusion** These results indicate that our SW therapy is also effective in ameliorating LV remodeling after myocardial I/R injury in pigs *in vivo*. *Coron Artery Dis* 21:304–311 © 2010 Wolters Kluwer Health | Lippincott Williams & Wilkins.

Coronary Artery Disease 2010, 21:304–311

Keywords: angiogenesis, reperfusion, shock wave, ventricular remodeling

Department of Cardiovascular Medicine, Tohoku University Graduate School of Medicine, Sendai, Japan

Correspondence to: Dr Kenta Ito, MD, PhD, Department of Cardiovascular Medicine, Tohoku University Graduate School of Medicine, 1-1 Seiryomachi, Aoba-ku, Sendai, 980-8574, Japan  
Tel: +81 22 717 7153; fax: +81 22 717 7156;  
e-mail: ito-kenta@cardio.med.tohoku.ac.jp

Received 5 January 2010 Revised 20 March 2010  
Accepted 2 April 2010

## Introduction

Ischemic heart disease is the leading cause of death in western countries. The development of left ventricular (LV) remodeling after acute myocardial infarction (AMI) leads to sudden cardiac death, heart failure, and poor prognosis. Thus, it is important to improve LV remodeling after AMI to improve prognosis and the quality of life. Several regenerative therapies, such as gene [1–3] and cell therapies [4–8], are currently under development; however, most of these are invasive in nature and their effectiveness and safety have not yet been fully established. Thus, more effective and less invasive therapies need to be developed.

We have shown earlier that low-energy extracorporeal cardiac shock wave (SW) therapy effectively induces angiogenesis and improves cardiac functions in a porcine model of chronic myocardial ischemia [9], and that SW therapy improves symptoms, reduces the use of nitroglycerin, and improves myocardial perfusion in patients with end-stage coronary artery disease [10,11]. Furthermore, we have recently shown that SW therapy improves LV

remodeling in a porcine model of AMI with permanent coronary ligation [12]. However, in the current clinical setting, most patients with AMI receive emergency reperfusion therapy with either percutaneous coronary intervention or thrombolytic agents. It remains to be determined whether our extracorporeal cardiac SW therapy also ameliorates myocardial ischemia-reperfusion (I/R) injury *in vivo*. Thus, in this study we examined whether our SW therapy also ameliorates LV remodeling after myocardial I/R injury in pigs *in vivo*, and if so, what mechanism(s) might be involved.

## Methods

The investigation conforms with the Guide for the Care and Use of Laboratory Animals established by the US National Institutes of Health (Publication No. 85-23, revised 1996). All procedures were performed according to the protocols approved by the Institutional Committee for Use and Care of Laboratory Animals at Tohoku University (20-Idou-151 and 21-Idou-156).

### Porcine model of myocardial I/R


A total of 30 domestic male pigs (25–30 kg in body weight) were used in this study. They underwent myocardial I/R injury with and without SW therapy. They were subjected to cardiac catheterization and histology study at 4 weeks after I/R ( $n = 8$  each) and to western blotting study at 1 week after I/R ( $n = 7$  each). The animals were anesthetized with ketamine hydrochloride (15 mg/kg, intramuscular), and after intubation, they were kept anesthetized with an inhalation of 2.0% sevoflurane for cardiac catheterization and euthanization. We inserted a 7F sheath into the left carotid artery for cardiac catheterization. A 5000-IU bolus of heparin was administered intravenously and 2000-IU was injected every hour. We performed a left ventriculography (LVG) and coronary angiography (CAG) in a left oblique view with the use of a cineangiography system (Toshiba Medical, Tochigi, Japan) [9,12]. LV volume and LV ejection fraction (LVEF) were calculated using Simpson's method. A coronary angioplasty balloon (2.5–3.5 mm in diameter depending on the vessel size) was then introduced into the left anterior descending coronary

artery (LAD) and inflated just distal to the first diagonal branch for 90 min, which has been shown earlier to effectively induce myocardial infarction [13,14], at the lowest pressure that completely occluded distal flow (Fig. 1). After 90 min of ischemia, the balloon was deflated and both CAG and LVG were reperformed to confirm the patency of distal LAD and the reduced LV wall motion, respectively. Cardiac catheterization was performed before ischemia, immediately after reperfusion, and 4 weeks after I/R. After the study, 4 weeks after I/R, the animals were euthanized by an overdose of pentobarbital.

### Extracorporeal cardiac SW therapy

On the basis of our earlier studies [9–12,15,16], we applied a low-energy SW (0.09 mJ/mm<sup>2</sup>, approximately 10% of the energy used for the lithotripsy treatment, 200 shots/spot for 27 spots) to the border zone around the infarcted myocardium with the guidance of an echocardiogram equipped within the specially designed SW generator (Storz Medical AG, Kreuzlingen, Switzerland) in an R-wave-triggered manner to avoid ventricular

Fig. 1



Porcine model of myocardial ischemia–reperfusion (I/R). Coronary angiograms at baseline (a), during balloon inflation in the left anterior descending coronary artery (b), immediately after reperfusion (c), and 4 weeks after the I/R in the same pig. The inflated balloon is shown by an arrowed line (b) and the ischemic myocardial area by the shaded area (c and d).

arrhythmias. In a preliminary study, we confirmed that no adverse effects, such as cardiac rupture or tamponade, were noted even if we applied a SW to the infarcted myocardium (data not shown). We examined LV wall motion by echocardiography during I/R and defined the border zone as the edge of the area where the LV wall motion was severely depressed after I/R. We were able to accurately focus a SW to any part of the heart under the guidance of echocardiography with a focus of approximately 2 mm [9–12]. We performed the SW treatment three times in the first week (day 1, 3, and 5), whereas the animals in the control group received the same procedures three times but without the SW treatment.

#### Cardiac enzymes

We measured serum concentrations of cardiac troponin T and creatinine kinase myocardial blood isoform (CK-MB) using an electrochemiluminescence immunoassay and a chemiluminescence immunoassay, respectively. Blood samples were serially collected before and 5, 12, 24, 48, and 72 h after the I/R injury, and the extent of myocardial infarction was expressed as the area under the curves of troponin T and CK-MB [12].

#### Echocardiography

We performed a transthoracic echocardiographic study (Aplio 80, Toshiba Medical). We calculated the wall thickening fraction (WTF, %) by using the following formula:  $WTF = 100 \times (\text{end-systolic wall thickness} - \text{end-diastolic wall thickness}) / \text{end-diastolic wall thickness}$  [9]. We measured the WTF in the infarcted area and the border zone when the animals were sedated.

#### Regional myocardial blood flow

We evaluated regional myocardial blood flow (RMBF) with colored microspheres (Dye-Trak VII+, Triton Technology, San Diego, USA) ( $n = 4$  each) [9,12]. We injected 6 million microspheres (diameter 15  $\mu\text{m}$ ) into the left atrium before the induction of myocardial ischemia and 4 weeks after I/R. We drew a reference arterial blood sample from the descending aorta at a constant rate of 12 ml/min for 90 s using a withdrawal pump. We extracted microspheres from the LV wall and blood samples by potassium hydroxide digestion, extracted the dyes from the microspheres with ethylene glycol monoethyl ether acetate (70  $\mu\text{l}$ ), and determined their concentrations by spectrophotometry. We calculated the change of myocardial blood flow (ml/min/g) in the infarcted region and border zone.

#### Myocardial capillary density

The heart was removed and 10% formaldehyde was injected into the left coronary artery with a pressure of 100–120 mmHg. After fixation, tissue specimens were obtained from the border zone of each animal. We treated the paraffin-embedded sections with a rabbit anti-factor VIII antibody (N1505, Dako, Copenhagen, Denmark), and counted the number of factor VIII-positive cells in 10 random fields of the

border zone and the remote area in each heart at  $\times 400$  magnification, and calculated capillary density [9,12]. Ten random fields of each sample were examined in a blinded manner. Each field covered 0.036 mm<sup>2</sup>.

#### Myocardial fibrosis

Masson-trichrome staining was performed using the paraffin-embedded sections. We evaluated the fibrosis area in 10 random fields of the border zone in each heart at  $\times 200$  magnification. A digital image processing software AxioVision 4.5.0.0 (Carl Zeiss, Göttingen, Germany) was used to detect the myocardial fibrosis area, and the ratio of the fibrosis area to the myocardial area was calculated.

#### Western blot analysis

To examine the mechanisms of the inhibitory effects of SW therapy on LV remodeling, another set of animals with I/R injury, with and without SW therapy, were made and they were euthanized at 1 week after the procedure. We performed western blot analysis for phosphorylated endothelial nitric oxide synthase (phospho-eNOS) and vascular endothelial growth factor (VEGF). Samples from the border zone were used and the extracted samples (50  $\mu\text{g}$  of protein) were subjected to SDS-PAGE/immunoblot analysis by using the specific antibody for phospho-eNOS at Ser1177 (No. 9571, Cell Signaling Technology, Danvers, Massachusetts, USA), total-eNOS (No. 610296, Becton Dickinson, Franklin Lakes, New Jersey, USA), and VEGF (sc-152, Santa Cruz Biotechnology, Santa Cruz, California, USA). The regions containing proteins were visualized by an electrochemiluminescence western blotting luminal reagent (RPN2132, GE Healthcare Bioscience, Waukesha, Wisconsin, USA). The extents of eNOS phosphorylation and VEGF expression were normalized by that of total-eNOS and  $\beta$ -actin, respectively [9,17].

#### Statistical analysis

Results were expressed as mean  $\pm$  SEM. We determined the statistical significance by an analysis of variance for multiple comparisons and the unpaired Student's *t*-test. Values of *P* less than 0.05 were considered to be statistically significant.

## Results

#### Extent of myocardial infarction

The extent of myocardial infarction, when evaluated by the area under the curve of troponin T or CK-MB, was comparable between the control and the SW groups (troponin T,  $296 \pm 32$  vs.  $319 \pm 30$  ng/ml<sup>h</sup>,  $P = 0.61$ ; CK-MB,  $1.641 \pm 301$  vs.  $1.993 \pm 353$  ng/ml<sup>h</sup>,  $P = 0.46$ ), indicating that the extent of myocardial infarction was comparable between the two groups.

#### Safety of SW therapy

No procedural complications or adverse effects related to SW therapy were noted throughout the experiments.

### Cardiac catheterization: CAG and LVG

At 4 weeks after the I/R injury, CAG confirmed the patency of reperfused LAD in all pigs (Fig. 1). Before and immediately after I/R (before the SW treatment), LV end-diastolic volume (LVEDV), LV end-systolic volume (LVESV), and LVEF were all comparable between the two groups (Fig. 2). Four weeks after I/R, LVG showed marked LV enlargement and reduced LVEF in the control group (Fig. 2). In contrast, LV enlargement and reduced LVEF were significantly ameliorated in the SW group. (LVEDV,  $100 \pm 7$  vs.  $131 \pm 9$  ml,  $P < 0.05$ ; LVESV,  $65 \pm 8$  vs.  $95 \pm 7$  ml,  $P < 0.05$ ; LVEF,  $36 \pm 3$  vs.  $28 \pm 2\%$ ,  $P < 0.05$ ) (Fig. 2). Although LV end-diastolic pressure was comparable between the two groups before and immediately after I/R (before the SW treatment), it remained elevated in the control group but was normalized in the SW group 4 weeks after I/R ( $11 \pm 2$  vs.  $4 \pm 1$  mmHg,  $P < 0.05$ ) (Fig. 2).

### Echocardiography

We measured the WTF of the infarcted region and the border zone by transthoracic echocardiography. The WTF in the infarcted region was significantly decreased to the same extent after I/R and was comparable between the control and the SW groups throughout the experimental period (before I/R,  $22 \pm 2$  vs.  $20 \pm 1\%$ ; immediately after I/R,  $2 \pm 1$  vs.  $1 \pm 0.3\%$ ; 4 weeks,  $4 \pm 2$  vs.  $5 \pm 2\%$ ) (Fig. 3a). In contrast, 4 weeks after I/R, the WTF was significantly improved at the border zone in the SW group as compared with the control

group (before I/R,  $24 \pm 2$  vs.  $22 \pm 2\%$ ,  $P = 0.54$ ; immediately after I/R,  $16 \pm 1$  vs.  $15 \pm 2\%$ ,  $P = 0.59$ ; and 4 weeks,  $15 \pm 2$  vs.  $24 \pm 4\%$ ,  $P < 0.05$ ) (Fig. 3b).

### Regional myocardial blood flow

In the infarcted region, RMBF was equally decreased in the control and the SW groups at 4 weeks after I/R as compared with before I/R ( $-0.52 \pm 0.22$  vs.  $-0.49 \pm 0.08$  ml/min/g,  $P = 0.89$ ), whereas RMBF at the border zone was significantly increased only in the SW group (control:  $-0.06 \pm 0.11$  vs. SW:  $0.36 \pm 0.13$  ml/min/g,  $P < 0.05$ ) (Fig. 4).


### Histopathology

Factor VIII staining showed that 4 weeks after I/R the number of factor VIII-positive blood vessels at the border zone was significantly higher in the SW group than in the control group ( $1.560 \pm 60$  vs.  $1.233 \pm 31/\text{mm}^2$ ,  $P < 0.001$ ) (Fig. 5a–c). In the remote area, the number of vessels was comparable between the two groups (Fig. 5d). Masson-trichrome staining showed that there was no difference in the extent of myocardial fibrosis at the border zone between the two groups (control,  $0.15 \pm 0.02$  vs. SW,  $0.13 \pm 0.03$ ,  $P = 0.72$ ).

### Western blot analysis


Western blot analysis showed that the ratio of phospho-eNOS to total-eNOS, a marker of eNOS activation, was

Fig. 2




The shock wave (SW) therapy ameliorates left ventricular (LV) remodeling after myocardial ischemia–reperfusion (I/R). The SW therapy significantly ameliorated LV remodeling as evaluated by LV end-diastolic volume (LVEDV) (a), LV end-systolic volume (LVESV) (b) and LV ejection fraction (LVEF) (c) and also normalized LV end-diastolic pressure (LVEDP) (d). Pre, before I/R; Post, immediately after I/R; 4 weeks, 4 weeks after I/R.

Fig. 3



The shock wave (SW) therapy ameliorates left ventricular (LV) systolic function. An echocardiographic study showed that the wall thickening fraction (WTF) in the infarcted region was comparable between the two groups throughout the study period (a), whereas the WTF in the border zone was normalized by the SW therapy at 4 weeks after the ischemia-reperfusion (I/R) injury (b). Pre, before I/R; Post, immediately after I/R; 4 weeks, 4 weeks after I/R.

Fig. 4



The shock wave (SW) therapy ameliorates myocardial blood flow. At 4 weeks after ischemia-reperfusion, although regional myocardial blood flow (RMBF) in the infarcted area was equally reduced in the control and SW groups, the flow in the border zone was significantly increased only in the SW group.

significantly increased in the SW group than in the control group 1 week after I/R ( $0.41 \pm 0.05$  vs.  $0.24 \pm 0.03$ ,  $P < 0.05$ ) (Fig. 6a). The protein expression of VEGF also tended to be increased in the SW group compared with the control group 1 week after I/R ( $0.78 \pm 0.26$  vs.  $0.40 \pm 0.12$ ,  $P = 0.22$ ) (Fig. 6b).

## Discussion

The novel finding of this study is that our extracorporeal cardiac SW therapy ameliorates LV remodeling after myocardial I/R injury in pigs *in vivo*. Importantly, no

procedural complications or adverse effects with SW therapy were noted in this study, a consistent finding with our earlier studies for chronic myocardial ischemia in pigs and humans, AMI in pigs, and hind limb ischemia in rabbits [9–12,15,16,18].


### Inhibitory effects of the SW therapy on LV remodeling after I/R

Although short-term and long-term outcomes of patients with AMI have improved during the last decades as reperfusion therapy became widely available in emergency care [19–22], LV remodeling after AMI still remains one of its major complications [23]. We have recently shown that our SW therapy ameliorates LV remodeling after AMI with permanent coronary ligation in pigs *in vivo* [12]. However, in the current clinical setting, most of the AMI patients are treated with emergency reperfusion therapy. In this study, to simulate the current situation with reperfusion therapy, we examined the possible beneficial effects of our SW therapy in a porcine model of myocardial I/R *in vivo*. In this model, severe LV remodeling characterized by marked LV enlargement and reduced LVEF was noted 4 weeks after I/R in the control group, which, on the other hand, was effectively ameliorated by SW therapy. Echocardiographic study also showed that regional LV wall motion was normalized at the border zone accompanied with increased RMBF and capillary density. These results suggest that SW-induced angiogenesis at the border zone substantially contributes to the suppression of LV remodeling *in vivo*.

### Mechanisms for the inhibitory effects of the SW therapy on LV remodeling after I/R


The precise mechanisms of SW-mediated suppression of LV remodeling after myocardial I/R remain to be fully elucidated. However, it is conceivable that multiple

Fig. 5



The shock wave (SW) therapy increases capillary density in the border zone. Representative factor VIII staining in the control group (a) and the SW group (b), and quantitative analysis of the vessel number in the border zone (c) and in the remote area (d). The SW therapy significantly increased the density of factor VIII-positive capillaries in the border zone, whereas the capillary density was comparable between two groups in the remote area.

Fig. 6



The shock wave (SW) enhances endothelial nitric oxide synthase (eNOS) activity. The SW therapy significantly enhanced eNOS phosphorylation, a marker of eNOS activation, in the border zone (a) and tended to do so for vascular endothelial growth factor (VEGF) protein expression (b) at 1 week after myocardial ischemia-reperfusion.

mechanisms are involved in the inhibitory effects of our SW therapy on LV remodeling after I/R. When a SW hits a tissue, the SW induces cavitation (a micrometer-sized violent collapse of bubbles) by the first compression by the positive pressure component and expansion with the tensile component of SW [24]. As the physical forces generated by cavitation are highly localized, the SW could induce localized stress on cell membranes, leading to a variety of biochemical effects including shear stress,

hyperpolarization, and Ras activation [25], and the induction of stress fibers and intercellular gaps [26]. In addition, the SW induces nonenzymatic NO synthesis from L-arginine and hydrogen peroxide [27], upregulates eNOS, and suppresses nuclear factor-κB activation in the cultured human umbilical venous endothelial cells [28]. NO exerts a wide variety of biological effects including the regulation of vascular tone and angiogenesis [29–31]. In this study, we confirmed that SW therapy increases

eNOS activity and capillary density at the border zone, associated with an improvement of LV remodeling and dysfunction. These results suggest that our SW therapy improves LV remodeling after I/R, at least in part, by enhancing NO production.

Enhanced expression of multiple angiogenic factors, such as VEGF and stromal-derived factor 1, is crucial for the recruitment and incorporation of endothelial progenitor cells [32–34]. We have shown earlier that our SW therapy upregulates myocardial VEGF/Flt-1 expression in pigs *in vivo* [9]. A SW has also been reported to promote mobilization and differentiation of bone marrow-derived cells in a rat model of chronic hindlimb ischemia [35] and in rat bone marrow-derived mononuclear cells *in vitro* [36]. In this study, we confirmed that SW therapy increases RMBF, capillary density, and eNOS activity, and tended to increase the VEGF expression at the border zone. Although the renin–angiotensin system plays an important role in the pathogenesis of LV remodeling after AMI, partly because of enhanced myocardial fibrosis [37–40], the extent of the fibrosis at the border zone was comparable between the two groups in this study.

#### Advantage of the non-invasive SW therapy

The gene [1–3] and cell therapies [4–8], although worthy of development, require invasive procedures such as general anesthesia, cardiac catheterizations, and open chest surgery [4–8,41,42] to deliver the genes or cells to the ischemic myocardium, which may limit the usefulness of these therapies in the clinical setting. Our extracorporeal cardiac SW therapy is quite noninvasive and safe without any adverse effects, which is a major advantage of our SW therapy. This is an important point in determining the clinical usefulness of angiogenic therapies, especially in elderly patients with severe ischemic heart disease.

#### Limitations of the study

Several limitations of this study should be mentioned. First, we observed a trend for but not a significant increase in the VEGF level in the SW group compared with the control group, although we have earlier shown a significant increase in the VEGF expression by SW therapy in a porcine model of chronic myocardial ischemia [9]. This discrepancy may be partly because of the different stage of myocardial ischemia examined. In our earlier study, the VEGF protein level was evaluated 8 weeks after creating a chronic myocardial ischemia, when the VEGF level might have returned to the normal level, and therefore the SW-induced enhancement of the VEGF expression was clearly detected [9]. On the other hand, in this study, the VEGF level was studied 1 week after creating a myocardial I/R, when the expression of VEGF was still strongly enhanced even in the control group. Importantly, however, we were able to show the significant upregulation of eNOS by the SW therapy in

this study, which we did not examine in the earlier study [9]. Second, although we were able to show that our SW therapy enhances angiogenesis at the border zone of the LV, the effects of the SW therapy on each component of the myocardial tissue, including vascular endothelial cells, vascular smooth muscle cells, cardiomyocytes, extracellular matrix, and inflammatory cells, remain to be clarified in future studies. Third, although we studied the expression of eNOS and VEGF in this study, there are many other growth factors and chemokines that could enhance angiogenesis such as stromal-derived factor 1/CXCR4 system and angiopoietin/Tie-2 system. This point also remains to be examined in future studies.

#### Conclusion

We were able to show that our low-energy extracorporeal cardiac SW therapy effectively induces angiogenesis and ameliorates LV remodeling after I/R in pigs *in vivo* without any adverse effects. Thus, our SW therapy could be a novel and safe strategy for the prevention of LV remodeling after AMI in humans.

#### Acknowledgements

The authors thank Dr Ernest H. Marlinghaus, Storz Medical AG, Switzerland, for valuable comments on our study. The authors also thank Fumie Tatebayashi and Naomi Yamaki for their excellent technical assistance. This study was supported in part by the Global COE Program and Grants-in-Aid from the Japanese Ministry of Education, Culture, Sports, Science, and Technology, Tokyo, Japan, and the Japanese Ministry of Health, Labor, and Welfare, Tokyo, Japan.

There is no conflict of interest.

#### References

- Rutanan J, Rissanen TT, Markkanen JE, Gruchala M, Silvennoinen P, Kiveli A, et al. Adenoviral calciferol-mediated intramyocardial gene transfer using the mature form of vascular endothelial growth factor-D induces transmurular angiogenesis in porcine heart. *Circulation* 2004; **109**:1029–1035.
- Kastrup J, Jørgensen E, Rück A, Tagli K, Glogar D, Ruzyllo W, et al. Direct intramyocardial plasmid vascular endothelial growth factor-A165 gene therapy in patients with stable severe angina pectoris. A randomized double-blind placebo-controlled study: the Euroinject One trial. *J Am Coll Cardiol* 2005; **45**:982–988.
- Gupta R, Tongers J, Losordo DW. Human studies of angiogenic gene therapy. *Circ Res* 2009; **105**:724–736.
- van Ramshorst J, Bax JJ, Beeres SL, Dibbets-Schneider P, Roes SD, Stokkel MP, et al. Intramyocardial bone marrow cell injection for chronic myocardial ischemia: a randomized controlled trial. *JAMA* 2009; **301**:1997–2004.
- Wollert KC, Meyer GP, Lotz J, Ringes-Lichtenberg S, Lippolt P, Breidenbach C, et al. Intracoronary autologous bone-marrow cell transfer after myocardial infarction: the BOOST randomised controlled clinical trial. *Lancet* 2004; **364**:141–148.
- Schächinger V, Assmus B, Britten MB, Honold J, Lehmann R, Teupe C, et al. Transplantation of progenitor cells and regeneration enhancement in acute myocardial infarction: final 1-year results of the TOPCARE-AMI Trial. *J Am Coll Cardiol* 2004; **44**:1690–1699.
- Schächinger V, Erbs S, Elsässer A, Haberbosch W, Hambrecht R, Hölschermann H, et al. Improved clinical outcome after intracoronary administration of bone-marrow-derived progenitor cells in acute myocardial

- infarction: final 1-year results of the REPAIR-AMI trial. *Eur Heart J* 2006; **27**:2775–2783.
- 8 Davani S, Deschaseaux F, Chalmers D, Tiberghien P, Kantelip JP. Can stem cells mend a broken heart? *Cardiovasc Res* 2005; **65**:305–316.
  - 9 Nishida T, Shimokawa H, Oi K, Tatewaki H, Uwatoku T, Abe K, *et al.* Extracorporeal cardiac shock wave therapy markedly ameliorates ischemia-induced myocardial dysfunction in pigs in vivo. *Circulation* 2004; **364**:183–192.
  - 10 Fukumoto Y, Ito A, Uwatoku T, Matoba T, Kishi T, Tanaka H, *et al.* Extracorporeal cardiac shock wave therapy ameliorates myocardial ischemia in patients with severe coronary artery disease. *Coron Artery Dis* 2006; **17**:63–70.
  - 11 Kikuchi Y, Ito K, Ito Y, Shiroto T, Tsuburaya R, Aizawa K, *et al.* Double-blind and placebo-controlled study of the effectiveness and safety of extracorporeal cardiac shock wave therapy for severe angina pectoris. *Circ J* 2010; **74**:589–591.
  - 12 Uwatoku T, Ito K, Abe K, Oi K, Hizume T, Sunagawa K, *et al.* Extracorporeal cardiac shock wave therapy improves left ventricular remodeling after acute myocardial infarction in pigs. *Coron Artery Dis* 2007; **18**:397–404.
  - 13 Klein HH, Schubothe M, Nebendahl K, Kreuzer H. Temporal and spatial development of infarcts in porcine hearts. *Basic Res Cardiol* 1984; **79**:440–447.
  - 14 Pich S, Klein HH, Lindert S, Nebendahl K, Kreuzer H. Cell death in ischemic, reperfused porcine hearts: a histochemical and functional study. *Basic Res Cardiol* 1988; **83**:550–559.
  - 15 Shimokawa H, Ito K, Fukumoto Y, Yasuda S. Extracorporeal cardiac shock wave therapy for ischemic heart disease. *Shock Waves* 2008; **17**:449–455.
  - 16 Ito K, Fukumoto Y, Shimokawa H. Extracorporeal shock wave therapy as a new and non-invasive angiogenic therapy. *Tohoku J Exp Med* 2009; **219**:1–9.
  - 17 Versteilen AM, Korstjens IJ, Musters RJ, Groeneveld AB, Sipkema P. Rho kinase regulates renal blood flow by modulating eNOS activity in ischemia-reperfusion of the rat kidney. *Am J Physiol Renal Physiol* 2006; **291**:F606–F611.
  - 18 Oi K, Fukumoto Y, Ito K, Uwatoku T, Abe K, Hizume T, *et al.* Extracorporeal shock wave therapy ameliorates hindlimb ischemia in rabbits. *Tohoku J Exp Med* 2008; **214**:151–158.
  - 19 Kushner FG, Hand M, Smith SC Jr, King SB III, Anderson JL, Antman EM, *et al.* 2009 Focused Updates: ACC/AHA Guidelines for the Management of Patients With ST-Elevation Myocardial Infarction (updating the 2004 Guideline and 2007 Focused Update) and ACC/AHA/SCAI Guidelines on Percutaneous Coronary Intervention (updating the 2005 Guideline and 2007 Focused Update): a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *Circulation* 2009; **120**:2271–2306.
  - 20 Van de Werf F, Bax J, Betriu A, Blomstrom-Lundqvist C, Crea F, Falk V, *et al.* Management of acute myocardial infarction in patients presenting with persistent ST-segment elevation: the Task Force on the Management of ST-Segment Elevation Acute Myocardial Infarction of the European Society of Cardiology. *Eur Heart J* 2008; **29**:2909–2945.
  - 21 Yasuda S, Shimokawa H. Acute myocardial infarction: the enduring challenge for cardiac protection and survival. *Circ J* 2009; **73**:2000–2008.
  - 22 Takii T, Yasuda S, Takahashi J, Ito K, Shiba N, Shirato K, *et al.* Trends in acute myocardial infarction incidence and mortality over 30 years in Japan: report from the MIYAGI-AMI registry study. *Circ J* 2010; **74**:93–100.
  - 23 Jessup M, Brozena S. Heart failure. *N Engl J Med* 2003; **348**:2007–2018.
  - 24 Apfel RE. Acoustic cavitation: a possible consequence of biomedical uses of ultrasound. *Br J Cancer* 1982; **45** (Suppl):140–146.
  - 25 Wang FS, Wang CJ, Huang HJ, Chung H, Chen RF, Yang KD. Physical shock wave mediates membrane hyperpolarization and Ras activation for osteogenesis in human bone marrow stromal cells. *Biochem Biophys Res Commun* 2001; **287**:648–655.
  - 26 Seidl M, Steinbach P, Wörle K, Hofstädter F. Induction of stress fibers and intercellular gaps in human vascular endothelium by shock-waves. *Ultrasonics* 1994; **32**:397–400.
  - 27 Gotte G, Amelio E, Russo S, Marlinghaus E, Musci G, Suzuki H. Short-time non-enzymatic nitric oxide synthesis from L-arginine and hydrogen peroxide induced by shock waves treatment. *FEBS Lett* 2002; **520**:153–155.
  - 28 Mariotto S, Cavaliere E, Amelio E, Ciampa AR, de Prati AC, Marlinghaus E, *et al.* Extracorporeal shock waves: from lithotripsy to anti-inflammatory action by NO production. *Nitric Oxide* 2005; **12**:89–96.
  - 29 Amano K, Matsubara H, Iba O, Okigaki M, Fujiyama S, Imada T, *et al.* Enhancement of ischemia-induced angiogenesis by eNOS overexpression. *Hypertension* 2003; **41**:156–162.
  - 30 Murohara T, Asahara T, Silver M, Bauters C, Masuda H, Kalka C, *et al.* Nitric oxide synthase modulates angiogenesis in response to tissue ischemia. *J Clin Invest* 1998; **101**:2567–2578.
  - 31 Shimokawa H. Primary endothelial dysfunction: atherosclerosis. *J Mol Cell Cardiol* 1999; **31**:23–37.
  - 32 Millauer B, Wizigmann-Voos S, Schnürch H, Martinez R, Möller NP, Risau W, *et al.* High affinity VEGF binding and developmental expression suggest Fk-1 as a major regulator of vasculogenesis and angiogenesis. *Cell* 1993; **72**:835–846.
  - 33 Grunewald M, Avraham I, Dor Y, Bachar-Lustig E, Itin A, Yung S, *et al.* VEGF-induced adult neovascularization: recruitment, retention, and role of accessory cells. *Cell* 2006; **124**:175–189.
  - 34 Askari AT, Unzek S, Popovic ZB, Goldman CK, Forudi F, Kiedrowski M, *et al.* Effect of stromal-cell-derived factor 1 on stem-cell homing and tissue regeneration in ischaemic cardiomyopathy. *Lancet* 2003; **362**:697–703.
  - 35 Aicher A, Heeschen C, Sasaki K, Urbich C, Zeiher AM, Dimmeler S. Low-energy shock wave for enhancing recruitment of endothelial progenitor cells: a new modality to increase efficacy of cell therapy in chronic hind limb ischemia. *Circulation* 2006; **114**:2823–2830.
  - 36 Yip HK, Chang LT, Sun CK, Youssef AA, Sheu JJ, Wang CJ. Shock wave therapy applied to rat bone marrow-derived mononuclear cells enhances formation of cells stained positive for CD31 and vascular endothelial growth factor. *Circ J* 2008; **72**:150–156.
  - 37 Pfeffer MA, McMurray JJ, Velazquez EJ, Rouleau JL, Køber L, Maggioni AP, *et al.* Valsartan, captopril, or both in myocardial infarction complicated by heart failure, left ventricular dysfunction, or both. *N Engl J Med* 2003; **349**:1893–1906.
  - 38 Latini R, Maggioni AP, Flather M, Sleight P, Tognoni G. ACE inhibitor use in patients with myocardial infarction. Summary of evidence from clinical trials. *Circulation* 1995; **92**:3132–3137.
  - 39 Weber KT. Extracellular matrix remodeling in heart failure: a role for de novo angiotensin II generation. *Circulation* 1997; **96**:4065–4082.
  - 40 Ertl G, Frantz S. Healing after myocardial infarction. *Cardiovasc Res* 2005; **66**:22–32.
  - 41 Dohmann HF, Perin EC, Takiya CM, Silva GV, Silva SA, Sousa AL, *et al.* Transendocardial autologous bone marrow mononuclear cell injection in ischemic heart failure: postmortem anatomicopathologic and immunohistochemical findings. *Circulation* 2005; **112**:521–526.
  - 42 Stamm C, Westphal B, Kleine HD, Petzsch M, Kittner C, Klinge H, *et al.* Autologous bone-marrow stem-cell transplantation for myocardial regeneration. *Lancet* 2003; **361**:45–46.

体外衝撃波を用いた非侵襲性血管新生治療

伊藤 健太 下川 宏明

要 旨

我が国では、人口の高齢化や生活習慣の欧米化に伴い、虚血性心疾患や閉塞性動脈硬化症といった動脈硬化性疾患患者が増加してきている。我々は、基礎研究の結果を基に、低出力の衝撃波を用いた血管新生療法（「低出力体外衝撃波治療」）を開発し、①重症狭心症、②急性心筋梗塞、③下肢閉塞性動脈硬化症を対象に臨床試験を行っている。重症狭心症に対しては、第1次臨床試験（オープン試験）と第2次臨床試験（二重盲検プラセボ対照試験）を行い、本治療法の有効性と安全性を確認し、論文報告している。本治療法は、麻酔や侵襲的な処置を伴わずに、体外から治療を行うことができる非侵襲的な治療法であり、繰り返し行うことも可能である。今後幅広い疾患への応用が期待される。

〔日内会誌 99：2846～2852, 2010〕

**Key words**：虚血性心疾患、閉塞性動脈硬化症、血管新生、衝撃波治療

はじめに

我が国では、人口の高齢化や生活習慣の欧米化・糖尿病患者の増加に伴い、虚血性心疾患や閉塞性動脈硬化症といった動脈硬化性疾患の患者数が増加の一途をたどっている。動脈硬化性疾患に対する治療は、生活習慣の改善をベースに、①薬物療法、②カテーテル治療、③バイパス手術の3本柱から成るが、近年これら従来の治療法では十分な治療効果を得られない重症例（例えば、びまん性狭窄病変を持つ症例）が増加してきている。このような症例は、胸痛や下肢痛・潰瘍形成のため生活の質（QOL）が低下するのみならず、予後も不良である。

近年、閉塞性動脈硬化症や虚血性心疾患に対して、遺伝子治療や細胞移植治療が試みられ、日本を含めた世界各国で臨床試験が行われてい

る。これらの治療では、骨髄細胞の採取や遺伝子・細胞の送達のために全身麻酔下での骨髄穿刺や開胸操作といった大きな侵襲を伴う。そのため患者への身体的負担は大きく、また繰り返し行うことは困難である。また、有効性・安全性が認められたとしても、費用が高額になることは避けられない。一方、遺伝子治療や細胞移植治療について、動物実験の結果から期待されたほどの有効性が、臨床試験において認められていないことも指摘されている。そのため、低侵襲で、かつ有効性の期待できる新しい治療法の開発が望まれている。我々は、培養細胞や大型動物（ブタ）を用いた検討の結果を基に、低出力の衝撃波を用いた血管新生療法（「低出力体外衝撃波治療」）を開発し、現在、①重症狭心症、②急性心筋梗塞、③下肢閉塞性動脈硬化症を対象に臨床試験を行っている<sup>1)</sup>。

いとう けんた、しもかわ ひろあき：東北大学大学院循環器病態学分野




図 1. 衝撃波による血管増殖因子の発現亢進：(A) VEGF, (B) Flt-1  
HUVEC に衝撃波を照射すると、VEGF と Flt-1 の発現が亢進した。発現の亢進は、結石破砕に用いる出力の約 10% という弱い出力 (0.09 mJ/mm<sup>2</sup>) の時に最大となった。

## 1. 衝撃波による血管新生作用

衝撃波とは音速を超えて伝わる圧力波で、同じような音響的特性を持つ媒体内を直線的に伝播していくことから、体外で発生させた衝撃波を皮膚表面から脂肪・筋肉などの体組織を通過して伝播させ、体内深部の一点に収束させることができる。衝撃波を用いた治療法としては、尿路結石などの結石破砕治療が確立しており、既に 20 年以上前から標準的治療の 1 つとして保険適応にもなっている。我々は培養細胞を用いた基礎実験により、ヒト臍帯静脈内皮細胞 (Human Umbilical Vein Endothelial Cells : HUVEC) に衝撃波を照射すると、結石破砕に用いる出力の約 10% という弱い出力 (0.09 mJ/mm<sup>2</sup>) をピークに、主要な血管新生因子の 1 つである血管内皮増殖因子 (Vascular Endothelial Growth Factor : VEGF) およびその受容体である Flt-1 の発現が増加することを確認した (図 1)<sup>2)</sup>。そこで、ブタ慢性心筋虚血モデルにおいて体外衝撃波治療の効果を検討した。慢性虚血心筋に 1 日おきに 3 回衝撃波を照射し、4 週間後に評価を行った。その結果、体外衝撃波治療により、虚血心筋組

織における VEGF の発現が遺伝子レベル・蛋白レベルのいずれにおいても亢進していた<sup>2)</sup>。さらに、毛細血管数の増加と冠血流の有意な増加、それに伴う左室壁運動の改善を認めた<sup>2)</sup>。衝撃波治療中および治療後 3 日間のホルター心電図では重篤な不整脈を認めず、突然死も認めなかった。組織学的検討においても、出血などの組織損傷は認めなかった。以上の結果から、低出力の衝撃波を用いた体外衝撃波治療は、安全で有効な血管新生療法であることが示唆された。

## 2. 狭心症に対する低出力体外衝撃波治療

上記の基礎的検討に基づき、我々は、重症狭心症に対して低出力体外衝撃波治療の臨床試験を行っている。対象は以下の 4 条件を満たす狭心症症例とした。① 20 歳以上、② 最大量の薬物治療に抵抗性で、かつ経皮的冠動脈形成術 (percutaneous coronary intervention : PCI) や冠動脈バイパス手術 (coronary artery bypass grafting : CABG) による完全な血行再建が不可能、③ カナダ心臓協会 (Canadian cardiovascular society : CCS) 分類で Class II~IV、④ 負荷心筋シンチグラムや負荷心エコーで明らかな虚血領域





図2. 体外衝撃波治療装置と治療風景：(A) 体外衝撃波治療装置，(B) 治療風景，(C) 心臓超音波検査画面，(D) 概念図

を認める。治療には、スイスのメーカーと共同開発した心臓病治療専用の衝撃波治療装置を用いている(図2)。衝撃波発生ヘッドを患者の前胸壁に当て、装置に内蔵された超音波プローブで心臓を観察しながら、虚血部位に照準を合わせ、衝撃波を照射する。1カ所につき200発の衝撃波を、虚血領域の広さに応じて20~120カ所照射する。1回の治療時間は約3時間で、1~2日おきに計3回治療を行う。痛みや苦痛を伴わないため、麻酔や鎮静薬の投与は必要ない。2003年から重症狭心症患者9名を対象に実施した第1次臨床試験では、個人差はあるものの全例で狭心症症状が軽減し、ニトログリセリンの使用量が激減するなどの効果を認め、その効果は1年以上にわたって持続した(図3)<sup>3)</sup>。また、負荷心筋シンチグラムで評価した心筋血流も、衝撃波を照射した部位においてのみ改善を認めた(図3)<sup>3)</sup>。この結果から、衝撃波を照射した部位のみ

で血管新生が生じ、心筋灌流が改善したと考えられた。一方、治療に伴う合併症や副作用は全く認めなかった。さらに、2005年から、低出力体外衝撃波治療とプラセボ治療を比較する第2次臨床試験を実施し、その結果、低出力体外衝撃波治療後には、CCS分類による狭心症の重症度、ニトログリセリンの使用頻度、6分間歩行距離が有意に改善し、MRIで測定した左室一回拍出量、左室駆出率も有意に増加した(図4)<sup>4)</sup>。これらの効果はプラセボ治療後では認められなかった。本年7月、狭心症に対する低出力体外衝撃波治療は、厚生労働省の高度医療に承認された。

### 3. 急性心筋梗塞に対する低出力体外衝撃波治療

我が国では、急性心筋梗塞の発症早期にPCIによる再灌流療法が行われ、心筋梗塞発症早期



(文献3より改変)

図3. 狭心症に対する第1次臨床試験の結果：(A) CCS分類による狭心症の重症度，(B) ニトログリセリンの使用頻度，(C) 負荷心筋シンチグラムによる心筋血流（青が虚血，黄～赤が血流良好を意味する）自覚症状の改善を認めた（A, B）。また，衝撃波治療を受けた領域（破線で示された領域）でのみ血流の改善を認めた（C）。

の死亡率低下に貢献している。しかし十分な薬物治療を受けていても，慢性期には左室リモデリング（心拡大と収縮能低下）が進行し，重篤な心不全へ進行する例が少なくない。再灌流療法成功後も梗塞巣周囲では組織の浮腫や炎症により微小循環障害が遷延しており，これが左室リモデリング進行の一因と考えられている。そこで我々は，ブタ急性心筋梗塞モデルを用いて，低出力体外衝撃波治療が，慢性期の左室リモデ

リングを抑制するか検討した。急性心筋梗塞作成急性期に梗塞境界領域に対して低出力の衝撃波を照射し，4週間後に評価を行ったところ，左室リモデリングが軽減されることが確認された<sup>5,6)</sup>。さらに，梗塞巣周囲の毛細血管数の増加していたことから，急性期の低出力体外衝撃波治療により境界領域での血管新生が促進されて梗塞サイズの拡大が抑制されたと考えられた。この結果を基に，2007年から急性心筋梗塞に対




図4. 狭心症に対する第2次臨床試験の結果：(A) CCS分類による狭心症の重症度，(B) ニトログリセリンの使用頻度，(C) 6分間歩行距離，(D) MRIで測定した左室一回拍出量，(E) MRIで測定した左室駆出率，(F) MRIで測定した左室拡張末期容積

自覚症状の改善 (A, B)，運動耐用能の改善 (C)，心機能の改善 (D, E) を認めた。

する体外衝撃波治療の臨床試験を行っている。この試験では、PCIによる再灌流療法が成功した症例に対して、心筋梗塞発症72時間以内に衝撃波治療を開始し、1~2日おきに計3回行い、慢性期の左室リモデリングや心不全発症の抑制効果を評価する。

#### 4. 下肢閉塞性動脈硬化症に対する低出力体外衝撃波治療

我々は、ウサギの大腿動脈~膝窩動脈を摘出することにより下肢虚血を作成し、下肢虚血作

成1週間後に低出力体外衝撃波治療を行った。治療3週間後に評価を行った結果、対照群に比して、低出力体外衝撃波治療群では虚血肢における側副血行路の発達が促進され、血管造影上の血流スコアや下肢血圧の改善効果を認めた(図5)<sup>7)</sup>。これらの結果を基に、2007年から、間歇性跛行症状を有する慢性閉塞性動脈硬化症症例を対象に、低出力体外衝撃波治療の臨床試験を行っている。




図5. 下肢虚血に対する体外衝撃波治療の効果

体外衝撃波治療群（下段）では、対照群（上段）に比して、虚血肢における側副血行路の発達が促進された。

## おわりに

我々が開発している低出力体外衝撃波治療は極めて低侵襲であり、血管新生を促進することにより組織灌流を改善することが期待される新しい治療法である。本治療法の着想や至適治療条件の決定、動物実験および臨床試験などの先駆的な研究は、すべて我々がトップランナーとして、世界で最初に論文発表を行ってきた。狭心症に対する低出力体外衝撃波治療については、欧州などで医療機器として認可され、既に世界10カ国以上で500名以上の狭心症症例の治療に用いられている。我が国においては、厚生労働省の高度医療評価制度へ申請中である。近年、整形外科領域でも肘や肩、足底等の疼痛治療や難治性の骨折の治療にも低出力の衝撃波を用い

た治療の有効性が報告されており、今後、幅広い疾患への応用が期待される。

本治療法の開発は、厚生労働科学研究費補助金を得て行っており、平成21年度には、研究課題「低侵襲性体外衝撃波治療法の実用化を目指したエビデンス確立のための拠点形成」が厚生労働省の先端医療開発特区（スーパー特区）に採択された。現在、幅広い疾患への適応拡大を目指して、様々な専門領域の研究者との共同研究による基礎研究および臨床試験を進めている。治療法の詳細並びに最新情報に関しては、以下のホームページを参照されたい (<http://www.cardio.med.tohoku.ac.jp/shockwave/index.html>)。

## 文 献

- 1) Ito K, et al: Extracorporeal shock wave therapy as a new