

- Mishra L, Shetty K, Tang Y, Stuart A, Byers SW. (2005). The role of TGF- β and Wnt signaling in gastrointestinal stem cells and cancer. *Oncogene* **24**: 5775–5789.
- Miyazawa K, Shinozaki M, Hara T, Furuya T, Miyazono K. (2002). Two major Smad pathways in TGF- β superfamily signalling. *Genes Cells* **7**: 1191–1204.
- Mizoi T, Ohtani H, Miyazono K, Miyazawa M, Matsuno S, Nagura H. (1993). Immunoelectron microscopic localization of transforming growth factor β 1 and latent transforming growth factor β 1 binding protein in human gastrointestinal carcinomas: qualitative difference between cancer cells and stromal cells. *Cancer Res* **53**: 183–190.
- Naka K, Hoshii T, Muraguchi T, Tadokoro Y, Ooshio T, Kondo Y *et al*. (2010). TGF- β -FOXO signalling maintains leukaemia-initiating cells in chronic myeloid leukaemia. *Nature* **463**: 676–680.
- Nishii T, Yashiro M, Shinto O, Sawada T, Ohira M, Hirakawa K. (2009). Cancer stem cell-like SP cells have a high adhesion ability to the peritoneum in gastric carcinoma. *Cancer Sci* **100**: 1397–1402.
- Patrawala L, Calhoun T, Schneider-Broussard R, Zhou J, Claypool K, Tang DG. (2005). Side population is enriched in tumorigenic, stem-like cancer cells, whereas ABCG2+ and ABCG2– cancer cells are similarly tumorigenic. *Cancer Res* **65**: 6207–6219.
- Peñuelas S, Anido J, Prieto-Sánchez RM, Folch G, Barba I, Cuatrecasas J *et al*. (2009). TGF- β increases glioma-initiating cell self-renewal through the induction of LIF in human glioblastoma. *Cancer Cell* **15**: 315–327.
- Piccirillo SG, Reynolds BA, Zanetti N, Lamorte G, Binda E, Broggi G *et al*. (2006). Bone morphogenetic proteins inhibit the tumorigenic potential of human brain tumour-initiating cells. *Nature* **444**: 761–765.
- Rabindran SK, Ross DD, Doyle LA, Yang W, Greenberger LM. (2000). Fumitremorgin C reverses multidrug resistance in cells transfected with the breast cancer resistance protein. *Cancer Res* **60**: 47–50.
- Scharenberg CW, Harkey MA, Torok-Storb B. (2002). The ABCG2 transporter is an efficient Hoechst 33342 efflux pump and is preferentially expressed by immature human hematopoietic progenitors. *Blood* **99**: 507–512.
- Sneddon JB, Zhen HH, Montgomery K, van de Rijn M, Tward AD, West R *et al*. (2006). Bone morphogenetic protein antagonist gremlin 1 is widely expressed by cancer-associated stromal cells and can promote tumor cell proliferation. *Proc Natl Acad Sci USA* **103**: 14842–14847.
- Stingl J, Caldas C. (2007). Molecular heterogeneity of breast carcinomas and the cancer stem cell hypothesis. *Nat Rev Cancer* **7**: 791–799.
- Takaishi S, Okumura T, Tu S, Wang SS, Shibata W, Vigneshwaran R *et al*. (2009). Identification of gastric cancer stem cells using the cell surface marker CD44. *Stem Cells* **27**: 1006–1020.
- Takemura S, Yashiro M, Sunami T, Tendo M, Hirakawa K. (2004). Novel models for human scirrhous gastric carcinoma *in vivo*. *Cancer Sci* **95**: 893–900.
- Vagenas K, Spyropoulos C, Gavala V, Tsamandas AC. (2007). TGF β 1, TGF β 2, and TGF β 3 protein expression in gastric carcinomas: correlation with prognostic factors and patient survival. *J Surg Res* **139**: 182–188.
- Velamakanni S, Wei SL, Janvilisri T, van Veen HW. (2007). ABCG transporters: structure, substrate specificities and physiological roles: a brief overview. *J Bioenerg Biomembr* **39**: 465–471.
- Visvader JE, Lindeman GJ. (2008). Cancer stem cells in solid tumours: accumulating evidence and unresolved questions. *Nat Rev Cancer* **8**: 755–768.
- Wakefield LM, Roberts AB. (2002). TGF- β signaling: positive and negative effects on tumorigenesis. *Curr Opin Genet Dev* **12**: 22–29.
- Weinberg RA. (1995). The retinoblastoma protein and cell cycle control. *Cell* **81**: 323–330.
- Wu C, Alman BA. (2008). Side population cells in human cancers. *Cancer Lett* **268**: 1–9.
- Yanagihara K, Kamada N, Tsumuraya M, Amano F. (1993). Establishment and characterization of a human gastric scirrhous carcinoma cell line in serum-free chemically defined medium. *Int J Cancer* **54**: 200–207.
- Yashiro M, Chung YS, Sowa M. (1994). Role of orthotopic fibroblasts in the development of scirrhous gastric carcinoma. *Jpn J Cancer Res* **85**: 883–886.
- Yashiro M, Chung YS, Nishimura S, Inoue T, Sowa M. (1996). Peritoneal metastatic model for human scirrhous gastric carcinoma in nude mice. *Clin Exp Metastasis* **14**: 43–54.
- Yashiro M, Yasuda K, Nishii T, Kaizaki R, Sawada T, Ohira M *et al*. (2009). Epigenetic regulation of the embryonic oncogene ERas in gastric cancer cells. *Int J Oncol* **35**: 997–1003.
- Yin L, Castagnino P, Assoian RK. (2008). ABCG2 expression and side population abundance regulated by a transforming growth factor β -directed epithelial-mesenchymal transition. *Cancer Res* **68**: 800–807.
- Zhou S, Schuetz JD, Bunting KD, Colapietro AM, Sampath J, Morris JJ *et al*. (2001). The ABC transporter Bcrp1/ABCG2 is expressed in a wide variety of stem cells and is a molecular determinant of the side-population phenotype. *Nat Med* **7**: 1028–1034.

Supplementary Information accompanies the paper on the Oncogene website (<http://www.nature.com/onc>)

