

**Table 2** Published reports comparing complication rates of EMR and ESD

Author	Site	EMR				ESD					
		Number of lesions	Removed en bloc %	Bleeding %	Perforation %	Recurrence %	Number of lesions	Removed en bloc %	Bleeding %	Perforation %	Recurrence %
Saito <sup>26</sup>	Colon	228	84.0	3.1 (minor)	1.3	14.5	145	33.0	1.4 (minor)	6.2	2.1
	Gastric	411	56.0	0.2	1.2	6.6	303	92.7	0.0 (major)	3.6	2.0
Shimura <sup>27</sup>	Gastric	48	31.3	12.5	0.0	35.4	59	88.1	13.6	3.4	1.7
	Gastric	125	63.6 (>10 mm)	1.8	3.2	5.6	120	91.3 (>10 mm)	0.0 (major)	4.2	2.5
Ishihara <sup>28</sup>	Oesophagus	52	10.9 (out of 46 lesions)	0.0	0.0	22.0	33	90.6 (out of 32 lesions)	0.0	0.0	3.1
	All	2987	57.7	5.8	1.0	5.2	1804	94.6	9.2	4.5	0.3

\*Meta-analysis of 15 studies of EMR and ESD.  
EMR, endoscopic mucosal resection; ESD, endoscopic submucosal dissection.

**Table 3** Indications for endoscopic resection of early gastrointestinal neoplasm

Lesion Position	Indication
Esophagus	Well- or moderately differentiated m1 or m2 SCC or AC < 20 mm, without venous or lymphatic involvement; less than a third of the circumference involved (to avoid risk of post-resection stricture formation)
Stomach	ER for Barrett's esophagus is still being studied Standard criteria: Well- or moderately differentiated AC and/or papillary carcinoma; cancer confined to mucosa IIa < 20 mm; cancer confined to the mucosa IIb, IIc < 10 mm, without evidence of lymphatic involvement Expanded criteria: Mucosal well/moderately differentiated AC, irrespective of size, without ulceration; ≤ 30 mm with ulceration; if minute submucosal invasion is found then the size of the lesion is ≤ 30 mm, without venous or lymphatic involvement; mucosal undifferentiated AC ≤ 20 mm, without lymphovascular involvement or ulceration
Colorectum	Laterally spreading tumors High-grade dysplasia The indication for resection of mucosal or AC invading slightly into the SM is still being studied. ESD has been reported for resection of: - well- or moderately differentiated AC; cancer confined to the mucosa: IIa < 20 mm, IIb, IIc < 10 mm, without evidence of venous or lymphatic involvement - superficially invading the SM (< 500 µm from the muscularis mucosa); without venous or lymphatic involvement

AC, adenocarcinoma; ER, endoscopic resection; IIa, slightly elevated superficial tumor; IIb, flat superficial tumor; IIc, slightly depressed superficial tumor; m, mucosal; SCC, squamous cell carcinoma; SM, submucosa.

Gastrectomy with regional lymph node dissection was formerly the only available curative treatment for early gastric cancer. In 1996, the National Cancer Center Hospital (Tokyo) published their data describing over 1000 patients with intramucosal early gastric cancer who underwent surgical resection. This study provided some of the first evidence to suggest that radical surgery with lymphadenectomy was unnecessary for certain gastric cancers due to the extremely low incidence of spread to lymph nodes.<sup>43</sup> Curative endoscopic resection of early intramucosal gastric cancers has since become a valid therapeutic option, but until recently was restricted to small lesions less than 2 cm in size with no evidence of surface ulceration. Although other publications suggested that certain lesions invading into the submucosa also carried a low risk of progression, these studies were limited by small patient cohorts.<sup>44-46</sup>

Gotoda and colleagues published extensive data in 2000 that provided a more robust evidence base for the expansion of endoscopic resection criteria. They examined the presence of lymph node metastasis in 5265 patients who underwent gastrectomy with

**Table 4** Early gastric cancer with no risk of lymph node metastasis

Tumor characteristics	Number of cases	95% confidence interval
Intramucosal	1230	0–0.3%
Well-/moderately differentiated		
No lymphovascular invasion		
Irrespective of ulcer findings	929	0–0.4%
Tumor less than 3 cm in size		
Intramucosal		
Well-/moderately differentiated	141	0–2.6%
No lymphovascular invasion		
No ulcer		
Irrespective of tumor size	145	0–2.5%
Intramucosal		
Poorly differentiated		
No lymphovascular invasion	145	0–2.5%
No ulcer		
Tumor less than 2 cm in size		
Minute submucosal penetration (SM1)	145	0–2.5%
Well-/moderately differentiated		
No lymphovascular invasion		
Tumor less than 3 cm in size		

lymph node dissection for early gastric cancer from two centers. Only 2.2% (65/3016) of intramucosal cancers were associated with regional lymph node metastasis. Of these lesions, lymph node metastasis was associated with poor differentiation, signet ring histology, lymphovascular invasion and lesions greater than 3 cm with surface ulceration. Specifically, intramucosal lesions without ulceration did not demonstrate lymph node metastasis irrespective of size. Gotoda *et al.* also showed that 18% of cancers with deeper invasion into the submucosal layer were associated with lymph node metastasis. However, lesions less than 3 cm in size with submucosal invasion less than 500  $\mu$ m, well- or moderately differentiated histology and no evidence of lymphovascular involvement demonstrated no lymph node metastasis. Table 4 summarizes data from this study, showing the lesion types that displayed no evidence of lymph node metastasis.<sup>47</sup>

In 2004, the Japanese Gastric Cancer Association issued expanded criteria for the treatment of early gastric cancer based on this study.<sup>48</sup> Hirasawa and colleagues have since explored undifferentiated early gastric cancers in a similar population of 3843 Japanese patients. Undifferentiated lesions confined to the mucosa, less than 20 mm in diameter, without lymphovascular involvement or ulcer presence showed no lymph node metastasis. They proposed that endoscopic resection should also be considered for these lesions, thus further expanding the criteria for endoscopic management of gastric cancer.<sup>49</sup> Other studies of the risk of lymph node metastasis in poorly differentiated lesions have produced similar results, although they involved smaller patient numbers.<sup>50–53</sup>

## Early lesions of the colorectum

Worldwide, colorectal cancer incidence ranks fourth in frequency in men and third in women. Despite a relatively good prognosis, rates of colorectal cancer are rising rapidly in countries such as

Japan where the risk was previously low.<sup>50</sup> Important work done in the 1980s demonstrated that specific genetic alterations occurred in adenomas and carcinomas, suggesting that colorectal cancer development involved mutational activation of an oncogene and loss of tumor suppressor genes. This evidence led to the development of a genetic model for colorectal tumorigenesis, and to the suggestion that most carcinomas arise from benign adenomatous precursors.<sup>54</sup> In contrast, a proportion of colorectal cancers appear to arise from normal mucosa and do not follow the adenoma–carcinoma sequence. These *de novo* carcinomas tend to be small, depressed-type lesions and may have an increased invasive tendency.<sup>55,56</sup> Originally, depressed-type colorectal neoplasms were thought to exist only in Eastern populations, but their existence and invasive potential in the West have since been proven by groups from the UK and the USA.<sup>57,58</sup>

Intramucosal colorectal lesions have no risk of lymph node metastasis and can be cured by endoscopic resection.<sup>59</sup> Once the submucosa has been breached, the incidence of lymphatic spread rises to around 10%, but this is dependent on depth of invasion. Lesions with submucosal invasion less than 1000  $\mu$ m have a low risk of lymph node metastasis and are good candidates for endoscopic therapy.<sup>6</sup> Kitajima *et al.* reported an overall incidence of lymph node metastasis in 865 submucosal invasive colorectal cancers of 10%. Poor differentiation, lymphatic invasion and venous invasion were significant risk factors for metastasis. They showed that pedunculated lesions with submucosal invasion less than 3000  $\mu$ m and no evidence of lymphatic invasion displayed no evidence of lymph node metastasis. All sessile cancers with lymph node metastasis had invaded the submucosal layer by more than 1000  $\mu$ m.<sup>60</sup>

Egashira and colleagues demonstrated a similar rate of lymph node metastasis of 9%, and identified submucosal invasion greater than 2000  $\mu$ m as an independent risk factor. Their study was smaller, involving only 140 cancers, and cases were not subdivided into pedunculated and non-pedunculated.<sup>61</sup> With regard to pedunculated lesions, Haggitt identified stalk invasion as an important factor in predicting clinical outcome. Tumors extending beyond the stalk into the submucosa, but not reaching the muscularis propria (Haggitt level 4) were associated with poor outcome. This study was limited by moderate patient numbers ( $n = 129$ ), a factor that should be taken into consideration in practical application.<sup>62</sup>

Special consideration should be given to LST of the colorectum. Uraoka *et al.* studied 511 colorectal LST and reported significant differences in depth of invasion between granular and non-granular lesions. LST-NG had a higher potential for malignancy compared to LST-G with frequency of submucosal invasion of 14% versus 7%. Whilst piecemeal resection was considered acceptable for LST-G type, en bloc resection was suggested as the best therapeutic approach for LST-NG type.<sup>63</sup>

The therapeutic approach to lesions of the colorectum is very much dependent on the accuracy of endoscopic diagnosis. Matsuda *et al.* recently carried out a large prospective study of 4215 lesions in 3029 consecutive patients between 1998 and 2005 at the National Cancer Center Hospital, Tokyo. All lesions were detected via the conventional endoscopic view and assessed using magnifying chromoendoscopy for evidence of invasive features according to pit pattern evaluation. They showed that 99.4% of lesions diagnosed endoscopically as 'non-invasive' were adenoma, high-grade dysplasia or adenocarcinoma with submucosal inva-

sion less than 1000  $\mu\text{m}$ . Among lesions diagnosed with 'invasive' pattern, 87% were cancers with submucosal invasion deeper than 1000  $\mu\text{m}$ . This is the first large-scale prospective study to validate the use of magnifying chromoendoscopy as a highly effective method in the prediction of invasion depth of colorectal neoplasms.<sup>64</sup>

## Application of ESD in countries other than Japan

ESD is an appealing prospect for treatment of certain lesions of the GIT in the West, such as superficial carcinomas of the esophagus, high-grade dysplasia in Barrett's mucosa and large flat non-granular tumors of the colorectum. There are, however, a number of limitations to widespread use of ESD outside Japan.

Firstly, selection of appropriate lesions for ESD is crucial, and the diagnostic skills to facilitate this, including determination of lesion characteristics, are of great importance. Whilst optical magnification is used in Japan allowing up to 150 $\times$  image enlargement, digital magnification is more commonly available in the West, providing views with less resolution. Chromoendoscopy is also a routine modality in GI lesion assessment in Japan, but rarely used outside specialist units in the West. Consequently, the ability to analyze lesion surface vascularity and pit pattern in detail and therefore lesion selection for ESD is limited. These assessment techniques are considered crucial in Japan to enable correct diagnosis of lesion type, depth and amenability to endoscopic treatment. Successful application of ESD in the West will certainly require a change in diagnostic technique and close reference to Japanese literature in selection of lesions for resection.

Secondly, ESD is a technically demanding procedure requiring a high level of endoscopic skill and intensive training. The learning curve is steep and involves animal model work in the first instance. Unlike Western countries, facilities for animal model training are readily available in Japan and materials such as the isolated pig stomach can be supplied at low cost. Initial ESD training in patients entails removal of small gastric lesions in the antrum under close expert supervision, and generally, at least 30 procedures are required to reach basic proficiency.<sup>65</sup> The likelihood of major complications for ESD of lesions in this position is low, even for endoscopists with less experience. The large lumen allows easy maneuvering and the risk of perforation is reduced due to the relative thickness of the gastric wall. Bleeding is common during ESD and safe hemostasis is one of the most important aspects of the procedure. However, acquiring skills for basic ESD maneuvers from the beginning of training is vital and the lower vascularity of the antral wall allows this due to reduced bleeding risk.

The incidence of early gastric cancer in the West is very low compared to Japan, so opportunities to perform training gastric ESD are few. Alternatively, rectal ESD is a comparatively safe procedure and may provide a useful training medium for Western endoscopists. Certain skills can be acquired during animal model training, but collaboration with expert Japanese endoscopists and training periods in their units may be helpful in order to reach the necessary skill level. Suzuki *et al.* recently reported their early experience of ESD as a modality to remove large sessile colorectal polyps at the Wolfson Endoscopy Unit, UK. Although only nine patients were enrolled in the study, en bloc resection was achieved

in seven patients, with only one major complication of post-procedural bleeding requiring blood transfusion. Importantly, the ESD technique was acquired under the supervision of an expert.<sup>66</sup> Dinis-Ribeiro *et al.* published a case series of 19 gastric ESD from Portugal reporting only one hemorrhage and no perforations.<sup>67</sup>

Thirdly, ESD is considered more economical and less invasive compared to surgery. Nevertheless, mean hospital inpatient stay for ESD is 5 days and this could prove logistically difficult in the West where bed availability is often limited. In addition, it could be argued that laparoscopic surgery and transanal resection for colorectal lesions in the West are more established techniques, requiring a shorter or similar length inpatient stay; thus, they may be a more viable option.

Finally, management of GIT lesions using ESD in the West will undoubtedly require a multidisciplinary team. During each procedure, several endoscopists are often present in Japan, either to assist or monitor patients, and propofol is frequently given without anesthetists being present. However, although conscious sedation is standard practice in the UK, anesthetists would be required to administer propofol.<sup>68</sup> Practice varies worldwide, with anesthetist- or nurse-administered propofol common in Australia and the USA.<sup>69</sup> Endoscopy nurse training would also need to be addressed in the West, as ESD requires highly trained assistants as well as skilled technicians. Introduction of ESD into Western countries could be of huge benefit to the management of GIT lesions. However, close and supportive working relationships between endoscopists, pathologists and surgeons would be vital for it to succeed as a viable therapeutic option.

## Acknowledgments

A. Conlin was awarded a travel scholarship from HCA International Foundation to fund training at the National Cancer Centre Hospital, Tokyo, Japan. T. Kaltenbach received the American College of Gastroenterology 2009 North American International Gastrointestinal Training Grant for externship at the National Cancer Center Hospital, Tokyo, Japan.

## Disclosure statement

The authors report no conflicts of interest in this work.

## References

- Itoh H, Oohata Y, Nakamura K *et al.* Complete ten-year postgastrectomy follow-up of early gastric cancer. *Am. J. Surg.* 1989; **158**: 14–16.
- Sano T, Sasako M, Kinoshita T, Maruyama K. Recurrence of early gastric cancer. Follow-up of 1475 patients and review of the Japanese literature. *Cancer* 1993; **72**: 3174–8.
- Soetikno R, Kaltenbach T, Yeh R, Gotoda T. Endoscopic mucosal resection for early cancers of the upper gastrointestinal tract. *J. Clin. Oncol.* 2005; **23**: 4490–8.
- Gotoda T, Kondo H, Ono H *et al.* A new endoscopic mucosal resection procedure using an insulation-tipped electrosurgical knife for rectal flat lesions: report of two cases. *Gastrointest. Endosc.* 1999; **50**: 560–3.
- Japanese Gastric Cancer Association. Japanese classification of gastric carcinoma—2nd English edition. *Gastric. Cancer* 1998; **1**: 10–24.

- 6 Dixon MF. Gastrointestinal epithelial neoplasia: Vienna revisited. *Gut* 2002; 51: 130–1.
- 7 Schlemper RJ, Riddell RH, Kato Y et al. The Vienna classification of gastrointestinal epithelial neoplasia. *Gut* 2000; 47: 251–5.
- 8 The Paris endoscopic classification of superficial neoplastic lesions: esophagus, stomach, and colon: November 30 to December 1, 2002. *Gastrointest. Endosc.* 2003; 58 (Suppl. 6): S3–43.
- 9 Kiesslich R, Jung M. Magnification endoscopy: does it improve mucosal surface analysis for the diagnosis of gastrointestinal neoplasias? *Endoscopy* 2002; 34: 819–22.
- 10 Kida M, Kobayashi K, Saigenji K. Routine chromoendoscopy for gastrointestinal diseases: indications revised. *Endoscopy* 2003; 35: 590–6.
- 11 Kudo S, Hirota S, Nakajima T et al. Colorectal tumours and pit pattern. *J. Clin. Pathol.* 1994; 47: 880–5.
- 12 Kudo S, Rubio CA, Teixeira CR, Kashida H, Kogure E. Pit pattern in colorectal neoplasia: endoscopic magnifying view. *Endoscopy* 2001; 33: 367–73.
- 13 Inoue H, Takeshita K, Hori H, Muraoka Y, Yoneshima H, Endo M. Endoscopic mucosal resection with a cap-fitted panendoscope for esophagus, stomach, and colon mucosal lesions. *Gastrointest. Endosc.* 1993; 39: 58–62.
- 14 Soetniko RM, Gotoda T, Nakanishi Y, Soehendra N. Endoscopic mucosal resection. *Gastrointest. Endosc.* 2003; 57: 567–79.
- 15 Tanabe S, Koizumi W, Kokutou M et al. Usefulness of endoscopic aspiration mucosectomy as compared with strip biopsy for the treatment of gastric mucosal cancer. *Gastrointest. Endosc.* 1999; 50: 819–22.
- 16 Akiyama M, Ota M, Nakajima H, Yamagata K, Munakata A. Endoscopic mucosal resection of gastric neoplasms using a ligating device. *Gastrointest. Endosc.* 1997; 45: 182–6.
- 17 Uno Y, Munakata A. The non-lifting sign of invasive colon cancer. *Gastrointest. Endosc.* 1994; 40: 485–9.
- 18 Kobayashi N, Saito Y, Sano Y et al. Determining the treatment strategy for colorectal neoplastic lesions: endoscopic assessment or the non-lifting sign for diagnosing invasion depth? *Endoscopy* 2007; 39: 701–5.
- 19 Saito Y, Uraoka T, Matsuda T et al. Endoscopic treatment of large superficial colorectal tumors: a case series of 200 endoscopic submucosal dissections (with video). *Gastrointest. Endosc.* 2007; 66: 966–73.
- 20 Saito Y, Uraoka T, Matsuda T et al. A pilot study to assess the safety and efficacy of carbon dioxide insufflation during colorectal endoscopic submucosal dissection with the patient under conscious sedation. *Gastrointest. Endosc.* 2007; 65: 537–42.
- 21 Uraoka T, Saito Y, Yamamoto K, Fujii T. Submucosal injection solution for gastrointestinal tract endoscopic mucosal resection and endoscopic submucosal dissection. *Drug Dev. Devel. Ther.* 2008; 2: 131–8.
- 22 Ishihara R, Iishi H, Takeuchi Y et al. Local recurrence of large squamous-cell carcinoma of the esophagus after endoscopic resection. *Gastrointest. Endosc.* 2008; 67: 799–804.
- 23 Hotta K, Fujii T, Saito Y, Matsuda T. Local recurrence after endoscopic resection of colorectal tumors. *Int. J. Colorectal. Dis.* 2009; 24: 225–30.
- 24 Kaltenbach T, Friedland S, Maheshwari A et al. Short- and long-term outcomes of standardized EMR of nonpolypoid (flat and depressed) colorectal lesions  $\geq$  or = 1 cm (with video). *Gastrointest. Endosc.* 2007; 65: 857–65.
- 25 Saito Y, Fukuzawa M, Matsuda T et al. Clinical outcome of endoscopic submucosal dissection versus endoscopic mucosal resection of large colorectal tumors as determined by curative resection. *Surg. Endosc.* 2009; 24: 343–52.
- 26 Oda I, Saito D, Tada M et al. A multicenter retrospective study of endoscopic resection for early gastric cancer. *Gastric. Cancer* 2006; 9: 262–70.
- 27 Shimura T, Sasaki M, Kataoka H et al. Advantages of endoscopic submucosal dissection over conventional endoscopic mucosal resection. *J. Gastroenterol. Hepatol.* 2007; 22: 821–6.
- 28 Watanabe K, Ogata S, Kawazoe S et al. Clinical outcomes of EMR for gastric tumors: historical pilot evaluation between endoscopic submucosal dissection and conventional mucosal resection. *Gastrointest. Endosc.* 2006; 63: 776–82.
- 29 Cao Y, Liao C, Tan A, Gao Y, Mo Z, Gao F. Meta-analysis of endoscopic submucosal dissection versus endoscopic mucosal resection for tumors of the gastrointestinal tract. *Endoscopy* 2009; 41: 751–7.
- 30 Parkin DM, Bray F, Ferlay J, Pisani P. Global cancer statistics. *CA Cancer J. Clin.* 2002; 55: 74–108.
- 31 Bousamra M Jr, Haasler GB, Parviz M. A decade of experience with transorally and transhiatal esophagectomy. *Am. J. Surg.* 2002; 183: 162–7.
- 32 Pech O, Gossner L, May A et al. Long-term results of photodynamic therapy with 5-aminolevulinic acid for superficial Barrett's cancer and high-grade intraepithelial neoplasia. *Gastrointest. Endosc.* 2005; 62: 24–30.
- 33 Ishihara R, Iishi H, Uedo N et al. Comparison of EMR and endoscopic submucosal dissection for en bloc resection of early esophageal cancers in Japan. *Gastrointest. Endosc.* 2008; 68: 1066–72.
- 34 Jemal A, Murray T, Ward E et al. Cancer statistics, 2005. *CA Cancer J. Clin.* 2005; 55: 10–30.
- 35 Sharma P, Sidorenko EI. Are screening and surveillance for Barrett's oesophagus really worthwhile? *Gut* 2005; 54 (Suppl. 1): i27–32.
- 36 Nigro JJ, Hagen JA, DeMeester TR et al. Prevalence and location of nodal metastases in distal esophageal adenocarcinoma confined to the wall: implications for therapy. *J. Thorac. Cardiovasc. Surg.* 1999; 117: 16–23. discussion 23–5.
- 37 Westertep M, Koppert LB, Buskens CJ et al. Outcome of surgical treatment for early adenocarcinoma of the esophagus or gastro-esophageal junction. *Virchows. Arch.* 2005; 446: 497–504.
- 38 Stein HJ, Feith M, Bruecher BL, Naeherig J, Sarbia M, Siewert JR. Early esophageal cancer: pattern of lymphatic spread and prognostic factors for long-term survival after surgical resection. *Ann. Surg.* 2005; 242: 566–73; discussion 573–5.
- 39 Ishihara R, Tanaka H, Iishi H et al. Long-term outcome of esophageal mucosal squamous cell carcinoma without lymphovascular involvement after endoscopic resection. *Cancer* 2008; 112: 2166–72.
- 40 Tajima Y, Nakanishi Y, Ochiai A et al. Histopathologic findings predicting lymph node metastasis and prognosis of patients with superficial esophageal carcinoma: analysis of 240 surgically resected tumors. *Cancer* 2000; 88: 1285–93.
- 41 Katada C, Muto M, Manabe T, Ohtsu A, Yoshida S. Local recurrence of squamous-cell carcinoma of the esophagus after EMR. *Gastrointest. Endosc.* 2005; 61: 219–25.
- 42 Inoue M, Sugane S. Epidemiology of gastric cancer in Japan. *Postgrad. Med. J.* 2005; 81: 419–24.
- 43 Yamao T, Shirao K, Ono H et al. Risk factors for lymph node metastasis from intramucosal gastric carcinoma. *Cancer* 1996; 77: 602–6.
- 44 Yasuda K, Shiraiishi N, Suematsu T, Yamaguchi K, Adachi Y, Kitano S. Rate of detection of lymph node metastasis is correlated with the depth of submucosal invasion in early stage gastric carcinoma. *Cancer* 1999; 85: 2119–23.
- 45 Tsujitani S, Oka S, Saito H et al. Less invasive surgery for early gastric cancer based on the low probability of lymph node metastasis. *Surgery* 1999; 125: 148–54.



- 46 Fujii K. A clinicopathological study on the indications of limited surgery for submucosal gastric cancer. *Jpn. J. Gastroenterol. Surg.* 1998; **31**: 2055–62.
- 47 Gotoda T, Yanagisawa A, Sasako M *et al.* Incidence of lymph node metastasis from early gastric cancer: estimation with a large number of cases at two large centers. *Gastric. Cancer* 2000; **3**: 219–25.
- 48 Japanese Gastric Cancer Association. *Gastric Cancer Treatment Guideline*, 2nd edn. Kyoto: Japanese Gastric Cancer Association, 2004 (in Japanese).
- 49 Hirasawa T, Gotoda T, Miyata S *et al.* Incidence of lymph node metastasis and the feasibility of endoscopic resection for undifferentiated-type early gastric cancer. *Gastric. Cancer* 2009; **12**: 148–52.
- 50 Li C, Kim S, Lai JF *et al.* Risk factors for lymph node metastasis in undifferentiated early gastric cancer. *Ann. Surg. Oncol.* 2008; **15**: 764–9.
- 51 Abe N, Watanabe T, Sugiyama M *et al.* Endoscopic treatment or surgery for undifferentiated early gastric cancer? *Am. J. Surg.* 2004; **188**: 181–4.
- 52 Li H, Lu P, Lu Y *et al.* Predictive factors for lymph node metastasis in poorly differentiated early gastric cancer and their impact on the surgical strategy. *World J. Gastroenterol.* 2008; **14**: 4222–6.
- 53 Ye BD, Kim SG, Lee JY *et al.* Predictive factors for lymph node metastasis and endoscopic treatment strategies for undifferentiated early gastric cancer. *J. Gastroenterol. Hepatol.* 2008; **23**: 46–50.
- 54 Fearon ER, Vogelstein B. A genetic model for colorectal tumorigenesis. *Cell* 1990; **61**: 759–67.
- 55 Shimoda T, Ikegami M, Fujisaki J, Matsui T, Aizawa S, Ishikawa E. Early colorectal carcinoma with special reference to its development de novo. *Cancer* 1989; **64**: 1138–46.
- 56 Bedenne L, Faivre J, Boutron MC, Piard F, Cauvin JM, Hillon P. Adenoma—carcinoma sequence or 'de novo' carcinogenesis? A study of adenomatous remnants in a population-based series of large bowel cancers. *Cancer* 1992; **69**: 883–8.
- 57 Soetikno RM, Kaltenbach T, Rouse RV *et al.* Prevalence of nonpolypoid (flat and depressed) colorectal neoplasms in asymptomatic and symptomatic adults. *Jama* 2008; **299**: 1027–35.
- 58 Rembacken BJ, Fujii T, Cairns A *et al.* Flat and depressed colonic neoplasms: a prospective study of 1000 colonoscopies in the UK. *Lancet* 2000; **355**: 1211–14.
- 59 Morson BC, Whiteway JE, Jones EA, Macrae FA, Williams CB. Histopathology and prognosis of malignant colorectal polyps treated by endoscopic polypectomy. *Gut* 1984; **25**: 437–44.
- 60 Kitajima K, Fujimori T, Fujii S *et al.* Correlations between lymph node metastasis and depth of submucosal invasion in submucosal invasive colorectal carcinoma: a Japanese collaborative study. *J. Gastroenterol.* 2004; **39**: 534–43.
- 61 Egashira Y, Yoshida T, Hirata I *et al.* Analysis of pathological risk factors for lymph node metastasis of submucosal invasive colon cancer. *Mod. Pathol.* 2004; **17**: 503–11.
- 62 Haggitt RC, Glotzbach RE, Soffer EE, Wruble LD. Prognostic factors in colorectal carcinomas arising in adenomas: implications for lesions removed by endoscopic polypectomy. *Gastroenterology* 1985; **89**: 328–36.
- 63 Uraoka T, Saito Y, Matsuda T *et al.* Endoscopic indications for endoscopic mucosal resection of laterally spreading tumours in the colorectum. *Gut* 2006; **55**: 1592–7.
- 64 Matsuda T, Fujii T, Saito Y *et al.* Efficacy of the invasive/non-invasive pattern by magnifying chromoendoscopy to estimate the depth of invasion of early colorectal neoplasms. *Am. J. Gastroenterol.* 2008; **103**: 2700–6.
- 65 Gotoda T, Friedland S, Hamanaka H, Soetikno R. A learning curve for advanced endoscopic resection. *Gastrointest. Endosc.* 2005; **62**: 866–7.
- 66 Suzuki N. Endoscopic submucosal dissection (ESD) for large, sessile colorectal polyps: early experience at a UK centre. *CME Gastroenterol. Hepatol. Nutr.* 2008; **9**: 121–6.
- 67 Dinis-Ribeiro M, Pimentel-Nunes P, Afonso M, Costa N, Lopes C, Moreira-Dias L. A European case series of endoscopic submucosal dissection for gastric superficial lesions. *Gastrointest. Endosc.* 2009; **69**: 350–5.
- 68 Gastroenterology, BSo. *Guidelines on Safety and Sedation During Endoscopic Procedures*, 2003.
- 69 Thomson A, Andrew G, Jones DB. Optimal sedation for gastrointestinal endoscopy: review and recommendations. *J. Gastroenterol. Hepatol* 2010; **25**: 469–78.

# The Natural History of Non-Polypoid Colorectal Neoplasms

Nozomu Kobayashi, MD<sup>a,\*</sup>, Takahisa Matsuda, MD, PhD<sup>b</sup>,  
Yasushi Sano, MD, PhD<sup>c</sup>

## KEYWORDS

• Non-polypoid colorectal neoplasms • Depressed lesion  
• De novo cancer • Colonoscopy • Natural history

The importance of non-polypoid colorectal neoplasms (NP-CRN) is now recognized throughout the world.<sup>1-9</sup> There is little information, however, known about the natural history of NP-CRN, perhaps because the initial reports of NP-CRN suggested that it had high risk of invasion and lymph node metastasis as compared with polypoid lesions of similar size.<sup>10,11</sup> Long-term follow-up of NP-CRN without resection was therefore not an accepted treatment strategy, and had been reported only based on analysis of interval neoplasms or sporadic case reports. In addition, outside Japan, many endoscopists viewed NP-CRN, especially the depressed lesion, as a uniquely Japanese phenomenon and thus, paid little attention to such lesions, limiting the data even more. This article will summarize the available data to gain some estimates of the natural history of NP-CRN.

## RADIOGRAPHIC ANALYSIS

Matsui and colleagues<sup>12</sup> reported a retrospective analysis of a series of colorectal cancers that were missed by double-contrast barium enema examinations. They found six depressed and seven flat lesions (41%) could be retrospectively identified as antecedent lesions that gave rise to 32 advanced cancers. The authors found that all depressed lesions developed into nonprotuberant-type advanced colorectal cancers, whereas flat or polypoid lesions had a possibility to develop into either protuberant or nonprotuberant-type advanced colorectal cancers.

<sup>a</sup> Department of Diagnostic Imaging, Tochigi Cancer Center, 4-9-13 Yonan, Utsunomiya, Tochigi 320-0834, Japan

<sup>b</sup> Endoscopy Division, National Cancer Center Hospital, 5-1-1 Tsukiji, Chuo-ku, Tokyo 104-0045, Japan

<sup>c</sup> Gastrointestinal Center, Sano Hospital, 2-5-1 Shimizugaoka, Tarumi-ku, Kobe, Hyogo, 655-0031, Japan

\* Corresponding author.

E-mail address: nkobayas@tcc.pref.tochigi.lg.jp

Umetani and colleagues<sup>13</sup> reported 11 cases of colorectal cancers that had more than two barium enema examinations that were at least 6 months apart. Five non-polypoid submucosal invasive cancers were studied; three developed from non-polypoid and two from polypoid lesions. The authors estimated tumor doubling time to evaluate the growth rate of each tumor, and suggested that NP-CRN grew slowly compared with polypoid lesion and maintained their macroscopic morphology. The data, however, are limited.

### COLONOSCOPIC ANALYSIS

Matsui and colleagues<sup>14</sup> reported eight early colorectal cancers that were incidentally followed by colonoscopy. Thirty-five cases, including 14 NP-CRN as initial lesions, were reviewed. They found that NP-CRN progressed to submucosally invasive cancer, retaining its non-polypoid configuration, and some flat lesions developed depressed areas during their progression. Although the authors also estimated the speed of growth compared with polypoid and non-polypoid lesions, they could not conclude which type grew more rapidly.

Sato and colleagues<sup>15</sup> prospectively followed 12 small flat adenomas. The size of the lesions ranged from 2 to 6 mm (median 4 mm), and the observation period ranged from 11 to 26 months (median 19 months). Although eight lesions showed various changes in their shape, only two lesions demonstrated an increase in diameter of the tumor. All of the flat lesions were subsequently removed endoscopically and found to be adenomas. The authors concluded small flat adenoma did not rapidly progress, and configuration change did not indicate tumor progression or invasion.

Watari and colleagues<sup>16</sup> conducted prospective colonoscopic study to elucidate the natural history of NP-CRN. The authors observed 75 colorectal tumors measuring less than 1 cm in diameter in 50 patients. The average follow-up period was 22 months, and 62 lesions (83%) were NP-CRN. They concluded similar observations as those of Sato and colleagues, although they found that 40% of small non-polypoid lesions had exophytic growth with time. This finding suggested some small non-polypoid lesions follow the adenoma-carcinoma sequence the same as polypoid lesions.

### THE IMPORTANCE OF THE DEPRESSED-TYPE NP-CRN

Matsuda and colleagues<sup>17</sup> analyzed 6638 colorectal neoplasms, excluding advanced cancers, treated in National Cancer Center Hospital, Tokyo, Japan. There were 4471 (67%) and 2167 (33%), polypoid and non-polypoid colorectal neoplasms, respectively. Among all non-polypoid lesions, there were 178 (2.7%) depressed lesions, 109 (61%) of which were diagnosed as high-grade dysplasia or submucosally invasive cancer. Among 5538 (83%) lesions that were identified as low- or high-grade dysplasia, the proportion of depressed lesions was 1.3%. On the other hand, depressed type was identified in 39% (Table 1) of submucosal cancers. This discrepancy may indicate of a rapid progression rate of depressed lesions into invasive cancers.

Sano and colleagues<sup>18</sup> described the incidence of depressed lesions among all of colorectal neoplasms, again excluding advanced cancers. Their multicenter retrospective study conducted in eight Japanese referral institutes revealed that the incidence of depressed lesions was 1.94% (1291 depressed lesions out of 66,670 neoplasms), and, in particular, 51.2% of intramucosal depressed lesions were diagnosed as high-grade dysplasia. These data also suggested that intramucosal depressed lesions showed more aggressive behavior and were perhaps more likely

Table 1  
Relationship between macroscopic type and histopathological findings

	LGD	HGD	SM-Ca
Polypoid	3781 (68.3)	578 (67.9)	112 (45.0)
Flat	1688 (30.5)	260 (30.6)	41 (16.5)
Depressed	69 (1.2)	13 (1.5)	96 (38.6)
Total	5538 (83.4)	851 (12.8)	249 (3.8)

*Abbreviations:* HGD, high-grade dysplasia; LGD, low-grade dysplasia; SM-Ca, submucosal invasive cancer.

*Data from* Matsuda T, Saito Y, Hotta K, et al. Prevalence and clinicopathological features of non-polypoid colorectal neoplasms: should we pay more attention to identifying flat and depressed lesions? *Dig Endosc* 2010;22(Suppl 1):S57-62.

to develop into invasive cancers as compared with the polypoid lesions. The depressed type of NP-CRN appears to be pathologically and molecular biologically distinct that other types of NP-CRN.

Several authors reported that depressed-type colorectal cancer does not arise from an adenomatous polyp. This theory was called de novo carcinogenesis, and lack of K-ras mutation was thought to be a distinctive genetic feature.<sup>10,11,19,20</sup> Goto and colleagues<sup>21</sup> reported the proportion of de novo cancers among all colorectal cancers in a cohort of 14,817 Japanese populations. The authors defined de novo cancers according to both criteria: (1) the absence of adenomatous components and (2) all lateral margins of the tumor covered with normal mucosa and non-polypoid growth pattern. They concluded that 22.9% of early colorectal cancers were de novo cancers. Chen and colleagues,<sup>22</sup> from Taiwan, also assessed the proportion of de novo carcinomas using the Markov model, and demonstrated about 30% of colorectal cancers arising from de novo sequence.

Table 2  
Description of 13 interval cancers diagnosed within 3 years of a initial colonoscopy

Number	Macroscopic Type	Size (mm)	Location	Depth of Lesion
1	lsp (semipedunculated)	13	Sigmoid	SM
2	lsp (semipedunculated)	15	Sigmoid	SM
3	ls (sessile)	8	Rectum	SM
4	ls (sessile)	10	Sigmoid	SM
5	ls (sessile)	20	Rectum	MP
6	ls (sessile)	6	Transverse	SM
7	lla (flat)	15	Transverse	SM
8	lla (flat)	20	Sigmoid	SM
9	lla + llc (depressed)	20	Cecum	SM
10	lla + llc (depressed)	20	Transverse	SM
11	lla + llc (depressed)	10	Rectum	MP
12	lla + llc (depressed)	6	Ascending	SM
13	lla + llc (depressed)	20	Sigmoid	SS

*Abbreviations:* MP, muscularis; SM, submucosa; SS, subserosa.

*Data from* Matsuda T, Fujii T, Sano Y, et al. Five-year incidence of advanced neoplasia after initial colonoscopy in Japan: a multicenter retrospective cohort study. *Jpn J Clin Oncol* 2009;39:435-42.

## THE JAPAN POLYP STUDY

To clarify the natural history of NP-CRN, a large cohort study focused on the detection of NP-CRN is required. Matsuda and colleagues<sup>23</sup> reported the results of multicenter retrospective cohort study to evaluate 5-year incidence of advanced neoplasia after initial colonoscopy. The authors studied 5309 patients with a median follow-up period of 5.1 years. Endoscopists diagnosed 13 invasive cancers on follow-up within 3 years. The initial colonoscopies were performed by Japanese endoscopists who had proficient technique with chromoendoscopy to diagnose NP-CRN, and the patients had good preparation quality, taking polyethylene glycol (PEG) solution in the morning on the day of colonoscopy. Out of the 13 incident cancer cases, seven were NP-CRN, and the mean size of these lesions was less than 15 mm in diameter (Table 2). These data suggested that NP-CRN is responsible for interval cancers, defined as colorectal cancers diagnosed within several years of a complete colonoscopy. Much prospective data, however, are necessary to elucidate the natural history and epidemiology of these lesions. The Japan Polyp Study (JPS) is a multicenter randomized controlled trial prospectively evaluating follow-up surveillance strategy for Japanese patients after removal of all polypoid and non-polypoid neoplasms.<sup>24</sup> This study is intended to continue until 2011 and hopefully will provide new information on the detection and progression of NP-CRN.

## SUMMARY

The natural history of NP-CRN is mostly unknown. The results of small observational studies suggest that NP-CRN lesions develop into invasive cancer with minimal size expansion. Among NP-CRN lesions, depressed lesions show more aggressive behavior and frequently develop into invasive cancers compared with polypoid lesions, regardless of their low incidence. A large prospective cohort study focused on the detection of NP-CRN is currently ongoing.

## REFERENCES

1. Muto T, Kamiya J, Sawada T, et al. Small flat adenoma of the large bowel with special reference to its clinicopathologic features. *Dis Colon Rectum* 1985;28: 847-51.
2. Kudo S. Endoscopic mucosal resection of flat depressed type of early colorectal cancer. *Endoscopy* 1993;25:455-61.
3. Fujii T, Rembacken BJ, Dixon MF, et al. Flat adenomas in the United Kingdom: are treatable cancers being missed? *Endoscopy* 1998;30:437-43.
4. Saitoh Y, Waxman I, West AB, et al. Prevalence and distinctive biologic features of flat colorectal adenomas in a North American population. *Gastroenterology* 2001; 120:1657-65.
5. Tsuda S, Veress B, Toth E, et al. Flat and depressed colorectal tumours in a southern Swedish population: a prospective chromoendoscopic and histopathological study. *Gut* 2002;51:550-5.
6. Rembacken BJ, Fujii T, Cairns A, et al. Flat and depressed colonic neoplasms: a prospective study of 1000 colonoscopies in the UK. *Lancet* 2000;355:1211-4.
7. Parra-Blanco A, Gimeno-Garcia AZ, Nicolas-Perez D, et al. Risk for high-grade dysplasia or invasive carcinoma in colorectal flat adenomas in a Spanish population. *Gastroenterol Hepatol* 2006;29:602-9.

8. Soetikno RM, Kaltenbach T, Rouse RV, et al. Prevalence of nonpolypoid (flat and depressed) colorectal neoplasms in asymptomatic and symptomatic adults. *JAMA* 2008;299:1027–35.
9. Chiu HM, Lin JT, Chen CC, et al. Prevalence and characteristics of nonpolypoid colorectal neoplasm in an asymptomatic and average-risk Chinese population. *Clin Gastroenterol Hepatol* 2009;7:463–70.
10. Kudo S, Kashida H, Tamura T, et al. Colonoscopic diagnosis and management of nonpolypoid early colorectal cancer. *World J Surg* 2000;24:1081–90.
11. Shimoda T, Ikegami M, Fujisaki J, et al. Early colorectal carcinoma with special reference to its development de novo. *Cancer* 1989;64:1138–46.
12. Matsui T, Yao T, Iwashita A. Natural history of early colorectal cancer. *World J Surg* 2000;24:1022–8.
13. Umetani N, Masaki T, Watanabe T, et al. Retrospective radiographic analysis of nonpedunculated colorectal carcinomas with special reference to tumour doubling time and morphological change. *Am J Gastroenterol* 2000;95:1794–9.
14. Matsui T, Tsuda S, Iwashita A, et al. Retrospective endoscopic study of developmental and configurational changes of early colorectal cancer: eight cases and a review of the literature. *Dig Endosc* 2004;16:1–8.
15. Sato T, Konishi F, Togashi K, et al. Prospective observation of small flat tumors in the colon through colonoscopy. *Dis Colon Rectum* 1999;42:1457–63.
16. Watari J, Saitoh Y, Obara T, et al. Natural history of colorectal nonpolypoid adenomas: a prospective colonoscopic study and relation with cell kinetics and K-ras mutations. *Am J Gastroenterol* 2002;97:2109–15.
17. Matsuda T, Saito Y, Hotta K, et al. Prevalence and clinicopathological features of nonpolypoid colorectal neoplasms: should we pay more attention to identifying flat and depressed lesions? *Dig Endosc* 2010;22(Suppl 1):S57–62.
18. Sano Y, Tanaka S, Teixeira CR, et al. Endoscopic detection and diagnosis of 0-Ic neoplastic colorectal lesions. *Endoscopy* 2005;37:261–7.
19. Fujimori T, Satonaka K, Yamamura-Idei Y, et al. Noninvolvement of ras mutations in flat colorectal adenomas and carcinomas. *Int J Cancer* 1994;57:51–5.
20. Kaneko K, Fujii T, Kato S, et al. Growth pattern and genetic changes of colorectal carcinoma. *Jpn J Clin Oncol* 1998;28:196–201.
21. Goto H, Oda Y, Murakami Y, et al. Proportion of de novo cancers among colorectal cancers in Japan. *Gastroenterology* 2006;131:40–6.
22. Chen CD, Yen MF, Wang WM, et al. A case-cohort study for the disease natural history of adenoma-carcinoma and de novo carcinoma and surveillance of colon and rectum after polypectomy: implication for efficacy of colonoscopy. *Br J Cancer* 2003;88:1866–73.
23. Matsuda T, Fujii T, Sano Y, et al. Five-year incidence of advanced neoplasia after initial colonoscopy in Japan: a multicenter retrospective cohort study. *Jpn J Clin Oncol* 2009;39:435–42.
24. Sano Y, Fujii T, Oda Y, et al. A multicenter randomized controlled trial designed to evaluate follow-up surveillance strategies for colorectal cancer: the Japan Polyp Study. *Dig Endosc* 2004;16:376–8.

## Diagnostic accuracy of narrow-band imaging and pit pattern analysis significantly improved for less-experienced endoscopists after an expanded training program

Reiji Higashi, MD, Toshio Uraoka, MD, PhD, Jun Kato, MD, PhD, Kenji Kuwaki, MD, Shin Ishikawa, MD, PhD, Yutaka Saito, MD, PhD, Takahisa Matsuda, MD, PhD, Hiroaki Ikematsu, MD, Yasushi Sano, MD, PhD, Seiyuu Suzuki, MD, PhD, Yoshitaka Murakami, PhD, Kazuhide Yamamoto, MD, PhD

Okayama, Tokyo, Chiba, Kobe, Niihama, Otsu, Japan

**Background:** Previous reports assessing diagnostic skill using narrow-band imaging (NBI) and pit pattern analysis for colorectal polyps involved only highly experienced endoscopists.

**Objective:** To evaluate diagnostic skills of less-experienced endoscopists (LEE group) for differentiation of diminutive colorectal polyps by using NBI and pit pattern analysis with and without magnification after an expanded training program.

**Design:** Prospective study.

**Patients:** This study involved 32 patients with 44 colorectal polyps (27 adenomas and 17 hyperplastic polyps) of  $\leq 5$  mm that were identified and analyzed by using conventional colonoscopy as well as non-magnification and magnification NBI and chromoendoscopy followed by endoscopic removal for histopathological analysis.

**Intervention:** Before a training course, 220 endoscopic images were distributed in randomized order to residents with no prior endoscopy experience (NEE group) and to the LEE group, who had performed colonoscopies for more than 5 years but had never used NBI. The 220 images were also distributed to highly experienced endoscopists (HEE group) who had routinely used NBI for more than 5 years. The images were distributed to the NEE and LEE groups again after a training class. Magnification NBI and chromoendoscopy images were assessed by using the Sano and Kudo classification systems, respectively.

**Main Outcome Measurements:** Diagnostic accuracy and interobserver agreement for each endoscopic modality in each group.

**Results:** Diagnostic accuracy was significantly higher, and kappa ( $\kappa$ ) values improved in the LEE group for NBI with high magnification after expanded training. Diagnostic accuracy and  $\kappa$  values when using high-magnification NBI were highest among endoscopic techniques for the LEE group after such training and the HEE group (accuracy 90% vs 93%;  $\kappa = 0.79$  vs 0.85, respectively).

**Limitations:** Study involved only polyps of  $\leq 5$  mm.

**Conclusion:** Using high-magnification NBI increased the differential diagnostic skill of the LEE group after expanded training so that it was equivalent to that of the HEE group. (*Gastrointest Endosc* 2010;72:127-35.)

*Abbreviations:* CC, conventional colonoscopy; CE-H, high-magnification chromoendoscopy; CE-L, low-magnification chromoendoscopy; HEE, highly experienced endoscopist; LEE, less-experienced endoscopist; NBI, narrow-band imaging; NBI-H, high-magnification narrow-band imaging; NBI-L, low-magnification narrow-band imaging; NEE, no-experience endoscopist; SSA, sessile serrated adenoma.

**DISCLOSURE:** All authors disclosed no financial relationships relevant to this publication.

Copyright © 2010 by the American Society for Gastrointestinal Endoscopy  
0016-5107/\$36.00

doi:10.1016/j.gie.2010.01.054

Received August 10, 2009. Accepted January 25, 2010.

Reprint requests: Toshio Uraoka, MD, PhD, Department of Endoscopy, Okayama University Hospital, 2-5-1 Shikata-cho, Okayama 700-8558, Japan.

Current affiliations: Department of Gastroenterology and Hepatology (R.H., J.K., K.K., S.I., K.Y.), Okayama University Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, Department of Endoscopy (T.U.), Okayama University Hospital, Okayama, Endoscopy Division (Y.S., T.M.), National Cancer Center Hospital, Tokyo, Division of Digestive Endoscopy and Gastrointestinal Oncology (H.I.), National Cancer Center Hospital East, Kashiwa, Gastrointestinal Center (Y.S.), Sano Hospital, Kobe, Department of Internal Medicine (S.S.), Sumitomo Besshi Hospital, Niihama, Department of Medical Statistics (Y.M.), Shiga University of Medical Science, Otsu, Japan.

It is widely accepted that adenomatous polyps are precursors of colorectal cancer, and performing polypectomies on such lesions can reduce the risk of subsequent colorectal cancer by up to 80% for a period that may exceed 10 years.<sup>1</sup> In addition, adenomas are a major factor in guidelines that have been developed for recommended colonoscopy surveillance intervals after polypectomies because they are a powerful predictor for future colorectal cancer risk.<sup>1-3</sup> Small colorectal adenomas as well as advanced adenomas<sup>4</sup> are precursors of colorectal cancer, and multiple genetic alterations have been implicated in the adenoma-carcinoma sequence.<sup>5</sup>

Endoscopic differentiation of small adenomas from non-adenomatous polyps is important because endoscopists should avoid performing any unnecessary procedure, including polypectomies, that can cause related complications such as bleeding and perforation.<sup>6,7</sup> The diagnostic accuracy of conventional colonoscopy for such colorectal polyps, however, has previously been reported to be unsatisfactory.<sup>8,9</sup> In contrast, chromoendoscopy with indigocarmine dye spraying has been shown to be an effective procedure for detecting and evaluating colorectal polyps<sup>10-15</sup> despite having several disadvantages, including a longer procedure time and the additional cost for dye spraying.

Narrow-band imaging (NBI) is an innovative optical technology providing a unique image that emphasizes the morphological and structural character of lesions as well as their surface capillary patterns.<sup>16-25</sup> It has been reported that this modality is a new non-dye tool for differentiating neoplastic from non-neoplastic polyps, with a diagnostic accuracy including pit pattern analysis equivalent to that of chromoendoscopy.<sup>17-21</sup> Such reports have been based on studies involving only highly experienced endoscopists, however, with few published articles concerned with the learning curve for NBI being dependent on an individual endoscopist's experience and ability. The aim of this study was to determine, therefore, whether expanded training in the effective use of NBI and pit pattern analysis with and without magnification would improve the diagnostic skill of less-experienced endoscopists in the differentiation of diminutive colorectal polyps.

## METHODS

### Patients

Patients scheduled for a total colonoscopy at Okayama University Hospital and Sumitomo Besshi Hospital between September and October 2008 were invited to participate in this study. Informed consent was obtained from all patients before their examinations. Patients with inflammatory bowel disease, familial adenomatous polyposis, an international normalized ratio greater than 2.0, or a platelet count less than 50,000/mm<sup>3</sup> were excluded from this study.

### Take-home Message

- Expanded interactive training in effective use of narrow-band imaging both with and without magnification as well as pit pattern analysis improved diagnostic accuracy and interobserver agreement of less-experienced colonoscopists in differentiating diminutive colorectal polyps. Using narrow-band imaging with high magnification increased the differential diagnostic skill of less-experienced colonoscopists who underwent such training to a level equivalent to that of highly experienced colonoscopists.

### Colonoscopy and polyp assessment protocol

Bowel preparation consisted of patients drinking 2 to 3 liters of polyethylene glycol solution in the morning before their procedures.<sup>13</sup> Total colonoscopies were prospectively performed by using a video endoscopic system (EVIS, Lucera Spectrum; Olympus Co, Tokyo, Japan) with CF-H260AZI or PCF-Q240ZI magnification colonoscopes (Olympus) by two highly experienced endoscopists (R.H., T.U.), each of whom had previously performed over 1000 colonoscopies annually.

When a lesion was detected by conventional colonoscopy examination, surface mucus was washed away with lukewarm water, and endoscopic images were taken in the following order: conventional colonoscopy (CC), low-magnification NBI (NBI-L), high-magnification NBI (NBI-H), low-magnification chromoendoscopy (CE-L) and high-magnification chromoendoscopy (CE-H). A standard optical filter was used for both CC and chromoendoscopy, with chromoendoscopic images taken after 0.2% indigocarmine dye was sprayed on the lesion surface. The enhanced surface structure function of the video image processor at the level A5 setting was used in taking all endoscopic images.<sup>23</sup> Location, size, and macroscopic type of each lesion were recorded, with size measured by using open forceps. Lesions were classified macroscopically based on the criteria of the Paris classification of superficial GI lesions.<sup>26</sup> A biopsy, polypectomy, or EMR was then performed, and the resulting specimen was analyzed histopathologically.

### Image evaluator categories

A total of 12 doctors with different levels of endoscopic experience were asked to independently evaluate endoscopic images. The doctors were separated into 3 groups: 4 residents with no prior endoscopy experience (NEE group); 4 less experienced endoscopists each of whom had performed colonoscopies for more than 5 years but had never used NBI (LEE group); and 4 highly experienced endoscopists each of whom had routinely used magnification colonoscopy with NBI for more than 5 years (HEE group).



## Assessment of endoscopic images

The best quality endoscopic images were selected for each modality and stored digitally in JPEG format. All images were distributed in randomized order to each group of evaluators. For the NEE and LEE groups, the same images were distributed in randomized order once before and once after those group members participated in an intensive, 1-hour, interactive training program on white-light endoscopy, NBI, and chromoendoscopy. The program included information on the Sano NBI classification<sup>20,21</sup> and the Kudo pit pattern classification<sup>27,28</sup> and used an atlas of endoscopic images of polyps produced by an independent group of highly experienced endoscopists. Although they were completely unaware of the histopathological results, every participant correctly diagnosed polyps as either neoplasms or non-neoplasms by using (1) chromoendoscopy based on the Kudo pit pattern classification, with types III (including IIIa and IIIb), IV, and V (including VI and VN) considered to be neoplasms and types I and II regarded as non-neoplasms,<sup>27,28</sup> (2) NBI-L that revealed a brownish area determined to be neoplastic; and (3) NBI-H with the Sano classification of meshed capillary vessel pattern, in which types II and III were considered neoplastic and type I, without meshed capillaries, was non-neoplastic (Figs. 1 and 2).<sup>20-21</sup> Patient information such as age, sex, and clinical diagnosis was not disclosed to any of the evaluators, and discussions were not permitted among the doctors individually or in groups.

## Statistical analysis

The diagnostic accuracy of each endoscopic modality was assessed in reference to histopathological results. Estimates of diagnostic accuracy were calculated based on the average diagnostic accuracy for each group of doctors as well as for each diagnostic modality. The upper and lower 95% confidence interval (CI) limits were calculated by using a normal model that consisted of symmetric CIs, with limits at a distance from the estimate equal to the product of 1.96 times the standard error. Interobserver agreement in diagnosing colorectal lesions in each group and by each modality was determined by calculation of the kappa statistic ( $\kappa$ ) and its 95% CI by using the Fleiss method. Diagnostic accuracies before training and after training in the NEE and LEE groups were compared with the McNemar test. As for differences in diagnostic accuracies after training among the NEE, LEE, and HEE groups, those findings were analyzed by using the Fisher exact test. Multiple statistical testing of outcome data was conducted in this study, therefore, a Bonferroni correction was applied, and differences with a  $P$  value of  $< .025$  were considered significant as the correction. A  $\kappa$  value of  $< 0.4$  was regarded as poor agreement, 0.41 to 0.60 fair agreement, 0.61 to 0.80 good agreement, and  $> 0.80$  excellent agreement. Statistical analyses were conducted by using

version 7.0 of the JMP statistical software package (SAS Institute, Cary, NC) and a Microsoft Excel 2007 spreadsheet (Microsoft, Renton, Wash).

## RESULTS

### Clinicopathological features of colorectal lesions


Seventy-two consecutive patients were enrolled in this study for prospective endoscopic evaluation. A total of 44 lesions of  $\leq 5$  mm were identified and analyzed in 32 patients (Table 1). Mean ( $\pm$  standard deviation [SD]) patient age was  $61.2 \pm 12.3$  years, and the male/female ratio was 2.2:1. Bowel preparation was considered adequate in all examinations, and complete colonoscopy was performed to the cecum in every case. There were no complications during any procedure. Of the 44 lesions, 37 were macroscopically classified as type 0-Is, 6 as type 0-IIa, and 1 as type 0-IIc. Mean ( $\pm$  SD) lesion size was  $3.4 \pm 1.1$  mm. As for location, 22 polyps (50%) were found in the right colon (cecum, ascending, and transverse colon), 14 (32%) in the left colon (descending and sigmoid colon), and 8 (18%) in the rectum. Histopathological assessments included 27 adenomas (61%) and 17 hyperplastic polyps (39%). A total of 220 images of the 44 lesions were collected as each lesion was photographed during CC, NBI-L, NBI-H, CE-L, and CE-H.

### Diagnostic accuracy of NBI and pit pattern analysis

Table 2 indicates diagnostic accuracy for each endoscopic modality. In the NEE group, diagnostic accuracies using CC, NBI-L, and NBI-H significantly improved after the training program (CC,  $P < .001$ ; NBI-L,  $P = .006$ ; and NBI-H,  $P < .001$ ), but the NEE group's diagnostic accuracies were still significantly lower in all modalities except CC compared with the HEE group (CC,  $P = .049$ ; NBI-L,  $P = .0023$ ; NBI-H,  $P < .001$ ; CE-L,  $P < .001$ ; and CE-H,  $P < .001$ ). Diagnostic accuracies in the LEE group for NBI-L, NBI-H, and CE-H also improved significantly after the training program ( $P = .001$ ,  $P < .001$ , and  $P = .001$ , respectively). In contrast with the NEE group's results, however, subsequent diagnostic accuracies of the LEE group were not significantly different from diagnostic accuracies of the HEE group with respect to the CC, NBI-L, NBI-H, CE-L, and CE-H modalities ( $P = 1.0$ ,  $P = .60$ ,  $P = .57$ ,  $P = .031$ , and  $P = .48$ , respectively).

### Assessment of interobserver agreement based on endoscopic experience

Interobserver agreements in the HEE group for NBI-H were  $> 0.80$  representing excellent agreement and  $> 0.60$  for NBI-L, CE-L, and CE-H, representing good agreement ( $\kappa$  value CC, 0.5; NBI-L, 0.62; NBI-H, 0.85; CE-L, 0.69; CE-H




**Figure 1.** Examples of colorectal neoplastic polyps viewed by different endoscopic modalities in this study. **A**, Conventional colonoscopy view. **B**, Low-magnification NBI showed a brownish area. **C**, High-magnification NBI revealed meshed capillary vessels indicative of Sano classification type II. **D**, Low-magnification chromoendoscopy using 0.2% indigo-carmin dye spraying clearly revealed demarcated margins and surface structure. **E**, High-magnification chromoendoscopy clearly indicated Kudo classification type III-L. *NBI*, narrow-band imaging.

0.7). Meanwhile,  $\kappa$  values for NBI-H in the LEE group improved to “good agreement” after the training program, whereas the  $\kappa$  values for NBI-L, CE-L, and CE-H in the LEE group improved to “fair agreement” after the training program ( $\kappa$  value before vs after training: NBI-H, 0.46 vs 0.79; NBI-L, 0.31 vs 0.54; CE-L, 0.32 vs 0.44; and CE-H, 0.33 vs 0.59). In contrast, however, none of the  $\kappa$  values for any of the modalities in the NEE group improved beyond “poor agreement” after the training program (CC, -0.068 vs 0.24;

NBI-L, 0.059 vs 0.25; NBI-H, 0.16 vs 0.39; CE-L, 0.28 vs 0.23; and CE-H, 0.12 vs 0.18) (Fig. 3).

When we compared diagnostic accuracy for each modality in the NEE group after the training program, the LEE group after the training program, and the HEE group, NBI-H had the highest accuracy rate among all 3 groups (Table 2). Similarly, the  $\kappa$  value for NBI-H was significantly higher in the NEE group after the training program, the LEE group after the training program, and the HEE group



**Figure 2.** Examples of colorectal non-neoplastic polyps viewed by different endoscopic modalities in this study. **A**, Conventional colonoscopy view. **B**, Low-magnification NBI showed a non-brownish area. **C**, High-magnification NBI in which meshed capillary vessels were not visible or only faintly visible, indicative of Sano classification type I. **D**, Low-magnification chromoendoscopy using 0.2% indigo-carmin dye spraying clearly revealed demarcated margins and surface structure. **E**, High-magnification chromoendoscopy clearly indicated Kudo classification type II. *NBI*, narrow-band imaging.

compared with the other endoscopic diagnostic modalities (Fig. 3).

## DISCUSSION

Endoscopic diagnostic tools and technology are expected to be accurate and provide reliably reproducible agreement as well as be easy to use, readily available, and relatively inexpensive, but sufficient skill on the part of the

endoscopist is still required for proper diagnosis. Our prospective study demonstrated significant improvement in the LEE group in diagnostic accuracy when using NBI and CE after undergoing limited but intensive training. The improved diagnostic accuracy of the LEE group was equivalent to that of the HEE group in terms of differential diagnosis using NBI-L, NBI-H, and CE-H. In addition, both higher diagnostic accuracy (>80%) and good interobserver agreement ( $\kappa$  value >0.6) for diminutive colorectal

**TABLE 1. Patient characteristics and histopathological features of lesions**

Patients (n = 32)	
Sex, male/female	22/10
Age, years, mean ( $\pm$ SD)	61.2 (12.3)
Lesions (n = 44)	
Macroscopic type	
0-Ic	37
0-IIa	6
0-IIc	1
Size, mm, mean ( $\pm$ SD)	3.4 (1.1)
Location (right*/left/rectum)	22/14/8
Histopathology	
Tubular adenoma	27
Hyperplastic polyp	17

SD, Standard deviation.

\*Right: cecum, ascending colon, and transverse colon.

†Left: descending colon and sigmoid colon.

polyps were achieved by the LEE group when using NBI-H after the training program.

The fact that the diagnostic accuracy and  $\kappa$  value of NBI-H were the highest among all the endoscopic techniques analyzed in this study for both the NEE and LEE groups after the expanded training program as well as for the HEE group leads us to suggest that NBI-H is more accurate and provides a higher level of reproducible agreement than the other diagnostic tools in differentiating diminutive neoplastic from non-neoplastic colorectal polyps. Chiu et al<sup>29</sup> earlier validated that diagnostic accuracy of NBI-H was equivalent to that of CE-H. Their study reported that diagnostic accuracies for two experienced endoscopists ranged from 91% to 92% using CE-H and from 87% to 90% using NBI-H. In our study, the diagnostic accuracy of the HEE group was 85% (95% CI, 79%-89%) using CE-H and 93% (95% CI, 88%-96%) with NBI-H, although the difference between the accuracy of the two modalities was not significant. Earlier reports analyzing the diagnostic accuracy rate based on polyp size indicated that differentiation using CE-H was more difficult with diminutive colorectal polyps of <6 mm in size.<sup>6-7</sup>

Our results indicated that it was possible to significantly improve the diagnostic skill for differentiating diminutive colorectal polyps by using NBI-L, NBI-H, CE-L, and CE-H in the LEE group following the limited but intensive 1-hour interactive training program. We believe that, of the various endoscopic modalities, the use of NBI-H by the LEE group subsequently became both statistically equivalent to that of the HEE group in terms of diagnostic accuracy and

closest to reaching "excellent agreement" compared with the other modalities in terms of  $\kappa$  value for two possible reasons. The first concerns the smaller size of the polyps examined in this study, because diagnostic accuracy of diminutive colorectal polyps by using CE-H has been reported to be lower than for polyps of >5 mm.<sup>6,9</sup> It is conceivable that differentiation of diminutive colorectal polyps could have been similarly affected, somehow reducing the diagnostic accuracy of CE-H while not affecting the diagnostic accuracy of NBI-H by the LEE group. Secondly, the possibility exists that members of the LEE group were able to recognize whether or not there were meshed capillary vessels on the surface of the mucosa easier than they could identify the pit patterns of diminutive colorectal polyps.

In the Rogart et al<sup>30</sup> report on the NBI learning curve, diagnostic accuracy by using NBI-L significantly improved as the experience level of endoscopists increased, with the diagnosis of approximately 130 lesions necessary for basic competency. Their findings indicated that educational sessions conducted before the assessment of lesions in combination with continual feedback regarding the accuracy of endoscopic diagnoses compared with histopathological results every 2 weeks for half a year were important factors in achieving a satisfactory learning curve. It has also been reported that use of the Kudo pit pattern classification required a longer learning curve, with experience from diagnosing at least 200 lesions needed to become competent.<sup>6,7,31</sup> In contrast, our study demonstrated that an expanded 1-hour intensive interactive training program conducted by a highly experienced endoscopist enabled the LEE group members in particular to accelerate their learning curve. In addition, the Sano classification with NBI-H appears to have had a shorter learning curve compared with using NBI-L or the Kudo pit pattern classification in the diagnostic differentiation of diminutive colorectal polyps.

Besides having a higher differential diagnosis accuracy and being easier to improve the necessary diagnostic skill for accurately differentiating diminutive colorectal polyps, NBI has other clinical advantages. First, the conventional endoscopic view can be switched almost instantaneously to the NBI view by pressing a single button on the control handle of the colonoscope, and, second, NBI does not require any dye or staining solution to detect and differentiate neoplastic lesions from non-neoplastic lesions.

In recent years, advancements in the quality of endoscopic images available from high-definition endoscopy and chromoendoscopy have considerably enhanced polyp detection. Although the risk of neoplasia in diminutive polyps is <50%, and the risk of high-grade dysplasia is <2%,<sup>7,32,33</sup> diminutive colorectal neoplasms as well as advanced neoplasms are among the precursors of colorectal cancer, and multiple genetic alterations have been implicated in the adenoma-carcinoma sequence.<sup>4</sup> It also has been reported that lesions of  $\leq 5$

TABLE 2. Effectiveness of training program on diagnostic accuracy

Modality	NEE group			LEE group			HEE group	
	Before-training accuracy (95% CI)	After-training accuracy (95% CI)	P value*	Before-training accuracy (95% CI)	After-training accuracy (95% CI)	P value*	Accuracy (95% CI)	P value†
CC	0.43 (0.35-0.50)	0.64 (0.57-0.71)	<.001	0.72 (0.65-0.78)	0.74 (0.67-0.80)	NS	0.74 (0.68-0.80)	NS
NBI-L	0.53 (0.46-0.61)	0.66 (0.59-0.73)	.006	0.65 (0.58-0.72)	0.78 (0.72-0.84)	=.001	0.81 (0.75-0.87)	NS
NBI-H	0.63 (0.56-0.70)	0.74 (0.68-0.80)	<.001	0.73 (0.66-0.79)	0.90 (0.85-0.94)	<.001	0.93 (0.88-0.96)	NS
CE-L	0.68 (0.60-0.74)	0.67 (0.60-0.74)	NS	0.68 (0.60-0.74)	0.76 (0.69-0.82)	NS	0.85 (0.79-0.90)	NS
CE-H	0.63 (0.56-0.70)	0.66 (0.59-0.73)	NS	0.67 (0.60-0.74)	0.81 (0.75-0.87)	=.001	0.85 (0.79-0.89)	NS

NEE, No endoscopy experience; LEE, less experienced endoscopist; HEE, highly experienced endoscopist; CI, confidence interval; CC, conventional colonoscopy; NS, not significant; NBI-L, low-magnification narrow-band imaging; NBI-H, high-magnification narrow-band imaging; CE-L, low-magnification chromoendoscopy; CE-H, high-magnification chromoendoscopy.

\*P values determined by McNemar test comparing before and after training.

†P values determined by Fisher exact test comparing LEE after training and HEE.




Figure 3. Comparison of 95% confidence interval of  $\kappa$  value for each endoscopic diagnostic modality according to endoscopy experience. Each bar represents the range of 95% confidence interval of the  $\kappa$  value. HEE, highly experienced endoscopist group; LEE, less experienced endoscopist group; Before, before participation in an intensive 1-hour interactive training program; After, after participation in an intensive 1-hour interactive training program; NEE, no endoscopy experience group; CC, conventional colonoscopy; NBI-L, low-magnification narrow-band imaging; NBI-H, high-magnification narrow-band imaging; CE-L, low-magnification narrow-band imaging; CE-H, high-magnification chromoendoscopy.

mm make up more than 80% of the colorectal polyps subjected to histopathological assessment.<sup>33</sup> Besides the primary consideration of reducing the risk of future colorectal cancer, the endoscopic differentiation of diminutive neoplastic polyps from non-neoplastic polyps is essential because endoscopists should avoid performing unnecessary procedures, including polypectomies on non-neoplastic polyps, and this also will reduce substantially the number of colorectal polyps requiring histopathological assessment.

There has been considerable interest recently in sessile serrated adenoma (SSA) and serrated adenoma (SA) polyps that also have been associated increasingly with an apparent increased risk of malignant transformation.<sup>34</sup> SSAs endoscopically appear as hyperplastic polyps, but there have not been any published reports as yet applying the Kudo pit pattern analysis to such SSA polyps. In the general population, the prevalence of SSAs has been estimated to range from only 1% to 7% of all polyps, and it further has been shown that most such SSA polyps can exceed 10 mm in size,<sup>35</sup> but we did not detect any SSAs or serrated adenomas in this study. Although it was recently reported that SSAs could be differentiated from hyperplastic polyps by combining NBI and autofluorescence imaging, the report in question had several limitations including the total number of SSAs being relatively small and the lack of any comparison between those two modalities and pit pattern analysis.<sup>36</sup> In addition, the actual prevalence of SSAs is difficult to assess because pathologists have been unable to reach a consensus on the diagnosis of either hyperplastic polyps or SSAs.<sup>37,38</sup> Further studies will be required, therefore, to clarify the endoscopic features and conduct histopathological and molecular-based analyses of SSAs and serrated adenomas.

The primary limitation of our study is that it involved only a small number of polyps. The power of the trial compared to the observed difference was lower because the observed difference was smaller than in the alternative hypothesis used in planning this study. The sample size that was set, however, was not much different from the sample size used in similar studies. Another limitation is that this study was conducted by using endoscopic images. During a "real-time" evaluation, an endoscopist can usually view a detected lesion by using multiple angles and light modalities at variable distances, but we digitally stored all the endoscopic images taken during each exam-

ination, selected the best image from each of the 5 endoscopic observation modalities, and then randomized the distribution order of the images for diagnosis. This process was intended to decrease the likelihood of observational bias and strengthen the reliability of our results, because separate findings based on NBI images and chromoendoscopic images might otherwise have been influenced by the other and made objective evaluation of the individual diagnostic modalities difficult. A third limitation was the relatively short interval between the intensive training program and the follow-up reviews by the NEE and LEE groups. The participants in both groups reviewed all the images for the second time within 24 hours of the training program so as to avoid any possible bias resulting from a feedback learning effect such as self-training. It has previously been reported that feedback received during the development of a diagnostic skill is effective.<sup>39</sup>

In conclusion, NBI, particularly high-magnification NBI, was shown to be a promising tool for diagnostic differentiation of diminutive colorectal neoplastic polyps from non-neoplastic polyps. Expanded training of the LEE group members improved their overall diagnostic ability so that it was equivalent in certain key respects to that of the participating HEE group.

## REFERENCES

- Winawer SJ, Zauber AG, Ho MN, et al. Prevention of colorectal cancer by colonoscopic polypectomy. The National Polyp Study Workgroup. *N Engl J Med* 1993;329:1977-81.
- Atkin WS, Morson BC, Cuzick J. Long-term risk of colorectal cancer after excision of rectosigmoid adenomas. *N Engl J Med* 1992;326:658-62.
- Citarida F, Tomaselli G, Capocaccia R, et al. Efficacy in standard clinical practice of colonoscopic polypectomy in reducing colorectal cancer incidence. *Gut* 2001;48:812-5.
- Winawer SJ, Zauber AG. The advanced adenoma as the primary target of screening. *Gastrointest Endosc Clin N Am* 2002;12:1-9.
- Vogelstein B, Fearon ER, Hamilton SR, et al. Genetic alterations during colorectal-tumor development. *N Engl J Med* 1988;319:525-32.
- Togashi K, Konishi F, Ishizuka T, et al. Efficacy of magnifying endoscopy in the differential diagnosis of neoplastic and non-neoplastic polyps of the large bowel. *Dis Colon Rectum* 1999;42:1602-8.
- Su MY, Ho YP, Chen PC, et al. Magnifying endoscopy with indigo carmine contrast for differential diagnosis of neoplastic and nonneoplastic colonic polyps. *Dig Dis Sci* 2004;49:1123-7.
- Chapuis PH, Dent OF, Goulston KJ. Clinical accuracy in the diagnosis of small polyps using the flexible fiberoptic sigmoidoscope. *Dis Colon Rectum* 1982;25:669-72.
- Neale AV, Demers RY, Budev H, et al. Physician accuracy in diagnosing colorectal polyps. *Dis Colon Rectum* 1987;30:247-50.
- Rembacken BJ, Fujii T, Cairns A, et al. Flat and depressed colonic neoplasms: a prospective study of 1000 colonoscopies in the UK. *Lancet* 2000;355:1211-4.
- Hurlstone DP, Cross SS, Slater R, et al. Detecting diminutive colorectal lesions at colonoscopy: a randomised controlled trial of pan-colonic versus targeted chromoscopy. *Gut* 2004;53:376-80.
- Hurlstone DP, Cross SS, Adam I, et al. Efficacy of high magnification chromoscopic colonoscopy for the diagnosis of neoplasia in flat and depressed lesions of the colorectum: a prospective analysis. *Gut* 2004;53:284-90.
- Konishi K, Kaneko K, Kurahashi T, et al. A comparison of magnifying and nonmagnifying colonoscopy for diagnosis of colorectal polyps: a prospective study. *Gastrointest Endosc* 2003;57:48-53.
- Matsuda T, Fujii T, Saito Y, et al. Efficacy of the invasive/non-invasive pattern by magnifying chromoendoscopy to estimate the depth of invasion of early colorectal neoplasms. *Am J Gastroenterol* 2008;103:2700-6.
- Fu KI, Sano Y, Kato S, et al. Chromoendoscopy using indigo carmine dye spraying with magnifying observation is the most reliable method for differential diagnosis between non-neoplastic and neoplastic colorectal lesions: a prospective study. *Endoscopy* 2004;36:1089-93.
- Sano Y, Muto M, Tajiri H. Optical/digital chromoendoscopy during colonoscopy using narrow-band imaging system. *Digestive Endoscopy* 2005;17:543-8.
- Machida H, Sano Y, Hamamoto Y, et al. Narrow-band imaging in the diagnosis of colorectal mucosal lesions: a pilot study. *Endoscopy* 2004;36:1094-8.
- East JE, Suzuki N, Stavrinidis M, et al. Narrow band imaging for colonoscopic surveillance in hereditary non-polyposis colorectal cancer. *Gut* 2008;57:65-70.
- Hirata M, Tanaka S, Oka S, et al. Magnifying endoscopy with narrow band imaging for diagnosis of colorectal tumors. *Gastrointest Endosc* 2007;65:988-95.
- Sano Y, Ikematsu H, Fu KI, et al. Meshed capillary vessels by use of narrow-band imaging for differential diagnosis of small colorectal polyps. *Gastrointest Endosc* 2009;69:278-83.
- Katagiri A, Fu KI, Sano Y, et al. Narrow band imaging with magnifying colonoscopy as diagnostic tool for predicting history of early colorectal neoplasia. *Aliment Pharmacol Ther* 2008;27:1269-74.
- Uraoka T, Saito Y, Matsuda T, et al. Detectability of colorectal neoplastic lesions using a narrow-band imaging system: a pilot study. *J Gastroenterol Hepatol* 2008;23:1810-5.
- Uraoka T, Sano Y, Saito Y, et al. Narrow-band imaging for improving colorectal adenoma detection: appropriate system function settings are required. *Gut* 2009;58:604-5.
- Yoshida T, Inoue H, Usui S, et al. Narrow-band imaging system with magnifying endoscopy for superficial esophageal lesions. *Gastrointest Endosc* 2004;59:288-95.
- Yao K, Oishi T, Matsui T, et al. Novel magnified endoscopic findings of microvascular architecture in intramucosal gastric cancer. *Gastrointest Endosc* 2002;56:279-84.
- The Paris endoscopic classification of superficial neoplastic lesions: esophagus, stomach, and colon: November 30 to December 1, 2002. *Gastrointest Endosc* 2003;58:53-43.
- Kudo S, Hirota S, Nakajima T, et al. Colorectal tumours and pit pattern. *J Clin Pathol* 1994;47:880-5.
- Kudo S, Rubio CA, Teixeira CR, et al. Pit pattern in colorectal neoplasia: endoscopic magnifying view. *Endoscopy* 2001;33:367-73.
- Chiu HM, Chang CY, Chen CC, et al. A prospective comparative study of narrow-band imaging, chromoendoscopy, and conventional colonoscopy in the diagnosis of colorectal neoplasia. *Gut* 2007;56:373-9.
- Rogart JN, Jain D, Siddiqui UD, et al. Narrow-band imaging without high magnification to differentiate polyps during real-time colonoscopy: improvement with experience. *Gastrointest Endosc* 2008;68:1136-45.
- Kudo S, Tamura S, Nakajima T, et al. Diagnosis of colorectal tumorous lesions by magnifying endoscopy. *Gastrointest Endosc* 1996;44:8-14.
- Buttler LF, Chase MP, Pohl H, et al. Prevalence of clinically important histology in small adenomas. *Clin Gastroenterol Hepatol* 2006;4:343-8.
- Chen SC, Mouchli A, Chadawalada V, et al. Histopathology of small polyps removed in the videoendoscopic era [abstract]. *Gastrointest Endosc* 2006;63:AB 199.
- Jass JR, Baker K, Zlobec I, et al. Advanced colorectal polyps with the molecular and morphological features of serrated polyps and adenoma.

- mas: concept of a 'fusion' pathway to colorectal cancer. *Histopathology* 2006;49:121-31.
35. Matsumoto T, Mizuno M, Shimizu M, et al. Clinicopathological features of serrated adenoma of the colorectum: comparison with traditional adenoma. *J Clin Pathol* 1999;52:513-6.
  36. Van den Broek FJ, Van Soest EJ, Naber AH, et al. Combining autofluorescence imaging and narrow-band imaging for the differentiation of adenomas from non-neoplastic colonic polyps among experienced and non-experienced endoscopists. *Am J Gastroenterol* 2009;104:1498-507.
  37. Farris AB, Misdraji J, Srivastava A, et al. Sessile serrated adenoma: challenging discrimination from other serrated colonic polyps. *Am J Surg Pathol* 2008;32:30-5.
  38. Sandmeier D, Seelentag W, Bouzourene H. Serrated polyps of the colorectum: is sessile serrated adenoma distinguishable from hyperplastic polyp in a daily practice? *Virchows Arch* 2007;450:613-8.
  39. Mahmood T, Darzi A. The learning curve for a colonoscopy simulator in the absence of any feedback: no feedback, no learning. *Surg Endosc* 2004;18:1224-30.

#### Receive tables of content by e-mail

To receive tables of content by e-mail, sign up through our Web site at [www.giejournal.org](http://www.giejournal.org).

#### Instructions

Log on and click "Register" in the upper right-hand corner. After completing the registration process, click on "My Alerts" then "Add Table of Contents Alert." Select the specialty category "Gastroenterology" or type *Gastrointestinal Endoscopy* in the search field and click on the Journal title. The title will then appear in your "Table of Contents Alerts" list.

Alternatively, if you are logged in and have already completed the Registration process, you may add tables of contents alerts by accessing an issue of the Journal and clicking on the "Add TOC Alert" link.

You will receive an e-mail message confirming that you have been added to the mailing list. Note that tables of content e-mails will be sent when a new issue is posted to the Web.

ORIGINAL ARTICLE

## Use of Gascon and Pronase either as a pre-endoscopic drink or as targeted endoscopic flushes to improve visibility during gastroscopy: A prospective, randomized, controlled, blinded trial

P. BHANDARI<sup>1</sup>, S. GREEN<sup>1</sup>, H. HAMANAKA<sup>2</sup>, T. NAKAJIMA<sup>2</sup>, T. MATSUDA<sup>2</sup>, Y. SAITO<sup>2</sup>, I. ODA<sup>2</sup> & T. GOTODA<sup>2</sup>

<sup>1</sup>Portsmouth Hospitals Trust, Portsmouth, UK, and <sup>2</sup>The National Cancer Centre Hospital, Tokyo, Japan

### Abstract

**Objective.** To assess whether endoscopic flushes of the bubble-bursting agent Gascon and the mucolytic agent Pronase are as effective in terms of improving endoscopic mucosal visibility as a pre-endoscopic drink of the same agents. **Material and methods.** A total of 112 patients attending a Japanese tertiary referral centre for upper gastrointestinal endoscopy were randomized to receive either the standard Japanese procedure of a pre-endoscopic drink of water containing Gascon and Pronase with endoscopic flushes of 20-ml aliquots of water, or no pre-endoscopic therapy but endoscopic flushes of 20-ml aliquots of water containing Gascon, with or without Pronase as necessary. **Results.** Visibility scores were significantly better in the pre-endoscopic drink group than in either of the endoscopic flush groups. The group receiving a pre-endoscopic drink required fewer flushes during the procedure and there was no difference in the endoscopic time between the three groups. **Conclusions.** Our results suggest that endoscopic spraying of these bubble-bursting and mucolytic agents is not able to offer equivalent improvements in endoscopic mucosal visibility when compared with the standard Japanese therapy of a pre-endoscopic drink of these agents. The addition of Pronase to the spray solution had no measurable benefit over Gascon alone. We therefore cannot recommend endoscopic spraying of mucous clearing agents over their use as a pre-endoscopic drink.

**Key Words:** Endoscopy, gascon, mucolytic, pronase, simethicone, visibility

### Introduction

Since the advent of gastrointestinal endoscopy, practitioners have been frustrated by foam and mucous obscuring the field of view. Mucosal toileting techniques with bubble-bursting agents such as Gascon (simethicone) have been used since the 1950s [1–3] and more recent studies have shown that the addition of a mucolytic such as Pronase further improves mucosal visualization [4,5]. These mucosal toileting techniques have become standard practice in Japan [6,7], where cancers tend to be detected earlier than in the West. Patients there are routinely asked to drink 100 ml of water containing 2 ml of Gascon and 20,000 units of Pronase 10 min prior to the endoscopy. These medications are freely available in Europe but it is not

usual practice for them to be used. One explanation for this is concern amongst Western endoscopists of an increased risk of aspiration during the procedure if a drink is taken beforehand.

Minimally invasive techniques such as photodynamic therapy and endoscopic mucosal resection (EMR) are now able to offer excellent results for cancers detected at early stages. EMR often offers complete cure but can only be considered for tumours that are well characterized at endoscopy. Detection and characterization of early changes can be achieved through a variety of diagnostic techniques, including chromoendoscopy, high-magnification endoscopy, confocal endoscopy and narrow-band imaging, but all depend upon optimized mucosal views. In addition, chromoendoscopy requires a clear field in order

Correspondence: Susannah Green, Gastroenterology Department, Queen Alexandra Hospital, Portsmouth, PO6 3LY, UK. E-mail: susi@doctors.org.uk

(Received 16 September 2009; accepted 11 November 2009)

ISSN 0036-5521 print/ISSN 1502-7708 online © 2010 Informa UK Ltd.  
DOI: 10.3109/00365520903483643