




Figure 2. (A) A 64-year-old man who enrolled in this study and classified into Group A was referred with positive fecal occult blood test. The lesion was detected in the transverse colon at first AFI examination. (B) AFI image revealed a magenta-colored flat elevated lesion, which was macroscopically diagnosed as laterally spreading tumor, non-granular type (LST-NG). The size of the lesion was 20 mm in diameter. (C) Conventional (WL) image. (D) Chromoendoscopy image (indigo carmine). (E) Magnifying view (crystal-violet); Non-invasive (Kudo's type III and III-L pit) pattern. The final endoscopic diagnosis was HGD without submucosal invasion. We performed endoscopic submucosal dissection (ESD) using B-knife. The final histopathological diagnosis was HGD (intramucosal carcinoma).

Table 2. Detected Lesions in Each Group

		Group A (AFI-WL) (N = 83)	Group B (WL-AFI) (N = 84)	P
Total number of lesions (%)				
First exam	AFI	50 (70)	WL 52 (51)	0.01
Second exam	WL	21 (30)	AFI 50 (49)	
Neoplastic lesions (%)				
First exam	AFI	47 (71)	WL 50 (53)	0.02
Second exam	WL	19 (29)	AFI 45 (47)	
Non-neoplastic lesions no. (%)				
First exam	AFI	3 (60)	WL 2 (29)	0.28
Second exam	WL	2 (40)	AFI 5 (71)	

dye spray group was almost twice as long as for the control group (4).

Another technology recently demonstrated to be effective for detecting neoplastic lesions is NBI. The NBI system has been shown to be helpful in visualizing such lesions by improving contrast and is considered to be a new type of optical/digital chromoendoscopy (32, 33). In particular, magnification using NBI colonoscopy for the observation of the presence of "meshed brown capillary vessels" is extremely useful for distinguishing between neoplastic and non-neoplastic lesions without any dye solution. Regarding polyp detection, however, it is controversial at this moment (34). Furthermore, during NBI colonoscopy examinations, intestinal fluid was seen as being reddish in color similar to blood. Therefore, proper bowel preparation is one of the limitations when using this system.

Meanwhile, the feasibility of AFI system use for gastrointestinal (GI) screening and surveillance has not been clarified previously. In 2005, Nakaniwa *et al.* (14) developed and reported a new AFI videoscope system. Images acquired by this new AFI system provided better brightness than old fiberoptic images. From this report, the sensitivity and specificity of differentiating adenomatous and hyperplastic polyps were

Table 3. Clinicopathologic Characteristics of Lesions Detected by AFI and WL Colonoscopy

	AFI	WL	P
No. of lesions	100	73	
Location no. (%)			
Cecum	9 (9)	8 (11)	0.31
Ascending	37 (37)	19 (26)	
Transverse	54 (54)	46 (63)	
Macroscopic type no. (%)			
Polypoid	23 (23)	26 (36)	0.07
Flat elevated	77 (77)	47 (64)	
Size no. (%)			
0–5 mm	84 (84)	53 (73)	0.19
6–10 mm	10 (10)	12 (16)	
> 11 mm	6 (6)	8 (11)	
Histopathology no. (%)			
Neoplastic LGD	85 (85)	63 (86)	0.92
HGD	6 (6)	5 (7)	
Inv.ca	1 (1)	1 (1)	
Non-neoplastic	8 (8)	4 (5)	

Table 4. Characteristics of the Missed Neoplastic Lesions by AFI and WL Colonoscopy

	AFI	WL	P
No. of lesions	19	45	
Location no. (%)			
Cecum	3 (16)	2 (4)	0.13
Ascending	4 (21)	19 (42)	
Transverse	12 (63)	24 (54)	
Macroscopic type no. (%)			
Polypoid	5 (26)	6 (13)	0.21
Flat elevated	14 (74)	39 (87)	
Size no. (%)			
0–5 mm	18 (95)	41 (91)	0.62
6–10 mm	1 (5)	4 (9)	
Histopathology no. (%)			
LGD	19 (100)	45 (100)	

89% and 81%, respectively. However, there are few prospective studies that have attempted to clarify the usefulness of the adenoma detection rate using AFI system.

In this study, a total of 173 lesions from 167 patients were detected and removed endoscopically. Among these lesions, the number of neoplastic lesion detected by AFI and WL was 92 (92%) and 69 (95%), respectively. In contrast, the number of non-neoplastic lesions recognized as a polyp and removed by AFI and WL colonoscopy was only 8 (8%) and 4 (5%), respectively. The lesions we diagnosed and resected in this study with AFI and WL systems were mostly neoplastic ones. Consequently, our results evaluate the diagnostic yield of adenomatous polyp detection. However, we consider further investigation is necessary to evaluate the efficiency for differential diagnosis with AFI system.

Diminutive flat elevated lesions are thought to be of little clinical significance because such lesions, especially less than 5 mm polyps, are low-grade dysplasia (LGD) in most cases. Meanwhile, depressed lesions are considered to have a high malignant potential compared to polypoid ones in similar size (35–38). In this present study, all detected lesions' macroscopic type was flat elevated or polypoid. Because of low incidence, there were no depressed lesions in this study. However, significantly more small and/or flat neoplastic lesions were detected by AFI compared with WL colonoscopy. Therefore, AFI colonoscopy is considered to be a promising modality to detect small depressed lesions.

There are several limitations in our study. First, we conducted this study using a single experienced colonoscopist. True, our data are precise, but it is uncertain whether it would be available for all examiners. Therefore, additional multicenter studies are necessary to clarify the usefulness of AFI system for all colonoscopists. Another point worth mentioning is that our study was conducted within the limits of the right colon. The higher prevalence of flat and diminutive lesions diagnosed in the right colon may be consistent with Woolfson (39) and Hofstad's (40) description. Furthermore, a higher miss rate of detection has been reported in the right colon compared to the left colon (3). Complete back-to-back colonoscopy may be painful for patients under no sedation.

Therefore, we defined from the cecum to the splenic flexure as the target area in our prospective study. In addition, it is suggested that proper bowel preparation is indispensable to achieve success to detect small colorectal lesions. In this study, the bowel preparation was described as excellent or good in 83% and adequate but imperfect in 17%.

In conclusion, AFI videoendoscopy system is useful for the detection of right-sided colonic polyps, especially flat and/or diminutive adenomatous lesions compared to conventional (WL) colonoscopy. In the near future, multicenter trials should be performed to validate the usefulness of this system.

ACKNOWLEDGMENTS

We would like to express our appreciation to Dr. Alejandro Jimenez (Hospital Universitario de Canarias, Tenerife, Spain) for his help in the statistical analysis of this manuscript. This study was presented at the ASGE, Digestive Disease Week (DDW), Washington DC, USA, 2007.

STUDY HIGHLIGHTS

What Is Current Knowledge

- Polyps can be missed with conventional white light (WL) colonoscopy.
- Efficacy of autofluorescence imaging (AFI) system is unclear.

What Is New Here

- AFI detects more polyps, especially flat and diminutive lesions, in the right-sided colon than conventional (WL) colonoscopy.
- Prospective multicenter studies are necessary to validate the usefulness of this system.

Reprint requests and correspondence: Takahisa Matsuda, M.D., National Cancer Center Hospital, Endoscopy Division, 5-1-1, Tsukiji, Chuo-ku, Tokyo, Japan.

Received September 2, 2007; accepted February 26, 2008.

REFERENCES

- Winawer SJ, Zauber AG, O'Brien MJ, et al. Randomized comparison of surveillance intervals after colonoscopic removal of newly diagnosed adenomatous polyps. The National Polyp Study Workgroup. *N Engl J Med* 1993;328:901-6.
- Winawer SJ, Zauber AG, Ho MN, et al. The National Polyp Study Workgroup. Prevention of colorectal cancer by colonoscopic polypectomy. *N Engl J Med* 1993;329:1977-81.
- Rex DK, Cutler CS, Lemmel GT, et al. Colonoscopic miss rates of adenomas determined by back-to-back colonoscopies. *Gastroenterology* 1997;112:24-8.
- Brooker JC, Saunders BP, Shah SG, et al. Total colonic dye-spray increases the detection of diminutive adenomas during routine colonoscopy: A randomized controlled trial. *Gastrointest Endosc* 2002;56:333-8.
- Hurlstone DP, Cross SS, Slater R, et al. Detecting diminutive colorectal lesions at colonoscopy: A randomized controlled trial of pan-colonic versus targeted chromoscopy. *Gut* 2004;53:376-80.
- Rex DK, Chadlawada V, Helper DJ. Wide angle colonoscopy with a prototype instrument: Impact on miss rates and efficiency as determined by back-to-back colonoscopies. *Am J Gastroenterol* 2003;98:2000-5.
- Deenadayalu VP, Chadlawada V, Rex DK. 170 degrees wide-angle colonoscopy: effect on efficiency and miss rates. *Am J Gastroenterol* 2004;99:2138-42.
- Matsushita M, Hajiro K, Okazaki K, et al. Efficacy of total colonoscopy with a transparent cap in comparison with colonoscopy without the cap. *Endoscopy* 1998;30:444-7.
- Takehana S, Kaneko M, Mizuno H. Endoscopic diagnostic system using autofluorescence. *Diagn Ther Endosc* 1999;5:59-63.
- Panjehpour M, Overholt BF, Vo-Dinh T, et al. Endoscopic fluorescence detection of high-grade dysplasia in Barrett's esophagus. *Gastroenterology* 1996;111:93-101.
- Mayering B, Neidhardt S, Reh H, et al. Fluorescence induced with 5-aminolevulinic acid for the endoscopic detection and follow-up of esophageal lesions. *Gastrointest Endosc* 2001;54:572-8.
- Kobayashi M, Tajiri H, Seike E, et al. Detection of early gastric cancer by a real-time autofluorescence imaging system. *Cancer Lett* 2001;165:155-9.
- Ohkawa A, Miwa H, Namihisa A, et al. Diagnostic performance of light-induced fluorescence endoscopy for gastric neoplasms. *Endoscopy* 2004;36:515-21.
- Nakanishi N, Namihisa A, Ogihara T, et al. Newly developed autofluorescence imaging videoendoscopy system for the detection of colonic neoplasms. *Dig Endosc* 2005;17:235-40.
- Uedo N, Iishi H, Tatsuta M, et al. A novel videoendoscopy system by using autofluorescence and reflectance imaging for diagnosis of esophagogastric cancers. *Gastrointest Endosc* 2005;62:521-8.
- Haringmsa J, Tytgat GN. The value of fluorescence techniques in gastrointestinal endoscopy: Better than the endoscopist's eye? The European experience. *Endoscopy* 1998;30:416-8.
- Haringmsa J, Tytgat GN, Yano H, et al. Autofluorescence endoscopy: Feasibility of detection of GI neoplasms unapparent to white light endoscopy with an evolving technology. *Gastrointest Endosc* 2001;53:642-50.
- Izushi K, Tajiri H, Fujii T, et al. The histological basis on detection of adenoma and cancer in the colon by autofluorescence endoscopic imaging. *Endoscopy* 1999;31:511-6.
- Schomacker KT, Frisoli JK, Compton CC, et al. Ultraviolet laser-induced fluorescence of colonic polyps. *Gastroenterology* 1992;102:1155-60.
- Kapadia CR, Cutruzzola FW, O'Brien KM, et al. Laser-induced fluorescence spectroscopy of human colonic mucosa. Detection of adenomatous transformation. *Gastroenterology* 1990;99:150-7.
- Cothren RM, Sivak MV Jr, Van Dam J, et al. Detection of dysplasia at colonoscopy using laser-induced fluorescence: A blinded study. *Gastrointest Endosc* 1996;44:168-76.
- Uedo N, Higashino K, Ishihara R, et al. Diagnosis of colonic adenomas by new autofluorescence imaging system: A pilot study. *Dig Endosc* 2007;19:134-8.

23. Hamilton SR, Aaltonen LA, eds. World Health Organization classification of tumours: Pathology and genetics of tumours of the digestive system. Lyon, France: IARC Press, 2000;104-19.
24. Fujii T, Hasegawa RT, Saitoh Y, et al. Chromoscopy during colonoscopy. *Endoscopy* 2001;33:1036-41.
25. Kudo S, Tamura S, Nakajima T, et al. Diagnosis of colorectal tumorous lesions by magnifying endoscopy. *Gastrointest Endosc* 1996;44:8-14.
26. Hurlstone DP, Cross SS, Adam I, et al. A prospective clinicopathological and endoscopic evaluation of flat and depressed colorectal lesions in the United Kingdom. *Am J Gastroenterol* 2003;98:2543-9.
27. Adachi M, Okinaka K, Muto T. Flat adenoma of the large bowel: Re-evaluation with special reference to central depression. *Dis Colon Rectum* 2000;43:782-7.
28. Tanaka S, Haruma K, Ito M, et al. Detailed colonoscopy for detecting early superficial carcinoma: Recent developments. *J Gastroenterol* 2000;35:121-5.
29. Kiesslich R, Von Bergh M, Hahn M, et al. Chromoendoscopy with indigo carmine improves the detection of adenomatous and nonadenomatous lesions in the colon. *Endoscopy* 2001;33:1001-6.
30. Lapalus MG, Helbert T, Napoleon B, et al. Does chromoendoscopy with structure enhancement improve the colonic adenoma detection rate? *Endoscopy* 2006;38:444-8.
31. Rembacken BJ, Fujii T, Cairns A, et al. Flat and depressed colonic neoplasms: a prospective study of 1000 colonoscopies in the UK. *Lancet* 2000;355:1211-4.
32. Sano Y, Muto M, Tajiri H, et al. Optical/digital chromoendoscopy during colonoscopy using narrow-band image system. *Dig Endosc* 2005;17:43-8.
33. Chiu HM, Chang CY, Chen CC, et al. A prospective comparative study of narrow-band imaging, chromoendoscopy, and conventional colonoscopy in the diagnosis of colorectal neoplasia. *Gut* 2007;56:373-9.
34. Rex DK, Helbig CC. High yields of small and flat adenomas with high-definition colonoscopes using either white light or narrow band imaging. *Gastroenterology* 2007;133:42-7.
35. Kudo S. Colonoscopic diagnosis and management of nonpolypoid early colorectal cancer. *World J Surg* 2000;24:1081-90.
36. Saitoh Y, Waxman I, West AB, et al. Prevalence and distinctive biologic features of flat colorectal adenomas in a North American population. *Gastroenterology* 2001;120:1657-65.
37. Tsuda S, Veress B, Toth E, et al. Flat and depressed colorectal tumours in a southern Swedish population: A prospective chromoendoscopic and histopathological study. *Gut* 2002;51:350-5.
38. Soetikno R, Friedland S, Kaltenbach T, et al. Nonpolypoid (flat and depressed) colorectal neoplasms. *Gastroenterology* 2006;130:566-76.
39. Woolfson IK, Eckholdt GJ, Wetzel CR, et al. Usefulness of performing colonoscopy one year after endoscopic polypectomy. *Dis Colon Rectum* 1990;33:389-93.
40. Hofstad B, Vatn M, Larsen S, et al. Growth of colorectal polyps: Recovery and evaluation of unresected polyps of less than 10 mm, 1 year after detection. *Scand J Gastroenterol* 1994;29:640-5.

CONFLICT OF INTEREST

Even though the prototype autofluorescence imaging scope was provided by Olympus Medical Systems Corp., Tokyo, Japan, this is not a collaborative study. Thus, this study being an absolutely independent investigation, there is neither financial support nor interest from Olympus Corporation.

Asia Pacific consensus recommendations for colorectal cancer screening

J J Y Sung,¹ J Y W Lau,¹ G P Young,² Y Sano,³ H M Chiu,⁴ J S Byeon,⁵ K G Yeoh,⁶ K L Goh,⁷ J Sollano,⁸ R Rerkinmitr,⁹ T Matsuda,¹⁰ K C Wu,¹¹ S Ng,¹ S Y Leung,¹² G Makharia,¹³ V H Chong,¹⁴ K Y Ho,¹⁵ D Brooks,¹⁶ D A Lieberman,¹⁷ F K L Chan,¹ for The Asia Pacific Working Group on Colorectal Cancer

For numbered affiliations see end of article

Correspondence to: Professor J Sung, Institute of Digestive Disease, Prince of Wales Hospital, The Chinese University of Hong Kong, Hong Kong; joesung@cuhk.edu.hk

Revised 21 January 2008
Accepted 22 January 2008

ABSTRACT

Colorectal cancer (CRC) is rapidly increasing in Asia, but screening guidelines are lacking. Through reviewing the literature and regional data, and using the modified Delphi process, the Asia Pacific Working Group on Colorectal Cancer and international experts launch consensus recommendations aiming to improve the awareness of healthcare providers of the changing epidemiology and screening tests available. The incidence, anatomical distribution and mortality of CRC among Asian populations are not different compared with Western countries. There is a trend of proximal migration of colonic polyps. Flat or depressed lesions are not uncommon. Screening for CRC should be started at the age of 50 years. Male gender, smoking, obesity and family history are risk factors for colorectal neoplasia. Faecal occult blood test (FOBT), guaiac-based and immunochemical tests), flexible sigmoidoscopy and colonoscopy are recommended for CRC screening. Double-contrast barium enema and CT colonography are not preferred. In resource-limited countries, FOBT is the first choice for CRC screening. Polyps 5–9 mm in diameter should be removed endoscopically and, following a negative colonoscopy, a repeat examination should be performed in 10 years. Screening for CRC should be a national health priority in most Asian countries. Studies on barriers to CRC screening, education for the public and engagement of primary care physicians should be undertaken. There is no consensus on whether nurses should be trained to perform endoscopic procedures for screening of colorectal neoplasia.

Colorectal cancer (CRC) is one of the most common cancers in Asia and its incidence is rising in a number of Asian countries, yet there are no national or regional guidelines on prevention and screening for early diagnosis of this important disease. The Asia Pacific Working Group on Colorectal Cancer was established in 2004. The group has since conducted several studies and accumulated/published local data on neoplasm of the colon. In 2007, the Working Group members felt that it was time to review regional data on CRC and colorectal neoplasia in Asia in order to draft guidelines and recommendations in the screening and prevention of CRC in Asia.

The aim of this Consensus Conference was to draw up recommendations for CRC screening suitable for Asia.

METHOD

Membership of the consensus group

Members of the Consensus Group were selected using the following criteria: (1) demonstrated knowledge/expertise in CRC by publication/research or participation in national or regional guidelines; (2) geographical representation of the Asia Pacific countries/region; (3) diversity of views and expertise in the healthcare system (including primary care doctor, surgeon, pathologist, health economist, epidemiologist, public health expert, nurse specialists); and (4) stakeholders representing different interest groups (including healthcare policy makers, representatives from patient groups and non-government organisations). Besides members from the Asia Pacific Working Group on Colorectal Cancer, the American Cancer Society (represented by D Brooks) and the Prevent Cancer Foundation of the United States (represented by C Aldige) as well as the International Digestive Cancer Alliance and OMED (represented by G Young) were invited to participate in this conference as overseas experts. D A Lieberman was invited on his personal capacity as an advisor in this conference. The voting members are listed in Appendix A.

Literature search

Comprehensive literature reviews were carried out by the Steering Committee on a number of topics, namely (1) epidemiology of CRC in Asia; (2) colorectal polyps; (3) methods for CRC screening; (4) risk stratification; and (5) policy in CRC screening. We identified relevant articles published in English using MEDLINE, EMBASE and the Cochrane Trials Register in human subjects from 1990 to 2007. National and international guidelines on CRC screening were solicited. Searches on meeting abstracts (Asia Pacific Digestive Week (APDW), American College of Gastroenterology (ACG), American Gastroenterological Association (AGA), American Society of Gastrointestinal Endoscopy (ASGE), British Society of Gastroenterology (BSG), United European Gastroenterology Week (UEGW)) and review articles were limited to the preceding 5 years. The panel members received a copy of the relevant articles before the first iteration. The reviews were presented at the Consensus Conference before the second iteration.

Modified Delphi process

The modified Delphi process was adopted to develop the consensus.¹ The Delphi process is a method for developing consensus using a combination of the principles of evidence-based medicine and anonymous voting. After a systematic literature review, change of views from the Consensus Panel was encouraged. The process was completed by grading of evidence and anonymous voting on a series of iterations of the statements. The Steering Committee (JJYS, JYWL and FKLC) drafted a list of statements and circulated it electronically in advance to panel members. After reading the reviews, each member rated the statements on a Likert scale anchored by 1–5 (1, accept completely; 2, accept with some reservation; 3, accept with major reservation; 4, reject with reservation; 5, reject completely). All votes are anonymous. The first vote was conducted for the entire Consensus Group electronically by e-mail, without explanation or justification of the statement. Feedback of the statements was collated. The results and comments were returned to the Steering Committee before the meeting. Consensus was considered to be achieved when $\geq 80\%$ of the voting members indicated "Accept completely" or "Accept with some reservation". A statement was refuted when $\geq 80\%$ of the voting members indicated "Reject completely" or "Reject with reservation".

A face-to-face meeting of the entire group was held on 15–16 September 2007 to review the evidence of statements that reached consensus and discuss those statements that did not reach consensus on the first iteration. A series of didactic lectures presented by members reviewed the literature on five topics in colorectal neoplasia, namely (1) Epidemiology of CRC in Asia; (2) Colorectal polyps; (3) Methods for CRC screening; (4) Risk stratification; and (5) Policy in CRC screening. The statements were discussed and debated based on feedback from the first vote. The second vote was held at the end of the talks, using electronic keypads to ensure anonymity.

For statements on which consensus could not be reached, further discussions were conducted. Statements were revised accordingly. Then, the third and last vote was taken electronically using the keypads. Each statement was graded to indicate the level of evidence available and the strength of recommendation by the whole group (table 1).

Funding sources

An unrestricted education grant was received from the Olympus Medical Systems Corporation and Boston Scientific. A donation was received from the Hong Kong Cancer Fund to support the Consensus Conference. The meeting was supported in part by the KC Wong Education Foundation and the Wei Lun Foundation of the Chinese University of Hong Kong. To avoid conflict of interest, industrial partners were not allowed to participate in the discussion and iteration in the Consensus Conference. None of the sponsors voted in the drawing up of the consensus statement. Some ethnic groups (eg, Japanese, Korean and Chinese) in Asia are more susceptible than others to CRC.

RESULTS

A 2-day Consensus Conference was held on 15–16 September 2007 under the auspices of the Asia Pacific Society of Digestive Endoscopy. Representatives from 14 Asian-Pacific countries/regions participated in the meeting: these included Australia, Brunei, China, Hong Kong, India, Indonesia, Japan, Malaysia, Philippines, Singapore, South Korea, Taiwan, Thailand and

Vietnam. A total of 25 statements were presented for the first vote. Fifty members participated in the voting.

EPIDEMIOLOGY OF COLORECTAL NEOPLASIA

Statement 1. Colorectal cancer is one of the most common cancers in Asia in both males and females

Level of agreement: a, 90%; b, 10%; c, 0%; d, 0%; e, 0%

Quality of evidence: II-3

Classification of recommendation: A

Reports from the World Health Organization (WHO) data set² and individual countries or cities in Asia show that the incidence of CRC is on a rapidly rising trend in regions within countries such as China, Japan, Korea and Singapore.^{3–9} The increase in number of new cases of CRC per year is witnessed in both men and women. However, not all countries in Asia witness the same degree of rise in incidence of CRC. For example, in East Asian countries such as Indonesia, Thailand, Vietnam and India, CRC is not the top cancer in either males or females. The group also recognised that there is a lack of adequate cancer registries in many Asian countries. Without such reliable figures, some reservations remain in certain countries in indicating an epidemic of CRC in the Asia Pacific Region.

Statement 2. The incidence of CRC is similar to that of the West

Level of agreement: a, 37%; b, 47%; c, 14%; d, 2%; e, 0%

Quality of evidence: II-3

Classification of recommendation: B

The group considered that in high incidence countries such as Japan, Korea, Singapore and Hong Kong, the incidence of CRC is comparable with or approaching that of Western countries.¹⁰ Direct comparison figures are available from a study comparing Japanese with the white population of the USA which showed that the rates of CRC of these two populations were very similar.¹¹ However, such direct comparison studies are few. In other countries such as India, Philippines and Vietnam, there is still a gap in the incidence of CRC between these countries and the West. There is a strong feeling that countries with an obviously rising CRC incidence are more "Westernised" in lifestyle, especially in dietary habit, with increased consumption of high fat and protein but less fibre in the diet. The change is more evident in urban cities than in rural areas of the same country.⁷ Yet, the effects of lifestyle and dietary habit modification on the changing epidemiology of CRC in Asia need to be more adequately studied to confirm this impression.

Statement 3. The incidence of advanced neoplasm in symptomatic and asymptomatic Asians is comparable with the West

Level of agreement: a, 37%; b, 43%; c, 16%; d, 4%; e, 0%

Quality of evidence: II-2

Classification of recommendation: B

Advanced neoplasia is defined as adenoma with a diameter of ≥ 10 mm, a villous adenoma (ie, at least 25% villous), an adenoma with high grade dysplasia or invasive cancer. There are a few studies in Asian populations investigating the incidence of advanced neoplasm in asymptomatic individuals in the Asia Pacific region. A study in Hong Kong which recruited asymptomatic subjects in a Chinese population showed that 4.4% had advanced neoplasia.¹² Similar figures have been reported in a screening colonoscopy study in asymptomatic subjects among Koreans (4.1%)¹³ and Chinese (3.0%).¹⁴ Two studies that involved multiple centres in Asia that studied

Guidelines

Table 1 Quality of evidence, classification of recommendation and voting on recommendation

Category and grade	Description
Quality of evidence	
I	Evidence obtained from at least 1 RCT
II-1	Evidence obtained from well-designed control trials without randomisation
II-2	Evidence obtained from well-designed cohort or case-control study
II-3	Evidence obtained from comparison between time or places with or without intervention
III	Opinion of respected authorities, based on clinical experience and expert committees
Classification of recommendation	
A	There is good evidence to support the statement
B	There is fair evidence to support the statement
C	There is poor evidence to support the statement but recommendation made on other grounds
D	There is fair evidence to refute the statement
E	There is good evidence to refute the statement
Voting on recommendation*	
a	Accept completely
b	Accept with some reservation
c	Accept with major reservation
d	Reject with reservation
e	Reject completely

*Statements for which >80% of participants voted a, b or c were accepted. RCT, randomised controlled trial.

symptomatic and asymptomatic populations have reported the incidence of advanced neoplasia as 7.8%¹⁵ and 4.5%, respectively.¹⁶ These figures are comparable with the larger Western series using colonoscopy as a screening tool for colorectal neoplasia.¹⁷⁻²⁰ Depending on the method of recruitment, studies enrolling asymptomatic individuals for screening may introduce selection bias by recruiting more health-conscious subjects and hence underestimate the true prevalence of the conditions. This phenomenon may occur in studies from both the East and the West.

Statement 4. While the death rate of CRC is declining in the West, Asia continues to see rising mortality

Level of agreement: a, 78%; b, 20%; c, 0%; d, 2%; e, 0%

Quality of evidence: III

Classification of recommendation: C

Reports from the American Cancer Society in 2007 showed that the number of Americans who died of cancer has dropped for a second consecutive year²¹ and it was probably caused by "a combination of factors including a decrease in cigarette smoking among men, wider screening for colon cancer..."²² "By far the greatest decrease in mortality has been in colorectal cancer".²³ A similar decline in CRC mortality has been reported in Europe.²⁴ On the contrary, according to the WHO mortality database, CRC mortality has doubled in both men and women over the last three decades in Taiwan.²⁵ In Korea, the National Cancer Center reported a decline in mortality from stomach and liver cancer but an increase in CRC.²⁶ In China, the National Census Data also demonstrated a decline in mortality related to cancer of the oesophagus, and gastric and liver cancer, but the age-adjusted mortality from CRC has increased in both urban and rural men.²⁷

Statement 5. There are some ethnic groups (eg, Japanese, Korean and Chinese) in Asia who are more susceptible to CRC

Level of agreement: a, 49%; b, 43%; c, 6%; d, 0%; e, 2%

Quality of evidence: II-c

Classification of recommendation: B

Existing evidence suggests that there are some ethnic difference in susceptibility to CRC. In Singapore, the incidence of CRC is significantly lower among Indians and Malays than among Chinese.²⁸⁻²⁹ In Malaysia, the same phenomenon has been reported in a population of mixed ethnic cultures.³⁰ In the multinational studies conducted by the Asia Pacific Working Group on CRC, Japanese, Korean and Chinese were found to have a higher risk of advanced neoplasia in the colon.¹⁵⁻¹⁶ If advanced neoplasia is considered a premalignant condition, these results will infer that the incidence of CRC is higher in these ethnic groups than in the others (eg, Indians, Thais and Filipinos). The higher incidence among Chinese and the lower incidence among Indians living in the same country mirror the incidence rates in their countries of origin even though both racial groups migrated more than three generations ago. These observations on racial differences suggest that genetic factors have an important aetiological role in CRC development, although differences in dietary habit and lifestyle might also contribute. An interesting study from Israel showed that Arabs born in Israel had a much lower CRC incidence than Israeli-born Jews, and this trend persisted over time.³¹ This observation again supports the notion of genetic influence on CRC development. However, the fact that the incidence of CRC among Jews rose concomitantly with Westernisation of their lifestyle hints that environmental influences cannot be neglected.

COLORECTAL POLYPS

Statement 6. Distribution of polyps between Asians and Caucasians is similar

Level of agreement: a, 22%; b, 61%; c, 8%; d, 8%; e, 2%

Quality of evidence: II-2

Classification of recommendation: B

There are very few direct comparisons of the incidence of CRC or polyps between Asian and Caucasian populations. A study comparing Chinese in Taiwan versus Caucasians in Seattle suggested that Asians are more likely to have distally located colorectal neoplasia.³² However, the distribution of advanced neoplasia (including advanced adenoma and invasive cancer) is not significantly different between the two studied populations. Comparing three studies from Caucasian populations¹²⁻¹⁹ with four studies from Asian populations^{12-14 16 33} and one from Australia,³⁴ there are more distally located polyps in the Asia Pacific studies. In Asia, 30% of polyps are proximal, 57% are distal and 13% are synchronous. In the West, 49% of polyps are proximal, 49% are distal and 2% are synchronous. However, the distribution of advanced neoplasia is not significantly different between the East^{12 13 16 34} and the West.^{17 18 35} The proximal, distal and synchronous advanced neoplasias are 29%, 52% and 19% in Asia, and 35%, 59% and 6% in the USA (table 2). Studies from Asia showed that 53-68% of proximal advanced neoplasias were found in patients without a distal lesion. This figure is also comparable with that reported in the West. The similar distribution of colorectal polyps implies that arguments used to recommend full colonoscopy instead of flexible sigmoidoscopy in CRC screening can be applied in Asia. However, it is worth pointing out that there are some variations in the definitions of distal colonic disease in the literature. Some use findings in the last 40 cm from the anal verge on

Table 2 Distribution of advanced colorectal neoplasm (ACRN) reported in studies in Asian vs Caucasian populations

		Male (%)	Mean age (years)	ACRN			
				Total (%)	Proximal (%)	Distal (%)	Both (%)
Byeon ¹⁶	Multicentre	860 (54.8)	54.4	39 (4.5)	17 (43.6)	19 (48.7)	3 (7.7)
Chiu ¹⁴	Taiwan, China	1708 (59.8)	52.5	51 (3.0)	10 (19.6)	32 (62.7)	9 (17.6)
Liu ²³	Taiwan, China	5973 (52.3)	56.6	199 (3.3)	56 (28.1)	95 (47.7)	48 (24.1)
Sung ¹²	Hong Kong	505 (44.4)	56.5	63 (12.5)	18 (28.6)	37 (58.7)	8 (12.7)
Distribution in Asian studies							
Imperiale ¹⁸	USA	1994 (58.9)	59.8	—	50 (45.0)	61 (55.0)	—
Lieberman ¹⁷	USA	3121 (96.8)	62.9	329 (10.5%)	101 (30.7)	201 (61.1)	27 (8.2)
Imperiale ¹⁸	USA	906 (61)	44.8	32 (3.5%)	14 (43.8)	17 (53.1)	1 (3.1)
Distribution of ACRN in Caucasian studies							
					23.7–45.0	53.1–64.1	3.1–12.2

withdrawal of the colonoscope¹² and others define distal lesions as findings beyond the splenic flexure,¹⁷ or lesions in the descending colon, sigmoid colon and rectum.¹⁴ These discrepant definitions of distal colon limit the interpretation of adenoma distributions reported in the literature.

Statement 7. There is a trend towards proximal migration of polyps in the colon in Asian subjects

Level of agreement: a, 41%; b, 39%; c, 18%; d, 2%; e, 0%

Quality of evidence: III

Classification of recommendation: C

Data from the Japan Society for Cancer of the Colon and Rectum from 1974 to 1994 reviewed a right shift of CRC within a period of two decades.³⁶ The increase in the percentage of right-sided CRC was accompanied by a continuous decline in the percentage of rectal cancer in both males and females in all age groups. A single-centre retrospective cohort study in Hong Kong showed that in the last 10 years there has been an age-adjusted increasing trend of colorectal polyps in the right colon and a decrease in incidence in the left colon.³⁰ However, this study was limited by its retrospective nature and by not representing a predefined population. In Australia, a study reviewed endoscopy reports on 2578 subjects and found that 51% of polyps are right-sided, 20% are left-sided and 29% are synchronous.³⁴ The incidence of right-sided adenoma increases with age, and hence evaluation of the proximal bowel is particularly important in older people. In Japan, a cohort study enrolling 23 444 consecutive asymptomatic subjects suggested that the right shift is a phenomenon resulting from ageing.³⁷ The Japan Polyp Study also reported that more than half of the advanced neoplasias are in the right colon.³⁸ A contradictory finding was reported from Singapore.³⁹ This study showed that, from 1968 to 1992, the age-standardised rate of cancer in the distal colon was doubled in the right colon (2–3% annually) but more than doubled in the distal colon (3–4% annually). The incidence of rectal cancer was stable in Singapore. A similar observation was reported in Malaysia.⁴⁰ The wider accessibility of screening colonoscopy in some Asian countries together with the ageing population would at least partly account for the apparent increase in proximal CRC. Further studies with a long timeline will be needed to substantiate this change in epidemiology.

Statement 8. Non-polyloid adenoma is not uncommon among Asians

Level of agreement: a, 82%; b, 16%; c, 0%; d, 0%; e, 2%

Quality of evidence: II-2

Classification of recommendation: A

Flat and depressed lesions were first reported by Muto.⁴¹ In Japan, it has been reported that the prevalence of flat depressed and flat elevated lesions constituted around 3% and 18% of neoplastic lesions found on colonoscopy.⁴² Submucosal invasion was found much more commonly in flat depressed lesions compared with elevated lesions. Kudo reported that around 1.8–2.3% of colonic neoplasias are depressed lesions.⁴³ In Japan, de novo cancers—that is, cancers not arising from pre-existing adenomas, are believed to develop from these non-polyloid lesions. It has been estimated that 18.6% of CRC in men and 27.4% of CRC in women are so-called de novo cancers in Japan.⁴⁴ Over 80% of de novo cancers were invasive cancers. With the increasing awareness of these lesions, the increasing use of chromoendoscopy and new endoscopic imaging technology, there are increasing reports of flat lesions. In Singapore, 91 flat lesions were found in a cohort of 491 236 patients without using chromoendoscopy or magnifying colonoscopy.⁴⁵ In Korea, 18 flat adenomas were identified using chromoendoscopy (indigocarmine) which would have been missed by conventional colonoscopy.⁴⁶ In Malaysia, 29 adenomas were identified in 12 patients, of which 14 were flat.⁴⁷ The flat adenomas found in this study were <5 mm in size. Despite the advancement in endoscopy imaging technology, the detection of non-polyloid adenoma and de novo cancer remains a challenge. However, the necessity of discovering these small lesions is yet to be determined. Small, polyloid adenomas without villous structure or high grade dysplasia are not associated with an increased risk for CRC. Whether small flat adenomas are of greater significance remains to be determined with certainty.

Statement 9. Certain types of hyperplastic polyps are associated with an increased risk of cancer

Level of agreement: a, 80%; b, 20%; c, 0%; d, 0%; e, 0%

Quality of evidence: II-3

Classification of recommendation: A

It is well known that hyperplastic polyposis syndrome is associated with an increased potential for developing into CRC whereas a typical small and distal hyperplastic polyp (with no dysplasia) has little malignant potential. However, subsets of hyperplastic polyps are now being described and the terminology is evolving. The ability to distinguish between hyperplastic polyp, admixed hyperplastic polyp/adenoma and serrated adenoma (a form of hyperplastic polyp with propensity for progression but without distinctive cytological dysplasia) is debated among pathologists. While the majority of CRCs develop through the adenoma–carcinoma sequence with APC, K-Ras, DCC and p53 mutations, it is now clear that an admixed hyperplastic polyp or serrated adenoma may have an alternative pathway for CRC carcinogenesis. Hyperplastic polyps

Guidelines

associated with CRC may be associated with MLH1 protein, MSI and MLH1 promoter methylation.⁴⁸ BRAF mutation and aberrant promoter methylation leading to microsatellite instability and methylation instability are common in serrated adenoma and admixed polyps.^{49–50} A large, right-sided, sessile hyperplastic polyp with certain architectural features (eg, branching of crypts, dilation of base of crypts, horizontal extension of crypts, etc.) should be completely removed and carefully monitored.

Statement 10. Polyps 5–9 mm in size should be removed

Level of agreement: a, 75%; b, 20%; c, 2%; d, 3%; e, 0%

Quality of evidence: III

Classification of recommendation: C

Reports from the early 1990s showed that screening sigmoidoscopy and removal of the polyp reduced CRC mortality.^{51–52} The National Polyp Study which included patients from the USA and the UK,⁵³ and the Italian Multicenter Study⁵⁴ provided the strongest evidence that removal of polyps reduced the risk of subsequent CRC. Recently, long-term follow-up figures from the National Polyp Study showed a reduction in mortality after polyps were removed during screening colonoscopy.⁵⁵ The Japan Polyp Study Group conducted a retrospective study of a cohort of 5309 subjects who underwent colonoscopy from 1990 to 1995 and followed-up for >3 years. In this period, polyps larger than 6 mm were removed.⁵⁶ The cumulative hazard of developing malignant disease for those who had a polyp <5 mm was comparable with that for those who had no polyp found on index colonoscopy. On the other hand, those with polyps measuring 6–9 mm had a cumulative hazard of developing invasive cancer comparable with those with intramucosal cancer.⁵⁸ These are important data from Asia which lend support to the removal of polyps 5–9 mm in size. The Japan Polyp Study Group is now conducting a study randomising polypectomised patients to be followed either at 3 years, or at 1 year and then 3 years to study the outcome of such surveillance intervals in the context of finding new colonic lesions.

SCREENING TESTS FOR COLORECTAL NEOPLASIA

Statement 11. Faecal occult blood test (FOBT; guaiac-based and immunochemical tests), flexible sigmoidoscopy and colonoscopy should be recommended for CRC screening

Level of agreement: a, 74%; b, 18%; c, 6%; d, 2%; e, 0%

Quality of evidence: I

Classification of recommendation: A

FOBT, flexible sigmoidoscopy and colonoscopy are recommended options for CRC screening in national guidelines from the USA, the UK and Canada.^{57–60} Annual or biennial screening with FOBT using a guaiac-based test or an immunochemical test has been shown to reduce both CRC and CRC-related mortality compared with no screening.^{61–63} Although the sensitivity of a single FOBT is low, in the range of 30–50%, repeated annual testing can detect as many as 92% of CRCs. It is perceived that FOBT is a “cancer test” instead of a test for polyps or adenoma. The advantage is that FOBT can be done at home and is non-invasive, but the test needs to be repeated every 1–2 years. Rehydration of the stool sample is not recommended. Although rehydration of the guaiac-based test increases sensitivity, the false-positive rate is also raised, leading to unnecessary anxiety and unnecessary performance of invasive tests. An immunochemical test may obviate the need for dietary restriction.^{65–66} Recently, faecal immunochemical

tests for haemoglobin have been shown to be more sensitive than the guaiac test for cancer and adenomas especially in Asian subjects, probably due to lack of dietary interference.⁶⁶

Flexible sigmoidoscopy performed every 5 years has been shown in case-controlled studies to reduce mortality from CRC.^{51–52} The preliminary findings of a randomised controlled trial of screening flexible sigmoidoscopy have been reported, but the result in terms of effectiveness on an intention-to-screen basis at the population level is not yet available.⁶⁷ The recommendation of a 5 year interval was based on a cohort study which showed that 5 years after a negative colonoscopy, new advanced neoplasias are rare.⁶⁸ The sensitivity of flexible sigmoidoscopy in detecting advanced neoplasia is reported to be 35–70% and reduced the cancer risk in the rectum and sigmoid by 50–60%.^{17–19–69} The recommended interval of screening is shorter than for colonoscopy because flexible sigmoidoscopy is less sensitive than colonoscopy even in the distal colon. This is because of the quality of bowel preparation, the varied experience of the examiners and the discomfort, which leads to colonic spasm which may affect the depth of sigmoidoscopy insertion and hence the adequacy of the examination. Since up to two-thirds of proximal advanced lesions in Asians are found in the absence of distal lesions, the disadvantage of creating a false sense of security using flexible sigmoidoscopy for screening is noted.

The use of colonoscopy for screening is not supported by a randomised controlled study but by indirect evidence. The National Polyp Study has demonstrated a reduced incidence of CRC⁵³ and recently a reduced mortality from CRC among those who underwent colonoscopy.⁵⁵ Colonoscopy is the only modality that allows removal of the adenoma and prevents CRC. A similar study in Europe has confirmed the benefit of the screening procedure.⁵⁴ The effectiveness of colonoscopy is dependent on the quality of the examination (see below).

Statement 12. Double-contrast barium enema (DCBE) is not a preferred CRC screening test

Level of recommendation: a, 78%; b, 20%; c, 0%; d, 2%; e, 0%

Quality of evidence: III

Classification of recommendation: C

DCBE every 5 years is listed as one of the options in CRC screening in national guidelines in North America. Like colonoscopy, there is no randomised trial evaluating whether screening DCBE reduces the incidence or mortality of CRC in the average-risk population, and there has been no actual report using barium enema in a true screening environment. The sensitivity of DCBE is lower than that of colonoscopy and it does not permit removal of polyps or biopsy of cancers. In a study comparing DCBE and colonoscopy, the sensitivity of DCBE for lesions >10 mm was 48% and for lesions of 6–9 mm it was 35%.⁷⁰ In the National Polyp Study, DCBE detected only 53% of adenomatous polyps 6–10 mm in size and 48% of those >10 mm in size compared with colonoscopy.⁷¹ Because of its lower sensitivity, even for large polyps, the Consensus Group does not recommend DCBE as a first-line option for CRC screening.

Statement 13. CT colonography is not currently a preferred CRC screening test

Level of recommendation: a, 90%; b, 8%; c, 0%; d, 2%; e, 0%

Quality of evidence: III

Classification of recommendation: C

Unlike DCBE, there is increasing evidence to suggest that CT colonography is an accurate screening method for the detection of colorectal neoplasia in asymptomatic average-risk adults.^{73 78} The sensitivity and specificity of the findings are also dependent on the size of the polyps. Meta-analysis showed that sensitivity is around 85% for polyps >9 mm, 70% for polyps 6–9 mm, and 50% for polyps <6 mm.^{74 75} Studies have also shown that large size (≥10 mm in size) is the best prediction of advanced neoplasia.⁷⁶ According to this study, high grade dysplasia and invasive cancer is very uncommon in medium sized (6–9 mm) lesions, which justifies the use of size alone as a surrogate measure for predicting advanced histological features. Since it is not clear whether small polyps should be removed by polypectomy, some radiologists recommended that polyps <5 mm in size should not be reported. Patients with polyps of 6–9 mm should have repeat CT at 1–2 yearly intervals.⁷⁷ Patients with polyps that are 6–9 mm can have a repeat CT colonography in 3 years, and polyps <6 mm need not be reported. In the literature, however, there is a discrepancy in the results of CT colonography as a result of a difference in CT collimation width, type of scanner and mode of imaging. The use of a multidetector scanner equipped with 3-D flythrough views that simulate colonoscopy may increase the sensitivity of CT colonography. While in the expert centre, a randomised trial of CT colonography compares favourably with conventional colonoscopy,⁷¹ the results in non-expert centres are less promising.⁷⁴ Furthermore, acceptability in a true screening population has not been fully explored. High cost, risk associated with radiation and requirement for bowel preparation are the other factors hindering the use of CT colonography as a primary screening method at this stage. In view of the inaccessibility of cutting-edge imaging technology in some Asian countries, the Consensus Group does not recommend CT colonography as a CRC screening tool at this stage. However, the group believes that with increased accessibility, CT colonography may become a recommended tool for CRC screening in the future.

Statement 14. In resource-limited countries, FOBT is the first choice for CRC screening

Level of recommendation: a, 72%; b, 18%; c, 6%; d, 4%; e, 0%

Quality of evidence: I

Classification of recommendation: C

FOBTs are used in screening to refine the likelihood of cancer being present and so direct scarce colonoscopy resources to those more likely to have neoplasia.⁷⁹ A large-scale case-control study in Japan using immunochemical FOBT has shown a decrease in CRC mortality by 70%.⁸⁰ This benefit was witnessed in both men and women in the cohort. Although the overall rate of incidence of CRC has not been significantly reduced, a reduction in advanced CRC was reported. A study from Australia showed that CRC screening by FOBT is cost-effective and comparable with other cancer screening programmes.⁸¹ More cost-effectiveness studies need to be done in Asia. Despite the fact that FOBTs (guaiac or immunochemical tests) are not diagnostically precise, many Western countries consider that they are the best approach to population screening because of their simplicity and high acceptance by asymptomatic subjects even in countries with a well-developed healthcare system. Clearly, in resource-limited countries in Asia, in order to have a population impact, FOBT is the most affordable test.

Statement 15. Following a negative colonoscopy, a repeat examination should be performed in 10 years

Level of recommendation: a, 27%; b, 53%; c, 8%; d, 12%; e, 0%

Quality of evidence: II-3

Classification of recommendation: C

The choice of a 10-year interval between screening examinations for average-risk subjects after a negative colonoscopy is based on estimates of the sensitivity of colonoscopy and the rate at which advanced neoplasia develops.^{82 83} Colonoscopy is not perfect and it can still miss colorectal adenoma or even cancer. The rate of new or missed CRC within 3 years after colonoscopy has been reported as around 5% in the proximal colon and around 2% in the distal colon.⁸⁴ The chance of missing a diagnosis is higher in older subjects, those with diverticular disease, right-sided or transverse lesions, suboptimal bowel preparation and when colonoscopy is performed by internists or family doctors in their office. A large series of colonoscopy screening showed that 0.3–0.9% of CRC can be missed.^{84–86} These so-called interval cancers after colonoscopy could be due to genuinely new and fast growing lesions,⁸⁷ incomplete removal of polyps⁸⁸ or missed lesions. Withdrawal time during colonoscopy is found to correlate with adenoma detection during screening colonoscopy.^{88 89} In essence, the impact and success of colonoscopy screening depend on the quality of the procedure. The potential benefit and risk of screening change in elderly patients of different life expectancies and the age for stopping screening should be considered.⁹⁰ Even though the prevalence of neoplasia increases with age, screening in elderly persons >80 years of age results in only a modest gain in life expectancy and thus may not be desirable.⁹¹

RISK STRATIFICATION IN CRC SCREENING

Statement 16. The age-adjusted incidence of CRC is higher in men than in women

Level of agreement: a, 82%; b, 16%; c, 0%; d, 0%; e, 2%

Quality of evidence: II-2

Classification of recommendation: A

In many Asian countries, the age-adjusted incidence of CRC is found to be higher in men than in women.^{8–10} While the exact mechanism of the hormonal effect on colorectal neoplasia is still unclear, in a prospective Japanese study pregnancy was found to be associated with reduced risk of CRC in women.⁹² It is postulated that female sex hormones reduce the risk of CRC. Indeed, this observation was also reported in the VA Cooperative Study 380 and suggested that women may start screening at a later age because of their relatively low incidence of colorectal neoplasia at the age of 50–55 years.¹⁹ The fact that the age-adjusted incidence in CRC is lower in women, however, does not imply that screening is less effective in women. In a large population-based Japanese cohort study using FOBT-selected cases for colonoscopy screening, mortality reduction was achieved in both men and women.⁸⁰

Statement 17. CRC screening should begin at the age of 50

Level of agreement: a, 35%; b, 57%; c, 6%; d, 2%; e, 0%

Quality of evidence: II-2

Classification of recommendation: B

The prevalence of colorectal neoplasia increases with age. As the risk of CRC starts to escalate at the age of 50 years, most national guidelines recommend that screening programmes should begin by this age.^{57–60} Screening colonoscopy studies in Asia also confirm that at the age of 50 the risk of finding advanced neoplasia is significantly increased from around 1% to

Guidelines

>3%.¹²⁻¹⁵ Therefore, the Consensus Group showed strong support for starting CRC screening at the age of 50 years. From the large population-based cohort study in Japan using FOBT for CRC screening, the best age to start screening was 50-59 years.⁷⁸ The group also noted that in each age group, however, other factors such as gender, family history and race may affect the outcome of CRC screening. Since Asia represents a very heterogeneous population, it is desirable to have a formula stratifying the risk according to age, gender, race and family history to select those who have the highest CRC risk for priority in a screening programme. A risk stratification strategy using colonoscopy and CT colonography has been described in the West.⁷⁹⁻⁸⁴ This would be a very useful way of making the best use of limited resources in many Asian countries. A two-tier approach has been proposed in Taiwan, which may reduce the workload on colonoscopy without jeopardising the efficacy of screening.⁷⁹ This kind of study would need further validation.

Statement 18. First-degree relatives of patients with CRC are at an increased risk and thus should receive screening earlier

Level of agreement: a, 78%; b, 20%; c, 0%; d, 2%; e, 0%

Quality of evidence: III

Classification of recommendation: C

A prospective study showed that among first-degree relatives of CRC patients, the age-adjusted risk of CRC was 1.72.⁸⁶ With two or more first-degree relatives, the risk is further escalated. For those under the age of 45 who had one or more affected first-degree relatives, the relative risk was increased by more than fivefold. Colonoscopy screening in first-degree relatives of patients with sporadic CRC has been demonstrated to yield a higher rate of finding colorectal neoplasia.⁹⁷ The odds ratios (ORs) reported were 1.5 for adenoma, 2.5 for large adenoma and 2.6 for high risk adenoma. A study from Italy also showed that compared with subjects with no family history, asymptomatic patients with one first-degree relative with CRC had nearly double the risk of developing adenomatous polyps.⁹⁸ A meta-analysis of pooled data from 27 case-control studies indicates that the first-degree relatives of a patient with CRC have an increased risk of colon cancer of 2.42 and of rectal cancer of 1.89.⁹⁹ This applies to both parents and siblings suffering from the disease. This phenomenon is also observed in a study from Taiwan in which 234 immediate family members of 186 CRC patients were screened.¹⁰⁰ The immediate family members were at increased risk for advanced neoplasia, with an OR of 4.5. Individuals with index cancer relatives diagnosed at <50 years or male relatives were found to have an even higher risk of advanced neoplasia.

Statement 19. Smoking increases the risk of CRC

Level of agreement: a, 51%; b, 31%; c, 12%; d, 4%; e, 2%

Quality of evidence: II-2

Classification of recommendation: B

There are studies in Asia, especially in Japan, China and Singapore, investigating the effects of cigarette smoking and the risk of CRC. A case-control study in Japan showed smoking in the past 10 years is significantly associated with risk of sigmoid and rectal adenoma.¹⁰¹ A larger and more recent study from Japan confirms the association of smoking and CRC.¹⁰² The effect of smoking has been observed to be related to the number of cigarettes consumed and the age of starting smoking.¹⁰³ Current smokers have a higher risk than ex-smokers, and men and women were equally affected. It has been estimated that approximately half of the CRC cases in Japan can be prevented

by tobacco and alcohol control in middle-aged and elderly Japanese men. The relationship of smoking and alcohol consumption to CRC was studied in Chinese living in Singapore.¹⁰⁴ In this population-based study, cigarette smoking was associated with an increased risk of rectal but not colonic cancer. Compared with non-smokers, light smokers have an increased risk of 1.43 and heavy smokers of 2.64 of developing rectal cancers. Smoking appears to interact with alcohol consumption in an additive manner in affecting the risk of rectal cancer.

Statement 20. Obesity increases the risk of CRC

Level of agreement: a, 47%

Quality of evidence: II-2

Classification of recommendation: A

Obesity has been found to increase the risk of CRC. A meta-analysis reported a relative risk of CRC of 1.37 for overweight and obese men, and the associated risk appears to be higher for men than for women.¹⁰⁵ Epidemiological studies have also shown in a Korean population that patients with metabolic syndrome had an increased risk of colorectal adenoma (OR 1.51).¹⁰⁶ The association with metabolic syndrome was more evident for proximal, multiple (>3) and advanced adenoma. In Japan, a nationwide prospective study which included >43 000 women and 58 000 men aged 40-70 showed, after adjustment for the lifestyle factors, a significant positive correlation of CRC with baseline body weight.¹⁰⁷ Women with baseline body mass index (BMI) >28 kg/m² had a relative risk of 2.4 for CRC compared with those with a BMI of 20-22 kg/m². BMI was also found to have a positive correlation with adenoma of the colon in Japan and Korea.¹⁰⁸⁻¹⁰⁹ This trend has not been demonstrated in men. Hyperinsulinaemia may be an important factor, but the role of oxidative stress initiated by hyperglycaemia is another possible mechanism. The lack of physical activity leading to overweight has been identified as a risk factor for CRC in a study from Shanghai.¹¹⁰ A recent study from Hong Kong also showed that those who underwent investigations for coronary heart disease are more likely to have CRC.¹¹¹ Metabolic syndrome among this group of patients is an independent risk factor for the condition.

Statement 21. Screening for CRC should be a national health priority in most Asian countries

Level of agreement: a, 57%; b, 33%; c, 8%; d, 2%; e, 0%

Quality of evidence: III

Classification of recommendation: C

In North America and Europe, as well as Australia and New Zealand, there is a widespread scientific agreement on the value of CRC screening. CRC screening is endorsed by the American Cancer Society, the US Preventive Services Task Force, the Multi-Society Taskforce on Colorectal Cancer, the American College of Gastroenterology, the American Gastroenterological Association, the American Society of Gastrointestinal Endoscopy, the American College of Physicians, the British Society of Gastroenterology, the Canadian Task Force on Preventive Health Care and the Royal Colleges of Physicians in the UK, to name just a few.⁵⁷⁻⁶⁰ Screening is available in different settings, but the vast majority of screening activities are still opportunistic and uncoordinated. In certain localities, large healthcare systems (e.g. Kaiser Permanente of North California and the Veteran's Administration) have developed organised screening for their populations. Local programmes are also found in other regions (eg, the State of Maryland and New

York City). In the USA, Medicare added coverage for CRC screening in 2000. Laws have been enacted in nearly half of the states in the USA requiring private insurers to pay for CRC screening tests. Legislation was introduced in the US Congress in 2007 to create a CRC screening, diagnostic and treatment programme for poor and/or uninsured citizens. In Europe, a public health programme has developed a comprehensive set of recommendations for cancer screening, and CRC screening was added in 2003. In a recent survey conducted by the International Digestive Cancer Alliance (IDCA) across Europe, 21 out of 39 nations have reported national screening guidelines promoted by medical and professional organisations.¹¹² Fifteen countries are currently performing some form of population screening programmes and seven others have feasibility studies underway. Respondents from 20 countries where screening is not taking place indicated a lack of official recognition of the importance and value of CRC screening. Lack of financial support is identified as the primary barrier to screening. Germany is the European country with the largest screening programme using guaiac-based FOBT. Since 2002, >2.1 million screening colonoscopies have been performed, detecting adenoma in about 20% of subjects and CRC in 0.6–0.8% of subjects.

In Asia, a national guideline is available only in Australia, Japan, Korea, Taiwan and Singapore.^{118–117} Despite these guidelines, the uptake of CRC screening is relatively low. In Korea, the government covers 50% of the cost of CRC screening and 100% for low-income individuals. Taiwan is the only country with free mass screening for CRC under the national health insurance scheme. CRC screening is endorsed but not funded in most Asian countries. In countries such as Brunei, China, India, Indonesia, Malaysia, Philippines, Thailand and Vietnam, a national guideline is not available. Government support for CRC screening is very limited. The Consensus Group urges strong support from Asian health authorities to promote CRC screening in the Asia Pacific region.

Statement 22. Research on barriers to CRC screening should be conducted in various Asian countries

Level of agreement: a, 86%; b, 14%; c, 0%; d, 0%; e, 0%

Quality of evidence: II-3

Classification of recommendation: B

Adherence to guidelines on screening is low even in Western countries. In Japan, around 17% of the eligible population participates in the immunochemical FOBT.¹¹⁸ In Canada, 23.5% of eligible respondents received screening.¹¹⁹ In the USA, the compliance rate was low and has only risen to 40–60% in recent years.¹²⁰ Only a limited number of studies have investigated the factors that play a major role in compliance/non-compliance with colon screen advice, and they have yielded somewhat inconsistent findings. A recent study from Hong Kong employed the Health Belief Model to study the knowledge, behavioural and psychological obstacles to CRC screening tests.¹²¹ Knowledge of CRC symptoms and risk factors, recommendation by a doctor and the availability of health insurance are positively associated with uptake of screening tests. On the other hand, health, psychological and access barriers, and perceived negative personal and family consequences of CRC are negatively associated with uptake of the screening test. The Asia Pacific Working Group on CRC is undertaking a similar study to compare the health-seeking behaviour and obstacles to screening tests in different cultures. This kind of study will provide important information for the successful implementation of CRC screening in the region.

Statement 23. Education of the public is essential in promoting CRC screening

Level of agreement: a, 96%; b, 4%; c, 0%; d, 0%; e, 0%

Quality of evidence: I

Classification of recommendation: A

In most Asian societies, public knowledge of CRC is poor and uptake of screening tests is expected to be low. Fewer than 10% of the Hong Kong Chinese are aware that CRC is the second most common cancer in their locality.¹²² Most Chinese believe that "screening" is needed only when they develop symptoms of cancer. Only one-third of the Singaporean Chinese know that they should go for screening even if asymptomatic.¹²³ Over 70% of individuals cannot name a single screening method for CRC. In both of these studies, there was little recommendation offered to intervene in existing low awareness and willingness to participate in CRC screening. A population survey suggested that male subjects above 50 years of age were significantly deficient in knowledge of CRC symptoms and the perceived benefits of screening.¹²¹ Educating this group would be important as they are the ones who may benefit from CRC screening.

Statement 24. Family doctors should be engaged in promoting CRC screening

Level of agreement: a, 82%; b, 18%; c, 0%; d, 0%; e, 0%

Quality of evidence: I

Classification of recommendation: A

Family doctors play a pivotal role in recommending asymptomatic individuals for CRC screening. A study in Iowa in the USA shows that the strongest predictors of patient's compliance with CRC screening, other than symptoms, were patient recollection of a doctor's recommendation and documentation by the doctor of advice to their patients.¹²⁴ In two population surveys in Hong Kong, a recommendation from the family doctor was found to have the highest impact on the patients' compliance with CRC screening.¹²¹ Recommendation by a doctor increases the likelihood of having a CRC screening test by 21 times.¹²¹ Why are some family doctors not interested in recommending CRC screening? Previous studies have shown that lack of knowledge and training, lack of time and opportunity, lack of financial support for the patients in participating in screening and inconsistency in recommendations are the most important reasons for their reluctance to advise their patients.¹²⁵ Education and training for family doctors should be an effective strategy to promote CRC screening.

Statement 25. Nurses in Asia should be trained to perform flexible sigmoidoscopy for CRC screening

Level of agreement: a, 6%; b, 4%; c, 30%; d, 30%; e, 30%

No consensus reached

The Consensus Group noted that in the UK, the USA and Canada, there are nurse-run flexible sigmoidoscopy programmes.^{126–128} There is at least one nurse practitioner-directed colonoscopy programme in the USA (Alaska). The advantages of recruiting nurses to perform endoscopy are to speed up the process of CRC screening and to relieve the endoscopy workload. The enhanced contributions of nurses will also strengthen the nursing profession to become more competent and independent. A stringent training programme and other academic qualifications may ensure the competence of nurse endoscopists. In order to implement nurse-led CRC screening by flexible sigmoidoscopy or colonoscopy, due recognition by the

Guidelines

nursing council and proper licensing criteria will be required. Among the members of the Consensus Group, there was a very divergent view. Issues of liability, third-party reimbursement, lack of medical support, lack of policies and guidelines on this have been discussed. The majority of members have reservations on this issue and a consensus view could not be reached in the meeting. However, despite the hurdles, the Consensus Group believes that other than endoscopy, nurses can play a very important role in patient education, coordination of service and tracking individuals tested positive by FOBT.

CONCLUSION

These are the first Asia Pacific Consensus statements formulated based on evidence in the literature, national registries and local data, input from international experts and thorough discussion among members of the Asia Pacific Working Group for Colorectal Cancer. It provides a basis for further elaboration and modification to suit the needs of each individual Asia Pacific country/region. There are areas which cannot be covered in the present statements. These include the recommendations for hereditary CRC and genetic counselling policy, lifestyle risk factors and intervention, cultural differences in health-seeking behaviour, among others. Future studies in the Asia Pacific region should aim at investigating the effects of culture on compliance with CRC screening, the effects of genetic and environmental factors on CRC development and the practical use of guaiac-based and immunochemical FOBTs in different Asian populations.

Author affiliations: ¹The Chinese University of Hong Kong, Prince of Wales Hospital, Shatin, NT, Hong Kong; ²Department of Medicine, School of Medicine, Flinders University, Adelaide, Australia; ³Sano Hospital, 2-5-1 Shimizu-cho, Kobe, Hyogo, Japan; ⁴Department of Internal Medicine, Health Management Center, National Taiwan University Hospital, Taipei, Taiwan; ⁵Division of Gastroenterology, Department of Internal Medicine, Asan Medical Center, Songpa-gu Seoul, Korea; ⁶Clinical Research Center, National University of Singapore, Singapore; ⁷Division of Gastroenterology, Department of Medicine, University of Malaya, Kuala Lumpur, Malaysia; ⁸Department of Medicine, University of Santo Tomas, Manila, Philippines; ⁹Gastroenterology Unit, Department of Internal Medicine, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand; ¹⁰Division of Endoscopy, The National Cancer Center Hospital, Chuo-ku, Tokyo, Japan; ¹¹Department of Gastroenterology, Xijing Hospital, Xi'an, Shaanxi, China; ¹²Department of Pathology, The University of Hong Kong, Queen Mary Hospital, Hong Kong; ¹³Department of Gastroenterology and Human Nutrition, All India Institute of Medical Sciences, Ansari Nagar, New Delhi, India; ¹⁴Gastroenterology Unit, Department of Medicine, Raja Isteri Pengiran Anak Saleha (RIPAS) Hospital, Bandar Seri Begawan, Negara Brunei Darussalam; ¹⁵Department of Medicine, Yong Loo Lin School of Medicine, National University of Singapore, Singapore; ¹⁶American Cancer Society, Dallas, Texas, USA; ¹⁷Division of Gastroenterology, Oregon Health and Science University, Portland VA Medical Center, Portland, Oregon, USA

Competing interests: None.

REFERENCES

1. **Linstone H, Turfitt M.** The Delphi method: techniques and application. <http://www.is.nyu.edu/pubs/delphibook/> (accessed 19 May 2008).
2. **Ferlay J, Bray F, Pisani P, et al.** *GLOBALCAN 2002: Cancer incidence, mortality and prevalence worldwide, version 2.0.* IARC CancerBase No. 5. Lyon: IARC Press, 2004.
3. **Yiu HY, Whittemore AS, Shibata A.** Increasing colorectal cancer incidence rate in Japan. *Int J Cancer* 2004;**109**:777–81.
4. **Tamura K, Ishiguro S, Munakata A, et al.** Annual changes in colorectal carcinoma incidence in Japan: analysis of survey data on incidence in Aomori Prefecture. *Cancer* 1996;**78**:1187–94.
5. **Chia KS, Lee HP, Seow A, et al.** Trends in cancer incidence in Singapore 1969–1992. *Singapore Cancer Registry Report* no. 4. Singapore: Singapore Cancer Registry, 1995.
6. **Hong Kong Cancer Registry.** Hospital Authority of Hong Kong Special Administrative Region. Available at <http://www3.ha.org.hk/cancerreg/> (accessed 19 May 2008).
7. **Jim BT, Devesa SS, Chow WH, et al.** Colorectal cancer incidence trends by subtype in urban Shanghai 1972–1994. *Cancer Epidemiol Biomarkers Prev* 1998;**7**:661–6.
8. **Lu JB, Sun XG, Dai DX, et al.** Epidemiology of gastroenterological cancer in Henan Province, China. *World J Gastroenterol* 2003;**9**:2400–3.
9. **Yang L, Parkin DM, Li LD, et al.** Estimation and projection of the national profile of cancer mortality in China: 1985 to 2005. *Br J Cancer* 2004;**90**:2157–66.
10. **Sung JJY, Lau JYW, Goh KL, Leung WK, Asia Pacific Working Group on Colorectal Cancer.** Increasing incidence of colorectal cancer in Asia: implications for screening. *Lancet Oncol* 2005;**6**:871–6.
11. **Yiu HY, Whittemore AS, Shibata A.** Increasing colorectal cancer incidence rate in Japan. *Int J Cancer* 2004;**109**:777–81.
12. **Sung JJY, Fok Chan, Leung WK, et al.** Screening for colorectal neoplasms in Chinese: faecal occult blood test, flexible sigmoidoscopy or colonoscopy? *Gastroenterology* 2003;**124**:608–14.
13. **Choe JW, Chang HS, Yang SK, et al.** Screening colonoscopy in asymptomatic average-risk Koreans: analysis in relation to age and sex. *J Gastroenterol Hepatol* 2007;**22**:1003–8.
14. **Chiu HM, Wang HP, Lee YC, et al.** A prospective study of the frequency and topographic distribution of colon neoplasia in asymptomatic average-risk Chinese adults as determined by colonoscopic screening. *Gastrointest Endosc* 2005;**61**:547–53.
15. **Leung WK, Ho KY, Kim WH, et al.** Asia Pacific Working Group on Colorectal Cancer. Colorectal neoplasia in Asia: a multicenter colonoscopy survey in symptomatic patients. *Gastrointest Endosc* 2006;**64**:751–9.
16. **Byeon JS, Yang SK, Kim TI, et al.** Asia Pacific Working Group on Colorectal Cancer. Colorectal neoplasia in asymptomatic Asians: a prospective multinational multicenter colonoscopy survey. *Gastrointest Endosc* 2007;**65**:1015–22.
17. **Lieberman DA, Weiss DG, Bond JH, Ahnen DJ, Garewal H, Cheifetz G, for Veterans Affairs Cooperative Study Group 380.** Use of colonoscopy to screen asymptomatic adults for colorectal cancer. *N Engl J Med* 2000;**343**:162–8.
18. **Imperiale TF, Wagner DR, Lin CY, et al.** Risk of advanced proximal neoplasm in asymptomatic adults according to the distal colorectal findings. *N Engl J Med* 2000;**343**:169–74.
19. **Schoenfeld P, Cash B, Flood A, et al.** CONCERN Study Investigators. Colonoscopic screening of average-risk women for colorectal neoplasia. *N Engl J Med* 2005;**352**:2061–8.
20. **Bates M, Munoz-Navas MA, Duque JM, et al.** Use of colonoscopy as a primary screening test for colorectal cancer in average risk people. *Am J Gastroenterol* 2003;**98**:2648–54.
21. **American Cancer Society.** Statistics for 2007. http://www.cancer.org/docroot/sty/sty_0.asp (accessed 19 May 2008).
22. **Stein R.** Cancer death decline for second straight year: few smokers, more screening credited. *Washington Post* 18 Jan 2007.
23. **Grady D.** Second drop in cancer deaths could point to a trend, researchers say. *The New York Times* 18 Jan 2007.
24. **Janout V, Kolarova H.** Epidemiology of colorectal cancer. *Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub* 2001;**145**:5–10.
25. **World Health Organization.** World Health Statistics Annual. World Health Organization (WHO) Database, Geneva, Switzerland. Available at <http://www.who.int/whodoc/whodoc.htm> (accessed 21 Jan 2008).
26. **Bae JM, Jung KW, Won YJ.** Estimation of cancer death in Korea for the upcoming years. *J Korean Med Sci* 2002;**17**:611–5.
27. **Yang L, Parkin DM, Li LD, et al.** Time trends in cancer mortality in China: 1987–1999. *Int J Cancer* 2003;**106**:771–783.
28. **Wang H, Seow A, Lee HP.** Trends in cancer incidence among Singapore Malays: a low-risk population. *Ann Acad Med Singapore* 2004;**33**:57–62.
29. **Lee HP, Lee J.** Shanmugaratnam K. Trends and ethnic variation in incidence and mortality from cancers of the colon and rectum in Singapore 1968 to 1982. *Ann Acad Med Singapore* 1987;**16**:397–401.
30. **Lim GCC, Lim TO, Yahaya H, eds.** *The first report of the National Cancer Registry: cancer incidence in Malaysia 2002.* Malaysia: National Cancer Registry, 2002.
31. **Fremgen Z, Neiman E, Abu Mouch S, et al.** Trends in incidence of colorectal cancer in Jewish and Arab populations in central Israel. *Digestion* 2005;**72**:223–7.
32. **Soon MS, Kozarek RA, Ayub K, et al.** Screening colonoscopy in Chinese and Western patients. *Am J Gastroenterol* 2005;**100**:2749–55.
33. **Li HH, Wu MC, Peng Y, et al.** Prevalence of advanced colonic polyps in asymptomatic Chinese. *World J Gastroenterol* 2005;**11**:4731–4.
34. **Patel K, Hoffman NE.** The anatomical distribution of colorectal polyps at colonoscopy. *J Clin Gastroenterol* 2001;**33**:222–5.
35. **Imperiale TF, Wagner DR, Lin CY, et al.** Results of screening colonoscopy among persons 40 to 49 years of age. *N Engl J Med* 2002;**346**:1781–5.
36. **Takada H, Ohnawa A, Iwamoto S, et al.** Changing site distribution of colorectal cancer in Japan. *Dis Colon Rectum* 2002;**45**:1249–54.
37. **Yamaji Y, Mitsuhashi T, Ikuma H, et al.** Right side shift of colorectal adenoma with ageing. *Gastrointest Endosc* 2006;**63**:453–8.
38. **Fujii T, Sano Y, Iishi H, et al.** Colorectal cancer screening in Japan: results of the multicenter retrospective cohort study. *Gastroenterology* 2002;**122**:A481.
39. **Huang J, Seow A, Shi CY, et al.** Colorectal carcinoma among ethnic Chinese in Singapore: trends in incidence rate by anatomic subsites from 1968 to 1992. *Cancer* 1999;**85**:2519–25.
40. **Goh KL, Quek KF, Yeo GTS, et al.** Colorectal cancer in Asians: a demographic and anatomic survey in Malaysian patients undergoing colonoscopy. *Aliment Pharmacol Ther* 2005;**22**:859–64.
41. **Muto T, Kamiya J, Sawada T, et al.** Small flat adenomas of the large bowel with special reference to its clinicopathologic features. *Dis Colon Rectum* 1985;**28**:847–51.

42. **Togashi K**, Konishi F, Koinuma K, et al. Flat and depressed lesions of the colon and rectum: pathogenesis and clinical management. *Ann Acad Med Singapore* 2003;**32**:152–8.
43. **Kudo SE**, Kashida H. Flat and depressed lesions of the colorectum. *Clin Gastroenterol Hepatol* 2005;**3**(7 Suppl 1):S33–6.
44. **Goto H**, Oda Y, Murakami Y, et al. Proportion of de novo cancers among colorectal cancers in Japan. *Gastroenterology* 2006;**131**:40–46.
45. **Wong KS**, Chua WC, Cheong DM, et al. Flat colorectal lesions: colonoscopic detection without dye spray or magnification and clinical significance. *Asian J Surg* 2004;**27**:299–302.
46. **Lee JH**, Kim JW, Cho YK, et al. Detection of colorectal adenoma by routine chromoendoscopy with indigocarmine. *Am J Gastroenterol* 2003;**98**:1284–1288.
47. **Rajendra S**, Kutty K, Karim N. Flat colonic adenoma in Malaysia: fact or fancy? *J Gastroenterol Hepatol* 2003;**18**:701–704.
48. **Hawkins MJ**, Ward RL. Sporadic colorectal cancer with microsatellite instability and their possible origin in hyperplastic polyps and serrated adenomas. *J Natl Cancer Inst* 2001;**93**:1307–13.
49. **Yuen ST**, Davies H, Chan TL, et al. Similarity of the phenotypic patterns associated with BRAF and KRAS mutations in colorectal neoplasia. *Cancer Res* 2002;**62**:6451–5.
50. **Chan TL**, Zhao W, Leung SY, Yuen ST, Cancer Genome Project. BRAF and KRAS mutation in colorectal hyperplastic polyps and serrated adenoma. *Cancer Res* 2003;**63**:4878–81.
51. **Selby JV**, Friedman GD, Quesenberry CP Jr, et al. A case control study of screening sigmoidoscopy and mortality from colorectal cancer. *N Engl J Med* 1992;**326**:653–7.
52. **Newcomb PA**, Norfleet RG, Storer BE, et al. Screening sigmoidoscopy and colorectal cancer mortality. *J Natl Cancer Inst* 1992;**84**:1572–75.
53. **Winawer SJ**, Zauber AG, Ho MN, et al. Prevention of colorectal cancer by colonoscopic polypectomy: The National Polyp Study Workgroup. *N Engl J Med* 1993;**329**:1877–81.
54. **Citarda G**, Tomasselli, Capocaccia R, Barchiesi S, Crespi M, the Italian Multicenter Study Group. Efficacy in standard clinical practice of colonoscopic polypectomy in reducing colorectal cancer incidence. *Gut* 2001;**48**:812–5.
55. **Zauber AG**, Winawer SJ, O'Brien MJ, et al. Significant long term reduction in colorectal cancer mortality with colonoscopic polypectomy: findings of the National Polyp Study. *Gastroenterol* 2007;**132**(Suppl 2):A–50.
56. **Sano Y**, Fujii T, Oda Y, et al. A multi-center randomized controlled trial designed to evaluate follow-up surveillance strategies for colorectal cancer: the Japan Polyp Study. *Digest Endo* 2004;**16**:376–8.
57. **US Preventive Service Task Force**. Screening for colorectal cancer: recommendation and rationale. *Ann Intern Med* 2002;**137**:129–31.
58. **Winawer S**, Fletcher R, Rex D, et al. Colorectal cancer screening and surveillance: clinical guideline and rationale—update based on new evidence. *Gastroenterology* 2003;**124**:544–60.
59. **Canadian Task Force on Preventive Health Care**. Colorectal cancer screening: Recommendation statement from the Canadian Task Force on Preventive Health Care. *CMAJ* 2001;**165**:205–8.
60. **Rhodes JM**. Colorectal cancer screening in the UK: Joint position statement by the British Society of Gastroenterology, the Royal Colleges of Physicians, and the Association of Coloproctology of Great Britain and Ireland. *Gut* 2000;**46**:746–8.
61. **Mandel JS**, Church TR, Bond JH, et al. The effect of fecal occult blood screening on the incidence of colorectal cancer. *N Engl J Med* 2000;**343**:1803–7.
62. **Mandel JS**, Church TR, Ederer F, et al. Colorectal cancer mortality: effectiveness of biennial screening for colorectal cancer with fecal occult-blood. *J Natl Cancer Inst* 1999;**91**:434–7.
63. **Young GP**, St John JB, Winawer SJ, et al. Choice of fecal occult blood tests for colorectal cancer screening: recommendations based on performance characteristics in population studies. A WHO and OMEP report. *Am J Gastroenterol* 2002;**97**:2499–507.
64. **Allison JE**, Tekawa IS, Ransom LJ, et al. A comparison of fecal occult blood tests for colorectal cancer screening. *N Engl J Med* 1996;**334**:155–9.
65. **Smith A**, Young GP, Cole SR, et al. Comparison of a brush-sample fecal immunochemical test for haemoglobin with a sensitive guaiac-based fecal occult blood test in detection of colorectal neoplasia. *Cancer* 2005;**107**:2152–9.
66. **Wong BCY**, Wong WM, Cheung KL, et al. A sensitive guaiac fecal occult blood test is less useful than an immunochemical test for colorectal cancer screening in a Chinese population. *Aliment Pharmacol Ther* 2003;**18**:941–6.
67. **Atkins WS**, Crook FC, Quirk J, et al. Single flexible sigmoidoscopy screening to prevent colorectal cancer: baseline findings of a UK multi-center randomized trial. *Lancet* 2002;**359**:1291–300.
68. **Rex DK**, Cutler CS, Lemmel GT, et al. Colonoscopic miss rates of adenoma determined by back-to-back colonoscopies. *Gastroenterology* 1997;**112**:24–8.
69. **Rabeneck L**, Lewis JD, Fazzari L, et al. Flexible sigmoidoscopy is associated with a reduced incidence of distal but not proximal colorectal cancer: a population based cohort study. *Gastroenterology* 2006;**130**:A44.
70. **Rockey DC**, Paulson E, Niedzwiedz D, et al. Analysis of air contrast barium enema, computed tomographic colonography and colonoscopy: prospective comparison. *Lancet* 2005;**365**:305–11.
71. **Winawer SJ**, Stewart ET, Zauber AG, et al. A comparison of colonoscopy and double-contrast barium enema for surveillance after polypectomy. National Polyp Study Working Group. *N Engl J Med* 2000;**342**:1766–72.
72. **Pickhardt PJ**, Choi JR, Hwang I, et al. Computed tomographic virtual colonoscopy to screen for colorectal neoplasia in asymptomatic adults. *N Engl J Med* 2003;**349**:2191–200.
73. **Kim DH**, Pickhardt PJ, Taylor AJ, et al. CT colonography versus colonoscopy for the detection of advanced neoplasia. *N Engl J Med* 2007;**357**:1403–12.
74. **Muthall BP**, Veerappan GR, Jackson JL. Meta-analysis: computed tomographic colonography. *Ann Intern Med* 2005;**142**:635–50.
75. **Halligan S**, Taylor SA. CT colonography: results and limitations. *Eur J Radiol* 2007;**61**:400–8.
76. **Kim DH**, Pickhardt PJ, Taylor AJ. Characteristics of advanced adenomas detected at CT colonographic screening: implications for appropriate polyp size thresholds for polypectomy versus surveillance. *Am J Roentgenol* 2007;**188**:940–4.
77. **Pickhardt PJ**. CT colonography (virtual colonoscopy) for primary colorectal screening: challenges facing clinical implementation. *Abdom Imaging* 2005;**30**:1–4.
78. **Cotton PB**, Durkalski VI, Flisau BC, et al. Computed tomographic colonography (virtual colonoscopy): a multicenter comparison with standard colonoscopy for detection of colorectal neoplasia. *JAMA* 2004;**291**:1713–9.
79. **Sung JJ**. Does FOBT have a place for colorectal cancer screening in China in 2007? *Am J Gastroenterol* 2006;**101**:1–4.
80. **Lee KJ**, Inoue M, Otani T, et al. Colorectal cancer screening using fecal occult blood test and subsequent risk of colorectal cancer: a prospective cohort study in Japan. *Cancer Detect Prev* 2007;**31**:3–11.
81. **Salkeld G**, Young G, Irwig L, et al. Cost-effectiveness analysis of screening by fecal occult blood testing for colorectal cancer in Australia. *Aust N Z J Public Health* 1996;**20**:138–43.
82. **Hofstad B**, Vatn M. Growth rate of colon polyps and cancer. *Gastrointest Endosc Clin North Am* 1997;**7**:345–363.
83. **Bressler B**, Passat LF, Chen Z, et al. Rates of new or missed colorectal cancer after colonoscopy and their risk factors: a population-based analysis. *Gastroenterology* 2007;**132**:96–102.
84. **Lieberman DA**, Weiss DG, Harford WV, et al. Five-year colon surveillance after screening colonoscopy. *Gastroenterology* 2007;**133**:1077–1085.
85. **Fabry A**, Schoen RE, Weissfeld JL, et al. Analysis of colorectal cancer occurrence during surveillance colonoscopy in the dietary Polyp Prevention Trial. *Gastrointest Endosc* 2005;**61**:385–91.
86. **Arber N**, Eagle CJ, Spicak J, et al. Celecoxib for the prevention of colorectal adenomatous polyps. *N Engl J Med* 2006;**355**:885–95.
87. **Sawhney MS**, Farrar WD, Gudivasa S, et al. Microsatellite instability in interval colon cancers. *Gastroenterology* 2006;**131**:1700–5.
88. **Barclay RL**, Vicari JJ, Doughty AS, et al. Colonoscopic withdrawal times and adenoma detection during screening colonoscopy. *N Engl J Med* 2006;**355**:2533–41.
89. **Simmons DT**, Harewood GC, Baron TH, et al. Impact of endoscopic withdrawal speed on polyp yield: implications for optimal colonoscopy withdrawal time. *Aliment Pharmacol Ther* 2006;**24**:965–71.
90. **Ko CW**, Sonnenberg A. Comparing risk and benefit of colorectal cancer screening in elderly patients. *Gastroenterology* 2005;**129**:1183–70.
91. **Lin OS**, Kozarek RA, Schreiber DB, et al. Screening colonoscopy in very elderly patients: prevalence of neoplasia and estimated impact on life expectancy. *JAMA* 2006;**295**:2357–65.
92. **Tamakoshi K**, Wakai K, Kojima M, et al. A prospective study on the possible association between having children and colon cancer risk: findings from the JACC study. *Cancer Sci* 2004;**95**:243–7.
93. **Lin OS**, Kozarek RA, Schreiber DB, et al. Risk stratification for colon neoplasia: screening strategies using colonoscopy and computerized tomographic colonography. *Gastroenterology* 2006;**131**:1011–9.
94. **Imperiale TF**, Wagner DR, Lin CY, et al. Using risk of advanced proximal colonic neoplasia to tailor endoscopic screening for colorectal cancer. *Ann Intern Med* 2003;**139**:959–65.
95. **Liao JM**, Lin JT, Huang SP, et al. Screening for colorectal cancer in average-risk Chinese population using a mixed strategy with sigmoidoscopy and colonoscopy. *Dis Colon Rectum* 2000;**43**:630–40.
96. **Fuchs CS**, Giovannucci EL, Colditz GA, et al. A prospective study of family history and the risk of colorectal cancer. *N Engl J Med* 1994;**331**:1669–74.
97. **Pariente A**, Milan C, Lafon J, et al. Colonoscopic screening in first-degree relatives of patients with sporadic colorectal cancer: a case control study. The Association Nationale des Gastroentérologues des Hôpitaux et Régistre Bourguignon des Cancers Digestifs. *Gastroenterol* 1998;**115**:7–12.
98. **Bazzoli F**, Fossi S, Sottili S, et al. The risk of adenomatous polyps in asymptomatic first-degree relatives of persons with colon cancer. *Gastroenterology* 1995;**109**:783–9.
99. **Johns LE**, Houlston RS. A systematic review and meta-analysis of familial colorectal cancer risk. *Am J Gastroenterol* 2001;**96**:2992–3003.
100. **Tung SY**, Wu CS. Risk factors for colorectal adenomas among immediate family members of patients with colorectal cancer in Taiwan—a case control study. *Am J Gastroenterol* 2000;**95**:3624–8.
101. **Honojo S**, Kono S, Shinchi K, et al. The relation of smoking, alcohol use and obesity to risk of sigmoidal and rectal adenoma. *Jpn J Cancer Res* 1995;**86**:1019–26.
102. **Otani T**, Iwasaki M, Yamamoto S, et al. Alcohol consumption, smoking and subsequent risk of colorectal cancer in middle-aged and elderly Japanese men and women: Japan Public Health Center-based prospective study. *Cancer Epidemiol Biomarkers Prev* 2003;**12**:1492–500.

Guidelines

103. Akhter M, Nishino Y, Nakaya N, et al. Cigarette smoking and the risk of colorectal cancer among men: a prospective study in Japan. *Eur J Cancer Prev* 2007;**16**:102-7.
104. Tsong WH, Koh WP, Yuan JM, et al. Cigarette and alcohol in relation to colorectal cancer: the Singapore Chinese Health Study. *Br J Cancer* 2007;**96**:821-7.
105. Dai Z, Xu YC, Niu L. Obesity and colorectal cancer risk: a meta-analysis of cohort studies. *World J Gastroenterol* 2007;**13**:4199-206.
106. Kim JH, Lim YJ, Kim YH, et al. Is metabolic syndrome a risk factor for colorectal adenoma? *Cancer Epidemiol Biomarkers Prev* 2007;**16**:1543-6.
107. Yamakoshi K, Wakai K, Kojima M, et al. A prospective study of body size and colon cancer mortality in Japan: the JACC study. *Int J Obes Relat Metab Disord* 2004;**28**:551-8.
108. Kono S, Shinichi K, Imanishi K. Body mass index and adenoma of the sigmoid colon in Japanese men. *Eur J Epidemiol* 1996;**12**:425-6.
109. Chung YW, Han DS, Park YK, et al. Association of obesity, serum glucose and lipids with the risk of advanced colorectal adenoma and cancer: a case control study in Korea. *Dig Liver Dis* 2006;**38**:668-72.
110. Wang YY, Lin SY, Lai WA, et al. Association between adenomas of rectosigmoid colon and metabolic syndrome features in a Chinese population. *J Gastroenterol Hepatol* 2005;**20**:1410-5.
111. Chan AO, Jim MH, Lam KF, et al. Prevalence of colorectal neoplasm among patients with newly diagnosed coronary artery disease. *JAMA* 2007;**298**:1412-9.
112. Pox C, Schmiegel W, Classen M. Current status of screening colonoscopy in Europe and in the United States. *Endoscopy* 2007;**39**:168-73.
113. National Health and Medical Research Council. Guidelines for the prevention, early detection and management of colorectal cancer. <http://www.cancerscreening.gov>.
114. Saito H. Colorectal cancer screening using immunochemical faecal occult blood testing in Japan. *J Med Screen* 2006;**13**(Suppl 1):S6-7.
115. Ministry of Health and Welfare ROK. National Cancer Control Guideline. Seoul: Ministry of Health and Welfare ROK, 2004.
116. Department of Health. National Cancer Control Five-year Program. Bureau of Health Promotion, Department of Health, Taiwan. <http://www.bhp.doh.gov.tw/english/category.php?table=research&page=detail&id=44&pid=96> (accessed 19 May 2008).
117. Ministry of Health. Clinical practice guidelines on health screening. Singapore: Ministry of Health, 2003.
118. Saito H. Colorectal cancer screening using immunochemical faecal occult blood test in Japan. *J Med Screen* 2006;**13**(Suppl 1):S6-7.
119. Zarychanski R, Chen Y, Bernstein CN, et al. Frequency of colorectal cancer screening and the impact of family physicians on screening behavior. *CMAJ* 2007;**177**:593-7.
120. Frazier AL, Colditz GA, Fuchs CS, et al. Cost-effectiveness of colorectal cancer in the general population. *JAMA* 2000;**284**:1954-61.
121. Sung JJY, Choi SYP, Chan FKL, et al. Obstacles to colorectal cancer screening in Chinese: a study based on the health belief model. *Am J Gastroenterol* 2008;**103**:974-81.
122. Wong BCY, Chan AOO, Wong WM, et al. Attitudes and knowledge of colorectal cancer and screening in Hong Kong: a population-based study. *J Gastroenterol Hepatol* 2006;**21**:42-6.
123. Ng EST, Tan CH, To DCL, et al. The knowledge and perceptions regarding colorectal screening are unique in the Chinese—a community-based study in Singapore. *Prev Med* 2007;**45**:332-5.
124. Levy BT, Dawson J, Hartz AJ, et al. Colorectal cancer testing among patients cared for by low family physicians. *Am J Prev Med* 2006;**31**:193-201.
125. Levy BT, Nordin T, Sinit S, et al. Why hasn't this patient been screened for colon cancer? An Iowa Research Network study. *J Am Board Fam Med* 2007;**20**:458-68.
126. Shapiro TF, Hoover J, Paszat LF, et al. Colorectal cancer screening with nurse-performed flexible sigmoidoscopy: result from a Canadian community-based program. *Gastrointest Endosc* 2007;**65**:640-5.
127. Brotherton H, Vance M, Edwards R, et al. Uptake of population-based flexible sigmoidoscopy screening for colorectal cancer: a nurse-led feasibility study. *J Med Screen* 2007;**14**:76-80.
128. Horton K, Reffel A, Rosen K, et al. Training of nurse practitioners and physician assistants to perform screening flexible sigmoidoscopy. *J Am Acad Nurse Pract* 2001;**13**:455-9.

Appendix A

Voting Members of the Consensus Group

Joseph Sung (Chair), Murdani Abdullah, Carolyn Aldige, Edgardo Bondoc, Durado Brooks, Jeong-Sik Byeon, Shan-rong Cai, Annie Chan, Francis Chan, Kelvin Cheng, Jessica Ching, Han-mo Chiu, Vui Heng Chong, Khean-Lee Goh, Lawrence KY Ho, Andrew Ip, Yasuo Kakugawa, Wing-man Ko, Ken Koo, Pinit Kulavanijaya, Philip Kwok, James Lau, Rupert Leong, Suet-yei Leung, Wai-keung Leung, Chu-jun Li, Peng Li, David Lieberman, Su-vui Lo, Vivian Lou, Susie Lum, Govind Makharja, Simon Ng, Yasushi Oda, Fei-chai Pang, Rungsun Rerknimitr, Yasushi Sano, Jose Sallano, Ka-fai To, Kelvin Tsoi, Martin Wong, Mei-Yin Wong, Kai-chun Wu, Deng-chyang Wu, Ming-shiang Wu, Eng-kiong Yeoh, Khay-quan Yeoh, Graeme Young, Su-tsan Yuen, Shu Zheng

Editor's quiz: GI snapshot

ANSWER

From the question on page 1101

During endoscopy, it was found that a wooden toothpick was embedded in the posterior wall of the distal antrum surrounded by a subtle, rounded bulge seen actively to exude a small amount of pus from its centre (fig 1 of the Question). An overtube was placed and a 33 mm long toothpick was recovered. Figure 1 shows the lesion after removal of the stick. In retrospect, the patient had no recollection of having swallowed a toothpick. There was marked diminution of the patient's pain postprocedure. A follow-up abdominal x ray and CT scan to rule out perforation and abscess were unremarkable. The patient was admitted to hospital for 1 day, and subsequently discharged in a stable condition.

Clinicians should include inadvertent foreign body ingestion in the differential diagnosis for abdominal pain and gastrointestinal bleed. Patients should be warned of the potential hazards of toothpicks and cocktails sticks, fragments of which may be left in club sandwiches which have been cut in half.




Figure 1 A subtle bulge with pus seen after the removed of the tooth pick from the distal antrum of the stomach.

Patient consent: Patient consent has been received for publication of the case details and the figures in this paper.

Gut 2008;**57**:1176. doi:10.1136/gut.2007.129981a

ORIGINAL CONTRIBUTIONS

Endoscopy

Efficacy of the Invasive/Non-invasive Pattern by Magnifying Chromoendoscopy to Estimate the Depth of Invasion of Early Colorectal Neoplasms

Takahisa Matsuda, M.D.,¹ Takahiro Fujii, M.D., Ph.D.,² Yutaka Saito, M.D., Ph.D.,¹ Takeshi Nakajima, M.D., Ph.D.,¹ Toshio Uraoka, M.D., Ph.D.,¹ Nozomu Kobayashi, M.D.,¹ Hisatomo Ikehara, M.D.,¹ Hiroaki Ikematsu, M.D.,³ Kuang-I Fu, M.D., Ph.D.,³ Fabian Emura, M.D., Ph.D.,¹ Akiko Ono, M.D.,¹ Yasushi Sano, M.D., Ph.D.,³ Tadakazu Shimoda, M.D., Ph.D.,⁴ and Takahiro Fujimori, M.D., Ph.D.⁵

¹Endoscopy Division, National Cancer Center Hospital, Tokyo, Japan; ²TF Clinic, Tokyo, Japan; ³Division of Digestive Endoscopy and Gastrointestinal Oncology, National Cancer Center Hospital East, Chiba, Japan; ⁴Clinical Laboratory Division, National Cancer Center Hospital, Tokyo, Japan; and ⁵Department of Surgical and Molecular Pathology, Dokkyo University School of Medicine, Shimotsuga, Tochigi, Japan

- OBJECTIVE:** During colonoscopy, estimation of the depth of invasion in early colorectal lesions is crucial for an adequate therapeutic management and for such task, magnifying chromoendoscopy (MCE) has been proposed as the best *in vivo* method. However, validation in large-scale studies is lacking. The aim of this prospective study was to clarify the effectiveness of MCE in the diagnosis of the depth of invasion of early colorectal neoplasms in a large series.
- METHODS:** A total of 4,215 neoplastic lesions were evaluated using MCE from October 1998 to September 2005 at the National Cancer Center Hospital, Tokyo, Japan. Lesions were prospectively classified according to the clinical classification of the pit pattern: invasive pattern or non-invasive pattern. All lesions were histopathologically evaluated.
- RESULTS:** There were 3,371 adenomas, 612 intramucosal cancers (m-ca), 232 submucosal cancers (sm-ca): 52 sm superficial (sm1) and 180 sm deep cancers (sm 2–3). Among lesions diagnosed as invasive pattern, 154 out of 178 (86.5%) were sm2–3, while among lesions diagnosed as non-invasive pattern, 4,011 out of 4,037 (99.4%) were adenomas, m-ca, or sm1. Sensitivity, specificity and diagnostic accuracy of the invasive pattern to differentiate m-ca or sm1 (<1000 μ m) from sm2–3 ($\geq 1000 \mu$ m) were 85.6%, 99.4%, and 98.8%, respectively.
- CONCLUSION:** The determination of invasive or non-invasive pattern by MCE is a highly effective *in vivo* method to predict the depth of invasion of colorectal neoplasms.

(Am J Gastroenterol 2008;103:2700–2706)

INTRODUCTION

It has been reported that intramucosal colorectal cancers (m-ca) show no lymph node (LN) metastasis (LNM) and are good candidates for endoscopic resection (1, 2). In contrast, submucosal cancers (sm-ca) have approximately 6–12% of LNM, which require surgical resection including LN dissection for curative treatment (3–6). Recently, there is growing evidence supporting the theory that lesions with sm invasion limited to less than 1000 μ m (sm1) without lymphovascular invasion and/or poorly differentiated component do not involve LNM (7). Therefore, it is crucial to estimate the depth

of invasion of early colorectal neoplasms accurately prior to therapeutic decisions.

Magnifying chromoendoscopy (MCE) has widely demonstrated its effectiveness to differentiate between colorectal neoplastic and non-neoplastic polyps (8–15). In relation to the depth of invasion of colorectal neoplastic lesions, Kudo's classification of colonic crypts suggests that type III and IV pit patterns are found on adenomatous polyps, while type VN is strongly suggestive of sm deep cancers (16–18). In practice, however, there are limitations using only the morphological classification of the pit pattern to discriminate between m-sm1 and sm2 or beyond. MCE with detailed analysis of the pit

pattern has been proposed as the best *in vivo* method to evaluate the depth of invasion, however, validation in large-scale studies is lacking. Herein, we report a clinical classification of the colonic pit pattern, which is useful to determine the proper treatment of colorectal lesions during colonoscopy.

METHODS

Patients

A total of 3,029 consecutive patients diagnosed with a neoplastic colorectal lesion at the National Cancer Center Hospital, Tokyo from October 1998 to September 2005 were enrolled. This study was conducted prospectively, and the study protocol was approved by our institutional review board. Written informed consents for diagnosis and treatment were obtained from all patients prior to the procedures. Exclusion criteria were advanced colorectal cancer, familial adenomatous polyposis (FAP), inflammatory bowel disease (IBD), and hereditary non-polyposis colorectal cancer (HNPCC).

Magnifying Colonoscope

All examinations were performed using magnifying colonoscopes (CF-Q240ZI, PCF-Q240ZI, and CF-200Z, Olympus

Optical Co., Tokyo, Japan), which enhance the image up to 80–100 times using a one-touch operation power system. These scopes have equal upward and downward bending range as well as sideways range and biopsy channel diameter but differ minimally in observation range diameters and view angles.

Definition of Terms

1. *Regular pit*: visible crypt orifice independently of shape (Fig. 1A).
2. *Irregular pit*: the orifice of each crypt is indented or jagged (Fig. 1B).
3. *Distorted pit*: the orifice of each crypt cannot be clearly traced, usually seen in desmoplastic areas (Fig. 1C).
4. *Demarcated area*: clearly visualized zone between two morphologically different types of pits, e.g., depression, large nodule, or reddened area (Fig. 1D).

Clinical Classification

1. *Non-neoplastic pattern*: normal mucosa and star-shaped crypts as observed in Kudo's type I or II, respectively (e.g., hyperplastic, juvenile and inflammatory polyps).




Figure 1. Definition of terms: (A) Regular pit. (B) Irregular pit. (C) Distorted pit. (D) Demarcated area: Depression/Reddened area.




Figure 2. Relationship between Kudo's classification and clinical classification.

2. *Non-invasive pattern*: regular crypts with or without demarcated area or irregular pits without a demarcated area. Usually observed in Kudo's type III_S, III_L, IV, and selected cases of VI (e.g., adenomatous polyps, intramucosal, and submucosal superficial cancers), where endoscopic resection is appropriate.
3. *Invasive pattern*: irregular and distorted crypts in a demarcated area as observed in Kudo's type VN and selected cases of VI (e.g., deep submucosal invasive can-


cers), where surgical resection is the appropriate treatment. Kudo's type VI is observed in both non-invasive and invasive patterns (Fig. 2, 3).

Endoscopic Examination

All patients were prepared for colonoscopy with 2–3 L of polyethylene glycol-electrolyte solution administered on the morning of examination day. Scopolamine butylbromide (10 mg) or Glucagon (0.5 mg) was administered intravenously in patients with no contraindication prior to examination to avoid bowel movement.

All procedures were carried out by experienced examiners who had performed more than 500 colonoscopies per year. When a lesion was detected by conventional view, its surface was washed out with proteinase to remove overlying mucous, after which 0.4% indigo carmine (IC) dye was sprayed to accentuate the contours of the lesions. When a colonoscopist intended to perform chromoendoscopy, a volume of 3 to 5 mL IC dye was flushed through the biopsy channel with 15 mL of air using a 20cc syringe and sprayed directly over the targeted lesion. The pit pattern was evaluated by magnifying view. When high magnification observation with IC dye was not enough for determining the surface structure (pit pattern analysis), 0.05% crystal violet (CV) was applied as a staining method (19). Lesions were evaluated under MCE in real time and categorized as non-neoplastic, neoplastic non-invasive and neoplastic invasive patterns. Those diagnosed as

Invasive pattern: Irregular/ Distorted pit with Demarcated area



Non-invasive pattern: Regular pit with or without demarcated area or irregular pits without a demarcated area




Figure 3. Definition of invasive/non-invasive pattern. Invasive pattern (A) Demarcated area (+): Depression, Irregular/distorted pit (B) Demarcated area (+): Reddened area, Irregular/distorted pit. Non-invasive pattern (C) Demarcated area (+): Depression, Regular pit (D) Demarcated area (+): Reddened area, Regular pit.

non-neoplastic were left untreated. If lesions were identified as neoplastic non-invasive lesions (adenomas or m-ca), hot biopsy, snare polypectomy, or endoscopic mucosal resection (EMR) was performed. Lesions <5 mm were resected by coagulation biopsy (hot biopsy), and flat lesions or those >5 mm were treated with loop snare polypectomy or EMR. If lesions were diagnosed as invasive pattern, biopsy specimens were taken and patients were basically referred for surgery. In cases where a polyp was not clearly diagnosed as either hyperplastic polyp (HP) or adenomatous polyp (AP), it was considered as AP and therefore removed for histopathological analysis.

Histopathology

Resected specimens were immediately fixed in 10% buffered formalin solution and subsequently stained with hematoxylin-eosin. Experienced gastrointestinal pathologists blinded to each endoscopic diagnosis evaluated all pathological specimens. Histopathological diagnosis was determined according to the Japanese Research Society for Cancer of the Colon and Rectum (JRSCCR) between 1998 and 2000. Since 2001, lesions were histopathologically evaluated according to JRSCCR and Vienna classification as well (20, 21). Non-pedunculated lesions with a vertical invasion length of less than 1000 μ m in the submucosal layer, and pedunculated lesions with head invasion, were classified as submucosal superficial invasive cancer (sm1). Non-pedunculated lesions with invasion of more than 1000 μ m and pedunculated lesions with stalk invasion were considered as submucosal deep invasive cancer (sm2-3) (7, 22). Regarding pedunculated lesions, level 2 according to Haggitt's classification (23) was used as the baseline to determine submucosal invasion.

RESULTS

A total of 4,215 neoplastic lesions in 3,029 patients were studied by MCE and removed endoscopically or surgically for analysis. Among lesions endoscopically diagnosed as having an invasive pattern, there were 45 (25.3%) right-sided colon lesions, 53 (29.8%) left-sided lesions, and 80 (44.9%) rectal lesions. On the other hand, among the lesions endoscopically diagnosed as having a non-invasive pattern, there were 2,032 (50.3%) right-sided, 1,475 (36.5%) left-sided and 530 (13.1%) rectal lesions. According to macroscopic type, there were 90 (50.6%) polypoid, 9 (5.1%) flat elevated, and 79 (44.4%) depressed lesions in the invasive pattern group. In contrast, there were 2,700 (66.9%) polypoid, 1,258 (31.2%) flat elevated, and 79 (2.0%) depressed lesions in the non-invasive pattern group (Table 1).

Histopathological analysis revealed adenoma: 3371 (80.0%), intramucosal cancer (m-ca): 612 (14.5%), and submucosal cancer (sm-ca): 232 (5.5%) [sm1: 52 (1.2%), sm2-3: 180 (4.3%)]. Among lesions diagnosed as invasive pattern, 154 out of 178 (86.5%) were submucosal deep invasive can-

Table 1. Characteristics of Lesions Diagnosed by Magnifying Chromoendoscopy

	Invasive Pattern (n = 178)	Non-Invasive Pattern (n = 4037)
Location—no. (%)		
Right colon*	45 (25)	2032 (50)
Left colon**	53 (30)	1475 (37)
Rectum	80 (45)	530 (13)
Macroscopic type—no. (%)		
Polypoid	90 (51)	2700 (67)
Flat elevated	9 (5)	1258 (31)
Depressed†	79 (44)	79 (2)
Size—no. (%)		
~5 mm	2 (1)	2024 (50)
6–10 mm	30 (17)	1396 (35)
11–20 mm	101 (57)	493 (12)
21 mm—	45 (25)	124 (3)

*Cecum-transverse colon.

**Descending-sigmoid colon.

†I1c, IIa+IIc, I+IIc.

cers (sm2-3), while 4,011 out of 4037 (99.4%) diagnosed as non-invasive pattern were adenomas, m-ca or sm1 (Table 2).

Diagnostic Accuracy

The calculated sensitivity, specificity, positive predictive value (PPV), negative predictive value (NPV), and accuracy of the invasive pattern to differentiate m-ca or sm1 (<1000 μ m) from sm2-3 ($\geq$ 1000 μ m) were 85.6%, 99.4%, 86.5%, 99.4%, and 98.8%, respectively. Based on the macroscopic appearance, the diagnostic sensitivity of the clinical pit pattern to determine the depth of invasion of polypoid, flat, and depressed lesions was 75.8% (75/99), 85.7% (6/7), and 98.6% (73/74), respectively. Meanwhile, the specificity for polypoid, flat, and depressed lesions was 99.4%, 99.8%, and 92.9%, respectively (Table 3).

Treatment Strategy

Among lesions endoscopically diagnosed as having an invasive pattern, 80.9% (144/178) were treated surgically of which sm2-3 cancer was found in 132 cases (91.7%), and

Table 2. Clinicopathologic Characteristics of Lesions Diagnosed by Magnifying Chromoendoscopy

	Invasive Pattern (n = 178)	Non-Invasive Pattern (n = 4037)
No. of lesions	178	4037
Histopathology—no. (%)		
Adenoma	0 (0)	3371 (83)
m-ca	12 (7)	600 (15)
sm-ca	166 (93)	66 (2)
~sm superficial (sm1*)	12 (7)	40 (1)
~sm deep (sm2-3)	154 (86)	26 (0.6)

*sm1: sm <1000 μ m.

Sensitivity: 85.6% (15/180).

Specificity: 99.4% (4011/4035).

Positive predictive value (PPV): 86.5% (154/178).

Negative predictive value (NPV): 99.4% (4011/4037).

Accuracy: 98.8% (4165/4215).

Table 3. Diagnostic Sensitivity and Specificity According to Macroscopic Type

	Polypoid	Flat	Depressed	Total
sm deep	75.8% (75/99)	85.7% (6/7)	98.6% (73/74)	85.6% (154/180)
Adenoma	99.4% (2676/2691)	99.8% (1257/1260)	92.9% (78/84)	99.4% (4011/4035)
m-ca				
sm superficial				

m-ca or sm1 in 12 cases (8.3%). Thirty-four lesions (19.1%) diagnosed as an invasive pattern were removed endoscopically. Based on lesion size, there were 31 out of 34 (91.2%) small lesions (<20 mm) and 25 out of 34 (73.5%) polypoid lesions. Among these, sm2-3 cancer was found in 22 cases (64.7%), and m-ca or sm1 in 12 cases (35.3%). Among 22 sm2-3 cancers, 15 cases underwent additional surgical treatment and in 7 cases, close follow-up was performed. In contrast, among 4,037 cases diagnosed endoscopically as having a non-invasive pattern, 4,024 (99.7%) were resected endoscopically. The remaining 13 cases (0.3%) were treated surgically. Based on size, there were 8 out of 13 (61.5%) large lesions ≥ 21 mm and 8 out of 13 (61.5%) polypoid lesions (Fig. 4).


DISCUSSION

This is the first large prospective study to assess the effectiveness of MCE for the endoscopic estimation of the depth of invasion of colorectal neoplasms. MCE is a standardized validated method that facilitates detailed analysis of the morphological architecture of colonic mucosal crypt orifices (pit pattern) in a simple and not time-consuming manner. Despite prospective randomized studies in large reference centers demonstrating the superiority of MCE over conventional colonoscopy, and studies that demonstrated no differences between magnifying and standard colonoscopes in the aver-

age time to reach the cecum (11), magnifying endoscopes are still rarely used in endoscopy units. Unrecognized necessity and lack of randomized studies validating the effectiveness of MCE are possible reasons for this. We believe that MCE is an essential armamentarium in gastrointestinal (GI) endoscopy units and that its main clinical significance is the *in vivo* diagnosis of the nature of colorectal lesions, which gives extremely useful information to determine the treatment modality.

The clinical classification of the colonic pit pattern (invasive and non-invasive) using MCE was originally described by Fujii in 1998 with the aim to discriminate between m-sm1 and sm2 or beyond (19). Contrary to the anatomic classification of Kudo *et al.*, the rationale for the clinical classification is based on the identification of irregular or distorted crypts in a demarcated area, (where the orifice of each crypt cannot be traced clearly) which highly suggests that the cancerous lesion is already invading deeply into the sm layer.

Some studies have already reported the clinical usefulness of detailed determination of the V pit pattern using MCE for predicting the depth of invasion of sm neoplasms. Kudo *et al.* reported that 11 of 22 (50%) lesions with a type V pit pattern with a bounded surface were found to be invasive cancers with involvement of the sm layer (16). Other studies have reported a diagnostic accuracy of type V pit for the diagnosis of sm invasive cancer of 85% (81/95) and 79% (11/14), respectively (24, 25). Recently, Ohta *et al.* reported that lesions with high-grade atypia have erosive change in the surface epithelium and histological appearance of a desmoplastic reaction. Due to these histopathological changes, lesions invading deeply into the sm layer usually show a demarcated area on its surface (26). However, protruded type lesions do not often show an invasive pattern, even if they invade deeply into sm. In such cases, it is sometimes difficult to predict the degree of invasion only evaluating the tumor surface. In this study, we found out that the clinical classification of the pit

**Figure 4.** Treatment strategy.