

associated with suicidal ideation in major depression among cancer patients. *Psychooncology* 19:384-9, 2010

学会発表

1. Uchida M, Akechi T, Okuyama T, Sagawa R, Nakaguchi T, Endo C, Yamashita H, Toyama T, Furukawa TA. Patients' supportive care needs and psychological distress in advanced breast cancer patients in Japan. Patients' supportive care needs and psychological distress in advanced breast cancer patients in Japan, 57th Psychosomatic Medicine, 2010 Nov
 2. Nakaguchi T, Akechi T, Okuyama T, Sagawa R, Uchida M, Ito Y, Arakawa A, Nishikawa H, Ishida T, Sugie C, Furukawa TA Usefulness of eye movement desensitization and reprocessing (EMDR) for psychological nausea, vomiting and learned food aversion experienced by cancer patients receiving repeated chemotherapy: a case series. Book Usefulness of eye movement desensitization and reprocessing (EMDR) for psychological nausea, vomiting and learned food aversion experienced by cancer patients receiving repeated chemotherapy: a case series, 57th Psychosomatic Medicine, 2010 Nov
 3. Akechi T, Okuyama T, Endo C, Sagawa R, Uchida M, Nakaguchi T, Akazawa T, Yamashita H, Toyama T, Furukawa TA. Patient's perceived need and psychological distress and/or quality of life in ambulatory breast cancer patients in Japan. Book Patient's perceived need and psychological distress and/or quality of life in ambulatory breast cancer patients in Japan, 57th Psychosomatic Medicine, 2010 Nov
 4. Okuyama T, Akechi T, Yamashita H, Toyama T, Endo C, Sagawa R, Uchida M, Furukawa TA. Nurses in outpatient chemotherapy center may have difficulty in assessing their patients' symptoms and supportive care needs. Nurses in outpatient chemotherapy center may have difficulty in assessing their patients' symptoms and supportive care needs, 12th World Congress of Psycho-Oncology, 2010 May
 5. Akechi T, Okuyama T, Endo C, Sagawa R, Uchida M, Nakaguchi T, Akazawa T, Yamashita H, Toyama T, Furukawa TA. Patient's perceived need and psychological distress and/or quality of life in ambulatory breast cancer patients in Japan. Book Patient's perceived need and psychological distress and/or quality of life in ambulatory breast cancer patients in Japan, 12th World Congress of Psycho-Oncology, 2010 May
 6. 中口智博, 明智龍男, 奥山 徹, 佐川竜一, 内田恵, 伊藤嘉規, 荒川敦志, 西川博, 石田高司, 杉江愛生, 古川壽亮: 化学療法に起因した予期性悪心嘔吐、食物嫌悪に奏功した短期心理療法-EMDR. 第15回日本緩和医療学会総会, 2010, 6.
- H. 知的財産権の出願・登録状況（予定を含む。）
1. 特許取得
なし。
 2. 実用新案登録
なし。
 3. その他
特記すべきことなし。

Ⅲ. 研究成果の刊行に関する一覧表

研究成果の刊行に関する一覧表

書籍（日本語）

| 著者氏名 | 論文タイトル名 | 書籍全体の編集者名 | 書籍名 | 出版社名 | 出版地 | 出版年 | ページ |
|------|---|---|------------------|-----------------------|-----|------|---------|
| 平井 啓 | 統合的ながんのケアにおける「こころと体」の医学 | Donald I. Abrams, Andrew T. Weil/ 伊藤壽記、上島悦子監訳 | がんの統合医療 | メディカル・サイエンス・インターナショナル | 東京 | 2010 | 246-261 |
| 平井 啓 | 問題解決療法の紹介 | Laurence Mynors-Walls / 明智龍男、平井 啓、本岡寛子（監訳） | 不安と抑うつに対する問題解決療法 | 金剛出版 | 東京 | 2009 | 7-16 |
| 伊藤壽記 | なぜ、がんに対して統合医療でなければならぬのか？ | Donald I. Abrams, Andrew T. Weil/ 伊藤壽記、上島悦子監訳 | がんの統合医療 | メディカル・サイエンス・インターナショナル | 東京 | 2010 | 1-12 |
| 伊藤壽記 | がん予防に対する統合的アプローチ | Donald I. Abrams, Andrew T. Weil/ 伊藤壽記、上島悦子監訳 | がんの統合医療 | メディカル・サイエンス・インターナショナル | 東京 | 2010 | 13-37 |
| 伊藤壽記 | 現代医療の新たな領域、補完代替医療から統合医療-さらなる医療の質の向上を求めて | 千代豪昭、黒田研二 | 学生のための医療概論（第3版） | 医学書院 | 東京 | 2009 | 212-224 |

雑誌（外国語）

| 発表者氏名 | 論文タイトル名 | 発表誌名 | 巻号 | ページ | 出版年 |
|---|---|--|-------|---------|----------|
| Yoshida S, Hirai K, Morita T, Miyashita M, et al: | Experience of families of Japanese patients with cancer for prognostic disclosure | Journal of Pain and Symptom Management | | | In press |
| Yoshida S, Otani H, Hirai K, et al | A qualitative study of decision-making by breast cancer patients about telling their children about their illness | Support Care Cancer | 18(4) | 439-447 | 2010 |

| | | | | | |
|--|---|-----------------------|--------|---------|------|
| <u>Hirai K</u> , et al | Self-efficacy, psychological adjustment and decisional-balance regarding decision making for outpatient chemotherapy in Japanese advanced lung cancer | Psychology and Health | 24 (2) | 149-60 | 2009 |
| <u>Hirai K</u> , Uchitomi Y, et al | Discrimination between worry and anxiety among cancer patients: development of a brief cancer-related worry inventory | Psychooncology | 17 | 1172-79 | 2008 |
| Shiozaki M, <u>Hirai K</u> , <u>Morita T</u> , et al | Measuring the regret of bereaved family members regarding the decision to admit cancer patients to palliative care units | Psychooncology | 17 | 926-931 | 2008 |
| Akechi T, <u>Hirai K</u> , et al | Problem-solving therapy for psychological distress in Japanese cancer patients: preliminary clinical experience from psychiatric consultations | Jpn J Clin Oncol | 38 | 867-70 | 2008 |
| Shinjo T, <u>Morita T</u> , <u>Miyashita M</u> , Sato K, Tsuneto S, Shima Y | Care for the Bodies of Deceased Cancer Inpatients in Japanese Palliative Care Units | J Palliat Med | 13(1) | 27-31 | 2010 |
| Shinjo T, <u>Morita T</u> , <u>Hirai K</u> , <u>Miyashita M</u> , Sato K, Tsuneto S, Shima Y | Care for imminently dying cancer patients: family members' experiences and recommendations | J Clin Oncol | 28(1) | 142-8 | 2010 |
| Okamoto T, Ando M, <u>Morita T</u> , <u>Hirai K</u> , Kawamura R, <u>Miyashita M</u> , Sato K, Shima Y | Religious care required for Japanese terminally ill patients with cancer from the perspective of bereaved family members | Am J Hosp Palliat Med | 27(1) | 50-4 | 2010 |
| Ando M, <u>Morita T</u> , Akechi T, Ifuku Y | A qualitative study of mindfulness-based meditation therapy in Japanese cancer patients | Support Care Cancer | | | 2010 |
| Ando M, Kawamura R, <u>Morita T</u> , <u>Hirai K</u> , <u>Miyashita M</u> , Okamoto T, Shima Y | Value of religious care for relief of psycho-existential suffering in Japanese terminally ill cancer patients: the perspective of bereaved family members | Psychooncology | 19(7) | 750-755 | 2010 |
| Akazawa T, Akechi T, <u>Morita T</u> , <u>Miyashita M</u> , Sato K, Tsuneto S, Shima Y, Furukawa TA | Self-perceived burden in terminally ill cancer patients: A categorization of care strategies based on bereaved family members' | J Pain Symptom Manage | 40(2) | 224-234 | 2010 |

| | | | | | |
|--|---|----------------------------|--------|----------|------|
| | perspectives | | | | |
| Yamagishi A, <u>Morita T</u> , <u>Miyashita M</u> , Sato K, Tsuneto S, Shima Y | The care strategy for families of terminally ill cancer patients who become unable to take nourishment orally: Recommendations from a nationwide survey of bereaved family members' experiences | J Pain Symptom Manage | 40(5) | 671-683 | 2010 |
| <u>Morita T</u> , <u>Miyashita M</u> , Tsuneto S, Sato K, Shima Y | Late Referrals to Palliative Care Units in Japan: Nationwide Follow-up Survey and Effects of Palliative Care Team Involvement After the Cancer Control Act | J Pain Symptom Manage | 38(2) | 191-6 | 2009 |
| <u>Morita T</u> , Murata H, Kishi E, <u>Miyashita M</u> , Yamaguchi T, Uchitomi Y | Meaninglessness in terminally ill cancer patients: a randomized controlled study | J Pain Symptom Manage | 37(4) | 649-58 | 2009 |
| Sanjo M, <u>Morita T</u> , <u>Miyashita M</u> , <u>Hirai K</u> , et al | Caregiving consequences inventory: a measure for evaluating caregiving consequences from the bereaved family member's perspective | Psychooncology | 18(6) | 657-666 | 2009 |
| Ando M, <u>Morita T</u> , et al | The efficacy of mindfulness-based meditation therapy on anxiety, depression, and spirituality in Japanese patients with cancer | J Palliat Med | 12(12) | 1091-94 | 2009 |
| Sanjo M, <u>Morita T</u> , <u>Hirai K</u> , et al | Perceptions of Specialized Inpatient Palliative Care: A Population-Based Survey in Japan | J Pain Symptom Management | 35 | 275-82 | 2008 |
| Akechi T, <u>Morita T</u> , et al | Psychotherapy for depression among incurable cancer patients | Cochrane Database Syst Rev | Apr 16 | CD005537 | 2008 |
| <u>Miyashita M</u> , Narita Y, Sakamoto A, Kawada M, Akiyama M, Kayama M, Suzukamo Y, Fukuhara S | Health-related quality of life among community-dwelling patients with intractable neurological diseases and their caregivers in Japan | Psychiat Clin Neuros | 65 | 30-8 | 2010 |

| | | | | | |
|---|---|-----------------------|-------|---------|------|
| Choi JE, <u>Miyashita M</u> , <u>Hirai K</u> , Sato K, <u>Morita T</u> , Tsuneto S, Shima Y | Preference of place for end-of-life cancer care and death among bereaved Japanese families who experienced home hospice care and death of a loved one | Support Care Cancer | 18 | 1445-53 | 2009 |
| <u>Miyashita M</u> , et al | Quality indicators of end-of-life cancer care from the bereaved family members' perspective in Japan | J Pain Symptom Manage | 37(6) | 1019-26 | 2009 |
| <u>Miyashita M</u> , Narita Y, Sakamoto A, et.al | Care burden and depression in caregivers caring for patients with intractable neurological diseases at home in Japan | J Neurol Sci | 276 | 148-52 | 2009 |
| Misawa T, <u>Miyashita M</u> , Kawa M, Abe K, Abe M, Nakayama Y, Given CW | Validity and reliability of the Japanese version of the Caregiver Reaction Assessment Scale (CRA-J) for community-dwelling cancer patients | Am J Hosp Palliat Med | 26(5) | 334-40 | 2009 |
| Sato K, <u>Miyashita M</u> , <u>Morita T</u> , Suzuki M | Long-term effect of a population-based educational intervention focusing on end-of-life home care, life prolongation treatment, knowledge about palliative care | J Palliat Care | 25(3) | 206-12 | 2009 |
| <u>Miyashita M</u> , <u>Morita T</u> , <u>Hirai K</u> , et al | Good death inventory: a measure for evaluating good death from the bereaved family member's perspective | J Pain Symptom Manage | 35 | 486-98 | 2008 |
| <u>Miyashita M</u> , <u>Morita T</u> , <u>Hirai K</u> , et al | Factors contributing to evaluation of a good death from the bereaved family member's perspective | Psychooncology | 17 | 612-620 | 2008 |
| <u>Miyashita M</u> , <u>Hirai K</u> , <u>Morita T</u> , et al | Barriers to referral to inpatient palliative care units in Japan: a qualitative survey with content analysis | Support Care Cancer | 16 | 217-22 | 2008 |
| <u>Miyashita M</u> , <u>Morita T</u> , <u>Hirai K</u> | Evaluation of end-of-life cancer care from the perspective of bereaved family members: the Japanese experience | J Clin Oncol | 26 | 3845-52 | 2008 |
| Sanjo M, <u>Miyashita M</u> , <u>Morita T</u> , <u>Hirai K</u> , et al | Perceptions of specialized inpatient palliative care: a population-based survey in Japan | J Pain Symptom Manage | 35 | 275-282 | 2008 |

| | | | | | |
|--|--|----------------------|-------------|----------|------|
| Miyashita M, et al | Quality of life, day hospice needs, and satisfaction of community-dwelling advanced cancer patients and their caregivers in Japan | J Palliat Med | 11(9) | 1203-7 | 2008 |
| Moritake H, Shimonodan H, Marutsuka K, Kamimura S, Kojima H, Nunoi H | C-MYC rearrangement may induce an aggressive phenotype in anaplastic lymphoma a kinase positive anaplastic large cell lymphoma: Identification of a novel fusion gene AL017/C-MYC | Am J Hematol | 86 | 5-8 | 2011 |
| Moritake H, et al | Cytomegalovirus infection mimicking juvenile myelomonocytic leukemia showing hypersensitivity to granulocyte-macrophage colony stimulating factor | Pediatr Blood Cancer | 53:1 | 1324-6 | 2009 |
| Funakoshi-Tago M, Pelletier S, Moritake H, Parganas E, Ihle JN | Jak2 FERM domain interaction with the Erythropoietin Receptor regulates Jak2 kinase activity | Mol Cell Biol | 28 | 1792-801 | 2008 |
| Ohta H, Iwamoto S, Kiyokawa N, Tsurusawa M, Deguchi T, Takase K, Fujimoto J, Horibe K, Komada Y | Flow cytometric analysis of de novo acute myeloid leukemia in childhood: report from the Japanese Pediatric Leukemia/ Lymphoma Study Group | Int J Hematol | 93 | 135-137 | 2011 |
| Ohta H, Kusuki S, Yoshida H, Sato E, Hashii Y, Ozono K | Allogeneic hematopoietic stem cell transplantation with reduced intensity conditioning for a child with recurrent anaplastic large cell lymphoma | Int J Hematol | 92 | 190-193 | 2010 |
| Ohta H, et al | Pediatric post-transplant lymphoproliferative disorder after cardiac transplantation | Int J Hematol | 90 | 127-136 | 2009 |
| Kusuki S, Ohta H, et al | Antifungal prophylaxis with micafungin in patients treated for childhood cancer | Pediatr Blood Cancer | 53 | 605-609 | 2009 |
| Akechi T, Okuyama T, Endo C, Sagawa R, Uchida M, Nakaguchi T, Sakamoto M, Komatsu H, Ueda R, Wada M, Furukawa TA | Anticipatory nausea among ambulatory cancer patients undergoing chemotherapy: Prevalence, associated factors, and impact on quality of life | Cancer Sci | Dec;101(12) | 2596-600 | 2010 |

| | | | | | |
|---|--|----------------|-----------|-------|----------|
| Akechi T, Okuyama T, Endo C, Sagawa R, Uchida M, Nakaguchi T, Akazawa T, Yamashita H, Toyama T, Furukawa TA | Patient's perceived need and psychological distress and/or quality of life in ambulatory breast cancer patients in Japan | Psychooncology | | | In press |
| Akechi T, Okamura H, Nakano T, Akizuki N, Okamura M, Shimizu K, Okuyama T, Furukawa TA, Uchitomi Y | Gender differences in factors associated with suicidal ideation in major depression among cancer patients | Psychooncology | Apr;19(4) | 384-9 | 2010 |

雑誌（日本語）

| 発表者氏名 | 論文タイトル名 | 発表誌名 | 巻号 | ページ | 出版年 |
|---|---|--------------------------|-------|---------|------|
| 平井 啓 | がん患者に対する問題解決療法とは？ | がん患者ケア | 4(3) | 48-52 | 2011 |
| 岩満優美, 平井 啓, 大庭 章, 塩崎麻里子, 浅井真理子, 尾形明子, 他 | 緩和ケアチームが求める心理士の役割に関する研究—フォーカスグループインタビューを用いて— | Palliative Care Research | 4 | 228-34 | 2009 |
| 平井 啓, 他 | 【緩和医療における精神症状への対策】 がん患者に対する問題解決療法 | 緩和医療学 | 10 | 37-42 | 2008 |
| 平井 啓 | がんの補完代替医療における心理学的研究の役割 | ストレス科学 | 22 | 216-223 | 2008 |
| 伊藤壽記, 井倉 技 | 補完代替医療（CAM）の現代医療における意義と展望 | 食品加工技術 | 28 | 16-19 | 2008 |
| 井倉 技, 伊藤壽記 | 特集「統合医療への期待—21世紀の予防医学と健康づくり」 米国における統合医療の発展 | 公衆衛生 | 72 | 95-98 | 2008 |
| 尾形明子 | 小児緩和ケアにおける心理士の取り組み | 小児看護 | 33 | 1541-5 | 2010 |
| 武井優子・尾形明子・小澤美和・真部淳・鈴木伸一 | 小児がん患者が退院後に抱える心理社会的問題に関する研究の現状と課題 | 小児がん | 47 | 84-90 | 2010 |
| 尾形明子 | 病気の子どもに対する心理的支援 | 育療 | 45 | 2-7 | 2009 |
| 尾形明子, 佐伯俊成 | 小児がん患者と家族に対する心理的ケア | 総合病院精神医学 | 20 | 26-32 | 2008 |
| 三善陽子, 太田秀明, 他 | 小児がん治療後長期フォローアップの現状 | 小児外科 | 40 | 697-702 | 2008 |
| 吉田沙蘭・天野功二・森田達也・尾形明子・平井 啓 | 難治性小児がん患児の家族が経験する困難の探索 | 小児がん | 47(1) | 91-97 | 2010 |

