

Umetani and colleagues¹³ reported 11 cases of colorectal cancers that had more than two barium enema examinations that were at least 6 months apart. Five non-polypoid submucosal invasive cancers were studied; three developed from non-polypoid and two from polypoid lesions. The authors estimated tumor doubling time to evaluate the growth rate of each tumor, and suggested that NP-CRN grew slowly compared with polypoid lesion and maintained their macroscopic morphology. The data, however, are limited.

COLONOSCOPIC ANALYSIS

Matsui and colleagues¹⁴ reported eight early colorectal cancers that were incidentally followed by colonoscopy. Thirty-five cases, including 14 NP-CRN as initial lesions, were reviewed. They found that NP-CRN progressed to submucosally invasive cancer, retaining its non-polypoid configuration, and some flat lesions developed depressed areas during their progression. Although the authors also estimated the speed of growth compared with polypoid and non-polypoid lesions, they could not conclude which type grew more rapidly.

Sato and colleagues¹⁵ prospectively followed 12 small flat adenomas. The size of the lesions ranged from 2 to 6 mm (median 4 mm), and the observation period ranged from 11 to 26 months (median 19 months). Although eight lesions showed various changes in their shape, only two lesions demonstrated an increase in diameter of the tumor. All of the flat lesions were subsequently removed endoscopically and found to be adenomas. The authors concluded small flat adenoma did not rapidly progress, and configuration change did not indicate tumor progression or invasion.

Watari and colleagues¹⁶ conducted prospective colonoscopic study to elucidate the natural history of NP-CRN. The authors observed 75 colorectal tumors measuring less than 1 cm in diameter in 50 patients. The average follow-up period was 22 months, and 62 lesions (83%) were NP-CRN. They concluded similar observations as those of Sato and colleagues, although they found that 40% of small non-polypoid lesions had exophytic growth with time. This finding suggested some small non-polypoid lesions follow the adenoma–carcinoma sequence the same as polypoid lesions.

THE IMPORTANCE OF THE DEPRESSED-TYPE NP-CRN

Matsuda and colleagues¹⁷ analyzed 6638 colorectal neoplasms, excluding advanced cancers, treated in National Cancer Center Hospital, Tokyo, Japan. There were 4471 (67%) and 2167 (33%), polypoid and non-polypoid colorectal neoplasms, respectively. Among all non-polypoid lesions, there were 178 (2.7%) depressed lesions, 109 (61%) of which were diagnosed as high-grade dysplasia or submucosally invasive cancer. Among 5538 (83%) lesions that were identified as low- or high-grade dysplasia, the proportion of depressed lesions was 1.3%. On the other hand, depressed type was identified in 39% (Table 1) of submucosal cancers. This discrepancy may indicate of a rapid progression rate of depressed lesions into invasive cancers.

Sano and colleagues¹⁸ described the incidence of depressed lesions among all of colorectal neoplasms, again excluding advanced cancers. Their multicenter retrospective study conducted in eight Japanese referral institutes revealed that the incidence of depressed lesions was 1.94% (1291 depressed lesions out of 66,670 neoplasms), and, in particular, 51.2% of intramucosal depressed lesions were diagnosed as high-grade dysplasia. These data also suggested that intramucosal depressed lesions showed more aggressive behavior and were perhaps more likely

	LGD	HGD	SM-Ca
Polypoid	3781 (68.3)	578 (67.9)	112 (45.0)
Flat	1688 (30.5)	260 (30.6)	41 (16.5)
Depressed	69 (1.2)	13 (1.5)	96 (38.6)
Total	5538 (83.4)	851 (12.8)	249 (3.8)

Abbreviations: HGD, high-grade dysplasia; LGD, low-grade dysplasia; SM-Ca, submucosal invasive cancer.

Data from Matsuda T, Saito Y, Hotta K, et al. Prevalence and clinicopathological features of non-polypoid colorectal neoplasms: should we pay more attention to identifying flat and depressed lesions? Dig Endosc 2010;22(Suppl 1):S57-62.

to develop into invasive cancers as compared with the polypoid lesions. The depressed type of NP-CRN appears to be pathologically and molecular biologically distinct that other types of NP-CRN.

Several authors reported that depressed-type colorectal cancer does not arise from an adenomatous polyp. This theory was called de novo carcinogenesis, and lack of K-ras mutation was thought to be a distinctive genetic feature.^{10,11,19,20} Goto and colleagues²¹ reported the proportion of de novo cancers among all colorectal cancers in a cohort of 14,817 Japanese populations. The authors defined de novo cancers according to both criteria: (1) the absence of adenomatous components and (2) all lateral margins of the tumor covered with normal mucosa and non-polypoid growth pattern. They concluded that 22.9% of early colorectal cancers were de novo cancers. Chen and colleagues,²² from Taiwan, also assessed the proportion of de novo carcinomas using the Markov model, and demonstrated about 30% of colorectal cancers arising from de novo sequence.

Number	Macroscopic Type	Size (mm)	Location	Depth of Lesion
1	lsp (semipedunculated)	13	Sigmoid	SM
2	lsp (semipedunculated)	15	Sigmoid	SM
3	ls (sessile)	8	Rectum	SM
4	ls (sessile)	10	Sigmoid	SM
5	ls (sessile)	20	Rectum	MP
6	ls (sessile)	6	Transverse	SM
7	lla (flat)	15	Transverse	SM
8	lla (flat)	20	Sigmoid	SM
9	lla + llc (depressed)	20	Cecum	SM
10	lla + llc (depressed)	20	Transverse	SM
11	lla + llc (depressed)	10	Rectum	MP
12	lla + llc (depressed)	6	Ascending	SM
13	lla + llc (depressed)	20	Sigmoid	SS

Abbreviations: MP, muscularis; SM, submucosa; SS, subserosa.

Data from Matsuda T, Fujii T, Sano Y, et al. Five-year incidence of advanced neoplasia after initial colonoscopy in Japan: a multicenter retrospective cohort study. Jpn J Clin Oncol 2009;39:435-42.

THE JAPAN POLYP STUDY

To clarify the natural history of NP-CRN, a large cohort study focused on the detection of NP-CRN is required. Matsuda and colleagues²³ reported the results of multicenter retrospective cohort study to evaluate 5-year incidence of advanced neoplasia after initial colonoscopy. The authors studied 5309 patients with a median follow-up period of 5.1 years. Endoscopists diagnosed 13 invasive cancers on follow-up within 3 years. The initial colonoscopies were performed by Japanese endoscopists who had proficient technique with chromoendoscopy to diagnose NP-CRN, and the patients had good preparation quality, taking polyethylene glycol (PEG) solution in the morning on the day of colonoscopy. Out of the 13 incident cancer cases, seven were NP-CRN, and the mean size of these lesions was less than 15 mm in diameter (**Table 2**). These data suggested that NP-CRN is responsible for interval cancers, defined as colorectal cancers diagnosed within several years of a complete colonoscopy. Much prospective data, however, are necessary to elucidate the natural history and epidemiology of these lesions. The Japan Polyp Study (JPS) is a multicenter randomized controlled trial prospectively evaluating follow-up surveillance strategy for Japanese patients after removal of all polypoid and non-polypoid neoplasms.²⁴ This study is intended to continue until 2011 and hopefully will provide new information on the detection and progression of NP-CRN.

SUMMARY

The natural history of NP-CRN is mostly unknown. The results of small observational studies suggest that NP-CRN lesions develop into invasive cancer with minimal size expansion. Among NP-CRN lesions, depressed lesions show more aggressive behavior and frequently develop into invasive cancers compared with polypoid lesions, regardless of their low incidence. A large prospective cohort study focused on the detection of NP-CRN is currently ongoing.

REFERENCES

1. Muto T, Kamiya J, Sawada T, et al. Small flat adenoma of the large bowel with special reference to its clinicopathologic features. *Dis Colon Rectum* 1985;28:847-51.
2. Kudo S. Endoscopic mucosal resection of flat depressed type of early colorectal cancer. *Endoscopy* 1993;25:455-61.
3. Fujii T, Rembacken BJ, Dixon MF, et al. Flat adenomas in the United Kingdom: are treatable cancers being missed? *Endoscopy* 1998;30:437-43.
4. Saitoh Y, Waxman I, West AB, et al. Prevalence and distinctive biologic features of flat colorectal adenomas in a North American population. *Gastroenterology* 2001;120:1657-65.
5. Tsuda S, Veress B, Toth E, et al. Flat and depressed colorectal tumours in a southern Swedish population: a prospective chromoendoscopic and histopathological study. *Gut* 2002;51:550-5.
6. Rembacken BJ, Fujii T, Cairns A, et al. Flat and depressed colonic neoplasms: a prospective study of 1000 colonoscopies in the UK. *Lancet* 2000;355:1211-4.
7. Parra-Blanco A, Gimeno-Garcia AZ, Nicolas-Perez D, et al. Risk for high-grade dysplasia or invasive carcinoma in colorectal flat adenomas in a Spanish population. *Gastroenterol Hepatol* 2006;29:602-9.

8. Soetikno RM, Kaltenbach T, Rouse RV, et al. Prevalence of nonpolypoid (flat and depressed) colorectal neoplasms in asymptomatic and symptomatic adults. *JAMA* 2008;299:1027–35.
9. Chiu HM, Lin JT, Chen CC, et al. Prevalence and characteristics of nonpolypoid colorectal neoplasm in an asymptomatic and average-risk Chinese population. *Clin Gastroenterol Hepatol* 2009;7:463–70.
10. Kudo S, Kashida H, Tamura T, et al. Colonoscopic diagnosis and management of nonpolypoid early colorectal cancer. *World J Surg* 2000;24:1081–90.
11. Shimoda T, Ikegami M, Fujisaki J, et al. Early colorectal carcinoma with special reference to its development de novo. *Cancer* 1989;64:1138–46.
12. Matsui T, Yao T, Iwashita A. Natural history of early colorectal cancer. *World J Surg* 2000;24:1022–8.
13. Umetani N, Masaki T, Watanabe T, et al. Retrospective radiographic analysis of nonpedunculated colorectal carcinomas with special reference to tumour doubling time and morphological change. *Am J Gastroenterol* 2000;95:1794–9.
14. Matsui T, Tsuda S, Iwashita A, et al. Retrospective endoscopic study of developmental and configurational changes of early colorectal cancer: eight cases and a review of the literature. *Dig Endosc* 2004;16:1–8.
15. Sato T, Konishi F, Togashi K, et al. Prospective observation of small flat tumors in the colon through colonoscopy. *Dis Colon Rectum* 1999;42:1457–63.
16. Watari J, Saitoh Y, Obara T, et al. Natural history of colorectal nonpolypoid adenomas: a prospective colonoscopic study and relation with cell kinetics and K-ras mutations. *Am J Gastroenterol* 2002;97:2109–15.
17. Matsuda T, Saito Y, Hotta K, et al. Prevalence and clinicopathological features of nonpolypoid colorectal neoplasms: should we pay more attention to identifying flat and depressed lesions? *Dig Endosc* 2010;22(Suppl 1):S57–62.
18. Sano Y, Tanaka S, Teixeira CR, et al. Endoscopic detection and diagnosis of 0-IIc neoplastic colorectal lesions. *Endoscopy* 2005;37:261–7.
19. Fujimori T, Satonaka K, Yamamura-Idei Y, et al. Noninvolvement of ras mutations in flat colorectal adenomas and carcinomas. *Int J Cancer* 1994;57:51–5.
20. Kaneko K, Fujii T, Kato S, et al. Growth pattern and genetic changes of colorectal carcinoma. *Jpn J Clin Oncol* 1998;28:196–201.
21. Goto H, Oda Y, Murakami Y, et al. Proportion of de novo cancers among colorectal cancers in Japan. *Gastroenterology* 2006;131:40–6.
22. Chen CD, Yen MF, Wang WM, et al. A case-cohort study for the disease natural history of adenoma-carcinoma and de novo carcinoma and surveillance of colon and rectum after polypectomy: implication for efficacy of colonoscopy. *Br J Cancer* 2003;88:1866–73.
23. Matsuda T, Fujii T, Sano Y, et al. Five-year incidence of advanced neoplasia after initial colonoscopy in Japan: a multicenter retrospective cohort study. *Jpn J Clin Oncol* 2009;39:435–42.
24. Sano Y, Fujii T, Oda Y, et al. A multicenter randomized controlled trial designed to evaluate follow-up surveillance strategies for colorectal cancer: the Japan Polyp Study. *Dig Endosc* 2004;16:376–8.

Diagnostic accuracy of narrow-band imaging and pit pattern analysis significantly improved for less-experienced endoscopists after an expanded training program

Reiji Higashi, MD, Toshio Uraoka, MD, PhD, Jun Kato, MD, PhD, Kenji Kuwaki, MD, Shin Ishikawa, MD, PhD, Yutaka Saito, MD, PhD, Takahisa Matsuda, MD, PhD, Hiroaki Ikematsu, MD, Yasushi Sano, MD, PhD, Seiyuu Suzuki, MD, PhD, Yoshitaka Murakami, PhD, Kazuhide Yamamoto, MD, PhD

Okayama, Tokyo, Chiba, Kobe, Niihama, Otsu, Japan

Background: Previous reports assessing diagnostic skill using narrow-band imaging (NBI) and pit pattern analysis for colorectal polyps involved only highly experienced endoscopists.

Objective: To evaluate diagnostic skills of less-experienced endoscopists (LEE group) for differentiation of diminutive colorectal polyps by using NBI and pit pattern analysis with and without magnification after an expanded training program.

Design: Prospective study.

Patients: This study involved 32 patients with 44 colorectal polyps (27 adenomas and 17 hyperplastic polyps) of ≤ 5 mm that were identified and analyzed by using conventional colonoscopy as well as non-magnification and magnification NBI and chromoendoscopy followed by endoscopic removal for histopathological analysis.

Intervention: Before a training course, 220 endoscopic images were distributed in randomized order to residents with no prior endoscopy experience (NEE group) and to the LEE group, who had performed colonoscopies for more than 5 years but had never used NBI. The 220 images were also distributed to highly experienced endoscopists (HEE group) who had routinely used NBI for more than 5 years. The images were distributed to the NEE and LEE groups again after a training class. Magnification NBI and chromoendoscopy images were assessed by using the Sano and Kudo classification systems, respectively.

Main Outcome Measurements: Diagnostic accuracy and interobserver agreement for each endoscopic modality in each group.

Results: Diagnostic accuracy was significantly higher, and kappa (κ) values improved in the LEE group for NBI with high magnification after expanded training. Diagnostic accuracy and κ values when using high-magnification NBI were highest among endoscopic techniques for the LEE group after such training and the HEE group (accuracy 90% vs 93%; $\kappa = 0.79$ vs 0.85, respectively).

Limitations: Study involved only polyps of ≤ 5 mm.

Conclusion: Using high-magnification NBI increased the differential diagnostic skill of the LEE group after expanded training so that it was equivalent to that of the HEE group. (*Gastrointest Endosc* 2010;72:127-35.)

Abbreviations: CC, conventional colonoscopy; CE-H, high-magnification chromoendoscopy; CE-L, low-magnification chromoendoscopy; HEE, highly experienced endoscopist; LEE, less-experienced endoscopist; NBI, narrow-band imaging; NBI-H, high-magnification narrow-band imaging; NBI-L, low-magnification narrow-band imaging; NEE, no-experience endoscopist; SSA, sessile serrated adenoma.

DISCLOSURE: All authors disclosed no financial relationships relevant to this publication.

Copyright © 2010 by the American Society for Gastrointestinal Endoscopy
0016-5107/\$36.00
doi:10.1016/j.gie.2010.01.054

Received August 10, 2009. Accepted January 25, 2010.

Reprint requests: Toshio Uraoka, MD, PhD, Department of Endoscopy, Okayama University Hospital, 2-5-1 Shikata-cho, Okayama 700-8558, Japan.

Current affiliations: Department of Gastroenterology and Hepatology (R.H., J.K., K.K., S.I., K.Y.), Okayama University Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, Department of Endoscopy (T.U.), Okayama University Hospital, Okayama, Endoscopy Division (Y.S., T.M.), National Cancer Center Hospital, Tokyo, Division of Digestive Endoscopy and Gastrointestinal Oncology (H.I.), National Cancer Center Hospital East, Kashiwa, Gastrointestinal Center (Y.S.), Sano Hospital, Kobe, Department of Internal Medicine (S.S.), Sumitomo Besshi Hospital, Niihama, Department of Medical Statistics (Y.M.), Shiga University of Medical Science, Otsu, Japan.

It is widely accepted that adenomatous polyps are precursors of colorectal cancer, and performing polypectomies on such lesions can reduce the risk of subsequent colorectal cancer by up to 80% for a period that may exceed 10 years.¹ In addition, adenomas are a major factor in guidelines that have been developed for recommended colonoscopy surveillance intervals after polypectomies because they are a powerful predictor for future colorectal cancer risk.¹⁻³ Small colorectal adenomas as well as advanced adenomas⁴ are precursors of colorectal cancer, and multiple genetic alterations have been implicated in the adenoma-carcinoma sequence.⁵

Endoscopic differentiation of small adenomas from non-adenomatous polyps is important because endoscopists should avoid performing any unnecessary procedure, including polypectomies, that can cause related complications such as bleeding and perforation.^{6,7} The diagnostic accuracy of conventional colonoscopy for such colorectal polyps, however, has previously been reported to be unsatisfactory.^{8,9} In contrast, chromoendoscopy with indigo-carmin dye spraying has been shown to be an effective procedure for detecting and evaluating colorectal polyps¹⁰⁻¹⁵ despite having several disadvantages, including a longer procedure time and the additional cost for dye spraying.

Narrow-band imaging (NBI) is an innovative optical technology providing a unique image that emphasizes the morphological and structural character of lesions as well as their surface capillary patterns.¹⁶⁻²⁵ It has been reported that this modality is a new non-dye tool for differentiating neoplastic from non-neoplastic polyps, with a diagnostic accuracy including pit pattern analysis equivalent to that of chromoendoscopy.¹⁷⁻²¹ Such reports have been based on studies involving only highly experienced endoscopists, however, with few published articles concerned with the learning curve for NBI being dependent on an individual endoscopist's experience and ability. The aim of this study was to determine, therefore, whether expanded training in the effective use of NBI and pit pattern analysis with and without magnification would improve the diagnostic skill of less-experienced endoscopists in the differentiation of diminutive colorectal polyps.

METHODS

Patients

Patients scheduled for a total colonoscopy at Okayama University Hospital and Sumitomo Besshi Hospital between September and October 2008 were invited to participate in this study. Informed consent was obtained from all patients before their examinations. Patients with inflammatory bowel disease, familial adenomatous polyposis, an international normalized ratio greater than 2.0, or a platelet count less than 50,000/mm³ were excluded from this study.

Take-home Message

- Expanded interactive training in effective use of narrow-band imaging both with and without magnification as well as pit pattern analysis improved diagnostic accuracy and interobserver agreement of less-experienced colonoscopists in differentiating diminutive colorectal polyps. Using narrow-band imaging with high magnification increased the differential diagnostic skill of less-experienced colonoscopists who underwent such training to a level equivalent to that of highly experienced colonoscopists.

Colonoscopy and polyp assessment protocol

Bowel preparation consisted of patients drinking 2 to 3 liters of polyethylene glycol solution in the morning before their procedures.¹³ Total colonoscopies were prospectively performed by using a video endoscopic system (EVIS, Lucera Spectrum; Olympus Co, Tokyo, Japan) with CF-H260AZI or PCF-Q240ZI magnification colonoscopes (Olympus) by two highly experienced endoscopists (R.H., T.U.), each of whom had previously performed over 1000 colonoscopies annually.

When a lesion was detected by conventional colonoscopy examination, surface mucus was washed away with lukewarm water, and endoscopic images were taken in the following order: conventional colonoscopy (CC), low-magnification NBI (NBI-L), high-magnification NBI (NBI-H), low-magnification chromoendoscopy (CE-L) and high-magnification chromoendoscopy (CE-H). A standard optical filter was used for both CC and chromoendoscopy, with chromoendoscopic images taken after 0.2% indigo-carmin dye was sprayed on the lesion surface. The enhanced surface structure function of the video image processor at the level A5 setting was used in taking all endoscopic images.²³ Location, size, and macroscopic type of each lesion were recorded, with size measured by using open forceps. Lesions were classified macroscopically based on the criteria of the Paris classification of superficial GI lesions.²⁶ A biopsy, polypectomy, or EMR was then performed, and the resulting specimen was analyzed histopathologically.

Image evaluator categories

A total of 12 doctors with different levels of endoscopic experience were asked to independently evaluate endoscopic images. The doctors were separated into 3 groups: 4 residents with no prior endoscopy experience (NEE group); 4 less experienced endoscopists each of whom had performed colonoscopies for more than 5 years but had never used NBI (LEE group); and 4 highly experienced endoscopists each of whom had routinely used magnification colonoscopy with NBI for more than 5 years (HEE group).

Assessment of endoscopic images

The best quality endoscopic images were selected for each modality and stored digitally in JPEG format. All images were distributed in randomized order to each group of evaluators. For the NEE and LEE groups, the same images were distributed in randomized order once before and once after those group members participated in an intensive, 1-hour, interactive training program on white-light endoscopy, NBI, and chromoendoscopy. The program included information on the Sano NBI classification^{20,21} and the Kudo pit pattern classification^{27,28} and used an atlas of endoscopic images of polyps produced by an independent group of highly experienced endoscopists. Although they were completely unaware of the histopathological results, every participant correctly diagnosed polyps as either neoplasms or non-neoplasms by using (1) chromoendoscopy based on the Kudo pit pattern classification, with types III (including III-L and III-S), IV, and V (including VI and VN) considered to be neoplasms and types I and II regarded as non-neoplasms;^{27,28} (2) NBI-L that revealed a brownish area determined to be neoplastic; and (3) NBI-H with the Sano classification of meshed capillary vessel pattern, in which types II and III were considered neoplastic and type I, without meshed capillaries, was non-neoplastic (Figs. 1 and 2).²⁰⁻²¹ Patient information such as age, sex, and clinical diagnosis was not disclosed to any of the evaluators, and discussions were not permitted among the doctors individually or in groups.

Statistical analysis

The diagnostic accuracy of each endoscopic modality was assessed in reference to histopathological results. Estimates of diagnostic accuracy were calculated based on the average diagnostic accuracy for each group of doctors as well as for each diagnostic modality. The upper and lower 95% confidence interval (CI) limits were calculated by using a normal model that consisted of symmetric CIs, with limits at a distance from the estimate equal to the product of 1.96 times the standard error. Interobserver agreement in diagnosing colorectal lesions in each group and by each modality was determined by calculation of the kappa statistic (κ) and its 95% CI by using the Fleiss method. Diagnostic accuracies before training and after training in the NEE and LEE groups were compared with the McNemar test. As for differences in diagnostic accuracies after training among the NEE, LEE, and HEE groups, those findings were analyzed by using the Fisher exact test. Multiple statistical testing of outcome data was conducted in this study, therefore, a Bonferroni correction was applied, and differences with a *P* value of $< .025$ were considered significant as the correction. A κ value of < 0.4 was regarded as poor agreement, 0.41 to 0.60 fair agreement, 0.61 to 0.80 good agreement, and > 0.80 excellent agreement. Statistical analyses were conducted by using

version 7.0 of the JMP statistical software package (SAS Institute, Cary, NC) and a Microsoft Excel 2007 spreadsheet (Microsoft, Renton, Wash).

RESULTS

Clinicopathological features of colorectal lesions

Seventy-two consecutive patients were enrolled in this study for prospective endoscopic evaluation. A total of 44 lesions of ≤ 5 mm were identified and analyzed in 32 patients (Table 1). Mean (\pm standard deviation [SD]) patient age was 61.2 ± 12.3 years, and the male/female ratio was 2.2:1. Bowel preparation was considered adequate in all examinations, and complete colonoscopy was performed to the cecum in every case. There were no complications during any procedure. Of the 44 lesions, 37 were macroscopically classified as type 0-Is, 6 as type 0-IIa, and 1 as type 0-IIc. Mean (\pm SD) lesion size was 3.4 ± 1.1 mm. As for location, 22 polyps (50%) were found in the right colon (cecum, ascending, and transverse colon), 14 (32%) in the left colon (descending and sigmoid colon), and 8 (18%) in the rectum. Histopathological assessments included 27 adenomas (61%) and 17 hyperplastic polyps (39%). A total of 220 images of the 44 lesions were collected as each lesion was photographed during CC, NBI-L, NBI-H, CE-L, and CE-H.

Diagnostic accuracy of NBI and pit pattern analysis

Table 2 indicates diagnostic accuracy for each endoscopic modality. In the NEE group, diagnostic accuracies using CC, NBI-L, and NBI-H significantly improved after the training program (CC, $P < .001$; NBI-L, $P = .006$; and NBI-H, $P < .001$), but the NEE group's diagnostic accuracies were still significantly lower in all modalities except CC compared with the HEE group (CC, $P = .049$; NBI-L, $P = .0023$; NBI-H, $P < .001$; CE-L, $P < .001$; and CE-H, $P < .001$). Diagnostic accuracies in the LEE group for NBI-L, NBI-H, and CE-H also improved significantly after the training program ($P = .001$, $P < .001$, and $P = .001$, respectively). In contrast with the NEE group's results, however, subsequent diagnostic accuracies of the LEE group were not significantly different from diagnostic accuracies of the HEE group with respect to the CC, NBI-L, NBI-H, CE-L, and CE-H modalities ($P = 1.0$, $P = .60$, $P = .57$, $P = .031$, and $P = .48$, respectively).

Assessment of interobserver agreement based on endoscopic experience

Interobserver agreements in the HEE group for NBI-H were > 0.80 representing excellent agreement and > 0.60 for NBI-L, CE-L, and CE-H, representing good agreement (κ value CC, 0.5; NBI-L, 0.62; NBI-H, 0.85; CE-L, 0.69; CE-H




Figure 1. Examples of colorectal neoplastic polyps viewed by different endoscopic modalities in this study. **A**, Conventional colonoscopy view. **B**, Low-magnification NBI showed a brownish area. **C**, High-magnification NBI revealed meshed capillary vessels indicative of Sano classification type II. **D**, Low-magnification chromoendoscopy using 0.2% indigo-carmin dye spraying clearly revealed demarcated margins and surface structure. **E**, High-magnification chromoendoscopy clearly indicated Kudo classification type III L. *NBI*, narrow-band imaging.

0.7). Meanwhile, κ values for NBI-H in the LEE group improved to “good agreement” after the training program, whereas the κ values for NBI-L, CE-L, and CE-H in the LEE group improved to “fair agreement” after the training program (κ value before vs after training: NBI-H, 0.46 vs 0.79; NBI-L, 0.31 vs 0.54; CE-L, 0.32 vs 0.44; and CE-H, 0.33 vs 0.59). In contrast, however, none of the κ values for any of the modalities in the NEE group improved beyond “poor agreement” after the training program (CC, -0.068 vs 0.24;

NBI-L, 0.059 vs 0.25; NBI-H, 0.16 vs 0.39; CE-L, 0.28 vs 0.23; and CE-H, 0.12 vs 0.18) (Fig. 3).

When we compared diagnostic accuracy for each modality in the NEE group after the training program, the LEE group after the training program, and the HEE group, NBI-H had the highest accuracy rate among all 3 groups (Table 2). Similarly, the κ value for NBI-H was significantly higher in the NEE group after the training program, the LEE group after the training program, and the HEE group




Figure 2. Examples of colorectal non-neoplastic polyps viewed by different endoscopic modalities in this study. **A**, Conventional colonoscopy view. **B**, Low-magnification NBI showed a non-brownish area. **C**, High-magnification NBI in which meshed capillary vessels were not visible or only faintly visible, indicative of Sano classification type I. **D**, Low-magnification chromoendoscopy using 0.2% indigo-carmin dye spraying clearly revealed demarcated margins and surface structure. **E**, High-magnification chromoendoscopy clearly indicated Kudo classification type II. *NBI*, narrow-band imaging.

compared with the other endoscopic diagnostic modalities (Fig. 3).

DISCUSSION

Endoscopic diagnostic tools and technology are expected to be accurate and provide reliably reproducible agreement as well as be easy to use, readily available, and relatively inexpensive, but sufficient skill on the part of the

endoscopist is still required for proper diagnosis. Our prospective study demonstrated significant improvement in the LEE group in diagnostic accuracy when using NBI and CE after undergoing limited but intensive training. The improved diagnostic accuracy of the LEE group was equivalent to that of the HEE group in terms of differential diagnosis using NBI-L, NBI-H, and CE-H. In addition, both higher diagnostic accuracy (>80%) and good interobserver agreement (κ value >0.6) for diminutive colorectal

TABLE 1. Patient characteristics and histopathological features of lesions

Patients (n = 32)	
Sex, male/female	22/10
Age, years, mean (\pm SD)	61.2 (12.3)
Lesions (n = 44)	
Macroscopic type	
0-Ia	37
0-IIa	6
0-IIc	1
Size, mm, mean (\pm SD)	3.4 (1.1)
Location (right*/left†/rectum)	22/14/8
Histopathology	
Tubular adenoma	27
Hyperplastic polyp	17

SD, Standard deviation.

*Right: cecum, ascending colon, and transverse colon.

†Left: descending colon and sigmoid colon.

polyps were achieved by the LEE group when using NBI-H after the training program.

The fact that the diagnostic accuracy and κ value of NBI-H were the highest among all the endoscopic techniques analyzed in this study for both the NEE and LEE groups after the expanded training program as well as for the HEE group leads us to suggest that NBI-H is more accurate and provides a higher level of reproducible agreement than the other diagnostic tools in differentiating diminutive neoplastic from non-neoplastic colorectal polyps. Chiu et al²⁹ earlier validated that diagnostic accuracy of NBI-H was equivalent to that of CE-H. Their study reported that diagnostic accuracies for two experienced endoscopists ranged from 91% to 92% using CE-H and from 87% to 90% using NBI-H. In our study, the diagnostic accuracy of the HEE group was 85% (95% CI, 79%-89%) using CE-H and 93% (95% CI, 88%-96%) with NBI-H, although the difference between the accuracy of the two modalities was not significant. Earlier reports analyzing the diagnostic accuracy rate based on polyp size indicated that differentiation using CE-H was more difficult with diminutive colorectal polyps of <6 mm in size.⁶⁻⁷

Our results indicated that it was possible to significantly improve the diagnostic skill for differentiating diminutive colorectal polyps by using NBI-L, NBI-H, CE-L, and CE-H in the LEE group following the limited but intensive 1-hour interactive training program. We believe that, of the various endoscopic modalities, the use of NBI-H by the LEE group subsequently became both statistically equivalent to that of the HEE group in terms of diagnostic accuracy and

closest to reaching "excellent agreement" compared with the other modalities in terms of κ value for two possible reasons. The first concerns the smaller size of the polyps examined in this study, because diagnostic accuracy of diminutive colorectal polyps by using CE-H has been reported to be lower than for polyps of >5 mm.⁶⁻⁹ It is conceivable that differentiation of diminutive colorectal polyps could have been similarly affected, somehow reducing the diagnostic accuracy of CE-H while not affecting the diagnostic accuracy of NBI-H by the LEE group. Secondly, the possibility exists that members of the LEE group were able to recognize whether or not there were meshed capillary vessels on the surface of the mucosa easier than they could identify the pit patterns of diminutive colorectal polyps.

In the Rogart et al³⁰ report on the NBI learning curve, diagnostic accuracy by using NBI-L significantly improved as the experience level of endoscopists increased, with the diagnosis of approximately 130 lesions necessary for basic competency. Their findings indicated that educational sessions conducted before the assessment of lesions in combination with continual feedback regarding the accuracy of endoscopic diagnoses compared with histopathological results every 2 weeks for half a year were important factors in achieving a satisfactory learning curve. It has also been reported that use of the Kudo pit pattern classification required a longer learning curve, with experience from diagnosing at least 200 lesions needed to become competent.^{6,7,31} In contrast, our study demonstrated that an expanded 1-hour intensive interactive training program conducted by a highly experienced endoscopist enabled the LEE group members in particular to accelerate their learning curve. In addition, the Sano classification with NBI-H appears to have had a shorter learning curve compared with using NBI-L or the Kudo pit pattern classification in the diagnostic differentiation of diminutive colorectal polyps.

Besides having a higher differential diagnosis accuracy and being easier to improve the necessary diagnostic skill for accurately differentiating diminutive colorectal polyps, NBI has other clinical advantages. First, the conventional endoscopic view can be switched almost instantaneously to the NBI view by pressing a single button on the control handle of the colonoscope, and, second, NBI does not require any dye or staining solution to detect and differentiate neoplastic lesions from non-neoplastic lesions.

In recent years, advancements in the quality of endoscopic images available from high-definition endoscopy and chromoendoscopy have considerably enhanced polyp detection. Although the risk of neoplasia in diminutive polyps is <50%, and the risk of high-grade dysplasia is <2%,^{7,32,33} diminutive colorectal neoplasms as well as advanced neoplasms are among the precursors of colorectal cancer, and multiple genetic alterations have been implicated in the adenoma-carcinoma sequence.⁴ It also has been reported that lesions of ≤ 5

TABLE 2. Effectiveness of training program on diagnostic accuracy

Modality	NEE group			LEE group			HEE group	
	Before-training accuracy (95% CI)	After-training accuracy (95% CI)	P value*	Before-training accuracy (95% CI)	After-training accuracy (95% CI)	P value*	Accuracy (95% CI)	P value†
CC	0.43 (0.35-0.50)	0.64 (0.57-0.71)	<.001	0.72 (0.65-0.78)	0.74 (0.67-0.80)	NS	0.74 (0.68-0.80)	NS
NBI-L	0.53 (0.46-0.61)	0.66 (0.59-0.73)	.006	0.65 (0.58-0.72)	0.78 (0.72-0.84)	=.001	0.81 (0.75-0.87)	NS
NBI-H	0.63 (0.56-0.70)	0.74 (0.68-0.80)	<.001	0.73 (0.66-0.79)	0.90 (0.85-0.94)	<.001	0.93 (0.88-0.96)	NS
CE-L	0.68 (0.60-0.74)	0.67 (0.60-0.74)	NS	0.68 (0.60-0.74)	0.76 (0.69-0.82)	NS	0.85 (0.79-0.90)	NS
CE-H	0.63 (0.56-0.70)	0.66 (0.59-0.73)	NS	0.67 (0.60-0.74)	0.81 (0.75-0.87)	=.001	0.85 (0.79-0.89)	NS

NEE, No endoscopy experience; LEE, less experienced endoscopist; HEE, highly experienced endoscopist; CI, confidence interval; CC, conventional colonoscopy; NS, not significant; NBI-L, low-magnification narrow-band imaging; NBI-H, high-magnification narrow-band imaging; CE-L, low-magnification chromoendoscopy; CE-H, high-magnification chromoendoscopy.

*P values determined by McNemar test comparing before and after training.

†P values determined by Fisher exact test comparing LEE after training and HEE.




Figure 3. Comparison of 95% confidence interval of κ value for each endoscopic diagnostic modality according to endoscopy experience. Each bar represents the range of 95% confidence interval of the κ value. HEE, highly experienced endoscopist group; LEE, less experienced endoscopist group; Before, before participation in an intensive 1-hour interactive training program; After, after participation in an intensive 1-hour interactive training program; NEE, no endoscopy experience group; CC, conventional colonoscopy; NBI-L, low-magnification narrow-band imaging; NBI-H, high-magnification narrow-band imaging; CE-L, low-magnification chromoendoscopy; CE-H, high-magnification chromoendoscopy.

mm make up more than 80% of the colorectal polyps subjected to histopathological assessment.³³ Besides the primary consideration of reducing the risk of future colorectal cancer, the endoscopic differentiation of diminutive neoplastic polyps from non-neoplastic polyps is essential because endoscopists should avoid performing unnecessary procedures, including polypectomies on non-neoplastic polyps, and this also will reduce substantially the number of colorectal polyps requiring histopathological assessment.

There has been considerable interest recently in sessile serrated adenoma (SSA) and serrated adenoma (SA) polyps that also have been associated increasingly with an apparent increased risk of malignant transformation.³⁴ SSAs endoscopically appear as hyperplastic polyps, but there have not been any published reports as yet applying the Kudo pit pattern analysis to such SSA polyps. In the general population, the prevalence of SSAs has been estimated to range from only 1% to 7% of all polyps, and it further has been shown that most such SSA polyps can exceed 10 mm in size,³⁵ but we did not detect any SSAs or serrated adenomas in this study. Although it was recently reported that SSAs could be differentiated from hyperplastic polyps by combining NBI and autofluorescence imaging, the report in question had several limitations including the total number of SSAs being relatively small and the lack of any comparison between those two modalities and pit pattern analysis.³⁶ In addition, the actual prevalence of SSAs is difficult to assess because pathologists have been unable to reach a consensus on the diagnosis of either hyperplastic polyps or SSAs.^{37,38} Further studies will be required, therefore, to clarify the endoscopic features and conduct histopathological and molecular-based analyses of SSAs and serrated adenomas.

The primary limitation of our study is that it involved only a small number of polyps. The power of the trial compared to the observed difference was lower because the observed difference was smaller than in the alternative hypothesis used in planning this study. The sample size that was set, however, was not much different from the sample size used in similar studies. Another limitation is that this study was conducted by using endoscopic images. During a “real-time” evaluation, an endoscopist can usually view a detected lesion by using multiple angles and light modalities at variable distances, but we digitally stored all the endoscopic images taken during each exam-

ination, selected the best image from each of the 5 endoscopic observation modalities, and then randomized the distribution order of the images for diagnosis. This process was intended to decrease the likelihood of observational bias and strengthen the reliability of our results, because separate findings based on NBI images and chromoendoscopic images might otherwise have been influenced by the other and made objective evaluation of the individual diagnostic modalities difficult. A third limitation was the relatively short interval between the intensive training program and the follow-up reviews by the NEE and LEE groups. The participants in both groups reviewed all the images for the second time within 24 hours of the training program so as to avoid any possible bias resulting from a feedback learning effect such as self-training. It has previously been reported that feedback received during the development of a diagnostic skill is effective.³⁹

In conclusion, NBI, particularly high-magnification NBI, was shown to be a promising tool for diagnostic differentiation of diminutive colorectal neoplastic polyps from non-neoplastic polyps. Expanded training of the LEE group members improved their overall diagnostic ability so that it was equivalent in certain key respects to that of the participating HEE group.

REFERENCES

- Winawer SJ, Zauber AG, Ho MN, et al. Prevention of colorectal cancer by colonoscopic polypectomy. The National Polyp Study Workgroup. *N Engl J Med* 1993;329:1977-81.
- Atkin WS, Morson BC, Cuzick J. Long-term risk of colorectal cancer after excision of rectosigmoid adenomas. *N Engl J Med* 1992;326:658-62.
- Citarda F, Tomaselli G, Capocaccia R, et al. Efficacy in standard clinical practice of colonoscopic polypectomy in reducing colorectal cancer incidence. *Gut* 2001;48:812-5.
- Winawer SJ, Zauber AG. The advanced adenoma as the primary target of screening. *Gastrointest Endosc Clin N Am* 2002;12:1-9.
- Vogelstein B, Fearon ER, Hamilton SR, et al. Genetic alterations during colorectal-tumor development. *N Engl J Med* 1988;319:525-32.
- Togashi K, Konishi F, Ishizuka T, et al. Efficacy of magnifying endoscopy in the differential diagnosis of neoplastic and non-neoplastic polyps of the large bowel. *Dis Colon Rectum* 1999;42:1602-8.
- Su MY, Ho YP, Chen PC, et al. Magnifying endoscopy with indigo carmine contrast for differential diagnosis of neoplastic and nonneoplastic colonic polyps. *Dig Dis Sci* 2004;49:1123-7.
- Chapuis PH, Dent OF, Goulston KJ. Clinical accuracy in the diagnosis of small polyps using the flexible fiberoptic sigmoidoscope. *Dis Colon Rectum* 1982;25:669-72.
- Neale AV, Demers RY, Budev H, et al. Physician accuracy in diagnosing colorectal polyps. *Dis Colon Rectum* 1987;30:247-50.
- Rembacken BJ, Fujii T, Cairns A, et al. Flat and depressed colonic neoplasms: a prospective study of 1000 colonoscopies in the UK. *Lancet* 2000;355:1211-4.
- Hurlstone DP, Cross SS, Slater R, et al. Detecting diminutive colorectal lesions at colonoscopy: a randomized controlled trial of pan-colonic versus targeted chromoscopy. *Gut* 2004;53:376-80.
- Hurlstone DP, Cross SS, Adam I, et al. Efficacy of high magnification chromoscopic colonoscopy for the diagnosis of neoplasia in flat and depressed lesions of the colorectum: a prospective analysis. *Gut* 2004;53:284-90.
- Konishi K, Kaneko K, Kurahashi T, et al. A comparison of magnifying and nonmagnifying colonoscopy for diagnosis of colorectal polyps: a prospective study. *Gastrointest Endosc* 2003;57:48-53.
- Matsuda T, Fujii T, Saito Y, et al. Efficacy of the invasive/non-invasive pattern by magnifying chromoendoscopy to estimate the depth of invasion of early colorectal neoplasms. *Am J Gastroenterol* 2008;103:2700-6.
- Fu KI, Sano Y, Kato S, et al. Chromoendoscopy using indigo carmine dye spraying with magnifying observation is the most reliable method for differential diagnosis between non-neoplastic and neoplastic colorectal lesions: a prospective study. *Endoscopy* 2004;36:1089-93.
- Sano Y, Muto M, Tajiri H. Optical/digital chromoendoscopy during colonoscopy using narrow-band imaging system. *Digestive Endoscopy* 2005;17:S43-8.
- Machida H, Sano Y, Hamamoto Y, et al. Narrow-band imaging in the diagnosis of colorectal mucosal lesions: a pilot study. *Endoscopy* 2004;36:1094-8.
- East JE, Suzuki N, Stavrinidis M, et al. Narrow band imaging for colonoscopic surveillance in hereditary non-polyposis colorectal cancer. *Gut* 2008;57:65-70.
- Hirata M, Tanaka S, Oka S, et al. Magnifying endoscopy with narrow band imaging for diagnosis of colorectal tumors. *Gastrointest Endosc* 2007;65:988-95.
- Sano Y, Ikematsu H, Fu KI, et al. Meshed capillary vessels by use of narrow-band imaging for differential diagnosis of small colorectal polyps. *Gastrointest Endosc* 2009;69:278-83.
- Katagiri A, Fu KI, Sano Y, et al. Narrow band imaging with magnifying colonoscopy as diagnostic tool for predicting histology of early colorectal neoplasia. *Aliment Pharmacol Ther* 2008;27:1269-74.
- Uraoka T, Saito Y, Matsuda T, et al. Detectability of colorectal neoplastic lesions using a narrow-band imaging system: a pilot study. *J Gastroenterol Hepatol* 2008;23:1810-5.
- Uraoka T, Sano Y, Saito Y, et al. Narrow-band imaging for improving colorectal adenoma detection: appropriate system function settings are required. *Gut* 2009;58:604-5.
- Yoshida T, Inoue H, Usui S, et al. Narrow-band imaging system with magnifying endoscopy for superficial esophageal lesions. *Gastrointest Endosc* 2004;59:288-95.
- Yao K, Oishi T, Matsui T, et al. Novel magnifying endoscopic findings of microvascular architecture in intramucosal gastric cancer. *Gastrointest Endosc* 2002;56:279-84.
- The Paris endoscopic classification of superficial neoplastic lesions: esophagus, stomach, and colon: November 30 to December 1, 2002. *Gastrointest Endosc* 2003;58:S3-43.
- Kudo S, Hirota S, Nakajima T, et al. Colorectal tumours and pit pattern. *J Clin Pathol* 1994;47:880-5.
- Kudo S, Rubio CA, Teixeira CR, et al. Pit pattern in colorectal neoplasia: endoscopic magnifying view. *Endoscopy* 2001;33:367-73.
- Chiu HM, Chang CY, Chen CC, et al. A prospective comparative study of narrow-band imaging, chromoendoscopy, and conventional colonoscopy in the diagnosis of colorectal neoplasia. *Gut* 2007;56:373-9.
- Rogart JN, Jain D, Siddiqui UD, et al. Narrow-band imaging without high magnification to differentiate polyps during real-time colonoscopy: improvement with experience. *Gastrointest Endosc* 2008;68:1136-45.
- Kudo S, Tamura S, Nakajima T, et al. Diagnosis of colorectal tumorous lesions by magnifying endoscopy. *Gastrointest Endosc* 1996;44:8-14.
- Butterly LF, Chase MP, Pohl H, et al. Prevalence of clinically important histology in small adenomas. *Clin Gastroenterol Hepatol* 2006;4:343-8.
- Chen SC, Mouchli A, Chadalaawada V, et al. Histopathology of small polyps removed in the videoendoscopic era [abstract]. *Gastrointest Endosc* 2006;63:AB 199.
- Jass JR, Baker K, Zlobec I, et al. Advanced colorectal polyps with the molecular and morphological features of serrated polyps and adeno-

- mas: concept of a 'fusion' pathway to colorectal cancer. *Histopathology* 2006;49:121-31.
35. Matsumoto T, Mizuno M, Shimizu M, et al. Clinicopathological features of serrated adenoma of the colorectum: comparison with traditional adenoma. *J Clin Pathol* 1999;52:513-6.
 36. Van den Broek FJ, Van Soest EJ, Naber AH, et al. Combining autofluorescence imaging and narrow-band imaging for the differentiation of adenomas from non-neoplastic colonic polyps among experienced and non-experienced endoscopists. *Am J Gastroenterol* 2009;104:1498-507.
 37. Farris AB, Misdraji J, Srivastava A, et al. Sessile serrated adenoma: challenging discrimination from other serrated colonic polyps. *Am J Surg Pathol* 2008;32:30-5.
 38. Sandmeier D, Seelentag W, Bouzourene H. Serrated polyps of the colorectum: is sessile serrated adenoma distinguishable from hyperplastic polyp in a daily practice? *Virchows Arch* 2007;450:613-8.
 39. Mahmood T, Darzi A. The learning curve for a colonoscopy simulator in the absence of any feedback: no feedback, no learning. *Surg Endosc* 2004;18:1224-30.

Receive tables of content by e-mail

To receive tables of content by e-mail, sign up through our Web site at www.giejournal.org.

Instructions

Log on and click "Register" in the upper right-hand corner. After completing the registration process, click on "My Alerts" then "Add Table of Contents Alert." Select the specialty category "Gastroenterology" or type *Gastrointestinal Endoscopy* in the search field and click on the Journal title. The title will then appear in your "Table of Contents Alerts" list.

Alternatively, if you are logged in and have already completed the Registration process, you may add tables of contents alerts by accessing an issue of the Journal and clicking on the "Add TOC Alert" link.

You will receive an e-mail message confirming that you have been added to the mailing list. Note that tables of content e-mails will be sent when a new issue is posted to the Web.

ORIGINAL ARTICLE

Use of Gascon and Pronase either as a pre-endoscopic drink or as targeted endoscopic flushes to improve visibility during gastroscopy: A prospective, randomized, controlled, blinded trial

P. BHANDARI¹, S. GREEN¹, H. HAMANAKA², T. NAKAJIMA², T. MATSUDA², Y. SAITO², I. ODA² & T. GOTODA²

¹Portsmouth Hospitals Trust, Portsmouth, UK, and ²The National Cancer Centre Hospital, Tokyo, Japan

Abstract

Objective. To assess whether endoscopic flushes of the bubble-bursting agent Gascon and the mucolytic agent Pronase are as effective in terms of improving endoscopic mucosal visibility as a pre-endoscopic drink of the same agents. **Material and methods.** A total of 112 patients attending a Japanese tertiary referral centre for upper gastrointestinal endoscopy were randomized to receive either the standard Japanese procedure of a pre-endoscopic drink of water containing Gascon and Pronase with endoscopic flushes of 20-ml aliquots of water, or no pre-endoscopic therapy but endoscopic flushes of 20-ml aliquots of water containing Gascon, with or without Pronase as necessary. **Results.** Visibility scores were significantly better in the pre-endoscopic drink group than in either of the endoscopic flush groups. The group receiving a pre-endoscopic drink required fewer flushes during the procedure and there was no difference in the endoscopic time between the three groups. **Conclusions.** Our results suggest that endoscopic spraying of these bubble-bursting and mucolytic agents is not able to offer equivalent improvements in endoscopic mucosal visibility when compared with the standard Japanese therapy of a pre-endoscopic drink of these agents. The addition of Pronase to the spray solution had no measurable benefit over Gascon alone. We therefore cannot recommend endoscopic spraying of mucous clearing agents over their use as a pre-endoscopic drink.

Key Words: Endoscopy, gascon, mucolytic, pronase, simethicone, visibility

Introduction

Since the advent of gastrointestinal endoscopy, practitioners have been frustrated by foam and mucous obscuring the field of view. Mucosal toileting techniques with bubble-bursting agents such as Gascon (simethicone) have been used since the 1950s [1–3] and more recent studies have shown that the addition of a mucolytic such as Pronase further improves mucosal visualization [4,5]. These mucosal toileting techniques have become standard practice in Japan [6,7], where cancers tend to be detected earlier than in the West. Patients there are routinely asked to drink 100 ml of water containing 2 ml of Gascon and 20,000 units of Pronase 10 min prior to the endoscopy. These medications are freely available in Europe but it is not

usual practice for them to be used. One explanation for this is concern amongst Western endoscopists of an increased risk of aspiration during the procedure if a drink is taken beforehand.

Minimally invasive techniques such as photodynamic therapy and endoscopic mucosal resection (EMR) are now able to offer excellent results for cancers detected at early stages. EMR often offers complete cure but can only be considered for tumours that are well characterized at endoscopy. Detection and characterization of early changes can be achieved through a variety of diagnostic techniques, including chromoendoscopy, high-magnification endoscopy, confocal endoscopy and narrow-band imaging, but all depend upon optimized mucosal views. In addition, chromoendoscopy requires a clear field in order

Correspondence: Susannah Green, Gastroenterology Department, Queen Alexandra Hospital, Portsmouth, PO6 3LY, UK. E-mail: susi@doctors.org.uk

(Received 16 September 2009; accepted 11 November 2009)

ISSN 0036-5521 print/ISSN 1502-7708 online © 2010 Informa UK Ltd.
DOI: 10.3109/00365520903483643

that the dye binds to the intended cells rather than the overlying mucous [8,9]. Effective and acceptable mucosal toileting techniques are therefore increasingly vital as advanced endoscopic techniques become used more frequently.

In an attempt to provide the proven benefits of Gascon and Pronase [9–12] without the theoretical increased risk of pulmonary aspiration associated with a pre-endoscopic drink, this study was designed to compare the effectiveness and practicality of spraying Gascon, with or without Pronase, directly onto the mucosa as intermittent flushes through the biopsy channel of the endoscope during the procedure, compared with identical treatment given as a drink prior to endoscopy (conventional Japanese mucosal toileting).

Material and methods

Patients

The Japanese national screening programme for gastric cancer involves the majority of people over the age of 40 years undergoing an annual barium swallow. The tertiary referral centre in which this trial was set accepts patients for gastroscopy either directly (patients with abnormal results on these tests or

with appropriate symptoms), or as referrals from other hospitals where early cancers have been detected that are thought to be suitable for EMR. This study was restricted to the screening population because there are differences in the endoscopy technique for those requiring a therapeutic procedure (e.g. the use of zoom scopes and special dyes requiring additional time). A total of 148 of these patients were recruited into this study over a 2-week period. Patients were excluded from the study if they had previously undergone oesophagectomy or gastrectomy, if the endoscopy revealed a lesion requiring a therapeutic procedure such as EMR or if there was active gastrointestinal bleeding or strictures in the upper gastrointestinal tract. The results from 112 patients were therefore available for analysis (Figure 1).

Pre-medication and endoscopic procedure

The study gained ethical approval and informed consent was obtained from all participants. Sealed envelopes were used to randomly allocate patients to one of three groups, as follows. Group S: standard Japanese procedure comprising a pre-endoscopic drink of 100 ml of water, 2 ml of Gascon and 20,000 units of Pronase. During the endoscopy, flushes of 20-ml




Figure 1. Flowchart showing the disposition of the study patients.

aliquots of water were used as required. Group G: no pre-endoscopic preparation. During the endoscopy, flushes of 20-ml aliquots of pre-mixed solution containing 100 ml of water and 2 ml of Gascon were used as required. Group GP: no pre-endoscopic preparation was given. During the endoscopy, flushes of 20-ml aliquots of pre-mixed solution containing 100 ml of water, 2 ml of Gascon and 20,000 units of Pronase were used as required.

All patients underwent routine gastroscopy, including chromoendoscopy, by one of 14 experienced unblinded endoscopists. The endoscopist was free to use as many flushes as deemed necessary to produce a satisfactory view. Once all flushes had been given, one extra photograph was taken from each of four pre-defined areas: the oesophagogastric junction, the antrum, the lower body and the upper body of the stomach. A record was kept of the total time taken to perform the procedure (from intubation to extubation) and the number of flushes required.

A single, blinded investigator who was experienced in endoscopy but had played no part in the endoscopic procedure then reviewed all of the pictures and assigned each of them a score between one and three for mucosal visibility: 1 = no adherent mucus and clear view of the mucosa; 2 = a thin coating of mucus but not obscuring vision; and 3 = adherent mucus obscuring vision.

The individual scores for each of the four photographs taken were then totalled for each patient to give an overall visibility score ranging from four to 12.

A second blinded investigator separately reviewed and scored the pictures from 20 patients and the results were compared with the original assigned scores.




Figure 2. Outcomes.

Statistical analysis

The sample-size calculations showed that 35 participants were required in each treatment group (105 patients overall) to detect a 20% improvement in visibility scores, from 7 to 5.6, assuming a standard deviation of 2 for each group and a power of 90%. Allowing for a 30% attrition rate, we aimed to recruit 150 participants.

Differences between the number of flushes and the time taken were analysed using ANOVA and Fisher's least significant difference. As visibility scores were non-normally distributed, the Kruskal-Wallis and Dwass-Steel-Chritchlow-Fligner tests were used for these results. All analyses used SPSS software (SPSS Inc, Chicago, IL). A P-value of 0.05 was taken to be significant throughout.

Results

A total of 112 patients were evaluable in the study, with a mean age of 61 years. The study population comprised 51 males (46%) and 61 females (54%). There were no significant differences between treatment groups (Table I) for a summary of outcome measures please see Figure 2.

Visibility

Visibility scores allocated by the two independent visibility score assessors correlated well (Cohen's weighted kappa 0.604, standard error 0.187, 95% CI 0.237-0.971).

There were significant differences in the visibility scores assigned between groups (H = 17.8, P = 0.0001). The photographs taken from the pre-medicated Group S scored significantly better for visibility than either of the endoscopic therapy groups GP and G (P = 0.0002 and P = 0.0008, respectively). There was no significant difference in visibility scores between Groups GP and G (P = 0.999).

Table I. Patient characteristics.

Characteristic	Group		
	S (n = 35)	G (n = 37)	GP (n = 40)
Gender; n (%)			
Male	18 (51)	14 (38)	19 (48)
Female	17 (49)	23 (62)	21 (52)
Age (years); mean (SD)	63 (1.9)	61 (1.6)	61 (2.1)

Number of flushes needed

There were significant differences in the mean number of flushes used between groups ($F = 12$, $P = 0.0001$). Significantly fewer flushes were used during the procedure in those patients receiving conventional Japanese pre-medication (Group S) than either of the other groups (Group GP, $P = 0.008$; Group G, $P < 0.001$). In the groups receiving endoscopic flush therapy only, significantly fewer flushes were used in the group with Pronase added to the Gascon mixture ($P = 0.023$).

Time taken for procedure

There was no significant difference in the time taken to complete the procedure between any of the three groups ($F = 2.23$, $P = 0.112$).

Safety

There were no complications in any of the groups. In particular, there were no clinically detectable cases of pulmonary aspiration.

Discussion

Optimal mucosal visualization is vital for thorough endoscopic inspection, particularly when using newer methods such as chromoendoscopy [13–16]. The use of bubble-bursting agents and mucolytics has been shown to improve mucosal visibility in previous trials [17–20], but safety concerns have discouraged generalized use in the West.

We assessed a potentially more acceptable technique of spraying these agents endoscopically. Gascon (simethicone or dimethicone) is silicone-based and non-absorbable, with an excellent safety record. It causes gas bubbles to burst by reducing their surface tension and is marketed for the relief of abdominal bloating. Pronase is a mixture of proteases isolated from *Streptomyces griseus*. These agents were chosen for the study as they both have proven efficacy and have been adopted as standard treatment at the trial centre.

Our results showed that spraying the anti-foam and mucolytic agents endoscopically was not as effective in terms of improved mucosal visibility as pre-endoscopic treatment with the same combination, despite the endoscopist using a greater number of flushes to attempt to clear the mucous. We would ideally have compared the endoscopic flushes with Western standard practice, which in the UK would be

to give no pre-endoscopic preparation and to use water endoscopic flushes, but were unable to do this in Japan as using mucous-clearing medication has become so accepted that it was considered unethical not to do so. Adding Pronase to the basic endoscopic flush mixture did not add any advantage in terms of mucosal visibility. The apparent superiority of a pre-endoscopic drink of mucous-clearing solution as compared to endoscopic flush therapy may reflect the more diffuse application of the solution or the 10-min delay between the drink and endoscopy.

No technique resulted in clinically detectable pulmonary aspiration but rates of aspiration during a standard gastroscopy are less than one in a thousand [21] and a larger trial would therefore be needed to properly evaluate this risk.

We conclude that the standard Japanese practice of administering a pre-endoscopic drink containing a mucolytic and anti-bubble agent is superior in terms of endoscopic mucosal visibility to endoscopic application of either both agents or an anti-bubble agent alone. We cannot recommend applying these agents as an endoscopic spray.

Whether improved mucosal visibility results in a higher detection rate of early cancers or improved clinical outcomes remains unknown and well-designed large clinical trials will be needed in the future to evaluate this.

Acknowledgement

We thank Mr Bernie Higgins for his statistical support.

Declaration of interest: The authors report no conflicts of interest. The authors alone are responsible for the content and writing of the paper.

References

- [1] Gasster M, Westwater JO, Molle WE. Use of a defoaming agent in gastroscopy. *Gastroenterology* 1954;27:652–5.
- [2] Hirschowitz BI, Bolt RJ, Pollard MH. Defoaming in gastroscopy with silicone. *Gastroenterology* 1954;27:649–51.
- [3] Dailey ME, Rider JA. Silicone anti-foam tablet in gastroscopy. *JAMA* 1954;155:859.
- [4] Fujii T, Iishi H, Tatsuta M, Hirasawa R, Uedo N, Hifumi K, et al. Effectiveness of premedication with pronase for improving visibility during gastroendoscopy: a randomized controlled trial. *Gastrointest Endosc* 1998;47:382–7.
- [5] Fujii T, Iishi H, Tatsuta M, Hirasawa R, Uedo N, Hifumi K, et al. Effectiveness of premedication with pronase for improving visibility during gastroendoscopy: a randomized controlled trial. *Gastrointest Endosc* 1998;47:382–7.

- [6] Misumi A, Harada K, Murakami A, Arima K, Kondo H, Akagi M, et al. Role of Lugol dye endoscopy in the diagnosis of early esophageal cancer. *Endoscopy* 1990;22:12-16.
- [7] Fujita R, Yamamura H. Preparation of patients for gastroscopic examination. In: Takemoto T, editor. *The manual of gastroenterological endoscopy*. Tokyo: Nankodo; 1982.
- [8] Shaw D, Blair V, Framp A, Harawira P, McLeod M, Guilford P, et al. Chromoendoscopic surveillance in hereditary diffuse gastric cancer: an alternative to prophylactic gastrectomy? *Gut* 2005;54:461-8.
- [9] Tamura S, Ookawauchi K, Onishi S, Yokoyama Y, Yamada T, Higashidani Y, et al. The usefulness of magnifying chromoendoscopy: pit pattern diagnosis can predict histopathological diagnosis precisely. *Am J Gastroenterol* 2002;97:2934-5.
- [10] Kuo CH, Sheu BS, Kao AW, Wu CH, Chuang CH. A defoaming agent should be used with pronase premedication to improve visibility in upper gastrointestinal endoscopy. *Endoscopy* 2002;34:531-4.
- [11] Yiangpruksawan A, Lightdale CJ, Gerdes H, Botet JF. Mucolytic-antifoam solution for reduction of artifacts during endoscopic ultrasonography: a randomized controlled trial. *Gastrointest Endosc* 1991;37:543-6.
- [12] Kuo CH, Sheu BS, Kao AW, Wu CH, Chuang CH. A defoaming agent should be used with pronase premedication to improve visibility in upper gastrointestinal endoscopy. *Endoscopy* 2002;34:531-4.
- [13] Shaw D, Blair V, Framp A, Harawira P, McLeod M, Guilford P, et al. Chromoendoscopic surveillance in hereditary diffuse gastric cancer: an alternative to prophylactic gastrectomy? *Gut* 2005;54:461-8.
- [14] Denis-Ribeiro M, da Costa-Pereira A, Lopes C, Lara-Santos L, Guilherme M, Moreira-Dias L, et al. Magnification chromoendoscopy for the diagnosis of gastric intestinal metaplasia and dysplasia. *Gastrointest Endosc* 2003;57:498-504.
- [15] Tamura S, Ookawauchi K, Onishi S, Yokoyama Y, Yamada T, Higashidani Y, et al. The usefulness of magnifying chromoendoscopy: pit pattern diagnosis can predict histopathological diagnosis precisely. *Am J Gastroenterol* 2002;97:2934-5.
- [16] Pathirana A, Poston GJ. Lessons from Japan—endoscopic management of early gastric and oesophageal cancer. *Eur J Surg Oncol* 2001;27:9-16.
- [17] Banerjee B, Parker J, Waits W, Davis B. Effectiveness of preprocedure simethicone drink in improving visibility during esophagogastroduodenoscopy: a double-blind, randomized study. *J Clin Gastroenterol* 1992;15:264-5.
- [18] Bertoni G, Gumina C, Conigliaro R, Ricci E, Staffetti J, Mortilla MG, et al. Randomized placebo-controlled trial of oral liquid simethicone prior to upper gastrointestinal endoscopy. *Endoscopy* 1992;24:268-70.
- [19] Fujii T, Iishi H, Tatsuta M, Hirasawa R, Uedo N, Hifumi K, et al. Effectiveness of premedication with pronase for improving visibility during gastroendoscopy: a randomized controlled trial. *Gastrointest Endosc* 1998;47:382-7.
- [20] Kuo CH, Sheu BS, Kao AW, Wu CH, Chuang CH. A defoaming agent should be used with pronase premedication to improve visibility in upper gastrointestinal endoscopy. *Endoscopy* 2002;34:531-4.
- [21] Clarke AC, Chiragakis L, Hillman LC, Kaye GL. Sedation for endoscopy: the safe use of propofol by general practitioner sedationists. *Med J Aust* 2002;176:158-61.

INDICATIONS FOR ENDOSCOPIC RESECTION OF COLORECTAL POLYPS AND SURVEILLANCE GUIDELINES

LOCAL RECURRENCE AND SURVEILLANCE AFTER ENDOSCOPIC RESECTION OF LARGE COLORECTAL TUMORS

KINICHI HOTTA,¹ YUTAKA SAITO,² TAKAHISA MATSUDA,² TOMOAKI SHINOHARA¹ AND TSUNEO OYAMA¹

¹Department of Gastroenterology, Saku Central Hospital, Nagano and ²Endoscopy Division, National Cancer Center Hospital, Tokyo, Japan

Local recurrence rates after endoscopic piecemeal mucosal resection (EPMR) typically range from 10 to 23%. In our previous study, the local recurrence rate after a piecemeal resection was significantly higher than that after an en bloc resection, irrespective of tumor size or macroscopic features. To reduce local recurrence after an EPMR, it is important to carefully note the circumferences of the edge and base of the ulcer. Recently, endoscopic submucosal dissection (ESD) was developed and recognized for its effectiveness in large, complete, en bloc resections and precise pathological assessments. ESD also showed lower local recurrence rates, ranging from 0 to 3% in previous, retrospective studies. However, ESD showed a higher perforation rate and longer procedure times; thus, it is necessary to improve ESD. An appropriate surveillance interval after EPMR was still controversial, and recommendations of some guidelines ranged from 2 to 9 months. In order to determine the appropriate interval, a randomized controlled study is necessary.

Key words: colorectal tumor, endoscopic mucosal resection, endoscopic submucosal dissection, local recurrence, surveillance.

INTRODUCTION

Both the incidence and mortality of colorectal tumors have increased recently; currently, colorectal cancers are the first and fourth leading causes of cancer mortality in Japanese women and men, respectively.¹ Large colorectal tumors are typically defined as ≥ 20 mm in diameter.^{2–4} Some large colorectal tumors are adenomas or non-invasive cancers that can be treated successfully with endoscopic resection. Laterally spreading tumor (LST) were described as extending laterally, rather than vertically, and tended to remain in the mucosa.^{5,6} Several issues need to be considered in the treatment of large colorectal tumors, including the indication for endoscopic resection, selection of an endoscopic treatment method, the risk of local recurrence, and the surveillance interval. Here, we present a review of the literature and discuss these issues.

INDICATION OF ENDOSCOPIC RESECTIONS

Endoscopic resection is indicated for early colorectal tumors that show negligible risk of lymph node metastases. The conditions for lymph node metastases were studied in a Japanese multicenter survey of colorectal cancers.⁷ Based on the report, the pathological conditions that indicated no or low

risk of lymph node metastasis included a shallow invasion depth ($< 1000 \mu\text{m}$), no lymphatic invasion, and no sprouting.⁷ Of these factors, only the invasion depth can be estimated before treatment. A biopsy is undesirable because it may complicate an endoscopic resection.⁸ Furthermore, it is difficult to precisely diagnose the depth of invasion based on a biopsy. In general, a conventional, white light, endoscopic evaluation is used to assess early colorectal cancers; findings of hardness, fold convergence, depression, and irregular shape are considered indicative of submucosal invasion.⁹ In pedunculated lesions, a thick stalk and jagged shape are important indications of stalk invasions. However, the accuracy of estimating the invasion depth with conventional endoscopy is insufficient for determining an appropriate therapeutic method that avoids excessive surgery. In our opinion, the most reliable method for predicting invasion depth is magnified chromoendoscopy with crystal violet staining. A pit pattern classification proposed by Kudo and Tsuruta has been adopted by Japanese endoscopists.¹⁰ Type V pit patterns, particularly the VN type pit pattern, are recognized as indications of submucosal invasion.¹⁰ Fujii and colleagues proposed a clinical classification of invasive or non-invasive patterns, taking into account the demarcated area with irregular or distorted pits.¹¹ We previously reported that the invasive pattern could differentiate between intramucosal or submucosal superficial cancers ($< 1000 \mu\text{m}$) and submucosal deep cancers ($\geq 1000 \mu\text{m}$) with sensitivity, specificity, and accuracy of 85.6%, 99.4%, and 98.8%, respectively (Fig. 1).¹¹ In a multivariable analysis of factors that predicted submucosal deep invasion of non-granular type LST, the invasive pattern was considered a risk factor, together with hardness and large tumor size (≥ 20 mm).¹² Narrow band

Correspondence: Kinichi Hotta, Saku Central Hospital, Department of Gastroenterology, 197 Usuda, Saku, Nagano, 384-0301, Japan. Email: kinichi1@janis.or.jp

Conflicts of interest: The authors declare no potential conflicts of interest.

Received 14 December 2009; accepted 8 January 2010.

© 2010 The Authors

Journal compilation © 2010 Japan Gastroenterological Endoscopy Society




Fig. 1. A case of an invasive pattern. (a) Conventional colonoscopy showed a laterally spreading granular type tumor, with a reddish depressed area, 30 mm in diameter, in the cecum. (b) Chromoendoscopy with indigo-carmin spray dyeing showed a demarcated area traced by yellow dotted line. (c,d) Magnified chromoendoscopy with crystal violet staining showed invasive pattern in the demarcated area. The lesion was treated with laparoscopic surgery and pathological diagnosis was well-differentiated adenocarcinoma with submucosal invasion (2000 μm).

imaging (NBI) was recently assessed for predicting tumor invasion depth.¹³ NBI offers the advantage of a simple, easy method; but currently, its diagnostic accuracy may be inferior to magnified endoscopy with crystal violet staining.¹³ Endoscopic ultrasonography was also used for predicting tumor invasion depth, but it requires a higher level of skill and diagnosis is difficult, even in good conditions. Our previous controlled evaluation showed that magnified endoscopy was superior to endoscopic ultrasonography for the estimation of tumor invasion depth.¹⁴

SELECTION OF ENDOSCOPIC TREATMENT METHODS

The choice of endoscopic treatment methods depends on lesion size and characteristics; they include snare polypectomy, endoscopic mucosal resection (EMR), endoscopic piecemeal mucosal resection (EPMR), or endoscopic submucosal dissection (ESD). We propose that ESD is indicated for non-granular type LST (≥ 20 mm), due to the relatively high rates of submucosal invasion and the difficulty in predicting the invasion site prior to treatment.¹⁵ Moreover, ESD is indicated for granular type LST, particularly mixed nodular types (≥ 40 mm), also due to relatively high rates of submucosal invasion.¹⁵ Alternatively, EPMR is indicated for granular type LST (homogeneous type), due to the similarity to adenoma or intramucosal cancer. Recently, a working group for the standardization of colorectal ESD proposed that ESD is indicated for colorectal tumors with the following features: large lesions (≥ 20 mm in diameter) that are difficult to resect en bloc with a snare EMR, but where an endoscopic treatment is indicated; mucosal lesions with fibrosis caused by

prolapse, due to biopsy or peristalsis of the lesion; sporadic localized tumors associated with chronic inflammation, e.g. ulcerative colitis; and local or residual early cancer after an endoscopic resection.¹⁶ For pedunculated lesions, snare polypectomy of an EMR, combined with looping or clipping, is indicated when there is no endoscopic finding of stalk invasion. One controlled trial suggested that a bleeding rate was lower with combination epinephrine injection plus endo-loop than epinephrine injection alone.¹⁷

LOCAL RECURRENCE AFTER ENDOSCOPIC RESECTION

After endoscopic resection local recurrence is an important issue in conventional EMR/EPMR methods (Fig. 2). Local recurrence rates after EPMR typically range from 10 to 23%^{2,4,18-20} (Table 1). In our previous study, the local recurrence rate after a piecemeal resection was significantly higher than that after an en bloc resection, irrespective of tumor size or macroscopic features.²⁰ To reduce local recurrence after an EPMR, it is important to carefully note the circumferences of the edge and base of the ulcer. Magnified observation is ideal for detecting a residual tumor. Tanaka *et al.* reported that magnified observation after an EPMR could effectively reduce the local recurrence rate.¹⁶ Argon plasma coagulation (APC) was also tested for reducing the local recurrence after EPMR. One controlled trial suggested that APC reduced the local recurrence after EPMR of large sessile tumors.²¹ However, in a prospective uncontrolled study, similar recurrence rates were found with or without APC after an EPMR.²² Thus, the usefulness of APC is controversial, but in some high-volume centers, it is routinely