

DHCA (cerebral circulatory arrest) on its own. Thus our two modifications, used alongside DHCA, minimized the risk of cerebral embolism and allowed for complete aortic resection, including the aortic cross-clamp site after thoracic aortic replacement through a left lateral approach. This method will be applicable particularly to patients at high risk of ischemic and embolic complications, such as those with severely athermanous aortas and complex aortic pathology prohibiting proximal cross-clamping.

The importance of antegrade aortic perfusion and the aortic cannulation site has been underestimated during thoracic and thoracoabdominal aortic surgery through the left lateral approach. Cerebral complications occur in 9% of patients after thoracic and thoracoabdominal aortic replacement using DHCA [1]. Although the femoral artery is generally chosen as an aortic cannulation site, aortic perfusion from the femoral artery results in retrograde aortic perfusion, which carries a risk of emboli entering into the arch branches and malperfusion of the dissected aorta. However, little is known about differences in cerebral complications in relation to the aortic cannulation site used in the left lateral approach. We have previously shown that antegrade aortic perfusion by AAC decreases the incidence of neurologic deficit to 7% compared with 28% using retrograde aortic perfusion by femoral artery cannulation [8]. Based on this evidence, we advocate using AAC in the left lateral approach.

In addition to AAC, under DHCA, we also established EEP during the open proximal anastomosis at the distal arch. Although, in part, EEP is similar to the original retrograde cerebral perfusion (RCP) reported by Ueda and colleagues [9] and Takamoto and colleagues [10], the method and degree of increase in CVP, and the purpose, all differ. Thus, in order to increase cerebral perfusion and prevent embolism, RCP is performed by elevating CVP over 15 mm Hg and this high level has been associated with postoperative cerebral edema leading to neurologic dysfunction. In contrast, the EEP used in this series was induced by aortic distal perfusion toward the lower torso from the single side branch of the distal graft. The aortic distal perfusion was also able to maintain antegrade aortic perfusion resulting in backward flow from the arch branches. The CVP levels were maintained between 3 and 7 mm Hg by controlling venous drainage and thus were relatively low compared with RCP. We would emphasize that this technique was used not to increase the cerebral blood flow during DHCA but to prevent emboli entering the arch branches. Importantly, there was no neurologic deficit in the brain postsurgery in our series. Although the degree of cerebral edema and neurologic dysfunction after EEP are still under investigation, this technique could be a safe, simple, and feasible method to prevent embolic stroke after aortic distal arch replacement.

In our surgical procedures we used two cannulation sites for aortic return: AAC and the side branch of the distal graft. The aortic distal perfusion toward the lower torso caused backward flow from the intercostal arteries as well as EEP. The backward flow comes through the cervicothoracic and thoracolumbar arterial networks [11,

12]. Thus, the aortic perfusion using these two cannulation sites alternately allowed us to maintain spinal cord perfusion without interruption of networks. Furthermore, based on the collateral network to the spinal cord, we were able to sacrifice all intercostal arteries without reimplantation. The region between T7 and L1, in which the Adamkiewicz artery is considered to exist, was included in 85% of patients. Etz and colleagues [13] reported postoperative paraplegia in 2 of 100 consecutive patients undergoing sacrifice of intercostal and lumbar arteries during thoracic and thoracoabdominal repair using various CPB techniques. Our results were similar to those in recent reports. In this study, although temporary neurologic deficit in the spinal cord occurred in 3 patients (7.7%), no patient had permanent neurologic deficit. Thus, alternate aortic perfusion from these two cannulation sites was effective in reducing the risk of permanent neurologic deficit in the spinal cord.

Crystalloid potassium and blood cardioplegic solutions are generally used for myocardial protection during ascending and aortic root surgery through a median sternotomy approach. However, little is known about myocardial protection during aortic surgery through the left lateral approach. Takamoto and colleagues [10] infused crystalloid potassium cardioplegic solution using an occlusion balloon in the left lateral approach. However, in our series we maintained hypothermic ventricular fibrillation during surgery without using cardioplegic solution. Cosseli and colleagues [1] have recently reported a method similar to ours. In their report, cardiac complications occurred at 2% and 24% in elective and emergent cases, respectively. In this study, 97.5% of the procedures were performed in elective settings and one patient (2.5%) had frequent ventricular tachycardia at the time of weaning from CPB. This accumulated evidence suggests that deep hypothermic ventricular fibrillation is an acceptable means of myocardial protection during thoracic aortic surgery through the left lateral approach. Further postoperative investigations, such as changes in biochemical markers and the analysis of arrhythmic events will be required to address the feasibility of hypothermic ventricular fibrillation myocardial protection.

Although ischemia-reperfusion injury is inevitable after surgery using DHCA, careful respiratory management should be taken throughout the postoperative period. Tracheotomy is required between 2% and 11% of cases [1, 5, 14] and was needed in two cases (5.0%) in this study. On average, it took 2.6 ± 1.7 days to achieve tracheal extubation; the length of time that the patients required mechanical ventilation was thus relatively longer than in other recent reports [14]. Several factors, such as prolonged CPB time, neurocognitive dysfunction, and chronic obstructive pulmonary disease, are considered as risk factors for prolonged mechanical ventilation after aortic surgery using DHCA [14, 15]. The CPB time in this study was also longer than in previous reports, although RCP time and DHCA time of the lower torso were less than 40 minutes. The size of the venous cannula through the femoral vein was usually 24 Fr, and this small

size may require much more time for cooling and re-warming of body temperature. The prolonged CPB time may thus be associated with the prolonged mechanical ventilation. On the other hand, permanent hemodialysis is required in 1.0% to 1.2% of elective patients who were not on chronic dialysis before surgery [1, 5]. In our series, no patients needed permanent hemodialysis.

The small number of patients was a limitation in this study. Furthermore, compared with previous reports the characteristics of our patients were relatively mild (eg, number on hemodialysis and those requiring emergency surgery) [1, 5]. These factors may be associated with the superior clinical results of this series. Accumulated clinical experience, as well as long-term follow-up, will be required to estimate the effect of minimizing the risk of cerebral and spinal cord embolism using this technique.

Combined with deep hypothermia and DHCA, AAC and EEP minimized the risk of cerebral embolism in aortic distal arch replacement through the left lateral approach.

References

1. Coselli JS, Bozinovski J, Cheung C. Hypothermic circulatory arrest: safety and efficacy in the operative treatment of descending and thoracoabdominal aortic aneurysms. *Ann Thorac Surg* 2008;85:956-64.
2. Gega A, Rizzo JA, Johnson MH, Tranquilli M, Farkas EA, Elefteriades JA. Straight deep hypothermic arrest: experience in 394 patients supports its effectiveness as a sole means of brain preservation. *Ann Thorac Surg* 2007;84:759-67.
3. Kouchoukos NT, Masetti P, Rokkas CK, Murphy SF. Hypothermic cardiopulmonary bypass and circulatory arrest for operations on the descending thoracic and thoracoabdominal aorta. *Ann Thorac Surg* 2002;74:S1885-7; discussion S1892-8.
4. Minatoya K, Karck M, Hagl C, et al. The impact of spinal angiography on the neurological outcome after surgery on the descending thoracic and thoracoabdominal aorta. *Ann Thorac Surg* 2002;74:S1870-2; discussion S1892-8.
5. Fehrenbacher JW, Hart DW, Huddleston E, Siderys H, Rice C. Optimal end-organ protection for thoracic and thoracoabdominal aortic aneurysm repair using deep hypothermic circulatory arrest. *Ann Thorac Surg* 2007;83:1041-6.
6. Westaby S, Katsumata T. Proximal aortic perfusion for complex arch and descending aortic disease. *J Thorac Cardiovasc Surg* 1998;115:162-7.
7. Katsumata T, Westaby S. Proximal aortic perfusion with passive cerebral flow: a method to prevent cerebral embolism in the arch and descending aortic operation. *J Card Surg* 1999;14:194-6.
8. Westaby S, Katsumata T, Vaccari G. Arch and descending aortic aneurysms: influence of perfusion technique on neurological outcome. *Eur J Cardiothorac Surg* 1999;15:180-5.
9. Ueda Y, Miki S, Kusuhara K, Okita Y, Tahata T, Yamanaka K. Surgical treatment of aneurysm or dissection involving the ascending aorta and aortic arch, utilizing circulatory arrest and retrograde cerebral perfusion. *J Cardiovasc Surg (Torino)* 1990;31:553-8.
10. Takamoto S, Matsuda T, Harada M, Shimamura Y, Miyata S. Simple hypothermic retrograde cerebral perfusion during aortic arch surgery. *J Cardiovasc Surg (Torino)* 1992;33:560-7.
11. Griep RB, Griep EB. Spinal cord perfusion and protection during descending thoracic and thoracoabdominal aortic surgery: the collateral network concept. *Ann Thorac Surg* 2007;83:S865-9; discussion S890-2.
12. Lazorthes G, Gouaze A, Zadeh JO, Santini JJ, Lazorthes Y, Burdin P. Arterial vascularization of the spinal cord; Recent studies of the anastomotic substitution pathways. *J Neurosurg* 1971;35:253-62.
13. Etz CD, Halstead JC, Spielvogel D, et al. Thoracic and thoracoabdominal aneurysm repair: is reimplantation of spinal cord arteries a waste of time? *Ann Thorac Surg* 2006;82:1670-7.
14. Etz CD, Di Luozzo G, Bello R, et al. Pulmonary complications after descending thoracic and thoracoabdominal aortic aneurysm repair: predictors, prevention, and treatment. *Ann Thorac Surg* 2007;83:S870-6; discussion S890-2.
15. Krahenbuhl ES, Immer FF, Stalder M, Englberger L, Eckstein FS, Carrel TP. Temporary neurological dysfunction after surgery of the thoracic aorta: a predictor of poor outcome and impaired quality of life. *Eur J Cardiothorac Surg* 2008;33:1025-9.

Sodium 4-phenylbutyrate protects against spinal cord ischemia by inhibition of endoplasmic reticulum stress

Taketomo Mizukami, MD, Kazumasa Orihashi, MD, Bagus Herlambang, MD, Shinya Takahashi, MD, Makoto Hamaishi, MD, Kenji Okada, MD, and Taijiro Sueda, PhD, *Hiroshima, Japan*

Objective: Delayed paraplegia after operation on the thoracoabdominal aorta is considered to be related to vulnerability of motor neurons to ischemia. Previous studies have demonstrated the relationship between neuronal vulnerability and endoplasmic reticulum (ER) stress after transient ischemia in the spinal cord. The aim of this study was to investigate whether sodium 4-phenylbutyrate (PBA), a chemical chaperone that reduces the load of mutant or unfolded proteins retained in the ER during cellular stress, can protect against ischemic spinal cord damage.

Methods: Spinal cord ischemia was induced in rabbits by direct aortic cross-clamping (below the renal artery and above the bifurcation) for 15 minutes at normothermia. Group A (n = 6) was a sham operation control group. In group B (n = 6) and group C (n = 6), vehicle or 15 mg/kg/h of sodium 4-PBA was infused intravenously, respectively, from 30 minutes before the induction of ischemia until 30 minutes after reperfusion. Neurologic function was assessed at 8 hours, and 2 and 7 days after reperfusion with a Tarlov score. Histologic changes were studied with hematoxylin-eosin staining. Immunohistochemistry analysis for ER stress-related molecules, including caspase12 and GRP78 were examined.

Results: The mean Tarlov scores were 4.0 in every group at 8 hours, but were 4.0, 2.5, and 3.9 at 2 days; and 4.0, 0.7, and 4.0 at 7 days in groups A, B, and C, respectively. The numbers of intact motor neurons at 7 days after reperfusion were 47.4, 21.5, and 44.9 in groups A, B, and C, respectively. There was no significant difference in terms of viable neurons between groups A and C. Caspase12 and GRP78 immunoreactivities were induced in motor neurons in group B, whereas they were not observed in groups A and C.

Conclusion: Reduction in ER stress-induced spinal cord injury was achieved by the administration of 4-PBA. 4-PBA may be a strong candidate for use as a therapeutic agent in the treatment of ischemic spinal cord injury. (*J Vasc Surg* 2010; 52:1580-6.)

Clinical Relevance: Spinal cord injury following surgical repair of thoracic or thoracoabdominal aorta is a disastrous complication. Previous studies have demonstrated the relationship between neuronal vulnerability and endoplasmic reticulum (ER) stress after transient ischemia of spinal cord. Sodium 4-phenylbutyrate (4-PBA) is a low molecular weight fatty acid that has been approved for clinical use as an ammonia scavenger in children with urea cycle disorders and for treatment of sickle cell disease and thalassemia. A number of investigators have recently reported utilization of 4-PBA as a chemical chaperone to reverse the mislocalization and/or aggregation of proteins associated with human disease. The 4-PBA of therapeutic dose has low toxicity and uniquely penetrates well into cerebrospinal fluid. In this study, we investigated whether intravenous administration of 4-PBA is beneficial to protecting the spinal cord against ischemic damage in a rabbit model. Spinal cord ischemia was induced by direct aortic cross-clamping while 4-PBA of therapeutic dose was infused intravenously. In this simple and less invasive model, reduction in ER stress-induced spinal cord damage was achieved by intravenous 4-PBA. Our results indicate that 4-PBA of therapeutic dose can be a promising strategy, which is clinically feasible without significant adverse effects. Although further investigations are mandatory before it is applied to the clinical practice, such as optimal timing or route for administration 4-PBA, intravenous administration of 4-PBA may be a new candidate as a therapeutic agent for protecting the spinal cord against ischemic damage in thoracoabdominal aortic surgery.

Paraplegia caused by ischemic spinal cord injury is one of the most serious complications of surgery performed on the thoracic and thoracoabdominal aorta, with a reported prevalence of 3% to 23% in different series.¹⁻³ Various adjunctive

procedures for spinal cord protection have been introduced, including cerebrospinal fluid drainage, distal perfusion of the aorta, or use of steroid, barbiturate, or free-radical scavengers, and have successfully reduced the incidence of postoperative paraplegia in recent years. However, postoperative paraplegia remains a devastating complication, and its occurrence is still somewhat unpredictable. Patients undergoing thoracic aneurysm repair who awake without neurologic deficit immediately after the operation may eventually develop paraplegia.^{4,5} The exact mechanism of such a delayed vulnerability is not fully understood. In a rabbit spinal cord ischemia model, delayed and selective motor neuron death after transient ischemia has been reported, and it was greatly associated with activated apoptotic signals.⁶

From the Department of Surgery, Division of Clinical Medical Science, Graduate School of Biomedical Sciences, Hiroshima University.

Competition of interest: none.

Reprint requests: Taketomo Mizukami, MD, Department of Surgery, Division of Clinical Medical Science, Graduate School of Biomedical Sciences, Hiroshima University, Kasumi 1-2-3, Minami-ku, Hiroshima, Japan (e-mail: tmizukamimd@yahoo.co.jp).

The editors and reviewers of this article have no relevant financial relationships to disclose per the JVS policy that requires reviewers to decline review of any manuscript for which they may have a competition of interest.

0741-5214/\$36.00

Copyright © 2010 by the Society for Vascular Surgery.

doi:10.1016/j.jvs.2010.06.172

1580

There is evidence to suggest that the activation of apoptosis is initiated by cell-surface receptors or by mitochondrial stress.⁷ Recently, another apoptotic-regulatory pathway involved in endoplasmic reticulum (ER) stress has received attention. In the case of mild ER stress, cells develop a self-protective, signal transduction pathway termed the unfolded protein response, which includes induction of molecular chaperones in the ER such as glucose regulated protein 78 (GRP78), translational attenuation, and ER-associated protein degradation.^{8,9} However, if the damage is severe, the unfolded protein response ultimately triggers the apoptosis pathway.¹⁰ One of the important molecules relevant to cell death is caspase12 which is regarded to be a representative cell death signal in ER stress.¹¹ Sodium 4-phenylbutyrate (4-PBA) is a low molecular weight fatty acid that has been approved for clinical use as an ammonia scavenger in children with urea cycle disorders and for treatment of sickle cell disease and thalassemia on the basis of its capacity to activate transcription of β - and γ -globin.^{12,13} 4-PBA has been found to prolong life and contribute to therapy for spinal muscular atrophy by altering the pattern of gene expression. In the last few years, a number of studies have described the use of 4-PBA as a chemical chaperone to reverse the mislocalization and/or aggregation of proteins associated with human disease. 4-PBA can act as a chemical chaperone by reducing the load of mutant or mislocated proteins retained in the ER.¹⁴⁻¹⁶ Evidence obtained in recent years has demonstrated that endoplasmic reticulum (ER)-mediated cell death plays an important role in the mechanisms underlying ischemia-reperfusion neuronal damage. Therefore, targeting the ER may provide a therapeutic approach for blocking the pathologic process induced by spinal cord ischemia. However, to our knowledge, no pharmacologic approach for treating spinal cord ischemia-induced ER dysfunction has been reported. In this study, we investigated the protective effects of 4-PBA after transient spinal cord ischemia, apoptotic signaling of caspase12, and induction of GRP78.

MATERIALS AND METHODS

Animal preparation. Animal care and all procedures were performed in compliance with the *Guide for the Care and Use of Laboratory Animals* prepared by the Institute of Laboratory Animal Resources, National Research Council, and published by the National Academy Press, revised 1996. This study protocol was approved by the Research Facilities for Laboratory Animal Science of the Hiroshima University School of Medicine.

Anesthesia and monitoring. We used Japanese female white rabbits (2.5-3.0 kg) in this experiment. General anesthesia was induced with an intramuscular injection of ketamine 50 mg/kg. An ear vein catheter was inserted for the continuous intravenous infusion of lactated Ringer solution. Cefazolin (10 mg/kg) was injected through the catheter. After a tracheostomy and endotracheal intubation, the rabbits were ventilated mechanically with a frac-

tion of inspired oxygen of 1.0 and isoflurane (1.5%-2.0%) to induce sufficient anesthesia. Adequacy of ventilation was monitored by blood gas analysis. The arterial blood pressure was measured by a catheter (JMS cut-down tube; C3, Hiroshima, Japan) placed into the right femoral artery. Electrocardiogram was also monitored continuously. The operation was performed at normothermia. The rectal temperature was continuously monitored and kept at $38.0 \pm 0.5^\circ\text{C}$ during the operation using a heating blanket.

Surgical procedures. With animals in the right lateral decubitus position, a flank skin incision parallel to the spine was made on the left side. From a retroperitoneal approach, the abdominal aorta was exposed below the left renal artery and above the iliac bifurcation, and was encircled. After the intravenous administration of heparin (100 U/kg), the abdominal aorta just distal to the left renal artery and just proximal to the iliac bifurcation was cross-clamped and isolated for 15 minutes to produce spinal cord ischemia. To determine the clamping time, we conducted preliminary experiments. Fifteen rabbits were divided into three groups with clamping times of 10, 12, and 15 minutes ($n = 5$ for each group). Only the abdominal aorta just distal to the left renal artery was clamped. At 7 days after reperfusion, the groups clamped for 15 minutes were all paraplegic.

For the formal experiments, female domesticated white rabbits, weighing 2.5 to 3.0 kg, were divided into three groups: sham operation control group (group A, with only aortic exposure but without transient ischemia), transient ischemia and treatment with vehicle group (group B), and transient ischemia and treatment with 4-PBA group (15 mg/kg/h, group C). Saline solution (vehicle, 1 mL/kg body weight) or 4-PBA (4 mg/mL in saline vehicle) was administered intravenously from 30 minutes before induction of ischemia until 30 minutes after reperfusion. Total operation time ranged from 1.5 hours to 2.5 hours, and the total amount of 4-PBA administration ranged from 46.9 mg to 56.3 mg. After 15 minutes, the aortic clamp was removed. The abdomen was closed, and all catheters were withdrawn. The animals were allowed to recover from anesthesia and extubated, and routine postoperative care was carried out. Immediately after euthanasia, the spinal cord was quickly removed through a laminectomy at L4 and the bone around the spinal cord was chopped off. All samples were frozen in powdered dry ice and stored at -80°C .

Neurologic and histopathologic studies. The neurologic status in groups A, B, and C ($n = 6$ in each group) was assessed at 8 hours, and 2 and 7 days after reperfusion by an observer blinded to the protocol used for each animal. The motor function of the hind limbs was graded by using the Tarlov score, in which 0 indicates no movement of the hind limbs, 1 indicates perceptible movement of the joints of the hind limbs, 2 indicates good movement of the joints but an inability to stand, 3 indicates the ability to stand and walk, and 4 indicates a complete recovery.¹⁷ The animals ($n = 6$ in each group) were then euthanized after the completion of the neurologic evaluation at 7 days. Sectioned specimens from the lumbar spinal cord at the L4 level were stained with hematoxylin and eosin and were

studied for ischemic pathology by using standard light microscopy. Neuronal ischemic injury was evaluated at a magnification of $\times 200$ by an observer blinded to the treatment groups. Normal motor neurons in the anterior horn of the spinal cord (anterior to a line drawn through the central canal perpendicular to the anterior median fissure) were counted for each animal and averaged. Ischemic neurons were defined by the following criteria: (1) shrinkage of the cell body, (2) an eosinophilic cytoplasm with a loss of Nissl granules, (3) triangular and pyknotic nuclei, and (4) a homogenizing change of the neuron.

Caspase12 and GRP78 immunohistochemistry.

Immunohistochemistry was used to evaluate the changes in expression of caspase12 and GRP78, at 8 hours and 1 day after reperfusion in groups A, B, and C ($n = 3$ for each group at each time point). After endogenous peroxidase activity was quenched by exposing the slides to 0.3% H_2O_2 and 10% methanol for 30 minutes, the spinal cord sections were blocked in 2% normal horse serum for 2 hours at room temperature. Next, they were incubated with primary antibodies for 20 hours at 4°C. The primary antibodies used were as follows: rat monoclonal anti-caspase12 antibody (SC-12747; Santa Cruz Biotechnology Inc, Santa Cruz, Calif) diluted 1:50 and goat polyclonal anti-GRP78 antibody (SC-1050; Santa Cruz Biotechnology) diluted 1:200. The slides were rinsed in PBS and incubated for 3 hours with biotinylated antirat IgG (PK-6104; Vector Laboratories, Burlingame, Calif) and biotinylated antigoat IgG (PK-6105; Vector Laboratories) at 1:200 dilution in PBS containing 0.018% normal horse serum, respectively. Subsequently, they were incubated with avidin-biotin-horseradish peroxidase complex (PK-6104 and PK-6105). The slices were colonized with DAB/ H_2O_2 solution and cytoplasm, and counterstained with hematoxylin. To ascertain specific binding of antibody for protein, a set of sections were stained in a similar way without primary antibody.

Statistical analysis. All results were expressed as means \pm SD. Statistical significance was assumed at a P value of $<.05$. The Mann-Whitney U test, one-way analysis of variance, and post hoc tests with the Fisher protected least significant difference test were used to adequately identify which group differences accounted for the significant P value.

RESULTS

Physiologic status. There were no significant changes in the distal mean arterial pressures and heart rates between the groups during the operation. All 18 animals survived for 7 days without significant intra- or postoperative complications.

Neurologic outcomes. Fig 1 shows the neurologic outcomes of rabbits in each group. The Tarlov scores in groups A, B, and C were 4.0 ± 0.0 , 4.0 ± 0.0 , and 4.0 ± 0.0 at 8 hours; 4.0 ± 0.0 , 2.5 ± 1.2 , and 3.9 ± 0.4 at 2 days; and 4.0 ± 0.0 , 0.7 ± 0.8 , and 4.0 ± 0.0 at 7 days after reperfusion, respectively. While all sham-operated control animals (group A) demonstrated normal neurologic func-




Fig 1. Changes in Tarlov score at 8 hours, and 2 and 7 days after reperfusion. In groups A and C, the Tarlov score was significantly higher at 2 and 7 days than in group B ($*P = .002$, $**P < .0001$). There was no significant difference in neurologic function between groups A and C.

tion, the group B rabbits presented with progressive deterioration of neurologic function, as was anticipated from the results of the preliminary experiments. Five out of six rabbits in group B demonstrated severe paraparesis by 7 days. In group C, neurologic function slightly deteriorated at 2 days but recovered by 7 days. There was no significant difference in Tarlov score between groups A and C across all time points. The score was significantly higher in group C than in group B at 2 and 7 days ($P < .0001$).

Histopathologic assessment. Fig 2 shows the typical histopathologic findings of the spinal cord from a group B rabbit with paraplegia and a group C rabbit with normal neurologic function in comparison with those in group A. In group B (Fig 2, B), the motor neurons in the anterior horn were shrunken or had pale homogenous cytoplasm compared with those in group A, consistent with ischemic change. In group C (Fig 2, C), no apparent histologic changes were detected. The number of motor neurons with a normal appearance in the anterior horn of the spinal cord at 7 days after reperfusion was 47.4 ± 5.7 , 21.5 ± 6.0 , and 44.9 ± 7.9 in groups A, B, and C, respectively (Fig 3). The number of normal neurons was significantly less in group B than in groups A and C ($P < .0001$). There was no significant difference in the number of normal neurons between groups A and C ($P = .019$).

Histochemical study. Immunohistochemistry results for caspase12 and GRP78 are shown in Figs 4 and 5. In the sham-operated animals, there was no evidence of caspase12 and GRP78 expression in any cells of the spinal cord. In group B, caspase12 and GRP78 were present in motor neurons after 8 hours of reperfusion (Fig 4, A and Fig 5, A) but were apparently reduced at 1 day (Fig 4, B and Fig 5, B). In group C, expression of caspase12 and GRP78 in motor neurons was not observed at either 8 hours or 1 day (Fig 4, C and D, and Fig 5, C and D).

DISCUSSION

This study has demonstrated that the peripheral administration of 4-PBA at therapeutic doses prevents both im-




Fig 2. Histologic findings of spinal cord 7 days after 15 minutes of ischemia (hematoxylin and eosin staining). A, Sham-operated control spinal cord was intact. B, In spinal cord of group B, about 60% of motor neurons were lost. C, In group C, no ischemic damage was detected, and there were many normal motor neurons.

mediate and delayed neuronal damage of the spinal cord following an ischemic insult. Neurologic function was preserved with little histologic change observed in motor neurons, most probably due to suppression of ER stress.

When cells are exposed to stress, ER function is disrupted. Because the ER regulates the folding and quality control systems of membrane and secretory proteins, ER stress leads to an accumulation of unfolded proteins in the ER lumen and ER-mediated cell death can be induced.¹⁸ In ER stress, several signaling pathways from the ER such as the unfolded protein response (UPR) are simultaneously activated. Accumulation of unfolded proteins in the ER is sensed by inositol-requiring protein 1 (IRE1), a serine/




Fig 3. The number of viable neurons in the gray matter of lumbar spinal cord sections stained with hematoxylin and eosin. Group C had significantly more viable neurons than group B ($P < .0001$ compared with group B).

threonine kinase with an endoribonuclease domain in the ER membrane, and transcription induction of chaperone genes such as glucose-regulated proteins (GRPs) including GRP78 is promoted to re-fold the unfolded and accumulated proteins.^{19,20} Caspase12, the first identified ER-associated member of the caspase family, is regarded as a representative molecule implicated in the cell death-pathway seen in ER stress.¹¹

ER stress has been reported to contribute to ischemia-reperfusion injury in the brain and spinal cord by inducing apoptosis of neuronal cells.^{8,10,21,22} We speculated that targeting the ER-associated apoptotic pathway might be an effective way to minimize neuronal cell damage, and explored the effect of an ER stress inhibitor, 4-PBA, on minimizing the ischemic injury using a rabbit spinal cord ischemia model.

A low molecular weight fatty acid, 4-PBA, can act as a chemical chaperone, assisting with protein folding and thus relieving the ER stress of the cell.^{16,23-25} In the present study, 4-PBA administration was associated with decreased expression of caspase12 and GRP78 as well as the number of apoptotic cells, suggesting it has beneficial effects on attenuating ER stress. It appears to effectively reduce the neuronal damage, which progresses after reperfusion. Treatment with 4-PBA 30 minutes before and after ischemia notably improved the neurologic outcome. We used the 4-PBA dose of 15 mg/kg/h, based on phase 1 and pharmacokinetic studies of 4-PBA by peripheral infusion.^{26,27} It has low toxicity and good penetrations into the cerebrospinal fluid.²⁸ It has been clinically used for treatment of urea cycle disorders in children, sickle cell disease, thalassemia, and cystic fibrosis.²⁹ Our study has suggested that 4-PBA at therapeutic doses can be a potential agent for preventing spinal cord ischemia. We found that 4-PBA seems to widely suppress ER stress-mediated apoptotic signaling under conditions of ischemia. Considering the findings of the present study, it is reasonable to conclude that the protective effect of 4-PBA on ER stress-mediated injury may involve the activity of chemical chaperone.




Fig 4. Immunohistochemical staining for caspase12 in spinal cords at 8 hours (A) and 1 day (B) in group B, and at 8 hours (C) and 1 day (D) in group C. Density of caspase12 immunoreactivity in group C was reduced relative to that in group B (black arrows).

There are two patterns of neuronal injury due to ischemia: immediate and delayed.³⁰ The former is often caused by a severe ischemic insult and rapidly leads to necrotic neuronal cells. The latter is caused either by apoptosis^{6,31,32} or necrosis.^{30,33,34} Disruption of the infrarenal abdominal aortic blood flow for 15 minutes causes immediate neuronal injury by 2 days and delayed paraplegia by 7 days.^{6,31-34} In the current study, 4-PBA prevented not only immediate neuronal injury but also delayed paraplegia. In the 4-PBA-treated groups, many intact motor neurons were preserved, whereas in the negative control group, the immunohistochemical findings showed expression of caspase12 in motor neurons. To date, 14 caspases have been discovered. They are supposed to be activated in a multiple cascade fashion, activated by caspase9,³⁵ an initiator caspase. Under appropriate stress signals, caspase12 is not only activated within the ER but also translocates to the cytosol, where it can activate caspase9 in a noncytochrome-c-dependent manner.³⁶ Activation of caspase 12 is considered to be one of the potential mechanisms by which the ER contributes to apoptosis. Expression of caspase12 precedes the appearance of neuronal damage and might, therefore, be implicated in the activation of apoptosis.

There are a few points to consider when 4-PBA is applied to clinical situation. First, histologic examination

shows that there is some loss of motor neurons in the 4-PBA-treated groups, indicating that the protective effect of 4-PBA is not perfect. One strategy that may enhance the effectiveness of 4-PBA without increasing its adverse effects is topical injection into the clamped segment of the aorta or combined use of 4-PBA with topical/core cooling. Although preservation of the feeding artery to the spinal cord is essential, such an artery cannot always be identified preoperatively. The protective effect of 4-PBA administered intravenously or the above-mentioned integrated "spinoplegia" protocol may be a potential option in such instances.

Second, the spinal cord anatomy and circulation in rabbits differ significantly from humans. The main source of blood in the rabbit spinal cord is a segmental supply from the abdominal aorta.³⁷ Thus, the lumbar spinal cord of rabbits becomes ischemic by clamping of the infrarenal aorta. Despite the different anatomy, rabbits are often used as the experimental model of spinal cord ischemia because of the simplicity and reproducibility of the model. Some studies have occluded the infrarenal aorta with an intra-aortic balloon.^{6,31,32} Although this procedure is easy and less invasive, it is unclear whether aortic occlusion is complete or not. We chose a direct clamping of the aorta through a retroperitoneal approach to precisely determine




Fig 5. Immunohistochemical staining for GRP78 in spinal cords at 8 hours (A) and 1 day (B) in group B, and at 8 hours (C) and 1 day (D) in group C. Density of GRP78 immunoreactivity in group C was reduced relative to that in group B (black arrows).

the ischemic time of the spinal cord for the reproducible development of paraplegia. On this basis, the current study showed the beneficial effect of 4-PBA.

Third, the drug toxicity of intravenous administration of 4-PBA and the peak time of ER stress at the onset of ischemia are other concerns that will likely limit the clinical application of 4-PBA, although reported side effects are not considerable.²⁶ An additional experiment that examines the timing of injection, injection method and dosage of 4-PBA administration would be required before any clinical applications.

In conclusion, administration of 4-PBA intravenously is feasible and effective without apparent neuronal damage of the spinal cord in a rabbit model. This study indicated that this method can potentially be a useful adjunct for spinal cord protection during thoracic and thoracoabdominal aortic operations.

The authors thank Mr Kazunori Iwase and Ms Emi Fukuda for their excellent technical assistance.

AUTHOR CONTRIBUTIONS

Conception and design: TM, KOrihashi, MH
Analysis and interpretation: TM, KOrihashi
Data collection: TM, BH, ST
Writing the article: TM

Critical revision of the article: KOrihashi, KOkada

Final approval of the article: TS

Statistical analysis: TM, KOrihashi

Obtained funding: Not applicable

Overall responsibility: TM

REFERENCES

1. Coselli JS, Conklin LD, LeMaire SA. Thoracoabdominal aortic aneurysm repair review and update of current strategies. *Ann Thorac Surg* 2002;74:1881-4.
2. Tabayashi K. Spinal cord protection during thoracoabdominal aneurysm repair. *Surg Today* 2005;35:1-6.
3. Cambria RP, Davison JK, Carter C, Brewster DC, Chang Y, Clark KA, et al. Epidural cooling for spinal cord protection during thoracoabdominal aneurysm repair a five-year experience. *J Vasc Surg* 2000;31:1093-102.
4. Matsui Y, Goh K, Shiiya N, Murashita T, Miyama M, Ohba J, et al. Clinical application of evoked spinal cord potentials elicited by direct stimulation of the cord during temporary occlusion of the thoracic aorta. *J Thorac Cardiovasc Surg* 1994;107:1519-27.
5. Moore WN, Hollier LY. The influence of severity of spinal cord ischemia in the etiology of delayed-onset paraplegia. *Ann Surg* 1991;213:427-32.
6. Sakurai M, Nagata T, Abe K, Horinouchi T, Itoyama Y, Tabayashi K, et al. Survival and death promoting events after transient spinal cord ischemia in rabbits: induction of Akt and caspase3 in motor neurons. *J Thorac Cardiovasc Surg* 2003;125:370-7.

7. Budihardjo I, Oliver H, Lutter M, Luo X, Wang X. Biochemical pathways of caspase activation during apoptosis. *Annu Rev Cell Dev Biol* 1999;15:269-90.
8. Cudna RE, Dickson AJ. Endoplasmic reticulum signaling as a determinant of recombinant protein expression. *Biotechnol Bioeng* 2003;81:56-65.
9. Hayashi T, Saito A, Okuno S, Ferrand-Drake M, Chian PH. Induction of GRP78 by ischemic preconditioning reduces endoplasmic reticulum stress and prevents delayed neuronal cell death. *J Cereb Blood Flow Metab* 2003;23:949-61.
10. Oyadomari S, Araki E, Mori M. Endoplasmic reticulum stress-mediated apoptosis in pancreatic beta-cells. *Apoptosis* 2002;7:335-45.
11. Nakagawa T, Zhu H, Morishima N, Li E, Xu J, Yankner BA, et al. Caspase-12 mediates endoplasmic-reticulum-specific apoptosis and cytotoxicity by amyloid-beta. *Nature* 2000;403:98-103.
12. Maestri NE, Brusilow SW, Clissold DB, Bassett SS. Long-term treatment of girls with ornithine transcarbamylase deficiency. *N Engl J Med* 1996;335:855-9.
13. Dover GJ, Brusilow S, Charache S. Induction of fatal hemoglobin production in subjects with sickle cell anemia by oral sodium phenylbutyrate. *Blood* 1994;84:339-43.
14. Burrows JA, Willis LK, Perlmutter DH. Chemical chaperones mediate increased secretion of mutant alpha-1 antitrypsin (α 1-AT)Z: a potential pharmacological strategy for prevention of liver injury and emphysema in α 1-AT deficiency. *Proc Natl Acad Sci USA* 2000;97:1796-801.
15. Rubenstein RC, Zeitlin PL. Sodium 4-phenylbutyrate downregulates Hsc70: implications for intracellular trafficking of DeltaF508-CFTR. *Am J Physiol* 2000;278:259-67.
16. Vilatoba M, Eckstein BA, Bilbao G, Smyth CA, Jenkins S, Thompson JA, et al. Sodium 4-phenylbutyrate protects against liver ischemia reperfusion injury by inhibition of endoplasmic reticulum-stress mediated apoptosis. *Surgery* 2005;138:342-51.
17. Tarlov IM. Spinal cord compression mechanism of paralysis and treatment. Springfield, IL: Charles C. Thomas; 1957. p. 147.
18. Kaufman RJ, Scheuner D, Schroder M, Shen X, Lee K, Liu CY, et al. The unfolded protein response in nutrient sensing and differentiation. *Nat Rev Mol Cell Biol* 2002;3:411-21.
19. Tirasophon W, Welihinda AA, Kaufman RJ. A stress response pathway from the endoplasmic reticulum to the nucleus requires a novel bifunctional protein kinase/endoribonuclease (Ire1p) in mammalian cells. *Genes Dev* 1998;12:1812-24.
20. Wang XZ, Harding HP, Zhang Y, Jolicoeur EM, Kuroda M, Ron D. Cloning of mammalian Ire1 reveals diversity in the ER stress responses. *EMBO J* 1998;17:5708-17.
21. Paschen W, Douthett J. Disturbances of the functioning of endoplasmic reticulum: a key mechanism underlying neuronal cell injury? *J Cereb Blood Flow Metab* 1999;19:1-18.
22. Kumar R, Krause GS, Yoshida H, Mori K, DeGracia DJ. Dysfunction of the unfolded protein response during global brain ischemia and reperfusion. *J Cereb Blood Flow Metab* 2003;23:462-71.
23. Qi X, Hosoi T, Okuma Y, Kaneko M, Nomura Y. Sodium 4-phenylbutyrate protects against cerebral ischemic injury. *Molecular Pharmacol* 2004;66:899-908.
24. Ryu H, Smith K, Camelo SI, Carreras I, Lee J, Iglesias AH, et al. Sodium phenylbutyrate prolongs survival and regulates expression of anti-apoptotic genes in transgenic amyotrophic lateral sclerosis mice. *J Neurochem* 2005;93:1087-98.
25. Kubota K, Niinuma Y, Kaneko M, Okuma Y, Sugai M, Omura T, et al. Suppressive effects of 4-phenylbutyrate on the aggregation of Pael receptors and endoplasmic reticulum stress. *J Neurochem* 2006;97:1259-68.
26. Carducci MA, Gilbert J, Bowling MK, Noc D, Eisenberger MA, Sini-baldi V, et al. A phase I clinical and pharmacological evaluation of sodium phenylbutyrate on a 120-h infusion schedule. *Clin Cancer Res* 2001;7:3047-55.
27. Camacho LH, Olson J, Tong WP, Young CW, Spriggs DR, Malkin MG. Phase I dose escalation clinical trial of phenylbutyrate sodium administered twice daily to patients with advanced solid tumors. *Invest New Drugs* 2007;2:131-8.
28. Berg S, Serabe B, Aleksic A, Bomgaars L, McGuffey L, Dauser R, et al. Pharmacokinetics and cerebrospinal fluid penetration of phenylacetate and phenylbutyrate in the nonhuman primate. *Cancer Chemother Pharmacol* 2001;47:385-90.
29. Perlmutter DH. Chemical chaperones: a pharmacological strategy for disorders of protein folding and trafficking. *Pediatr Res* 2002;52:832-6.
30. Papakostas JC, Matsagas MI, Toumpoulis IK, Malamou-Mitsi VD, Pappa LS, Gkrepi C, et al. Evolution of spinal cord injury in a porcine model of prolonged aortic occlusion. *J Surg Res* 2006;133:159-66.
31. Sakurai M, Nagata T, Abe K, Horinouchi T, Itoyama Y, Tabayashi K. Oxidative damage and reduction of redox factor-1 expression after transient spinal cord ischemia in rabbits. *J Vasc Surg* 2003;37:446-52.
32. Sakurai M, Takahashi G, Abe K, Horinouchi T, Itoyama Y, Tabayashi K. Endoplasmic reticulum stress induced in motor neurons by transient spinal cord ischemia in rabbits. *J Thorac Cardiovasc Surg* 2005;130:640-5.
33. Kiyoshima T, Fukuda S, Matsumoto M, Iida Y, Oka S, Nakakimura K, et al. Lack of evidence for apoptosis as a cause of delayed onset paraplegia after spinal cord ischemia in rabbits. *Anesth Analg* 2003;96:839-46.
34. Lee JC, Hwang IK, Park SK, Yoo KY, Seo K, Kang TC, et al. Histochemical and electron microscopic study on motor neuron degeneration following transient spinal cord ischaemia at normothermic conditions in rabbits. *Anat Histol Embryol* 2005;34:252-7.
35. Slee EA, Harte MT, Kluck RM, Wolf BB, Casiano CA, Newmeyer DD, et al. Ordering the cytochrome c-initiated caspase cascade: hierarchical activation of caspases-2, -3, -6, -7, -8, and -10 in a caspase-9-dependent manner. *J Cell Biol* 1999;144:281-92.
36. Rao RV, Hermel E, Castro-Obregon S, del Rio G, Ellerby LM, Ellerby IIM, et al. Coupling endoplasmic reticulum stress to the cell death program. Mechanism of caspase activation. *J Biol Chem* 2001;276:33869-74.
37. Zivin JA, DeGirolami U. Spinal cord infarction: a highly reproducible stroke model. *Stroke* 1980;11:200-2.

Submitted Mar 23, 2010; accepted Jun 27, 2010.

Cold blood spinoplegia under motor-evoked potential monitoring during thoracic aortic surgery

Shinya Takahashi, MD, Kazumasa Orihashi, MD, Katsuhiko Imai, MD, Taketomo Mizukami, MD, Taiichi Takasaki, MD, and Taijiro Sueda, MD

Objective: Motor-evoked potential monitoring is used to prevent paraplegia during thoracic aortic surgery. Multidetector computed tomography has been used preoperatively to detect the Adamkiewicz artery, but the hemodynamic significance of the Adamkiewicz artery is controversial. This study aims to evaluate whether the multidetector computed tomography–defined Adamkiewicz artery is hemodynamically essential and needs to be reconstructed with cold blood spinoplegia under motor-evoked potential monitoring.

Methods: From 2005 to 2008, both preoperative multidetector computed tomographic analysis and intraoperative neurogenic motor-evoked potential monitoring with cold blood infusion into the clamped segment of the aorta were done in 15 patients. A motor-evoked potential decrease to less than 50% of the initial value at 3 minutes after cold blood infusion determined the hemodynamic significance of the multidetector computed tomography–defined Adamkiewicz artery. Adamkiewicz arteries determined to be essential were reconstructed, and those determined to be nonessential were sacrificed.

Results: The Adamkiewicz artery was involved in the clamped segment of the aorta in 11 cases. After cold blood infusion, 8 patients experienced no significant motor-evoked potential decrease, and Adamkiewicz artery ligation was undertaken, whereas a moderate motor-evoked potential decrease was noted in 1 patient, prompting reconstruction. None of these 9 patients had permanent neurologic deficits. In 2 patients, the Adamkiewicz artery was reconstructed based on motor-evoked potential findings, with paraparesis occurring in 1 patient. In 4 patients without Adamkiewicz artery involvement in the clamped segment, there was no neurologic deficit.

Conclusions: Cold blood infusion accelerates motor-evoked potential changes and might enable decision making regarding the need for reconstruction of multidetector computed tomography–defined Adamkiewicz arteries. Cold blood–loaded motor-evoked potential is beneficial to minimize Adamkiewicz artery reconstruction time and limit spinal cord ischemia. (*J Thorac Cardiovasc Surg* 2011;141:755-61)

Paraplegia, paraparesis, or both remain serious complications of thoracic and thoracoabdominal aortic aneurysm repair and occur in 2% to 16% of patients¹⁻⁵ despite the implementation of various adjunctive procedures, including distal perfusion, mild hypothermia, cerebrospinal fluid drainage, and protective drugs. Two strategies that are commonly accepted for preserving essential spinal cord perfusion are (1) preoperative identification of the Adamkiewicz artery (AKA) by means of 64-row multidetector computed tomographic angiographic (MDCTA) or magnetic resonance angiographic analysis and (2) intraoperative monitoring of the motor-evoked potential (MEP).

The former modalities enable visualization of the continuity of the intercostal or lumbar arteries, the greater radicular

artery, and the anterior spinal artery in more than 95% of patients.⁶⁻⁸ Despite preoperatively identified AKAs being elaborately reconstructed in many institutions to minimize the risk of spinal cord injury, they are often found to be occluded on postoperative assessment, suggesting that they were not necessarily essential.

MEP monitoring has proved to be a sensitive monitor of spinal cord ischemia and is beneficial to improving surgical outcomes.^{9,10} However, it can take longer than 10 minutes to determine ischemic changes of MEP in normothermia. For this reason, we introduced cold blood infusion into the clamped segment of the aorta, aiming to prompt MEP changes by topically cooling the spinal cord.¹¹ The MEP amplitude significantly decreased within 3 minutes after cold blood infusion, whereas no patient showed delayed onset of MEP decrease. Cold blood–loaded MEP appeared to clearly delineate the presence or absence of MEP changes within 3 minutes.

Recently, Griep and Griep¹² postulated the “collateral network theory” and reported excellent results. The intercostal and lumbar arteries between T7 and L1 were sacrificed while the MEP and somatosensory-evoked potential (SSEP) were monitored during thoracic and thoracoabdominal repair.¹³ These reports make us reconsider the significance of the AKA.

From the Department of Cardiovascular Surgery, Hiroshima University Hospital, Hiroshima, Japan.

Disclosures: Authors have nothing to disclose with regard to commercial support. Received for publication May 14, 2010; revisions received Aug 18, 2010; accepted for publication Sept 12, 2010; available ahead of print Nov 22, 2010.

Address for reprints: Shinya Takahashi, MD, 1-2-3, Kasumi, Minamiku, Hiroshima, Japan 732-0062 (E-mail: shiny@nte.biglobe.ne.jp).

0022-5223/536.00

Copyright © 2011 by The American Association for Thoracic Surgery

doi:10.1016/j.jtcvs.2010.09.056

Abbreviations and Acronyms

AKA	= Adamkiewicz artery
CSF	= cerebrospinal fluid
CT	= computed tomography
MDCTA	= multidetector computed tomographic angiography
MEP	= motor-evoked potential
SSEP	= somatosensory-evoked potential
TEE	= transesophageal echocardiography
TEVAR	= thoracic endovascular aneurysm repair

Thus we hypothesize that (1) a prompt MEP decrease indicates hemodynamic significance of the AKA and need for reconstruction and (2) hemodynamic insignificance is responsible for an occlusion of the reconstructed AKA. The purpose of this article is to report the preliminary results of this study.

MATERIALS AND METHODS**Subjects**

From 2005 to 2008, 29 patients underwent aortic repair for descending thoracic or thoracoabdominal aortic aneurysms in our institute, excluding emergency cases. In 15 of these patients, preoperative assessment of the AKA was done with MDCTA, and MEP was monitored intraoperatively. In the remaining 14 patients, MEP monitoring was not used because the aneurysm was located proximal to the T6 level or far distal to the AKA visualized with MDCTA, or cold blood infusion was not performed because of aortic dissection with a large false lumen, and these patients were excluded from the study. Demographics of the patients are shown in Table 1. There were 11 male and 4 female patients with ages ranging from 59 to 86 years (mean age, 72.7 ± 7.6 years). All patients had several comorbidities, including hypertension. Seven patients had descending thoracic aortic aneurysms, and 7 patients had Crawford type I and 1 patient had Crawford type IV thoracoabdominal aortic aneurysms.

Preoperative MDCTA Study

Helical computed tomographic (CT) angiographic analysis was performed with a 64-row multidetector CT scanner (Light Speed VCT; GE Healthcare, Waukesha, Wis). After 100 mL of contrast medium was administered intravenously, scanning was started when the CT value in the lumen of the thoracic aorta (T7) reached 200 Hounsfield units. Three-dimensional image reconstruction was performed with analysis software (Virtual Place Rajjin; AZE, Ltd, Tokyo, Japan). The AKA was defined as a continuous vessel from the aorta through the intercostal or lumbar artery and the radiculomedullary artery to the anterior spinal cord artery to eliminate venous signatures from being mistaken as arterial.

MEP Monitoring

Neurogenic MEP monitoring was used in every case to obtain discrete waveforms during adequate muscle relaxation. On the day before the operation, a pair of bipolar, platinum electrode catheters (UKG-100-5PM; Unique Medical, Tokyo, Japan) were placed in the dorsal epidural space for intraoperative evoked spinal cord potential monitoring: a stimulating electrode was placed at the C5-C6 level, and a recording electrode was placed at the lumbar level. A cerebrospinal fluid (CSF) drainage catheter was inserted at a low lumbar location (L3-L4).

After induction of anesthesia with fentanyl/remifentanyl and propofol with vecuronium bromide, a pair of bipolar, screw-type platinum electrodes was placed on the bilateral anterior parietal region for transcranial electrical stimulation of the cerebral motor cortices. A pair of bipolar surface electrodes was placed on the bilateral peroneal nerves for the SSEP. The MEPs and SSEPs were recorded at the lumbar electrodes. Stimulation and recordings were performed with the EpochXp system (Nihon Koden, Tokyo, Japan; Figure 1). The stimuli (MEP: intensity, 100 mA; duration, 0.2–1.0 ms, single pulse; SSEP: intensity, 25–50 mA; duration, 0.2 ms, single pulse) were delivered at a rate of 3.3 to 7.3 Hz. Each evoked potential was filtered between 20 and 1500 Hz and was amplified by averaging a total of 20 to 50 responses. All data were stored in the hard disc of this system.

Surgical Procedures

The aorta was exposed through a left thoracotomy with or without a paramedian retroperitoneal approach. Femorofemoral partial cardiopulmonary bypass was established with a pump flow of 2.2 to $2.6 \text{ L} \cdot \text{min}^{-1} \cdot \text{m}^{-2}$ after intravenous injection of steroids and the oxygen radical scavenger edaravone. The rectal temperature was maintained at 34°C . CSF drainage was continued for 2 days to maintain the CSF pressure at less than 10 mm Hg. The distal perfusion pressure was maintained at a level higher than 60 mm Hg. The intercostal arteries, including the preoperatively identified AKA, were exposed.

After aortic crossclamping, cold blood (4°C) was infused into the clamped segment of the aorta through a 14-gauge needle at a flow rate of 100 to 200 mL/min for 3 minutes (Figure 2) while MEP amplitudes were monitored. The puncture site was carefully determined based on meticulous assessment by means of transesophageal echocardiographic (TEE) analysis, and cold blood inflow was visualized with TEE analysis to avoid unintended detachment of a mural thrombus. The amount of infused cold blood depended on the size of the aneurysm and ranged from 300 to 600 mL.

Decision Making Based on MEP Monitoring

The AKA in the clamped segment was diagnosed as essential when the MEP amplitude decreased to a level less than 50% of the original amplitude after cold blood infusion within 3 minutes (Figure 3). In patients with a critical decrease in MEP, the AKA was immediately reconstructed after aortotomy with a vascular prosthesis (10 mm in diameter) or a saphenous vein and was perfused with warm blood at a flow rate of 40 mL/min and a perfusion pressure of up to 150 mm Hg. After the aorta was repaired, this reconstructed intercostal graft was anastomosed to the main graft. When there was no decrease of MEP amplitude after cold blood infusion, the AKA identified by means of preoperative MDCTA study was determined to be not essential and was ligated extraluminally before opening the aneurysm. If the other intercostal arteries were exposed, they were also ligated extraluminally before opening the aneurysm; otherwise, they were ligated intraluminally.

RESULTS

A total of 17 AKAs were identified in 15 patients undergoing preoperative MDCTA analysis: a single AKA in 13 patients and 2 AKAs in 2 patients. The AKA entered the spinal canal at T7 (2 AKAs), T9 (3 AKAs), T10 (5 AKAs), T11 (3 AKAs), T12 (2 AKAs), or L1 (2 AKAs). Fifteen AKAs were on the left side, and 2 AKAs were on the right side. No complications related to the placement of epidural electrodes for MEP monitoring were encountered in any case.

Duration of the operation, duration of extracorporeal circulation, and aortic clamp time were 418 ± 182 , 92 ± 40 ,

TABLE 1. Patients' characteristics

No. of patients	15
Age, y (range)	72.7 ± 7.6 (59 - 86)
Sex (male/female)	11/4
Preoperative variables	
Hypertension	13 (87%)
Hyperlipidemia	4 (27%)
Diabetes mellitus	3 (20%)
COPD	1 (7%)
Coronary artery disease	4 (27%)
Renal insufficiency	2 (13%)
Cerebral infarction	1 (7%)
Smoking	4 (27%)
Marfan syndrome	0 (0%)
Coactation of aorta	0 (0%)
Previous chest pain	3 (20%)
Aortic dissection	4 (27%)
Traumatic	0 (0%)
Aortic infection	0 (0%)
Aortic rupture	0 (0%)

COPD, Chronic obstructive pulmonary disease.

and 77 ± 31 minutes, respectively. There was no paraplegia, but 1 patient experienced paraparesis caused by cerebral infarction, probably secondary to multiple thromboembolism during the proximal aortic clamping. There were no operative or hospital deaths.

Table 2 shows the level of the AKAs, involvement of the AKA in the aneurysm, changes in MEP amplitudes, and surgical outcomes. Although the AKA was not involved in the clamped segment of the aorta in 4 patients (patients 1-4),

the origin of the AKA was within the clamped segment in 11 patients (patients 5-15). In the former 4 patients, there were no MEP changes after cold blood infusion. No neurologic deficit was encountered in any of these patients. Among the latter 11 patients, there was no significant MEP decrease within 3 or 20 minutes in 7 patients (patients 5-11). Although every intercostal artery in the clamped segment, including the supposed AKA, was ligated, none of them presented with neurologic deficit. The MEP amplitude decreased but remained at a level greater than 50% of the initial value in 2 patients (patients 12 and 13). Delayed MEP decrease within 20 minutes was not appreciated in either of these patients. The AKA was ligated in both patients, but it was finally reconstructed in 1 patient because of the reason described below. The MEP amplitude decreased to a level less than 50% of the initial value in only 2 patients (patients 14 and 15), both of whom underwent reconstruction of the AKAs.

Although patient 12 presented with a gradual decrease of MEP to 70% of the initial value during the operation, motor function of the lower limb was temporarily restricted after the operation and was recovered during the hospital stay. Patient 13, who had previous repair of the proximal thoracic aorta for acute aortic dissection, underwent replacement of the distal descending aorta. The MEP amplitude decreased to 70% at 3 minutes of cold blood infusion and gradually recovered to the initial level. However, the MEP amplitude decreased to 70% again during a period of hypotension caused by sustained bleeding and did not recover to the




FIGURE 1. Evoked potential (EP) monitoring system and conduction pathway of evoked potential (dotted line). Transcranial electrical stimulation passed through motor cortices and the spinal cord was recorded by using the epidural electrode (L1-L2; transcranial motor-evoked potential). Descending evoked spinal cord potentials (C5-C6 to L1-L2) and somatosensory-evoked potentials (peroneal nerve to L1-L2) were also recorded for multilateral observation.




FIGURE 2. Procedure of cold blood infusion. A, A 14-gauge needle was punctured into the aorta. B, Schema of the picture. Cold blood was infused into the clamped segment of the aorta.

initial level. In this case, we initially ligated the AKA but decided to reconstruct the AKA at T10 to ensure adequate blood flow because the spinal cord perfusion appeared to be partially dependent on the AKA. This patient did not have a neurologic deficit. Postoperative CT scanning showed occlusion of the reconstructed intercostal artery.

In patients 14 and 15, the intercostal arteries were reconstructed based on the significant and immediate MEP changes. In patient 14, the MEP recovered after reperfusion, and no neurologic deficit developed. In patient 15, however, MEP disappeared at one point and later recovered, although only to 50% of the initial level. Aortic perfusion pressure was maintained at greater than 100 mm Hg during the operation and during the postoperative period. The patient in this case had paraparesis postoperatively. In this patient, the aorta was diffusely and severely atherosclerotic, and fragile intima was likely to have caused left cerebral infarction from a multiple thromboembolism caused by proximal aortic clamping. Furthermore, selective perfusion of the intercostal artery was not feasible because of atherosclerotic stenosis of the AKA, and ischemic time was prolonged.

Postoperatively CSF drainage pressure increased to greater than 15 mm Hg, and CSF drainage was continued. The cranial magnetic resonance image showed multiple left cerebral infarctions. The left upper limb function was normal, but the left lower limb movement was impaired, which was unexplainable based on the cranial magnetic resonance image. Therefore we concluded that the patient had lower limb paraparesis with right hemiplegia caused by both stroke and spinal cord ischemia.

DISCUSSION

Despite the fact that thoracic endovascular aneurysm repair (TEVAR) has achieved a lower incidence of paraplegia and paraparesis than open surgical intervention, neurologic complications have not been completely eliminated, suggesting a possibility of occluding the essential AKA by the endovascular graft.¹⁴ Siegenthaler and associates¹⁵ reported delayed temporary paraparesis that appeared after successful TEVAR, but the deficit was corrected by increasing the blood pressure. These findings indicate that spinal cord perfusion depends not solely on AKAs but partially




FIGURE 3. Changes in motor-evoked potential (MEP). A, MEP amplitude did not change. B, MEP amplitude decreased after cold blood infusion within 3 minutes and was restored after Adamkiewicz artery (AKA) reconstruction and declamping of the aorta.

PM

TABLE 2. MDCTA-defined AKA and results

Patients no.	Type of aneurysm	AKA (MDCTA)	AKA in aneurysm	MEP at 3 min	Amplitude (%)		Reconstruction of branch artery	Neurologic outcome
					Lowest	At end		
1	DTAA	T9L, T12L	N	100%	100%	100%	None	No deficit
2	DTAA	T11L	N	100%	100%	100%	None	No deficit
3	I	T12L	N	100%	100%	100%	None	No deficit
4	IV	T11L	N	100%	100%	100%	None	No deficit
5	DTAA	T7L	Y	100%	100%	100%	None	No deficit
6	DTAA	T7L	Y	100%	100%	100%	None	No deficit
7	I	T9R	Y	100%	100%	100%	None	No deficit
8	DTAA	T10R	Y	100%	100%	100%	None	No deficit
9	I	T11L	Y	100%	100%	100%	None	No deficit
10	I	T9L	Y	100%	80%	100%	None	No deficit
11	DTAA	T10L	Y	100%	70%	100%	None	No deficit
12	I	L1L	Y	80%	70%	70%	None	No deficit
13	I	T10L, L1L	Y	70%	70%	80%	T10L	No deficit
14	DTAA	T10L	Y	45%	45%	90%	T10L	No deficit
15	I	T10L	Y	40%	0%	50%	T10L	Paraparesis

The patients are in order in relation to motor-evoked potential change and not in chronologic order. Reconstructed branch arteries were patent in patients 13 and 14 and occluded in patient 15. MDCTA, Multidetector computed tomographic angiography; AKA, Adamkiewicz artery; MEP, motor-evoked potential; DTAA, descending thoracic aortic aneurysm.

on collateral circulation, although the latter can be inadequate under low perfusion pressure.

The role of collateral circulation has been emphasized in the collateral network concept of Griep and Griep.¹² Etz and colleague¹³ reported that sacrifice of the intercostal and lumbar arteries during thoracic and thoracoabdominal aortic aneurysm repair resulted in low rates of immediate and delayed paraplegia. They described a patient with immediate paraplegia who experienced precipitating intraoperative dissection resulting in 6 hours of lower body ischemia. This case indicates the importance of collateral circulation.

Two questions arise. First, should every AKA be elaborately reconstructed at a risk of prolonged ischemia of the spinal cord? Second, is it the best strategy to create a situation in which spinal cord perfusion depends only on collateral circulation? A possible solution is to reconstruct only the essential AKAs and sacrifice the other nonessential AKAs to minimize the ischemic insult on the spinal cord.

Improved quality of 64-row MDCTA and magnetic resonance angiography has enabled visualization of AKAs in more than 95% of cases.⁶⁻⁸ Three-dimensional CT images discretely visualize the continuity from the intercostal artery to the anterior spinal cord artery and facilitate identification of the AKA in the operative field. This development has led to a situation that AKAs need to be reconstructed in most cases, and the majority of grafts are found to be occluded postoperatively.¹⁶ It is unclear whether the occluded graft was effective in avoiding spinal cord ischemia in the perioperative period or whether reconstruction was even necessary. A considerably lower incidence of paraplegia after TEVAR, despite occlusion of every branch artery, appears to support the latter. In another respects, however, TEVAR might create a new Achilles heel: persistent susceptibility to hypotension, even if the patient can survive the

neurologic deficit in the acute phase. Thus it is desirable to identify and reconstruct only the essential AKAs among the morphologically defined AKAs.

We previously reported that cold blood infusion into the clamped segment of the aorta induces a rapid change in MEP amplitude.¹¹ A decrease in MEP amplitude might be related to local hypothermia by cold blood.⁵ Ueno and co-workers¹⁷ reported that intra-aortic infusion of cold lactated Ringer's solution (3°C) cooled the spinal cord by 6°C to 7°C. Because hypothermia induces a decreased electrophysiologic response in cerebral and peripheral neurologic activity,^{18,19} an immediate MEP decrease is likely to be caused by cooling of the spinal cord. Furthermore, because cold blood saturated with oxygen is infused, infusion of cold blood does not induce spinal cord ischemia and just accelerates MEP changes. We hypothesized that cold blood infused into the essential AKA topically cools the spinal cord and decreases MEP amplitude. We did not perform a morphologic assessment but rather a functional assessment.

In this study, as well as a previous study, the MEP was recorded at the intrathecal electrode (neurogenic MEP).^{11,20} The advantage of a neurogenic MEP is a simple waveform compared with a compound waveform recorded in a myogenic MEP.^{13,21} However, it takes a longer time to detect ischemic changes in a neurogenic MEP than in a myogenic MEP. de Haan and associates²² reported that the median time to a 50% reduction in epidural and myogenic MEP amplitudes after aortic occlusion was 11.3 minutes and within 2 minutes, respectively. We previously showed that an MEP decrease was caused by cooling of the spinal cord in a dog model.²³ Despite the longer time to detect changes in neurogenic MEP during ischemia, cold blood infusion might be helpful for identifying

hemodynamically significant AKAs based on an early change in the neurogenic MEP.

We routinely monitor both afferent and efferent conduction to check for damage of the spinal cord or in case the MEP recording is unexpectedly interrupted for some unknown reason. SSEPs were used to distinguish between spinal cord ischemia and systemic factors or technical problems. The monitoring of both MEPs and SSEPs is considered to be a safety system to prevent spinal cord ischemia.

Our current strategy for descending aortic repair is described as follows:

- (1) If there is no AKA in the clamped segment, the branched arteries are not preserved.
- (2) If the AKA is present in the clamped segment, MEP changes after cold blood infusion into the clamped segment of aorta are monitored.
 - (a) Without a significant MEP change, the AKA is not reconstructed.
 - (b) When MEP amplitude decreases to a level less than 50% of the initial value, the AKA is determined to be essential and is reconstructed.

The cutoff value of MEP amplitude as 50% of the baseline value is based on the report of van Dongen and coworkers¹⁰ that the relative risks of paraplegia in patients with an MEP amplitude of 50% or less than the baseline value after 5 minutes of reperfusion and during closure of the skin were 21.7 and 30.9 times higher, respectively, compared with those of patients with an MEP of greater than 50% of the baseline value. We evaluated the MEP changes at 3 minutes after cold blood infusion based on the findings in our previous studies.^{11,20} The MEP amplitude did not decrease to a level less than 50% of the baseline value after 3 minutes of cold blood infusion in any patient unless there was an apparent cause, such as considerable systemic hypotension. If our hypothesis was wrong, neurologic deficit should have occurred in several patients in whom the AKA in the clamped segment of the aorta was sacrificed (patients 5–12). In this series, MEP amplitude was unchanged after cold blood infusion in 7 of 11 patients, and this might explain the high incidence of graft occlusion of reconstructed AKAs.

It remains uncertain whether the AKA should be revascularized when the decrease in MEP amplitude is mild.²⁰ Kawanishi and colleagues²⁴ reported that 8 of 9 patients whose MEP amplitude decreased to less than 75% of baseline value at the end of the procedure had a neurologic deficit postoperatively. Because our study had a small number of patients, it is difficult to make a definitive conclusion, especially when the MEP is between 50% and 99% of the baseline value. In 2 patients, the MEP amplitude decreased to 70% to 80% of the baseline value. Although

1 patient had no paraplegia with the immediate decision to reconstruct the AKA, postoperative CT scanning revealed occlusion of the reconstructed intercostal artery, and this patient often noticed discomfort and slight palsy of the lower limbs during periods of hypotension associated with daily hemodialysis. The other patient had a transient paraparesis, probably as a result of sustained hypotension, which was caused by a large amount of oozing from the broadly dissected surface in this redo case. The events in these patients appear to be comparable with those in the patient undergoing successful TEVAR with delayed paraplegia caused by hypotension. These results suggest that the AKA should be reconstructed if there is any decrease in MEP amplitude. Hypotension during the operation and the postoperative period should be avoided. This will be our new strategy in the future based on the results of the current study.

Neurologic deficit in surgical cases is likely to be caused by (1) occlusion of the essential AKA, (2) intraoperative hypotension that leads to hypoperfusion of the spinal cord for a prolonged period of time, or both. Acher and associates²⁵ reported desirable outcomes of descending aortic surgery by maintaining the blood pressure at greater than 90 mm Hg and mentioned that the mean arterial pressure during aortic clamping was a significant factor for paralysis. By using a multidisciplinary strategy in aortic surgery, the incidence of neurologic deficit can be decreased to a level comparable with that of TEVAR. Reconstruction of all AKAs detected in the preoperative assessment can adversely prolong the duration of cardiopulmonary bypass and increase the amount of bleeding. It might be beneficial to limit the number of reconstructions by identifying the essential functional AKA by means of MEP monitoring under cold blood infusion. If the essential AKA can be successfully reconstructed without neurologic deficit, it can be beneficial in reducing the future risk against unpredicted hypotension in various situations.

In 4 patients (patients 1–4), the AKA was located close to the aneurysm on the preoperative CT scan, and there was a risk of including this artery in the clamped segment. The level of the aortic clamp might be altered when the anastomosis is difficult. In such a situation, the nearby AKA can be included in the clamped segment. Thus we have used our method in these patients and included it in this study.

The small number of patients in the current series is a limitation of this study. Nevertheless, our preliminary results demonstrate that the majority of MDCTA-defined AKAs can be sacrificed without neurologic sequelae. The need for puncture and cold blood infusion into the aneurysm can be another limitation because it might raise a concern of possible thromboembolism. However, careful TEE assessment for an appropriate puncture site and subsequent observation during infusion has shown that this procedure can be safely performed.

In conclusion, cold blood-loaded MEP monitoring might enable identification of the hemodynamically essential AKAs among the commonly visualized AKAs with preoperative MDCTA analysis. Although the nonessential AKAs thus defined by our criteria could be sacrificed without significant neurologic complications, we need further investigation in a case of an MEP decreasing to a level greater than 50%. This method might be useful for efficiently repairing the aorta with AKA reconstruction to mitigate against the effects of further unpredicted hypotension.

References

- Griep RB, Ergin MA, Galla JD, Lansman S, Khan N, Quintana C, et al. Looking for the artery of Adamkiewicz: a quest to minimize paraplegia after operations for aneurysms of the descending thoracic and thoracoabdominal aorta. *J Thorac Cardiovasc Surg.* 1996;112:1202-15.
- Svensson LG, Crawford ES, Hess KR, Coselli JS, Safi HJ. Experience with 1509 patients undergoing thoracoabdominal aortic operations. *J Vasc Surg.* 1993;17:357-70.
- Grabitz K, Sandmann W, Stuhmeier K, Mainzer B, Godehardt E, Ohle B, et al. The risk of ischemic spinal cord injury in patients undergoing graft replacement for thoracoabdominal aortic aneurysms. *J Vasc Surg.* 1996;23:230-40.
- Safi HJ, Campbell MP, Miller CC 3rd, Iliopoulos DC, Khojenezhad A, Letsou GV, et al. Cerebral spinal fluid drainage and distal aortic perfusion decrease the incidence of neurological deficit: the results of 343 descending and thoracoabdominal aortic aneurysm repairs. *Eur J Vasc Endovasc Surg.* 1997;14:118-24.
- Coselli JS. The use of left heart bypass in the repair of thoracoabdominal aortic aneurysms: current techniques and results. *Semin Thorac Cardiovasc Surg.* 2003;15:326-32.
- Nakayama Y, Awai K, Yanaga Y, Nakaura T, Funama Y, Hirai T, et al. Optimal contrast medium injection protocols for the depiction of the Adamkiewicz artery using 64-detector CT angiography. *Clin Radiol.* 2008;63:880-7.
- Boll DT, Bulow H, Blackham KA, Aschoff AJ, Schmitz BL. MDCT angiography of the spinal cord vasculature and the artery of Adamkiewicz. *Am J Roentgenol.* 2006;187:1054-60.
- Nijenhuis RJ, Jacobs MJ, Schurink GW, Kessels AG, van Engelsehoven JM, Backes WH. Magnetic resonance angiography and neuromonitoring to assess spinal cord blood supply in thoracic and thoracoabdominal aortic aneurysm surgery. *J Vasc Surg.* 2007;45:71-7.
- Jacobs MJ, Mess WH. The role of evoked potential monitoring in operative management of type I and type II thoracoabdominal aortic aneurysms. *Semin Thorac Cardiovasc Surg.* 2003;15:353-64.
- van Dongen EP, Schepens MA, Morshuis WJ, ter Beek HT, Aarts LP, de Boer A, et al. Thoracic and thoracoabdominal aortic aneurysm repair: use of evoked potential monitoring in 118 patients. *J Vasc Surg.* 2001;34:1035-40.
- Sueda T, Okada K, Orihashi K, Sugawara Y, Kouchi K, Imai K. Cold blood spinal cord plegia for prediction of spinal cord ischemia during thoracoabdominal aneurysm repair. *Ann Thorac Surg.* 2002;73:1155-9.
- Griep RB, Griep EB. Spinal cord perfusion and protection during descending thoracic and thoracoabdominal aortic surgery: the collateral network concept. *Ann Thorac Surg.* 2007;83:S865-92.
- Etz CD, Halstead JC, Spielvogel D, Shahani R, Lazala R, Homann TM, et al. Thoracic and thoracoabdominal aneurysm repair: is reimplantation of spinal cord arteries a waste of time? *Ann Thorac Surg.* 2006;82:1670-7.
- Kotelis D, Geisbusch P, von Tengg-Kobligk H, Allenberg JR, Bockler D. [Paraplegia after endovascular repair of the thoracic and thoracoabdominal aorta]. *Zentralbl Chir.* 2008;133:338-43.
- Siegenthaler MP, Weigang E, Brehm K, Euringer W, Baumann T, Uhl M, et al. Endovascular treatment for thoracoabdominal aneurysms: outcomes and results. *Eur J Cardiothorac Surg.* 2008;34:810-9.
- Svensson LG. Intraoperative identification of spinal cord blood supply during repairs of descending aorta and thoracoabdominal aorta. *J Thorac Cardiovasc Surg.* 1996;112:1455-61.
- Ueno T, Furukawa K, Katayama Y, Itoh T. Protection against ischemic spinal cord injury: one-shot perfusion cooling and percutaneous topical cooling. *J Vasc Surg.* 1994;19:882-7.
- Stecker MM, Cheung AT, Pochettino A, Kent GP, Patterson T, Weiss SJ, et al. Deep hypothermic circulatory arrest: I. Effects of cooling on electroencephalogram and evoked potentials. *Ann Thorac Surg.* 2001;71:14-21.
- Stecker MM, Cheung AT, Pochettino A, Kent GP, Patterson T, Weiss SJ, et al. Deep hypothermic circulatory arrest: II. Changes in electroencephalogram and evoked potentials during rewarming. *Ann Thorac Surg.* 2001;71:22-8.
- Hamaishi M, Orihashi K, Takahashi S, Isaka M, Okada K, Sueda T. Transcranial motor-evoked potentials following intra-aortic cold blood infusion facilitates detection of critical supplying artery of spinal cord. *Eur J Cardiothorac Surg.* 2008;33:695-9.
- de Haan P, Kalkman CJ, de Mol BA, Ubags LH, Veldman DJ, Jacobs MJ. Efficacy of transcranial motor-evoked myogenic potentials to detect spinal cord ischemia during operations for thoracoabdominal aneurysms. *J Thorac Cardiovasc Surg.* 1997;113:87-101.
- de Haan P, Kalkman CJ, Ubags LH, Jacobs MJ, Drummond JC. A comparison of the sensitivity of epidural and myogenic transcranial motor-evoked responses in the detection of acute spinal cord ischemia in the rabbit. *Anesth Analg.* 1996;83:1022-7.
- Kumagai H, Sugawara Y, Isaka M, Okada K, Orihashi K, Sueda T. Cold saline injection attenuates motor-evoked potential in the spinal cord by cortical electrical stimulation in the dog. *Hiroshima J Med Sci.* 2005;54:77-82.
- Kawanishi Y, Munakata H, Matsumori M, Tanaka H, Yamashita T, Nakagiri K, et al. Usefulness of transcranial motor evoked potentials during thoracoabdominal aortic surgery. *Ann Thorac Surg.* 2007;83:456-61.
- Acher CW, Wynn MM, Mell MW, Tefera G, Hoch JR. A quantitative assessment of the impact of intercostal artery reimplantation on paralysis risk in thoracoabdominal aortic aneurysm repair. *Ann Surg.* 2008;248:529-40.

腹部大動脈瘤に対する自作ステントグラフト治療後の migration に対して企業デバイスを使用した 1 例

坂下 英樹 田中 厚寿 鬼塚 誠二 廣松 伸一 明石 英俊 青柳 成明

要 旨: 症例は 74 歳女性. 71 歳時腹部大動脈瘤に対して自作ステントグラフト (tapered type)+ 大腿動脈-大腿動脈交差バイパスによる治療を施行され外来経過観察を行っていた. 術後 31 カ月目に type I エンドリークを認め, 再手術となった. ステントグラフトが瘤内に落ち込むように migration し, 中枢からの type I エンドリークを認めたため中枢端にパーブを有する Zenith を選択し, Converter を用いて aorta-uniiliac type とすることとした. 手術は問題なく終了し, CT (computed tomography) でもエンドリークは消失した. 本症例で用いた自作ステントグラフトはパーブやナイチノールアンカーといった固定装置を有しておらず, エンドリークの原因の一因となったことが考えられる. (日血外会誌 2010;19:561-564)

索引用語: EVAR, migration

はじめに

腹部大動脈瘤(AAA)に対して低侵襲治療である EVAR(endovascular aneurysm repair)は増加傾向にあり当科でも 1999 年より自作ステントグラフトによる治療を行ってきた. 企業製ステントグラフトとしては 2006 年 7 月に米国 Cook 社製ゼニス AAA エンドバスキュラーグラフト(Zenith)を皮切りに 2007 年 5 月に Gore Excluder, 2008 年 2 月には Powerlink の 3 つが国内で認可されている. 今回 AAA に対する自作ステントグラフト治療後, 遠隔期のエンドリークに対して企業デバイスを使用することにより良好な結果を得たので報告する.

症 例

症 例: 74 歳, 女性

既往歴: 29 歳: 子宮筋腫手術, 63 歳: 急性大動脈解離にて上行, 弓部人工血管置換術を施行.

現病歴: 2006 年 4 月に腹部大動脈瘤(中枢ネック径 20×長さ 20 mm)に対して自作ステントグラフト内挿術(UBE+Z-stent, tapered type 径 24 → 14 mm×長さ 100 mm)+ 大腿動脈-大腿動脈交差バイパスを施行した(Fig. 1). 術後 computed tomography(CT)にてエンドリークは認めず定期的に単純 X 線撮影や CT を用いた外来経過観察を行っていたが 2008 年 12 月の CT にてステントグラフト中枢端が中枢ネックから瘤内へ落ち込むように migration し, type I エンドリークが発生していた. 今回, 追加治療目的にて入院となった.

現 症: 身長 145.2 cm, 体重 78.5 kg, BMI 38

胸骨正中・下腹部手術痕(+), 腹部拍動性腫瘍-触知(-)

CT: 瘤の最大径は 50 mm で瘤内に落ち込むようにステントグラフトが migration しており, 中枢からの type I エンドリークを認めた(Fig. 2). 大腿動脈-大腿動脈交差バイパスは開存していた.

BMI 38 と高度肥満で開腹歴も有することより再度ステントグラフトでの治療を行うこととした. 中枢側

久留米大学外科(Tel: 0942-35-3311)
〒830-0011 福岡県久留米市旭町 67
受付: 2009 年 10 月 7 日
受理: 2010 年 3 月 10 日




Fig. 1 The custom-made stent graft was placed under the renal artery.




Fig. 2 Preoperative computed tomography (CT) showed that the stent-graft dropped into abdominal aortic aneurysm.

ネックの固定を確実にするためにパーブを有する Zenith を選択し、Converter を用いて aorta-uniiliac type とすることとした。

手術：仰臥位，全身麻酔下，後腹膜アプローチにて右外腸骨動脈を露出，ヘパリン投与後 ACT (activated clotting time) が 200 秒を超えたのを確認した後に DSA (digital subtraction angiography) を行った。Stiff guide wire 下にステントグラフトメインボディー (TFB 24-103) を右外腸骨動脈より前回留置したステントグラフトを通過させて腎動脈上腹部大動脈まで誘導，再度 DSA にて腎動脈分岐部を確認後，留置した。次に Converter (ESC 24-12-80) を留置し aorta-uniiliac type へ変更し，最後に iliac leg (TFLE 14-54) をメインボディーと 1.5 連重なるように留置，バルーンカテーテル (TMP ロックバルーンカテーテル 7Fr.) にてステントグラフト中枢，末梢，連結部を圧着した。DSA にてエンドリークがないことを確認し，手術を終了した。

術後経過：肥満による合併症防止のため早期離床をすすめた。また術後 7 日目の造影 CT (Fig. 3) にてエンドリークは認めず，経過良好にて退院となった。2009 年 6 月 9 日現在，とくに問題なく経過している。

考 察

動脈瘤に対する低侵襲治療として 1991 年に Parodi ら¹⁾が手作りのステントグラフトを用いて腹部大動脈瘤の治療を行ったのが近年における EVAR の始まりで，当科でも 1999 年より自作ステントグラフトを使用した EVAR を行ってきた。2006 年 7 月からは企業製ステントグラフトも認可され，現在 3 種類のデバイスを使用することができる。EVAR は瘤を空置したまま大動脈病変の中核と末梢を密着し病変部を血流から隔離する治療であるため，エンドリークや migration といった特有の不具合が生じる可能性がある。初期の臨床的成功を得ても遠隔期に migration を生じて，2 次的エンドリークを招き，瘤径拡大や瘤破裂にいたる場合もあり^{2,3)}，Harris ら⁴⁾によると 2 次的エンドリーク率は 18% と報告されている。

本症例も術後 2 年 7 カ月でステントグラフト中枢端が瘤内に落ち込むように migration し，type I エンドリークを生じた。デバイスは aorta-uniiliac type で末梢に行くにしたがって taper 状に作製されており，大動脈分岐部で固定されたような状態で留置されている。よって永続的な血流の圧力が，瘤内に位置するステントグラフト部にかかることになり瘤内に落ち込む原因のひとつ

つになったと推測される。Malinaら⁵⁾は、2000年にすでにEVAR術後23例の病理学的検討を行い、内挿されたステントグラフトと大動脈壁との間は、数年経ても堅固には癒合しておらず、容易に引き離すことができたと報告している。また鬼塚ら⁶⁾もEVAR(woven Dacron+Z-stent)術後1年半を経てmigrationによるtype Iエンドリークから人工血管置換術に移行した症例で、ステントグラフトは容易に大動脈壁から取り外すことができたと報告している。Malinaら⁵⁾の病理学的検討ではDacron人工血管周囲には粗雑な血球成分があるのみで線維の間に入り込んでいるような細胞は認めなかったと報告されており、大動脈壁に生化学的に癒着しない人工血管が使用されたことも原因のひとつと考えられる。

自作デバイスは、個々の大動脈形状に合わせて作製したり、彎曲部の大彎側は比較的強固な骨格となるようステント同士を連結させ、瘤内への逸脱を防止してきた⁷⁾。一方、企業デバイスは個々の大動脈形状に合わせて作製することは困難で1種類で多数の症例に適合しなければならないため、ステントグラフト自体柔軟で、さまざまな大動脈形状に追随しなければならない。このような柔軟なステントグラフトではステントの拡張力だけではmigrationが危惧されるため、多くは中樞や末梢にhookやバープといった機械的な固定機器を装備している⁸⁾。本症例では中樞端にバープを有するZenithを選択したが、Zenithは中樞側ベアステントが腎動脈分岐部前後の大動脈に留置されるため十分な圧着部位が確保でき、また中樞端にバープを有することにより大動脈壁に強固に固定されることから遠隔期のmigrationを防止できるものと考えた。

またエンドリークに対する処置であるが、再EVARやsurgical conversionといった方法がある。Dattiloら⁹⁾は362例のEVAR症例のうち遠隔期で再EVAR 10.8%、surgical conversion 2.2%であったと報告している。当科ではtype IおよびIVエンドリークに対して再EVAR 2例(1.3%)、surgical conversion 3例(2.0%)を施行しているが、2期的治療成績は100%と良好であり、個々の症例に応じて再EVARあるいはsurgical conversionかを選択すれば良好な結果を得るものと考えられる。

結 語

腹部大動脈瘤に対する自作ステントグラフト治療後




Fig. 3 Postoperative computed tomography (CT) showed no endoleak.

のmigrationに対して企業製ステントグラフトであるZenithを使用し、良好な結果を得た症例を経験したので報告した。

文 献

- 1) Parodi JC, Palmaz JC, Barone HD. Transfemoral intraluminal graft implantation for abdominal aortic aneurysms. *Ann Vasc Surg* 1991;5:491-499.
- 2) Dake MD, Miller DC, Semba CP, et al. Transluminal placement of endovascular stent-grafts for the treatment of descending thoracic aortic aneurysms. *N Engl J Med* 1994;331:1729-1734.
- 3) Mitchell RS, Miller DC, Dake MD, et al. Thoracic aortic aneurysm repair with an endovascular stent-graft: the "first generation". *Ann Thorac Surg* 1999;67:1971-1974.
- 4) Harris PL, Dimitri S. Predicting failure of endovascular aneurysm repair. *Eur J Vasc Endovasc Surg* 1999;17:1-2.
- 5) Malina M, Brunkwall J, Ivancev K, et al. Endovascular healing is inadequate for fixation of Dacron stent-grafts in human aortoiliac vessels. *Eur J Vasc Endovasc Surg* 2000;19:5-11.
- 6) 鬼塚誠二, 田中厚寿, 明石英俊, 他. Stent-graft migration. *日血管内治療会誌* 2008;9:37-41.
- 7) 川口 聡, 石丸 新. 胸部大動脈瘤に対するステントグラフト内挿術. *Cardiac Practice* 2004;15:297-301.
- 8) Veerapen R, Dorandeu A, Serre I, et al. Improvement in

proximal aortic endograft fixation: An experimental study using different stent-grafts in human cadaveric aortas. *J Endovasc Ther* 2003;**10**:1101-1109.

9) Dattilo JB, Brewster DC, Fan CM, et al. Clinical failures

of endovascular abdominal aortic aneurysm repair: incidence, causes, and management. *J Vasc Surg* 2002;**35**:1137-1144.

A Case of Re-EVAR with a Zenith Device after Custom-made Stent Graft Placement for Abdominal Aortic Aneurysm

Hideki Sakashita, Atsuhisa Tanaka, Seiji Onitsuka, Shinnichi Hiromatsu, Hidetoshi Akashi and Shigeaki Aoyagi

Department of Surgery, Kurume University

Key words: Endovascular aneurysm repair, Migration

74-year-old woman had undergone endovascular aneurysm repair with a homemade stent graft at age 71 because of obesity. Follow-up computed tomography 31 months after this first operation showed a type I endoleak due to stent graft migration. A Zenith device, with a barb at the edge of proximal bare stent was chosen for re-EVAR. The patient was discharged with good course, with no evidence of endoleak. The reason for stent graft migration was thought to be that the custom-made stent graft had no device such as a barb or nitinol anchor to secure it in place.

(*Jpn J Vasc Surg* 2010;**19**:561-564)