

Kitoh H, Kaneko H, <u>Ishiguro N.</u>	Radiographic analysis of movements of the acetabulum and the femoral head after Salter innominate osteotomy.	J Pediatr Orthop.	29(8)	879-884	2009
Kitoh H, Kawasumi M, Kaneko H, <u>Ishiguro N.</u>	Differential effects of culture-expanded bone marrow cells on the regeneration of bone between the femoral and the tibial lengthenings.	J Pediatr Orthop.	29(6)	643-649	2009
Yuji Hirano, Toshihisa Kojima, Yasuhide Kanayama, Hisato Ishikawa, <u>Naoki Ishiguro.</u>	A case of lung tuberculosis in a patient with rheumatoid arthritis treated with infliximab after anti-tuberculosis chemoprophylaxis with isoniazid.	Modern Rheumatology.	19(3)	323-328	2009
Masayo Kojima, Toshihisa Kojima, <u>Naoki Ishiguro.</u> Takeshi Oguchi, Michinari Oba, Hiroki Tsuchiya, Fumiaki Sugiura, Toshiaki A Furukawa, Sadao Suzuki, Shinkan Tokudome.	Psychosocial factors, disease status, and quality of life in patients with rheumatoid arthritis.	Journal of Psychosomatic Research.	67(5)	425-431	2009
Hiroshi Kitoh, Takahiko Kitakoji, Motoaki Kawasumi, <u>Naoki Ishiguro.</u>	A histological and ultrastructural study of the iliac crest apophysis in legg-calve-perthes disease.	J Pediatr Orthop.	28	435-439	2008
T Kojima, M Kojima, K Noda, <u>N Ishiguro</u> , AR Poole.	Influences of menopause, aging, and gender on the cleavage of type II collagen in cartilage in relationship to bone turnover.	Menopause.	15	133-137	2008
<u>石黒直樹</u>	《特集／関節リウマチの新しい治療方針》 II 関節リウマチの最新薬物療法と理学療法 1. 薬物療法の基本原則と効果判定法.	整形外科	59	870-875	2008

石黒直樹	VI,リウマチ性疾患	整形外科	59	740-747	2008
------	------------	------	----	---------	------

書籍

著者氏名	論文タイトル名	書籍全体の編集者名	出版社名	出版年
		書籍名	出版地	ページ
石黒直樹	第2部各疾患別 NSAIDs の使い方 第2章整形外科領域 総論	佐野 統	羊土社	2010
		NSAIDs の選び方・使い方ハンドブック	東京	151-153
石黒直樹	序幕四場：変形性膝関節症を俯瞰する 軟骨の一生を振り返る-運動器を支える 気丈夫な組織	井原秀俊	全日本病院出版 会	2010
		老いを内包する膝～早期診断と 早期治療～	東京	7-13
石黒直樹	II 疾患編 16 運動器疾患 変形性膝関節 症	金澤一郎、永井良三	医学書院	2010
		今日の診断指針 第6版	東京	1526-1527
小嶋俊久、 石黒直樹	4、膝痛の補助診断 膝痛の血液検査	中村耕三	中山書店	2010
		整形外科臨床パサージュ2 膝 の痛みクリニカルプラクティス	東京	116-121
石黒直樹	XII.運動器系疾患 1.関節リウマチ	井樋栄二、大田健、岡本美孝、 河野 茂、武田雅俊、直江知樹、 春間 賢、本間之夫、森脇久隆、 山本哲也、吉村泰典	メディカルレビ ュー社	2010
		ポケット判 診療ガイドライン UP-TO DATE2010-2011	東京	553-562
石黒直樹	第7章リウマチ性疾患とその他類縁疾患 A.関節リウマチ	長野 昭、松下 隆、戸山芳昭、 安田和則、石黒直樹	南江堂	2010
		整形外科専門医テキスト	東京	303-314
石黒直樹	第7章リウマチ性疾患とその他類縁疾患 B.悪性関節リウマチ	長野 昭、松下 隆、戸山芳昭、 安田和則、石黒直樹	南江堂	2010
		整形外科専門医テキスト	東京	314-315
石黒直樹	5.関節リウマチ、慢性関節疾患および 骨壊死症 血友病性関節症	国分正一、岩谷 力、落合直之、 佛淵孝夫	医学書院	2010
		今日の整形外科治療指針第6版	東京	178-179
石黒直樹	5.関節リウマチ、慢性関節疾患および 骨壊死症 神経病性関節症 (Charcot 関 節)	国分正一、岩谷 力、落合直之、 佛淵孝夫	医学書院	2010
		今日の整形外科治療指針第6版	東京	179-180

<u>石黒直樹</u>	5.関節リウマチ、慢性関節疾患および骨壊死症 糖尿病性関節症	国分正一、岩谷 力、落合直之、佛淵孝夫	医学書院	2010
		今日の整形外科治療指針第6版	東京	180-181
<u>石黒直樹</u>	5.関節リウマチ、慢性関節疾患および骨壊死症 ステロイド性関節症	国分正一、岩谷 力、落合直之、佛淵孝夫	医学書院	2010
		今日の整形外科治療指針第6版	東京	181-182
<u>石黒直樹</u>	VI. 他科とのコンサルテーション 4. 整形外科	三森径世	文光堂	2010
		リウマチ・膠原病内科クリニカルスタンダード	東京	292-299

(竹内 勤)

雑誌

発表者氏名	論文タイトル名	発表誌名	巻号	ページ	出版年
Yamanaka H, Tanaka Y, Hoshi D, Inoue E, Saito K, Amano K, Kameda H, and <u>Takeuchi T.</u>	Efficacy of tocilizumab for rheumatoid arthritis patients in daily practice in Japan –message from REACTION study.	Mod Rheum.	in press		
Okuyama A, Nagasawa H, Suzuki K, Kameda H, Kondo H, Amano K, and <u>Takeuchi T.</u>	Fc gamma receptor IIIb polymorphism and usage of glucocorticoids at baseline are associated with infusion reactions to infliximab in patients with rheumatoid arthritis.	Ann Rheum Dis.	70	299-304	2011
Nagasawa H, Kameda H, Sekiguchi N, Amano K, and <u>Takeuchi T.</u>	Differences between the Health Assessment Questionnaire Disability Index (HAQ-DI) and the modified HAQ (mHAQ) score before and after infliximab treatment in patients with rheumatoid arthritis.	Mod Rheum.	20	337-342	2010
Kameda H, Ueki Y, Saito K, Nagaoka S, Hidaka T, Atsumi T, Tsukano M, Kasama T, Shiozawa S, Tanaka Y, <u>Takeuchi T.</u> and Japan Biological Agent Integrated Consortium (J-BASIC).	The comparison of efficacy and safety between continuation and discontinuation of methotrexate (MTX) at the commencement of etanercept in patients with active rheumatoid arthritis despite MTX therapy: 24-week results from the JESMR study.	Mod Rheum.	20	531-538	2010

Tsuzaka K, Itami Y, <u>Takeuchi T</u> , Shinozaki N, and Morishita T.	ADAMTS5 is a biomarker for prediction of the response to Infliximab in patients with rheumatoid arthritis.	J Rheum.	37	1454-1460	2010
<u>Takeuchi T</u> and Kameda H.	The Japanese experiences with biologic therapies for Rheumatoid Arthritis.	Nat Rev Rheum.	6	544-562	2010
Tanaka Y, <u>Takeuchi T</u> , Mimori T, Saito K, Nawata M, Kameda H, Nojima T, Miyasaka N, and Koike T.	Discontinuation of infliximab after attaining low disease activity in patients with rheumatoid arthritis, RRR (remission induction by Remicade in RA) study.	Ann Rheum Dis.	69	1286-1291	2010
Nagasawa H, Kameda H, Sekiguchi N, Amano K, and <u>Takeuchi T</u> .	Normalization of physical function by infliximab in RA patients: Factors associated with normal physical function.	Clin Exp Rheum.	26	365-372	2010
Komano Y, Harigai M, Koike R, Sugiyama H, Ogawa J, Saito K, Sekiguchi N, Inoo M, Onishi I, Ohashi H, Amamoto F, Miyata M, Kageyama G, Imaizumi K, Tokuda H, Okochi Y, Tanaka Y, <u>Takeuchi T</u> , and Miyasaka N.	Pneumocystis pneumonia in patients with rheumatoid arthritis treated with infliximab: a retrospective review and case-control study of 21 patients.	Arthritis & Rheum.	61	305-312	2009
Nishimoto N, Miyasaka N, Yamamoto K, Kawai S, <u>Takeuchi T</u> , and Azuma J.	Long-term safety and efficacy of tocilizumab, an anti-interleukin(IL)-6 receptor monoclonal antibody, in monotherapy, in patients with rheumatoid arthritis (the STREAM study): evidence of safety and efficacy in a 5-year extension study.	Ann Rheum Dis.	68	1580-1584	2009

Nishimoto N, Miyasaka N, Yamamoto K, Kawai S, <u>Takeuchi T</u> , Azuma J, and Kishimoto T.	Study of Active controlled Tocilizumab monotherapy for Rheumatoid Arthritis Patients with an Inadequate Response to Methotrexate (SATORI) : significantly reduction in disease activity and serum vascular endothelial growth factor by IL-6 receptor inhibition therapy.	Mod Rheum.	19	12-19	2009
Suzuki K, Kameda H, Amano K, Nagasawa H, Sekiguchi H, Nishi E, Ogawa H, Tsuzaka K, and <u>Takeuchi T</u> .	Single Center Prospective Study for Efficacy and Safety of Tacrolimus in Rheumatoid Arthritis.	Rheumatology Int.	29	431-436	2009
Nagasawa H, Kameda H, Sekiguchi N, Amano K, and <u>Takeuchi T</u> .	Improvement of the HAQ score by infliximab treatment in patients with RA: its association with disease activity and joint destruction.	Mod Rheum.	19	166-172	2009
<u>Takeuchi T</u> , Miyasaka N, Inoue K, Abe T, and Koike T.	Impact of through serum level on Radiographic and Clinical Response to Infliximab plus Methotrexate in Patients with Rheumatoid Arthritis: results from the RISING Study.	Mod Rheum.	19	478-487	2009
Hashimoto J, Garnero P, van der Heijde, Miyasaka N, Yamamoto K, Kawai S, <u>Takeuchi T</u> Yoshikawa H, and Nishimoto N.	A combination of biochemical markers of cartilages and markers of cartilage and bone turnover, radiographic image and body mass index to predict progression of joint destruction in patients with rheumatoid Arthritis treated with Disease modifying Anti-Rheumatic Drugs.	Mod Rheum.	19	273-282	2009
Tanino M, Matoba R, Nakamura S, Kameda H, Amano K, Okayama T, Nagasawa H, Suzuki K, Matsubara K, and <u>Takeuchi T</u> .	Prediction of efficacy of anti-TNF biologic agent, infliximab, for rheumatoid arthritis patients using a comprehensive transcriptome analysis of white blood cells.	Biochem Biophys Research Comm.	387	261-265	2009

Koike T, Harigai M, Inokuma S, Inoue K, Ishiguro N, Ryu J, <u>Takeuchi T</u> , Tanaka Y, Yamanaka H, Fujii K, Freundlich B, and Suzukawa M.	Post-marketing surveillance of the safety and efficacy of etanercept in Japan.	J Rheum.	36	898-906	2009
Sawada T, Inokuma S, Sato T, Ohtsuka T, Saeki Y, <u>Takeuchi T</u> , Matsuda T, Takemura T, and Sagawa A.	Leflunomide-induced interstitial lung disease: Prevalence and risk factors in Japanese patients with rheumatoid arthritis.	Rheumatology	48	1069-1072	2009
Sato T, Inokuma S, Sagawa A, Matsuda T, Takemura T, Ohtsuka T, Saeki Y, <u>Takeuchi T</u> , and Sawada T.	Factors associated with fatal outcome of leflunomide-induced lung injury in Japanese patients with rheumatoid arthritis.	Rheumatology	48	1265-1268	2009
Kameda H, Amano K, Nagasawa H, Ogawa H, Sekiguchi N, Takei H, Suzuki K, and <u>Takeuchi T</u> .	Notable difference between the development of vertebral fracture and osteonecrosis of femoral head in patients treated with high-dose glucocorticoids for systemic rheumatic diseases.	Internal Med.	48	1931-1938	2009
<u>Takeuchi T</u> , Tatsuki Y, Nogami Y, Ishiguro N, Tanaka Y, Yamanaka H, Harigai Y, Ryu J, Inoue K, Kondo H, Inokuma S, Kamatani N, Ochi T, and koike T. Post-marketing Surveillance of the Safety Profile of Tanaka Y, Takeuchi T, Inoue E, Saito K, Sekiguchi N, Iikuni N, Nawata M, Kameda H, Shinozaki M, Iwata S, and Amano K, and Yamanaka H.	Retrospective clinical study on the notable efficacy and related factors of infliximab therapy in a rheumatoid arthritis management group in Japan: One-year clinical and radiographic outcomes (RECONFIRM-II). Mod Rheum 18:146-52, 2008 Infliximab in 5,000 Japanese Patients with Rheumatoid Arthritis.	Ann Rheum Dis.	67	189-194	2008

Sekiguchi N, Kawauchi S, Furuya T, Matsuda K, Ando S, Ogasawara M, Inaba N, Abe T, Ito S, and <u>Takeuchi T.</u>	Monitoring of cDNA microarray profile in peripheral blood during infliximab treatment of Rheumatoid Arthritis patients.	Rheumatology.	47	780-788	2008
Tanaka Y, <u>Takeuchi T.</u> , Inoue E, Saito K, Sekiguchi N, Iikuni N, Nawata M, Kameda H, Shinozaki M, Iwata S, and Amano K, and Yamanaka H.	Retrospective clinical study on the notable efficacy and related factors of infliximab therapy in a rheumatoid arthritis management group in Japan: One-year clinical and radiographic outcomes (RECONFIRM-II).	Mod Rheum.	18	146-152	2008
Tokuda H, Sakai F, Yamada H, Johkoh T, Imamura A, Dohi M, Hirakata M, Yamada T, Kamatani N, Kikuchi Y, Sugii S, <u>Takeuchi T.</u> , Tateda K, and Goto H.	Clinical radiological features of pneumocystis pneumonia in patients with Rheumatoid Arthritis, in comparison with Methotrexate pneumonitis and Pneumocystis pneumonia in acquired immunodeficiency syndrome: A multicenter study.	Intern Med.	47	915-923	2008
<u>Takeuchi T.</u> , Yamanaka H, Inoue E, Nagasawa Y, Nawata M, Ikari K, Saito K, Sekiguchi N, Saito E, Kameda H, Iwata S, Mochizuki T, Amano K, and Tanaka Y.	Retrospective clinical study on the notable efficacy and related factors of infliximab therapy in a rheumatoid arthritis management group in Japan: One-year outcome of joint destruction (RECONFIRM-2J).	Mod Rheum.	18	447-454	2008
Inokuma S, Sato T, Sagawa A, Matsuda T, Takemura T, Ohtsuka T, Saeki Y, <u>Takeuchi T.</u> , and Sawada T.	Proposals for leflunomide use to avoid lung injury in patients with rheumatoid arthritis.	Mod Rheum.	18	442-446	2008

(桃原 茂樹)

雑誌

発表者氏名	論文タイトル名	発表誌名	巻号	ページ	出版年
Kochi Y, Okada Y, Suzuki A, Ikari K, Terao C, Takahashi A, Yamazaki K, Hosono N, Myouzen K, Tsunoda T, Kamatani N, Furuichi T, Ikegawa S, Ohmura K, Mimori T, Matsuda F, Iwamoto T, <u>Momohara S</u> , Yamanaka H, Yamada R, Kubo M, Nakamura Y, Yamamoto K.	A regulatory variant in CCR6 is associated with rheumatoid arthritis susceptibility.	Nat Genet.	42(6)	515-519	2010
Kawakami K, Ikari K, Yano K, <u>Momohara S</u> .	Adductor muscle strain in the elderly patient suspected the femoral neck fracture.	Arthritis Rheum.	60 (11)	3446	2009
Saito S, <u>Momohara S</u> , Taniguchi A, Yamanaka H.	The intra-articular efficacy of hyaluronate injections in the treatment of rheumatoid arthritis.	Mod Rheumatol.	19(6)	643-651	2009
<u>Momohara S</u> , Ikari K, Mochizuki T, Kawamura K, Tsukahara S, Toki H, Hara M, Kamatani N, Yamanaka H, Tomatsu T.	Declining use of synovectomy surgery for rheumatoid arthritis patients in Japan.	Ann Rheum Dis.	68(2)	291-292	2009
Niki Y, Mochizuki T, <u>Momohara S</u> , Saito S, Toyama Y, Matsumoto H.	Is Minimally Invasive Surgery in Total Knee Arthroplasty Really Minimally Invasive Surgery?	J Arthroplasty.	24(4)	499-504	2009
Tsukahara S, Ikari K, Sato E, Yamanaka H, Hara M, Tomatsu T, <u>Momohara S</u> , Kamatani N.	A polymorphism in Fc gamma IIIA receptor coding gene is a possible genetic marker to predict primary response to infliximab in Japanese rheumatoid arthritis patients.	Ann Rheum Dis.	67(12)	1791-1792	2008

<u>Momohara S</u> , Okamoto H, Yamanaka H.	Chondrocyte of rheumatoid arthritis serve as a source of intra-articular acute-phase serum amyloid A protein.	Clin Chim Acta.	398(1-2)	155-156	2008
<u>Momohara S</u> , Inoue E, Ikari K, Tsukahara S, Kawamura K, Hara M, Kamatani N, Yamanaka H, Tomatsu T.	Risk Factors for Wrist Surgery in Rheumatoid Arthritis.	Clin Rheumatol.	27(11)	1387-1391	2008
Kobayashi S, <u>Momohara S</u> , Kamatani N, Okamoto H.	Molecular aspects of rheumatoid arthritis: role of environmental factors.	FEBS J.	275(18)	4456-4462	2008
Iwamoto T, Okamoto H, Toyama Y, <u>Momohara S</u> .	Molecular aspects of rheumatoid arthritis: chemokines in the joints of patients.	FEBS J.	275(18)	4448-4455	2008
Kobayashi S, Ikari K, Kaneko H, Kochi Y, Yamamoto K, Shimane K, Nakamura Y, Toyama Y, Mochizuki T, Tsukahara S, Kawaguchi Y, Terai C, Hara M, Tomatsu T, Yamanaka H, Horiuchi T, Tao K, Yasumoto K, Hamada D, Yasui N, Inoue H, Itakura M, Okamoto H, Kamatani N, <u>Momohara S</u> .	Association of STAT4 with susceptibility to rheumatoid arthritis and systemic lupus erythematosus in Japanese.	Arthritis Rheum.	58(7)	1940-1946	2008
Kobayashi S, Okamoto H, Iwamoto T, Toyama Y, Tomatsu T, Yamanaka H, <u>Momohara S</u> .	A role for the aryl hydrocarbon receptor and the dioxin TCDD in rheumatoid arthritis.	Rheumatology (Oxford).	47(9)	1317-1322	2008
Nishimoto K, Ikari K, Mochizuki T, Tomatsu T, Toyama Y, Hara M, Yamanaka H, Kamatani N, <u>Momohara S</u> .	Lack of association between PADI4 and functional severity in Japanese rheumatoid arthritis patients.	Ann Rheum Dis.	67(3)	431-432	2008

Tsukahara S, Iwamoto T, Ikari K, Inoue E, Tomatsu T, Hara M, Yamanaka H, Kamatani N, <u>Momohara S.</u>	CTLA-4 CT60 polymorphism is not an independent genetic risk marker of rheumatoid arthritis in Japanese.	Ann Rheum Dis.	67(3)	428-429	2008
Tsukahara S, Shinozaki M, Ikari K, Mochizuki T, Inoue E, Tomatsu T, Hara M, Yamanaka H, Kamatani N, <u>Momohara S.</u>	Effect of matrix metalloproteinase-3 functional SNP on serum matrix metalloproteinase-3 level and outcome measures in Japanese RA patients.	Rheumatology (Oxford).	47(1)	41-44	2008

(津谷 喜一郎)

### 雑誌

発表者氏名	論文タイトル名	発表誌名	巻号	ページ	出版年
津谷喜一郎、五十嵐中、 菊田健太郎	リウマチ治療の薬剤経済学	治療学	44 (10)	1135-1138	2010
津谷喜一郎、五十嵐中	生物学的製剤と薬剤経済評価	日本臨床	68 (suppl.)	653-657	2010
津谷喜一郎、五十嵐中、 白岩 健	分子標的薬の薬剤経済学	日本臨床	68 (10)	1923-1928	2010
五十嵐中、 <u>津谷喜一郎</u>	生物学的製剤の効果と費用負担の バランス	Journal of Clinical Rehabilitation	18 (2)	124-129	2009

### 書籍

著者氏名	論文タイトル名	書籍全体の編集者名	出版社名	出版年
		書籍名	出版地	ページ
津谷喜一郎、 五十嵐中	関節リウマチの薬剤経済学	宮坂信之	永井書店	2009
		よくわかる関節リウマチのすべ て	東京	273-278

(中島 亜矢子)

雑誌

発表者氏名	論文タイトル名	発表誌名	巻号	ページ	出版年
Shidara K, Inoue E, Tanaka E, Hoshi D, Seto Y, <u>Nakajima A</u> , Momohara S, Taniguchi A, Yamanaka H.	Comparison of the second and third generation anti-cyclic citrullinated peptide antibody assays in the diagnosis of Japanese patients with rheumatoid arthritis.	Rheumatol Int.		in press	
Toru Yamada, <u>Ayako</u> <u>Nakajima</u> , Eisuke Inoue, Eiichi Tanaka, Atsuo Taniguchi, Shigeki Momohara, Naoyuki Kamatani, Hisashi Yamanaka.	Incidence of malignancy in Japanese patients with rheumatoid arthritis in IORRA cohort.	Rheumatol Int.		in press	
<u>Ayako Nakajima</u> , Eisuke Inoue, Gurkirpal Singh, Eri Sato, Kumi Shidara, Daisuke Hoshi, Akiko Kiire, Masako Hara, Shigeki Momohara, Atsuo Taniguchi, Naoyuki Kamatani, Hisashi Yamanaka.	Mortality of and cause of death in Japanese patients with rheumatoid arthritis based on a large observational cohort, IORRA.	Scand J Rheumatol.	39	360-369	2010
Shidara K, Hoshi D, Inoue E, Yamada T, <u>Nakajima</u> <u>A</u> , Taniguchi A, Hara M, Momohara S, Kamatani N, Yamanaka H.	Incidence of and risk factors for interstitial pneumonia in patients with rheumatoid arthritis in a large Japanese observational cohort, IORRA.	Mod Rheumatol.	20(3)	280-286	2010

