

- 24 Fujino S, Andoh A, Bamba S, Ogawa A, Hata K, Araki Y *et al*. Increased expression of interleukin 17 in inflammatory bowel disease. *Gut* 2003; **52**: 65–70.
- 25 Zrioual S, Ecochard R, Tournadre A, Lenief V, Cazalis MA, Miossec P. Genome-wide comparison between IL-17A- and IL-17F-induced effects in human rheumatoid arthritis synovocytes. *J Immunol* 2009; **182**: 3112–3120.
- 26 Antonysamy MA, Fanslow WC, Fu F, Li W, Qian S, Troutt AB *et al*. Evidence for a role of IL-17 in organ allograft rejection: IL-17 promotes the functional differentiation of dendritic cell progenitors. *J Immunol* 1999; **162**: 577–584.
- 27 Vanaudenaerde BM, Dupont LJ, Wuyts WA, Verbeke EK, Meyts I, Bullens DM *et al*. The role of interleukin-17 during acute rejection after lung transplantation. *Eur Respir J* 2006; **27**: 779–787.
- 28 Van Kooten C, Boonstra JG, Paape ME, Fossiez F, Banchereau J, Lebecque S *et al*. Interleukin-17 activates human renal epithelial cells *in vitro* and is expressed during renal allograft rejection. *J Am Soc Nephrol* 1998; **9**: 1526–1534.
- 29 Loong CC, Hsieh HG, Lui WY, Chen A, Lin CY. Evidence for the early involvement of interleukin 17 in human and experimental renal allograft rejection. *J Pathol* 2002; **197**: 322–332.
- 30 Yi T, Chen Y, Wang L, Du G, Huang D, Zhao D *et al*. Reciprocal differentiation and tissue-specific pathogenesis of Th1, Th2, and Th17 cells in graft-versus-host disease. *Blood* 2009; **114**: 3101–3112.
- 31 Yi T, Zhao D, Lin CL, Zhang C, Chen Y, Todorov I *et al*. Absence of donor Th17 leads to augmented Th1 differentiation and exacerbated acute graft-versus-host disease. *Blood* 2008; **112**: 2101–2110.
- 32 Tawara I, Maeda Y, Sun Y, Lowler KP, Liu C, Toubai T *et al*. Combined Th2 cytokine deficiency in donor T cells aggravates experimental acute graft-vs-host disease. *Exp Hematol* 2008; **36**: 988–996.
- 33 Kappel LW, Goldberg GL, King CG, Suh DY, Smith OM, Ligh C *et al*. IL-17 contributes to CD4-mediated graft-versus-host disease. *Blood* 2009; **113**: 945–952.
- 34 Icozian C, Yu Y, Liu C, Liang Y, Yi T, Anasetti C *et al*. Th17 cells are sufficient but not necessary to induce acute graft-versus-host disease. *Biol Blood Marrow Transplant* 2009; **16**: 170–178.
- 35 Carlson MJ, West ML, Coghill JM, Panoskaltis-Mortari A, Blazar BR, Serody JS. *In vitro*-differentiated TH17 cells mediate lethal acute graft-versus-host disease with severe cutaneous and pulmonary pathologic manifestations. *Blood* 2009; **113**: 1365–1374.
- 36 Arisawa T, Tahara T, Shibata T, Nagasaka M, Nakamura M, Kamiya Y *et al*. The influence of polymorphisms of interleukin-17A and interleukin-17F genes on the susceptibility to ulcerative colitis. *J Clin Immunol* 2008; **28**: 44–49.
- 37 Furuya T, Hakoda M, Ichikawa N, Higami K, Nanke Y, Yago T *et al*. Associations between HLA-DRB1, RANK, RANKL, OPG, and IL-17 genotypes and disease severity phenotypes in Japanese patients with early rheumatoid arthritis. *Clin Rheumatol* 2007; **26**: 2137–2141.
- 38 Nordang GB, Viken MK, Hollis-Moffatt JE, Merriman TR, Forre OT, Helgetveit K *et al*. Association analysis of the interleukin 17A gene in Caucasian rheumatoid arthritis patients from Norway and New Zealand. *Rheumatology (Oxford)* 2009; **48**: 367–370.
- 39 Southam L, Heath O, Chapman K, Loughlin J. Association analysis of the interleukin 17 genes IL17A and IL17F as potential osteoarthritis susceptibility loci. *Ann Rheum Dis* 2006; **65**: 556–557.
- 40 Kawase T, Morishima Y, Matsuo K, Kashiwase K, Inoko H, Saji H *et al*. High-risk HLA allele mismatch combinations responsible for severe acute graft-versus-host disease and implication for its molecular mechanism. *Blood* 2007; **110**: 2235–2241.
- 41 Sasazuki T, Juji T, Morishima Y, Kinukawa N, Kashiwabara H, Inoko H *et al*. Effect of matching of class I HLA alleles on clinical outcome after transplantation of hematopoietic stem cells from an unrelated donor. Japan Marrow Donor Program. *N Engl J Med* 1998; **339**: 1177–1185.
- 42 Morishima Y, Yabe T, Matsuo K, Kashiwase K, Inoko H, Saji H *et al*. Effects of HLA allele and killer immunoglobulin-like receptor ligand matching on clinical outcome in leukemia patients undergoing transplantation with T-cell-replete marrow from an unrelated donor. *Biol Blood Marrow Transplant* 2007; **13**: 315–328.
- 43 Livak KJ. Allelic discrimination using fluorogenic probes and the 5' nuclease assay. *Genet Anal* 1999; **14**: 143–149.
- 44 Przepiorka D, Weisdorf D, Martin P, Klingemann HG, Beatty P, Hows J *et al*. 1994 Consensus Conference on Acute GVHD Grading. *Bone Marrow Transplant* 1995; **15**: 825–828.
- 45 Shulman HM, Sullivan KM, Weiden PL, McDonald GB, Striker GE, Sale GE *et al*. Chronic graft-versus-host syndrome in man. A long-term clinicopathologic study of 20 Seattle patients. *Am J Med* 1980; **69**: 204–217.
- 46 Scrucca L, Santucci A, Aversa F. Competing risk analysis using R: an easy guide for clinicians. *Bone Marrow Transplant* 2007; **40**: 381–387.
- 47 Gooley TA, Leisenring W, Crowley J, Storer BE. Estimation of failure probabilities in the presence of competing risks: new representations of old estimators. *Stat Med* 1999; **18**: 695–706.
- 48 Shibata T, Tahara T, Hirata I, Arisawa T. Genetic polymorphism of interleukin-17A and -17F genes in gastric carcinogenesis. *Hum Immunol* 2009; **70**: 547–551.
- 49 Alexandra HF, Daniel W, Steven P, Gerard S, John RW, Stephanie JL *et al*. National Institutes of Health Consensus Development Project on Criteria for Clinical Trials in Chronic Graft-versus-Host Disease: I. Diagnosis and Staging Working Group Report. *Biol Blood and Marrow Transplantation* 2005; **11**: 945–956.
- 50 Murashige N, Kami M, Mori S, Katayama Y, Kobayashi K, Onishi Y *et al*. Characterization of acute graft-versus-host disease following reduced-intensity stem-cell transplantation from an HLA-identical related donor. *Am J Hematol* 2008; **83**: 630–634.
- 51 Vigorito AC, Campregher PV, Storer BE, Carpenter PA, Moravec CK, Kiem H-P *et al*. Evaluation of NIH consensus criteria for classification of late acute and chronic GVHD. *Blood* 2009; **114**: 702–708.
- 52 Shlomchik WD, Couzens MS, Tang CB, McNiff J, Robert ME, Liu J *et al*. Prevention of graft versus host disease by inactivation of host antigen-presenting cells. *Science* 1999; **285**: 412–415.
- 53 Teshima T, Ordemann R, Reddy P, Gagin S, Liu C, Cooke KR *et al*. Acute graft-versus-host disease does not require alloantigen expression on host epithelium. *Nat Med* 2002; **8**: 575–581.
- 54 Duffner UA, Maeda Y, Cooke KR, Reddy P, Ordemann R, Liu C *et al*. Host dendritic cells alone are sufficient to initiate acute graft-versus-host disease. *J Immunol* 2004; **172**: 7393–7398.
- 55 Goldstone AH, Richards SM, Lazarus HM, Tallman MS, Buck G, Fielding AK *et al*. In adults with standard-risk acute lymphoblastic leukemia, the greatest benefit is achieved from a matched sibling allogeneic transplantation in first complete remission, and an autologous transplantation is less effective than conventional consolidation/maintenance chemotherapy in all patients: final results of the International ALL

- Trial (MRC UKALL XII/ECOG E2993). *Blood* 2008; **111**: 1827–1833.
- 56 Thomas E, Buckner C, Banaji M, Clift R, Fefer A, Flournoy N et al. One hundred patients with acute leukemia treated by chemotherapy, total body irradiation, and allogeneic marrow transplantation. *Blood* 1977; **49**: 511–533.
- 57 Boeckh M, Nichols WG. The impact of cytomegalovirus serostatus of donor and recipient before hematopoietic stem cell transplantation in the era of antiviral prophylaxis and preemptive therapy. *Blood* 2004; **103**: 2003–2008.
- 58 Kollman C, Howe CW, Anasetti C, Antin JH, Davies SM, Filipovich AH et al. Donor characteristics as risk factors in recipients after transplantation of bone marrow from unrelated donors: the effect of donor age. *Blood* 2001; **98**: 2043–2051.
- 59 Miller RA. The aging immune system: primer and prospectus. *Science* 1996; **273**: 70–74.

## ORIGINAL ARTICLE

# A single-nucleotide polymorphism of the Fcγ receptor type IIIA gene in the recipient predicts transplant outcomes after HLA fully matched unrelated BMT for myeloid malignancies

A Takami<sup>1,13</sup>, JL Espinoza<sup>1,13</sup>, M Onizuka<sup>2</sup>, K Ishiyama<sup>1</sup>, T Kawase<sup>3</sup>, Y Kanda<sup>4</sup>, H Sao<sup>5</sup>, H Akiyama<sup>6</sup>, K Miyamura<sup>7</sup>, S Okamoto<sup>8</sup>, M Inoue<sup>9</sup>, S Ohtake<sup>1</sup>, T Fukuda<sup>10</sup>, Y Morishima<sup>11</sup>, Y Kodera<sup>12</sup> and S Nakao<sup>1</sup>, for the Japan Marrow Donor Program

<sup>1</sup>Department of Hematology and Oncology, Kanazawa University Hospital, Kanazawa, Japan; <sup>2</sup>Department of Hematology and Oncology, Tokai University School of Medicine, Isehara, Japan; <sup>3</sup>Division of Epidemiology, Aichi Cancer Center Research Institute, Nagoya, Japan; <sup>4</sup>Division of Hematology, Saitama Medical Center, Jichi Medical University, Saitama, Japan; <sup>5</sup>Department of Hematology, Meitetsu Hospital, Nagoya, Japan; <sup>6</sup>Hematology Division, Tokyo Metropolitan Cancer and Infectious Diseases Center, Komagome Hospital, Tokyo, Japan; <sup>7</sup>Department of Hematology, Japanese Red Cross Nagoya First Hospital, Nagoya, Japan; <sup>8</sup>Division of Hematology, Department of Medicine, Keio University School of Medicine, Tokyo, Japan; <sup>9</sup>Department of Hematology and Oncology, Osaka Medical Center and Research Institute for Maternal and Child Health, Osaka, Japan; <sup>10</sup>Hematopoietic Stem Cell Transplantation Unit, National Cancer Center Hospital, Tokyo, Japan; <sup>11</sup>Department of Hematology and Cell Therapy, Aichi Cancer Center, Nagoya, Japan and <sup>12</sup>Department of Promotion for Blood and Marrow Transplantation, Aichi Medical University, Nagoya, Japan

Fcγ receptor type IIIA (FCGR3A) has a functional single-nucleotide polymorphism (rs396991), at which a G-to-T-point mutation results in an amino acid substitution at position 158 (valine to phenylalanine; V158F). This study examined the effect of the FCGR3A polymorphism in donors and recipients on the clinical outcomes in unrelated HLA fully matched myeloablative BMT. The FCGR3A-V158F genotype was retrospectively analyzed in a total of 99 recipients with myeloid malignancies, and their unrelated donors. The presence of the 158V genotype in recipients showed a statistically better OS (adjusted hazard ratio (HR) 0.49; 95% confidence interval (CI) 0.26–0.93;  $P=0.03$ ) and TRM (HR 0.30; 95% CI 0.14–0.67;  $P=0.003$ ) without significant influence on the relapse rate. The recipient 158V genotype was also associated with a significantly reduced risk of chronic GVHD (HR 0.45; 95% CI 0.20–0.99;  $P=0.049$ ) and a trend toward a reduced risk of grade II–IV acute GVHD (HR 0.55; 95% CI 0.27–1.10;  $P=0.09$ ), leading to a significantly reduced GVHD-related mortality (HR 0.22; 95% CI 0.06–0.77;  $P=0.02$ ). The donor FCGR3A polymorphism did not have any effect on the transplant outcomes. These results suggest an association between the recipient FCGR3A genotype and the clinical outcomes after BMT.

*Bone Marrow Transplantation* (2011) **46**, 238–243; doi:10.1038/bmt.2010.88; published online 19 April 2010  
**Keywords:** FCGR3A; unrelated donor; single-nucleotide polymorphism

## Introduction

Hematopoietic SCT is a potentially curative therapy in a range of malignant and nonmalignant diseases. However, its utility is limited because of transplant-related life-threatening complications, including GVHD, infections and disease relapse.<sup>1</sup> Although HLA matching represents the major genetic determinant in clinical outcome after allo-SCT, recent evidence suggests that non-HLA immune-associated genes are also implicated.<sup>2</sup> Previous investigations have revealed that several single-nucleotide polymorphisms (SNPs) that effect individual immune response to infections and inflammatory reactions are associated with transplant outcomes.<sup>3–9</sup>

Fcγ receptor type IIIA (FCGR3A), a low-affinity receptor capable of interaction with complexed or monomeric IgG, is expressed on neutrophils, eosinophils, natural killer cells, macrophages, monocytes, DC, γδ-positive T cells and keratinocytes.<sup>10–13</sup> FCGR3A mediates Ab-dependent cell-mediated cytotoxicity, phagocytosis, cytokine production and regulation of Ig production. FCGR3A has a functional SNP (rs396991), at which a G-to-T point mutation results in an amino acid substitution at position 158 (valine to phenylalanine; V158F) in the second Ig-like domain.<sup>14</sup> The cells bearing the FCGR3A-158V genotype show a higher affinity for IgG1 and IgG3 than those without 158V, and are capable of binding IgG4,<sup>15</sup> and thus

Correspondence: Dr A Takami, Department of Hematology and Oncology, Kanazawa University Hospital, 13-1 Takaramachi, Kanazawa, 920-8641, Japan.

E-mail: takami@med3.m.kanazawa-u.ac.jp

<sup>13</sup>These authors contributed equally to this work.

Received 7 December 2009; revised 15 January 2010; accepted 23 January 2010; published online 19 April 2010

can exert Ab-dependent cell-mediated cytotoxicity more efficiently.<sup>16</sup> The 158V genotype is associated with susceptibility to rheumatoid arthritis<sup>17</sup> and immune-mediated thrombocytopenic purpura,<sup>18</sup> better clinical response to rituximab in B-cell lymphomas<sup>19,20</sup> and a lower risk of recurrent periodontitis.<sup>21</sup> In contrast, systemic lupus erythematosus and better clinical outcome of cetuximab against metastatic colorectal cancer correlates with the 158F genotype.<sup>22,23</sup> This study analyzed the effect of donor and recipient SNP (rs396991) in the FCGR3A gene on the clinical outcomes in patients after allogeneic myeloablative BMT using an HLA allele-matched unrelated donor. The data show that the presence of the FCGR3A-158V genotype in the recipient was associated with significantly better transplant outcomes on the OS, TRM and GVHD.

## Patients and methods

### Patients

FCGR3A genotyping was performed on a total of 99 recipients with myeloid malignancies and their unrelated donors who underwent transplantation after myeloablative conditioning through the JMDP (Japan Marrow Donor Program) with T-cell-replete marrow from an HLA-A, -B, -C and -DRB1 allele-matched donor between November 1995 and March 2000. HLA genotypes of HLA-A, -B, -C and -DRB1 allele of patient and donor were determined by the Luminex microbead method described previously (Luminex 100 System; Luminex, Austin, TX, USA).<sup>24,25</sup> No patient had a history of any previous transplantation. The final clinical survey of these patients was completed by 1 November 2007. Diagnoses were AML in 47 (47%), CML in 42 (42%) and myelodysplastic syndrome in 10 patients (10%; Table 1). The recipients were defined as having standard risk disease if they had AML in first CR, CML in any chronic phase or myelodysplastic syndrome. All others were designated as high-risk disease. CYA- or tacrolimus-based regimens were used in all patients for GVHD prophylaxis, and anti-T-cell therapy, such as antithymocyte globulin and *ex vivo* T-cell depletion, was not. All patients and donors gave their written informed consent to participate in molecular studies of this nature according to the declaration of Helsinki at the time of transplantation. The project was approved by the institutional review board of Kanazawa University Graduate School of Medicine and JMDP.

### FCGR3A rs3969913 genotyping

Genotyping of FCGR3A was performed using the TaqMan-Allelic discrimination method<sup>26</sup> with a 7900-HT Real Time PCR system (Applied Biosystems, Foster City, CA, USA), and results were analyzed using the Allelic Discrimination software program (Applied Biosystems). The genotyping assay was conducted in 96-well PCR plates. The amplification reaction contained template DNA, the TaqMan universal master mix and the specific probe rs396991 designed for SNP of FCGR3A (product No C\_25815666\_10; Applied Biosystems).

**Table 1** Donor and recipient characteristics

Variable	Myeloid malignancies (n = 99)				P-value
	Recipient FCGR3A genotype				
	158V positive n = 46, 46%		158V negative n = 53, 54%		
	No.	Ratio	No.	Ratio	
<i>Age, years</i>					
Recipient					0.89
Median		31		36	
Range		8–49		1–50	
Donor					0.45
Median		33		31	
Range		21–50		22–48	
<i>Donor FCGR3A genotype</i>					0.15
With 158V (V/F or V/V)	24	86%	20	83%	
Without 158V (F/F)	22	79%	33	138%	
<i>Sex, male</i>					0.68
Recipient	30	65%	35	66%	
Donor	29	63%	39	74%	
<i>Recipient/donor sex</i>					0.69
Sex matched	25	54%	31	58%	
Male/female	11	24%	9	17%	
Female/male	10	22%	13	25%	
<i>Disease</i>					0.48
AML	23	50%	24	45%	
MDS	6	13%	4	8%	
CML	17	37%	25	47%	
<i>Disease risk</i>					0.26
Low	30	65%	40	75%	
High	16	35%	13	25%	
<i>ABO matching</i>					0.69
Match	26	57%	32	60%	
Major mismatch	10	22%	13	25%	
Minor mismatch	10	22%	8	15%	
<i>Conditioning regimen</i>					0.25
With TBI	39	85%	40	75%	
Without TBI	7	15%	13	25%	
<i>Pre-transplant CMV serostatus</i>					0.28
CMV-negative recipient	9	20%	13	25%	
Missing	7	15%	3	6%	
<i>GVHD prophylaxis</i>					0.20
With CYA	40	87%	50	94%	
With tacrolimus	6	13%	3	6%	
<i>TNC, × 10<sup>8</sup>/kg</i>					0.45
Median		5.4		5.6	
Range		2.3–16.2		2.4–52.6	
Engraftment	1	2%	2	4%	0.64

Abbreviations: FCGR3A = Fcγ receptor type IIIA; MDS = myelodysplastic syndrome; TNC = total nucleated cell count harvested.

### Data management and statistic analysis

Data were collected by the JMDP using a standardized report form. Follow-up reports were submitted at 100 days, 1 year and annually after transplantation. The pre-transplant CMV serostatus was routinely tested only for

**Table 2** Univariate analysis of the association between the FCGR3A genotype and the clinical outcomes after transplantation

	Donor FCGR3A genotype			Recipient FCGR3A genotype		
	With 158V	Without 158V	P-value	With 158V	Without 158V	P-value
No.	44	55		46	53	
5-year OS	50%	60%	0.52	65%	49%	0.15
5-year TRM	36%	27%	0.28	22%	40%	0.07
5-year relapse	16%	16%	0.92	17%	15%	0.79
II–IV aGVHD	39%	40%	0.81	33%	45%	0.19
cGVHD	34%	36%	0.81	26%	43%	0.07
GVHD-related mortality	15%	18%	0.63	7%	25%	0.01
Infection-related mortality	27%	20%	0.46	20%	26%	0.33

Abbreviations: aGVHD = acute GVHD; cGVHD = chronic GVHD; FCGR3A = Fcγ receptor type IIIA.

patients but not for their donors. Engraftment was confirmed by an ANC of  $>0.5 \times 10^9/L$  for at least 3 consecutive days. Acute and chronic GVHD were diagnosed and graded using established criteria.<sup>27,28</sup> The OS was defined as the number of days from transplantation to death from any cause. Disease relapse was defined as the number of days from transplantation to disease relapse. TRM was defined as death without relapse. Any patients who were alive at the last-follow-up date were censored. The data on the causative microbes of infections, post-mortem changes in the cause of death and staging of acute GVHD, as well as the data on supportive care including infection prophylaxis and therapy of GVHD, which were given on institution basis, were not available in this cohort. The analysis was performed using the Excel 2007 (Microsoft Corp., Redmond, WA, USA), OriginPro version 8.0J (Lightstone Inc., Tokyo, Japan) and R (The R Foundation for Statistical Computing, Perugia, Italy) software programs.<sup>29</sup> The probability of OS was calculated using the Kaplan–Meier method and compared using the log-rank test. The probabilities of TRM, disease relapse, acute GVHD, chronic GVHD and each cause of death were compared using the Grey test<sup>30</sup> and analyzed using the cumulative incidence analysis,<sup>29</sup> considering relapse, death without disease relapse, death without acute GVHD, death without chronic GVHD and death without each cause as respective competing risks. The variables included the recipient age at the time of transplantation, sex, CMV serostatus before transplantation, disease characteristic (disease type and disease risk at transplantation), donor characteristics (age, sex, sex compatibility and ABO compatibility), transplant characteristics (TBI-containing regimen, tacrolimus vs CYA and total nucleated cell count harvested per recipient weight). The median values were used as the cutoff point for continuous variables. The  $\chi^2$  test and Mann–Whitney test were used to compare two groups. The Hardy–Weinberg equilibrium for the FCGR3A gene polymorphism was tested using the Haploview program.<sup>5</sup> Multivariate Cox models were used to evaluate the hazard ratio (HR) associated with the FCGR3A polymorphism. Covariates found to be significant in univariate analyses ( $P \leq 0.10$ ) were included in the models. The *P*-values were two sided and outcomes were considered to be significant with  $P \leq 0.05$  in both the univariate and multivariate analyses.

**Results**

*Frequencies of the FCGR3A genotyping*

The FCGR3A gene polymorphism (rs396991) was analyzed in 99 unrelated BM donor–myeloablative transplant recipient pairs (Table 1). The genotype frequencies of 158V/V, 158V/F and 158F/F were 4, 42 and 54% in donors, and 3, 41 and 56% in recipients. These were similar to a previous report<sup>14,31</sup> in Japanese populations and were in accord with the Hardy–Weinberg equilibrium ( $P = 0.91$ ).


*Transplant outcome according to the FCGR3A genotype*

The median follow-up duration in the cohort was 109 months among the survivors (range 43–134 months), and 16 recipients (16%) had relapsed or progressed and 47 (47%) had died. Three patients (3%) died before undergoing engraftment.

The transplant outcomes according to the FCGR3A genotype are summarized in Table 2. The recipient 158V genotype was associated with a significantly reduced incidence of GVHD-related mortality (7 vs 25%,  $P = 0.01$ ; Figure 1b), and a trend toward a reduced incidence of chronic GVHD (33 vs 45%,  $P = 0.07$ ) and reduced 5-year TRM (22 vs 40%,  $P = 0.07$ ; Figure 1a). The donor genotype had no significant effects on the transplant outcomes.

*Multivariate analysis*

All factors that were found to be significant in univariate analyses were included in the model. The presence of the 158V genotype in recipients were statistically significant in the multivariate analyses for better OS (HR 0.49; 95% confidence interval (CI) 0.26–0.93;  $P = 0.03$ ; Table 3) and TRM (HR 0.30; 95% CI 0.14–0.67;  $P = 0.003$ ). In addition, the recipient 158V genotype was associated with a significantly reduced incidence of chronic GVHD (HR 0.45; 95% CI 0.20–0.99;  $P = 0.049$ ) and a trend toward a lower incidence of grade II–IV acute GVHD (HR 0.55; 95% CI 0.27–1.10;  $P = 0.09$ ), resulting in a significantly reduced GVHD-related mortality (HR 0.22; 95% CI 0.06–0.77;  $P = 0.02$ ). A correlation between the recipient 158V genotype and low infection-related death was also observed (HR 0.42; 95% CI 0.17–1.01;  $P = 0.05$ ). The donor 158V genotype did not significantly influence the transplant outcomes.



**Figure 1** Estimated cumulative incidence curves of TRM (a) and GVHD-related mortality (b) according to the recipient FCGR3A genotype. Solid lines and dashed lines represent the 158V-positive recipient and the 158V-negative recipient, respectively.

**Table 3** A multivariate analysis of the association of the FCGR3A genotype with the clinical outcomes after transplantation

Variable	Donor 158V positivity			Recipient 158V positivity			Minor ABO incompatibility		
	HR	95% CI	P-value	HR	95% CI	P-value	HR	95% CI	P-value
OS	1.29	0.70–2.38	0.42	0.49	0.26–0.93	0.03	—	—	—
TRM	1.73	0.84–3.54	0.14	0.30	0.14–0.67	0.003	—	—	—
Relapse	0.82	0.27–2.55	0.74	0.67	0.22–2.02	0.47	—	—	—
II–IV aGVHD	1.24	0.63–2.46	0.53	0.55	0.27–1.10	0.09	—	—	—
cGVHD	0.55	0.25–1.21	0.14	0.45	0.20–0.99	0.049	0.26	0.03–1.88	0.18
GVHD-related mortality	1.65	0.61–4.45	0.32	0.22	0.06–0.77	0.02	—	—	—
Infection-related mortality	1.36	0.58–3.20	0.48	0.42	0.17–1.01	0.05	—	—	—
	High-risk disease			Age (>27 years)			Donor age (>32 years)		
	HR	95% CI	P-value	HR	95% CI	P-value	HR	95% CI	P-value
OS	2.17	1.17–4.04	0.01	—	—	—	—	—	—
TRM	—	—	—	—	—	—	2.56	1.22–5.37	0.01
Relapse	6.44	1.92–21.59	0.003	0.43	0.13–1.43	0.17	—	—	—
II–IV aGVHD	—	—	—	0.38	0.19–0.74	0.005	2.84	1.42–5.69	0.003
cGVHD	—	—	—	—	—	—	—	—	—
GVHD-related mortality	0.43	0.10–1.90	0.27	—	—	—	—	—	—
Infection-related mortality	2.78	1.19–6.46	0.02	—	—	—	3.27	1.31–8.12	0.01

Abbreviations: aGVHD = acute GVHD; CI = confidence interval; cGVHD = chronic GVHD; FCGR3A = Fcγ receptor type IIIA; HR = hazard ratio.

## Discussion

This study showed a considerable effect of the recipient FCGR3A-158V genotype on GVHD development and GVHD-related mortality, thus positively contributing to a significantly better TRM and OS for patients with myeloid malignancies receiving HLA-matched myeloablative BMT. The presence of 158V in recipients did not influence disease relapse. Therefore, recipients with the 158V genotype may be capable of avoiding GVHD without compromising a GVL effect. The recipient 158V genotype also showed a trend toward reduced infection-related mortality, which might result from a reduced need for immunosuppressive therapy due to a low incidence of GVHD, although the data regarding treatment for GVHD were unavailable in this cohort. This is the first report to show that the FCGR3A-V158F polymorphism influences transplant outcomes.

Little is known about the involvement of FCGR3A in the pathogenesis of GVHD. Indirect evidence in animal and human studies showed that B cells have an important

role in the immunopathophysiology of acute and chronic GVHD,<sup>12,32–35</sup> in which FCGR3A is involved through Ab-mediated immune responses, including Ab-dependent cell-mediated cytotoxicity and FCGR3A-mediated endocytosis. Previous reports showing the effectiveness of B-cell depletion in the treatment of acute<sup>33</sup> and chronic GVHD<sup>35</sup> prompted a working hypothesis that the presence of the FCGR3A-158V genotype, which potentially mediates Ab-dependent cell-mediated cytotoxicity more efficiently in comparison to its absence,<sup>16</sup> may be a risk factor contributing to the development of acute and chronic GVHD. However, the current results contradict this hypothesis. One explanation of this conflict may be observed in reports that B-cell-deficient mice experience more exacerbated acute GVHD than wild-type mice,<sup>36</sup> and that a high number of B-cell progenitors in the stem cell graft or in patients after allo-SCT is associated with a significantly lower rate of acute and chronic GVHD.<sup>37,38</sup> A study in HIV-infected men showed the presence of the FCGR3A-158V genotype to be associated with susceptibility to Kaposi's sarcoma and human herpesvirus-8

infection,<sup>39</sup> thus suggesting that the 158V genotype may possess an immunologically recessive nature in immunocompromised patients similar to transplant recipients. However, precisely how the recipient FCGR3A polymorphism influences the pathophysiology of GVHD is still unknown. The first assumption is that in the light of the pathogenesis of GVHD,<sup>40,41</sup> differential binding of IgG to FCGR3A on recipient DCs might alter downstream events, such as the release of cytokines or chemokines, which consecutively could influence pathway essential for the development of GVHD. The second assumption is that the expression of FCGR3A on keratinocytes<sup>42</sup> may have a role in neutralizing the autoantibodies contributing to GVHD.

An alternative explanation is that the association between the recipient FCGR3A polymorphism and the transplant outcomes could develop from polymorphisms in other genes in linkage disequilibrium with the FCGR3A gene. One possible candidate is the Fcγ receptor IIA (FCGR2A) H131R polymorphism shown to be in linkage disequilibrium with the FCGR3A polymorphism in the Caucasian populations.<sup>31,43</sup> However, the absence of a linkage disequilibrium between FCGR3A and FCGR2A genes in Japanese population<sup>31</sup> suggests that the FCGR3A polymorphism is more likely responsible for the transplant outcomes.

The current data are not consistent with a previous French study<sup>44</sup> that found no significant association between the FCGR3A-V158F SNP polymorphism and transplant outcomes. One of the reasons for this discrepancy may be that the population in the initial study included only HLA-identical sibling BMT recipients and was not stratified into myeloid and lymphoid malignancies to examine the effect of gene polymorphism on transplant outcomes. There is a possibility that the influence of the FCGR3A polymorphism is restricted exclusively to unrelated BMT and myeloid malignancies. Alternatively, these conflicting findings may result from ethnic differences in the study population. Verification of the present data in other cohorts has still to be made so that the implications of the findings can be fully accepted.

This study suggests that the genotyping of FCGR3A in transplant recipients before transplantation may provide a recipient bearing the 158F/F genotype an opportunity to avoid the risk of GVHD by favoring a BM or cord blood, and an HLA-matched graft, and planning more immunosuppressive regimens. Further studies are required to ascertain whether the findings of this study can be extended to other disease groups or other stem sources or HLA-mismatched transplantation.

#### Conflict of interest

The authors declare no conflict of interest.

#### Acknowledgements

We are indebted to Drs Hiroko Oshima, Masanobu Oshima and Atsushi Hirao, Ms Kayoko Yamada, Mayu Yamada and Yuki Motohashi at Kanazawa University, and Dr Keitaro Matsuo at

Aichi Cancer Center Research Institute for their technical assistance. We thank all the Japan Marrow Donor Program (JMDF) transplant teams who have contributed patients and donors to this study. This study was supported by grants from the Ministry of Health, Labor and Welfare, and the Ministry of Education, Culture, Sports and Technology, and Funds from the Mitani Research and Development Assistance Organization (Kanazawa, Japan) and by the Japan Leukemia Research Fund (Tokyo, Japan).

#### References

- 1 Gratwohl A, Brand R, Frassoni F, Rocha V, Niederwieser D, Reusser P *et al*. Cause of death after allogeneic haematopoietic stem cell transplantation (HSCT) in early leukaemias: an EBMT analysis of lethal infectious complications and changes over calendar time. *Bone Marrow Transplant* 2005; **36**: 757–769.
- 2 Dickinson AM, Middleton PG, Rocha V, Gluckman E, Holler E. Genetic polymorphisms predicting the outcome of bone marrow transplants. *Br J Haematol* 2004; **127**: 479–490.
- 3 Elmaagacli AH, Koldehoff M, Landt O, Beelen DW. Relation of an interleukin-23 receptor gene polymorphism to graft-versus-host disease after hematopoietic-cell transplantation. *Bone Marrow Transplant* 2008; **41**: 821–826.
- 4 Gerbitz A, Hillemanns P, Schmid C, Wilke A, Jayaraman R, Kolb HJ *et al*. Influence of polymorphism within the heme oxygenase-I promoter on overall survival and transplantation-related mortality after allogeneic stem cell transplantation. *Biol Blood Marrow Transplant* 2008; **14**: 1180–1189.
- 5 Kim DH, Jung HD, Lee NY, Sohn SK. Single nucleotide polymorphism of CC chemokine ligand 5 promoter gene in recipients may predict the risk of chronic graft-versus-host disease and its severity after allogeneic transplantation. *Transplantation* 2007; **84**: 917–925.
- 6 Noori-Daloi MR, Rashidi-Nezhad A, Izadi P, Hossein-Nezhad A, Sobhani M, Derakhshandeh-Peykar P *et al*. Transforming growth factor-beta1 codon 10 polymorphism is associated with acute GVHD after allogeneic BMT in Iranian population. *Ann Transplant* 2007; **12**: 5–10.
- 7 Viel DO, Tsuneto LT, Sossai CR, Lieber SR, Marques SB, Vigorito AC *et al*. IL2 and TNFA gene polymorphisms and the risk of graft-versus-host disease after allogeneic haematopoietic stem cell transplantation. *Scand J Immunol* 2007; **66**: 703–710.
- 8 Sugimoto K, Murata M, Onizuka M, Inamoto Y, Terakura S, Kuwatsuka Y *et al*. Decreased risk of acute graft-versus-host disease following allogeneic hematopoietic stem cell transplantation in patients with the 5,10-methylenetetrahydrofolate reductase 677TT genotype. *Int J Hematol* 2008; **87**: 451–458.
- 9 Espinoza JL, Takami A, Onizuka M, Sao H, Akiyama H, Miyamura K *et al*. NKG2D gene polymorphism has a significant impact on transplant outcomes after HLA-fully-matched unrelated bone marrow transplantation for standard risk hematologic malignancies. *Haematologica* 2009; **94**: 1427–1434.
- 10 Nimmerjahn F, Ravetch JV. Fcγ receptors: old friends and new family members. *Immunity* 2006; **24**: 19–28.
- 11 Nimmerjahn F, Ravetch JV. Fcγ receptors as regulators of immune responses. *Nat Rev Immunol* 2008; **8**: 34–47.
- 12 Shimabukuro-Vornhagen A, Hallek MJ, Storb RF, von Bergwelt-Baildon MS. The role of B cells in the pathogenesis of graft-versus-host disease. *Blood* 2009; **114**: 4919–4927: blood-2008-10-161638.
- 13 Cauza K, Grassauer A, Hinterhuber G, Horvat R, Rappersberger K, Wolff K *et al*. FcγRIII expression on cultured

- human keratinocytes and upregulation by interferon-gamma. *J Invest Dermatol* 2002; **119**: 1074–1079.
- 14 Leppers-van de Straat FG, van der Pol WL, Jansen MD, Sugita N, Yoshie H, Kobayashi T et al. A novel PCR-based method for direct Fc gamma receptor IIIa (CD16) allotyping. *J Immunol Methods* 2000; **242**: 127–132.
  - 15 Koene HR, Kleijer M, Algra J, Roos D, von dem Borne AE, de Haas M. Fc gammaRIIIa-158V/F polymorphism influences the binding of IgG by natural killer cell Fc gammaRIIIa, independently of the Fc gammaRIIIa-48L/R/H phenotype. *Blood* 1997; **90**: 1109–1114.
  - 16 Wu J, Edberg JC, Redecha PB, Bansal V, Guyre PM, Coleman K et al. A novel polymorphism of Fc gammaRIIIa (CD16) alters receptor function and predisposes to autoimmune disease. *J Clin Invest* 1997; **100**: 1059–1070.
  - 17 Tutuncu Z, Kavanaugh A, Zvaifler N, Corr M, Deutsch R, Boyle D. Fc gamma receptor type IIIA polymorphisms influence treatment outcomes in patients with inflammatory arthritis treated with tumor necrosis factor alpha-blocking agents. *Arthritis Rheum* 2005; **52**: 2693–2696.
  - 18 Foster CB, Zhu S, Erichsen HC, Lehrnbecher T, Hart ES, Choi E et al. Polymorphisms in inflammatory cytokines and Fc gamma receptors in childhood chronic immune thrombocytopenic purpura: a pilot study. *Br J Haematol* 2001; **113**: 596–599.
  - 19 Cartron G, Dacheux L, Salles G, Solal-Celigny P, Bardos P, Colombat P et al. Therapeutic activity of humanized anti-CD20 monoclonal antibody and polymorphism in IgG Fc receptor Fc gammaRIIIa gene. *Blood* 2002; **99**: 754–758.
  - 20 Weng WK, Levy R. Two immunoglobulin G fragment C receptor polymorphisms independently predict response to rituximab in patients with follicular lymphoma. *J Clin Oncol* 2003; **21**: 3940–3947.
  - 21 Sugita N, Yamamoto K, Kobayashi T, Van Der Pol W, Horigome T, Yoshie H et al. Relevance of Fc gamma RIIIa-158V-F polymorphism to recurrence of adult periodontitis in Japanese patients. *Clin Exp Immunol* 1999; **117**: 350–354.
  - 22 Dijkstra HM, Bijl M, Fijnheer R, Scheepers RH, Oost WW, Jansen MD et al. Fc gamma receptor polymorphisms in systemic lupus erythematosus: association with disease and *in vivo* clearance of immune complexes. *Arthritis Rheum* 2000; **43**: 2793–2800.
  - 23 Zhang W, Gordon M, Schultheis AM, Yang DY, Nagashima F, Azuma M et al. FCGR2A and FCGR3A polymorphisms associated with clinical outcome of epidermal growth factor receptor expressing metastatic colorectal cancer patients treated with single-agent cetuximab. *J Clin Oncol* 2007; **25**: 3712–3718.
  - 24 Kawase T, Morishima Y, Matsuo K, Kashiwase K, Inoko H, Saji H et al. High-risk HLA allele mismatch combinations responsible for severe acute graft-versus-host disease and implication for its molecular mechanism. *Blood* 2007; **110**: 2235–2241.
  - 25 Sasazuki T, Juji T, Morishima Y, Kinukawa N, Kashiwabara H, Inoko H et al. Effect of matching of class I HLA alleles on clinical outcome after transplantation of hematopoietic stem cells from an unrelated donor. Japan Marrow Donor Program. *N Engl J Med* 1998; **339**: 1177–1185.
  - 26 Livak KJ. Allelic discrimination using fluorogenic probes and the 5' nuclease assay. *Genet Anal* 1999; **14**: 143–149.
  - 27 Przepiora D, Weisdorf D, Martin P, Klingemann HG, Beatty P, Hows J et al. 1994 Consensus conference on acute GVHD grading. *Bone Marrow Transplant* 1995; **15**: 825–828.
  - 28 Shulman HM, Sullivan KM, Weiden PL, McDonald GB, Striker GE, Sale GE et al. Chronic graft-versus-host syndrome in man. A long-term clinicopathologic study of 20 Seattle patients. *Am J Med* 1980; **69**: 204–217.
  - 29 Scrucca L, Santucci A, Aversa F. Competing risk analysis using R: an easy guide for clinicians. *Bone Marrow Transplant* 2007; **40**: 381–387.
  - 30 Gooley TA, Leisenring W, Crowley J, Storer BE. Estimation of failure probabilities in the presence of competing risks: new representations of old estimators. *Stat Med* 1999; **18**: 695–706.
  - 31 van der Pol WL, Jansen MD, Sluiter WJ, van de Sluis B, Leppers-van de Straat FG, Kobayashi T et al. Evidence for non-random distribution of Fc gamma receptor genotype combinations. *Immunogenetics* 2003; **55**: 240–246.
  - 32 Schultz KR, Paquet J, Bader S, HayGlass KT. Requirement for B cells in T cell priming to minor histocompatibility antigens and development of graft-versus-host disease. *Bone Marrow Transplant* 1995; **16**: 289–295.
  - 33 Kamble R, Oholendt M, Carrum G. Rituximab responsive refractory acute graft-versus-host disease. *Biol Blood Marrow Transplant* 2006; **12**: 1201–1202.
  - 34 Zhang C, Todorov I, Zhang Z, Liu Y, Kandeel F, Forman S et al. Donor CD4+ T and B cells in transplants induce chronic graft-versus-host disease with autoimmune manifestations. *Blood* 2006; **107**: 2993–3001.
  - 35 Cutler C, Miklos D, Kim HT, Treister N, Woo SB, Bienfang D et al. Rituximab for steroid-refractory chronic graft-versus-host disease. *Blood* 2006; **108**: 756–762.
  - 36 Rowe V, Banovic T, MacDonald KP, Kuns R, Don AL, Morris ES et al. Host B cells produce IL-10 following TBI and attenuate acute GVHD after allogeneic bone marrow transplantation. *Blood* 2006; **108**: 2485–2492.
  - 37 Storek J, Wells D, Dawson MA, Storer B, Maloney DG. Factors influencing B lymphopoiesis after allogeneic hematopoietic cell transplantation. *Blood* 2001; **98**: 489–491.
  - 38 Michonneau D, Peffault de Latour R, Porcher R, Robin M, Benbunan M, Rocha V et al. Influence of bone marrow graft B lymphocyte subsets on outcome after HLA-identical sibling transplants. *Br J Haematol* 2009; **145**: 107–114.
  - 39 Lehrnbecher TL, Foster CB, Zhu S, Venzon D, Steinberg SM, Wyvill K et al. Variant genotypes of Fc gamma RIIIa influence the development of Kaposi's sarcoma in HIV-infected men. *Blood* 2000; **95**: 2386–2390.
  - 40 Ferrara JL, Cooke KR, Teshima T. The pathophysiology of acute graft-versus-host disease. *Int J Hematol* 2003; **78**: 181–187.
  - 41 Teshima T, Ordemann R, Reddy P, Gagin S, Liu C, Cooke KR et al. Acute graft-versus-host disease does not require alloantigen expression on host epithelium. *Nat Med* 2002; **8**: 575–581.
  - 42 Cauza K, Grassauer A, Hinterhuber G, Horvat R, Rappersberger K, Wolff K et al. Fc[gamma]RIII expression on cultured human keratinocytes and upregulation by interferon-[gamma]. *J Invest Dermatol* 2002; **119**: 1074–1079.
  - 43 Lejeune J, Thibault G, Ternant D, Cartron G, Watier H, Ohresser M. Evidence for linkage disequilibrium between Fc gamma RIIIa-V158F and Fc gamma RIIa-H131R polymorphisms in white patients, and for an Fc gamma RIIIa-restricted influence on the response to therapeutic antibodies. *J Clin Oncol* 2008; **26**: 5489–5491.
  - 44 Rocha V, Franco RF, Porcher R, Bittencourt H, Silva Jr WA, Latouche A et al. Host defense and inflammatory gene polymorphisms are associated with outcomes after HLA-identical sibling bone marrow transplantation. *Blood* 2002; **100**: 3908–3918.



# blood

2010 115: 4664-4670  
Prepublished online Mar 24, 2010;  
doi:10.1182/blood-2009-10-251157

## Impact of highly conserved HLA haplotype on acute graft-versus-host disease

Satoko Morishima, Seishi Ogawa, Aiko Matsubara, Takakazu Kawase, Yasuhito Nannya, Koichi Kashiwase, Masahiro Satake, Hiroo Saji, Hidetoshi Inoko, Shunichi Kato, Yoshihisa Koderu, Takehiko Sasazuki, Yasuo Morishima and for the Japan Marrow Donor Program

---

Updated information and services can be found at:

<http://bloodjournal.hematologylibrary.org/cgi/content/full/115/23/4664>

Articles on similar topics may be found in the following *Blood* collections:

- Transplantation (1555 articles)
- Free Research Articles (1020 articles)
- Clinical Trials and Observations (2999 articles)

---

Information about reproducing this article in parts or in its entirety may be found online at:

[http://bloodjournal.hematologylibrary.org/misc/rights.dtl#repub\\_requests](http://bloodjournal.hematologylibrary.org/misc/rights.dtl#repub_requests)

Information about ordering reprints may be found online at:

<http://bloodjournal.hematologylibrary.org/misc/rights.dtl#reprints>

Information about subscriptions and ASH membership may be found online at:

<http://bloodjournal.hematologylibrary.org/subscriptions/index.dtl>

Blood (print ISSN 0006-4971, online ISSN 1528-0020), is published semimonthly by the American Society of Hematology, 1900 M St, NW, Suite 200, Washington DC 20036.  
Copyright 2007 by The American Society of Hematology; all rights reserved.



## Impact of highly conserved HLA haplotype on acute graft-versus-host disease

\*Satoko Morishima,<sup>1</sup> \*Seishi Ogawa,<sup>2</sup> Aiko Matsubara,<sup>2</sup> Takakazu Kawase,<sup>3</sup> Yasuhito Nannya,<sup>2</sup> Koichi Kashiwase,<sup>4</sup> Masahiro Satake,<sup>4</sup> Hiroo Saji,<sup>5</sup> Hidetoshi Inoko,<sup>6</sup> Shunichi Kato,<sup>7</sup> Yoshihisa Kodera,<sup>8</sup> Takehiko Sasazuki,<sup>9</sup> and Yasuo Morishima,<sup>1</sup> for the Japan Marrow Donor Program

<sup>1</sup>Department of Hematology and Cell Therapy, Aichi Cancer Center Hospital, Nagoya; <sup>2</sup>21st Century COE Program, Graduate School of Medicine, University of Tokyo, Tokyo; <sup>3</sup>Division of Epidemiology and Prevention, Aichi Cancer Center Research Institute, Nagoya; <sup>4</sup>Tokyo Metropolitan Red Cross Blood Center, Tokyo; <sup>5</sup>HLA Laboratory, NPC, Kyoto; <sup>6</sup>Division of Molecular Science, Tokai University School of Medicine, Isehara; <sup>7</sup>Department of Cell Transplantation and Regenerative Medicine, Tokai University School of Medicine, Isehara; <sup>8</sup>Aichi Medical University, Aichi; and <sup>9</sup>International Medical Center of Japan, Tokyo, Japan

Although the effects of human leukocyte antigen (HLA) locus matching on clinical outcome in unrelated hematopoietic stem cell transplantations have been characterized, the biologic implications of HLA haplotypes have not been defined. We demonstrated the genetic fixity of Japanese conserved extended haplotypes by multi-single nucleotide polymorphism analysis in 1810 Japanese donor-recipient pairs matching with HLA-A, -B, -C, -DRB1, and -DQB1 alleles. Three major Japanese con-

served extended haplotypes (named HP-P1, HP-P2, and HP-P3) were essentially completely conserved at least in the 3.3-Mb HLA region from HLA-A to -DPB1, and extended far beyond HLA-A. The risk of acute graft-versus-host disease (GVHD) of these HLA haplotypes was assessed with multivariate Cox regression in 712 patients transplanted from HLA fully (HLA-A, B, C, DRB1, DQB1, and DPB1) matched unrelated donors. HP-P2 itself reduced the risk of grade 2 to 4 acute GVHD (hazard ratio

[HR] = 0.63;  $P = .032$  compared with HP-P2-negative), whereas HP-P3 tended to increase the risk (HR = 1.38;  $P = .07$ ). Among 381 patients with HP-P1, HP-P1/P3 (HR = 3.35;  $P = .024$ ) significantly increased the risk of acute GVHD compared with homozygous HP-P1. This study is the first to demonstrate that a genetic difference derived from HLA haplotype itself is associated with acute GVHD in allogeneic hematopoietic stem cell transplantation. (*Blood*. 2010;115(23):4664-4670)

### Introduction

Allogeneic hematopoietic stem cell transplantation (HSCT) from a human leukocyte antigen (HLA)-matched unrelated (UR) donor has been established as a mode of curative therapy for hematologic malignancies and other hematologic or immunologic disorders, when an HLA-identical sibling donor is unavailable. Although the effect of donor and recipient HLA locus matching on the clinical outcome of UR-HSCT has been well elucidated,<sup>1-4</sup> the biologic implications of HLA haplotype itself have not been explored for HSCT.

HLA-identical sibling shares 2 identical major histocompatibility complex (MHC) haplotypes by descent, including non-HLA polymorphic genes, and it has been generally accepted that transplantation between these related pairs provides a superior outcome. On the other hand, there is no guarantee of matching for non-HLA genes between HLA-allele matched UR donor and recipient pairs, and mismatching of haplotype block in MHC has been suggested to lead to severe acute GVHD<sup>5</sup> and an inferior outcome<sup>6,7</sup> in UR-HSCT.

In human population, multiple DNA blocks in the MHC are strongly associated with each other, and these relatively long stretches of conserved DNA sequence in the MHC have been named conserved extended haplotypes (CEHs) or ancestral haplotypes.<sup>8,9</sup> CEHs are often population-specific and have been investigated as markers for disease susceptibility, particularly in autoim-

mune diseases.<sup>10</sup> However, the relation between CEHs and clinical outcome of UR-HSCT has not been yet reported.

Using the large-scale Japan Marrow Donor Program (JMDP) data, we evaluated the conservation of common HLA haplotypes among a Japanese population and elucidated its impact on acute graft-versus-host disease (GVHD) and other clinical outcomes in UR-HSCT.

### Methods

#### Study population

A total of 5210 donor-recipient pairs who underwent transplantation through the JMDP between January 1993 and January 2006 were retrospectively genotyped for HLA-A, -B, -C, -DRB1, -DQB1, and -DPB1 alleles as described elsewhere.<sup>4</sup> For the genome-wide association studies, 1810 pairs (3620 persons) who matched HLA-A, -B, -C, -DRB1, and -DQB1 alleles and were available for DNA sample were selected from these 5210 pairs. For the analysis of acute GVHD, 712 patients who had received T cell-replete bone marrow from an HLA-A, -B, -C, -DRB1, -DQB1, and -DPB1 allele-matched donor were selected from these 5210 pairs. The characteristics of these 712 patients are shown in Table 1. A final clinical survey of the patients was completed by June 2007. Informed consent was obtained from patients and donors in accordance with the Declaration of Helsinki, and approval of the study was obtained from the Institutional Review Board of Aichi Cancer Center and JMDP.

Submitted October 26, 2009; accepted March 10, 2010. Prepublished online as *Blood* First Edition paper, March 24, 2010; DOI 10.1182/blood-2009-10-251157.

\*S.M. and S.O. contributed equally to this paper.

The online version of this article contains a data supplement.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 USC section 1734.

© 2010 by The American Society of Hematology

**Table 1. Clinical characteristics of patients according to HLA haplotype**

Characteristic	Haplotype P1		Haplotype P2		Haplotype P3	
	Negative	Positive	Negative	Positive	Negative	Positive
No. of patients	381	331	601	111	608	104
Median patient age, y (range)	33 (0-70)	33 (1-68)	33 (0-70)	35 (1-65)	33 (1-68)	36 (0-70)
<b>Sex (donor/ patient)</b>						
Male/male	160	152	250	62	274	38
Male/female	71	63	117	17	112	22
Female/male	66	49	103	12	100	15
Female/female	84	67	131	20	122	29
<b>Disease</b>						
ALL	91	71	138	24	144	18
ANLL	90	106	167	29	166	30
CML	65	54	95	24	101	18
Hereditary disease	5	9	11	3	12	2
MDS	52	39	76	15	78	13
Malignant lymphoma	39	30	61	8	52	17
Multiple myeloma	4	4	7	1	7	1
Severe aplastic anemia	24	9	28	5	31	2
Other	11	9	18	2	17	3
<b>Risk of leukemia relapse*</b>						
Standard	137	112	211	38	214	35
High	109	119	189	39	197	31
Disease other than leukemia	135	100	201	34	197	38
<b>GVHD prophylaxis</b>						
Cyclosporine-based	199	203	345	57	339	63
Tacrolimus-based	182	128	256	54	269	41
<b>ATG</b>						
ATG	25	23	38	10	40	8
Non-ATG	356	308	563	101	568	96
<b>Preconditioning</b>						
TBI regimen	298	241	456	83	463	76
Non-TBI regimen	83	90	145	28	145	28

HLA indicates human leukocyte antigen; ALL, acute lymphoblastic leukemia; ANLL, acute nonlymphoblastic leukemia; CML, chronic myeloid leukemia; MDS, myelodysplastic syndrome; GVHD, graft-versus-host disease; ATG, anti-human thymocyte globulin; and TBI, total body irradiation.

\*Standard risk for leukemia relapse was defined as the status of the first complete remission of ALL and ANLL and the first chronic phase of CML at transplantation. High risk was defined as a more advanced status than standard risk in AML, ANLL, and CML. Disease other than leukemia was defined as other than ALL, ANLL, and CML.

### SNP typing and HLA haplotype analysis

The single nucleotide polymorphism (SNP) array experiments were performed according to the standard protocol of Affymetrix GeneChip Mapping 500K Array (Affymetrix). After excluding those SNPs showing less than 95% call rate and deviation from Hardy-Weinberg equilibrium ( $P < .001$ ), 10.8% of SNPs for the HLA region failed. And 4761 SNPs in the region spanning the MHC (20-46 Mb from the telomere in chromosome 6p) were analyzed to evaluate the conservation of common HLA haplotypes.

Persons who were homozygous for HLA-A, -B, -C, -DRB1, -DQB1, and -DPB1 were isolated, and the 3 major HLA haplotypes were named HP-P1, HP-P2, and HP-P3. The homozygosity of consecutive SNPs of these HLA haplotypes was analyzed to assess the region of conservation. SNP alleles of the extended homozygous region in each HLA haplotype were analyzed to determine allele frequencies, and a consensus sequence of major alleles in each haplotype was established. Then, the SNP sequence of persons who carried at least one copy of HLA haplotype (shared the same HLA alleles as common HLA haplotype at the HLA-A, -B, -C, -DRB1, -DQB1, and -DPB1 loci) was compared with this consensus sequence. Missing alleles in a sample were not accounted for in this analysis.

### Statistical analysis

Cumulative incidences of acute GVHD and relapse were assessed by the method described elsewhere to eliminate the effect of competing risk.<sup>4</sup> Overall survival was calculated using the Kaplan-Meier method. The competing event regarding acute GVHD was defined as death without acute GVHD. A log-rank test was applied to assess the impact by the factor of interest. Multivariable Cox regression analyses were conducted to evaluate the impact of the specific haplotype on acute GVHD, leukemia relapse, and


mortality after transplantation. Confounders considered were sex (donor-recipient pair), patient age (linear), donor age (linear), transplantation year, type of disease, risk of leukemia relapse (standard, high, and diseases other than leukemia), GVHD prophylaxis (cyclosporine-based regimen vs tacrolimus-based regimen), anti-thymocyte globulin (anti-thymocyte globulin vs no anti-thymocyte globulin), and preconditioning (total body irradiation vs non-total body irradiation).

## Results

### Highly conserved common HLA haplotypes among Japanese

To evaluate for conservation of Japanese common HLA haplotypes, persons who were homozygous HLA haplotype (having homozygous alleles in HLA-A, -B, -C, -DRB1, -DQB1, and -DPB1 loci) were selected from a total 3620 persons (1810 donor-recipient pairs) for whom genome-wide association study was performed. Among those, 72 persons were homozygous HLA-A\*2402 -Cw\*1202 -B\*5201 -DRB1\*1502 -DQB1\*0601 -DPB1\*0901 (named HP-P1), 10 persons were homozygous HLA-A\*3303 -Cw\*1403 -B\*4403 -DRB1\*1302 -DQB1\*0604 -DPB1\*0401 (named HP-P2), and 8 persons were homozygous HLA-A\*2402 -Cw\*0702 -B\*0702 -DRB1\*0101 -DQB1\*0501 -DPB1\*0402 (named HP-P3).


Homozygosity at consecutive SNP loci of persons with homozygous HLA haplotype was shown in Figure 1. The extended



**Figure 1. Representation of genotypes in persons with homozygous HLA haplotype.** Data from chromosome 6p (24-37 Mb) of persons with homozygous HLA haplotype are shown. Each column indicates 1 person (72 persons with homozygous HP-P1, 10 persons with homozygous HP-P2, and 8 persons with homozygous HP-P3). Each of the 2389 evenly spaced rows represents 1 SNP locus. Blue row represents homozygous genotype; and red row, heterozygous genotype. Missing genotypes were not counted.

homozygous region of HP-P1 gradually broke up the region of HLA-A. The longest homozygous region in persons with HP-P1 was 18.7 Mb. Of 72 persons with homozygous HP-P1, 32 persons (nos. 1-32 of HP-P1 in Figure 1) had more than 99.0% of 1395 consecutive homozygous SNPs throughout the 6.9-Mb region from rs806971 to rs6937061 (nucleotides 26252770-33187790). Although haplotypes of all those 32 persons were identical centromeric rs9257745 (nucleotide 29414635), the telomeric region was clearly divided into 2 different haplotypes (Figure 2). A total of 26 of 32 persons had one of the homozygous haplotypes (named subtype A of HP-P1), and the remaining 6 persons (nos. 3, 9, 19, 23, 24, and 25 of HP-P1 in Figure 1) had another homozygous haplotype (subtype B of HP-P1). A total of 65 of 72 persons with homozygous HP-P1 (nos. 1-65 of HP-P1 in Figure 1) had more than 99.0% homozygous alleles for 888 consecutive SNPs throughout the 3.3-Mb region from rs1610630 to rs6937061 (nucleotides 29837265-33187790). Seven other persons (nos. 66-72 in Figure 1) had an apparently lower conserved region, with 1.0% to 10.0% heterozygous alleles within the 3.3-Mb region.

All 10 persons with homozygous HP-P2 had more than 99.0% homozygous alleles for consecutive SNPs throughout the 3.3-Mb region. Furthermore, 9 of 10 persons with homozygous HP-P2 showed homozygosity extending across the 7.7-Mb region from rs6912426 to rs6937061 (nucleotides 25517764-33187790), and those 9 persons had identical genotypes in almost all the 1540 consecutive SNPs throughout the 7.7-Mb region.



**Figure 2. Subtypes of HLA haplotype P1.** Data from chromosome 6p (from nucleotides 26252770-33187790) of 32 persons with consecutive homozygous SNPs throughout the 6.9-Mb region. The SNP sequence of persons was compared with consensus sequence across the 6.9-Mb region. Each column indicates 1 person. Each of 1395 evenly spaced rows represents 1 SNP locus. Gray row represents SNPs identical to the consensus alleles; and red row, SNPs different from the consensus alleles. Missing SNPs were not counted. A total of 26 of 32 persons had alleles identical to consensus alleles across 6.9 Mb (subtype A of HP-P1), whereas the remaining 6 persons had apparently different alleles in the telomeric region from nucleotide 29414635 (subtype B of HP-P1). These data indicated that the telomeric region of HP-P1 was clearly divided into 2 different haplotypes.

Among 8 persons with homozygous HP-P3, 5 persons had more than 99.0% homozygous alleles throughout the 3.3-Mb region, and the other 3 persons had 1.7% to 8.0% heterozygous alleles within that region. One person with homozygous HP-P3 showed an extraordinary long stretch of homozygosity across the 25.4-Mb region (nucleotides 20162518-45595922).


Consensus sequence of each haplotype was determined using analysis of persons who were homozygous for almost all the SNPs, ie, 6.9-Mb region in HP-P1 (subtypes A and B), 7.7-Mb region in HP-P2, and 3.3-Mb region HP-P3. The person with the longest homozygous telomeric region of HLA-A served to determine a further extended haplotype (Table 2). These persons had alleles identical to the consensus sequence of major alleles in each haplotype described above in this section.

We ascertained whether the consensus sequence of each HLA haplotype was present in the persons carrying at least one copy of HLA haplotype, that is, sharing the same HLA-A, -B, -C, -DRB1, -DQB1, and -DPB1 alleles of common HLA haplotypes (Figure 3; supplemental Figure 1, available on the *Blood* Web site; see the Supplemental Materials link at the top of the online article). Among 3620 persons analyzed for multi-SNP, 1000 of 1045 persons (95%) with HP-P1 had identical alleles for more than 99.5% of consecutive 888 SNPs as a consensus sequence across 3.3-Mb region from 181 kb telomeric HLA-A to 25 kb centromeric HLA-DPB1.

**Table 2. Longest homozygous region of common HLA haplotype**

Type	Start of homozygous region		End of homozygous region		Region length, Mb
	SNP	Position, kb	SNP	Position, kb	
<b>HP-P1</b>					
Subtype A	rs199026	23441.813	rs2395801	42231.646	18.8
Subtype B	rs573863	24080.365	rs1536501	33835.863	9.7
HP-P2	rs1175427	24387.055	rs1873254	34134.467	9.7
HP-P3	rs1688325	20162.518	rs6905847	45595.922	25.4





**Figure 3. Conservation of common HLA haplotypes.** The SNP sequence of persons who carried at least 1 copy of HLA haplotype (shared the same HLA alleles as common HLA haplotype) was compared with consensus sequence of common HLA haplotypes, and conserved regions in each HLA haplotype were illustrated schematically. The majority of persons who share the same HLA alleles in HLA-A, -B, -C, -DRB1, -DQB1, and -DPB1 as common HLA haplotypes possess at least a 3.3-Mb conserved region from HLA-A to DPB1. HP-P1 splits into the A and B subtypes. The length of conservation in HP-P1, HP-P2, and HP-P3 is more extensive to the telomeric region of HLA-A. ID and position of SNPs are: \*1 rs199026 (nucleotide 23551813), \*2 rs573863 (nucleotide 24080365), \*3 rs1397843 (nucleotide 24111433), \*4 rs11754278 (nucleotide 24387055), \*5 rs303031 (nucleotide 25144959), \*6 rs806971 (nucleotide 26252770), \*7 rs9257745 (nucleotide 29414635), \*8 rs1610630 (nucleotide 29837265), and \*9 rs6937061 (nucleotide 33187790).

Furthermore, 654 of 1045 (62%) persons with HP-P1 had identical alleles for more than 99.5% of consecutive 1395 SNPs as subtype A of HP-P1 across the 6.9-Mb region, and 231 persons (22%) had identical alleles as subtype B of HP-P1 across the 6.9-Mb region. Fewer persons showed the conserved region extending up to 9.0 Mb. Among 385 persons with HP-P2, 378 (98%) had identical alleles for 888 consecutive SNPs across the 3.3-Mb region. Furthermore, 305 (79%) had identical alleles for more than 99.5% of 1395 consecutive SNPs as a consensus sequence across the 6.9-Mb region, and 205 (53%) also did across the 8.0-Mb region (nucleotides 24111433-33187790). Among 438 persons with HP-P3, 381 (87%) had identical alleles across the 3.3-Mb region, and 257 (59%) had identical alleles for consecutive SNPs as a consensus sequence of HP-P3 across the 6.9-Mb region.

These results indicate that most of the persons with a common HLA haplotype had a conserved region at least 3.3 Mb from HLA-A to HLA-DPB1. Furthermore, a considerable number of unrelated persons with common HLA haplotype had a more extended conserved telomeric region of HLA-A.

**Effect of HLA haplotype on acute GVHD**

To elucidate the effect of specific HLA haplotype on acute GVHD, we analyzed 712 patients who underwent transplantation from HLA fully matched (12 of 12 HLA alleles) donor with T cell-replete marrow (Table 1). We excluded HLA-mismatched transplantation to avoid obscuring the relationship between HLA haplotype itself and acute GVHD by powerful allogeneic immune responses caused by HLA allele disparities. Among those patients, 331 (46.4%) had HP-P1, 111 (15.0%) had HP-P2, and 104 (14.6%) had HP-P3.

At first, grade 2 to 4 acute GVHD in patients with specific haplotype was compared with those without specific haplotype using multivariate analysis (Table 3). There was no significant difference in the hazard ratio (HR) in grades 2 to 4 acute GVHD between HP-P1-positive and -negative patients, and also no significant difference in the cumulative incidence of acute GVHD between HP-P1-positive and HP-P1-negative patients (31.5% vs 30.7%; Figure 4). Of note, HR of grades 2 to 4 acute GVHD in HP-P2-positive patients was 0.63 (95% confidence interval [CI], 0.41-0.96, *P* = .032) compared with HP-P2-negative patients, and the cumulative incidence of acute GVHD in patients with HP-P2 was significantly lower than HP-P2-negative patients (22.3% vs 33.7%, *P* = .031; Figure 4). On the other hand, a trend of increasing risk of acute GVHD was observed in HP-P3-positive patients (HR = 1.38; 95% CI, 0.97-1.95; *P* = .07). The cumulative incidence of acute GVHD in HP-P3-positive patients was 39.2% and HP-P3-negative patients 29.5% (*P* = .064; Figure 4).


A total of 331 patients with HP-P1 in HLA fully matched transplantation made it possible to elucidate the effect of another HLA haplotype on acute GVHD (supplemental Table 1 for patient characteristics). We did not determine unique haplotypes other than HP-P1, HP-P2, and HP-P3 in this study, so all the persons with HP-P1 and unknown haplotype were lumped together with HP-P1/other. The incidence of grade 2 to 4 acute GVHD in patients with HP-P1/P3 (49.9%) was significantly higher than those with homozygous HP-P1 (16.0%), and there was no significant difference between patients with homozygous HP-P1 and those with HP-P1/P2 (12.0%; Figure 5). Multivariate analyses of HR for acute GVHD showed the same results (Table 3). There was no significant difference in the risk of acute GVHD between patients with homozygous HP-P1 and those with HP-P1/P2 (*P* = .64). Compared with patients with homozygous HP-P1, patients with HP-P1/P3 (HR = 3.35; 95% CI, 1.18-9.55; *P* = .024) had a significantly higher risk of acute GVHD.

As for grade 3 to 4 acute GVHD, there were no significant differences between HP-positive and -negative patients. When 331 patients with HP-P1 in HLA fully match transplantation

**Table 3. Hazard ratio of HLA haplotype on acute GVHD (grade 2-4)**

HLA haplotype	Negative/positive	No.	Hazard ratio (95% CI)	<i>P</i>
P1	Negative	381	Referent 1.00	
P1	Positive	331	1.06 (0.81-1.39)	.665
P2	Negative	601	Referent 1.00	
P2	Positive	111	0.63 (0.41-0.96)	.032
P3	Negative	608	Referent 1.00	
P3	Positive	104	1.38 (0.97-1.95)	.07
P1/P1		36	Referent 1.00	
P1/P2		25	0.71 (0.17-2.93)	.64
P1/P3		19	3.35 (1.18-9.55)	.024
P1/other		251	2.49 (1.06-5.85)	.036

Multivariate analysis adjusted by clinical factors (see Table 2 and supplemental Table 1).



**Figure 4. Cumulative incidence of grade 2 to 4 acute GVHD by common HLA haplotype.** The endpoint was the time to diagnosis of grade 2 to 4 acute GVHD with censoring of date of death until 100 days after transplantation. *P* value was calculated with the log-rank test. (A) Patients with or without HP-P1. (B) Patients with or without HP-P2. (C) Patients with or without HP-P3.

were analyzed, grade 3 to 4 acute GVHD showed the same tendency with grade 2 to 4 acute GVHD. Incidence in patients with homozygous HP-P1 was 2.7%, HP-P1/P2 8.0%, and HP-P1/P3 23.6%.

The cumulative incidence of relapse showed a higher trend in HP-P2-positive patients compared with -negative patients (37.3% vs 29.6%, *P* = .051), and HR was 1.34 (95% CI, 0.94-1.91, *P* = .108). There were no significant differences in the relapse rate between HP-P1-positive and -negative patients, and also between HP-P3 -positive and -negative patients.

Overall survival showed no significant differences between HP-positive and -negative patients. When 331 patients with HP-P1 in HLA fully matched transplantation were analyzed, HR of mortality in HP-P1/P3 was 2.03 (95% CI, 0.92-4.49 *P* = .08) compared with patients with homozygous HP-P1, and HR of HP-P1/P2 1.69 (95% CI, 0.78-3.70, *P* = .186).

## Discussion


First, we demonstrated that Japanese common HLA haplotypes were extraordinarily conserved. Preferential selection of HLA-A-,

-B-, and -DR-matched donor through JMDP made it easy to identify a considerable number of persons with homozygous common HLA haplotype extending HLA-DPB1. HP-P1, HP-P2, and HP-P3 have been previously reported as common HLA haplotypes in the Japanese population.<sup>11-13</sup> The haplotype frequency of HP-P1 was 0.054 to 0.062, that of HP-P2 was 0.029 to 0.036, and that of HP-P3 was 0.016 to 0.040.

CEHs have been mainly identified by blocks of fragment in the MHC region, such as complement genes, alleles of HLA class I/II gene, and tumor necrosis factor- $\alpha$  gene. Recently, high-density SNP analysis in the HLA region made it possible to confirm the genetic fixity of HLA haplotype.<sup>14-16</sup>

We determined the consensus sequence of these HLA haplotypes using multi-SNP data of unrelated persons with homozygous HLA haplotype, and the majority of persons who share the same HLA alleles in HLA-A, -B, -C, -DRB1, -DQB1, and -DPB1 as common HLA haplotypes possess at least a 3.3-Mb conserved region from HLA-A to -DPB1. Furthermore, we showed that those Japanese common HLA haplotypes extend far beyond the HLA-A. We also found, for the first time, that HP-P1 was divided into 2 subtypes based on the telomeric region from nucleotide 29414635 (subtypes A and B). The A1-B8-DR3 CEH, which is one of the most frequent haplotypes in northern European populations, showed that the region of conservation extended 6 Mb telomeric to HLA-A.<sup>16,17</sup> Caucasian common HLA haplotypes have often reportedly shown a lack of nonrandom association between HLA-DR, -DQ, and -DP,<sup>18</sup> whereas Japanese common HLA haplotypes have been subdivided into haplotypes with only a very limited number of HLA-DPB1.<sup>11,12</sup> The highly conserved HLA haplotype that extends to HLA-DPB1 might be attributable to ethnic isolation in the Japanese.

For the analysis of comparison between haplotype-positive and -negative patients (Figure 4; Table 3), patients with HP-P2 significantly reduced the risk of acute GVHD. On the other hand, patients with HP-P3 showed a tendency to increase the risk. Although the relapse rate showed a higher trend in patients with HP-P2 compared with HP-P2-negative patients, the grade 3 to 4 acute GVHD and overall survival did not differ between HP-P2-positive and -negative patients, nor between HP-P3-positive and -negative patients. We suspected the differences with weak power might be attributed to the effects of various other haplotypes combined with a particular haplotype. To confirm the differences in the effects on acute GVHD among a particular HLA haplotype, we analyzed the effect of another



**Figure 5. Cumulative incidence of grade 2 to 4 acute GVHD by common HLA haplotype in patients with HP-P1.** The endpoint was the time to the diagnosis of grade 2 to 4 acute GVHD with censoring of date of death until 100 days after transplantation. *P* value was calculated with the log-rank test. A total of 331 patients with HP-P1 were analyzed for the effect of another HLA haplotype by HP-P1, HP-P2, HP-P3, and the other haplotypes, which were lumped together.

haplotype on acute GVHD among patients with HP-P1 (Figure 5). Patients with HP-P1/P3 showed a significantly higher risk of grade 2 to 4 acute GVHD compared with patients with homozygous HP-P1, a tendency to increase the incidence of grade 3 to 4, and also a tendency to decrease overall survival. Patients with homozygous HP-P1 and patients with HP-P1/P2 showed an extremely lower incidence of grade 2 to 4 acute GVHD (16.2% and 12.0%). Therefore, we could not detect any difference between HP-P1 and HP-P2.

Thus, we demonstrated, for the first time, that the HLA haplotype itself affected the occurrence of acute GVHD. These findings suggest that the genetic factor of specific haplotypes would contribute to reducing or increasing the risk of acute GVHD. Alternatively, we should consider the effect of donor and recipient mismatch SNPs because HP-P3 showed more variation than HP-P1 or HP-P2.

There are several possible explanations for the reduced or increased risk of acute GVHD in patients with the specific haplotype. In HSCT, GVHD has been known to result mainly from donor T cells recognizing minor histocompatibility antigens presented on HLA molecules of a recipient's organs.<sup>19</sup> Different HLA haplotypes possess a different combination of HLA alleles. Therefore, various HLA alleles in each haplotype might present different immunodominant peptides to T cells and evoke different alloreactivity in HLA-matched UR-HSCT. Presumably, the critical but as yet unidentified minor histocompatibility antigens linked to major histocompatibility antigen should be explored based on HLA haplotype, such as common gene deletion polymorphisms.<sup>20</sup> Distinct forms of GVHD were found in different MHC haplotypes in mice, and it has been argued that genes in the MHC locus can dominantly determine the forms of GVHD, probably through MHC-based selection of immunodominant antigens.<sup>21</sup>

In human retrospective analysis, several single-center studies have shown a reduced incidence of acute GVHD,<sup>22</sup> reduced relapse rate,<sup>23</sup> and improved overall survival<sup>24</sup> for HLA-DR15-positive patients transplanted from HLA-matched donors. HLA-DR15 has been known to be a marker of disease susceptibility and clinical response to immunosuppressive therapy in autoimmune-mediated bone marrow failure,<sup>25,26</sup> and it is speculated that immune responses specific to HLA-DR15 are induced. However, our analysis of the effect of HLA-DR15 using the same database showed no effect on acute GVHD (data not shown).

HLA haplotype serves as a model system for studies of disease association, especially in autoimmune disease or infection, and several candidate genes in the HLA region associated with specific haplotype have the potential to modulate immune or inflammatory responses.<sup>10</sup> The observed effect of HLA haplotypes on GVHD development could be explained by particular SNPs that are closely associated with those HLA haplotypes. Within the region of conserved HLA haplotypes, there exist several candidate genes whose SNPs may be related to immune responses. Genetic variants of tumor necrosis factor- $\alpha$  gene located in the HLA region might influence the risk of developing GVHD.<sup>27,28</sup> In addition, *TAP1/TAP2* and *LMP2/LMP7* genes encode subunit components of the proteasomes implicated in the processing of class I HLA-bound peptides,<sup>29,30</sup> and polymorphisms of these genes may affect antigen presentation on recipient tissues, leading to different susceptibility to GVHD. However, they probably do not explain the observed effects of haplotypes; correlations of each haplotype with known SNPs in these genes are generally weak, although

*D'* among these alleles is high (> 0.98). Moreover, currently no haplotype-specific non-HLA polymorphisms have been identified in our series, although we could not exclude the possibility that there may exist some nonobserved SNPs that are closely associated with relevant HLA haplotypes.

We showed that Japanese common HLA haplotypes were conserved from HLA-DPB1 to extensively telomeric HLA-A, so it might be possible that the responsible gene is located in the telomeric region of the classic HLA. Interestingly, HP-P1 was divided into 2 subtypes based on the telomeric region from nucleotide 29414635 (subtypes A and B). Although we analyzed the effect of those subtypes on grade 2 to 4 acute GVHD among patients with HP-P1, we could not detect significant differences between patients with HP-P1 subtype A and subtype B. We also could not detect the differences between patients transplanted from a donor with HP-P1 subtype A and subtype B (data not shown).

In conclusion, in the present study, we demonstrated that highly conserved HLA haplotype might contribute to the occurrence of acute GVHD in HSCT. Although the clinical implications of our results should be considered cautiously, these results imply the proof of principle for an association between one or another HLA haplotype and GVHD. Our findings on the conservation of the MHC region may also provide background for exploring not only genetic factors associated with acute GVHD but also genetic disease susceptibility in our population. More extensive studies are warranted to identify specific genes associated with a particular haplotype contributing to acute GVHD.

## Acknowledgments

The authors thank the staff members of the transplantation centers, donor centers, and the Japan Marrow Donor Program office for their generous cooperation.

This work was supported in part by the Research on Allergic Disease and Immunology (Health and Labor Science Research Grant), the Ministry of Health, Labor, and Welfare of Japan (Grant-in-Aid for Cancer Research), and the Core Research for Evolutional Science and Technology of Japan (Grant-in-Aid).

## Authorship

Contribution: S.M., Y.M., S.O., H.S., M.S., H.I., and T.S. participated in the design of this study; S.O., K.K., A.M., and Y.N. performed histocompatibility analysis; Y.M., S.K., and Y.K. organized data collection for transplantation; T.K. performed statistical data analysis; S.M. and Y.M. performed analysis and wrote this paper; and all authors checked the final version of the paper.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

Correspondence: Seishi Ogawa, The 21st Century COE Program, Graduate School of Medicine, University of Tokyo, 7-3-1 Hongo Bunkyo-ku, Tokyo, 113-0033 Japan; e-mail: sogawa-tyk@umin.net; or Yasuo Morishima, Department of Hematology and Cell Therapy, Aichi Cancer Center Hospital, 1-1 Kanokoden, Chikusa-ku, Nagoya 464-8681, Japan; e-mail: ymorisim@aichi-cc.jp.

## References

- Sasazuki T, Juji T, Morishima Y, et al. Effect of matching of class I HLA alleles on clinical outcome after transplantation of hematopoietic stem cells from an unrelated donor: Japan Marrow Donor Program. *N Engl J Med*. 1998;339(17):1177-1185.
- Morishima Y, Sasazuki T, Inoko H, et al. The clinical significance of human leukocyte antigen (HLA) allele compatibility in patients receiving a marrow transplant from serologically HLA-A, HLA-B, and HLA-DR matched unrelated donors. *Blood*. 2002;99(11):4200-4206.
- Lee SJ, Klein J, Haagenson M, et al. High-resolution donor-recipient HLA matching contributes to the success of unrelated donor marrow transplantation. *Blood*. 2007;110(13):4576-4583.
- Kawase T, Morishima Y, Matsuo K, et al. High-risk HLA allele mismatch combinations responsible for severe acute graft-versus-host disease and implication for its molecular mechanism. *Blood*. 2007;110(7):2235-2241.
- Petersdorf EW, Malkki M, Gooley TA, Martin PJ, Guo Z. MHC haplotype matching for unrelated hematopoietic cell transplantation. *PLoS Med*. 2007;4(1):e8.
- Tay GK, Witt CS, Christiansen FT, et al. Matching for MHC haplotypes results in improved survival following unrelated bone marrow transplantation. *Bone Marrow Transplant*. 1995;15(3):381-385.
- Kitcharoen K, Witt CS, Romphruk AV, Christiansen FT, Leelayuwat C. MICA, MICB, and MHC beta block matching in bone marrow transplantation: relevance to transplantation outcome. *Hum Immunol*. 2006;67(3):238-246.
- Alper CA, Awdeh Z, Yunis EJ. Conserved, extended MHC haplotypes. *Exp Clin Immunogenet*. 1992;9(2):58-71.
- Degli-Esposti MA, Leaver AL, Christiansen FT, Witt CS, Abraham LJ, Dawkins RL. Ancestral haplotypes: conserved population MHC haplotypes. *Hum Immunol*. 1992;34(4):242-252.
- Price P, Witt C, Allcock R, et al. The genetic basis for the association of the 8.1 ancestral haplotype (A1, B8, DR3) with multiple immunopathological diseases. *Immunol Rev*. 1999;167:257-274.
- Tokunaga K, Ishikawa Y, Ogawa A, et al. Sequence-based association analysis of HLA class I and II alleles in Japanese supports conservation of common haplotypes. *Immunogenetics*. 1997;46(3):199-205.
- Saito S, Ota S, Yamada E, Inoko H, Ota M. Allele frequencies and haplotypic associations defined by allelic DNA typing at HLA class I and class II loci in the Japanese population. *Tissue Antigens*. 2000;56(6):522-529.
- Shiina T, Ota M, Shimizu S, et al. Rapid evolution of major histocompatibility complex class I genes in primates generates new disease alleles in humans via hitchhiking diversity. *Genetics*. 2006;173(3):1555-1570.
- Bilbao JR, Calvo B, Aransay AM, et al. Conserved extended haplotypes discriminate HLA-DR3-homozygous Basque patients with type 1 diabetes mellitus and celiac disease. *Genes Immun*. 2006;7(7):550-554.
- Romero V, Larsen CE, Duke-Cohan JS, et al. Genetic fixity in the human major histocompatibility complex and block size diversity in the class I region including HLA-E. *BMC Genet*. 2007;8:14.
- Aly TA, Baschal EE, Jahromi MM, et al. Analysis of single nucleotide polymorphisms identifies major type 1A diabetes locus telomeric of the major histocompatibility complex. *Diabetes*. 2008;57(3):770-776.
- Worwood M, Raha Chowdhury R, Robson KJ, Pointon J, Shearman JD, Darke C. The HLA A1-B8 haplotype extends 6 Mb beyond HLA-A: associations between HLA-A, B, F and 15 microsatellite markers. *Tissue Antigens*. 1997;50(5):521-526.
- Yunis EJ, Larsen CE, Fernandez-Vina M, et al. Inheritable variable sizes of DNA stretches in the human MHC: conserved extended haplotypes and their fragments or blocks. *Tissue Antigens*. 2003;62(1):1-20.
- Akatsuka Y, Morishima Y, Kuzushima K, Kodera Y, Takahashi T. Minor histocompatibility antigens as targets for immunotherapy using allogeneic immune reactions. *Cancer Sci*. 2007;98(8):1139-1146.
- McCarroll SA, Bradner JE, Turpeinen H, et al. Donor-recipient mismatch for common gene deletion polymorphisms in graft-versus-host disease. *Nat Genet*. 2009;41(12):1341-1344.
- Kaplan DH, Anderson BE, McNiff JM, Jain D, Shlomchik MJ, Shlomchik WD. Target antigens determine graft-versus-host disease phenotype. *J Immunol*. 2004;173(9):5467-5475.
- Battiwalla M, Hahn T, Radovic M, et al. Human leukocyte antigen (HLA) DR15 is associated with reduced incidence of acute GVHD in HLA-matched allogeneic transplantation but does not impact chronic GVHD incidence. *Blood*. 2006;107(5):1970-1973.
- Stern M, Passweg J, Tiercy JM, et al. Human leukocyte antigen DR15 is associated with reduced relapse rate and improved survival after human leukocyte antigen-identical sibling hematopoietic stem cell transplantation. *Biol Blood Marrow Transplant*. 2006;12(11):1169-1175.
- Davidson JA, Tate DG, Poulton KV, et al. HLA-DR15, reduced relapse rate and improved survival after HLA-identical sibling hematopoietic stem cell transplantation. *Biol Blood Marrow Transplant*. 2007;13(4):493-494.
- Maciejewski JP, Follmann D, Nakamura R, et al. Increased frequency of HLA-DR2 in patients with paroxysmal nocturnal hemoglobinuria and the PNH/aplastic anemia syndrome. *Blood*. 2001;98(13):3513-3519.
- Sauntharajah Y, Nakamura R, Nam JM, et al. HLA-DR15 (DR2) is overrepresented in myelodysplastic syndrome and aplastic anemia and predicts a response to immunosuppression in myelodysplastic syndrome. *Blood*. 2002;100(5):1570-1574.
- Ishikawa Y, Kashiwase K, Akaza T, et al. Polymorphisms in TNFA and TNFR2 affect outcome of unrelated bone marrow transplantation. *Bone Marrow Transplant*. 2002;29(7):569-575.
- Hansen JA, Petersdorf EW, Lin MT, et al. Genetics of allogeneic hematopoietic cell transplantation: role of HLA matching, functional variation in immune response genes. *Immunol Res*. 2008;41(1):56-78.
- Trowsdale J, Ragoussis J, Campbell RD. Map of the human MHC. *Immunol Today*. 1991;12(12):443-446.
- York IA, Rock KL. Antigen processing and presentation by the class I major histocompatibility complex. *Annu Rev Immunol*. 1996;14:369-396.



## ORIGINAL ARTICLE

# Donor single nucleotide polymorphism in the *CCR9* gene affects the incidence of skin GVHD

Y Inamoto<sup>1,2</sup>, M Murata<sup>1</sup>, A Katsumi<sup>1</sup>, Y Kuwatsuka<sup>2</sup>, A Tsujimura<sup>2</sup>, Y Ishikawa<sup>1</sup>, K Sugimoto<sup>1</sup>, M Onizuka<sup>2</sup>, S Terakura<sup>1</sup>, T Nishida<sup>2</sup>, T Kanie<sup>2</sup>, H Taji<sup>2</sup>, H Iida<sup>2</sup>, R Suzuki<sup>2</sup>, A Abe<sup>1</sup>, H Kiyoi<sup>3</sup>, T Matsushita<sup>1</sup>, K Miyamura<sup>2</sup>, Y Kodera<sup>2</sup> and T Naoe<sup>1</sup>

<sup>1</sup>Department of Hematology and Oncology, Nagoya University Graduate School of Medicine, Nagoya, Japan; <sup>2</sup>Department of Hematology, Japanese Red Cross Nagoya First Hospital, Nagoya, Japan and <sup>3</sup>Department of Infectious Diseases, Nagoya University Graduate School of Medicine, Nagoya, Japan

The interactions between chemokines and their receptors may have an important role in initiating GVHD after allogeneic hematopoietic SCT (allo-HSCT). CCL25 and CCR9 are unique because they are exclusively expressed in epithelial cells and in Peyer's patches of the small intestine. We focused on rs12721497 (G926A), one of the non-synonymous single nucleotide polymorphisms (SNPs) in the *CCR9* gene, and analyzed the SNP of donors in 167 consecutive patients who received allo-HSCT from an HLA-identical sibling donor. Genotypes were tested for associations with acute and chronic GVHD in each organ and transplant outcome. Multivariate analyses showed that the genotype 926AG was significantly associated with the incidence of acute stage  $\geq 2$  skin GVHD (hazard ratio: 3.2; 95% confidence interval (95% CI): 1.1–9.1;  $P=0.032$ ) and chronic skin GVHD (hazard ratio: 4.1; 95% CI: 1.1–15;  $P=0.036$ ), but not with GVHD in other organs or with relapse, non-relapse mortality or OS. To clarify the functional differences between genotypes, each SNP in retroviral vectors was transfected into Jurkat cells. In chemotaxis assays, the 926G transfectant showed greater response to CCL25 than the 926A transfectant. In conclusion, more active homing of *CCR9*-926AG T cells to Peyer's patches may produce changes in Ag presentation and result in increased incidence of skin GVHD.

*Bone Marrow Transplantation* (2010) 45, 363–369; doi:10.1038/bmt.2009.131; published online 15 June 2009

**Keywords:** allogeneic transplantation; CCR9; chemokine; gene polymorphism; GVHD

## Introduction

Acute GVHD is a severe complication of allogeneic hematopoietic SCT (allo-HSCT).<sup>1</sup> After Ag presentation in secondary lymphoid tissues, migration of activated donor T lymphocytes to target organs has a central role in its induction. Recent studies have shown that the migration of lymphocytes to secondary lymphoid tissues or target organs, such as the skin, liver and gut, is regulated by specific chemokines.<sup>2,3</sup> Chemokines are a group of small molecules that regulate the trafficking of leukocytes through interactions with a subset of seven transmembrane, G protein-coupled receptors (chemokine receptors).<sup>4,5</sup> Their interactions may have an important role in initiating organ-specific GVHD.

Sites of expression are ubiquitous in many chemokines. For example, CCL17, which is well known as a skin-homing chemokine, is also expressed in many other organs, including the adrenal gland, bronchus, cerebellum, colon, heart and liver.<sup>4</sup> CCL28 is expressed by epithelial cells in several mucosal tissues, including the trachea, small intestine, colon, rectum, salivary gland and mammary gland.<sup>6,7</sup> By contrast, CCL25 (thymus-expressed chemokine) and its receptor CCR9 are unique because, outside the thymus, they are almost exclusively expressed by epithelial cells and Peyer's patches in the small intestine.<sup>8–10</sup> Therefore, we focused on CCR9 because it may influence the onset of intestinal GVHD or Ag presentation in Peyer's patches.

The *CCR9* gene is located on chromosome 3p21.3. A variety of single nucleotide polymorphisms (SNPs) in the *CCR9* gene have been reported, although their functional differences are not yet known. Within these SNPs, rs12721497 (G926A) is non-synonymous in exons, and it is the sole SNP whose frequency and linkage disequilibrium have been reported. This SNP alters the CCR9 amino acid sequence of the third exoloop from Val272 to Met272. We hypothesize that this SNP may have an effect on the onset of GVHD and transplant outcome because of the differences in the tissue-specific migration of T lymphocytes.

Correspondence: Dr A Katsumi, Department of Hematology and Oncology, Nagoya University Graduate School of Medicine, 65 Tsurumai-cho, Showa-ku, Nagoya 466-8560, Japan.

E-mail: katsumi@med.nagoya-u.ac.jp

Received 3 February 2009; revised 22 April 2009; accepted 24 April 2009; published online 15 June 2009

## Materials and methods

### Patients

A total of 186 consecutive patients received allogeneic BM or PBSC transplantation from an HLA-identical sibling donor at the Nagoya University Hospital and the Japanese Red Cross Nagoya First Hospital between 1987 and 2006. HLA matching among donor–recipient pairs was confirmed by either family study or genotyping in all patients. Of these 186 patients, 167 who received T-cell-replete transplantation and CYA in combination with short-term MTX as a GVHD prophylaxis were selected to participate in the study. CYA was administered daily at 3.0 mg/kg from day 1 as an i.v. infusion, and then switched to an oral dose at twice the i.v. dose when oral intake resumed. MTX was administered at 10 mg/m<sup>2</sup> on day 1 and, on days 3 and 6, was administered at 7 mg/m<sup>2</sup>. Informed consent was obtained from all patients and donors, and the study was approved by the ethics committees at the Nagoya University Hospital and Japanese Red Cross Nagoya First Hospital.

### Allelic discrimination of the polymorphism G926A in the CCR9 gene

The CCR9-G926A polymorphism (rs12721497) was determined by the PCR-RFLP method using genomic DNA obtained from donor PBMCs. The primers used for PCR were 5'-CACACCCTGATACAAGCCAA (forward) and 5'-CTCCAGCAACATAGACGACA (reverse). Sequences of interest were amplified by PCR, using Advantage II Polymerase Mix (Clontech Laboratories, Mountain View, CA, USA) in reaction mixtures containing 0.5 µl of genomic DNA and 10 pmol of each primer in a volume of 20 µl. Amplifications were performed using 35 cycles of denaturation at 95 °C for 30 s, annealing at 65 °C for 15 s and elongation at 72 °C for 30 s on a model 9600 thermocycler (Perkin-Elmer, Norwalk, CT, USA). After amplification, the 369-bp CCR9 fragment was digested for 2 h at 37 °C with 5 U of *NLAIII* (New England BioLabs, Ipswich, MA, USA) in a 20 µl reaction mixture. The digested products were analyzed by electrophoresis on a 1.5% agarose gel. Wild-type (AA) individuals were identified by the presence of only a 369-bp fragment, heterozygotes (AG) by the presence of both 231/138- and 369-bp fragments and homozygotes (GG) by the presence of only the 231- and 138-bp fragments. To rule out the incomplete digestion of the AG genotype, PCR products of this genotype were directly sequenced using the Applied Biosystems 310 automated DNA sequencer (Applied Biosystems, Foster City, CA, USA) following the manufacturer's instructions.

### Site-directed mutagenesis and construction of CCR9-926A and 926G expression vectors

Site-directed mutagenesis of the human wild-type CCR9 cDNA in pFIK vector (purchased from Kazusa DNA Research Institute, Kisarazu, Chiba, Japan) was carried out using the Quickchange Kit (Stratagene, La Jolla, CA, USA). Synthetic oligonucleotide primers containing the corresponding 926G point mutation had the following

sequences: 5'-CCATTGACGCCTATGCCGTGTTTCATCTCCAACGTG (forward) and 5'-ACAGTTGGAGATGAACACGGCATAGGCGTCAATGG (reverse). The oligonucleotide was amplified with Pfu turbo DNA polymerase (Stratagene), and the template plasmid was digested by *DpnI*. Each sequence of CCR9 cDNA was amplified by PCR with primers containing the following *EcoRI/NotI* sites: 5'-CGCGGAATTCATGACACCCACAGACTTCACA (forward) and 5'-ATCGGCGGCCGC TCAGAGGGAGAGTGCTCCTGAGGT (reverse). Each product was cut at *EcoRI/NotI* sites, and ligated into pMX-IRES-Puro (a kind gift from Dr Toshio Kitamura, University of Tokyo), which had been digested with *EcoRI* and *NotI*. The final construct used for cell transfection was sequenced entirely to verify the presence of the mutation and to ensure that no other variant was accidentally introduced during DNA amplification.

### Retrovirus transfection

PLAT-A packaging cells (a kind gift from Dr Toshio Kitamura, University of Tokyo) were used to produce recombinant retrovirus particles.<sup>11</sup> PLAT-A cells were transfected with retroviral vectors using FuGENE6 (Roche, Indianapolis, MN, USA). Jurkat cells were infected with each of the pMX-CCR9-926A-IRES-Puro, pMX-CCR9-926G-IRES-Puro and pMX-IRES-Puro (control) retroviruses. The cells were washed once and resuspended in the fresh selection medium containing 500 ng/ml of puromycin (Cayla, Toulouse, France), 48 h after transfection.

### Flow cytometric analysis

Phycoerythrin-labeled monoclonal anti-CCR9 (112509) was purchased from R&D Systems (Minneapolis, MN, USA). Analyses were carried out on FACSAria (BD Biosciences, San Jose, CA, USA) using the FlowJo software (Treestar, San Carlos, CA, USA).

### Chemotaxis assays

Chemotaxis assays were carried out as previously described<sup>12</sup> using 6.5-mm-thick Transwell tissue culture inserts with a 5-µm pore size (Corning, Corning, NY, USA). The transfected cell lines were starved overnight in the plain RPMI 1640 medium, suspended at  $1 \times 10^7$  cells per ml in this medium with 0.1% of BSA, and 100 µl of cell suspension was added to an upper insert in a lower well with 600 µl of the medium. After equilibration at 37 °C for 2 h, various concentrations (0–2000 ng/ml) of recombinant human CCL25 (R&D Systems) were added to the lower wells, and the plates were incubated for an additional 90 min before migrated cells in the lower well were counted.

### Statistical analysis

OS was calculated from the date of transplantation to the date of death from any cause using the Kaplan–Meier method, and *P*-values were calculated using a log-rank test. Non-relapse mortality (NRM) was defined as mortality due to any cause other than relapse or disease progression. Cumulative incidences of NRM and relapse were estimated

using Gray's method, with relapse and NRM, respectively, as a competing risk. Acute GVHD was graded by established criteria.<sup>13</sup> Chronic GVHD was evaluated in patients who survived beyond day +100, and was classified as limited or extensive according to the Seattle criteria.<sup>14</sup> A multivariate Cox model was created for grade II–IV acute GVHD, organ stages of acute GVHD, chronic GVHD, OS, NRM and relapse using stepwise selection at a significance level of 5%. Age, conditioning, disease risk, remission state, donor–recipient sex combination and graft source were used as covariates. For chronic GVHD analysis, a history of acute GVHD was included as covariates. Hazard ratios of the *CCR9* genotype were adjusted by these models. In acute GVHD analysis, patients who died before day +30 were censored. Analysis was carried out using STATA (StataCorp. 2007; Stata Statistical Software: Release 10.0. Special Edition. Stata Corporation, College Station, TX, USA). Data analyses were completed as of January 2007 using the most updated database at each institute.

## Results

### Patient characteristics

Patient characteristics are summarized in Table 1. Ninety-four male patients and 73 female patients, with a median age of 38 years, were included in the study. Our patients were afflicted with various diseases, including myeloid malignancies ( $n=106$ ), lymphoid malignancies ( $n=42$ ) and benign diseases ( $n=19$ ). Disease risk was standard in 97 patients and high in 70 patients. Standard risk included malignancies in the first and second remission, chronic myelogenous leukemia in the chronic phase, myelodysplastic syndrome with refractory anemia with or without ringed sideroblasts and benign diseases. High risk included all others. Ninety-eight patients with malignant disease received transplantation with their disease in remission. The graft source was BM in 130 patients and PBSCs in 37 patients. The conditioning regimen was myeloablative in 147 patients and reduced-intensity conditioning in 20 patients. TBI was used as part of the conditioning in 98 patients. Myeloablative conditioning regimens for malignancy included BU 16 mg/kg + CY 120 mg/kg ( $n=32$ ), CY 120 mg/kg + TBI 12 Gy ( $n=9$ ), CY 120 mg/kg + TBI 10 Gy + another agent of BU 8 mg/kg, cytarabine 8 g/m<sup>2</sup>, etoposide 50 mg/kg or melphalan 140 mg/m<sup>2</sup> ( $n=62$ ), BU 8 mg/kg + melphalan 180 mg/m<sup>2</sup> + TBI 10 Gy ( $n=12$ ) and melphalan 180 mg/m<sup>2</sup> + TBI 10 Gy ( $n=13$ ). Reduced-intensity conditioning regimens for malignancy included fludarabine 125 mg/m<sup>2</sup> + melphalan 100–180 mg/m<sup>2</sup> ( $n=20$ ). Conditioning regimens for aplastic anemia included CY 200 mg/kg + TLI 7.5 Gy ( $n=17$ ) and CY 200 mg/kg + TLI 5 Gy + TBI 5 Gy ( $n=1$ ) as previously described.<sup>15</sup> One patient with paroxysmal nocturnal hemoglobinuria was conditioned with CY 120 mg/kg + TBI 12 Gy. At a median follow-up of 42 months (range: 2–220 months), 104 patients were still alive. The estimated 4-year OS, NRM and relapse rates were 55, 18 and 30%, respectively. Causes of 26 non-relapse mortalities included bronchiolitis obliterans ( $n=2$ ), idiopathic pneumonia syndrome ( $n=8$ ), hepatic failure ( $n=1$ ), veno-occlusive

**Table 1** Patient characteristics

No. of patients	167
Median age in years (range)	38 (15–62)
Sex (M/F)	94/73
Race (Japanese/other)	167/0
<i>Disease</i>	
AML	50
ALL	30
CML	40
MDS	16
ML	6
ATL	1
MM	5
AA	18
PNH	1
<i>Disease risk</i>	
Standard	97
High	70
<i>Status at transplant among patients with malignant disease</i>	
Remission	98
Non-remission	50
<i>Graft source</i>	
Bone marrow	130
PBSC	37
<i>Gender compatibility</i>	
Female donor in male recipient	45
Others	122
<i>Conditioning</i>	
TBI-containing conditioning	98
Myeloablative conditioning for malignancy	147
BU + CY	32
CY + TBI	9
CY + TBI + another agent	62
BU + melphalan + TBI	12
Melphalan + TBI	13
Reduced-intensity conditioning for malignancy	20
Fludarabine + melphalan	20
Conditioning for aplastic anemia/PNH	
CY + TLI	17
CY + TLI + TBI	1
CY + TBI	1
<i>GVHD prophylaxis</i>	
Cyclosporine + MTX	167
Overall survival at 4 years	55%
Non-relapse mortality at 4 years	18%
Relapse rate at 4 years	30%
<i>Acute GVHD</i>	
Grade (0/1/II/III/IV)	99/37/20/8/3
Skin stage (0/1/2/3/4)	104/33/6/22/2
Liver stage (0/1/2/3/4)	159/3/1/3/1
Gut stage (0/1/2/3/4)	153/4/2/5/3
<i>Chronic GVHD (n = 155 evaluable)</i>	
None	91
Limited/extensive	13/51
Organ (eye/oral/skin/lung/liver)	21/55/31/2/29

Abbreviations: AA = aplastic anemia; ATL = adult T-cell leukemia/lymphoma; F = female; M = male; MDS = myelodysplastic syndrome; ML = malignant lymphoma; MM = multiple myeloma; PNH = paroxysmal nocturnal hemoglobinuria.

**Table 2** Proportion of patients who developed GVHD in each genotype

Events	Genotype 926AG	Genotype 926AA
<i>Acute GVHD</i>		
Grade II–IV	2/10 (20%)	29/157 (18%)
Skin stage 2–4	4/10 (40%)	26/157 (17%)
Liver stage 2–4	1/10 (10%)	4/157 (2.5%)
Gut stage 2–4	1/10 (10%)	9/157 (5.7%)
<i>Chronic GVHD</i>		
Limited/extensive	4/10 (40%)	60/145 (41%)
Eye	2/10 (20%)	19/145 (13%)
Oral	4/10 (40%)	51/145 (35%)
Skin	3/10 (30%)	28/145 (19%)
Lung	1/10 (10%)	1/145 (0.6%)
Liver	2/10 (20%)	27/145 (19%)


disease ( $n=2$ ), hepatitis ( $n=2$ ), intestinal bleeding ( $n=1$ ), transplant-associated microangiopathy ( $n=3$ ), acute GVHD ( $n=3$ ), scleroderma ( $n=1$ ) and bacterial or fungal pneumonia ( $n=3$ ). The incidence rates of grade II–IV, III–IV acute GVHD and chronic limited/extensive GVHD were 18.6, 6.6 and 41%, respectively. The incidence rates of stage 2–4 skin, liver and gut involvement were 18, 3 and 6%, respectively.

*Frequency of CCR9 genotypes*

Ten donors had genotype 926AG by the RFLP method, which was subsequently confirmed using direct sequence. The frequencies of the 926AA, 926AG and 926GG genotypes among the donors were 94, 6 and 0%, respectively, which were comparable with those reported in the HapMap-JPT database (<http://www.hapmap.org>).

*Hazard analysis and the effect of CCR9 genotypes on transplant outcome*

The proportion of patients who developed GVHD in each genotype is summarized in Table 2. Grade II–IV GVHD, stage 2–4 skin GVHD, stage 2–4 liver GVHD, stage 2–4 gut GVHD and chronic limited/extensive GVHD developed in 2, 4, 1, 1 and 4 patients, respectively, among patients whose donor had the genotype 926AG, whereas they developed in 29, 26, 4, 9 and 60 patients, respectively, among patients whose donor had the genotype 926AA. The estimated 4-year OS, NRM and relapse rates were not significantly different between G926A genotypes (56 vs 55%,  $P=0.78$ ; 33 vs 17%,  $P=0.32$ ; 10 vs 32%,  $P=0.19$ , respectively) (Figure 1). Multivariate analyses showed that PBSC transplantation was a risk factor for grade II–IV GVHD, stage 2–4 skin GVHD and stage 2–4 gut GVHD; high risk disease was a risk factor for OS and relapse; age of more than 40 years was a risk factor for OS and NRM; and female-to-male transplantation was a risk factor for chronic liver GVHD (Table 3, middle column). Hazard ratios of the genotype 926AG, adjusted by these factors, are listed in Table 3 (right column). The genotype 926AG was significantly associated with acute stage 2–4 skin GVHD (hazard ratio: 3.2; 95% confidence interval (95% CI): 1.1–9.1;  $P=0.032$ ) and chronic skin GVHD (hazard ratio: 4.1; 95% CI: 1.1–15;  $P=0.036$ ), whereas it was not



**Figure 1** Overall survival between G926A genotypes. At a median follow-up of 42 months (range: 2–220 months), the estimated 4-year overall survival was not statistically different between G926A genotypes (56 vs 55%,  $P=0.78$ ).

associated with grade II–IV GVHD or with stage 2–4 liver GVHD, stage 2–4 gut GVHD, limited or extensive chronic GVHD, chronic GVHD in organs other than skin, OS, NRM or relapse. PBSCs were used in only 1 of 10 patients who received transplantation from 926AG donors.

*Functional comparison between CCR9-926A and 926G*

To clarify the functional differences between genotypes of 926A and 926G, we created cDNA constructs with each genotype using the Quickchange Kit. Each construct was transfected into Jurkat cells with retroviral vectors. After 3 weeks of selection with puromycin, we analyzed the expression of CCR9 in each of the stably transfected cells. Cells were stained with phycoerythrin-labeled monoclonal anti-CCR9. The level of CCR9 expression was higher in CCR9-transfected cells compared with that in control-transfected cells, and equivalent between the 926A and 926G genotypes (Figure 2a).

We next analyzed the migration of Jurkat cells transfected with control vectors, 926A and 926G, as well as plain Jurkat cells in response to varying concentrations of CCL25, using porous Transwell tissue culture inserts to separate the cells in the upper chambers from the chemokine-containing medium in the lower chambers. As shown in Figure 2b, CCR9-transfected cells, but not the control-transfected or plain Jurkat cells, migrated in response to CCL25 in a dose-dependent manner, producing a bell-shaped curve. It is noted that CCR9-926G-expressing cells were more responsive to CCL25 compared with those expressing CCR9-926A.

**Discussion**

Several genetic polymorphisms of inflammatory cytokine genes are reported to affect the outcome of allo-HSCT.<sup>16–21</sup> Chemokines are another group of cytokines that control the trafficking of leukocytes through interactions with chemokine receptors. We hope to clarify the role of these