

Figure 4 Effects of gain and loss of function of HIF-2 α on endochondral ossification parameters in cultures of chondrogenic cells. **(a)** mRNA levels of *Col10a1*, *Mmp13* and *Vegfa*, alkaline phosphatase (ALP) activity (relative to control) and Alizarin red staining in stable lines of ATDC5 cells retrovirally transfected with GFP, HIF-2 α , ARNTL or both HIF-2 α and ARNTL and in nontransfected parental cells (-) after culture for 3 weeks with ITS and 2 d with Pi. HIF-2 α and ARNTL levels were confirmed by western blotting, with the actin level as the internal control. **(b)** Analyses of the parameters in **a** in stable ATDC5 lines transfected with GFP or HIF-2 α mutants at the oxygen-dependent hydroxylation residues causing enhancement (N847A and P531A) and abrogation (P849A) of HIF-2 α transactivation activity under the culture conditions used in **a**. Gene expression was confirmed by RT-PCR with the *EPAS1* primer set inside the coding sequence, with the *Gapdh* level as the internal control. **(c)** Analyses of the same read-outs in a in stable ATDC5 lines transfected with GFP or HA-tagged dominant-negative HIF-2 α (HA-dnHIF-2 α) (left) or siRNA specific for *GFP* or *Epas1* mRNA (right) under the culture conditions used in **a**. HA-dnHIF-2 α and Hif-2 α amounts were confirmed by western blotting. **(d)** mRNA levels of *Col10a1*, *Mmp13* and *Vegfa* and *Epas1* in the pellet cultures of primary chondrocytes derived from wild-type (WT) and *Epas1*^{+/-} littermate embryos for 2 weeks. For the rescue experiment, adenoviral transfection with HIF-2 α (Ax-HIF-2 α) or the control GFP (Ax-GFP) was performed before the pellet formation. All data are expressed as means \pm s.d. **P* < 0.05 versus GFP unless otherwise indicated.

central three steps of endochondral ossification (chondrocyte hypertrophy, cartilage degradation and vascularization) and found that all were the direct transcriptional targets. However, there are other factors related to endochondral ossification, including in the earlier cartilage formation step and in the later osteogenesis step, which might also be targets of HIF-2 α . We therefore examined the expression patterns of the following factors: type II collagen (COL2A1) and aggrecan (AGC1) as cartilage matrix proteins; RUNX2, Indian hedgehog (IHH), and type 1 PTH/PTHrP receptor (PTH1R) as chondrocyte hypertrophy markers; MMP-3, MMP-9, a disintegrin and metalloproteinase with thrombospondin type 1 motif-4 (ADAMTS4) and ADAMTS5 as cartilage-degradable proteinases; and type I collagen (COL1A1), bone sialoprotein (BSP), osteocalcin, alkaline phosphatase, bone morphogenetic protein-2 (BMP-2), BMP-4 and BMP-7 as osteogenic markers. In cultured ATDC5 cells, expression of chondrocyte hypertrophy markers, cartilage degradable proteinases and osteogenic markers increased in accordance with the cell differentiation and *Epas1* expression (Supplementary Fig. 3a). The chondrocyte hypertrophy markers and cartilage degradable proteinases were localized mainly in the hypertrophic zone of the mouse limb cartilage, similarly to Hif-2 α (Supplementary Fig. 3b). The mRNA levels of cartilage matrix proteins, chondrocyte hypertrophy markers and most of the osteogenic markers were increased in ATDC5 cells overexpressing HIF-2 α or HIF-2 α -ARNTL (Supplementary




Fig. 4a). Among cartilage-degradable proteinases, expression of *Mmp3* and *Mmp9* was increased, whereas neither *Adamts4* nor *Adamts5* was affected (Supplementary Fig. 4a). In contrast, mRNA levels of the chondrocyte hypertrophy markers *Mmp3* and *Mmp9* were decreased after overexpression of a dominant-negative mutant form of HIF-2 α and siRNA specific for *Epas1* mRNA in ATDC5 cells (Supplementary Fig. 4b,c); however, cartilage matrix proteins, *Adamts4*, *Adamts5* and most of osteogenic markers were little affected. Primary chondrocytes derived from *Epas1*^{+/-} mice reproducibly showed suppression of the chondrocyte hypertrophy markers, *Mmp3* and *Mmp9*, but not cartilage matrix proteins, *Adamts4*, *Adamts5* or osteogenic factors, and the suppression was restored to wild-type levels by HIF-2 α overexpression (Supplementary Fig. 5). Further *in vivo* analyses of embryonic limbs (Supplementary Fig. 6a) and osteoarthritic knee joints (Supplementary Fig. 6b) confirmed the decreases in expression of the chondrocyte hypertrophy markers, *Mmp-3* and *Mmp-9*, but not cartilage matrix proteins *Adamts4* or *Adamts5*, by Hif-2 α insufficiency. HIF-2 α enhanced the promoter activities of the chondrocyte hypertrophy markers *MMP3* and *MMP9*, as well as *Col10a1*, *Mmp13*, and *Vegfa* mRNA levels, much more strongly than HIF-1 α , and the stimulation of the mRNA levels by HIF-2 α was not altered by cotransfection of




Figure 5 Contribution of HIF-2 α to osteoarthritis development in mice and humans. (a) Cartilage degradation assessed by safranin O staining and expression of Hif-2 α , Col10a1, Mmp-13 and Vegf by immunostaining (brown) and immunofluorescence (green) in mouse knee joints 0 and 8 weeks after creating a surgical osteoarthritis model in 8-week-old wild-type (WT) and *Epas1*^{+/-} littermates. Boxed areas in each safranin O-stained or each immunostained image indicate the regions shown in the enlarged safranin O-stained or immunofluorescent image immediately below. Scale bars, 100 μ m. (b) Quantification of osteoarthritis development by our (two left graphs) and OARS1 (two right graphs) grading systems. Data are expressed as means \pm s.d. #*P* < 0.05, **P* < 0.01 versus WT. (c) Safranin O staining, H&E staining and immunofluorescence with an antibody to HIF-2 α in human tibial cartilages of various degradation stages, subchondral bone (beneath the cartilage with Mankin score = 8) and synovium (around the cartilage with Mankin score = 8), obtained as surgical specimens of total knee arthroplasty. Scale bars, 100 μ m. (d) Top, the identified SNP, rs17039192, and primers used for genotyping (red lines) in the human *EPAS1* gene. CDS, coding sequence. Bottom, association of the rs17039192 (+18C/T) SNP with knee osteoarthritis (OA) diagnosed on radiographs using the Kellgren/Lawrence grade in a Japanese population. The odds ratio of the susceptibility allele was 1.44 (95% confidence interval: 1.08–1.92). **P* = 0.05, ***P* = 0.013. (e) Luciferase activities in chondrogenic SW1353, OUMS27 and ATDC5 cells and nonchondrogenic HeLa cells transfected with a luciferase reporter gene construct ligated to a fragment (-1,000 bp to +488 bp) containing +18C or +18T. Data are shown as means \pm s.d. **P* < 0.05 versus 18C.

HIF-1 α (Supplementary Fig. 7a,b). Furthermore, the endochondral ossification parameters stimulated by HIF-2 α overexpression in ATDC5 cells were not inhibited by suppression of RUNX2 through overexpression of a dominant-negative mutant RUNX2 (Supplementary Fig. 7c).

Finally, to identify the upstream mechanism that regulates HIF-2 α , we performed a screen of transcription factors with the *EPAS1* promoter fragment including the +18C SNP described above. Among candidate molecules that are known to regulate chondrocyte differentiation, such as sex-determining region Y box (SOX), RUNX, CCAAT/enhancer binding protein (C/EBP), MEF2, SP/Kruppel-like factor (KLF), activating transcription factor (ATF), cAMP responsive element-binding protein (CREB), Notch and NF- κ B family members, we found that v-rel reticuloendotheliosis viral oncogene homolog A (RELA or NF- κ B p65), an essential molecule

of the NF- κ B signal, showed the strongest activation in all cells (Fig. 6a). In the *EPAS1* promoter we identified an NF- κ B motif and found that site-directed mutagenesis in the motif caused suppression of transactivation by RELA (Fig. 6b). The allelic difference (+18C/T) of the rs17039192 SNP described above also altered the activation of *EPAS1* promoter by RELA but did not affect it when the NF- κ B motif was mutated, suggesting the involvement of this SNP in the *EPAS1* transactivation and osteoarthritis development caused by the NF- κ B signal. We further confirmed that the proinflammatory cytokines tumor necrosis factor- α (TNF- α) and interleukin-1 β (IL-1 β), putative inducers of the NF- κ B signal²⁸, increased *EPAS1* expression in cultured chondrogenic cells (Fig. 6c). In mouse knee joint cartilage, the expression of *Rela* was increased during osteoarthritis development, similarly to Hif-2 α expression (Fig. 6d).




Figure 6 Upstream mechanism that regulates HIF-2 α . (a) Luciferase activities after transfections of putative chondrocyte-related transcription factors into chondrogenic SW1353, OUMS27 and ATDC5 cells and nonchondrogenic HeLa cells with a reporter construct containing a fragment (−1,000 bp to +488 bp) of the *EPAS1* gene. OSX, osterix; AP2, transcription factor AP-2 α ; Notch1-ICD, intercellular domain of Notch1; HES1, hairy and enhancer of split 1. Data are shown as means \pm s.d. (b) Top, depiction of the NF- κ B motif (−983 to −973) in the human *EPAS1* gene. Bottom, site-directed mutagenesis analyses of the luciferase assay in the three chondrogenic cell lines transfected with GFP or RELA. Luciferase activities were compared with or without mutation in the NF- κ B motif and with +18C or +18T of the rs17039192 SNP. Data are shown as means \pm s.d. * P < 0.05 versus wild-type NF- κ B and 18C with RELA. (c) mRNA levels of *EPAS1* in the three chondrogenic cells cultured with or without TNF- α or IL-1 β (each 1 ng ml^{−1}) for 2 d. Data are expressed as means \pm s.d. * P < 0.05 versus control. (d) Time course of degradation in mouse knee joint cartilage, as shown by Safranin O staining and expression of Rel α and Hif-2 α by immunostaining and immunofluorescence, respectively, in a surgical osteoarthritis model in 8-week-old mice. Boxed areas in each of the top images are enlarged in the bottom images directly beneath. Scale bar, 100 μ m.

DISCUSSION

Among the sequential steps of endochondral ossification—cartilage formation, chondrocyte hypertrophy, cartilage degradation, vascularization and osteogenesis—this study reveals that HIF-2 α functions as an extensive transcriptional regulator of the central three steps. HIF-2 α shares about 50% amino acid homology with HIF-1 α (ref. 20), a potent regulator of cartilage homeostasis^{14–16}; however, accumulating evidence has shown distinct expression patterns and functions between the two HIF proteins^{17–21}. HIF-1 α is expressed mainly in hypovascular and hypoxic tissues^{16,19,29}, whereas HIF-2 α is expressed even in vascularized tissues^{11,29}. In cartilage as well, previous studies and our current study show that HIF-1 α is expressed from the early stage of cartilage formation, and its activity is enhanced by hypoxia^{13–16,30}. In contrast, HIF-2 α is expressed mainly in highly differentiated chondrocytes, and its function is independent of oxygen-dependent hydroxylation. Likewise, although cartilage-specific knockout of HIF-1 α leads to defects in the earlier stage of cartilage formation and the

later stage of chondrocyte survival and osteogenesis^{14–16}, *Epas1*^{+/-} mice show growth retardation with defects in the central steps of endochondral ossification. Hence, HIF-1 α and HIF-2 α may have distinct roles via different mechanisms: hypoxia-dependent cartilage formation and maintenance by HIF-1 α and less hypoxia-dependent endochondral ossification by HIF-2 α . We have also confirmed that neither gain nor loss of function of HIF-2 α alters HIF-1 α expression during chondrocyte differentiation, nor does HIF-1 α transfection affect *EPAS1* promoter activity. Furthermore, HIF-1 α hardly stimulates the expression of markers of the central steps of endochondral ossification in the presence or absence of HIF-2 α , indicating independent functions of HIF-1 α and HIF-2 α at least in the central steps. However, we do not deny the possibility of interactions between HIF-1 α and HIF-2 α in earlier and later stages of endochondral ossification. The HIF-2 α function may possibly be compensated by HIF-1 α in this earlier stage, as cartilage matrix proteins are not altered by HIF-2 α suppression. In the later or severe stage of osteoarthritic

cartilage, expression of HIF-2 α decreases after reaching a maximum at the initiation of cartilage degradation in mice and humans, as was reported in a previous study³¹. In this terminal stage, the decreased HIF-2 α expression may enhance autophagy in mature chondrocytes of osteoarthritic cartilage, as HIF-2 α is known to antagonize the autophagy-accelerator function of HIF-1 α ³².

Our study reveals that *COL10A1*, *MMP13* and *VEGFA* are the direct transcriptional targets of HIF-2 α . The functional relationships *in vivo* are supported by previous reports that a deficiency of *Mmp13* or *Vegfa* in mice causes a skeletal phenotype similar to that in *Epas1*^{+/-} mice, with elongation of the hypertrophic zone and delay of ossification in the limb cartilage^{9,10}, although the skeletal phenotypes of *Col10a1*-knockout mice differs among the reports^{33–35}. Our further studies identify *RUNX2*, *IHH*, *PTH1R*, *MMP3* and *MMP9* as possible transcriptional targets of HIF-2 α . We have recently reported that HIF-2 α enhances *Runx2* promoter activity³⁶ and that *Runx2*^{+/-} mice show resistance to osteoarthritis development under mechanical instability, similarly to *Epas1*^{+/-} mice⁵. However, HIF-2 α and *RUNX2* may promote endochondral ossification via independent mechanisms.

There are two mechanisms of osteoarthritis protection: induction of anabolism or inhibition of catabolism in joint cartilage. The protection in *Epas1*^{+/-} mice is not likely to be due to induction of anabolism, as the anabolic markers *COL2A1* and *AGC1* are unaffected in joint cartilage. Although recent studies have identified *ADAMTS5* and related molecules as key catabolic regulators of osteoarthritis development^{37–40}, neither *ADAMTS4* nor *ADAMTS5* is regulated by HIF-2 α , implicating another pathway. Because *Mmp13*^{-/-} mice are reported to be protected from cartilage degradation despite considerable aggrecan loss after surgical osteoarthritis induction⁴¹, similarly to *Epas1*^{+/-} mice, the osteoarthritis protection caused by the Hif-2 α insufficiency might occur principally through regulation of *Mmp-13*.

Suppression of osteoarthritis development was obvious in *Epas1*^{+/-} mice, whereas skeletal growth retardation was mild and transient, suggesting that pathological endochondral ossification is more dependent on HIF-2 α than is physiological endochondral ossification. As a trigger of osteoarthritis, mechanical stress may induce the upstream NF- κ B signal and HIF-2 α expression in joint cartilage, which causes endochondral ossification by transactivation of *COL10A1*, *MMP13*, *VEGFA* and other factors. Recent comprehensive profiling analyses of not only genes and proteins, but also microRNAs, is unraveling the molecular network underlying osteoarthritis development⁴²; however, we hereby propose that signals in the HIF-2 α axis from NF- κ B signaling to endochondral ossification-related molecules may represent a rational therapeutic target for osteoarthritis with minimal effects on physiological skeletal homeostasis.

METHODS

Methods and any associated references are available in the online version of the paper at <http://www.nature.com/naturemedicine/>.

Note: Supplementary information is available on the Nature Medicine website.

ACKNOWLEDGMENTS

We thank R. Yamaguchi and H. Kawahara for technical assistance. This study was supported by a grant-in-aid for Scientific Research from the Japanese Ministry of Education, Culture, Sports, Science and Technology (19109007 and 20689028). The sponsor had no role in study design, data collection, data analysis, data interpretation or writing of the manuscript.

AUTHOR CONTRIBUTIONS

T.S., T.I. and H.K. performed project planning; T.S., A.F., A.M., F.Y. and S.O. performed the experiments; T.S., A.M., N.N., T.A., N.Y., T.N., K.N., K.T., U.-i.C. and H.K. conducted data analysis; T.S. and H.K. wrote the manuscript.

COMPETING FINANCIAL INTERESTS

The authors declare no competing financial interests.


Published online at <http://www.nature.com/naturemedicine/>.

Reprints and permissions information is available online at <http://npg.nature.com/reprintsandpermissions/>.

- Kronenberg, H.M. Developmental regulation of the growth plate. *Nature* **423**, 332–336 (2003).
- Kühn, K., D'Lima, D.D., Hashimoto, S. & Lotz, M. Cell death in cartilage. *Osteoarthritis Cartilage* **12**, 1–16 (2004).
- Kawaguchi, H. Endochondral ossification signals in cartilage degradation during osteoarthritis progression in experimental mouse models. *Mol. Cells* **25**, 1–6 (2008).
- Kamekura, S. *et al.* Osteoarthritis development in novel experimental mouse models induced by knee joint instability. *Osteoarthritis Cartilage* **13**, 632–641 (2005).
- Kamekura, S. *et al.* Contribution of runt-related transcription factor 2 to the pathogenesis of osteoarthritis in mice after induction of knee joint instability. *Arthritis Rheum.* **54**, 2462–2470 (2006).
- Yamada, T. *et al.* Carminerin contributes to chondrocyte calcification during endochondral ossification. *Nat. Med.* **12**, 665–670 (2006).
- Drissi, H., Zuscik, M., Rosier, R. & O'Keefe, R. Transcriptional regulation of chondrocyte maturation: potential involvement of transcription factors in OA pathogenesis. *Mol. Aspects Med.* **26**, 169–179 (2005).
- Ortega, N., Behonick, D.J. & Werb, Z. Matrix remodeling during endochondral ossification. *Trends Cell Biol.* **14**, 86–93 (2004).
- Stickens, D. *et al.* Altered endochondral bone development in matrix metalloproteinase 13-deficient mice. *Development* **131**, 5883–5895 (2004).
- Zelzer, E. *et al.* VEGFA is necessary for chondrocyte survival during bone development. *Development* **131**, 2161–2171 (2004).
- Semenza, G.L. HIF-1 and human disease: one highly involved factor. *Genes Dev.* **14**, 1983–1991 (2000).
- Schofield, C.J. & Ratcliffe, P.J. Oxygen sensing by HIF hydroxylases. *Nat. Rev. Mol. Cell Biol.* **5**, 343–354 (2004).
- Lando, D., Peet, D.J., Whelan, D.A., Gorman, J.J. & Whitelaw, M.L. Asparagine hydroxylation of the HIF transactivation domain a hypoxic switch. *Science* **295**, 858–861 (2002).
- Pfander, D., Cramer, T., Schipani, E. & Johnson, R.S. HIF-1 α controls extracellular matrix synthesis by epiphyseal chondrocytes. *J. Cell Sci.* **116**, 1819–1826 (2003).
- Schipani, E. Hypoxia and HIF-1 α in chondrogenesis. *Ann. NY Acad. Sci.* **1068**, 66–73 (2006).
- Schipani, E. *et al.* Hypoxia in cartilage: HIF-1 α is essential for chondrocyte growth arrest and survival. *Genes Dev.* **15**, 2865–2876 (2001).
- Patel, S.A. & Simon, M.C. Biology of hypoxia-inducible factor-2 α in development and disease. *Cell Death Differ.* **15**, 628–634 (2008).
- O'Rourke, J.F., Tian, Y.M., Ratcliffe, P.J. & Pugh, C.W. Oxygen-regulated and transactivating domains in endothelial PAS protein 1: comparison with hypoxia-inducible factor-1 α . *J. Biol. Chem.* **274**, 2060–2071 (1999).
- Jain, S., Maltepe, E., Lu, M.M., Simon, C. & Bradfield, C.A. Expression of ARNT, ARNT2, HIF1 α , HIF2 α and Ah receptor mRNAs in the developing mouse. *Mech. Dev.* **73**, 117–123 (1998).
- Tian, H., Hammer, R.E., Matsumoto, A.M., Russell, D.W. & McKnight, S.L. The hypoxia-responsive transcription factor EPAS1 is essential for catecholamine homeostasis and protection against heart failure during embryonic development. *Genes Dev.* **12**, 3320–3324 (1998).
- Scortegagna, M. *et al.* Multiple organ pathology, metabolic abnormalities and impaired homeostasis of reactive oxygen species in *Epas1*^{-/-} mice. *Nat. Genet.* **35**, 331–340 (2003).
- Komori, T. Regulation of skeletal development by the Runx family of transcription factors. *J. Cell. Biochem.* **95**, 445–453 (2005).
- Arnold, M.A. *et al.* MEF2C transcription factor controls chondrocyte hypertrophy and bone development. *Dev. Cell* **12**, 377–389 (2007).
- Magee, C., Nurminskaya, M., Faverman, L., Galera, P. & Linsenmayer, T.F. SP3/SP1 transcription activity regulates specific expression of collagen type X in hypertrophic chondrocytes. *J. Biol. Chem.* **280**, 25331–25338 (2005).
- Pescador, N. *et al.* Identification of a functional hypoxia-responsive element that regulates the expression of the egl nine homologue 3 (*eglN3/phd3*) gene. *Biochem. J.* **390**, 189–197 (2005).
- Pritzker, K.P. *et al.* Osteoarthritis cartilage histopathology: grading and staging. *Osteoarthritis Cartilage* **14**, 13–29 (2006).
- Muraki, S. *et al.* Prevalence of radiographic knee osteoarthritis and its association with knee pain in the elderly of Japanese population-based cohorts: the ROAD study. *Osteoarthritis Cartilage* **17**, 1137–1143 (2009).
- Li, Q. & Verma, I.M. NF- κ B regulation in the immune system. *Nat. Rev. Immunol.* **2**, 725–734 (2002).
- Stewart, A.J., Houston, B. & Farquharson, C. Elevated expression of hypoxia inducible factor-2 α in terminally differentiating growth plate chondrocytes. *J. Cell. Physiol.* **206**, 435–440 (2006).
- Amarillo, R. *et al.* HIF1 α regulation of Sox9 is necessary to maintain differentiation of hypoxic prechondrogenic cells during early skeletogenesis. *Development* **134**, 3917–3928 (2007).

ARTICLES

31. Bohensky, J. *et al.* Regulation of autophagy in human and murine cartilage: hypoxia-inducible factor 2 suppresses chondrocyte autophagy. *Arthritis Rheum.* **60**, 1406–1415 (2009).
32. Srinivas, V., Bohensky, J., Zahm, A.M. & Shapiro, I.M. Autophagy in mineralizing tissues: microenvironmental perspectives. *Cell Cycle* **8**, 391–393 (2009).
33. Gress, C.J. & Jacenko, O. Growth plate compressions and altered hematopoiesis in collagen X null mice. *J. Cell Biol.* **149**, 983–993 (2000).
34. Kwan, K.M. *et al.* Abnormal compartmentalization of cartilage matrix components in mice lacking collagen X: implications for function. *J. Cell Biol.* **136**, 459–471 (1997).
35. Rosati, R. *et al.* Normal long bone growth and development in type X collagen-null mice. *Nat. Genet.* **8**, 129–135 (1994).
36. Tamiya, H. *et al.* Analysis of the Runx2 promoter in osseous and non-osseous cells and identification of HIF2A as a potent transcription activator. *Gene* **416**, 53–60 (2008).
37. Echtermeyer, F. *et al.* Syndecan-4 regulates ADAMTS-5 activation and cartilage breakdown in osteoarthritis. *Nat. Med.* **15**, 1072–1076 (2009).
38. Glasson, S.S. *et al.* Deletion of active ADAMTS5 prevents cartilage degradation in a murine model of osteoarthritis. *Nature* **434**, 644–648 (2005).
39. Lin, A.C. *et al.* Modulating hedgehog signaling can attenuate the severity of osteoarthritis. *Nat. Med.* **15**, 1421–1425 (2009).
40. Stanton, H. *et al.* ADAMTS5 is the major aggrecanase in mouse cartilage in vivo and in vitro. *Nature* **434**, 648–652 (2005).
41. Little, C.B. *et al.* Matrix metalloproteinase 13-deficient mice are resistant to osteoarthritic cartilage erosion but not chondrocyte hypertrophy or osteophyte development. *Arthritis Rheum.* **60**, 3723–3733 (2009).
42. Iliopoulos, D., Malizos, K.N., Oikonomou, P. & Tsezou, A. Integrative microRNA and proteomic approaches identify novel osteoarthritis genes and their collaborative metabolic and inflammatory networks. *PLoS One* **3**, e3740 (2008).

© 2010  nature publishing group

To order reprints, please contact:

Americas: Tel 212 726 9278; Fax 212 679 0843; author-reprints@nature.com

Europe/UK/ROW: Tel + 44 (0)20 7833 4000; Fax + 44 (0)20 7843 4500; author-reprints@nature.com

Japan & Korea: Tel +81 3 3267 8751; Fax +81 3 3267 8746; reprints@naturejpn.com

IOF position statement: vitamin D recommendations for older adults

B. Dawson-Hughes · A. Mithal · J.-P. Bonjour ·
S. Boonen · P. Burckhardt · G. E.-H. Fuleihan ·
R. G. Josse · P. Lips · J. Morales-Torres · N. Yoshimura

Received: 9 March 2010 / Accepted: 16 April 2010 / Published online: 27 April 2010
© International Osteoporosis Foundation and National Osteoporosis Foundation 2010

Abstract This position paper of the International Osteoporosis Foundation makes recommendations for vitamin D nutrition in elderly men and women from an evidence-based perspective.

Keywords Musculoskeletal health · Requirement · Vitamin D

This material is based in part upon work supported by the US Department of Agriculture, Agricultural Research Service, under agreement no. 58-1950-7-707. Any opinions, findings, conclusion, or recommendations expressed in this publication are those of the author(s) and do not necessarily reflect the view of the US Department of Agriculture, the Centers for Disease Control and Prevention or the US Department of Health and Human Services.

This statement has been endorsed by the IOF Committee of Scientific Advisors

B. Dawson-Hughes (✉)
Jean Mayer USDA Human Nutrition Research Center on Aging,
Tufts University,
711 Washington Street,
Boston, MA 02111, USA
e-mail: bess.dawson-hughes@tufts.edu

A. Mithal
Indraprastha Apollo Hospitals, Sarita Vihar,
Delhi-Mathura Road,
110044 New Delhi, India

J.-P. Bonjour
University Hospital, Division of Bone Diseases,
University Hospitals and Faculty of Medicine,
Rue Micheli-du-Crest 24,
1211 Geneva, Switzerland

Introduction

Vitamin D is important for bone and muscle development, function, and preservation. The serum 25OHD concentration is the best available clinical indicator of vitamin D status. Until recently, optimal serum 25OHD concentration was considered to be that level associated with maximal parathyroid hormone (PTH) suppression. Estimates of that threshold level have clustered around 32–50 nmol/L (12.8–20 ng/ml) and 68–75 nmol/L (27.2–30 ng/ml), depending upon analytical approach used [1]. In the last decade, however, the evidence base for older men and women has grown to include many randomized, controlled clinical trials (RCTs) with falls and fracture endpoints. Because the RCTs have for the most part been conducted in men and women over the age of 60 to 65 years, our recommendations are directed at this large and growing older segment of the adult population. Our objective is to use available evidence to support recommendations for optimal vitamin D status. We approach this by examining

S. Boonen
Centre for Metabolic Bone Diseases and Division of Geriatric
Medicine, University of Leuven,
Herestraat 49,
3000 Leuven, Belgium

P. Burckhardt
Association Suisse contre l'Ostéoporose,
Clinique Bois-Cerf,
Avenue d'Ouchy 31,
1006 Lausanne, Switzerland

G. E.-H. Fuleihan
Calcium Metabolism and Osteoporosis Program,
American University of Beirut-Medical Center,
Riad El Solh, PO BOX: 11-0236, 1107 2020 Beirut, Lebanon

the efficacy of different vitamin D doses administered and levels of 25OHD achieved in reducing risk of falls and fractures. In this process, it is important to consider other factors that influence serum 25OHD levels and responses to oral vitamin D supplementation.

Determinants of serum 25OHD levels and of the serum 25OHD response to oral vitamin D

Vitamin D intake and effective sun exposure are the major determinants of the serum 25OHD level. Several factors influence the increment in serum 25OHD in response to a given dose of vitamin D₃, including the starting level of 25OHD. At a dose of 2.5 µg (100 IU/d), the mean increment ranges from 2.75 nmol/l (1.1 ng/ml) at low starting 25OHD levels to 1.75 nmol/l (0.7 ng/ml) at higher (near optimal) starting levels [2]. The increment in 25OHD in response to a given dose of vitamin D also varies with body size. It is smaller in subjects with high BMI than in individuals with normal BMI [3, 4]. Other factors affect 25OHD levels but have no known impact on 25OHD responses to supplemental vitamin D. Estrogen use increases measured serum 25OHD levels by increasing levels of vitamin D binding protein [5] but does not alter the serum 25OHD increment achieved with supplementation. Serum 25OHD levels decline with aging, but the serum 25OHD response to a given dose of supplemental vitamin D₃ is not affected by age [6]. Similarly, the dietary calcium intake, within the range usually consumed, does not affect the serum 25OHD response to vitamin D supplementation [7]. (The latter statement should not be confused with the observations that the *calcium* requirement may be dependent upon vitamin D status and that an adequate calcium intake is important for bone health [8].) Finally, serum 25OHD levels vary widely across commonly used assays. Until this problem is addressed by widespread use of standard reference material such as the NIST standards [9] and participation in the DEQAS quality control program (www.deqas.org), assay variability will continue to complicate the process of determining the desired 25OHD level and the impact of a given dose on serum 25OHD levels.

R. G. Josse
Division of Endocrinology and Metabolism,
University of Toronto, St Michael's Hospital Health Centre,
61 Queen Street East,
Toronto, ON M5C 2T2, Canada

P. Lips
Division of Internal Medicine, Endocrine Section,
VU University Medical Center,
PO Box 7057, 1007MB Amsterdam, Netherlands

Falls

Vitamin D is thought to act on myocyte vitamin D receptors to exert its effect on muscle tissue. In prospective studies, lower serum 25OHD levels have been associated with decreased grip strength and appendicular muscle mass in older men and women [10, 11]. Supplementation with vitamin D has improved lower extremity muscle performance and reduced risk of falling in several high-quality double blind RCTs [12]. These trials have employed doses up to 25 µg (1,000 IU) of vitamin D per day, with and without calcium. Supplementation in amounts of 17.5 to 25 µg/day (700–1,000 IU/day) lowered risk of falling by 20% in older individuals, independent of their calcium intake level. In contrast, supplementation with doses of <17.5 µg/day (<700 IU/day) had no detectable effect on falls. From this meta-analysis of available data, it appears that a mean serum 25OHD level of at least 60 nmol/L (24 ng/ml) is needed for optimal fall risk reduction. Observational studies suggest that there may be benefit to increasing serum 25OHD levels beyond 60 nmol/L (24 ng/ml), but higher levels (and doses) have not been evaluated in RCTs.

Fractures

Vitamin D affects fracture risk through its effects on bone metabolism and on risk of falling. Randomized controlled trials indicate that supplementation with vitamin D reduces rates of bone loss in older women [13]. The impact of supplemental vitamin D on fracture risk has been examined mainly in men and women age 65 and older. A recent meta-analysis revealed that vitamin D in doses in the range of more than 10 through 20 µg/day (>400–800 IU/day) reduced risk of non-vertebral and hip fracture by approximately 20% whereas doses up through 10 µg/day (400 IU/day) had no evident effect [14]. Doses above 20 µg/day (800 IU/day) have not been studied. The mean serum 25OHD level associated with reduction in non-vertebral fracture risk was 66 nmol/L (26.4 ng/ml). Hip fracture risk reduction was observed at a mean 25OHD level of 74 nmol/L (29.6 ng/ml) and higher. Based on this

J. Morales-Torres
Hospital Aranda de la Parra,
Hidalgo 329-704,
León 37000 GTO, Mexico

N. Yoshimura
Department of Joint Disease Research,
22nd Century Medical and Research Center,
University of Tokyo,
7-3-1 Hongo, Bunkyo-ku,
Tokyo 113-8655, Japan

and other evidence, eight of ten IOF Working Group members felt that 75 nmol/L (30 ng/ml) is the appropriate target level of serum 25OHD for older individuals; in contrast, two members felt that the target should be 50 to 75 nmol/L (20 to 30 ng/ml). The estimate of 75 nmol/L (30 ng/ml) is close to the higher cluster of 25OHD levels associated with maximal PTH suppression [1].

Other potential benefits

Vitamin D insufficiency has been implicated as a contributing factor in a growing number of important chronic diseases including type 2 diabetes, cardiovascular disease, selected cancers, and autoimmune diseases as well as infections, and also to increased mortality. RCTs are needed before causal relationships can be determined and optimal 25OHD levels for prevention can be established. The doses and mean serum levels needed to achieve optimal impact on these non-classical outcomes are not clear.

Global vitamin D status

Vitamin D insufficiency, whether defined as 25OHD levels <75 or <50 nmol/L (<30 or <20 ng/ml), is prevalent worldwide [15]. For instance, the prevalence of levels <75 nmol/L (<30 ng/ml) in postmenopausal women has been reported to be approximately 50% in Thailand and Malaysia, 75% in the USA, and 90% in Japan and South Korea. Vitamin D deficiency, defined as a level below 25 nmol/L (10 ng/ml) is very common in the Middle East and South Asia where mean levels range from 10 to 30 nmol/L (4 to 12 ng/ml) [15, 16]. The high prevalence of suboptimal 25OHD levels in older men and women around the world raises the possibility that many falls and fractures can be prevented with vitamin D supplementation.

Vitamin D preparations

Vitamin D is available in two forms, vitamin D₂ (ergocalciferol) and vitamin D₃ (cholecalciferol). Some find that the two forms are equally effective in raising the serum 25OHD level [17] whereas others find that vitamin D₃ gives a larger increment [18]. We generally recommend that vitamin D₃ be used when available. In some parts of the world, active metabolites are available for use in the treatment of osteoporosis. These metabolites are not a substitute for adequate vitamin D intake. Vitamin D is the substrate for 25OHD and the circulating 25OHD level may be important to support the non-renal production of 1,25-dihydroxyvitamin D. Local production of 1,25-dihydrox-

vitamin D appears to mediate some of the non-classical effects of vitamin D.

Recommendations

The estimated average vitamin D requirement for older adults to reach a serum 25OHD level of 75 nmol/L (30 ng/ml) is 20 to 25 µg/day (800 to 1,000 IU/day). Considerably higher doses would be needed to ensure that almost all older adults reached 75 nmol/L (30 ng/ml). Efficacy of doses higher than 20 µg/day (800 IU/day) for fractures and 25 µg/day (1,000 IU/day) for falls however have not been evaluated in RCTs. It is therefore premature to recommend higher intakes for all older adults at this time.

The repletion dose will vary among individuals according to their starting level, their BMI, their effective sun exposure, and other unidentified factors. An intake lower than 20 µg/day (800 IU/day) may be adequate for individuals with regular effective sun exposure. Intake may need to be adjusted upward to as much as 50 µg/day (2,000 IU/day) in individuals who are obese, and in those with osteoporosis, limited sun exposure (institutionalized, homebound), and malabsorption, and in non-European populations known to be at high risk for vitamin D deficiency such as those in the Middle East and South Asia, or immigrants from such regions living in Europe. In these and other high-risk individuals, we recommend measuring the serum 25OHD level. The required dose to reach 75 nmol/L can be estimated from the measured level. Each 2.5 µg (100 IU) of added vitamin D will increase the serum 25OHD level by about 2.5 nmol/L (range 1.75–2.75 nmol/L) or 1.0 ng/ml (range 0.7 to 1.1 ng/ml) [2]. Because of the variability in individual 25OHD responses to supplemental vitamin D, however, in high-risk individuals, the serum 25OHD levels should be retested after about 3 months of supplementation to confirm that the target 25OHD level has been reached.

Conflicts of interest None.

References

1. Durazo-Arvizu RA, Dawson-Hughes B, Sempos CT, Yetley EA, Looker AC, Cao G, Harris SS, Burt VL, Carriquiry AL, Picciano MF (2010) Three-phase model harmonizes estimates of the maximal suppression of parathyroid hormone by 25-hydroxyvitamin D in persons 65 years of age and older. *J Nutr*. doi:10.3945/jn.109.116681
2. Heaney RP, Davies KM, Chen TC, Holick MF, Barger-Lux MJ (2003) Human serum 25-hydroxycholecalciferol response to

- extended oral dosing with cholecalciferol. *Am J Clin Nutr* 77:204–210
3. Wortsman J, Matsuoka LY, Chen TC, Lu Z, Holick MF (2000) Decreased bioavailability of vitamin D in obesity. *Am J Clin Nutr* 72:690–693 [erratum appears in *Am J Clin Nutr*. 2003 May;77(5):1342]
 4. Blum M, Dallal GE, Dawson-Hughes B (2008) Body size and serum 25 hydroxy vitamin D response to oral supplements in healthy older adults. *J Am Coll Nutr* 27:274–279
 5. Harris SS, Dawson-Hughes B (1998) The association of oral contraceptive use with plasma 25-hydroxyvitamin D levels. *J Am Coll Nutr* 17:282–284
 6. Harris SS, Dawson-Hughes B (2002) Plasma vitamin D and 25OHD responses of young and old men to supplementation with vitamin D3. *J Am Coll Nutr* 21:357–362
 7. Gousous R, Song L, Dallal G, Dawson-Hughes B (2005) The effect of calcium intake on the vitamin D requirement. *J Clin Endocrinol Metab* 90:707–711
 8. Boonen S, Lips P, Bouillon R, Bischoff-Ferrari HA, Vanderschueren D, Haentjens P (2007) Need for additional calcium to reduce the risk of hip fracture with vitamin D supplementation: evidence from a comparative metaanalysis of randomized controlled trials. *J Clin Endocrinol Metab* 92:1415–1423
 9. May W, Parris R, Beck C, Fassett J, Greenberg R, Guenther F, Kramer G, Wise S, Gills T, Colbert J, Gettings R, MacDonald B (2000) Definitions of terms and modes used at NIST for value-assignment of reference materials for chemical measurements. In *Technology NIOSa* (ed). US Government Printing Office, Gaithersburg
 10. Flicker L, MacInnis RJ, Stein MS, Scherer SC, Mead KE, Nowson CA, Thomas J, Lowndes C, Hopper JL, Wark JD (2005) Should older people in residential care receive vitamin D to prevent falls? Results of a randomized trial. *J Am Geriatr Soc* 53:1881–1888
 11. Visser M, Deeg DJ, Lips P (2003) Low vitamin D and high parathyroid hormone levels as determinants of loss of muscle strength and muscle mass (sarcopenia): the Longitudinal Aging Study Amsterdam. *J Clin Endocrinol Metab* 88:5766–5772
 12. Bischoff-Ferrari HA, Dawson-Hughes B, Staehelin HB, Orav EJ, Stuck AE, Theiler R, Wong JB, Egli A, Kiel DP, Henschkowski J (2009) Fall prevention with supplemental and active forms of vitamin D: a meta-analysis of randomised controlled trials. *Br Med J* 339:b3692
 13. Ooms ME, Roos JC, Bezemer PD, van der Vijgh WJ, Bouter LM, Lips P (1995) Prevention of bone loss by vitamin D supplementation in elderly women: a randomized double-blind trial. *J Clin Endocrinol Metab* 80:1052–1058
 14. Bischoff-Ferrari HA, Willett WC, Wong JB, Giovannucci E, Dietrich T, Dawson-Hughes B (2005) Fracture prevention by vitamin D supplementation: a meta-analysis of randomized controlled trials. *J Am Med Assoc* 293:2257–2264
 15. Mithal A, Wahl DA, Bonjour JP, Burckhardt P, Dawson-Hughes B, Eisman JA, El-Hajj Fuleihan G, Josse RG, Lips P, Morales-Torres J (2009) Global vitamin D status and determinants of hypovitaminosis D. *Osteoporos Int* 20:1807–1820
 16. El-Hajj Fuleihan G (2009) Vitamin D deficiency in the Middle East and its health consequences. *Clin Rev Bone Miner Metab* 7:77–93
 17. Holick MF, Biancuzzo RM, Chen TC, Klein EK, Young A, Bibuld D, Reitz R, Salameh W, Ameri A, Tannenbaum AD (2008) Vitamin D2 is as effective as vitamin D3 in maintaining circulating concentrations of 25-hydroxyvitamin D. *J Clin Endocrinol Metab* 93:677–681
 18. Trang HM, Cole DE, Rubin LA, Pierratos A, Siu S, Vieth R (1998) Evidence that vitamin D3 increases serum 25-hydroxyvitamin D more efficiently than does vitamin D2. *Am J Clin Nutr* 68:854–858

Health-related quality of life with vertebral fracture, lumbar spondylosis and knee osteoarthritis in Japanese men: the ROAD study

Shigeyuki Muraki · Toru Akune · Hiroyuki Oka · Yoshio En-yo · Munehito Yoshida · Akihiko Saika · Takao Suzuki · Hideyo Yoshida · Hideaki Ishibashi · Fumiaki Tokimura · Seizo Yamamoto · Kozo Nakamura · Hiroshi Kawaguchi · Noriko Yoshimura

Received: 30 July 2009 / Accepted: 15 February 2010 / Published online: 24 April 2010
© International Osteoporosis Foundation and National Osteoporosis Foundation 2010

Abstract

Summary This study examined associations of VFx, lumbar spondylosis and knee OA with QOL in 767 men over 40 years old from the ROAD study (mean, 69.7 years.). Multiple regression analysis showed VFx and knee OA as significantly associated with lower PCS scores, but lumbar spondylosis was not.

Electronic supplementary material The online version of this article (doi:10.1007/s11657-010-0039-3) contains supplementary material, which is available to authorized users.

S. Muraki (✉) · T. Akune
Department of Clinical Motor System Medicine,
22nd Century Medical and Research Center, Faculty of Medicine,
the University of Tokyo,
Hongo 7-3-1, Bunkyo,
Tokyo 113-8655, Japan
e-mail: murakis-ort@h.u-tokyo.ac.jp

H. Oka · N. Yoshimura
Department of Joint Disease Research,
22nd Century Medical and Research Center, Faculty of Medicine,
the University of Tokyo,
Tokyo, Japan

Y. En-yo · M. Yoshida · A. Saika
Department of Orthopaedic Surgery,
Wakayama Medical University,
Wakayama, Japan

T. Suzuki · H. Yoshida · H. Ishibashi · F. Tokimura · S. Yamamoto
Tokyo Metropolitan Institute of Gerontology,
Tokyo, Japan

K. Nakamura · H. Kawaguchi
Department of Sensory and Motor System Medicine,
Faculty of Medicine, the University of Tokyo,
Tokyo, Japan

Purpose Little data have been accumulated regarding associations of vertebral fracture (VFx), lumbar spondylosis and knee osteoarthritis (OA) with quality of life (QOL) in men. The purpose of the present study is to clarify the impact of these diseases on QOL parameters such as the Medical Outcomes Study Short Form 8 (SF-8) and the EuroQOL (EQ-5D). In addition, to provide greater insight into the magnitude of QOL loss, a comparison was made with cerebral stroke.

Methods From the 3,040 participants in the ROAD study, this study analyzed 767 men over 40 years who had completed the questionnaires (mean, 69.7 years.). Vertebral fracture was assessed by lateral radiography of the lumbar spine. Lumbar spondylosis and knee OA were defined as Kellgren/Lawrence grade ≥ 3 . Cerebral stroke was assessed by self-report.

Results Multiple regression analysis after adjustments for age, body mass index and presence of the above four diseases showed VFx was significantly associated with lower scores in physical function (PF), role physical (RP), bodily pain (BP) and vitality (VT) domains as well as physical component summary (PCS). Knee OA were significantly associated with lower scores in PF, RP, BP and PCS scores. Lumbar spondylosis was not associated with any domains of the SF-8. Lumbar spondylosis and knee OA were significantly associated with EQ-5D utility scores, but VFx was not. The impact for VFx on BP, VT and PCS scores was larger than cerebral stroke.

Conclusions This study revealed that VFx and knee OA impaired physical QOL in men, rather than lumbar spondylosis.

Keywords Quality of life · Vertebral fracture · Lumbar spondylosis · Knee osteoarthritis · Men

Introduction

Vertebral fracture (VFX) is reportedly associated with functional impairment [1], back pain, kyphosis [2, 3], esophageal reflux [4], depressive mood [5], respiratory dysfunctions [6] and mortality [7]. Lumbar spondylosis and knee osteoarthritis (OA), characterized by pathological features including disk or joint space narrowing and osteophytosis, are also major public health issues causing chronic pain and disability among the elderly [8–12]. In fact, prevalences of lumbar spondylosis and knee OA are quite high in the elderly in Japan [13–15], and 37,900,000 and 25,300,000 people ≥ 40 years old would be affected by radiographic lumbar spondylosis and knee OA, respectively [15]. Furthermore, according to the recent National Livelihood Survey of the Ministry of Health, Labour and Welfare in Japan, OA and fracture represent the fourth and fifth among diseases that cause disabilities, respectively, subsequently requiring support with regard to activities of daily living (ADL) [16].

Gender differences have been observed in these bone and joint diseases. The prevalence of knee OA is higher in women than men [14, 15], while that of lumbar spondylosis is higher in men [13, 15]. For VFX, prevalence is higher in women [17], while mortality is higher in men [18], so the impact of these diseases on quality of life (QOL) may also differ between genders. Although several studies have examined associations of VFX [19–27] and knee OA [28–32] with QOL, men and women were not separated [20, 21, 28, 29, 31, 32] or only women were focused [22, 23], and few large-scale population-based studies have examined bone and joint diseases in men [19, 24, 27, 30]. Furthermore, the association of VFX, lumbar spondylosis and knee OA with ADL and QOL may not be independent, but no studies have examined VFX, lumbar spondylosis and knee OA simultaneously in the same population using the same tools.

The objective of the present study is to clarify the impact of VFX, lumbar spondylosis and knee OA on QOL among 767 men using the cohorts of the ROAD study. In addition, to provide greater insight into the magnitude of QOL loss with VFX, lumbar spondylosis and knee OA, we made a comparison with cerebral stroke. Cerebral stroke is ranked first among diseases causing disabilities according to the recent National Livelihood Survey of the Ministry of Health, Labour and Welfare in Japan [16], and several studies have already reported that QOL is greatly affected after cerebral stroke [33]. Thus, such information can provide the health care physician with insights into the influence of VFX, lumbar spondylosis and knee OA on QOL.

Methods

Subjects

The ROAD study is a nationwide prospective study for bone and joint diseases (with OA and osteoporosis as the representative bone and joint diseases) constituting population-based cohorts established in several communities in Japan. As detailed profile of the ROAD study has already been described elsewhere [13–15, 34], the brief summary is provided here. To date, we have completed creation of a baseline database including clinical and genetic information for 3,040 inhabitants (1,061 men, 1,979 women) in the age range of 23 to 95 years (mean, 70.6 years), recruited from listings of resident registrations in three communities: an urban region in Itabashi, Tokyo, a mountainous region in Hidakagawa, Wakayama, and a seacoast region in Taiji, Wakayama. All participants provided written informed consent, and the study was conducted with the approval of ethical committees of the University of Tokyo and the Tokyo Metropolitan Institute of Gerontology. Participants completed an interviewer-administered questionnaire of 400 items that included lifestyle information such as smoking habits, alcohol consumption, family history, history, physical activity, reproductive variables and health-related QOL. Anthropometric measurements included height, weight, bilateral grip strength and body mass index (BMI) (weight [kg]/height² [m²]). In the present study, to compare the magnitude of QOL loss in VFX, lumbar spondylosis and knee OA with another chronic disease, we assessed medical history of cerebral stroke by self-report. The following question was asked by an interviewer: “Have you ever experienced cerebral stroke?” Furthermore, to assess the impact of these bone and joint diseases according to symptoms, all participants were also interviewed regarding low back pain (LBP) by asking, “Have you experienced pain on most days in the past month, in addition to now?” Subjects who answered “yes” were defined as having LBP.

From the baseline data of the overall participants, the present study analyzed 767 men ≥ 40 years old who had completed questionnaires for the Medical Outcomes Study Short Form 8 (SF-8) and the EuroQOL (EQ-5D).

Radiographic assessment

All participants underwent radiographic examination of the lumbar spine including intervertebral levels from L1/2 to L5/S with anteroposterior and lateral views and both knees using anteroposterior and lateral views with weight-bearing and foot map positioning. Vertebral fracture, lumbar

spondylosis and knee OA were determined by a single well-experienced orthopaedist blinded to participant clinical status (S.M.). Vertebral fracture was assessed by lateral radiographs of the lumbar spine (L1–L5) in terms of a wedge, biconcave, or crush appearance according to the Japanese Society for Bone and Mineral Research (JSBMR) criteria [35]. The films were marked up, and morphometric measurements of anterior, middle and posterior heights on lateral radiography of the thoracic and lumbar spine were made. Wedge appearance was defined as a site where anterior height of the vertebra was $\leq 75\%$ than posterior height. Biconcave appearance was where the height of the central part of the vertebra was $\leq 80\%$ than that of the anterior or posterior parts of the vertebra. Crush appearance was where the height of the anterior, central and posterior parts of an axial vertebra was all reduced to $\leq 80\%$ of the normal value (Fig. 1). Lumbar spondylosis and knee OA were assessed using the Kellgren/Lawrence (KL) radiographic atlas, and severity by KL grading was determined [36]. We have defined lumbar spondylosis and knee OA as KL ≥ 3 in at least one knee and in one intervertebral level, respectively. To evaluate intraobserver agreement of the JSBMR criteria for VFx and the KL grade (0–4) for the lumbar spine and knee, 100 randomly selected radiographs were scored by the same observer at >1 month after the first reading. Furthermore, 100 other radiographs were scored by two experienced orthopaedic surgeons (S.M. and H.O.) using the same radiographic atlas for interobserver agreement. Intra- and interobserver agreements were evaluated by kappa analysis. Intra- and interobserver agreements in JSBMR criteria for VFx and KL grade for lumbar spine and knee have been shown to be sufficient for assessment (0.93 and 0.91 for VFx, 0.84 and 0.76 for lumbar spine and 0.86 and 0.80 for knee, respectively).

Instruments

To carry out the QOL assessment, we used the SF-8 Health Survey (SF-8) scale. The SF-8 is an alternate form of the




Fig. 1 Diagnostic criteria for VFXs according to the JSBMR

SF-36 Health Survey (SF-36) [37], the most widely used patient-based health status survey. The SF-8 was constructed to provide an even shorter alternative to the SF-36 for use in large population-based surveys of general and specific populations. The SF-8 uses one question to measure each of the eight SF-36 domains. Although none of the SF-8 items are identical to SF-36 items, the item pool including the SF-36 survey, SF-8 single-item scales and summary measures is scored on the same metric as the SF-36 scales and summary measures. The SF-8 and SF-36 measure eight concepts: general health (GH), physical function (PF), role physical (RP), bodily pain (BP), vitality (VT), social function (SF), mental health (MH) and role emotional (RE). Each domain includes questions regarding overall health, limitations to usual physical activities due to physical health problems, difficulties in performing daily work due to physical health, severity of pain, energy levels, limitations to usual social activities due to physical health or emotional problems, severity of emotional problems and difficulties with usual work, school or other daily activities due to personal or emotional problems, respectively. The SF-8 was scored by assigning the mean SF-36 scale score from the 2002 general Japanese population to each response category of the SF-8 measuring the same concept and then weighting each SF-8 item to compute aggregate physical (PCS) and mental (MCS) summary scale measures. The SF-8 may be scored using a published algorithm for Japanese versions of the SF-8, which have been well validated [38]. We also used the EuroQOL (EQ-5D) questionnaire [39] translated into Japanese [40]. This five-dimensional health care classification included questions on the status of morbidity, self-care, usual activities, pain/discomfort and anxiety/depression. Participants were asked to indicate current health status by ticking the most appropriate of three statements about each of five QOL dimensions. Each statement represents an increasing degree of severity. These results were coded and converted to a score of utility using the tables of values [40].

Statistical analysis

We performed nonpaired Student's *t* test to compare mean scores of QOL parameters between subjects with and without each chronic disease. Associations of VFX, lumbar spondylosis and knee OA with QOL parameters were determined by multiple regression analysis after adjustment for age and BMI. Next, to determine the independent impact of these bone and joint diseases, multiple regression analysis was used by age, BMI and presence of VFX, lumbar spondylosis and knee OA as independent variables. Furthermore, to compare the magnitude of QOL loss of the

three bone and joint diseases to cerebral stroke, multiple regression analysis was performed by age, BMI and presence of cerebral stroke in addition to VFx, lumbar spondylosis and knee OA as independent variables. Tukey honestly significant difference (HSD) test after adjustment for age and BMI was used to determine the differences of PCS values among VFx with LBP, VFx without LBP and no VFx and the differences among lumbar spondylosis with LBP, lumbar spondylosis without LBP and no lumbar spondylosis. Data analyses were performed using SAS version 9.0 (SAS Institute, Cary, NC).

Results

Impact of VFx, lumbar spondylosis and knee OA on QOL scores

Characteristics of the 767 men ≥ 40 years old in the ROAD study are shown in Table 1. Prevalence of lumbar

Table 1 Characteristics of participants

N	767
Age, years	69.7 \pm 10.5
Height, cm	162.8 \pm 6.7
Weight, kg	61.5 \pm 10.8
BMI, kg/m ²	23.1 \pm 3.4
Medical history (%)	
Cerebral stroke	5.8
Prevalence (%)	
VFx	11.6
LS	41.6
KOA	12.0
LBP	15.4
SF-8	
GH	50.2 \pm 5.5
PF	49.9 \pm 6.2
RP	50.2 \pm 6.7
BP	50.4 \pm 9.2
VT	50.4 \pm 6.3
SF	52.4 \pm 5.5
MH	54.4 \pm 5.3
RE	52.0 \pm 5.2
PCS	47.4 \pm 6.8
MCS	53.4 \pm 5.3
EQ-5D utility score	0.91 \pm 0.14

Except where indicated otherwise, values represent mean \pm SD.

BMI body mass index; *VFx* vertebral fracture; *LS* lumbar spondylosis; *KOA* knee osteoarthritis; *LBP* low back pain; *GH* general health; *PF* physical function; *RP* role physical; *BP* bodily pain; *VT* vitality; *SF* social function; *MH* mental health; *RE* role emotional; *PCS* physical component summary; *MCS* mental component summary.

spondylosis and knee OA were approximately 42% and 12%, respectively, compared to 12% for VFx. Six percent of all subjects had a medical history of cerebral stroke.

Table 2 shows scores for all domains in the SF-8 and the EQ-5D utility score according to the presence of chronic diseases. Scores for PF, RP, BP, VT and PCS in the SF-8 were significantly lower in subjects with VFx compared to those in subjects without VFx, but GH, SF and RE were not. Physical function, RP, SF and PCS were significantly lower in subjects with lumbar spondylosis compared to subjects without lumbar spondylosis. For knee OA, scores of PF, RP, BP and PCS were significantly lower compared to subjects without knee OA. For MCS, the score was higher in subjects with VFx and knee OA compared to those in subjects without them. EQ-5D utility score was significantly lower in subjects with lumbar spondylosis and knee OA compared to those without them, respectively, while no significant difference was apparent in subjects with or without VFx.

We next performed further multiple regression analyses to examine the independent association of VFx, lumbar spondylosis and knee OA with QOL parameters after adjusting for age, BMI and all other bone and joint diseases. Beta values in each domain of SF-8 and EQ-5D utility score after adjusting for age, BMI and all other bone and joint diseases are shown in Table 3. Vertebral fracture was significantly associated with lower scores in PF, RP, BP, VT and PCS, while not in GH, SF and RE. Knee OA was significantly associated with lower PF, BP and PCS scores. For MCS, VFx and knee OA were associated with higher scores, but lumbar spondylosis was not associated. Lumbar spondylosis and knee OA were significantly associated with EQ-5D utility scores, while VFx was not. The Tukey HSD test after adjustment for age and BMI showed that PCS score was significantly lower in subjects having VFx with LBP than in subjects having VFx without LBP (Fig. 2). Vertebral fracture with or without LBP was significantly associated with lower PCS scores compared with no VFx. However, PCS scores were significantly lower in subjects having lumbar spondylosis with LBP than in subjects having lumbar spondylosis without LBP. There were no significant differences in PCS scores between subjects having lumbar spondylosis without LBP and those having no lumbar spondylosis.

Comparison of the magnitude of QOL loss in VFx, lumbar spondylosis and knee OA with that in cerebral stroke

To compare the magnitude of QOL loss in VFx, lumbar spondylosis and knee OA with another chronic disease, we analyzed the association of medical history of cerebral stroke with QOL (Supplementary Table). Multiple regression analysis showed that cerebral stroke was significantly associated with lower QOL scores measured by PF, RP, BP, SF, MH and PCS in the SF-8, along with EQ-5D utility

Table 2 Mean scores of all domains, PCS and MCS in the SF-8 and EQ-5D by VFx, LS and KOA

	VFx		LS		KOA		Japanese general population*
	No	Yes	No	Yes	No	Yes	
SF-8							
GH	50.3 (5.5)	49.4 (5.4)	50.3 (5.4)	50.1 (5.6)	50.2 (5.4)	50.6 (6.2)	50.3 (6.6)
PF	50.2 (5.8)	48.1 [†] (8.2)	50.5 (5.7)	49.2 [†] (6.8)	50.3 (5.7)	47.7 [†] (8.7)	49.8 (5.0)
RP	50.5 (6.2)	47.9 [†] (9.1)	50.7 (6.3)	49.6 [†] (7.2)	50.5 (6.3)	48.4 [†] (8.6)	50.3 (5.0)
BP	50.8 (9.2)	47.0 [†] (9.1)	50.7 (9.2)	49.9 (9.3)	50.7 (9.1)	48.3 [†] (10.0)	50.8 (7.9)
VT	50.6 (6.2)	49.2 [†] (6.4)	50.4 (6.1)	50.4 (6.4)	50.5 (6.2)	49.7 (6.4)	52.1 (5.6)
SF	52.4 (5.5)	52.4 (5.6)	52.8 (5.0)	51.9 [†] (6.0)	52.3 (5.4)	53.0 (5.6)	50.3 (6.3)
MH	54.2 (5.4)	55.9 [†] (3.7)	54.5 (5.2)	54.3 (5.3)	54.2 (5.2)	55.5 [†] (5.4)	52.9 (5.9)
RE	51.9 (5.1)	52.6 (5.4)	52.2 (4.9)	51.7 (5.6)	52.0 (5.0)	52.4 (6.1)	51.3 (4.8)
PCS	47.9 (6.5)	43.9 [†] (8.4)	47.8 (6.7)	46.8 [†] (7.5)	47.8 (6.5)	44.7 [†] (8.3)	48.3 (6.2)
MCS	53.1 (5.3)	55.6 [†] (4.8)	53.4 (5.3)	53.4 (5.2)	53.1 (5.2)	55.3 [†] (5.5)	51.9 (5.8)
EQ-5D	0.91 (0.14)	0.89 (0.16)	0.93 (0.13)	0.89 [†] (0.16)	0.92 (0.14)	0.87 [†] (0.17)	

Unless otherwise indicated, values represent mean (SD).

* Reference data derived from the 2002 general Japanese men at the age of 60 to 69 years [38]

[†] $p < 0.05$ vs. subjects without the corresponding disease by non-paired Student's *t* test

score (Table 4). Adjusted beta values for PF and RP in VFx were lower than those in cerebral stroke, while these for BP, VT and PCS were higher in VFx. For knee OA, adjusted beta values for PF and RP were lower than those in cerebral stroke, while those for BP and PCS were higher. For EQ-5D utility score, lumbar spondylosis and knee OA was significantly associated with lower scores, but the adjusted beta values were lower than that in cerebral stroke.

Discussion

This is the first population-based study to examine the effects of a variety of bone and joint diseases including

VFx, lumbar spondylosis and knee OA on QOL as measured by both SF-8 and EQ-5D in Japanese men. In the present study, we performed multiple regression analysis to determine independent associations of QOL parameters with each bone and joint disease after adjustment for age, BMI and all other bone and joint diseases. Vertebral fracture and knee OA were significantly associated with lower PCS scores, while they were associated with higher MCS. Lumbar spondylosis was associated with the EQ-5D utility scores, while not with any domains in the SF-8. The impact of diseases on PCS was largest in VFx among the three bone and joint diseases in men. Furthermore, to provide greater insight into the magnitude of QOL loss with the bone and joint diseases, we compared the

Table 3 Beta values for VFx, LS and KOA in all domains, PCS and MCS in the SF-8 and EQ-5D

	VFx		LS		KOA	
	Beta ^a	Adjusted beta ^b	Beta ^a	Adjusted beta ^b	Beta ^a	Adjusted beta ^b
SF-8						
GH	-0.054	-0.053	-0.015	-0.009	0.018	0.020
PF	-0.094*	-0.088*	-0.051	-0.032	-0.087*	-0.082*
RP	-0.113*	-0.109*	-0.038	-0.016	-0.073*	-0.069
BP	-0.133*	-0.131*	-0.035	-0.010	-0.081*	-0.078*
VT	-0.074*	-0.074*	-0.003	0.011	-0.056	-0.055
SF	0.011	0.018	-0.051	0.060	0.066	0.071
MH	0.106*	0.109*	-0.013	-0.035	0.076*	0.076*
RE	0.038	0.046	-0.052	-0.062	0.032	0.035
PCS	-0.181*	-0.178*	-0.042	-0.007	-0.118*	-0.113*
MCS	0.149*	0.153*	-0.015	-0.047	0.121*	0.122*
EQ-5D	-0.050	-0.035	-0.107*	-0.096*	-0.081*	-0.073*

^a Beta values are shown using multiple regression analysis after adjustment for age and BMI

^b Adjusted beta values are shown using multiple regression analysis after adjustment for age, BMI and all other diseases

* $p < 0.05$




Fig. 2 Physical component summary values among subjects having VFX with LBP, VFX without LBP and no VFX, and those having lumbar spondylosis (LS) with LBP, LS without LBP and no LS. The number of subjects in each group is shown in parentheses. * $p < 0.05$ by Tukey HSD test adjustment for age and BMI

impact of these bone and joint diseases on QOL loss with that of cerebral stroke. The impact of VFX and knee OA on BP and PCS loss was larger than that of cerebral stroke.

Few population-based studies have examined relationships between radiographic VFX and QOL [19–21, 24, 27], and genders were only adjusted, not separated in almost all these studies, although the impact of vertebral deformities on QOL may differ between genders. In the Canadian Multicentre Osteoporosis Study [27], no strong fracture-related associations of subclinical vertebral deformity with QOL were found in men as measured by the Health Utilities Index (HUI) Mark II and III Systems [41]. In the present study, radiographic VFX showed a significant association with lower PCS scores in men, with the largest impact among the three bone and joint diseases. Furthermore, multiple regression analysis showed that the magnitude of PCS loss was stronger than that of cerebral stroke, which is ranked first among the diseases causing disabilities according to the recent National Livelihood Survey of the Ministry of Health, Labour and Welfare in Japan [16]. Furthermore, VFX with LBP had a strong effect on PCS, while VFX without LBP had a moderate but significant effect. Reasons for the discrepancy between the present study and the Camos study can be partly attributed to differences in QOL measurements. In HUI scores, multi-attribute utility score reflects global health, but PCS reflects

only physical QOL. In fact, for pain domains, subclinical vertebral deformities tend to be associated with lower HUI scores as seen for the SF-8 in the present study. Another reason may be racial differences. Racial variations exist in the prevalence of vertebral deformities [42, 43], so differences may also exist in the impact of QOL. For GH, SF and RE in the SF-8, VFX was not associated with the scores by multiple regression analysis. In this study, VFX was diagnosed according to lumbar radiographs, so many fractures in men may have occurred years previously and may have been related to trauma, which must be one reason why some domains of QOL were not impaired.

The significant effect of knee OA on QOL is shown by poorer scores in PF, BP and PCS domains in the SF-8 in multiple regression analysis. A previous survey in Chinese using a GH-related QOL measure also showed that OA has comparable impact compared with stroke, asthma and chronic obstructive pulmonary disease [32], but men and women were not separated in that analysis. The present study is the first population-based study to clarify that knee OA is significantly associated with lower PCS scores in men. Although multiple regression analysis showed that the magnitude of the impact on PCS is lower than VFX, prevalence is much higher for knee OA than for VFX, so the total burdens of these diseases might be similar.

Likewise, for lumbar spondylosis, few population-based studies have examined QOL [20]. Unlike VFX or knee OA, multiple regression analysis in the present study showed that lumbar spondylosis was not associated with PCS in men, supporting previous findings [20], although gender was only adjusted in that analysis. These results may be

Table 4 Comparison of adjusted beta values for the three bone and joint diseases such as VFX, LS and KOA with that for cerebral stroke in the SF-8 and EQ-5D

	VFX	LS	KOA	Stroke
SF-8				
GH	-0.054	0.007	0.004	-0.070
PF	-0.092*	-0.031	-0.097*	-0.107*
RP	-0.115*	-0.015	-0.083*	-0.125*
BP	-0.136*	-0.008	-0.086*	-0.079*
VT	-0.078*	0.004	-0.063	-0.029
SF	0.013	-0.061	0.061	-0.100*
MH	0.106*	-0.033	0.066	-0.096*
RE	0.042	-0.065	0.028	-0.053
PCS	-0.183*	0.007	-0.128*	-0.107*
MCS	0.151*	-0.049	0.118*	-0.036
EQ-5D	-0.039	-0.090*	-0.081*	-0.122*

Adjusted beta values are shown using multiple regression analysis after adjustment for age, BMI and all other diseases

* $p < 0.05$

explained by the fact that associations between lumbar spondylosis and LBP are not so strong [13, 44, 45]. In fact, the domain of BP score in SF-8 was not associated with lumbar spondylosis in this study.

In the present study, VFx and knee OA were significantly associated with lower PCS, while they were associated with higher MCS. Past literatures also showed the dissociation between PCS and MCS in VFx and knee OA [20, 46]. Several factors may contribute to the dissociation between MCS and PCS for VFx and knee OA. First, MCS questions within the SF-8 include generic questions about energy levels, feelings of being “downhearted and blue,” and interference in daily activities as a result of emotional problems. These questions are less sensitive to the presence of MH issues than disease-specific scales such as the Kessler psychological distress scale [47]. In fact, Hill et al. [48] showed that psychological distress has been shown to be significantly more frequent in those with arthritis than those without, although scores on the MCS were not significantly different between these two groups. Second, the dissociation may be due to a disability paradox [49], which suggests that people with chronic disabilities report serious limitations in ADL, problems in performing social roles, yet state that they have excellent or good QOL. Many subjects with VFx and knee OA had LBP or knee pain, which leads to functional impairment. This may be associated with lower scores of PCS, but the individual may not feel that the impairment of social activity or ADL was due to mental factors. Particularly in elderly individuals, pain or functional impairment may be considered a natural consequence of being elderly. Vertebral fracture and knee OA were thus not associated with lower scores for SF or RE domains in the SF-8 and, thus, showed no associations with MCS. Conversely, elderly individuals may think that having cerebral stroke is not a natural consequence of being elderly, potentially contributing to the differences between VFx, knee OA and cerebral stroke.

The present study showed that the association of chronic diseases with QOL differed between the SF-8 and EQ-5D. For VFx, PCS of the SF-8 was reduced, while EQ-5D utility score was not, while for lumbar spondylosis, both PCS and MCS of the SF-8 was not associated, but the EQ-5D utility scores were significantly reduced. The reason may be explained by the fact that in the EQ-5D, all five domains are combined together to analyze the association with chronic diseases, while PCS and MCS are analyzed separately in the SF-8. In fact, associations of VFx differed between PCS and MCS of the SF-8, so when all domains were combined together, the results may differ. Lumbar spondylosis reduced both the PCS and MCS scores, although they were not significant, so when combined, the association may be significant. For VFx and knee OA, the

SF-8 may be more useful to examine associations with QOL than the EQ-5D.

There are several limitations in the present study. First, this was a large-scale, population-based study, but a cross-sectional study of baseline data. Causal relationships could therefore not be determined. The ROAD study is a longitudinal survey, so further progress may help to elucidate any causal relationships. Second, among the 1,047 men ≥ 40 years old in the ROAD study, 767 men had completed questionnaires for both the SF-8 and the EQ-5D, so the response rate was 73.7%. Subjects who completed questionnaires may have had better QOL than those who did not, so our results regarding QOL may have represented overestimations. Third, we only used semi-quantitative methods to assess VFx. Furthermore, we used the KL system for lumbar spondylosis and knee OA. Since the KL system emphasizes osteophytosis, it is unclear how to handle lumbar spondylosis or knee OA with disc or joint space narrowing but no osteophytosis. We are currently developing a computer-aided diagnostic program to enable fully automatic measurement of the major features of VFx, lumbar spondylosis and knee OA, including joint and disc space narrowing and osteophytosis on plain radiography [50]. Fourth, cerebral stroke was assessed by self-report, so severity could not be examined. Furthermore, cerebral stroke is a serious disease, so participants are considered highly likely to know if they have been diagnosed with cerebral stroke, although some participants may not; thus, strict comparison with bone and joint diseases was limited.

Conclusions

The present cross-sectional study using a large-scale population from the ROAD study revealed that VFx and knee OA were significantly associated with lower PCS scores of the SF-8 in men, while lumbar spondylosis was not. The impact of diseases on PCS was the largest for VFx in men. Further progress, along with continued longitudinal survey in the ROAD study, will elucidate the environmental and genetic backgrounds of VFx, lumbar spondylosis and knee OA and the relationship with QOL.

Acknowledgements This work was supported by Grants-in-Aid for Scientific Research (B20390182, C20591737, C20591774), for Young Scientists (A18689031) and for Exploratory Research (19659305) from the Japanese Ministry of Education, Culture, Sports, Science and Technology, H17-Men-eki-009, H18-Choujyu-037 and H20-Choujyu-009 from the Ministry of Health, Labor and Welfare, Research Aid from the Japanese Orthopaedic Association (JOA-Subsidized Science Project Research 2006-1), and grant no. 166 from the Japan Orthopaedics and Traumatology Foundation.

The authors thank Dr. Anamizu, members of the department of orthopaedics, Mr. Kutsuma, other members of the department of

radiology at Tokyo Metropolitan Geriatric Medical Center, Mrs. Tomoko Takijiri, other members of the Public Office in Hidakagawa Town, Mrs. Tamako Tsutsumi, Mrs. Kanami Maeda, and other members of the Public Office in Taiji Town for their assistance in the location and scheduling of participants for examinations.

Conflict of Interest All authors have no conflicts of interest.

References

- Burger H, Van Daele PL, Grashuis K, Hofman A, Grobbee DE, Schutte HE, Birkenhager JC, Pols HA (1997) Vertebral deformities and functional impairment in men and women. *J Bone Miner Res* 12:152–157
- Ross PD (1997) Clinical consequences of vertebral fractures. *Am J Med* 103:30S–42S, discussion 42S–43S
- Nevitt MC, Ettinger B, Black DM, Stone K, Jamal SA, Ensrud K, Segal M, Genant HK, Cummings SR (1998) The association of radiographically detected vertebral fractures with back pain and function: a prospective study. *Ann Intern Med* 128:793–800
- Yamaguchi T, Sugimoto T, Yamada H, Kanzawa M, Yano S, Yamauchi M, Chihara K (2002) The presence and severity of vertebral fractures is associated with the presence of esophageal hiatal hernia in postmenopausal women. *Osteoporos Int* 13:331–336
- Gold DT, Lyles KW, Shipp KM, Drezner MK (2001) Osteoporosis and its nonskeletal consequences: their impact on treatment decisions. In: Marcus R, Feldman D, Kelsey J (eds) *Osteoporosis*, 2nd edn. Academic Press, San Diego, pp 819–829
- Leech JA, Dulberg C, Kellie S, Pattee L, Gay J (1990) Relationship of lung function to severity of osteoporosis in women. *Am Rev Respir Dis* 141:68–71
- Johnell O, Kanis JA, Oden A, Sernbo I, Redlund-Johnell I, Pettersson C, De Laet C, Jonsson B (2004) Mortality after osteoporotic fractures. *Osteoporos Int* 15:38–42
- Sharma L, Kapoor D (2007) Epidemiology of osteoarthritis. In: Moskowitz RW, Altman RD, Hochberg MC, Buckwalter JA, Goldberg VM (eds) *Osteoarthritis: diagnosis and medical/surgical management*, 4th edn. Lippincott Williams & Wilkins, Philadelphia, pp 3–26
- Guccione AA, Felson DT, Anderson JJ, Anthony JM, Zhang Y, Wilson PW, Kelly-Hayes M, Wolf PA, Kreger BE, Kannel WB (1994) The effects of specific medical conditions on the functional limitations of elders in the Framingham Study. *Am J Public Health* 84:351–358
- Felson DT, Zhang Y (1998) An update on the epidemiology of knee and hip osteoarthritis with a view to prevention. *Arthritis Rheum* 41:1343–1355
- Hadjipavlou AG, Simmons JW, Pope MH, Necessary JT, Goel VK (1999) Pathomechanics and clinical relevance of disc degeneration and annular tear: a point-of-view review. *Am J Orthop* 28:561–571
- Emery SE, Ringus VM (2007) Osteoarthritis of the spine. In: Moskowitz RW, Altman RD, Hochberg MC, Buckwalter JA, Goldberg VM (eds) *Osteoarthritis: diagnosis and medical/surgical management*, 4th edn. Lippincott Williams & Wilkins, Philadelphia, pp 427–452
- Muraki S, Oka H, Akune T, Mabuchi A, En-Yo Y, Yoshida M, Saika A, Suzuki T, Yoshida H, Ishibashi H, Yamamoto S, Nakamura K, Kawaguchi H, Yoshimura N (2008) Prevalence of radiographic lumbar spondylosis and its association with low back pain in the elderly of population-based cohorts: the ROAD study. *Ann Rheum Dis* 68:1401–1406
- Muraki S, Oka H, Akune T, Mabuchi A, En-Yo Y, Yoshida M, Saika A, Suzuki T, Yoshida H, Ishibashi H, Yamamoto S, Nakamura K, Kawaguchi H, Yoshimura N (2009) Prevalence of radiographic knee osteoarthritis and its association with knee pain in the elderly of Japanese population-based cohorts: the ROAD study. *Osteoarthritis Cartilage* 17:1137–1143
- Yoshimura N, Muraki S, Oka H, Mabuchi A, En-yo Y, Yoshida M, Saika A, Yoshida H, Suzuki T, Kawaguchi H, Nakamura K, Akune T. Prevalence of knee osteoarthritis, lumbar spondylosis and osteoporosis in Japanese men and women: the Research on Osteoarthritis/osteoporosis Against Disability (ROAD). *J Bone Miner Metab* 27:620–628
- Ministry of Health, Labour and Welfare (2007) The outline of the results of National Livelihood Survey 2007 available online at <http://www.mhlw.go.jp/toukei/list/20-19-1.html>.
- Yoshimura N, Kinoshita H, Oka H, Muraki S, Mabuchi A, Kawaguchi H, Nakamura K (2006) Cumulative incidence and changes in the prevalence of vertebral fractures in a rural Japanese community: a 10-year follow-up of the Miyama cohort. *Arch Osteoporos* 1:43–49
- Bliuc D, Nguyen ND, Milch VE, Nguyen TV, Eisman JA, Center JR (2009) Mortality risk associated with low-trauma osteoporotic fracture and subsequent fracture in men and women. *JAMA* 301:513–521
- Masanari N, Fujiwara S, Nakata Y, Nakashima E, Nakamura T (2007) Historical height loss, vertebral deformity, and health-related quality of life in Hiroshima cohort study. *Osteoporos Int* 18:1493–1499
- van Schoor NM, Smit JH, Twisk JW, Lips P (2005) Impact of vertebral deformities, osteoarthritis, and other chronic diseases on quality of life: a population-based study. *Osteoporos Int* 16:749–756
- Cockerill W, Lunt M, Silman AJ, Cooper C, Lips P, Bhalla AK, Cannata JB, Eastell R, Felsenberg D, Gennari C, Johnell O, Kanis JA, Kiss C, Masaryk P, Naves M, Poor G, Raspe H, Reid DM, Reeve J, Stepan J, Todd C, Woolf AD, O'Neill TW (2004) Health-related quality of life and radiographic vertebral fracture. *Osteoporos Int* 15:113–119
- Badia X, Diez-Perez A, Alvarez-Sanz C, Diaz-Lopez B, Diaz-Curiel M, Guillen F, Gonzalez-Macias J (2001) Measuring quality of life in women with vertebral fractures due to osteoporosis: a comparison of the OQLQ and QUALEFFO. *Qual Life Res* 10:307–317
- Oleksik A, Lips P, Dawson A, Minshall ME, Shen W, Cooper C, Kanis J (2000) Health-related quality of life in postmenopausal women with low BMD with or without prevalent vertebral fractures. *J Bone Miner Res* 15:1384–1392
- Adachi JD, Ioannidis G, Berger C, Joseph L, Papaioannou A, Pickard L, Papadimitropoulos EA, Hopman W, Poliquin S, Prior JC, Hanley DA, Olszynski WP, Anastassiades T, Brown JP, Murray T, Jackson SA, Tenenhouse A (2001) The influence of osteoporotic fractures on health-related quality of life in community-dwelling men and women across Canada. *Osteoporos Int* 12:903–908
- Lips P, Cooper C, Agnusdei D, Caulin F, Egger P, Johnell O, Kanis JA, Kellingray S, Leplege A, Liberman UA, McCloskey E, Minne H, Reeve J, Reginster JY, Scholz M, Todd C, de Vernejoul MC, Wiklund I (1999) Quality of life in patients with vertebral fractures: validation of the Quality of Life Questionnaire of the European Foundation for Osteoporosis (QUALEFFO). Working party for quality of life of the European Foundation for Osteoporosis. *Osteoporos Int* 10:150–160
- Tosteson AN, Gabriel SE, Grove MR, Moncur MM, Kneeland TS, Melton LJ 3rd (2001) Impact of hip and vertebral fractures on quality-adjusted life years. *Osteoporos Int* 12:1042–1049
- Adachi JD, Ioannidis G, Pickard L, Berger C, Prior JC, Joseph L, Hanley DA, Olszynski WP, Murray TM, Anastassiades T, Hopman

- W, Brown JP, Kirkland S, Joyce C, Papaioannou A, Poliquin S, Tenenhouse A, Papadimitropoulos EA (2003) The association between osteoporotic fractures and health-related quality of life as measured by the Health Utilities Index in the Canadian Multicentre Osteoporosis Study (CaMos). *Osteoporos Int* 14:895–904
28. Brazier JE, Harper R, Munro J, Walters SJ, Snaith ML (1999) Generic and condition-specific outcome measures for people with osteoarthritis of the knee. *Rheumatology (Oxford)* 38:870–877
 29. Xie F, Li SC, Luo N, Lo NN, Yeo SJ, Yang KY, Fong KY, Thumboo J (2007) Comparison of the EuroQol and short form 6D in Singapore multiethnic Asian knee osteoarthritis patients scheduled for total knee replacement. *Arthritis Rheum* 57:1043–1049
 30. Woo J, Lau E, Lee P, Kwok T, Lau WC, Chan C, Chiu P, Li E, Sham A, Lam D (2004) Impact of osteoarthritis on quality of life in a Hong Kong Chinese population. *J Rheumatol* 31:2433–2438
 31. Bachmeier CJ, March LM, Cross MJ, Lapsley HM, Tribe KL, Courtenay BG, Brooks PM (2001) A comparison of outcomes in osteoarthritis patients undergoing total hip and knee replacement surgery. *Osteoarthritis Cartilage* 9:137–146
 32. Lam CL, Lauder IJ (2000) The impact of chronic diseases on the health-related quality of life (HRQOL) of Chinese patients in primary care. *Fam Pract* 17:159–166
 33. Carod-Artal FJ, Egido JA (2009) Quality of life after stroke: the importance of a good recovery. *Cerebrovasc Dis* 27(Suppl 1):204–214
 34. Yoshimura N, Muraki S, Oka H, Kawaguchi H, Nakamura K, Akune T. Cohort profile: Research on Osteoarthritis/osteoporosis Against Disability (ROAD) study *Int J Epidemiol* (in press)
 35. Inoue T (1990) Clinical features and findings: osteoporosis. *Bone* 4:39–47 (in Japanese)
 36. Kellgren JH, Lawrence JS (eds) (1963) *The epidemiology of chronic rheumatism: atlas of standard radiographs of arthritis*. Blackwell Scientific, Oxford, England
 37. Ware JE, Snow KK, Kosinski M, Gandek B (1993) *SF-36 health survey manual and interpretation guide*. The Health Institute, New England Medical Centre, Boston, USA
 38. Fukuhara S, Suzukamo Y (2004) *Manual of the SF-8 Japanese version*. Institute for Health Outcome and Process Evaluation Research, Kyoto, Japan (in Japanese)
 39. Dolan P (1997) Modeling valuations for EuroQol health states. *Med Care* 35:1095–1108
 40. Japanese EuroQol Translation Team (1997) The development of the Japanese EuroQol instrument. *J Health Care Soc* 8:109–123 (in Japanese)
 41. Feeny DH, Torrance GW, Furlong WJ (1996) Health utilities index. In: Spilker B (ed) *Quality of life and pharmacoeconomics in clinical trials*, 2nd edn. Lippincott, Philadelphia, pp 239–252
 42. Duan Y, Wang XF, Evans A, Seeman E (2005) Structural and biomechanical basis of racial and sex differences in vertebral fragility in Chinese and Caucasians. *Bone* 36:987–998
 43. Tracy JK, Meyer WA, Grigoryan M, Fan B, Flores RH, Genant HK, Resnik C, Hochberg MC (2006) Racial differences in the prevalence of vertebral fractures in older men: the Baltimore Men's Osteoporosis Study. *Osteoporos Int* 17:99–104
 44. Symmons DP, van Hemert AM, Vandenbroucke JP, Valkenburg HA (1991) A longitudinal study of back pain and radiological changes in the lumbar spines of middle aged women. II. Radiographic findings. *Ann Rheum Dis* 50:162–166
 45. O'Neill TW, McCloskey EV, Kanis JA, Bhalla AK, Reeve J, Reid DM, Todd C, Woolf AD, Silman AJ (1999) The distribution, determinants, and clinical correlates of vertebral osteophytosis: a population based survey. *J Rheumatol* 26:842–848
 46. Boutron I, Rannou F, Jardinaud-Lopez M, Meric G, Revel M, Poiraudou S (2008) Disability and quality of life of patients with knee or hip osteoarthritis in the primary care setting and factors associated with general practitioners' indication for prosthetic replacement within 1 year. *Osteoarthritis Cartilage* 16:1024–1031
 47. Kessler RC, Andrews G, Colpe LJ, Hiripi E, Mroczek DK, Normand SL, Walters EE, Zaslavsky AM (2002) Short screening scales to monitor population prevalences and trends in non-specific psychological distress. *Psychol Med* 32:959–976
 48. Hill CL, Gill T, Taylor AW, Daly A, Grande ED, Adams RJ (2007) Psychological factors and quality of life in arthritis: a population-based study. *Clin Rheumatol* 26:1049–1054
 49. Albrecht G, Devlieger P (1999) The disability paradox: high quality of life against all odds. *Soc* 48:977–988
 50. Oka H, Muraki S, Akune T, Mabuchi A, Suzuki T, Yoshida H, Yamamoto S, Nakamura K, Yoshimura N, Kawaguchi H (2008) Fully automatic quantification of knee osteoarthritis severity on plain radiographs. *Osteoarthritis Cartilage* 16:1300–1306

Osteoarthritis and Cartilage



Association of radiographic and symptomatic knee osteoarthritis with health-related quality of life in a population-based cohort study in Japan: the ROAD study

S. Muraki †*, T. Akune †, H. Oka ‡, Y. En-yo §, M. Yoshida §, A. Saika §, T. Suzuki ||, H. Yoshida ||, H. Ishibashi ||, F. Tokimura ||, S. Yamamoto ||, K. Nakamura ¶, H. Kawaguchi ¶, N. Yoshimura ‡

† Department of Clinical Motor System Medicine, 22nd Century Medical and Research Center, Faculty of Medicine, The University of Tokyo, Tokyo, Japan

‡ Department of Joint Disease Research, 22nd Century Medical and Research Center, Faculty of Medicine, The University of Tokyo, Tokyo, Japan

§ Department of Orthopaedic Surgery, Wakayama Medical University, Wakayama, Japan

|| Tokyo Metropolitan Institute of Gerontology, Tokyo, Japan

¶ Department of Sensory & Motor System Medicine, Faculty of Medicine, The University of Tokyo, Tokyo, Japan

ARTICLE INFO

Article history:

Received 4 August 2009

Accepted 17 June 2010

Keywords:

Osteoarthritis

Knee

Quality of life

Cohort

Epidemiology

SUMMARY

Objective: Knee osteoarthritis (OA) is a major public health issue causing chronic pain and disability. However, there is little information on the impact of this disease on quality of life (QOL) in Japanese men and women. The objective of the present study was to clarify the impact of radiographic and symptomatic knee OA on QOL in Japan.

Methods: This study examined the association of radiographic and symptomatic knee OA with QOL parameters such as the Medical Outcomes Study Short Form-8 (SF-8), EuroQOL (EQ-5D) and Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC). Radiographic knee OA was defined according to Kellgren/Lawrence (KL) grades, and symptomatic knee OA was defined as KL = 3 or 4 with knee pain. We also examined the independent association of symptomatic knee OA and grip strength with QOL.

Results: From the 3040 participants in the Research on Osteoarthritis Against Disability (ROAD) study, the present study analyzed 2126 subjects older than 40 years who completed the questionnaires (767 men and 1359 women; mean age, 68.9 ± 10.9 years). Subjects with KL = 3 or 4 had significantly lower physical QOL as measured by the physical component summary (PCS) score of the SF-8 and pain domains of the WOMAC, whereas mental QOL, as measured by the mental component summary (MCS) score of the SF-8, was higher in subjects with KL = 3 or 4 than KL = 0 or 1. Symptomatic knee OA was significantly more likely than radiographic knee OA without pain to be associated with physical QOL loss as measured by the PCS score and physical domains of the WOMAC. Symptomatic knee OA and grip strength were independently associated with physical QOL.

Conclusion: This cross-sectional study revealed that subjects with symptomatic knee OA had significantly lower physical QOL than subjects without it.

© 2010 Osteoarthritis Research Society International. Published by Elsevier Ltd. All rights reserved.

Introduction

Knee osteoarthritis (OA) is a major public health issue that causes chronic pain and disability^{1–3}. The prevalence of radiographic knee OA is high in Japan⁴, with 25,300,000 subjects aged 40

years and older estimated to experience radiographic knee OA⁵. According to the recent National Livelihood Survey of the Ministry of Health, Labour and Welfare in Japan, OA is ranked fourth among diseases that cause disabilities that subsequently require support with activities of daily living⁶.

Quality of life (QOL) measurements in patients with chronic diseases are useful tools for estimating disease impact; these QOL scales may be generic or disease specific. Among the generic scales, the EuroQOL (EQ-5D) has been widely used to measure health-related QOL (HRQOL) in patients with OA^{7,8}, and several studies have used the Medical Outcomes Study Short Form-36 (SF-36) in

* Address correspondence and reprint requests to: Shigeyuki Muraki, Department of Clinical Motor System Medicine, 22nd Century Medical and Research Center, Faculty of Medicine, The University of Tokyo, Hongo 7-3-1, Bunkyo-ku, Tokyo 113-8655, Japan. Tel: 81-3-5800-9178; Fax: 81-3-5800-9179.

E-mail address: murakis-ort@h.u-tokyo.ac.jp (S. Muraki).