

the expression ratio of *COL1A2* to *COL1A1* was compared between OA cartilage and other normal tissues containing type I collagen as a major component (Figure 3C). While the expression ratio of *COL1A2* to *COL1A1* was between 0.7 and 1.7 in the bone, ligament, or meniscus tissues obtained from nonarthritic joints, the ratio in OA cartilage ranged widely from 0.2 to 44. The poor coordination in expression suggests that the expression of type I collagen genes could be induced by an aberrant mechanism(s) in OA cartilage.

In contrast to type I collagen, the induction of type III collagen messenger RNA (mRNA) was consistently observed in OA samples. Within OA cartilage, the expression of type III collagen was most intense in the upper region of degenerated cartilage (Figure 3D). The expression of another gene, fibronectin, was consistently induced in OA cartilage. The regional change of fibronectin expression was very similar to that of type III collagen expression (Figure 3E).

Unlike type I or type III collagen, the induction of type X collagen was observed primarily in the deep zone (Figure 3F). The induction was weaker than that of type I or type III collagen as judged by the ratios of expression to that of *GAPDH*, and the level of induction was considerably different among OA samples; the expression was virtually absent in approximately half of the samples. Interestingly, the expression of type X collagen was more obvious in the less degenerated areas than in the more degenerated areas where the superficial zone was lost to the disease.

Consistent with previous reports, the expression of exon 2 of the *COL2A1* gene was obviously increased in OA cartilage when evaluated by the ratio of its expression to that of *GAPDH* (Figure 3G). However, the expression of exon 2 relative to total *COL2A1* expression was rather reduced in OA cartilage (Figure 3H). Thus, it was assumed that the appearance of type IIA procollagen might not be the result of a phenotypic change in the chondrocytes as previously speculated (11,12), but is more likely to be associated with the up-regulation of type II collagen expression.

Chondrocytes at the upper part of degenerated cartilage undergo a phenotypic change. Next, we compared gene expression between preserved areas and degenerated areas in the respective cartilage zones of the respective OA joints. In the superficial zone, the expression was compared in each sample between the preserved and degenerated areas (i.e., between the 2 regions in the superficial zone without and with macroscopic degeneration). In the middle and deep zones, the comparison was performed in each sample between the

preserved areas and the degenerated areas where the zones were directly exposed to the joint cavity.

The result clearly indicated that a shift occurred in the pattern of gene expression at the upper region of degenerated cartilage (Figure 4). In the degenerated areas in the middle and deep zones, the expression of cartilage matrix genes (type II collagen and aggrecan) was suppressed, while the expression of minor cartilaginous genes (type III collagen and fibronectin) was enhanced. In the superficial zone, the expression of minor cartilaginous genes was induced similarly in the degenerated areas, although the suppression of cartilage matrix gene expression was not apparent. In spite of considerable differences in expression levels among the samples, the shift of gene expression was consistently observed in almost all OA samples. Thus, the chondrocytes are considered to undergo a phenotypic change at the upper region of degenerated cartilage, no matter in which cartilage zone the cells reside.

Expression of *SOX* genes in OA and control cartilage. During chondrogenic differentiation, the expression of cartilage matrix genes is regulated by the transcriptional factors *SOX5*, *SOX6*, and *SOX9* (24). In order to estimate the involvement of these molecules in the change of chondrocyte metabolism in OA, their expression was investigated (Figure 5). In OA cartilage, the expression of *SOX* genes tended to be reduced in the degenerated areas, particularly in the upper region of the degenerated cartilage. The reduction was most obvious with *SOX6*, followed by *SOX9*, and was least apparent with *SOX5*. In the preserved areas, the expression of *SOX5* and *SOX6* tended to be increased above control levels, although this trend was not observed with *SOX9*. These regional changes of *SOX* expression within OA cartilage suggested that the altered *SOX* gene expression might be related to the change in matrix gene expression in OA.

Correlation of gene expression in OA cartilage. In an attempt to understand the mechanism(s) underlying the altered gene expression in OA cartilage, a possible correlation of gene expression was investigated (Figure 6A). The expression of 3 cartilage matrix genes correlated significantly. The expression of type II collagen was significantly correlated with that of aggrecan ($r = 0.110$, $P = 0.0081$) (Figure 6B), and a stronger correlation was observed between aggrecan and link protein ($r = 0.512$, $P < 0.0001$) (Figure 6C). A significant correlation was also observed between type II collagen and link protein ($r = 0.294$, $P < 0.0001$) (Figure 6D), implying that the expression of these genes might be modulated by a common factor(s) in OA cartilage.

In contrast, no significant correlation was found between the expression of cartilage matrix genes and minor cartilaginous genes induced by the disease in any combination (from $P = 0.102$ to $P = 0.991$) (Figure 6A).

Among the 5 minor cartilaginous genes evaluated, a significant correlation was observed only between type III collagen and fibronectin ($r = 0.764$, $P < 0.0001$) (Figure 6E). Therefore, the expression of minor cartilaginous genes was assumed to occur without any association in OA cartilage, except for that of type III collagen and fibronectin. Interestingly, the correlation between type III collagen and fibronectin was stronger than any other relationship observed in this study, suggesting the presence of certain link(s) in their expression. In fact, we have obtained data indicating that the expression of type III collagen in human OA cartilage could be induced, at least partly, through the activation of $\alpha 5\beta 1$ integrin by fibronectin (Fukui N: unpublished observation).

Next, a possible correlation of expression was investigated between the *SOX* genes and the 3 cartilage matrix genes. Although no significant correlation was found between *SOX5* or *SOX6* and the matrix genes (from $P = 0.072$ to $P = 0.857$) (Figure 6A), the expression of all 3 matrix genes was significantly correlated with that of *SOX9* (Figures 6F–H). The correlation was strongest with aggrecan ($r = 0.627$, $P < 0.0001$), followed by link protein ($r = 0.560$, $P < 0.0001$), and was weakest with type II collagen ($r = 0.270$, $P = 0.013$). The expression of *SOX* genes was not correlated with that of the minor cartilaginous genes in any combination (from $P = 0.436$ to $P = 0.959$) (Figure 6A). Meanwhile, the expression of *SOX* genes was mutually correlated. Significant correlations were observed between *SOX5* and *SOX6* ($r = 0.527$, $P < 0.0001$) (Figure 6I) and between *SOX5* and *SOX9* ($r = 0.468$, $P = 0.001$) (Figure 6J), although the correlation between *SOX6* and *SOX9* was not significant ($P = 0.728$).

DISCUSSION

The result of this study has provided a comprehensive view of the change in metabolic activity of the chondrocytes in OA. The profile of gene expression differed considerably with the site, depending on the cartilage zone and the extent of cartilage degeneration. In the macroscopically intact areas of OA cartilage, the expression of cartilage matrix genes was markedly enhanced, particularly in the middle and deep zones. This observation was consistent with the results of previous studies using in situ hybridization (3,4,6,9), in which the

enhanced matrix synthesis was considered to be a reparative response that attempts to reconstitute the impaired cartilage matrix (2–4). Meanwhile, the up-regulation of cartilage matrix genes was less obvious in the degenerated areas, particularly in the upper regions. Instead, at those regions, the expression of type III collagen and fibronectin was most enhanced. The shift in gene expression was apparent when the profile of gene expression was compared between preserved and degenerated areas in each OA joint (Figure 4).

This shift in gene expression could be significantly involved in the progression of the disease. First, in OA, cartilage matrix is lost primarily from the surface of degenerated cartilage (25), and that loss of matrix could be accelerated by the reduced cartilage matrix synthesis in the surface region (4,9). Second, matrix loss may be facilitated by the induction of type III collagen synthesis. Although this collagen could be a minor component of normal articular cartilage (26–28), it may diminish the quality of cartilage matrix when expressed in excess through the inhibition of proper matrix organization (28,29). Third, fibronectin is known to cause an intense catabolic response in chondrocytes and synoviocytes when cleaved into fragments (30). Therefore, the induction of this protein at the site of enhanced catabolism may be even more significant in the progression of the disease. Taking these findings together, the shift in matrix gene expression at the upper region of degenerated cartilage could be a critical event in OA pathology. Since the shift of gene expression was observed in virtually all OA samples, the regulation of cellular metabolism at that site may be an effective strategy in the future to delay or inhibit disease progression.

Compared with type III collagen and fibronectin, the expression of the other minor cartilaginous genes was less pronounced in OA cartilage in terms of areas, intensities, and frequencies. The induction of type I collagen mRNA was highly variable among OA samples, and, even when expressed, *COL1A1* and *COL1A2* mRNA were often induced at different intensities. The expression of type I collagen in human OA cartilage has remained controversial in previous studies. Although our result of *COL1A1* expression was consistent with several reports (4,6,9), it was discordant with another report regarding the area of expression (31). Further, while we observed the expression of *COL1A2* in human OA cartilage, it was not detected in an earlier study (9). The revealed discrepancy between *COL1A1* and *COL1A2* expression may account for these contradictions in the literature. Likewise, there has been a controversy regarding the induction of *COL10A1* ex-

pression in human OA cartilage: some investigators observed the expression in the upper part of OA cartilage (12,32), whereas others reported it in the deep zone (13,14,33–35). Our result is consistent with the latter finding, in that we identified its expression primarily in the deep zone. However, because the expression of *COL10A1* was relatively weak and fairly inconsistent among OA cartilage samples, we assume that the expression of type X collagen in OA cartilage might be of limited significance in the pathology of OA.

Previously, the appearance of type IIA procollagen mRNA or exon 2 of *COL2A1* in OA cartilage was considered to be the result of a phenotypic reversal of chondrocytes (11,12). However, this speculation is not supported by the present result. Since a result consistent with our own was reported in another recent study (23), a phenotypic reversal of chondrocytes may not be a dominant event in OA cartilage.

In light of these findings, the metabolic change of the chondrocytes in OA may be understood as follows. In the degenerated areas, a major change in the metabolism occurs in the upper region of degenerated cartilage. Such a change resembles that of the dedifferentiation process in the decline of type II collagen and aggrecan expression and the induction of type III collagen expression (Figures 2 and 3) (an illustration of the sequential changes of gene expression in articular chondrocytes during dedifferentiation is available at <http://www.hosp.go.jp/~sagami/rinken/crc/index.html>). However, the change is different from that process in the expression of link protein, fibronectin, and type I collagen genes. Thus, the metabolic change in the degenerated areas of OA cartilage was considered to be unique and not closely related to the one during the dedifferentiation process. Meanwhile, in the preserved areas, the expression of cartilage matrix genes is highly up-regulated. Although the phenotypic deviation is less obvious in those areas, the expression of type I collagen and type X collagen genes may be induced there in the superficial and deep zones, respectively.

Although the mechanism(s) for these metabolic changes remains entirely unknown, the change in *SOX9* expression may be related to the altered chondrocyte metabolism in OA. As shown in the correlation study, the regional difference in matrix gene expression within OA cartilage could be ascribed, at least partly, to the change in *SOX9* expression. However, the present result also indicates that the general up-regulation of matrix gene expression in OA chondrocytes was not associated with the increase in *SOX9* expression. In this study, the amounts of SOX proteins were not assessed. Further-

more, the transcriptional activity of *SOX9* is known to be modulated by the level of phosphorylation (36) and by the presence of coregulators (37,38). Thus, taking these factors into account may provide a better explanation of the significance of SOX proteins in the altered chondrocyte metabolism in OA.

Although the present study has clarified the metabolic change of chondrocytes in OA cartilage, it also has several limitations. First, the metabolic change was evaluated primarily by mRNA expression, and protein synthesis was not determined. The major difference in mRNA expression levels among the samples posed another problem. A large variation among human cartilage samples has been reported repeatedly in previous studies (7,8,23). For OA samples, this might reflect the diversity of the pathology, while the variation among the controls might have stemmed from differences in joint physiology that could be related to the donor's condition before death. These points should be clarified by future studies. Despite these limitations, we believe that our study has revealed several novel aspects of OA pathology. We hope that the current results may offer another clue to eventually establishing a novel strategy to treat this tenacious disease.

AUTHOR CONTRIBUTIONS

Dr. Fukui had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Study design. Fukui.

Acquisition of data. Ikeda, Ohnuki, Tanaka, Hikita, Mitomi, Juji, Katsuragawa, Yamamoto, Sawabe, Yamane, Suzuki.

Analysis and interpretation of data. Fukui, Mori, Sandell, Ochi.

Manuscript preparation. Fukui.

Statistical analysis. Fukui.

REFERENCES

- Collins DH, McElligott TF. Sulphate ($^{35}\text{SO}_4$) uptake by chondrocytes in relation to histological changes in osteoarthritic human articular cartilage. *Ann Rheum Dis* 1960;19:318–30.
- Sandell LJ, Heinegard D, Hering TM. Cell biology, biochemistry, and molecular biology of articular cartilage in osteoarthritis. In: Moskowitz RW, Altman RD, Hochberg MC, Buckwalter JA, Goldberg VM, editors. *Osteoarthritis: diagnosis and medical/surgical management*. 4th ed. Philadelphia: Lippincott Williams & Wilkins; 2007. p. 73–106.
- Aigner T, Gluckert K, von der Mark K. Activation of fibrillar collagen synthesis and phenotypic modulation of chondrocytes in early human osteoarthritic cartilage lesions. *Osteoarthritis Cartilage* 1997;5:183–9.
- Aigner T, Vornehm SI, Zeiler G, Dudhia J, von der Mark K, Bayliss MT. Suppression of cartilage matrix gene expression in upper zone chondrocytes of osteoarthritic cartilage. *Arthritis Rheum* 1997;40:562–9.
- Adam M, Deyl Z. Altered expression of collagen phenotype in osteoarthrosis. *Clin Chim Acta* 1983;133:25–32.

6. Miosge N, Hartmann M, Maelicke C, Herken R. Expression of collagen type I and type II in consecutive stages of human osteoarthritis. *Histochem Cell Biol* 2004;122:229–36.
7. Aigner T, Fundel K, Saas J, Gebhard PM, Haag J, Weiss T, et al. Large-scale gene expression profiling reveals major pathogenetic pathways of cartilage degeneration in osteoarthritis. *Arthritis Rheum* 2006;54:3533–44.
8. Aigner T, Zien A, Gehrsitz A, Gebhard PM, McKenna L. Anabolic and catabolic gene expression pattern analysis in normal versus osteoarthritic cartilage using complementary DNA–array technology. *Arthritis Rheum* 2001;44:2777–89.
9. Aigner T, Bertling W, Stoss H, Weseloh G, von der Mark K. Independent expression of fibril-forming collagens I, II, and III in chondrocytes of human osteoarthritic cartilage. *J Clin Invest* 1993;91:829–37.
10. Nerlich AG, Wiest I, von der Mark K. Immunohistochemical analysis of interstitial collagens in cartilage of different stages of osteoarthrosis. *Virchows Arch B Cell Pathol Incl Mol Pathol* 1993;63:249–55.
11. Aigner T, Zhu Y, Chansky HH, Matsen FA III, Maloney WJ, Sandell LJ. Reexpression of type IIA procollagen by adult articular chondrocytes in osteoarthritic cartilage. *Arthritis Rheum* 1999;42:1443–50.
12. Nah HD, Swoboda B, Birk DE, Kirsch T. Type IIA procollagen: expression in developing chicken limb cartilage and human osteoarthritic articular cartilage. *Dev Dyn* 2001;220:307–22.
13. Girkontaite I, Frischholz S, Lammi P, Wagner K, Swoboda B, Aigner T, et al. Immunolocalization of type X collagen in normal fetal and adult osteoarthritic cartilage with monoclonal antibodies. *Matrix Biol* 1996;15:231–8.
14. Von der Mark K, Kirsch T, Nerlich A, Kuss A, Weseloh G, Gluckert K, et al. Type X collagen synthesis in human osteoarthritic cartilage: indication of chondrocyte hypertrophy. *Arthritis Rheum* 1992;35:806–11.
15. Poole RA, Guilak F, Abramson SB. Etiopathogenesis of osteoarthritis. In: Moskowitz RW, Altman RD, Hochberg MC, Buckwalter JA, Goldberg VM, editors. *Osteoarthritis: diagnosis and medical/surgical management*. 4th ed. Philadelphia: Lippincott Williams & Wilkins; 2007. p. 27–49.
16. Emmert-Buck MR, Bonner RF, Smith PD, Chuaqui RF, Zhuang Z, Goldstein SR, et al. Laser capture microdissection. *Science* 1996;274:998–1001.
17. Altman R, Asch E, Bloch D, Bole G, Borenstein D, Brandt K, et al. Development of criteria for the classification and reporting of osteoarthritis: classification of osteoarthritis of the knee. *Arthritis Rheum* 1986;29:1039–49.
18. Buckwalter JA, Mankin HJ, Grodzinsky AJ. Articular cartilage and osteoarthritis. In: Pellegrini VD Jr, editor. *AAOS instructional course lectures*. Vol. 54. Rosemont (IL): American Academy of Orthopedic Surgeons; 2005. p. 465–80.
19. Poole RA. Cartilage in health and disease. In: Koopman WJ, Moreland LW, editors. *Arthritis and allied conditions: a textbook of rheumatology*. 15th ed. Philadelphia: Lippincott Williams & Wilkins; 2004. p. 223–69.
20. Miller DR, Mankin HJ, Shoji H, D'Ambrosia RD. Identification of fibronectin in preparations of osteoarthritic human cartilage. *Connect Tissue Res* 1984;12:267–75.
21. Goldwasser M, Astley T, van der Rest M, Glorieux FH. Analysis of the type of collagen present in osteoarthritic human cartilage. *Clin Orthop Relat Res* 1982;(167):296–302.
22. Burton-Wurster N, Horn VJ, Lust G. Immunohistochemical localization of fibronectin and chondronectin in canine articular cartilage. *J Histochem Cytochem* 1988;36:581–8.
23. Gebhard PM, Gehrsitz A, Bau B, Soder S, Eger W, Aigner T. Quantification of expression levels of cellular differentiation markers does not support a general shift in the cellular phenotype of osteoarthritic chondrocytes. *J Orthop Res* 2003;21:96–101.
24. De Crombrughe B, Lefebvre V, Behringer RR, Bi W, Murakami S, Huang W. Transcriptional mechanisms of chondrocyte differentiation. *Matrix Biol* 2000;19:389–94.
25. Hollander AP, Pidoux I, Reiner A, Rorabeck C, Bourne R, Poole AR. Damage to type II collagen in aging and osteoarthritis starts at the articular surface, originates around chondrocytes, and extends into the cartilage with progressive degeneration. *J Clin Invest* 1995;96:2859–69.
26. Young RD, Lawrence PA, Duance VC, Aigner T, Monaghan P. Immunolocalization of type III collagen in human articular cartilage prepared by high-pressure cryofixation, freeze-substitution, and low-temperature embedding. *J Histochem Cytochem* 1995;43:421–7.
27. Wotton SF, Duance VC. Type III collagen in normal human articular cartilage. *Histochem J* 1994;26:412–6.
28. Eyre DR, Weis MA, Wu JJ. Articular cartilage collagen: an irreplaceable framework? *Eur Cell Mater* 2006;12:57–63.
29. Young RD, Lawrence PA, Duance VC, Aigner T, Monaghan P. Immunolocalization of collagen types II and III in single fibrils of human articular cartilage. *J Histochem Cytochem* 2000;48:423–32.
30. Homandberg GA. Cartilage damage by matrix degradation products: fibronectin fragments. *Clin Orthop Relat Res* 2001;(391 Suppl):S100–7.
31. Wagner S, Hofstetter W, Chiquet M, Mainil-Varlet P, Stauffer E, Ganz R, et al. Early osteoarthritic changes of human femoral head cartilage subsequent to femoro-acetabular impingement. *Osteoarthritis Cartilage* 2003;11:508–18.
32. Kirsch T, Swoboda B, Nah H. Activation of annexin II and V expression, terminal differentiation, mineralization and apoptosis in human osteoarthritic cartilage. *Osteoarthritis Cartilage* 2000;8:294–302.
33. Gibson GJ, Verner JJ, Nelson FR, Lin DL. Degradation of the cartilage collagen matrix associated with changes in chondrocytes in osteoarthrosis: assessment by loss of background fluorescence and immunodetection of matrix components. *J Orthop Res* 2001;19:33–42.
34. Boos N, Nerlich AG, Wiest I, von der Mark K, Ganz R, Aebi M. Immunohistochemical analysis of type-X-collagen expression in osteoarthritis of the hip joint. *J Orthop Res* 1999;17:495–502.
35. Aigner T, Reichenberger E, Bertling W, Kirsch T, Stoss H, von der Mark K. Type X collagen expression in osteoarthritic and rheumatoid articular cartilage. *Virchows Arch B Cell Pathol Incl Mol Pathol* 1993;63:205–11.
36. Huang W, Zhou X, Lefebvre V, de Crombrughe B. Phosphorylation of SOX9 by cyclic AMP-dependent protein kinase A enhances SOX9's ability to transactivate a Col2a1 chondrocyte-specific enhancer [published erratum appears in *Mol Cell Biol* 2000;20:7838]. *Mol Cell Biol* 2000;20:4149–58.
37. Tsuda M, Takahashi S, Takahashi Y, Asahara H. Transcriptional co-activators CREB-binding protein and p300 regulate chondrocyte-specific gene expression via association with Sox9. *J Biol Chem* 2003;278:27224–9.
38. Kawakami Y, Tsuda M, Takahashi S, Taniguchi N, Esteban CR, Zemmyo M, et al. Transcriptional coactivator PGC-1 α regulates chondrogenesis via association with Sox9. *Proc Natl Acad Sci U S A* 2005;102:2414–9.

Research

Open Access

Proinflammatory role of amphiregulin, an epidermal growth factor family member whose expression is augmented in rheumatoid arthritis patients

Shoji Yamane*^{1,2}, Satoru Ishida^{1,2}, Yukie Hanamoto¹, Ken-ichi Kumagai¹, Riako Masuda¹, Konagi Tanaka¹, Noriyuki Shiobara¹, Noriko Yamane², Toshihito Mori¹, Takuo Juji¹, Naoshi Fukui¹, Tsunetoshi Itoh³, Takahiro Ochi¹ and Ryuji Suzuki¹

Address: ¹Clinical Research Center for Allergy and Rheumatology, National Hospital Organization, Sagamihara National Hospital, Sakuradai 18-1, Sagamihara, Kanagawa 228-8522, Japan, ²Discovery Research Laboratories, Shionogi & Co., Ltd., 3-1-1, Futaba-cho, Toyonaka, Osaka 561-0825, Japan and ³Department of Immunology and Embryology, Tohoku University School of Medicine, 2-1 Seiry-Machi, Aoba-ku, Sendai 980-8575, Japan

Email: Shoji Yamane* - shoji.yamane@shionogi.co.jp; Satoru Ishida - s-ishida@sagamihara-hosp.gr.jp; Yukie Hanamoto - y-hanamoto@sagamihara-hosp.gr.jp; Ken-ichi Kumagai - k-kumagai@sagamihara-hosp.gr.jp; Riako Masuda - r-masuda@sagamihara-hosp.gr.jp; Konagi Tanaka - k-tanaka@sagamihara-hosp.gr.jp; Noriyuki Shiobara - n-shiobara@sagamihara-hosp.gr.jp; Noriko Yamane - noriko.yamane@shionogi.co.jp; Toshihito Mori - t-mori@sagamihara-hosp.gr.jp; Takuo Juji - t-juji@sagamihara-hosp.gr.jp; Naoshi Fukui - n-fukui@sagamihara-hosp.gr.jp; Tsunetoshi Itoh - itoh@immem.med.tohoku.ac.jp; Takahiro Ochi - r-suzuki@sagamihara-hosp.gr.jp; Ryuji Suzuki - t-ochi@sagamihara-hosp.gr.jp

* Corresponding author

Published: 27 April 2008

Received: 11 October 2007

Journal of Inflammation 2008, 5:5 doi:10.1186/1476-9255-5-5

Accepted: 27 April 2008

This article is available from: <http://www.journal-inflammation.com/content/5/1/5>

© 2008 Yamane et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: The epidermal growth factor (EGF) and EGF receptor (EGFR) families play important roles in the hyperplastic growth of several tissues as well as tumor growth. Since synovial hyperplasia in rheumatoid arthritis (RA) resembles a tumor, involvement of the EGF/EGFR families in RA pathology has been implied. Although several reports have suggested that ErbB2 is the most important member of the EGFR family for the synovitis in RA, it remains unclear which members of the EGF family are involved. To clarify the EGF-like growth factors involved in the pathology of RA, we investigated the expression levels of seven major EGF-like growth factors in RA patients compared with those in osteoarthritis (OA) patients and healthy control subjects.

Methods: The expression levels of seven EGF-like growth factors and four EGFR-like receptors were measured in mononuclear cells isolated from bone marrow and venous blood, as well as in synovial tissues, using quantitative RT-PCR. Further evidence of gene expression was obtained by ELISAs. The proinflammatory roles were assessed by the growth-promoting and cytokine-inducing effects of the corresponding recombinant proteins on cultured fibroblast-like synoviocytes (FLS).

Results: Among the seven EGF-like ligands examined, only amphiregulin (AREG) was expressed at higher levels in all three RA tissues tested compared with the levels in OA tissues. The AREG protein concentration in RA synovial fluid was also higher than that in OA synovial fluid. Furthermore, recombinant human AREG stimulated FLS to proliferate and produce several proinflammatory cytokines, including angiogenic cytokines such as interleukin-8 and vascular

endothelial growth factor (VEGF), in a dose-dependent manner. The VEGF mRNA levels in RA synovia and VEGF protein concentrations in RA synovial fluid were significantly higher than those in the corresponding OA samples and highly correlated with the levels of AREG.

Conclusion: The present findings suggest that AREG functions to stimulate synovial cells and that elevated levels of AREG may be involved in the pathogenesis of RA.

Background

Rheumatoid arthritis (RA) is a chronic inflammatory disease that is mainly characterized by synovial hyperplasia and progressive destruction of the affected joints. Activated synoviocytes in the hypertrophic synovia induce angiogenesis, and play pivotal roles in the recruitment and differentiation of inflammatory cells. However, the driving force of the synovial hyperplasia remains obscure.

The granulomatous tissues of RA synovia, referred to as pannuses, resemble tumors. Cultured fibroblast-like synoviocytes (FLS) from these pannuses share some features with transformed cells, *i.e.* anchorage-independent growth [1,2] and downregulation of tumor suppressors [3-5]. Similar to transformed cells, tyrosine-phosphorylated proteins are augmented in RA-FLS, and several growth factors whose receptors possess tyrosine kinase activities have been reported to promote the tumor-like behavior of RA synovial membranes [6-9]. Since platelet-derived growth factor (PDGF) and fibroblast growth factor (FGF) stimulate DNA synthesis and proliferation of FLS cultured in medium containing low concentrations of serum [10] and histochemical studies have revealed upregulated expression levels of PDGF and FGF and their receptors in RA synovial tissues [11-13], these molecules are considered to be the major contributors to synovial hyperplasia [2,14].

On the other hand, the proto-oncogene *c-erb-B*, referred to as epidermal growth factor (EGF) receptor (EGFR), is a well-known tyrosine kinase growth factor receptor. Four members of the EGFR family have been identified to date, namely *c-erb-B/EGFR* and its related products ErbB2, ErbB3 and ErbB4. The family members form homodimers or heterodimers in various combinations, and exhibit different ligand specificities for the 13 members of the EGF family [15]. Although expression of ErbB2, but not the other ErbB-related receptors, has been reported to be augmented in RA synovial tissues [7,16,17], it remains unknown which members of the EGF family are expressed in the affected joints and involved in the pathology of RA.

In previous studies, we investigated the involvement of bone marrow in the pathology of RA. An increase in myeloid cells expressing abnormal surface antigens in bone marrow was associated with the severity of RA [18-23]. Pathogenic synovial fibroblasts may be derived from bone

marrow CD34⁺ cells in RA [24]. Recently, we identified RA-associated genes in bone marrow cells using a cDNA subtraction technique [25]. In that report, we demonstrated that two EGF-like growth factors, amphiregulin (AREG) and epiregulin (EREG), were upregulated in RA bone marrow.

In the present study, we examined the extents of involvement of EGF family members in RA pathology by investigating the expression of seven major EGF-like growth factors, namely EGF, AREG, EREG, transforming growth factor α (TGF α), heparin-binding EGF-like growth factor (HB-EGF), betacellulin (BTC) and neuregulin-1 (NRG1), in synovial tissues and mononuclear cells isolated from bone marrow and venous blood. The results revealed that AREG expression was augmented in all RA tissues and cells examined. Moreover, the AREG protein concentration in RA synovial fluid was significantly higher than that in osteoarthritis (OA) synovial fluid. Recombinant human AREG stimulated RA-FLS to proliferate and express several proinflammatory cytokines. These findings suggest that AREG may play a role in the pathogenesis of RA.

Methods

Patients and samples

Bone marrow fluid, venous blood and/or synovial tissues were intraoperatively obtained from 15 RA patients (all women; mean age \pm SD: 59.3 \pm 8.7 years) and 12 OA patients (all women; mean age \pm SD: 64.5 \pm 11.8 years) undergoing joint arthroplasty. None of the patients had taken any medication for at least 1 week before the operation. The RA and OA patients fulfilled the 1987 revised criteria of the American College of Rheumatology for the classification of RA [26] and the diagnostic criteria for OA [27], respectively. Bone marrow fluid and venous blood were mixed with heparin and separated by centrifugation at 1700 g for 15 min. After removal of the plasma, the blood cells and bone marrow cell fractions were adjusted to their original volumes with Hank's balanced salt solution (HBSS) and fractionated by density-gradient centrifugation at 3000 g for 30 min on Ficoll-Hypaque (GE Healthcare Bioscience, Tokyo, JPN). Mononuclear cells were collected from both the bone marrow and peripheral blood and used for the experiments described below. For further separation, the collected mononuclear cells were fractionated by magnetic beads coated with immobilized

CD14, CD3 or CD19 antibodies (Miltenyi Biotec, Tokyo, JPN), since CD14, CD3 and CD19 are lineage-specific markers for monocytes, T lymphocytes and B lymphocytes, respectively. The cell populations fractionated by these antibodies were measured using flow cytometry, and confirmed to be > 95% pure. Synovial fluid was obtained from 24 RA patients and 10 OA patients and venous blood was obtained from 57 RA patients and 12 OA patients attending the outpatient clinic of our hospital. Synovial fluid was separated from cells and debris by centrifugation, and the clear supernatant was collected. Plasma was collected by centrifuging heparinized blood as described above. The synovial fluid and plasma samples were analyzed by ELISAs. All patients and healthy volunteers provided informed consent for participation in the study, which was approved by the Ethical Committee of the National Hospital Organization, Sagami National Hospital.

Isolation of FLS and establishment of cell lines

Synovial membranes were minced aseptically and then digested enzymatically with 1 mg/ml collagenase (Wako, Osaka, JPN) in Dulbecco's modified Eagle's medium (DMEM; GIBCO, Grand Island, NY, USA) for 2 h at 37°C. Single cell suspensions were filtered through a nylon mesh, seeded in culture dishes containing DMEM supplemented with 100 units/ml penicillin, 0.1 mg/ml streptomycin (GIBCO) and 10% fetal bovine serum (FBS; Hyclone, Logan, UT, USA), and cultured at 37°C in humidified air containing 7.5% CO₂. Since freshly isolated FLS contain many lymphocytes, monocytes and granulocytes, we used homogeneous fibroblastic cell populations after more than 4 passages. In proliferation assays, FLS cultures were stimulated by various concentrations of recombinant human AREG (R&D Systems, Tokyo, JPN) for 2 days. Prior to cell harvesting onto glass fiber disks, FLS were cultured with ³H-thymidine for 18 h. The radioactivities on the disks were measured using a liquid scintillation counter.

RNA extraction and cDNA synthesis

Total cellular RNAs were extracted using the TRIZOL™ reagent (Invitrogen, Tokyo, JPN) according to the manufacturer's instructions. For RNA extraction from synovia, minced tissues were homogenized in TRIZOL using a Polytron homogenizer and the extracted RNAs were further purified using an RNeasy micro kit (QIAGEN, Tokyo, JPN). In cytokine induction assays, FLS cultures were stimulated by various concentrations of recombinant human AREG and/or genistein (SIGMA, Tokyo, JPN) for 3 h and subjected to RNA extraction using the RNeasy micro kit. First-strand cDNAs were synthesized from 2 µg of total RNAs by priming with oligo dT and Omniscript™ reverse transcriptase (QIAGEN) according to the manufacturer's instructions.

Quantitative RT-PCR

Using real-time PCR, we estimated the mRNA expression levels of four EGFR family members and seven EGF family members. In subsequent investigations, the mRNA expression levels of five proinflammatory cytokines, namely interleukin (IL)-1β, IL-6, IL-8, tumor necrosis factor-α (TNF-α) and granulocyte-macrophage colony-stimulating factor (GM-CSF) were measured. The mRNA expression levels of vascular endothelium growth factor (VEGF), platelet-derived growth factor (PDGF), basic fibroblast growth factor (bFGF), a disintegrin and metalloproteinase (ADAM) 10 and ADAM17 were also measured. The primer sequences used were: 5'-GTGATTCCATCATGTATCCCAGGAG-3', 5'-AGATGCCTGTCCATGCAAACA-3' (EREG); 5'-CTTCACTGTGTGGTGGCAGATG-3', 5'-ATGCAGTAATGCTTGTATTGCTTGG-3' (BTC); 5'-CAAC-CAGTGGCTGGTGGAGGA-3', 5'-GAGCCCTTATCACTGGATACTGGAA-3' (EGF); 5'-GTGGTGGCTGTCCGCTCTTGATACTC-3', 5'-TCAAATCCATCAGCACTGTGGTC-3' (AREG); 5'-GGCCATGACTAAT-TCCCCTGA-3', 5'-GCCCAATCCTAGACGGCAAC-3' (HB-EGF); 5'-AGATAGACAGCAGCCAAACCCTGA-3', 5'-CTAGGGCCATCTCTGCCATC-3' (TGFα); 5'-AGAATGTGCCATGAAAGTCCAA-3', 5'-GCAGATGCCGGTTATGGTCAG-3' (NRG1); 5'-GGTGGCAATGACAGTAGCATTATGA-3', 5'-AAAGGTGGGCTCCTAAGTAGCTGAA-3' (EGFR); 5'-CAGGCACCGCAGCTCATCTA-3', 5'-TCCCAGGTCACCATCAAATACATC-3' (ErbB2); 5'-CCCAGCATCTGAGCAAGGGTA-3', 5'-TTTAGGCGGGCATAATGGACA-3' (ErbB3); 5'-TGATAGGCCGTTGGTTGTCTGA-3', 5'-CCAGGTAGACATACCCAATCCACTG-3' (ErbB4); 5'-CCCCTGAGGAGTCCAACAT-3', 5'-AAATGCTTTCTC-CGCTCTGA-3' (VEGF); 5'-CTCTGATCATGCTAATGGCTCGA-3', 5'-GCTGCAGTTAGCGTCTCATGTGT-3' (ADAM10); 5'-GTGACATGAATGGCAAATGTGAG-3', 5'-AGACCCAACGATGTTGTCTGCTA-3' (ADAM17); 5'-CCCCTGCCATTCCGGAGGAAGAG-3', 5'-TTGGCCACCTTGACGCTGCGGTG-3' (PDGF); 5'-GTTGTGACAAC-CACAAGCAC-3', 5'-CTCTCACACTATCCACTGGT-3' (bFGF); 5'-ACACTGCGCCAACACAGAAATTA-3', 5'-TTT-GCTTGAAGTTCACTGGCAGC-3' (IL-8); 5'-AAGCCA-GAGCTGTGCAGATGAGTA-3', 5'-TGTCCTGCAGCCACTGGTTC-3' (IL-6); 5'-CCAGGGACAGGATATGGAGCA-3', 5'-TTCAACACCCAGGACAGGTACAG-3' (IL-1β); 5'-CATGTGGCCAGCCACTACAA-3', 5'-ACTGGCTCCCAG-CAGTCAAAG-3' (GM-CSF); 5'-GACAAGCCTGTAGCCCATGTTGTA-3', 5'-CAGCCTTGGCCCTTGAAGA-3' (TNF-α). Real-time PCR was performed using a Light-Cycler 2.0 (Roche Diagnostics, Tokyo, JPN) and SYBR Premix Ex Taq (Takara, Kyoto, JPN) following the manufacturers' protocols. The amounts of PCR products were

assessed by the fluorescence of SYBR Green intercalated in the DNA fragments, and melting curves were routinely recorded to verify the singularity of the products. The amplified products using each primer pair were cloned into the pGEM-T vector (Promega, Tokyo, JPN) and plasmids linearized by enzymatic digestion were used as quantification standards. A reference cDNA was used in every assay to control the precision among assays. The cDNA levels among the samples were normalized by the expression level of the internal control gene GAPDH (5'-GCACCGTCAAGGCTGAGAAC-3', 5'-ATGGTGGTGAAGACGCCAGT-3').

ELISAs

The AREG protein concentrations in plasma samples from 57 RA patients, 12 OA patients and 9 healthy volunteers and synovial fluid samples from 24 RA patients and 8 OA patients were determined using an AREG Duo-set ELISA kit (R&D Systems). The protein concentrations of IL-1 β , IL-6, IL-8, TNF- α , GM-CSF and VEGF in culture supernatants of RA-FLS stimulated with recombinant human AREG (R&D Systems) and those of VEGF and IL-8 in synovial fluid samples from 9 RA patients and 7 OA patients were determined using a Quantikine ELISA kit (R&D Systems).

Statistical analysis

Statistical analysis was carried out using the StatView statistical analysis software (SAS, Cary, NC, USA). Differences between RA specimens and controls were determined to be significant when $P < 0.05$ by the Mann-Whitney U-test. The effects of AREG on RA-FLS were analysed by the Mann-Whitney U-test following the Kruskal-Wallis test. Correlation coefficients (ρ) were calculated by Spearman's rank correlation method and tested for statistical significance at the 0.05 (two-tailed) level.

Results

Expression profiles of EGF family members in bone marrow mononuclear cells (BMMCs)

First, we determined the mRNA expression levels of seven EGF family members in BMMCs obtained from 9 RA patients and 10 OA patients (Fig. 1A). EREG was the most abundantly expressed, and its mRNA level in RA-BMMCs was significantly higher than that in OA-BMMCs ($P = 0.0060$). The expression levels of AREG, TGF α and EGF were about 10-fold lower than that of EREG in OA-BMMCs, but significantly upregulated in RA-BMMCs ($P = 0.0258$, $P = 0.000045$ and $P = 0.0140$, respectively). Although the expression of HB-EGF was the next most abundant in RA-BMMCs, there was no significant difference between its expression levels in RA- and OA-BMMCs. The BTC and NRG1 mRNA expression levels were almost undetectable in both RA- and OA-BMMCs.

Expression profiles of EGF family members in peripheral blood mononuclear cells (PBMCs)

Next, we determined the mRNA expression levels of the seven EGF family members in PBMCs obtained from 8 RA patients and 7 OA patients (Fig. 1B). EREG, TGF α and EGF were highly abundantly expressed in PBMCs, and their mRNA levels in RA- and OA-PBMCs did not differ. The expression levels of AREG, HB-EGF and NRG1 were about 10-fold lower than the levels of the highly abundant members in OA-PBMCs. Although AREG and HB-EGF were markedly upregulated in RA-PBMCs ($P = 0.0017$ and $P = 0.0367$, respectively), NRG1 was not upregulated in RA-PBMCs. BTC was not detected in either type of PBMCs.

Expression profiles of EGF-like growth factors in synovial tissues

Next, we determined the mRNA expression levels of the seven EGF family members in synovial tissues obtained from 14 RA patients and 5 OA patients (Fig. 1C). Although HB-EGF was the most abundantly expressed and EREG was the next most abundantly expressed in synovial tissues from both RA and OA joints, their mRNA levels in RA synovia were somewhat lower than those in OA synovia. On the other hand, AREG expression, which was 1000-fold lower than HB-EGF expression in OA synovia, was markedly upregulated in RA synovial tissues ($P = 0.0110$). Expression of EGF, BTC or NRG1 was not detected in either OA or RA joints. Since only AREG expression was augmented in BMMCs, PBMCs and synovia of RA patients compared with the levels in control samples among the seven EGF-related growth factors examined, we narrowed the focus of the study to AREG.

Determination of plasma and synovial fluid concentrations of AREG

To confirm whether the protein concentration of AREG was augmented in RA patients, the AREG concentrations in plasma and synovial fluid samples were examined by ELISA. As shown in Fig. 2A, there were no significant differences among the AREG protein concentrations in the RA, OA and healthy control (HC) plasma samples, whereas the AREG concentration in RA synovial fluid samples was significantly higher than that in OA synovial fluid samples. Evaluation of BMMC, PBMC and plasma samples from 5 RA patients revealed that the AREG mRNA levels in RA-PBMCs were highly correlated with those in RA-BMMCs (data not shown), but not correlated with the plasma concentrations of this protein (Fig. 2B). Besides transformed cells, AREG-producing cells were previously reported to be activated monocytes [28] and activated T lymphocytes [29]. To clarify which lineage of blood cells expressed AREG in RA, PBMCs from RA patients were fractionated using magnetic beads coated with immobilized CD14, CD3 or CD19 antibodies. As shown in Fig. 2C,




Figure 1
mRNA expression levels of EGF-related growth factors in BMMCs (A), PBMCs (B) and synovial tissues (C). The results of real-time PCR are shown as box-plots. The log ratio of the mRNA quantities relative to the total RNA amount (A) or GAPDH mRNA (B, C) is plotted on the y-axis of each graph. The upper and lower error bars indicate the 90th and 10th percentiles, respectively. The upper and lower edges of each box indicate the 75th and 25th percentiles, respectively, and the line inside the box shows the median. Genes not detected are shown as ND. The differences between the mRNA levels in the RA and control samples were analyzed by the Mann-Whitney U-test, and significant differences are shown by asterisks (*P < 0.05; **P < 0.01). RA: samples from RA patients; OA: samples from OA patients.




Figure 2

Amphiregulin expression in peripheral blood and synovial fluid samples. (A) The concentrations of AREG in plasma and synovial fluid samples are plotted as log-values on the y-axis of box-plots. Significant differences are shown by asterisks (** $P < 0.01$). RA: samples from RA patients; OA: samples from OA patients; HC: samples from healthy volunteers. (B) The AREG mRNA expression levels in PBMCs and AREG protein concentrations in plasma are shown. Venous blood samples from 6 RA patients and 6 healthy volunteers (HC) were separated into plasma and PBMCs. Total RNAs were extracted from PBMCs and subjected to cDNA synthesis. The AREG mRNA levels were measured by real-time PCR and normalized by the GAPDH mRNA levels. The relative AREG mRNA level relative to the GAPDH mRNA level is plotted as the log ratio on the primary y-axis (left), while the plasma concentration of AREG protein measured by ELISA is plotted as the log value on the secondary y-axis (right). The correlation coefficient (ρ) of the protein level in plasma to the mRNA level in PBMCs is -0.378 ($P = 0.2104$). (C) PBMCs from 4 RA patients and 2 HCs were separated by CD14 microbeads, and the AREG mRNA level in each fraction was measured by real-time PCR. The log ratio of the AREG mRNA level relative to the GAPDH mRNA level is plotted on the y-axis.

both CD14-positive and CD14-negative fractions of RA-PBMCs expressed equal amounts of AREG mRNA, and their levels were markedly higher than that in control PBMCs. The CD3 and CD19 separations yielded similar results (data not shown).

Effects of AREG on the proliferation of RA-FLS

To investigate the biological activity of AREG in joints affected by RA, we assessed the effects of recombinant human AREG on RA-FLS. Since AREG is a member of the EGF-like growth factor family, its growth-promoting activity was measured first. As shown in Fig. 3A, recombinant human AREG enhanced *de novo* DNA synthesis by RA-FLS in a dose-dependent manner. Fig. 3B shows the expression levels of the four EGFR family members in the cell lines used in the proliferation assay. In all FLS cell lines, ErbB2 and EGFR were the predominantly expressed receptors and ErbB3 and ErbB4 were expressed at about 100-fold lower levels than the most abundant ErbB2 level. There were no differences among the three RA-FLS lines and the one OA-FLS line. Although the amounts of radioactivity incorporated into the RA-FLS lines were higher than that incorporated into the OA-FLS line, the issue of whether RA-FLS are more sensitive to AREG than OA-FLS requires further examination. EGF-like growth factors are expressed as transmembrane-type precursors, and ectodomain shedding by ADAMs is essential for their effects as well as the expression of their receptors [30,31]. Since ADAM10 and ADAM17 are known to be sheddases for EGF-like growth factors, we measured the expression levels of the four EGFR-like receptors and ADAM10 and ADAM17 in synovial tissues from 10 RA patients and 6 OA patients (Fig. 3C). Similar to the findings for FLS, EGFR and ErbB2 were the predominantly expressed receptors in synovial tissues and their expression levels were not augmented in RA samples. Although AREG is supposed to be mainly processed by ADAM17, ADAM17 was expressed at a lower level than ADAM10, and neither ADAM10 nor ADAM17 was upregulated in RA synovia.

Effects of AREG on cytokine production by RA-FLS

Next, we tested the expression levels of five proinflammatory cytokines (IL-1 β , TNF- α , IL-8, GM-CSF and IL-6) in RA-FLS stimulated by recombinant AREG. To clarify whether the recombinant AREG stimulated RA-FLS to proliferate via the induction of other growth factors, we also tested the expression levels of PDGF, bFGF and VEGF, which are involved in synovial hyperplasia. Recombinant AREG upregulated the mRNA expression levels of VEGF, IL-8, GM-CSF and IL-6 (Fig. 4A), but not those of PDGF, bFGF, IL-1 β or TNF- α (data not shown). Recombinant AREG stimulated RA-FLS to express these cytokines in a dose-dependent manner, and the EGFR-tyrosine kinase inhibitor genistein suppressed the AREG-dependent expression in a dose-dependent manner. ELISA analysis

revealed elevated levels of VEGF, IL-8, GM-CSF and IL-6 proteins in culture supernatants of AREG-stimulated RA-FLS (Fig. 4B), consistent with the results of the real-time PCR. We also tested the expression levels of ADAM10 and ADAM17 in AREG-stimulated RA-FLS. Contrary to the effect on the cytokine induction, AREG downregulated the expression of ADAM17 in a dose-dependent manner, and the AREG-dependent suppression was abolished by genistein (Fig. 4C). Analyses of ADAM10 expression produced similar results to those for ADAM17 (data not shown).

Correlation between VEGF and AREG expression levels

Since higher inductions of IL-6, IL-8 and GM-CSF than those induced by AREG have been observed and induction of VEGF has not yet been observed in our previous studies [32,33], we hypothesized that AREG would be closely related to VEGF in RA joints. To examine the relationship of AREG with this angiogenic factor in affected joints, VEGF expression was assessed in synovial tissues from 10 RA patients and 6 OA patients. Fig. 5A shows the mRNA levels of VEGF measured by real-time PCR. Ikeda et al. reported that the VEGF₁₆₅ transcript may be augmented in RA synovial tissues, and that the products of this transcript may be associated with RA pathology [34]. The primers for the real-time PCR amplification of VEGF used in our study were also designed to detect VEGF₁₆₅. VEGF expression in RA synovia was significantly higher than that in OA synovia (Fig. 5A, left panel), and highly correlated with AREG expression (Fig. 5A, right panel). Fig. 5B shows the VEGF protein concentrations in synovial fluid samples measured by ELISA. The ELISA system is able to detect all isoforms of VEGF-A, although it was designed for VEGF₁₆₅. Consistent with the results of the mRNA expression analyses, the VEGF protein levels in RA synovial fluid samples were significantly higher than those in OA synovial fluid samples (Fig. 5B, left panel), and highly correlated with the AREG concentration (Fig. 5B, right panel).

Discussion

Several previous studies have reported the involvement of c-erb-B family members, especially ErbB2, in the pathology of RA. Hallbech et al. showed immunohistochemically that the expression levels of ErbB2 and TGF α were augmented in RA synovia [17]. Satoh et al. demonstrated that ErbB2 was predominant in RA synovia and primary RA-FLS, but not in OA synovia or primary OA-FLS, and that a neutralizing antibody against ErbB2 suppressed the proliferation of primary RA-FLS, but not primary OA-FLS [16]. In the present study, we investigated which members of the ErbB family are predominantly expressed in RA-FLS and RA synovial tissues. Among the four ErbB family members, the mRNA level of ErbB2 was the highest, followed by that of EGFR, while the others were expressed at




Figure 3

Stimulatory activity of AREG on the proliferation of RA-FLS. (A) Effect of AREG on the proliferation of RA-FLS.

Three RA-FLS lines and one OA-FLS line were cultured with the indicated concentrations of AREG. After 24 h of stimulation, the cells were labeled with ³H-thymidine for 18 h and then harvested on glass filters with a cell harvester. The incorporated radioactivity was measured by liquid scintillation counting. The values are shown as the means ± SD of three independent experiments. Significant differences are shown by asterisks (*P < 0.05; **P < 0.01). (B) Expression profiles of EGFR family members in FLS. cDNA samples of the four FLS lines used in the proliferation assay were subjected to real-time PCR analysis. (C) Expression profiles of the receptors and sheddases of the EGF family in synovia. cDNAs of synovial tissues from 10 RA patients and 6 OA patients were subjected to real-time PCR analysis.




Figure 4
Stimulatory activity of AREG on cytokine production by RA-FLS. (A) Effects of AREG on cytokine expression in RA-FLS. Four RA-FLS lines were cultured with the indicated concentrations of recombinant human AREG and/or genistein. After 4 h of stimulation, total RNAs were extracted and the mRNA levels of PDGF, bFGF, VEGF, IL-1 β , IL-6, IL-8, TNF- α and GM-CSF were measured by real-time PCR. The results for VEGF, IL-8, GM-CSF and IL-6 are shown. The results for the other molecules were omitted from the figure, since AREG had no effect on their expressions. (B) Effects of AREG on cytokine production by RA-FLS. Four RA-FLS lines were cultured with the indicated concentrations of AREG for 24 h. The GM-CSF, IL-6, IL-8 and VEGF concentrations in the supernatants were measured by ELISA, and are shown as means \pm SD. (C) Effects of AREG on the expression of sheddases. The same cDNA samples used in panel A were subjected to real-time PCR analysis for ADAM10 and ADAM17. The results for ADAM10 were omitted from the figure, since they were similar to those for ADAM17. Each panel shows a representative result of three independent experiments. Significant differences from unstimulated cells are shown by asterisks (* $P < 0.05$; ** $P < 0.01$).

almost undetectable levels in FLS and synovial membranes. While these results are consistent with the previous report [16], there were no differences in the expression levels between RA and OA samples. This discrepancy may reflect differences in the synovial specimens, although it will be necessary to confirm this hypothesis by assessing the ErbB2 protein concentrations in RA and OA samples. The OA synovial samples used in the present study were obtained from synovia with villous formation, rather than from the joint capsule, and thus our OA samples may be more activated than those used in the previous study. In any case, ErbB2 and EGFR were confirmed to be predominantly expressed in RA-FLS and RA synovia.

On the other hand, there have been very few reports of the expression profiles of EGF-like growth factors in RA synovia. In the present study, we found that AREG expression, which was 1000-fold lower than the most abundantly expressed HB-EGF in OA synovia, was markedly upregulated in RA synovia. Since it was correlated with the expression of AREG ($\rho = 0.532$, $P = 0.0241$), the expression of TGF α may tend to be augmented in RA synovia, as reported previously [17].

Since we recently reported augmented expression of AREG in BMMCs and PBMCs from RA patients [25], we examined the expression levels of other EGF family members in the present study. In RA-BMMCs, EGF and TGF α were also




Figure 5
Correlation between AREG and VEGF expression levels. (A) The VEGF mRNA expression levels in synovia from 10 RA patients and 6 OA patients were measured by real-time PCR, and are shown by box-plots (left). Significant differences from unstimulated cells are shown by asterisks (* $P < 0.05$). The correlation between the mRNA levels of VEGF and AREG is shown by a distribution chart (right). The linear regression coefficient (R^2) is 0.496 and the correlation coefficient (ρ) is 0.700 ($P = 0.0067$). (B) The concentrations of VEGF in synovial fluids from 7 RA patients and 6 OA patients were measured by ELISA, and are shown by box-plots (left). Significant differences from unstimulated cells are shown by asterisks (* $P < 0.05$). The correlation between the protein concentrations of VEGF and AREG is shown by a distribution chart (right), in which R^2 is 0.577 and ρ is 0.772 ($P = 0.0075$).

upregulated, similar to AREG and EREG whose augmented expressions were reported in our previous study. Although there was no significant difference between the levels of HB-EGF in the RA and OA samples, its high correlations with AREG and EREG ($\rho = 0.788$, $P = 0.0008$ and $\rho = 0.823$, $P = 0.0005$, respectively) imply that HB-EGF may be upregulated in RA. Although EGF and TGF α were also upregulated in RA, their expression levels showed no correlations with those of other members. These results suggest that AREG, EREG and HB-EGF may be regulated by a common expression-controlling system.

In RA-PBMCs, AREG and HB-EGF were significantly upregulated, and their expression levels were correlated with each other ($\rho = 0.600$, $P = 0.0305$). Among the seven EGF family members examined, only AREG expression was augmented in all three tissues tested in the present study. To confirm which lineage of blood cells expresses AREG, PBMCs were further separated into monocyte-, T lymphocyte- and B lymphocyte-rich fractions. None of these fractions was enriched in AREG-expressing cells in healthy controls or RA patients. In our recent report, we speculated that bone marrow-derived abnormal monocytes expressing AREG may migrate via the blood circula-

tion, and bring about disease in synovia and/or other tissues they infiltrated [25]. Although our present findings strongly support that hypothesis, the abnormal cells expressing AREG among RA-PBMCs were not restricted to monocytes. We conclude that they are mononuclear leukocytes and not of a particular lineage.

Herceptin, a specific inhibitor of ErbB2, was reported to suppress the proliferation of RA-FLS, but not OA-FLS, and augmented expression of ErbB2 was considered to be a major contributor to the autonomous proliferation of RA-FLS [16]. In the present study, EGFR and ErbB2 were found to be predominantly expressed in synovial tissues and cultured FLS, with no differences between their expression levels in RA and OA. Furthermore, the expression levels of ADAM10 and ADAM17, which are also important for the functions of EGF-like growth factors, showed no differences between RA and OA. On the other hand, the mRNA and protein levels of AREG were upregulated in RA synovial tissues. Furthermore, recombinant human AREG enhanced the proliferation of FLS in a dose-dependent manner. In our study, differences were detected for EGF-like growth factors between RA and OA synovia, but not for their receptors or sheddases. AREG induces tyrosine phosphorylation of EGFR and transduces a stronger signal when bound to EGFR/ErbB2 heterodimers [15]. It has been reported that synovitis with granulomatous hyperplasia occurs in AREG transgenic mice [35]. These findings suggest that overexpression of AREG may promote the proliferation of synoviocytes in affected joints of RA patients. We investigated whether recombinant AREG induced the expression of PDGF and bFGF, which are well-known growth factors for hyperplastic proliferation of RA-FLS. We found that AREG had no effects on the expression of these factors, suggesting that AREG did not stimulate RA-FLS to proliferate via these growth factors. However, AREG stimulated RA-FLS to express VEGF, an angiogenic factor involved in synovial hyperplasia.

A large number of reports have shown that RA-FLS produce proinflammatory cytokines when stimulated by various stimuli [36,37]. In addition, we previously reported that RA-FLS produce proinflammatory cytokines, such as IL-6, IL-8, GM-CSF, IL-1 β and/or TNF α , when co-cultured with monocytes or lymphocytes [32,33]. Our analyses of cytokine production in the present study revealed that AREG enhanced the production of several proinflammatory cytokines (IL-6, IL-8 and GM-CSF) and VEGF in RA-FLS. Since they were suppressed by an EGFR tyrosine kinase inhibitor, the AREG-dependent induction of these cytokines seemed to occur via activation of EGFR/ErbB2. Interestingly, AREG downregulated the expression of ADAM10 and ADAM17 in a dose-dependent manner. These results suggest the presence of negative feedback

regulation of ADAMs via AREG/EGFR signaling. To assess the involvement of AREG in the elevated expression of VEGF in affected joints of RA patients, the correlations between their mRNA levels and protein levels in synovial fluid samples were analyzed, and good correlations were found for both the mRNA levels ($\rho = 0.700$, $P = 0.0067$) and protein levels ($\rho = 0.772$, $P = 0.0075$). There have been very few reports about cytokine induction by AREG to date. The present results suggest that the increased level of AREG may be involved in the upregulation of VEGF in RA joints, and demonstrate, for the first time, that AREG stimulates RA-FLS to produce proinflammatory cytokines, including angiogenic factors. IL-8 has been reported to be an angiogenic factor as well as a chemoattractant factor [38,39]. Although several studies have recently shown that AREG plays important roles in hyperplasia or angiogenesis of skin diseases or tumors [40-43], the role of AREG in RA pathology remains unknown. Ma et al. speculated on the involvement of AREG in the angiogenesis of tumors [40]. Although it is known that EGF and TGF α are potent angiogenic mediators [44,45], the proangiogenic activity of AREG has not been directly determined to date. Its induction of angiogenic factors, such as IL-8 and VEGF, strongly suggests that AREG may be involved in the angiogenesis of synovial hyperplasia in affected joints of RA patients.

Myeloid cells expressing abnormal cell surface markers have been observed in RA bone marrow [18,19] and reported to be correlated with the disease severity [20-23]. Consistent with these reports, the present study revealed that several EGF-like growth factors were upregulated in RA bone marrow cells, suggesting that the onset and/or progression of chronic synovitis may be influenced by alterations to the bone marrow in RA patients.

Conclusion

Among the seven EGF-like growth factors, AREG was upregulated in synovial tissues of RA patients. Recombinant human AREG stimulated RA-FLS to proliferate and produce several proinflammatory cytokines, including angiogenic factors. These results suggest that the elevated expression of AREG in synovial tissues may be involved in RA pathology containing synovial hyperplasia. AREG-expressing cells were observed in both the blood and bone marrow of RA patients as well as in RA synovial tissues. Abnormal leukocytes may lead to the upregulated expression of AREG in affected joints of RA patients.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

SY designed the study, carried out the experiments, analyzed the data and drafted the manuscript. SI, KK and YH

carried out the RNA extractions and cDNA syntheses. RM and KT carried out the quantitative real-time PCR. NS performed the measurements of ^3H -TdR incorporation into fibroblast-like synoviocytes. TM, TJ and TO participated in the study design and collection of clinical samples. NY, NE, TI and RS participated in the study design and coordination as well as editing of the manuscript. All authors have read and approved the final manuscript.

References

- Hamilton JA: **Hypothesis: in vitro evidence for the invasive and tumor-like properties of the rheumatoid pannus.** *J Rheumatol* 1983, **10**:845-851.
- Imamura F, Aono H, Hasunuma T, Sumida T, Tateishi H, Maruo S, Nishioka K: **Monoclonal expansion of synoviocytes in rheumatoid arthritis.** *Arthritis Rheum* 1998, **41**:1979-1986.
- Aupperle KR, Boyle DL, Hendrix M, Sefter EA, Zvaifler NJ, Barbosa M, Firestein GS: **Regulation of synoviocyte proliferation, apoptosis, and invasion by the p53 tumor suppressor gene.** *Am J Pathol* 1998, **152**:1091-1098.
- Pap T, Franz JK, Hummel KM, Jeisy E, Gay R, Gay S: **Activation of synovial fibroblasts in rheumatoid arthritis: lack of expression of the tumour suppressor PTEN at sites of invasive growth and destruction.** *Arthritis Res* 2000, **2**:59-64.
- Schedel J, Distler O, Woencckhaus M, Gay RE, Simmen B, Michel BA, Muller-Ladner U, Gay S: **Discrepancy between mRNA and protein expression of tumour suppressor maspin in synovial tissue may contribute to synovial hyperplasia in rheumatoid arthritis.** *Ann Rheum Dis* 2004, **63**:1205-1211.
- Williams WV, VonFeldt JM, Ramanujam T, Weiner DB: **Tyrosine kinase signal transduction in rheumatoid synovitis.** *Semin Arthritis Rheum* 1992, **21**:317-329.
- Takeuchi T, Abe T: **Tyrosine phosphorylated proteins in synovial cells of rheumatoid arthritis.** *Int Rev Immunol* 1998, **17**:365-381.
- Migita K, Eguchi K, Tsukada T, Kawabe Y, Aoyagi T, Nagataki S: **The role of protein kinase in human synovial fibroblast growth.** *Biochem Biophys Res Commun* 1995, **210**:1066-1075.
- Remmers EF, Lafyatis R, Kumkumian GK, Case JP, Roberts AB, Sporn MB, Wilder RL: **Cytokines and growth regulation of synoviocytes from patients with rheumatoid arthritis and rats with streptococcal cell wall arthritis.** *Growth Factors* 1990, **2**:179-188.
- Butler DM, Leizer T, Hamilton JA: **Stimulation of human synovial fibroblast DNA synthesis by platelet-derived growth factor and fibroblast growth factor. Differences to the activation by IL-1.** *J Immunol* 1989, **142**:3098-3103.
- Rubin K, Terracio L, Ronnstrand L, Heldin CH, Klareskog L: **Expression of platelet-derived growth factor receptors is induced on connective tissue cells during chronic synovial inflammation.** *Scand J Immunol* 1988, **27**:285-294.
- Byrd V, Zhao XM, McKeehan WL, Miller GG, Thomas JW: **Expression and functional expansion of fibroblast growth factor receptor T cells in rheumatoid synovium and peripheral blood of patients with rheumatoid arthritis.** *Arthritis Rheum* 1996, **39**:914-922.
- Sano H, Engleka K, Mathern P, Hla T, Crofford LJ, Remmers EF, Jelsema CL, Goldmuntz E, Maciag T, Wilder RL: **Coexpression of phosphotyrosine-containing proteins, platelet-derived growth factor-B, and fibroblast growth factor-I in situ in synovial tissues of patients with rheumatoid arthritis and Lewis rats with adjuvant or streptococcal cell wall arthritis.** *J Clin Invest* 1993, **91**:553-565.
- Watanabe N, Ando K, Yoshida S, Inuzuka S, Kobayashi M, Matsui N, Okamoto T: **Gene expression profile analysis of rheumatoid synovial fibroblast cultures revealing the overexpression of genes responsible for tumor-like growth of rheumatoid synovium.** *Biochem Biophys Res Commun* 2002, **294**:1121-1129.
- Riese DJ 2nd, Stern DF: **Specificity within the EGF family/Erbb receptor family signaling network.** *Bioessays* 1998, **20**:41-48.
- Satoh K, Kikuchi S, Sekimata M, Kabuyama Y, Homma MK, Homma Y: **Involvement of Erbb-2 in rheumatoid synovial cell growth.** *Arthritis Rheum* 2001, **44**:260-265.
- Hallbeck AL, Walz TM, Briheim K, Wasteson A: **TGF-alpha and Erbb2 production in synovial joint tissue: increased expression in arthritic joints.** *Scand J Rheumatol* 2005, **34**:204-211.
- Ochi T, Hakomori S, Adachi M, Owaki H, Okamura M, Ono Y, Yamasaki K, Fujimoto M, Wakitani S, Ono K: **The presence of a myeloid cell population showing strong reactivity with monoclonal antibody directed to difucosyl type 2 chain in epiphyseal bone marrow adjacent to joints affected with rheumatoid arthritis (RA) and its absence in the corresponding normal and non-RA bone marrow.** *J Rheumatol* 1988, **15**:1609-1615.
- Kawanaka N, Yamamura M, Aita T, Morita Y, Okamoto A, Kawashima M, Iwahashi M, Ueno A, Ohmoto Y, Makino H: **CD14+, CD16+ blood monocytes and joint inflammation in rheumatoid arthritis.** *Arthritis Rheum* 2002, **46**:2578-2586.
- Ohtsu S, Yagi H, Nakamura M, Ishii T, Kayaba S, Soga H, Gotoh T, Rikimaru A, Kokubun S, Itoh T: **Enhanced neutrophilic granulopoiesis in rheumatoid arthritis. Involvement of neutrophils in disease progression.** *J Rheumatol* 2000, **27**:1341-1351.
- Berthelot JM, Bataille R, Maugars Y, Prost A: **Rheumatoid arthritis as a bone marrow disorder.** *Semin Arthritis Rheum* 1996, **26**:505-514.
- Tomita T, Shimaoka Y, Kashiwagi N, Hashimoto H, Kawamura S, Lee SB, Nakagawa S, Shiho O, Hayashida K, Ochi T: **Enhanced expression of CD14 antigen on myeloid lineage cells derived from the bone marrow of patients with severe rheumatoid arthritis.** *J Rheumatol* 1997, **24**:465-469.
- Hirohata S, Yanagida T, Itoh K, Nakamura H, Yoshino S, Tomita T, Ochi T: **Accelerated generation of CD14+ monocyte-lineage cells from the bone marrow of rheumatoid arthritis patients.** *Arthritis Rheum* 2005, **39**:836-843.
- Hirohata S, Yanagida T, Nagai T, Sawada T, Nakamura H, Yoshino S, Tomita T, Ochi T: **Induction of fibroblast-like cells from CD34+ progenitor cells of the bone marrow in rheumatoid arthritis.** *J Leukocyte Biol* 2001, **70**:413-421.
- Nakamura N, Shimaoka Y, Tougan T, Onda H, Okuzaki D, Zhao H, Fujimori A, Yabuta N, Nagamori I, Tanigawa A, Sato J, Oda T, Hayashida K, Suzuki R, Yukioka M, Nojima H, Ochi T: **Isolation and expression profiling of genes upregulated in bone marrow-derived mononuclear cells of rheumatoid arthritis patients.** *DNA Res* 2006, **13**:169-183.
- Arnett FC, Edworthy SM, Bloch DA, McShane DJ, Fries JF, Cooper NS, Healey LA, Kaplan SR, Liang MH, Luthra HS, Medsger TA Jr, Mitchell DM, Neustadt DH, Pinals RS, Schaller JG, Sharp JT, Wilder RL, Hunder GG: **The American Rheumatism Association 1987 revised criteria for the classification of rheumatoid arthritis.** *Arthritis Rheum* 1988, **31**:315-324.
- Altman RD, Asch E, Bloch D, Bole G, Borenstein D, Brandt K, Christy W, Cooke TD, Greenwald R, Hochberg M, Howell D, Kaplan D, Koopman W, Longley S III, Mankin H, McShane DJ, Medsger T Jr, Meenan R, Mikkelsen W, Moskowitz R, Murphy W, Rothschild B, Segal M, Sokoloff L, Wolfe F: **Diagnostic and Therapeutic Criteria Committee of the American Rheumatism Association. Development of criteria for the classification and reporting of osteoarthritis: classification of osteoarthritis of the knee.** *Arthritis Rheum* 1986, **29**:1039-1049.
- Mograbi B, Rochet N, Invert V, Bourget I, Bocciardi R, Emiliozzi C, Rossi B: **Human monocytes express amphiregulin and heregulin growth factors upon activation.** *Eur Cytokine Netw* 1997, **8**:73-81.
- Johansson CC, Yndestad A, Enserink JM, Ree AH, Aukrust P, Tasken K: **The epidermal growth factor-like growth factor amphiregulin is strongly induced by the adenosine 3',5'-monophosphate pathway in various cell types.** *Endocrinology* 2004, **145**:5177-5184.
- Sahn U, Weskamp G, Kelly K, Zhou HM, Higashiyama S, Peschon J, Hartmann D, Saftiq P, Blobel CP: **Distinct roles for ADAM10 and ADAM17 in ectodomain shedding of six EGFR ligands.** *J Cell Biol* 2004, **164**:769-779.
- Horiuchi K, Le Gall S, Schulte M, Yamaguchi T, Reiss K, Murphy G, Toyama Y, Hartmann D, Saftiq P, Blobel CP: **Substrate selectivity of epidermal growth factor-receptor ligand sheddases and their regulation by phorbol esters and calcium influx.** *Mol Biol Cell* 2007, **18**:176-188.
- Tanaka K, Mori T, Juji T, Suzuki S, Watanabe J, Goto A, Shiobara N, Yamane S, Fukui N, Suzuki R, Ochi T: **Production of interleukin-6 and interleukin-8 by nurse-like cells from rheumatoid arthritis.**

- tis patients after stimulation with monocytes. *Mod Rheumatol* 2005, **15**:415-422.
33. Takeuchi E, Tomita T, Toyosaki-Maeda T, Kaneko M, Takano H, Hashimoto H, Sugamoto K, Suzuki R, Ochi T: **Establishment and characterization of nurse cell-like stromal cell lines from synovial tissues of patients with rheumatoid arthritis.** *Arthritis Rheum* 1999, **42**:221-228.
 34. Ikeda M, Hosoda Y, Hirose S, Okada Y, Ikeda E: **Expression of vascular endothelial growth factor isoforms and their receptors Flt-1, KDR, and neuropilin-1 in synovial tissues of rheumatoid arthritis.** *J Pathol* 2000, **191**:426-433.
 35. Cook PVW, Brown JR, Cornell KA, Pittelkow MR: **Suprabasal expression of human amphiregulin in the epidermis of transgenic mice induces a severe, early-onset, psoriasis-like skin pathology: expression of amphiregulin in the basal epidermis is also associated with synovitis.** *Exp Dermatol* 2004, **13**:347-356.
 36. Jeong JG, Kim JM, Cho H, Hahn W, Yu SS, Kim S: **Effects of IL-1beta on gene expression in human rheumatoid synovial fibroblasts.** *Biochem Biophys Res Commun* 2004, **324**:3-7.
 37. Ritchlin C: **Fibroblast biology: effector signals released by the synovial fibroblast in arthritis.** *Arthritis Res* 2000, **2**:356-360.
 38. Koch AE, Polverini PJ, Kunkel SL, Harlow LA, DiPietro LA, Elnor VM, Elnor SG, Strieter RM: **Interleukin-8 as a macrophage-derived mediator of angiogenesis.** *Science* 1992, **258**:1798-1801.
 39. Koch AE, Volin MV, Woods JM, Kunkel SL, Connors MA, Harlow LA, Woodruff DC, Burdick MD, Strieter RM: **Regulation of angiogenesis by the C-X-C chemokines interleukin-8 and epithelial neutrophil activating peptide 78 in the rheumatoid joint.** *Arthritis Rheum* 2001, **44**:31-40.
 40. Ma L, Gauville C, Berthois Y, Millot G, Johnson GR, Calvo F: **Antisense expression for amphiregulin suppress tumorigenicity of a transformed human breast epithelial cell line.** *Oncogene* 1999, **18**:6513-6520.
 41. Desruisseau S, Palmari J, Giusti C, Romain S, Martin PM, Berthois Y: **Clinical relevance of amphiregulin and VEGF in primary breast cancers.** *Int J Cancer* 2004, **111**:733-740.
 42. Riedel F, Gotte K, Li M, Hormann K, Grandis JR: **RGFR antisense treatment of human HNSCC cell lines down-regulates VEGF expression and endothelial cell migration.** *Int J Oncol* 2002, **21**:11-16.
 43. Bhagavathula N, Nerusu KC, Fisher GJ, Liu G, Thakur AB, Gemmill L, Kumar S, Xu ZH, Hinton P, Tsurushita N, Landolfi NF, Voorhees JJ, Warani J: **Amphiregulin and epidermal hyperplasia - amphiregulin is required to maintain the psoriatic phenotype of human skin grafts on severe combined immunodeficient mice.** *Am J Pathol* 2005, **166**:1009-1016.
 44. Schreiber AB, Winkler ME, Derynck R: **Transforming growth factor-alpha: a more potent angiogenic mediator than epidermal growth factor.** *Science* 1986, **232**:1250-1253.
 45. Detmar M, Brown LF, Claffey KP, Yeo KT, Kocher O, Jackman RW, Berse B, Dvorak HF, Van de Water L: **Overexpression of vascular permeability factor/vascular endothelial growth factor and its receptors in psoriasis.** *J Exp Med* 1994, **180**:1141-1146.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:

http://www.biomedcentral.com/info/publishing_adv.asp



LIGHT Induces Cell Proliferation and Inflammatory Responses of Rheumatoid Arthritis Synovial Fibroblasts via Lymphotoxin β Receptor

SATORU ISHIDA, SHOJI YAMANE, SAORI NAKANO, TOSHIHITO MORI, TAKUO JUJI, NAOSHI FUKUI, TSUNETOSHI ITOH, TAKAHIRO OCHI, and RYUJI SUZUKI

ABSTRACT. *Objective.* To investigate the effects of LIGHT (lymphotoxin-like, exhibits inducible expression and competes with herpes simplex virus glycoprotein D for herpes virus entry mediator, a receptor expressed by T lymphocytes) on the proliferation and gene expression of fibroblast-like synoviocytes (FLS) from patients with rheumatoid arthritis (RA).

Methods. We measured LIGHT levels in RA synovial fluids (SF) by ELISA, and compared them with those in osteoarthritis (OA) SF. Levels of LIGHT and its receptors in RA-FLS and synovium were assessed using real-time quantitative polymerase chain reaction (PCR). RA-FLS proliferation was examined by a bromodeoxyuridine assay. Expression of intercellular adhesion molecule-1 (ICAM-1) and several chemokines, such as interleukin 8 (IL-8), monocyte chemoattractant protein-1 (MCP-1), and macrophage inflammatory protein-1 α (MIP-1 α), was examined by real-time quantitative PCR, ELISA, and flow cytometry. The effects of LIGHT on nuclear factor- κ B (NF- κ B) activation were investigated using immunofluorescence and Western blotting.

Results. LIGHT was upregulated in both SF and synovium of RA patients compared with OA patients. Herpes virus entry mediator (HVEM) and lymphotoxin β receptor (LT β R), but not LIGHT, were detected in RA-FLS. LIGHT significantly promoted RA-FLS proliferation and induced expression of MCP-1, IL-8, MIP-1 α , and ICAM-1 by RA-FLS. As well, LT β R small interfering RNA (siRNA), but not HVEM siRNA, inhibited these effects of LIGHT. LIGHT induced I κ B α degradation and NF- κ B translocation, and a NF- κ B inhibitor suppressed the effects of LIGHT on RA-FLS.

Conclusion. Our findings suggest that LIGHT signaling via LT β R plays an important role in the pathogenesis of RA by affecting key processes such as the proliferation and activation of RA-FLS. Regulation of LIGHT-LT β R signaling may represent a new therapeutic target for RA treatment. (First Release April 15 2008; J Rheumatol 2008;35:960-8)

Key Indexing Terms:

RHEUMATOID ARTHRITIS

INFLAMMATION

SYNOVIOCYTES

Rheumatoid arthritis (RA) is a chronic inflammatory disease characterized by synovial hyperplasia and progressive destruction of cartilage and bone. Fibroblast-like synovio-

cytes (FLS), an important component of the synovial lining in joints, proliferate aggressively to form a pannus causing irreversible joint damage. In RA synovial tissue, activated FLS and infiltrating macrophages and lymphocytes produce inflammatory cytokines, including tumor necrosis factor- α (TNF- α), interleukin 1 β (IL-1 β), and IL-6, that play important roles in the pathogenesis of RA^{1,2}. These cytokines have been shown to not only directly promote FLS proliferation leading to pannus formation³, but also to induce the expression of inflammatory cytokines, chemokines, and adhesion molecules, which further recruit inflammatory leukocytes and perpetuate inflammatory responses.

LIGHT (lymphotoxin-like, exhibits inducible expression and competes with herpes simplex virus glycoprotein D for herpes virus entry mediator, a receptor expressed by T lymphocytes) is a recently identified type-2 transmembrane glycoprotein of the TNF ligand superfamily (TNFSF14)⁴. LIGHT is expressed on activated T lymphocytes^{4,5}, monocytes⁶, granulocytes⁶, and immature dendritic cells⁷. LIGHT signaling is transduced via 2 members of the TNFR family,

From the Clinical Research Center for Allergy and Rheumatology, National Hospital Organization, Sagami National Hospital, Sagami, Kanagawa; Discovery Research Laboratories, Shionogi & Co., Ltd., Toyonaka, Osaka; and Department of Immunology and Embryology, Tohoku University School of Medicine, Aoba-ku, Sendai, Japan.

S. Ishida, MSc; S. Yamane, PhD, Clinical Research Center for Allergy and Rheumatology, National Hospital Organization, Sagami National Hospital, Discovery Research Laboratories, Shionogi & Co., Ltd.; S. Nakano, BSc; T. Mori, MD, PhD; T. Juji, MD, PhD; N. Fukui, MD, PhD, Clinical Research Center for Allergy and Rheumatology, National Hospital Organization, Sagami National Hospital; T. Itoh, MD, PhD, Department of Immunology and Embryology, Tohoku University School of Medicine; T. Ochi, MD, PhD; R. Suzuki, DVM, PhD, Clinical Research Center for Allergy and Rheumatology, National Hospital Organization, Sagami National Hospital.

Address reprint requests to S. Ishida, Clinical Research Center for Allergy and Rheumatology, National Hospital Organization, Sagami National Hospital, Sakuradai 18-1, Sagami, Kanagawa, 228-8522, Japan. E-mail: satoru.ishida@shionogi.co.jp

Accepted for publication January 10, 2008.

herpes virus entry mediator (HVEM, TNFRSF14) and lymphotoxin β receptor (LTBR, TNFRSF3). HVEM is expressed prominently on monocytes, dendritic cells, and lymphocytes^{5,8-10}, whereas LTBR is expressed on many cell types with the exception of lymphocytes^{4,6,11}. LIGHT has been shown to regulate cell proliferation^{7,12,13} and apoptosis^{6,14} to induce the secretion of various cytokines, and to augment the expression of adhesion molecules^{12,15-17}. Recently, Fava, *et al* reported that LTBR-Ig protein blocked the induction of experimental arthritis in mice¹⁸. Moreover, LIGHT induced the expression of inflammatory cytokines in macrophages from RA synovial fluid (SF)¹⁹. These studies suggest that LIGHT may be an important inflammatory cytokine in the development of RA. However, the effect of LIGHT on RA-FLS has not yet been analyzed.

Our aim was to clarify the role of LIGHT in the proliferation and inflammatory response of RA-FLS. We observed that the concentrations of LIGHT in both SF and synovium were higher in patients with RA than in those with osteoarthritis (OA). In addition, LIGHT signaling via LTBR, but not HVEM, enhanced RA-FLS proliferation and induced the expression of inflammatory cytokines, chemokines, and adhesion molecules in RA-FLS through a nuclear factor- κ B (NF- κ B)-dependent signal transduction pathway. We suggest that activation of RA-FLS by LIGHT/LTBR signaling may play an important role in the pathogenesis of RA.

MATERIALS AND METHODS

Chemicals. Recombinant human LIGHT and platelet-derived growth factor (PDGF)-AB were obtained from R&D Systems (Minneapolis, MN, USA). Monoclonal antibodies (mAb) against human actin and NF- κ B p65 were purchased from Sigma-Aldrich (St. Louis, MO, USA) and BD Biosciences (Palo Alto, CA, USA), respectively. The mAb against I κ B α was from Santa Cruz Biotechnology (Santa Cruz, CA, USA). Pyrrolidine dithiocarbamate (PDTC) was purchased from Calbiochem (La Jolla, CA, USA).

Patients and tissue samples. All patients with RA fulfilled the 1987 American College of Rheumatology (formerly, the American Rheumatism Association) criteria²⁰ for the diagnosis of RA. Patients with RA ranged in age from 41 to 74 years (mean \pm SD 66.0 \pm 12.0 yrs). Patients with OA ranged in age from 39 to 90 years (mean \pm SD 64.1 \pm 14.7 yrs). All patients were women. Synovial tissues were obtained from 27 patients with RA and 11 patients with OA at the time of knee prosthetic replacement surgery. RA-FLS were established from the synovia of RA patients as described²¹. RA-FLS were cultured in Dulbecco's modified Eagle's medium (DMEM; Invitrogen, Carlsbad, CA, USA) supplemented with 10% fetal calf serum, penicillin, streptomycin, and L-glutamine. RA-FLS from passages 4-9 were used for each experiment. SF were obtained by arthrocentesis from 23 RA patients and 10 OA patients and, after centrifugation at 20,000 \times g for 10 min, the supernatants were collected and frozen at -80°C until used. All specimens were obtained from patients who gave written informed consent, according to the protocol approved by the institutional review board of the National Hospital Organization, Sagami National Hospital.

LIGHT in synovial fluids. The amount of LIGHT in SF was measured using an ELISA kit (R&D Systems) according to the manufacturer's instructions. The minimum and maximum detection levels of the ELISA were 7.8 pg/ml and 2000 pg/ml, respectively.

RNA extraction, cDNA synthesis, and real-time quantitative polymerase chain reaction (PCR) analysis. Total RNA was extracted from synovium and FLS using an RNeasy Micro kit (Qiagen). cDNA was generated from

RNA using Omniscript Reverse Transcriptase (Qiagen) and used as a template for real-time quantitative PCR on a LightCycler (Roche Diagnostics). PCR was performed using SYBR Premix Ex Taq (Takara). The primers used for real-time PCR were as follows: for IL-6, 5'-AAG CCA GAG CTG TGC AGA TGA GTA-3' and 5'-TGT CCT GCA GCC ACT GGT TC-3'; for IL-8, 5'-ACA CTG CGC CAA CAC AGA AAT TA-3' and 5'-TTT GCT TGA AGT TTC ACT GGC ATC-3'; for granulocyte macrophage-colony stimulating factor (GM-CSF), 5'-CAT GAT GGC CAG CCA CTA CAA-3' and 5'-ACT GGC TCC CAG CAG TCA AAG-3'; for monocyte chemoattractant protein-1 (MCP-1), 5'-GCT CAT AGC AGC CAC CTT CAT TC-3' and 5'-GGA CAC TTG CTG CTG GTG ATT C-3'; for RANTES, 5'-ACC AGT GGC AAG TGC TCC AAC-3' and 5'-CTC CCA AGC TAG GAC AAG AGC AAG-3'; for MIP-1 α , 5'-TCC GTC ACC TGC TCA GAA TCA-3' and 5'-AGC ACT GGC TGC TCG TCT CA-3'; for vascular cell adhesion molecule-1 (VCAM-1), 5'-CGT GAT CCT TGG AGC CTC AAA TA-3' and 5'-GAC GGA GTC ACC AAT CTG AGC A-3'; for intercellular adhesion molecule-1 (ICAM-1), 5'-CCT GAT GGG CAG TCA ACA GCT A-3' and 5'-ACA GCT GGC TCC CGT TTC A-3'; for GAPDH, 5'-GCA CCG TCA AGG CTG AGA AC-3' and 5'-ATG GTG GTG AAG ACG CCA GT-3'; for LIGHT, 5'-TCA CGA GGT CAA CCC AGC AG-3' and 5'-CCC AGC TGC ACC TTG GAG TAG-3'; for HVEM, 5'-TTT GCT CCA CAG TTG GCC TAA TC-3' and 5'-CAA TGA CTG TGG CCT CAC CTT C-3'; and for LTBR, 5'-ATG CTG ATG CTG GCC GTT C-3' and 5'-AGG CTC CCA GCT TCC AGC TA-3'.

PCR was performed under the following conditions: initial denaturation at 95°C for 10 s, then 40 cycles of 95°C for 5 s and 60°C for 20 s. When SYBR Green dye was used to monitor PCR, melting curves were routinely recorded to verify the singularity of the PCR product. In each sample, the level of cDNA was normalized to the level of GAPDH.

Proliferation assay. RA-FLS were seeded into a 96-well plate at a density of 5.0 \times 10³ cells/well. After 24 h of preculture, the cells were stimulated for 48 h by the addition of LIGHT or PDGF used as a positive control. A previous report described stimulation of RA-FLS proliferation by PDGF²². Bromodeoxyuridine (BrdU) was added for the last 24 h of culture, then incorporation of BrdU was measured using a cell proliferation ELISA (Roche Diagnostics) according to the manufacturer's instructions.

Measurement of cytokine and chemokine levels in culture supernatants. TNF- α , IL-1 β , IL-6, IL-8, and GM-CSF levels were measured in the supernatants of RA-FLS cultures using a Human Inflammatory Five-Plex Antibody bead kit (Biosource, Camarillo, CA, USA) according to the manufacturer's instructions, on a Luminex 100 instrument (Luminex, Austin, TX, USA). The levels of MIP-1 α , MIP-1 β , MCP-1, eotaxin, and RANTES in the supernatant were measured using a Human Chemokine Five-Plex Antibody bead kit (Biosource).

Transfection of RA-FLS with small interfering RNA (siRNA). All siRNA were purchased from Qiagen. The sense strand sequences of the RNA duplexes were as follows: HVEM, 5'-GGC ACU GCC UCA CAG CCA AdTdT-3'; LTBR, 5'-CAU CUA CAA UGG ACC AGU AdTdT-3'; and control siRNA 5'-UUC UCC GAA CGU GUC ACG UdTdT-3'. The day before transfection, RA-FLS were seeded into 6-well culture plates at a density of 4 \times 10⁴ cells/well, or 96-well plates at 5 \times 10³ cells/well, in complete medium without antibiotics. The next day, siRNA (at final concentration of 50 nM) were introduced into cells using Lipofectamine 2000 (Invitrogen) following the manufacturer's instructions; 24 h after transfection, media were replaced with regular culture media. The cells were then cultured for 96 h before analysis of the gene-silencing effects. mRNA levels were measured by real-time quantitative PCR analysis.

Immunofluorescence assay for NF- κ B localization. To examine the nuclear translocation of NF- κ B, RA-FLS were seeded at a density of 5 \times 10³ cells/well in 8-well Lab-Tek chamber slides (Nalgen Nunc International, Naperville, IL, USA). Cells were stimulated with 10 ng/ml LIGHT for 40 min, washed with cold PBS, and then fixed in PBS with 4% paraformaldehyde for 10 min. The cells were permeabilized with PBS and 0.1% Triton-X100 for 10 min. Nonspecific binding was prevented with blocking buffer

containing 2% goat serum diluted in PBS. The cells were incubated with mouse monoclonal anti-NF- κ B p65 antibody or an isotype control for 1 h at room temperature, then with AlexaFluor 488-conjugated goat anti-mouse antibody for 30 min at room temperature. Slides were coverslipped and examined using a fluorescence microscope (Olympus, Tokyo, Japan).

Western blotting analysis. For measurement of I κ B α by Western blotting, RA-FLS at a density of 1.5×10^6 /well were seeded into 6-well plates in culture medium for 24 h. After incubation with 10 ng/ml LIGHT for 40 min, cells were washed twice in ice-cold PBS and lysed in 100 μ l of sodium dodecyl sulfate (SDS) sample buffer (62.5 mM Tris HCl, pH 6.8, 10% glycerol, 2% SDS, 5% mercaptoethanol, and 0.001% bromophenol blue). Cell lysates were separated by SDS-polyacrylamide gel electrophoresis in 12% polyacrylamide gels, and transferred onto nitrocellulose membranes (Invitrogen). After blocking, membranes were incubated with either anti- β -actin or anti-I κ B α antibody, overnight at 4°C, and then with secondary antibody conjugated to horseradish peroxidase (Dako), at room temperature for 1 h. The signals were visualized using chemiluminescence reagent (ECL; Amersham Biosciences, Little Chalfont, UK).

Statistical analysis. Comparisons of ≥ 3 populations were made using the Kruskal-Wallis test. Comparisons of 2 independent data sets were by Mann-Whitney U-test. P values less than 0.05 were considered statistically significant.

RESULTS

Increased expression of LIGHT in SF of patients with RA. To examine whether LIGHT is involved in the pathogenesis of RA, we analyzed the level of LIGHT in SF from 23 RA patients and 10 OA patients by ELISA. SF from OA patients were used as controls, because they were not available from healthy individuals. The concentration of LIGHT in SF from RA patients was significantly higher than in those from OA patients (Figure 1). The median levels of LIGHT in SF from RA and OA patients were 108.5 pg/ml and 7.8 pg/ml, respectively.

Expression of LIGHT and its receptors in RA synovial tissue and RA-FLS. Because RA patients had more LIGHT in their SF than OA patients, we investigated whether LIGHT and its membrane-bound receptors HVEM and LTBR were expressed in the RA and OA synovial tissues. Although




Figure 1. Levels of LIGHT in synovial fluid from RA patients and OA patients, determined by ELISA. Box plots represent 25th to 75th percentiles. Error bars represent 10th to 90th percentiles. Lines inside boxes represent the median. * $p < 0.05$.

quantitative real-time PCR analysis revealed that mRNA expression of LIGHT in synovial tissue was significantly higher in RA patients than in OA patients (Figure 2A), HVEM and LTBR levels were not different between RA and OA patients.

Further, we investigated the mRNA expression of LIGHT, HVEM, and LTBR in RA-FLS by quantitative real-time PCR. RA-FLS from all 7 patients expressed HVEM and LTBR mRNA, and the level of LTBR mRNA was significantly higher than that of HVEM mRNA, whereas no LIGHT expression was detected (Figure 2B).

Induction of RA-FLS proliferation by LIGHT. Previous studies reported that LIGHT induces cell proliferation in T lymphocytes^{7,12} and vascular smooth muscle cells¹³. Since the expression of HVEM and LTBR in RA-FLS had been confirmed, we next evaluated the effect of LIGHT on the proliferation of RA-FLS using a BrdU assay. As shown in




Figure 2. Expression of LIGHT, herpes virus entry mediator (HVEM), and lymphotoxin β receptor (LTBR) mRNA in synovial tissues and fibroblast-like synoviocytes (FLS): A. In synovial tissues from RA patients and OA patients. B. In RA-FLS. Level was evaluated by real-time quantitative PCR; results are represented as relative ratios to GAPDH levels. * $p < 0.05$.