



## ORIGINAL ARTICLE

# Relationship between number of metabolic syndrome components and dietary factors in middle-aged and elderly Japanese subjects

Rei Otsuka<sup>1</sup>, Tomoko Imai<sup>1,2</sup>, Yuki Kato<sup>1</sup>, Fujiko Ando<sup>1,3</sup> and Hiroshi Shimokata<sup>1</sup>

Metabolic syndrome (MetS) represents a cluster of risk factors for atherosclerosis and is considered a risk factor for cardiovascular disease. The role of diet in the etiology of MetS is poorly understood, especially among Asian subjects. This cross-sectional study assessed the relationship between diet and the number of MetS components among Japanese men ( $n=609$ ) and women ( $n=631$ ). Mean (s.d.) age and body mass index were 57.1 (12.1) years and 22.8 (2.8)  $\text{kg m}^{-2}$  for men and 55.5 (12.0) years and 22.0 (3.0)  $\text{kg m}^{-2}$  for women, respectively. Diet was assessed by a 3-day dietary record that included photographs: 16 nutrients, 11 food groups, and energy % of protein and dietary fat were selected as a dietary index. The definition of MetS was based on modified National Cholesterol Education Program, Adult Treatment Panel III criteria, and the number of clustering MetS components was calculated by adding the presence of each five MetS components. A total of 61 men (10.0%) and 46 women (7.3%) were determined to have MetS. After adjusting for age, energy intake, alcohol intake, smoking status and physical activity, a lower intake of vitamin B6 and dietary fiber in men, and lower intake of calcium, milk and dairy products and higher intake of cereal in women were related to the number of MetS components. These results suggest that some dietary factors were related to the number of MetS components among community-dwelling Japanese men and women. *Hypertension Research* (2010) 33, 548–554; doi:10.1038/hr.2010.29; published online 12 March 2010

**Keywords:** cross-sectional study; dietary record; Japanese; metabolic syndrome

## INTRODUCTION

Metabolic syndrome (MetS) represents a cluster of risk factors for atherosclerosis, including visceral obesity, hypertension, dyslipidemia and hyperglycemia; MetS is considered a risk factor for cardiovascular disease.<sup>1</sup> The National Nutrition Survey in Japan, a population-based study among 40- to 74-year-olds, revealed that 24% of men and 12% of women were strongly suspected of having MetS, and 27% of men and 8% of women were suspected of having MetS.<sup>2</sup>

MetS has become a major public health challenge in Japan.<sup>3</sup> The pathophysiology of MetS appears to be largely attributable to insulin resistance with an excessive flux of fatty acids,<sup>1</sup> although this disorder presumably exists as a function of a complex interaction between environmental factors, including diet or physical activity and genetic factors.<sup>4,5</sup>

Although dietary aspects have been linked to individual features of MetS,<sup>6–8</sup> the role of diet in the etiology of this syndrome is poorly understood. Asians have different lifestyle and genetic factors compared with Caucasians,<sup>9,10</sup> but only a few epidemiologic studies examining diet and MetS among Asians have been conducted.<sup>11–13</sup> The aim of this study was to examine the relations between diet and

the number of MetS factors among community-dwelling Japanese men and women.

## METHODS

### Study subjects

Data for this survey were collected as part of the National Institute for Longevity Sciences Longitudinal Study of Aging (NILS-LSA). In this project, the normal aging process has been assessed over time using detailed questionnaires and medical checkups, anthropometrical measurements, physical fitness tests and nutritional examinations. Participants in the NILS-LSA included randomly selected age- and sex-stratified individuals from the pool of residents in the NILS neighborhood areas, Obu City and Higashiura Town of Aichi Prefecture. Details of the NILS-LSA study are reported elsewhere.<sup>14</sup>

Subjects in this study included 1189 men and 1194 women aged 40–86 years who participated in the fourth wave of the NILS-LSA from June 2004 to July 2006. Subjects under non-pharmacological and/or pharmacological treatment for hypertension, hypertriglyceridemia or diabetes were excluded, as were subjects who indicated that they were aware of having these disorders but were not undergoing treatment. There were 334 men and 327 women with hypertension, 241 men and 174 women with hypertriglyceridemia and 104 men and 69 women with diabetes. As some subjects had multiple disorders, a total

<sup>1</sup>Department of Epidemiology, National Institute for Longevity Sciences, National Center for Geriatrics and Gerontology, Aichi, Japan; <sup>2</sup>Department of Registered Dietitians, Faculty of Human Wellness, Tokaigakuen University, Aichi, Japan; <sup>3</sup>Department of Community Care Philanthropy, Faculty of Medical Welfare, Aichi Shukutoku University, Aichi, Japan  
Correspondence: Dr R Otsuka, Department of Epidemiology, National Institute for Longevity Sciences, National Center for Geriatrics and Gerontology, 36-3 Gengo, Morioka-cho, Obu, Aichi 454-8522, Japan.  
E-mail: otsuka@ncgg.go.jp

Received 26 May 2009; revised 17 December 2009; accepted 1 February 2010; published online 12 March 2010

of 489 men and 492 women were excluded from the study. Additionally, subjects who did not fast overnight for venipuncture (11 men and 12 women), those who did not participate or complete the nutrition survey (54 men and 47 women) and those whose energy intake was <1200 kcal per day (2 men and 11 women) or more than 3000 kcal per day (24 men and 1 woman) were also excluded. After these exclusions, 609 men and 631 women remained in the study.

The study protocol was approved by the Committee of Ethics of Human Research of the National Center for Geriatrics and Gerontology. Written informed consent was obtained from all subjects.

#### Nutritional assessments

Nutritional intakes were assessed by a 3-day dietary record. The dietary record was completed over 3 continuous days (both weekend days and one weekday).<sup>15</sup> Food was weighed separately on a scale (1 kg kitchen scale, Sekisui Jushi, Tokyo, Japan) before being cooked or portion sizes were estimated. Subjects used a disposable camera (27 shots, Fuji Film, Tokyo, Japan) to take photos of meals before and after eating. Dietitians used the photos to complete missing data, and telephoned subjects to resolve any discrepancies or obtain further information when necessary. The averages of the 3-day food and nutrient intakes were calculated according to the fifth edition of the Standard Tables of Foods Composition in Japan and other sources.<sup>15</sup> Alcohol intake in the previous year was assessed by a food frequency questionnaire; trained dietitians interviewed subjects using this questionnaire. According to previous large epidemiological studies,<sup>7,11</sup> we selected 16 nutrients and 11 food groups as a dietary index, along with energy % of protein and dietary fat.

#### Other measurements

Anthropometric measurements included waist circumference, height and body weight. Waist circumference was measured at the umbilicus<sup>16,17</sup> and body mass index was calculated as weight/height<sup>2</sup> (kg m<sup>-2</sup>). Blood pressure was measured by an automated sphygmomanometer (BP-203RVII, Omron Colin, Tokyo, Japan) after participants had been comfortably seated for at least 5 min. All venous blood samples were obtained after an overnight fast. The serum was separated promptly, and all lipid analyses were conducted at the clinical laboratory in the health examination center. Serum glucose and triglycerides were measured using enzymatic methods. HDL-cholesterol was measured after dextran sulfate-magnesium precipitation.

Medical history (past and current) and smoking status (yes/no) were collected using questionnaires. Physical activity was assessed by trained interviewers using the Met Score (a multiple of the resting metabolic rate). Participants were interviewed using a semi-quantitative assessment to determine their level of habitual physical activity during leisure time, on the job and sleeping hours,<sup>18</sup> and we calculated the total MetS\*minutes score per day (MetS\*1000 min per day). For example, walking for pleasure was assigned a 2.5 MetS intensity, and thus the leisure-time physical activity score was 50 MetS\*min per day in the case of a 20 min walk every day.<sup>18</sup>

#### Definition of MetS and components

The definition of MetS was based on modified National Cholesterol Education Program, Adult Treatment Panel III (NCEP-ATP III) criteria.<sup>19</sup> Only the criteria of abdominal obesity was different from that of NCEP-ATP III criteria.<sup>20</sup> For abdominal obesity, we used the International Obesity task Force central obesity criteria for Asia, which defined abdominal obesity as a waist circumference of at least 90 cm for men and at least 80 cm for women.<sup>21</sup>

As stated earlier, study subjects under treatment for hypertension, hypertriglyceridemia and diabetes were excluded before the analyses. Thus, MetS was defined as the presence of three or more of the following five components: (1) abdominal obesity, defined as a waist circumference of at least 90 cm for men and at least 80 cm for women; (2) elevated blood pressure, defined as blood pressure  $\geq 130/85$  mm Hg; (3) hypertriglyceridemia, defined as triglycerides  $\geq 150$  mg per 100 ml ( $\geq 1.70$  mmol l<sup>-1</sup>); (4) low HDL-cholesterol, defined as HDL-cholesterol  $< 40$  mg per 100 ml ( $< 1.0$  mmol l<sup>-1</sup>) in men and  $< 50$  mg per 100 ml ( $< 1.3$  mmol l<sup>-1</sup>) in women; and (5) elevated blood glucose levels, defined as fasting blood glucose  $\geq 100$  mg per 100 ml. The number of clustering MetS components was calculated by adding the number of MetS components.

#### Statistical analyses

All statistical analyses were conducted with Statistical Analysis System, release 9.1.3 (SAS Institute, Cary, NC, USA). Subjects were categorized into four groups according to the number of clustering MetS components (0, 1, 2, 3–5). Values of 3–5 were combined because only a few or no subjects had 4 or 5 MetS components (4 components: 12 men and 5 women; 5 components: 0 men and 2 women). Associations between categorical variables were tested by  $\chi^2$  test and 95% confidence interval (CI) was estimated using the PROC FREQ procedure. Comparisons between continuous variables were performed by analysis of variance and trend test. Linear regression models were constructed using the PROC GLM procedure to examine the association between the number of MetS components and dietary indexes, that is, the 11 food groups, 16 nutrients and energy % of protein and dietary fat. Mean nutritional intakes were calculated by the number of MetS components (0, 1, 2, 3–5) after multivariate adjustment for potential confounding factors, which included age, energy intake, alcohol intake, smoking status and physical activity. Additionally, demographic differences between subjects (609 men and 631 women) and those excluded from the study (580 men and 563 women) were analyzed by *t*-test.

All reported *P*-values were two-sided, and a *P*-value  $< 0.05$  was considered significant.

#### RESULTS

Subject characteristics are presented Table 1. Mean (s.d.) age and BMI were 57.1 (12.1) years and 22.8 (2.8) kg m<sup>-2</sup> for men and 55.5 (12.0) years and 22.0 (3.0) kg m<sup>-2</sup> for women, respectively. Age and BMI gradually increased with the number of MetS components in both men and in women, whereas mean age of subjects with 2 or 3–5 MetS components was similar in men (59.5 years and 59.5 years, respectively) and women (60.7 and 60.9 years, respectively).

Abdominal obesity, defined as waist circumference  $\geq 90$  cm, was noted in 20.0% of men (95% CI: 16.8–23.2), with a prevalence of 14.4, 41.9 and 70.5% among men with 1, 2 or 3–5 MetS components, respectively. Abdominal obesity, defined as waist circumference  $\geq 80$  cm, was noted in 44.1% of women (95% CI: 40.2–48.0), with a prevalence of 69.7, 84.5 and 97.8% among those with 1, 2 or 3–5 MetS components, respectively. Elevated blood glucose levels were seen in 38.1% of men and 17.9% of women according to the number of MetS components (1, 2 or 3–5) was 49.8, 61.5 and 91.8% in men and 17.1, 41.8 and 73.9% in women, respectively. Hypertriglyceridemia and hypertension were seen in 18.2 and 22.5% of men and 7.5 and 12.8% of women, respectively. Low HDL-C levels occurred in 5.9% of men and 7.0% of women. A total of 61 men (10.0%) and 46 women (7.3%) met the NCEP-ATP III modified criteria for MetS, that is, they had 3–5 MetS components.

In sub-analyses, demographic differences between subjects included in the study (609 men and 631 women) and those excluded from the study (580 men and 563 women) were analyzed. Age, BMI and the other anthropometric variables (for example, blood pressure, waist circumference or fasting glucose levels) were lower among subjects than those excluded (mean (s.d.) age: 56.9 (12.1) vs. 64.8 (11.7) years, mean BMI: 22.5 (2.9) vs. 23.6 (3.2) kg m<sup>-2</sup>, respectively).

Table 2 shows multivariate adjusted mean food and nutrient intake according to the number of MetS components in men. Among the dietary indexes, daily intakes of vitamin B6 decreased from 1.36 to 1.21 mg, and dietary fiber decreased from 16.2 to 14.5 mg as the number of MetS components increased. Although analysis of covariance did not reach statistical significance, intake of vegetables was lower among men with higher number of MetS components, and decreased from 301.1 to 271.9 g as the number of MetS components increased (ANCOVA *P*=0.07, trend test *P*=0.03). In addition, although analysis of covariance or trend tests did not reach statistical

Table 1 Subject characteristics according to the number of metabolic syndrome (MetS) components

Variable	All	Number of MetS components					P <sup>a</sup>	Trend P
		0	1	2	3-5			
<b>Men</b>								
n, %	609 (100%)	222 (36.5%)	209 (34.3%)	117 (19.2%)	61 (10.0%)			
Age (year)	57.1 ± 12.1	55.1 ± 12.5	57.1 ± 11.7	59.5 ± 12.0	59.5 ± 11.0	<0.01	<0.01	
Body mass index (kg m <sup>-2</sup> )	22.8 ± 2.8	21.7 ± 2.4	22.5 ± 2.5	24.0 ± 2.7	25.6 ± 2.6	<0.01	<0.01	
Alcohol intake (g per day)	14.2 ± 17.3	12.7 ± 16.1	14.6 ± 17.1	15.9 ± 17.8	15.3 ± 21.2	0.38	0.24	
Physical activity (MetS*1000 min per day)	2116 ± 302	2107 ± 290	2109 ± 266	2124 ± 334	2159 ± 389	0.65	0.21	
Energy intake (kcal/)	2247 ± 335	2247 ± 308	2233 ± 342	2283 ± 337	2227 ± 399	0.59	0.95	
Current smoker (n, %)	186 (30.5%)	62 (27.9%)	67 (32.1%)	38 (32.5%)	19 (31.2%)	0.76		
<b>Metabolic abnormalities (n, %)</b>								
Waist circumference ≥ 90 (cm)	122 (20.0%)	0 (-%)	30 (14.4%)	49 (41.9%)	43 (70.5%)			
Triglyceride ≥ 150 (mg per 100 ml)	111 (18.2%)	0 (-%)	24 (11.5%)	44 (37.6%)	43 (70.5%)			
HDL-cholesterol < 40 (mg per 100 ml)	36 (5.9%)	0 (-%)	7 (3.4%)	17 (14.5%)	12 (19.7%)			
Blood pressure ≥ 130/85 mm Hg	137 (22.5%)	0 (-%)	44 (21.1%)	52 (44.4%)	41 (67.2%)			
Fasting glucose ≥ 100 mg per 100 ml	232 (38.1%)	0 (-%)	104 (49.8%)	72 (61.5%)	56 (91.8%)			
<b>Women</b>								
n, %	631 (100%)	272 (43.1%)	210 (33.3%)	103 (16.3%)	46 (7.3%)			
Age (year)	55.5 ± 12.0	51.5 ± 10.9	56.8 ± 12.1	60.7 ± 11.8	60.9 ± 10.9	<0.01	<0.01	
Body mass index (kg m <sup>-2</sup> )	22.0 ± 3.0	20.2 ± 1.9	22.7 ± 2.7	23.6 ± 3.0	25.4 ± 3.2	<0.01	<0.01	
Alcohol intake (g per day)	3.4 ± 7.6	3.7 ± 7.3	3.2 ± 7.7	2.3 ± 5.3	4.6 ± 12.1	0.27	0.60	
Physical activity (MetS*1000 min per day)	2161 ± 160	2187 ± 152	2144 ± 169	2132 ± 131	2143 ± 201	<0.01	0.07	
Energy intake (kcal/)	1884 ± 288	1884 ± 299	1878 ± 293	1893 ± 275	1895 ± 229	0.97	0.72	
Current smoker (n, %)	39 (6.2%)	21 (7.7%)	9 (4.3%)	4 (3.9%)	5 (10.9%)	0.17		
<b>Metabolic abnormalities (n, %)</b>								
Waist circumference ≥ 80 (cm)	278 (44.1%)	0 (-%)	146 (69.7%)	87 (84.5%)	45 (97.8%)			
Triglyceride ≥ 150 (mg per 100 ml)	47 (7.5%)	0 (-%)	6 (2.9%)	21 (20.4%)	20 (43.5%)			
HDL-cholesterol < 50 (mg per 100 ml)	44 (7.0%)	0 (-%)	8 (3.8%)	19 (18.5%)	17 (37.0%)			
Blood pressure ≥ 130/85 mm Hg	81 (12.8%)	0 (-%)	14 (6.7%)	36 (35.0%)	31 (67.4%)			
Fasting glucose ≥ 100 mg per 100 ml	113 (17.9%)	0 (-%)	36 (17.1%)	43 (41.8%)	34 (73.9%)			

Values shown are mean ± s.d.

<sup>a</sup>Statistical significance was determined by analysis of variance or  $\chi^2$  test.

significance, higher daily intakes of cholesterol (379.2–404.6 mg) and eggs (52.5–58.2 g) were related to an increased number of MetS components (trend test  $P < 0.1$ ).

In women (Table 3), intake of calcium decreased from 609.2 to 549.9 mg daily, and intake of milk and dairy food decreased from 181.2 to 134.9 mg daily as the number of MetS components increased. Cereal intake increased from 382.3 to 418.4 g daily as the number of MetS components increased. Although analysis of covariance and trend test did not reach statistical significance, lower intake of saturated fat was related to an increased number of MetS components, and decreased from 15.7 to 14.8 g daily as MetS components increased (trend test  $P < 0.1$ ).

**DISCUSSION**

Our findings suggest that lower intakes of vitamin B6 and dietary fiber in men, and lower intakes of calcium, milk and dairy products and higher intake of cereal in women are related to the number of MetS components. To our knowledge, this is the first observational study to examine relations between dietary factors and the number of clustering MetS components among Japanese men and women.

In previous epidemiologic studies, dietary fiber, fruits, vegetables and moderate alcohol intake were negatively associated<sup>22-24</sup> and fat

and red meat were positively associated<sup>25-27</sup> with MetS in Caucasian or Japanese-Brazilian subjects. Consistent with previous studies,<sup>22,24,28</sup> intake of dietary fiber in men was negatively related to clustering MetS components. Diets rich in dietary fiber are associated with a reduced risk of diabetes and cardiovascular disease. Dietary fiber has a higher satiety value compared with digestible complex carbohydrates and simple sugars because of its bulk and relatively low energy. Fiber may also affect secretion of gut hormones or peptides, such as cholecystokinin or glucagon-like peptid-1, which may act as satiety factors or alter glucose homeostasis.<sup>28,29</sup> Thus, an increased fiber intake may prevent MetS.

Although carbohydrate intake in women did not correlate with the number of MetS components, cereals that mainly consisted of carbohydrates showed a negative effect on the number of MetS components in this study. Carbohydrates are also implicated in changes in blood glucose and insulin concentrations and are known to affect satiety.<sup>30</sup> The beneficial effect of a high carbohydrate diet on glucose tolerance has been reported;<sup>31</sup> however, contradicting reports have also been published.<sup>32-36</sup> Dietary carbohydrate through cereal intake is thought to modulate lipolysis, and a low-carbohydrate diet reduces cardiovascular risk through improvement in hepatic, intravascular and peripheral processing of lipoproteins.<sup>32</sup> Although no positive relation was

**Table 2** Energy and multivariate adjusted<sup>a,b</sup> mean food and nutrient intake according to the number of metabolic syndrome (MetS) components in men (n=609)

Variable	Number of MetS components				P <sup>c</sup>	Trend P
	0	1	2	3-5		
n (%)	222 (36.5%)	209 (34.3%)	117 (19.2%)	61 (10.0%)		
Energy (kcal)	2247 ± 23	2233 ± 23	2283 ± 31	2227 ± 43	0.59	0.95
<b>Nutrients<sup>a</sup></b>						
Protein (energy %)	3.71 ± 0.03	3.71 ± 0.03	3.71 ± 0.05	3.70 ± 0.06	0.99	0.93
Fat (energy %)	2.71 ± 0.03	2.67 ± 0.04	2.66 ± 0.05	2.79 ± 0.07	0.35	0.36
Carbohydrate (energy %)	13.8 ± 0.1	14.0 ± 0.1	14.0 ± 0.2	13.8 ± 0.2	0.62	0.98
<b>Nutrients<sup>b</sup></b>						
Protein (g)	82.9 ± 0.7	82.8 ± 0.8	83.5 ± 1.0	82.5 ± 1.4	0.93	0.92
Fat (g)	61.2 ± 0.8	60.3 ± 0.8	60.4 ± 1.1	62.8 ± 1.5	0.46	0.33
Carbohydrate (g)	310.3 ± 2.4	312.7 ± 2.5	312.1 ± 3.3	309.8 ± 4.6	0.89	0.88
Calcium (mg)	619.5 ± 12.4	604.2 ± 12.6	590.1 ± 17.0	588.8 ± 23.5	0.47	0.20
β-Carotene (μg)	3269 ± 118	3274 ± 121	2804 ± 163	3003 ± 225	0.08	0.11
Vitamin E (mg)	9.8 ± 0.2	9.5 ± 0.2	9.1 ± 0.2	9.3 ± 0.3	0.130	0.10
Vitamin B6 (mg)	1.36 ± 0.02	1.35 ± 0.02	1.29 ± 0.03	1.21 ± 0.04	0.006	0.001
Vitamin B12 (μg)	8.8 ± 0.4	8.2 ± 0.4	9.4 ± 0.5	8.0 ± 0.7	0.23	0.70
Folate (μg)	343.6 ± 7.3	352.6 ± 7.5	339.9 ± 10.1	340.4 ± 13.9	0.70	0.65
Vitamin C (mg)	126.4 ± 7.1	127.2 ± 7.2	135.1 ± 9.7	125.1 ± 13.4	0.89	0.93
Saturated fat (g)	16.5 ± 0.3	15.9 ± 0.3	16.0 ± 0.4	16.1 ± 0.5	0.58	0.57
Monounsaturated fat (g)	21.2 ± 0.4	20.8 ± 0.4	20.8 ± 0.5	21.5 ± 0.7	0.66	0.70
Polyunsaturated fat (g)	13.3 ± 0.2	13.4 ± 0.2	13.5 ± 0.3	14.0 ± 0.4	0.55	0.15
Cholesterol (mg)	379.2 ± 8.8	369.0 ± 9.0	396.7 ± 12.2	404.6 ± 16.8	0.15	0.08
Dietary fiber (g)	16.2 ± 0.3	15.9 ± 0.3	14.8 ± 0.4	14.5 ± 0.5	0.002	0.001
Salt (g)	11.8 ± 0.1	12.1 ± 0.1	11.9 ± 0.2	11.6 ± 0.3	0.29	0.28
<b>Foods</b>						
Cereals (g)	512.2 ± 7.5	522.6 ± 7.7	538.3 ± 10.3	512.9 ± 14.2	0.21	0.72
Beans (g)	76.5 ± 4.7	86.9 ± 4.8	86.6 ± 6.4	82.8 ± 8.9	0.41	0.55
Nuts and seeds (g)	6.0 ± 0.6	4.0 ± 0.6	3.9 ± 0.8	5.4 ± 1.2	0.07	0.61
Vegetables (g)	301.1 ± 7.9	304.5 ± 8.1	277.0 ± 10.9	271.9 ± 15.0	0.07	0.03
Fruits (g)	137.1 ± 7.1	130.8 ± 7.3	122.0 ± 9.8	134.3 ± 13.5	0.66	0.72
Fish and shellfish (g)	111.2 ± 3.5	107.1 ± 3.6	109.3 ± 4.9	108.3 ± 6.7	0.88	0.79
Meats (g)	86.2 ± 3.1	89.4 ± 3.2	93.9 ± 4.2	95.0 ± 5.9	0.38	0.13
Eggs (g)	52.5 ± 1.9	53.5 ± 1.9	59.5 ± 2.6	58.2 ± 3.6	0.11	0.07
Milk and dairy food (g)	163.6 ± 8.0	144.3 ± 8.1	140.7 ± 10.9	129.8 ± 15.1	0.12	0.049
Fats and oils (g)	11.3 ± 0.4	11.6 ± 0.4	11.6 ± 0.6	11.6 ± 0.8	0.97	0.77
Confectioneries (g)	45.1 ± 3.3	46.6 ± 3.4	43.2 ± 4.5	46.7 ± 6.3	0.94	0.95

Values shown are mean ± s.e.  
<sup>a</sup>Adjusted for age, alcohol intake, smoking and physical activity.  
<sup>b</sup>Adjusted for age, energy intake, alcohol intake, smoking and physical activity.  
<sup>c</sup>Statistical significance was determined by analysis of covariance.

shown between carbohydrate intake and the number of MetS components in this study, recent nutritional reviews indicate that the quantity and type of carbohydrate affect metabolic outcomes.<sup>35</sup> Among cereals, whole grain products that have a lower glycemic index and are richer in fiber and antioxidant vitamins than refined grain products were suggested to improve insulin sensitivity, probably by blunting postprandial glycemic and insulinemic responses.<sup>36</sup> Thus, control of these factors in future studies will be important to determine the most effective dietary approach to prevent metabolic disorders.

In Japan, there has been a significant reduction in the intake of cereals, and rice in particular, in recent decades.<sup>2,37</sup> On the other hand, dietary fat intake is increasing, and consumption of a more

Westernized diet is thought to be associated with the evident increase in diabetes mellitus and obesity. In women in this study, saturated fat intake weakly decreased as the number of MetS components increased. Dietary intervention studies show that total fat is not associated with the risk of MetS, although saturated fats increase the risk of MetS, whereas monounsaturated and polyunsaturated fats reduce this risk.<sup>38-40</sup> Our results do not agree with studies that show that saturated fat-rich lipid infusion reduces the insulin sensitivity index more than polyunsaturated fat infusion.<sup>41</sup> Although the reason of this inverse relation was shown in this study is not clear, two possibilities can be considered. First, some intermediate event, such as dietary counseling, could have lead to changes in diet and might have confounded the association between saturated fat intake and metabolic

**Table 3** Energy and multivariate adjusted<sup>a,b</sup> mean food and nutrient intake according to the number of metabolic syndrome (MetS) components in women (n=631)

Variable	Number of MetS components				P <sup>c</sup>	Trend P
	0	1	2	3-5		
n (%)	272 (43.1%)	210 (33.3%)	103 (16.3%)	46 (7.3%)		
Energy (kcal)	1884±18	1878±20	1893±29	1896±43	0.97	0.72
<b>Nutrients<sup>a</sup></b>						
Protein (energy%)	3.75±0.03	3.84±0.03	3.79±0.05	3.70±0.07	0.15	0.45
Fat (energy%)	2.97±0.03	2.94±0.03	2.93±0.05	2.91±0.07	0.85	0.46
Carbohydrate (energy%)	14.2±0.1	14.2±0.1	14.1±0.2	14.2±0.2	0.97	0.99
<b>Nutrients<sup>b</sup></b>						
Protein (g)	70.2±0.5	71.9±0.6	71.4±0.9	69.7±1.3	0.16	0.64
Fat (g)	56.4±0.6	55.7±0.7	55.5±0.9	54.7±1.4	0.67	0.28
Carbohydrate (g)	267.4±1.8	266.2±2.0	265.4±2.9	268.2±4.3	0.91	0.93
Calcium (mg)	609.2±11.0	604.8±12.3	556.3±17.9	549.9±26.4	0.024	0.01
β-Carotene (μg)	3358±112	3263±125	2931±181	3508±268	0.18	0.90
Vitamin E (mg)	9.1±0.1	9.0±0.2	9.0±0.2	9.4±0.3	0.70	0.35
Vitamin B6 (mg)	1.13±0.02	1.19±0.02	1.13±0.03	1.15±0.04	0.11	0.98
Vitamin B12 (μg)	6.9±0.3	7.0±0.4	7.4±0.5	6.9±0.8	0.87	0.95
Folate (μg)	333.9±6.8	342.7±7.6	334.2±11.0	360.0±16.3	0.45	0.21
Vitamin C (mg)	121.7±4.2	125.6±4.7	111.0±6.9	131.7±10.1	0.24	0.66
Saturated fat (g)	15.7±0.2	15.5±0.3	15.0±0.4	14.8±0.6	0.23	0.07
Monounsaturated fat (g)	19.1±0.3	19.0±0.3	19.1±0.4	18.8±0.6	0.96	0.74
Polyunsaturated fat (g)	12.1±0.2	12.0±0.2	12.2±0.3	12.2±0.4	0.91	0.67
Cholesterol (mg)	321.1±6.7	339.2±7.4	343.1±10.8	318.7±15.9	0.16	0.95
Dietary fiber (g)	15.2±0.2	15.7±0.3	15.2±0.4	15.5±0.5	0.49	0.82
Salt (g)	10.4±0.1	10.4±0.1	10.6±0.2	10.8±0.3	0.45	0.11
<b>Foods<sup>b</sup></b>						
Cereals (g)	382.3±5.2	385.8±5.8	400.3±8.4	418.4±12.4	0.03	0.004
Beans (g)	75.4±4.1	80.7±4.6	76.0±6.6	83.1±9.8	0.77	0.58
Nuts and seeds (g)	6.3±0.7	6.9±0.7	6.2±1.1	5.3±1.6	0.81	0.48
Vegetables (g)	285.4±6.5	285.1±7.3	275.2±10.6	322.6±15.6	0.09	0.06
Fruits (g)	152.1±6.7	160.9±7.5	145.3±10.9	131.0±16.1	0.32	0.15
Fish and shellfish (g)	81.9±2.5	89.3±2.8	90.7±4.1	84.2±6.1	0.16	0.69
Meats (g)	65.5±2.0	70.8±2.3	71.1±3.3	66.1±4.9	0.27	0.90
Eggs (g)	48.2±1.6	52.1±1.8	53.1±2.6	50.6±3.9	0.33	0.54
Milk and dairy food (g)	181.2±6.7	177.0±7.4	153.7±10.8	134.9±16.0	0.018	0.003
Fats and oils (g)	11.2±0.3	10.7±0.4	10.6±0.6	10.2±0.8	0.64	0.28
Confectioneries (g)	65.5±2.7	61.6±3.0	60.1±4.4	56.9±6.5	0.52	0.21

Values shown are mean ± s.e.  
<sup>a</sup>Adjusted for age, alcohol intake, smoking and physical activity.  
<sup>b</sup>Adjusted for age, energy intake, alcohol intake, smoking and physical activity.  
<sup>c</sup>Statistical significance was determined by analysis of covariance.

risks. We tried to exclude subjects from the analysis who were aware of their potential risks and who may have made dietary changes based on perceived dangers. Second, Japanese subjects consume a relatively large amount of fish containing abundant polyunsaturated fatty acids. The ratio of saturated fat to polyunsaturated fat might be more important than the absolute intake considering the physiologic dietary effect on developing MetS. These factors might have affected our results. Further studies are needed to clarify the role of fat quality in the prevention of MetS.

Calcium, milk and dairy food intake in women decreased with the increase in the number of MetS components. Additionally, although findings were not statistically significant, a similar relation between milk and dairy products and MetS was shown in men (ANCOVA

$P=0.12$ , trend  $P=0.049$ ). Diets rich in calcium, particularly calcium derived from dairy products, have been shown to be associated with a low prevalence of MetS.<sup>42</sup> The mechanism by which calcium intake can reduce MetS is unclear, but Scholz-Ahrens and Schrenzenmeir<sup>42</sup> implied that dietary calcium intake has benefits on traits of MetS, specifically on weight reduction and fat loss. Meijl *et al.*<sup>43</sup> reviewed the physiological effects of three main dairy constituents (calcium, protein and fat) on MetS, and indicated that the effects of calcium might be related to intestinal binding to fatty acids or bile acids or to changes in intracellular calcium metabolism by suppressing calciotropic hormones. In an epidemiologic study, Otsuka *et al.*<sup>44</sup> reported that higher milk consumption was associated with a lower incidence of MetS after 5 years among middle-aged

Japanese male workers, suggesting that calcium derived from dairy products might help prevent MetS.

Vitamin B6 in men was also negatively related to the number of MetS components. Esmailzadeh *et al.*<sup>45</sup> discussed the favorable effect of whole grain on MetS through the rich content of viscous fiber and showed that intake of whole grains was positively associated not only with dietary fiber ( $r=0.43$ ) but also with vitamin B6 ( $r=0.48$ ). Consistent with their study, the results of our study showed favorable effects of dietary fiber and vitamin B6 on the number of MetS components. Hayden and Tyagi<sup>46</sup> reported on the role of water soluble B vitamins, including vitamin B6, in lowering plasma total homocysteine, a risk marker of MetS, through remethylation. Although the precise mechanism of vitamin B6 on MetS is unclear, a favorable association of vitamin B6 to MetS in men may be attributed to a healthier diet that contained rich fiber, milk and dairy products or vegetables.

Although not statistically significant, lower intake of vegetables in men was related to an increased number of MetS components. Higher vegetable intake has been previously reported as a protective factor of MetS or inflammation in women.<sup>7,22</sup> Vegetables rich in dietary fiber are thought to reduce the risk of developing MetS by improving glucose control, and minerals, antioxidants or vitamins contained in vegetables are thought to have a favorable effect on glucose tolerance.<sup>47,48</sup> Our results regarding dietary fiber on MetS may represent the positive effect of vegetable intake.

In men, higher intakes of eggs and cholesterol were related to an increased number of MetS components, although this was not statistically significant (trend  $P<0.1$ ). Excess consumption of eggs should be avoided from the standpoint of preventing hypercholesterolemia among Japanese subjects.<sup>49</sup> Increased dietary cholesterol intake is associated with atherosclerosis.<sup>50</sup> The result of this study may suggest that men with a lower number of MetS components tend to abstain from eating eggs or cholesterol-rich foods. As a result, a dose-response relationship between egg consumption and the number of MetS components was shown in our study.

The lack of other correlations between the number of MetS components and dietary factors, such as fruits, or meats in this study is thought to be due to differences in dietary intake<sup>51</sup> or metabolic responses<sup>9,10</sup> between Caucasian and Japanese subjects.

The relation of diet to the risk of MetS has been examined in many studies, but few have addressed the association with the number of MetS components.<sup>52-54</sup> We considered that the number of MetS components is more closely assessed through the effects of diet, that is, through a dose-response relationship, rather than the prevalence of MetS. In addition, to focus on the dose-response relationship, trend test by general linear model was used, and further *post hoc* analyses were not performed. MetS is a cluster of atherosclerotic cardiovascular disease risk factors<sup>20</sup> but there are some different criteria of MetS in Japan. The ideal threshold for waist circumference used to define abdominal obesity among Japanese men and women is still under discussion.<sup>55</sup> Recently, Kokubo *et al.*<sup>19</sup> reported that the number of MetS components (modified NCEP-ATP III criteria) might be more strongly associated with the incidence of cardiovascular disease than the presence of abdominal obesity (the Japanese criteria) in a general urban Japanese population. We used a modified NCEP-ATP III criteria, that is, a cutoff point of waist circumference that was different from NCEP ATP III (102 cm in men and 88 cm in women, modified: 90 cm in men and 80 cm in women), in this study.<sup>20,21</sup>

Several limitations of this study warrant consideration. First, the cross-sectional nature of the study did not permit the assessment of causality. Subjects under treatment for hypertension,

hypertriglyceridemia or diabetes or those aware of morbidities might have modified their food intake, and these subjects were excluded from analysis. However, some plausible relationships between foods or nutrients and MetS have been identified in this study. NILS-LSA, which is a population-based prospective cohort study, followed participants for more than 10 years. Future analyses should examine the associations between the dietary index and MetS.

Second, we used foods and nutrients as an indicator of diet, although these items are consumed in combination, and their complex effects are likely to be interactive or synergistic.<sup>56</sup> Consumption of several foods or food groups might be a more comprehensive variable to assess the impact of diet on disease risk than any single nutrient or food. Third, nutritional intakes were assessed by a 3-day dietary records. The 3-day dietary record is one of the most reliable methods for nutritional assessment; however, it is limited because individual food intake varies greatly from day to day,<sup>15</sup> and it is not clear whether short-term records adequately reflect long-term dietary intake.<sup>57</sup> On the basis of this limitation, we preliminarily decided on 3 continuous days (both weekend days and one weekday) to avoid events or special days such as trips, long vacations or out-of-the-ordinary events and thus minimize food variations. Although the 3-day dietary record is not the best way to assess long-term dietary intake, it can be considered to have a certain level of accuracy that reflects the usual nutrient intakes in this cohort.

Fourth, food consumption and nutrient intake among Japanese have dramatically changed during the past five decades. For example, Westernization of the Japanese diet has led to decreased consumption of carbohydrates, especially rice, and increased consumption of fat and meat.<sup>2,37</sup> In particular, younger Japanese subjects tend to eat a more Westernized diet.<sup>37</sup> As a result, not only food intake but also the number of MetS components might differ among different generations of Japanese subjects. Thus, it may be difficult to detect statistically significant relationships between diet and the number of MetS components in the age group we studied.

In summary, this study showed that some foods and nutritional components are related to the number of MetS components. These results suggest the potential effect of diet on the prevention of MetS among community-dwelling Japanese men and women.

#### CONFLICT OF INTEREST

The authors declare no conflict of interest.

#### ACKNOWLEDGEMENTS

We express our sincere appreciation to study participants and colleagues in the NILS-LSA in conducting the survey for this study. This work was supported in part by grants to Rei Otsuka from the Research Fellowships of Japan Society for the Promotion of Science for Young Scientists. This work was supported in part by grants to Rei Otsuka from the Research Fellowships of the Japan Society for the Promotion of Science for Young Scientists.

- 1 Lakka HM, Laaksonen DE, Lakka TA, Niskanen LK, Kumpusalo E, Tuomilehto J, Salonen JT. The metabolic syndrome and total and cardiovascular disease mortality in middle-aged men. *JAMA* 2002; 288: 2709-2716.
- 2 Ministry of Health Welfare. *Annual Report of the National Nutrition Survey in 2006*. [in Japanese] Daiichi Publishing Co: Tokyo, Japan, 2009.
- 3 Udagawa K, Miyoshi M, Yoshikae N. Mid-term evaluation of 'Health Japan 21': focus area for the nutrition and diet. *Asia Pac J Clin Nutr* 2008; 17(Suppl 2): 445-452.
- 4 Eckel RH, Grundy SM, Zimmet PZ. The metabolic syndrome. *Lancet* 2005; 365: 1415-1428.
- 5 Park YW, Zhu S, Palaniappan L, Heshka S, Carnethon MR, Heymsfield SB. The metabolic syndrome—prevalence and associated risk factor findings in the US popula-



- tion from the Third National Health and Nutrition Examination Survey, 1988-1994. *Arch Intern Med* 2003; 163: 427-436.
- 6 Esposito K, Marfella R, Ciotola M, Di Palo C, Giugliano F, Giugliano G, D'Armiento M, D'Andrea F, Giugliano D. Effect of a Mediterranean-style diet on endothelial dysfunction and markers of vascular inflammation in the metabolic syndrome—a randomized trial. *JAMA* 2004; 292: 1440-1446.
- 7 Millen BE, Pencina MJ, Kimokoti RW, Zhu L, Meigs JB, Ordovas JM, D'Agostino RB. Nutritional risk and the metabolic syndrome in women: opportunities for preventive intervention from the Framingham Nutrition Study. *Am J Clin Nutr* 2006; 84: 434-441.
- 8 Maki KC. Dietary factors in the prevention of diabetes mellitus and coronary artery disease associated with the metabolic syndrome. *Am J Cardio* 2004; 93 (11A): 12C-17C.
- 9 Kanazawa M, Yoshitake N, Osaka T, Numba Y, Zimmet P, Inoue S. Criteria and classification of obesity in Japan and Asia-Oceania. *Asia Pac J Clin Nutr* 2002; 11(Suppl 8): 732-737.
- 10 Ferreira SR, Lavarro DB, Gimeno SG, Sanudo A, Franco LJ. Japanese-Brazilian Diabetes Study Group. Obesity and central adiposity in Japanese immigrants: role of the Western dietary pattern. *J Epidemiol* 2002; 12: 431-438.
- 11 Damilo R, Castro TG, Cardoso MA, Gimeno SG, Ferreira SR. Dietary intakes associated with metabolic syndrome in a cohort of Japanese ancestry. *Br J Nutr* 2006; 96: 532-538.
- 12 Fujimoto WY, Bergstrom RW, Boyko EJ, Chen K, Kahn SE, Leonetti DL, McNeely MJ, Newell LL, Shofer JB, Wahl PW. Type 2 diabetes and the metabolic syndrome in Japanese Americans. *Diabetes Res Clin Pract* 2000; 50: 73-76.
- 13 Kawada T, Okada K, Amazawa M. Components of the metabolic syndrome and lifestyle factors in Japanese male workers. *Metab Syndr Relat Disord* 2008; 6: 263-266.
- 14 Shimokata H, Ando F, Niino N. A new comprehensive study on aging—the National Institute for Longevity Sciences, Longitudinal Study of Aging (NILS-LSA). *J Epidemiol* 2000; 10(Suppl 1): S1-S9.
- 15 Imai T, Sakai S, Mori K, Ando F, Niino N, Shimokata H. Nutritional assessments of 3-day dietary records in National Institute for Longevity Sciences—Longitudinal Study of Aging (NILS-LSA). *J Epidemiol* 2000; 10(Suppl 1): S70-S76.
- 16 Examination Committee of Criteria for 'Obesity Disease' in Japan; Japan Society for the Study of Obesity. New criteria for 'obesity disease' in Japan. *Circ J* 2002; 66: 987-992.
- 17 Yoshizumi T, Nakamura T, Yamane M, Islam AH, Menju M, Yamasaki K, Arai T, Kotani K, Funahashi T, Yamashita S, Matsuzawa Y. Abdominal fat: standardized technique for measurement at CT. *Radiology* 1999; 211: 283-286.
- 18 Iwai N, Yoshiike N, Saiboh S, Nose T, Kushiro T, Tanaka H. Japan Lifestyle Monitoring Study Group: leisure-time physical activity and related lifestyle characteristics among middle-aged Japanese. *J Epidemiol* 2000; 10: 226-233.
- 19 Kokubo Y, Okamura T, Yoshimasa Y, Miyamoto Y, Kawanishi K, Kotani Y, Okayama A, Tomoike H. Impact of metabolic syndrome components on the incidence of cardiovascular disease in a general urban Japanese population: the Suita study. *Hypertens Res* 2008; 31: 2027-2035.
- 20 Grundy SM, Cleeman JI, Daniels SR, Donato KA, Eckel RH, Franklin BA, Gordon DJ, Krauss RM, Savage PJ, Smith Jr SC, Spertus JA, Costa F. Diagnosis and management of the metabolic syndrome: an American Heart Association/National Heart, Lung, and Blood Institute Scientific Statement. *Circulation* 2005; 112: 2735-2752.
- 21 James PT, Leach R, Kalamara E, Shayeghi M. The worldwide obesity epidemic. *Obes Res* 2001; 9: 228S-233S.
- 22 Giugliano D, Ciriello A, Esposito K. The effects of diet on inflammation: emphasis on the metabolic syndrome. *J Am Coll Cardio* 2006; 48: 677-685.
- 23 Williams DE, Prevost AT, Whitchelov MJ, Cox BD, Day NE, Wareham NJ. A cross-sectional study of dietary patterns with glucose intolerance and other features of the metabolic syndrome. *Br J Nutr* 2000; 83: 257-266.
- 24 McKeown NM, Meigs JB, Liu S, Salzman E, Wilson PW, Jacques PF. Carbohydrate nutrition, insulin resistance, and the prevalence of the metabolic syndrome in the Framingham Offspring Cohort. *Diabetes Care* 2004; 27: 538-546.
- 25 Millen BE, Pencina MJ, Kimokoti RW, Zhu L, Meigs JB, Ordovas JM, D'Agostino RB. Nutritional risk and the metabolic syndrome in women: opportunities for preventive intervention from the Framingham Nutrition Study. *Am J Clin Nutr* 2006; 84: 434-441.
- 26 Sonnenberg L, Pencina M, Kimokoti R, Quatromoni P, Nam BH, D'Agostino R, Meigs JB, Ordovas J, Cobain M, Millen B. Dietary patterns and the metabolic syndrome in obese and non-obese Framingham women. *Obes Res* 2005; 13: 153-162.
- 27 Freire RD, Cardoso MA, Gimeno SG, Ferreira SR. Dietary fat is associated with metabolic syndrome in Japanese Brazilians. *Diabetes Care* 2005; 28: 1779-1785.
- 28 Davy BM, Mealy CL. The effect of fiber-rich carbohydrates on features of Syndrome X. *J Am Diet Assoc* 2003; 103: 86-96.
- 29 Gelfisio M, Duarte J, Zarzuelo A. Effects of dietary fibers on disturbances clustered in the metabolic syndrome. *J Nutr Biochem* 2008; 19: 71-84.
- 30 Fainle C, O'Donovan D, Horowitz M. Carbohydrate and satiety. *Nutr Rev* 2002; 60: 155-169.
- 31 Howard BV, Abbott WG, Swinburn BA. Evaluation of metabolic effects of substitution of complex carbohydrates for saturated fat in individuals with obesity and NIDDM. *Diabetes Care* 1991; 14: 786-795.
- 32 Volek JS, Fernandez ML, Feinman RD, Phinney SD. Dietary carbohydrate restriction induces a unique metabolic state positively affecting atherogenic dyslipidemia, fatty acid partitioning, and metabolic syndrome. *Prog Lipid Res* 2008; 47: 307-318.
- 33 Kim K, Yun SH, Choi BY, Kim MK. Cross-sectional relationship between dietary carbohydrate, glycaemic index, glycaemic load and risk of the metabolic syndrome in a Korean population. *Br J Nutr* 2008; 100: 576-584.
- 34 Lara-Castro C, Garvey WT. Diet, insulin resistance, and obesity: zoning in on data for Atkins dieters living in South Beach. *J Clin Endocrinol Metab* 2004; 89: 4197-4205.
- 35 Wood RJ, Fernandez ML. Carbohydrate-restricted versus low-glycemic-index diets for the treatment of insulin resistance and metabolic syndrome. *Nutr Rev* 2009; 67: 179-183.
- 36 Pereira MA, Jacobs DR Jr, Pins JJ, Raatz SK, Gross MD, Slavin JL, Seaquist ER. Effect of whole grains on insulin sensitivity in overweight hyperinsulinemic adults. *Am J Clin Nutr* 2002; 75: 848-855.
- 37 Shinbo S, Kawamura S, Yamamoto K, Kimura K, Imai Y, Yasumoto M, Watanabe T, Iwai O, Ikeda M. Reduced carbohydrate intake in past 10 years in 2 rural-areas in Japan. *Ecol Food Nutr* 1994; 33: 123-130.
- 38 Phillips C, Lopez-Miranda J, Perez-Jimenez F, McManus R, Roche HM. Genetic and nutrient determinants of the metabolic syndrome. *Curr Opin Cardio* 2006; 21: 185-193.
- 39 Riccardi G, Giacco R, Rivellese AA. Dietary fat, insulin sensitivity and the metabolic syndrome. *Clin Nutr* 2004; 23: 447-456.
- 40 Chen X, Pang Z, Li K. Dietary fat, sedentary behaviors and the prevalence of the metabolic syndrome among Qingdao adults. *Nutr Metab Cardiovasc Dis* 2009; 19: 27-34.
- 41 Stefan N, Wahl HG, Fritsche A, Haring H, Stumvoll M. Effect of the pattern of elevated free fatty acids on insulin sensitivity and insulin secretion in healthy humans. *Horm Metab Res* 2001; 33: 432-438.
- 42 Scholz-Ahrens KE, Schrezenmeir J. Milk minerals and the metabolic syndrome. *Int Dairy J* 2006; 16: 1399-1407.
- 43 Meiji LE, Wrolix R, Mensink RP. Dairy product consumption and the metabolic syndrome. *Nutr Res Rev* 2008; 21: 148-157.
- 44 Otsuka R, Yamakoshi K, Shimokata H, Toyoshima H, Yatsuya H. Dietary habits and incidence of metabolic syndrome among middle-aged Japanese male workers. *J Jpn Soc Nutr Food Sci* 2009; 62: 123-129. (in Japanese).
- 45 Esmailizadeh A, Mirmiran P, Azizi F. Whole-grain consumption and the metabolic syndrome: a favorable association in Tehranian adults. *Eur J Clin Nutr* 2005; 59: 353-362.
- 46 Hayden MR, Tyagi SC. Homocysteine and reactive oxygen species in metabolic syndrome, type 2 diabetes mellitus, and atherosclerosis: the pleiotropic effects of folate supplementation. *Nutr J* 2004; 10: 3-4.
- 47 Yoo S, Nicklas T, Baranowski T, Zakeri IF, Yang SJ, Srinivasan SR, Berenson GS. Comparison of dietary intakes associated with metabolic syndrome risk factors in young adults: the Bogalusa Heart Study. *Am J Clin Nutr* 2004; 80: 841-848.
- 48 Ford ES, Mokdad AH, Giles WH, Brown DW. The metabolic syndrome and antioxidant concentrations: findings from the Third National Health and Nutrition Examination Survey. *Diabetes* 2003; 52: 2346-2352.
- 49 Nakamura Y, Iso H, Kita Y, Ueshima H, Okada K, Konishi M, Inoue M, Tsugana S. Egg consumption, serum total cholesterol concentrations and coronary heart disease incidence: Japan Public Health Center-based prospective study. *Br J Nutr* 2006; 96: 921-928.
- 50 Kleemann R, Verschuren L, van Erk MJ, Nikolsky Y, Crubben NH, Verheij ER, Smilde AK, Hendriks HF, Zedelaar S, Smith GJ, Kaznatcheev V, Nikolakaya T, Melnikov A, Hurt-Camejo E, van der Greef J, van Ommen B, Kooistra T. Atherosclerosis and liver inflammation induced by increased dietary cholesterol intake: a combined transcriptomics and metabolomics analysis. *Genome Biol* 2007; 8: R200.
- 51 Kagawa M, Saito Y, Kerr D, Uchida H, Binns CW. Differences in nutrient intakes and physical activity levels of Japanese and Australian Caucasian males living in Australia and Japanese males living in Japan. *Asia Pac J Clin Nutr* 2006; 15: 208-216.
- 52 McKeown NM, Meigs JB, Liu S, Salzman E, Wilson PW, Jacques PF. Carbohydrate nutrition, insulin resistance, and the prevalence of the metabolic syndrome in the Framingham Offspring Cohort. *Diabetes Care* 2004; 27: 538-546.
- 53 Ford ES, Giles WH, Dietz WH. Prevalence of the metabolic syndrome among US adults: findings from the third National Health and Nutrition Examination Survey. *JAMA* 2002; 287: 356-359.
- 54 Evangelopoulos AA, Vailianou NG, Panagiotakos DB, Georgiou A, Zacharias GA, Alevis AN, Zisakosta GJ, Vogiatzakis ED, Angelopoulos PC. An inverse relationship between circulating components of the metabolic syndrome and serum magnesium levels. *Nutr Res* 2008; 28: 659-663.
- 55 Kashiwara H, Lee JS, Kawakubo K, Tamura M, Akabayashi A. Criteria of waist circumference according to computed tomography-measured visceral fat area and the clustering of cardiovascular risk. *Circ J* 2009; 73: 1881-1886.
- 56 Hu FB. Dietary pattern analysis: a new direction in nutritional epidemiology. *Curr Opin Lipidol* 2002; 13: 3-9.
- 57 Hu FB. *Dietary Assessment Methods: Obesity Epidemiology*, Vol. 86. Oxford University Press: New York, 2008, pp 95-98.

Received: 2009.12.18  
Accepted: 2010.03.12  
Published: 2010.07.01

## Diabetes reduces auditory sensitivity in middle-aged listeners more than in elderly listeners: A population-based study of age-related hearing loss

PH

### Authors' Contribution:

- A** Study Design
- B** Data Collection
- C** Statistical Analysis
- D** Data Interpretation
- E** Manuscript Preparation
- F** Literature Search
- G** Funds Collection

Yasue Uchida<sup>1ACDEI</sup>, Salko Suglura<sup>1BE</sup>, Fujiko Ando<sup>2,3DE</sup>, Tsutomu Nakashima<sup>4DE</sup>, Hiroshi Shimokata<sup>3BCEG</sup>

<sup>1</sup> Department of Otorhinolaryngology, National Center for Geriatrics and Gerontology, Aichi, Japan

<sup>2</sup> Faculty of Medical Welfare, Aichi Shukutoku University, Aichi, Japan

<sup>3</sup> Department of Epidemiology, National Center for Geriatrics and Gerontology, Aichi, Japan

<sup>4</sup> Department of Otorhinolaryngology, Nagoya University School of Medicine, Nagoya, Japan

**Source of support:** This work was partially supported by a Grant-in-Aid for Comprehensive Research on Aging and Health (H17-choju-ippan-033), and by the Research Grant for Longevity Sciences (21A-17), from the Ministry of Health, Labour and Welfare of Japan

### Summary

#### Background:

Diabetes mellitus (DM) and hearing impairment are both highly prevalent in older adult populations, but how the impact of diabetes on hearing varies by age is not well-studied.

#### Material/Methods:

The subjects were 2306 adults aged 40 to 86 years who participated in a population-based study of aging, and were divided into 2 age groups, 40–64 years and 65–86 years, for cross-sectional analysis. Air-conduction pure-tone thresholds at octave intervals from 125 to 8000 Hz were obtained. Outcomes were categorized in relation to presence or absence of DM. Hearing levels at 7 frequencies were set in the general linear model as objective variables with adjustment for confounders. Explanatory variables were age (<65 years vs. >65 years), DM (absence vs. presence), and interaction between age and DM.

#### Results:

A statistically-significant adverse effect of DM on hearing was observed. This effect varied by age at the higher frequencies. The DM-age interaction was not synergistic at any test frequencies. No significant effects of the DM-age interaction were observed below 4000 Hz. In contrast, significant reciprocal effects of the DM-by-age interaction were found at 4000 Hz and 8000 Hz. Diabetes may accordingly affect higher-frequency hearing more strongly in the younger age-bracket.

#### Conclusions:

This study demonstrated that diabetes detrimentally affected hearing in community-dwelling middle-aged and elderly people, and that the effect of diabetes on higher-frequency hearing might be stronger in middle age. Screening for hearing impairment in diabetic patients may provide benefits for intervention or prevention of early presbycusis, particularly in this age group.

#### key words:

diabetes • aging • hearing • interaction • population-based study

#### Full-text PDF:

<http://www.medscimonit.com/fulltxt.php?ICID=880914>

#### Word count:

2508

#### Tables:

3

#### Figures:

1

#### References:

25

#### Author's address:

Yasue Uchida, Department of Otorhinolaryngology, National Center for Geriatrics and Gerontology, 36-3 Gengo, Morioka, Obu, Aichi 474-8511, Japan, e-mail: yasueu@ncgg.go.jp



## BACKGROUND

Diabetes mellitus (DM) is a prevalent disease in older people. DM in older adults has become a major public health problem, affecting an increasing number of individuals worldwide. According to a 2006 national nutrition survey in Japan, the rates of likely DM (HbA1c >6.1% or in treatment) were 3.2%, 10.2%, 13.6%, and 18.0% for people in their 40s, 50s, 60s, and 70s or over, respectively. The rates of subclinical DM (HbA1c 5.6–6.1%) were 14.2%, 19.0%, 23.1%, and 24.7% for people in their 40s, 50s, 60s, and 70s or over, respectively [1].

Hearing loss is also highly prevalent in older adult populations [2,3]. Research has shown that presbycusis affects about 30% of people aged 65 and over [4,5], and that about half of the population aged over 75 years has significant hearing loss [6].

A relationship between DM and hearing loss was first proposed in a case report by Jordao in 1857 [7]. Although this potential association has been investigated ever since then [8–13], it has not been as firmly established as the association between DM and its known complications affecting the renal, visual, and peripheral nervous systems. Several factors, such as noise exposure, presbycusis, and syndromic hearing loss, might confound the association between DM and hearing impairment. Although aging is thought to be a key factor in both glucose metabolism and cochlear function, it is difficult to demonstrate the interrelated contribution of DM and aging to hearing impairment.

We aimed to investigate the impact of DM on hearing, and the relationship between this and age, in middle-aged and elderly community-dwelling individuals.

## MATERIAL AND METHODS

### Subjects

The present study was conducted as part of the comprehensive 'Longitudinal Study of Aging (NILS-LSA)' conducted by the National Institute for Longevity Sciences of the National Center for Geriatrics and Gerontology. The NILS-LSA is a population-based biennial survey of a cohort of approximately 2200 people, which started in November 1997. The subjects of the NILS-LSA were randomly selected from resident registrations, stratified by both age and sex. Details of the methodology used in the NILS-LSA have been reported elsewhere [14]. The study protocol was reviewed by the Ethical Committee of the National Center for Geriatrics and Gerontology, and written informed consent was obtained from all participants.

Data obtained from the 4<sup>th</sup> wave examination of NILS-LSA were analyzed cross-sectionally in the present study. Participants were 2383 adults aged 40 to 86 years who took part in the NILS-LSA between May 2004 and July 2006. Demographic characteristics, personal history, family history, lifestyle habits, and medical history were obtained from detailed questionnaires filled out before the examination visit. Venous blood was collected early in the morning after at least 12 hours' fasting.

The definition of DM was based on medical history obtained from questionnaires, or defined as a fasting plasma glucose

concentration greater than 126 mg/dl and an HbA1c of more than 6.5%, or the taking of medication to lower the blood glucose level. In this way, participants were classified into a DM (+) group and a DM (-) group. Histories of ear disease and occupational noise exposure were obtained from the self-reported questionnaire. Occupational noise was defined in our questionnaires as background noise in a work environment over which the worker could not hold a conversation in a normal voice. Former and current noise exposures were combined. History of ear disease and history of occupational noise exposure were treated as binary variables (presence =1, absence =0).

Those who did not undergo blood testing, had incomplete hearing measurements, or provided invalid questionnaire responses were eliminated from the analysis. Accordingly, 2306 participants with complete data were selected for the present analysis. The subjects were divided into 2 groups for analysis: 40–64 and 65–86 years (Table 1).

### Audiometric measurements

Audiometric measurements were examined in a soundproof compartment by laboratory technicians on the same day of the blood draw. Air-conduction pure-tone thresholds at octave intervals from 125 to 8000 Hz were obtained using diagnostic audiometers (AA-73A and AA-78; Rion, Tokyo, Japan). The thresholds over the predetermined output level, according to the Japanese Industrial Standards T 1201 calibration, were treated as that level plus an additional 5 dB; that is to say, over 70 dB at 125 Hz was treated as 75 dB. For analyses of supraliminal levels we used 90 dB at 250 Hz, 105 dB at 500 to 4000 Hz, and 100 dB at 8000 Hz. Pure-tone averages (PTAs) were calculated for the better ear (BE) and the worse ear (WE) in order not to overlook subjects with at least 1 affected ear. The low-frequency PTA was calculated as the average threshold across the 125-, 250- and 500-Hz thresholds. The high-frequency PTA was calculated as the average across the 2000-, 4000- and 8000-Hz thresholds. The mid-frequency PTA was calculated as the average across the 500-, 1000-, 2000- and 4000-Hz thresholds. Hearing impairment was defined as PTAs greater than 25 dB.

### Statistical analyses

Statistical analyses were conducted using the Statistical Analysis System (SAS) version 9.13 (SAS Institute, Cary, NC, USA). All values are expressed as mean  $\pm$  standard error if not specified otherwise. Comparisons of hearing impairment rates between the DM (-) group and DM (+) group by age were performed using the chi-square test. In multivariable analyses, general linear model (GLM) analyses were performed to assess both the individual and the interactive impacts of age and DM on hearing of the BE and WE, based on the mid-frequency PTA. Hearing levels at 7 frequencies were set in the GLM as objective variables. Explanatory variables were age (binary variable; <65 years vs. >65 years), DM (binary variable; absence vs. presence), and interaction between age and DM. Moderator variables were sex, history of ear disease, and history of occupational noise exposure.

## RESULTS

Table 1 shows the clinicodemographic profile of the subjects by age group and DM status. A total of 2306 participants

**Table 1.** Clinicodemographic profile of the subjects by DM status and age group.

(mean ± SE)	40–64 years			65–86 years		
	DM (–)	DM (+)	p	DM (–)	DM (+)	p
N	1349	67		806	84	
Male (%)	50.1	64.2	0.0246	48.4	56.0	NS
Fasting blood glucose (mg/dl)	96.3±0.4	159.7±1.6	<.0001	98.9±0.6	146.2±1.8	<.0001
HbA1c (%)	5.2±0.01	7.4±0.06	<.0001	5.3±0.02	7.2±0.06	<.0001
History of ear disease (%)	41.4	34.3	NS	33.3	33.3	NS
History of occupational noise exposure (%)	19.4	16.4	NS	18.9	25.0	NS

DM – diabetes mellitus.

**Table 2.** The prevalence of hearing impairment based on the low-frequency, high-frequency and mid-frequency PTA.

Hearing impairment (%)	40–64 years			65–86 years			
	DM (–)	DM (+)	p	DM (–)	DM (+)	p	
Better ear	Low-frequency PTA <sub>125,250,500</sub> >25 dB	5.6	13.4	0.0087	37.0	44.1	NS
	High-frequency PTA <sub>2000,4000,8000</sub> >25 dB	16.8	44.8	<.0001	75.1	79.8	NS
	Mid-frequency PTA <sub>500,1000,2000,4000</sub> >25 dB	7.3	13.4	NS	52.0	59.5	NS
Worse ear	Low-frequency PTA <sub>125,250,500</sub> >25 dB	19.2	31.3	0.0149	60.3	66.7	NS
	High-frequency PTA <sub>2000,4000,8000</sub> >25 dB	29.0	59.7	<.0001	87.3	88.1	NS
	Mid-frequency PTA <sub>500,1000,2000,4000</sub> >25 dB	15.7	40.3	<.0001	66.4	75.0	NS

PTA pure tone averages; DM – diabetes mellitus; NS – not significant.

ages 40 through 86 years were analyzed in 4 subject groups: in the younger age bracket (40-64 years), 67 had diabetes and 1349 did not; in the older age-bracket (65–86 years) 84 had diabetes and 806 did not. The prevalence of DM was 4.7% in the younger participants, and 9.4% in the older participants.

Table 2 provides the results from chi-square analysis regarding the prevalence of hearing impairments (based on low-frequency, high-frequency and mid-frequency PTAs) according to age group and DM status. In the younger group, participants with DM had a significantly higher prevalence of hearing impairment than those without DM, for all frequency criteria except mid-frequency in the BE. In contrast, in the older age bracket, no significant differences were observed in the prevalence rates of any defined hearing impairments according to DM status.

The individual and interactive effects of age and DM on the hearing levels at 7 frequencies, from the GLM analysis, are shown in Table 3. The statistically significant main effect of DM was more moderate than that of age, but it was observed from low to high frequencies in both the BE and WE. The direction of each main effect was confirmed for the BE and WE in the lower sections of Table 3, which provide the mean hearing levels in the 4 groups: DM (–) and DM (+) in the younger and the older age-brackets. The main

effect of DM was to impair hearing levels at all frequencies except 125 Hz in the BE and 500Hz in the WE. Moreover, the interactive effect of age and DM was statistically significant at 4000 Hz and 8000 Hz in both the BE and WE. The adverse effect of DM on hearing varied according to age.

For the purpose of visual comparison, the interactive effect of age and DM in the BE at 8000 Hz are graphically presented in the Figure 1. The solitary main effect of DM on hearing level was strong in the younger age bracket but less obvious in the older age bracket.

**DISCUSSION**

We found a statistically significant harmful effect of DM on hearing and noted that this effect varied by age in the high frequencies. The effects of the DM-age interaction were not synergistic at any of the test frequencies; in other words, aging did not intensify the deleterious effects of DM on hearing. The DM-age interaction was additive at below 4000 Hz, but reciprocal at 4000 Hz and 8000 Hz. The impact of DM on hearing at 4000 Hz and 8000 Hz was more severe in the younger than the older participants. Hearing sensitivity at the higher frequencies is particularly vulnerable to ototoxic insults such as aging [4, 6], hazardous noise exposure [14], and ototoxic agents. Accordingly the present results can be explained by the fact that the effect of DM on



This copy is for personal use only - distribution prohibited.

**Table 3.** Relationship between hearing level, age, and DM status as assessed by general linear model analyses.

		Hearing level at 125 Hz		Hearing level at 250 Hz		Hearing level at 500 Hz		Hearing level at 1000 Hz		Hearing level at 2000 Hz		Hearing level at 4000 Hz		Hearing level at 8000 Hz	
<b>Better ear based on the mid-frequency</b>		F value	p value	F value	p value	F value	p value	F value	p value	F value	p value	F value	p value	F value	p value
	DM	3.13	NS	4.41	0.0359	6.41	0.0114	6.98	0.0083	5.35	0.0208	11.68	0.0006	18.96	<0.0001
	Age	496.5	<0.0001	445.3	<0.0001	509.1	<0.0001	648.4	<0.0001	907.7	<0.0001	1094	<0.0001	1536	<0.0001
	DM × age	1.49	NS	1.21	NS	1.39	NS	1.04	NS	0.89	NS	5.62	0.0179	8.39	0.0038
		Mean	95% CI	Mean	95% CI	Mean	95% CI	Mean	95% CI	Mean	95% CI	Mean	95% CI	Mean	95% CI
40–64 years	DM (–)	20.5	19.9–21.1	18.3	17.7–18.9	14.3	13.7–14.9	10.9	10.3–11.6	15.3	14.5–16.0	18.0	17.1–19.0	23.2	22.0–24.3
	DM (+)	22.9	20.7–25.0	20.9	18.6–23.2	17.4	15.1–19.8	14.3	11.7–16.8	18.6	15.7–21.6	25.7	22.0–29.3	34.7	30.3–39.2
65–86 years	DM (–)	29.3	28.6–30.0	26.9	26.2–27.7	23.8	23.1–24.6	22.6	21.8–23.4	30.9	30.0–31.9	39.8	38.7–41.0	54.5	53.1–55.9
	DM (+)	29.8	27.8–31.7	27.8	25.8–29.8	25.0	22.9–27.1	24.1	21.8–26.3	32.4	29.8–35.0	41.4	38.2–44.7	57.1	53.2–61.0
<b>Worse ear based on the mid-frequency</b>		F value	p value	F value	p value	F value	p value	F value	p value	F value	p value	F value	p value	F value	p value
	DM	6.93	0.0085	9.74	0.0018	3.56	NS	4.26	0.039	9.14	0.0025	11.88	0.0006	16.49	<0.0001
	Age	460.6	<0.0001	408.8	<0.0001	411.9	<0.0001	523.6	<0.0001	753	<0.0001	926.32	<0.0001	1364	<0.0001
	DM × age	0.21	NS	0.18	NS	1.29	NS	1.01	NS	2.47	NS	6.61	0.0102	10.47	0.0012
		Mean	95% CI	Mean	95% CI	Mean	95% CI	Mean	95% CI	Mean	95% CI	Mean	95% CI	Mean	95% CI
40–64 years	DM (–)	22.8	22.1–23.6	21.1	20.3–22.0	18.4	17.5–19.3	15.6	14.7–16.6	20.4	19.5–21.4	24.2	23.1–25.3	23.2	26.3–28.8
	DM (+)	26.0	23.1–28.9	25.2	21.9–28.4	22.2	18.7–25.8	19.6	16.0–23.3	26.6	22.8–30.3	33.6	29.3–37.9	34.7	35.7–45.5
65–86 years	DM (–)	34.0	33.1–34.9	32.7	31.7–33.8	31.3	30.2–32.4	30.5	29.4–31.7	38.8	37.7–40.0	48.1	46.7–49.4	54.5	59.1–62.2
	DM (+)	36.3	33.7–38.8	35.8	33.0–38.7	32.3	29.2–35.5	32.0	28.7–35.2	40.9	37.6–44.2	49.7	45.9–53.5	57.1	58.1–66.9

Objective variables: Hearing level at respective frequencies; Explanatory variables: DM, age, DM × age; Moderator variables: sex, history of ear disease (presence=1), history of occupational noise-exposure (presence=1).

The degrees of freedom were 1 for all F values. DM – diabetes mellitus; presence vs. absence; age: <65 years vs. ≥65 years; CI – confidence interval; NS – not significant.

hearing in the higher frequencies might be more emphatic in the younger age bracket because this type of hearing is generally better preserved in younger than in older people.

Regarding the association of diabetes with hearing, Bainbridge et al. reported the risk for hearing loss in people with self-reported DM in a recent study examining a large sample of 5140 non-institutionalized adults in the US National Health and Nutrition Examination Surveys [15]. They found that people with DM were at increased risk for hearing loss. The literature also contains some discussion about the cross-contribution of DM and aging to hearing impairment [9,15–17]. Although Bainbridge et al. mentioned that the relative contribution of DM to hearing impairment might have been stronger among their study group (age 20 to 69 years) than in a previously reported older group, they did not analyze the DM-by-age interaction as it relates to hearing in a straightforward manner. They demonstrated that the prevalence of hearing impairment among people with diagnosed DM statistically exceeded the prevalence among those without DM in all groups except people aged 60 to 69 years. They therefore speculated

that hearing of younger people, before the cumulative effects of aging, noise exposure, and other factors have made substantial contributions to hearing impairment, is potentially affected by DM more than in older people. Vaughan et al. tested audiometric measures including ultra-high-frequency range in 342 veterans with DM and 352 without DM. They concluded that patients aged 60 or younger with DM may show early high-frequency hearing loss similar to early presbycusis, while there was less difference in hearing loss between patients with and without DM after age 60 [17]. They found a significant effect of DM in both ears after adjusting for age, but only in the ultra-high-frequency range (10–16 kHz), and the DM-age interaction was significant in the right ear but not in the left at this frequency range. They also demonstrated that differences between patients with and without DM diminished or disappeared at 10, 12.5, 14 and 16 kHz in the right ear as age increased. Vasilyeva et al. prospectively assessed hearing abilities in middle-aged mice with Type 1 DM or Type 2 DM, and found that induction of diabetes in middle-aged CBA/CaJ mice promoted amplification of age-related peripheral hearing loss [18].



2. Jerger J, Chmiel R, Wilson N, Luchi R: Hearing impairment in older adults: new concepts. *J Am Geriatr Soc*, 1995; 43: 928-35
3. Wiley TL, Cruickshanks KJ, Nondahl DM, Tweed TS: Self-reported hearing handicap and audiometric measures in older adults. *J Am Acad Audiol*, 2000; 11: 67-75
4. Gates GA, Cooper JC Jr, Kannel WB, Miller NJ: Hearing in the elderly: the Framingham cohort, 1983-1985, part I—basic audiometric test results. *Ear Hear*, 1990; 11: 247-56
5. Pleis JR, Lethbridge-Cejku M: Summary health statistics for U.S. adults: National Health Interview Survey, 2006. *Vital Health Stat*, 2007; 10: 1-153
6. Cruickshanks KJ, Wiley TL, Tweed TS et al: Prevalence of hearing loss in older adults in Beaver Dam, Wisconsin. *The Epidemiology of Hearing Loss Study*. *Am J Epidemiol*, 1998; 148: 879-86
7. Jordao AMD: Consideration sur un cas du diabete. *Union Med. Du Paris*, 1857; 11: 446
8. Cullen JR, Cinnamon MJ: Hearing loss in diabetics. *J Laryngol Otol*, 1993; 107: 179-82
9. Dalton DS, Cruickshanks KJ, Klein R et al: Association of NIDDM and hearing loss. *Diabetes Care*, 1998; 21: 1540-44
10. Frisina ST, Mapes F, Kim S et al: Characterization of hearing loss in aged type II diabetics. *Hear Res*, 2006; 211: 103-13
11. Kakarlapudi V, Sawyer R, Staecker H: The effect of diabetes on sensorineural hearing loss. *Otol Neurotol*, 2003; 24: 382-86
12. Ma F, Gomez-Mari'n O, Lee DJ, Balkany T: Diabetes and hearing impairment in Mexican American adults: a population-based study. *J Laryngol Otol*, 1998; 112: 835-39
13. Ren J, Zhao P, Chen L et al: Hearing loss in middle-aged subjects with type 2 diabetes mellitus. *Arch Med Res*, 2009; 40: 18-23
14. Uchida Y, Nakashima T, Ando F et al: Is there a relevant effect of noise and smoking on hearing? A population-based aging study. *Int J Audiol*, 2005; 44: 86-91
15. Bainbridge KE, Hoffman HJ, Cowie CC: Diabetes and hearing impairment in the United States: audiometric evidence from the National Health and Nutrition Examination Surveys, 1999 to 2004. *Ann Intern Med*, 2008; 149: 1-10
16. Helzner EP, Cauley JA, Pratt SR et al: Race and sex differences in age-related hearing loss: the Health, Aging and Body Composition Study. *J Am Geriatr Soc*, 2005; 53: 2119-27
17. Vaughan N, James K, McDermott D et al: A 5-year prospective study of diabetes and hearing loss in a veteran population. *Otol Neurotol*, 2006; 27: 37-43
18. Vasilyeva ON, Frisina ST, Zhu X et al: Interactions of hearing loss and Diabetes Mellitus in the middle age CBA/CaJ mouse model of presbycusis. *Hear Res*, 2009; 249: 44-53
19. Rust KR, Prazma J, Triana RJ et al: Inner ear damage secondary to diabetes mellitus, II: changes in aging SHR/N-cp rats. *Arch Otolaryngol Head Neck Surg*, 1992; 118: 397-400
20. Makishima K, Tanaka K: Pathological changes of the inner ear and central auditory pathway in diabetics. *Ann Otol Rhinol Laryngol*, 1971; 80: 218-28
21. Fukushima H, Cureoglu S, Schachern PA et al: Effects of type 2 diabetes mellitus on cochlear structure in humans. *Arch Otolaryngol Head Neck Surg*, 2006; 132: 934-38
22. Chan L, Terashima T, Fujimiya M, Kojima H: Chronic diabetic complications: the body's adaptive response to hyperglycemia gone awry? *Trans Am Clin Climatol Assoc*, 2006; 117: 341-52
23. Jiang H, Talaska AE, Schacht J, Sha SH: Oxidative imbalance in the aging inner ear. *Neurobiol Aging*, 2007; 28: 1605-12
24. Sha SH, Taylor R, Forge A, Schacht J: Differential vulnerability of basal and apical hair cells is based on intrinsic susceptibility to free radicals. *Hear Res*, 2001; 155: 1-8
25. Yoshikawa T, Miyazaki A, Fujimoto S: Decrease in serum levels of advanced glycation end-products by short-term lifestyle modification in non-diabetic middle-aged females. *Med Sci Monit*, 2009; 15: PH65-73

## 一般住民における動脈硬化と骨粗鬆症の関連

竹村真里枝 松井康素 原田 敦  
安藤富士子 下方浩史

 **ライフサイエンス出版**  
TEL (03) 3664-7900 (代表)

【禁 無断転載・複製】

## 一般住民における動脈硬化と骨粗鬆症の関連

竹村真里枝<sup>1)</sup> 松井康素<sup>1)</sup> 原田 敦<sup>1)</sup>  
安藤富士子<sup>2)</sup> 下方浩史<sup>3)</sup>

## はじめに

骨粗鬆症と動脈硬化の進行は、高齢者の自立を障害して quality of life (QOL) を低下させ、健康寿命に大きく影響する。これら両疾病はともに、加齢に伴い発症・増加する病態である。

骨密度 (bone mineral density: BMD) と循環器疾患の関連については、これまでも多方面からの研究報告が数多くあるが、日本人地域住民男女を対象としての疫学研究はまだ少ない。本研究では、地域在住中高年者を対象に、骨粗鬆症と動脈硬化の関連について横断的に検討を行った。

## 1 対象と方法

国立長寿医療センター研究所疫学研究所では、老化に関する包括的な疫学調査である「国立長寿医療センター研究所・老化に関する長期縦断疫学研究 (NILS-LSA: National Institute for Longevity Sciences-Longitudinal Study of Aging)」を、1997年11月から縦断的(2年ごと)に実施している<sup>1)</sup>。調査対象は、センター周辺(愛知県大府市, 知多郡東浦町)の地域住民から、年齢、性別で層化した無作為抽出法で選出された、ベースライン調査時年齢が40~79歳の2,267名である。

本研究では、第1次調査(1997年11月~2000

年4月)の参加者のうち、BMD測定および頸動脈超音波検査を受けた女性1,050名(平均年齢±SD: 59.0±10.9)、男性1,063名(59.2±10.9)を対象として横断研究を行った。表1に対象者特性を示す。

骨粗鬆症の評価は、dual energy X-ray absorptiometry (DXA: Hologic, QDR-4500) で第2~4腰椎と右大腿骨頸部のBMD測定を行い、日本骨代謝学会の「原発性骨粗鬆症の診断基準」<sup>2)</sup>に準じ、BMDが若年成人平均値(young adult mean: YAM)の70~80%を骨量減少、

表1 対象者特性

	男性	女性
対象者数(名)	1,063	1,050
40歳代	274	266
50歳代	267	266
60歳代	255	263
70歳代	267	255
年齢(歳)	59.2±10.9	59.0±10.9
BMI(kg/m <sup>2</sup> )	22.9±2.8	22.9±3.3
腰椎BMD(g/cm <sup>2</sup> )	1.0±0.2	0.9±0.2
大腿骨頸部BMD(g/cm <sup>2</sup> )	0.8±0.1	0.7±0.1
IMT(mm)	0.9±0.5	0.8±0.3

平均値±標準偏差

BMI: body mass index, BMD: bone mineral density, IMT: intima-media thickness

Association between Arteriosclerosis and Osteoporosis in Community Dwelling Population

Marie Takemura: Department of Orthopedics National Center for Geriatrics and Gerontology, et al.

Key words: Osteoporosis, Arteriosclerosis, Epidemiology

<sup>1)</sup> 国立長寿医療センター整形外科 <sup>2)</sup> 愛知淑徳大学医療福祉学部医療貢献学科 <sup>3)</sup> 国立長寿医療センター研究所疫学研究所





図1 骨粗鬆症/骨量減少の有病率



70%未満を骨粗鬆症と判定した。

また、動脈硬化の評価手段として頸動脈超音波検査(日立メディコ電子走査形超音波断層装置EUB-655, 電子リニア形探触子EUP-L3 10MHz)を行った。頸動脈内膜中膜複合体厚(intima-media thickness: IMT)を左右総頸動脈および左右頸動脈分岐部で計測し、その最肥厚部をIMTの代表値とした。IMTが1.1mm以上を異常肥厚とし、動脈硬化ありと判定した。

統計学的検討として、骨粗鬆症および動脈硬化の地域在住中高年者の有病率を性別、年代別に求め、Cochran-Mantel-Haenszel法によるトレンド検定を行った。次に動脈硬化と骨粗鬆症の関連について検討するために、動脈硬化の有無を説明変数とし、年齢およびbody mass index(BMI)を調整した骨粗鬆症有病率についての多重ロジスティック回帰分析を性別に行った。解析には、統計プログラムSAS release 9.1.3を使用した。

## 2 結 果

### 1) 骨粗鬆症/骨量減少の有病率(性別, 年代別)

腰椎BMD判定での40歳以上の骨粗鬆症/骨量減少の有病率は、女性37.9%, 男性16.1%であった。女性の有病率は加齢で有意に高くなり、特に60歳代以降は急速に高くなった。男性でも、有病率は加齢で有意に高くなった。大腿骨頸部BMD判定でも、女性30.4%, 男性15.1%で、男




図2 動脈硬化有病率

女とも加齢で有意に高くなった(図1)。

### 2) 動脈硬化有病率(性別, 年代別)

40歳以上の女性の動脈硬化有病率は16.6%, 男性は21.4%であった。動脈硬化の有病率は、男女とも加齢で有意に高くなった(図2)。

### 3) 動脈硬化と骨粗鬆症との関連

女性で骨粗鬆症診断を腰椎BMDで判定した場合、動脈硬化のある者は、ない者に比べて骨粗鬆症/骨量減少の有病の割合が高かった(オッズ比1.97, 95%信頼区間1.03~2.99,  $p = 0.0014$ )。また、女性を未閉経群と閉経群に分けて検討したところ、閉経群のみで同様の結果が得られた(1.78, 1.19~2.67,  $p = 0.00052$ )。一方、大腿骨頸部BMD判定の場合には、いずれも有意な関連は認められなかった。男性での解析では、両部位BMDと動脈硬化のあいだに有意な



表2 多重ロジスティック回帰分析による動脈硬化と骨粗鬆症有病の関連

		オッズ比(95%信頼区間)	p値
腰椎BMD判定の場合	女性全体	1.97(1.03~2.99)	p=0.0014
	閉経女性	1.78(1.19~2.67)	p=0.0052
	男性全体	0.98(0.63~1.55)	NS
大腿骨頸部BMDの場合	女性全体	0.96(0.63~1.46)	NS
	閉経女性	0.94(0.60~1.43)	NS
	男性全体	0.74(0.27~1.17)	NS

NS: not significant

関連は認められなかった(表2)。

### 3 考 察

BMDと心血管系疾患については、低BMDや骨密度減少が、心血管疾患による死亡リスク上昇と関連するという報告や<sup>3,4)</sup>、骨粗鬆症の閉経後女性は、年齢や心血管疾患の危険因子を考慮しても、心・血管系イベントの発生リスクが有意に高い<sup>5)</sup>など、これまでに多方面からの研究が行われている。日本人を対象とした疫学研究においても、骨粗鬆症や動脈硬化の評価手法はそれぞれの研究で異なるが、BMDと動脈硬化の程度とのあいだに関連を認めたと報告されている<sup>6~9)</sup>。本研究では、脳・心血管疾患の予知因子として有用とされるIMT<sup>10)</sup>とBMDのデータを用いて、地域住民男女における両疾病間の関連について解析を行った。その結果、骨粗鬆症評価を腰椎BMDで行った場合、女性で動脈硬化のある者は、ない者比べて骨粗鬆症有病の割合が高くなった。この結果は、これまでの先行研究と矛盾するものではなかった。

女性の骨粗鬆症と心血管疾患は、どちらも閉経後より罹患率が高くなることが臨床的に広く知られている。またエストロゲン受容体は、骨芽細胞や破骨細胞、血管内皮細胞、血管平滑筋細胞に存在することが確認されており、両疾患の進行に共通して関与する因子としてエストロゲンがあげられる。本研究において、女性で両疾病間に有意な関連が認められた要因の一つに、エストロゲンの関与が示唆される。

またエストロゲン以外にも、酸化脂質やビタ

ミンD、副甲状腺ホルモン、オステオカルシン、オステオポンチン、ホモシステイン、アンジオテンシン、マトリックスグラブプロテイン、オステオプロテジェリン、一酸化窒素、インターロイキン(IL-6)などは、骨と血管の相互に関与する共通因子として近年検証が進み、いわゆる「骨・血管相関」の機序が解明されつつある。

腰椎は大腿骨頸部に比べ、構成組織として海綿骨の占める割合が高い。エストロゲン減少による骨代謝への影響は、皮質骨よりも海綿骨のほうがより反映されやすいと考えられている。今回の結果において、腰椎と大腿骨に相違を認めた要因の一つとしてエストロゲンの関与が示唆される。

一方、変形性脊椎症では椎体に骨棘形成を認めるが、その割合は、特に男性において加齢に伴い高くなるため、BMDが高く判定されてしまう。また腹部の大動脈石灰化が存在する場合にも、腰椎BMDが過大評価される可能性があり、結果への影響は否定できない。

両疾患の進行には性ホルモン以外にも喫煙、糖尿病、高脂血症など共通の危険因子が存在する。今後、これらの交絡因子を考慮しての検討を要すると思われる。

患者のQOL低下を招くおそれのある骨粗鬆症性骨折予防の観点から、動脈硬化の基盤となる高血圧や高脂血症など生活習慣病の日常診療において、骨粗鬆症についても評価や治療の必要性がある。また逆に閉経後女性の骨粗鬆症有病者の日常診療において、脳・心血管疾患発症の基盤となる動脈硬化の存在にも留意すること

が必要と考えられた。

### ま と め

地域在住中高年者を対象に、骨粗鬆症と動脈硬化症の有病の関連について横断的解析を行った。女性では年齢とBMIで調整しても、腰椎BMDとIMTとのあいだに有意な関連を認め、骨粗鬆症と動脈硬化進展とのあいだに密接な関連が示唆された。

### 文 献

- 1) Shimokata H, Ando F, Niino N. A new comprehensive study on aging- the- National Institute for Longevity Sciences, Longitudinal Study of Aging (NILS-LSA). *J Epidemiol* 2000;10:S1-9.
- 2) 折茂肇, 林泰史, 福永仁夫, 曾根照喜, 藤原佐枝子, 白木正孝ほか, 原発性骨粗鬆症の診断基準(2000年度改訂版). *日骨代謝誌* 2001;18:76-82.
- 3) von der Recke P, Hansen MA, Hassager C. The association between low bone mass at the menopause and cardiovascular mortality. *Am J Med* 1999;106:273-8.
- 4) Kado DM, Browner WS, Blackwell T, et al. Rate of bone loss is associated with mortality in older women ; a prospective study. *J Bone Miner Res* 2000;15:1974-80.
- 5) Tankó LB, Christiansen C, Cox DA, Geiger MJ, McNabb MA, Cummings SR. Relationship between osteoporosis and cardiovascular disease in postmenopausal women. *J Bone Miner Res* 2005;20:1912-20.
- 6) 鈴木隆雄, 吉田英世, 金憲経. 高齢者における骨密度と脈は速度の関連性についての前向き追跡研究-骨粗鬆症は動脈硬化の促進に関与するか-. *Osteoporosis Jpn* 2005;13:75-7.
- 7) Uyama O, Yoshimoto Y, Yamamoto Y, Kawai A. Bone changes and carotid atherosclerosis in postmenopausal women. *Stroke* 1997;28:1730-2.
- 8) Sumino H, Ichikawa S, Kasama S, Takahashi T, Sakamoto H, Kumakura H, et al. Relationship between carotid atherosclerosis and lumbar spine bone mineral density in postmenopausal women. *Hypertens Res* 2008;31:1191-7.
- 9) Tamaki J, Iki M, Hirano Y, Sato Y, Kajita E, Kagamimori S, Kagawa Y, Yoneshima H. Low bone mass is associated with carotid atherosclerosis in postmenopausal women: the Japanese Population-based Osteoporosis (JPOS) Cohort Study. *Osteoporos Int* 2009;20:53-60.
- 10) Lorenz MW, Markus HS, Bots ML, Rosvall M, Sitzer M. Prediction of clinical cardiovascular events with carotid intima-media thickness: a systematic review and meta-analysis. *Circulation* 2007;115:459-67.

# ロコモティブ シンドローム ——運動器科学の新時代

企画 中村耕三 東京大学大学院医学系研究科感覚運動機能医学(整形外科学)

- 運動器疾患の疫学
- ロコモティブシンドローム
- 運動器の保存治療
- 介護保険制度と在宅医療
- 運動器障害の診断と評価
- 歩行と姿勢の評価
- 主な疾患
- 運動器疾患の基礎



医歯薬出版株式会社  
<http://www.ishiyaku.co.jp/>

# ロコモティブシンドローム

## —運動器科学の新時代

企画 中村耕三(東京大学大学院医学系研究科感覚運動機能医学<整形外科学>)



Vol.236 No.5 2011/1/29

311 “ロコモティブシンドローム”特集にあたって……………中村耕三

### 運動器疾患の疫学

315 大規模住民調査からみえてきた運動器疾患の実態——ROAD study ……吉村典子

319 運動器疾患の長期縦断疫学研究……………下方浩史・安藤富士子

325 転倒の疫学と予防対策——ロコモティブシンドロームの視点から……………鈴木隆雄

333 DPCデータベースからみた日本整形外科の現状  
——超高齢社会を迎えて……………門野夕峰

339 地域在住高齢者の歩行能力に関する横断的・縦断的分析……………坂田悞教

### ロコモティブシンドローム

347 ロコモティブシンドロームの概念……………中村耕三

353 ロコチェックの運動機能低下の予見性と、ロコトレの運動機能改善効果  
……………石橋英明

360 病院における転倒予防教室——その運営・効果・課題……………岡田知佐子

366 臨床現場におけるロコモティブシンドローム……………大江隆史

371 運動器障害診断ツール(足腰指数25)の開発……………星野雄一・星地亜都司

### 運動器の保存治療

379 中高年者の背筋力維持・向上のための運動……………宮腰尚久

383 変形性膝関節症に対する運動療法・体操——ロコモ予防への取組み……………帖佐悦男・他