

CA), and Goat anti-mouse IgG conjugated with Alex Fluor (AF) 488 or 594 and goat anti-rat IgG conjugated with AF 488 (Molecular Probes, Eugene, OR).

Immunopositive signals were visualized using an observation system (Compix Imaging System, Lake Oswego, OR) linked to an Olympus microscope BX50 through a highly sensitive CCD camera or a confocal laser scanning microscope (Carl Zeiss LSM510).

Fluoro-Jade B was purchased from Chemicon (now part of Millipore, Schwalbach, Germany). 7-AAD was purchased from Invitrogen (Carlsbad, CA).

### ***Histochemistry***

For Fluoro-Jade B histochemistry (1:50000), 5- $\mu$ m-thick paraffin-embedded sections were deparaffinized and stained following the manufacturer's instruction (Chemicon, now part of Millipore, Schwalbach, Germany). The Fluoro-Jade B-stained product fluoresces when excited at 488 nm and staining was imaged using a confocal laser scanning microscope (Carl Zeiss LSM510). The same procedure was applied for 30- $\mu$ m-thick cryostat sections.

### ***Cell culture and cellular uptake***

Human neuroblastoma SH-SY5Y (SY5Y) cells were cultured in Dulbecco's modified

Eagle's medium/Ham's F-12 medium supplemented with 10% fetal bovine serum. To investigate the fate of extracellular A $\beta$ , SY5Y cells were exposed to HiLyte Fluor<sup>TM</sup> 488-labeled A $\beta$ M<sub>s</sub> (10 kDa-filtrate), HiLyte Fluor<sup>TM</sup> 488-labeled A $\beta$ O<sub>s</sub> (30 kDa-retentate) at 5  $\mu$ M (AnaSpec, San Jose, CA), or Fluor<sup>TM</sup> 488 alone for 10 min, 30 min, and 180 min. In a separate set of experiment, cultures were treated at 37°C for 0, 3, 6, and 24 h with 5  $\mu$ M Fluor<sup>TM</sup> 488-labeled A $\beta$ O<sub>s</sub>, and for 24 h with 5  $\mu$ M Fluor<sup>TM</sup> 488-labeled A $\beta$ M<sub>s</sub> and synthetic A $\beta$ 42-1 (AnaSpec, San Jose, CA). Toxicity was assessed by CytoTox 96 Non-Radioactive Cytotoxicity Assay Kit in accordance with the manufacturer's instructions (Promega, Madison, WI) as described previously [22].

#### ***Protein extraction and immunoblotting***

Saline-soluble, saline-insoluble, SDS-soluble fractions, or SDS-insoluble, formic acid (FA)-extractable fractions were prepared from the Tg2576 mouse brains as described previously [32]. Briefly, frozen brain samples were homogenized with a motor-driven Teflon/glass homogenizer (20 strokes) in TBS containing a cocktail of protease inhibitors (150 mg/ml), followed by centrifugation at 100,000 g for 1 h. The resultant supernatant (soluble fraction) was subjected to dot blot immunoanalysis or western blotting employing the same antibodies as those used for immunohistochemical staining.

The pellet was further extracted with 2% sodium dodecylsulphate (SDS), followed by 70% FA, and the homogenate was ultracentrifuged as described above. The resultant supernatant (insoluble fraction) was also subjected to dot blot immunoanalysis and western blot analysis. For western blotting, aliquots of isolated fractions were separated using NuPAGE 4-12% bis-tris-glycine gels and transblotted to nitrocellulose membrane or Immobilon P (Millipore) for 1 h at 400 mA using 10 mM 3-cyclohexylamino-1-propanesulphonic acid (pH 11) containing 10% methanol. Membranes were blocked for 3 h at room temperature with 5% low-fat milk and 1% BSA in PBST and incubated with either the polyclonal anti-A11 (1:1000) or anti-PSD95 antibody (1:250), and monoclonal anti-drebrin (1:100), anti-SYP (1:2000), and anti- $\beta$  actin antibodies (1:1000), followed by HRP-labeled goat anti-rabbit or anti-mouse F(ab')<sub>2</sub> antibody (1:3000; Amersham). Immunoblots were visualized with an ECL kit using LAS3000 mini (Fujitsu, Tokyo, Japan). Densitometric analysis of immunoblot was performed using Multi Gauge v3.0 software (Fuji Film, Tokyo), and bands of interest were normalized to the corresponding actin bands indicated.

### ***Immunization and behavioral analyses***

All animal procedures were performed in accordance with a protocol approved by the

Animal Care Committee of the National Institute for Longevity Sciences. Several 3-month-old female nontransgenic (non-Tg) and Tg2576 mice that carry and overexpress the human APP gene with the Swedish double mutation (K670N; M671L) of familial AD were purchased from Taconics (Germantown, NY, USA) and maintained at our animal care facility until 13 months of age. To determine whether immunization prevents the development of Alzheimer-like phenotype, 4-month-old Tg2576 mice were administered 1A9 or 2C3 (0.4 mg/kg/week), or PBS intravenously via the tail vein until 13 months of age. Memory functions were measured at 13 months of age in the following four behavioral paradigms, as described previously [28]: (1) spontaneous alternation in the Y-maze test; (2) novel object recognition test; (3) Morris water maze test; and (4) cued and fear conditioning tests. Mice were sacrificed 3 days after the termination of the behavioral tests for biochemical and histological assessments. Experimental results were analyzed by one-way ANOVA and two-way ANOVA, with Fisher's test for post hoc analysis.

#### *Spontaneous alternation in Y-maze test*

The maze was made of black painted wood; each arm was 40 cm long, 12 cm high, 3 cm wide at the bottom, and 10 cm wide at the top. The arms converged at an

equilateral triangular central area that was 4 cm long at its longest axis. Each mouse was placed at the center of the apparatus, the sequence and number of arm entries were recorded for each mouse over an 8 min period. Alternation was defined as successive entry into the three arms on overlapping triplet sets. Alternation behavior (%) was calculated as the ratio of actual alternations to possible alternations (defined as the number of arm entries minus two) multiplied by 100.

#### *Novel object recognition test*

The test procedure consisted of three sessions: habituation, training, and retention. Each mouse was habituated to the box (30 x 30 x 35 cm), with 10 min of exploration in the absence of objects for 3 days (habituation session). During the training session, two objects were placed at the back corner of the box. A mouse was then placed midway at the front of the box and the total time spent exploring the two objects was recorded for 10 min. During the retention session, the mice were placed back in the same box 24 hr after the training session, in which one of the familiar objects used during the training was replaced with a novel object. The animals were then allowed to explore freely for 10 min, and the time spent exploring each object was recorded. Throughout the experiments, the objects used were counterbalanced in terms of their

physical complexity and emotional neutrality. A preference index, a ratio of the amount of time spent exploring any one of the two objects (training session) or the novel object (retention session) over the total time spent exploring both objects, was used to measure cognitive function. To eliminate the influence of the last behavioral test, the objects were changed each time.

#### ***Morris water maze test***

The Morris water maze test was conducted in a circular pool (120 cm in diameter) filled with water at a temperature of  $22 \pm 1^\circ\text{C}$ . A hidden platform (one block) (7 cm in diameter) was used. The mice underwent two trials (one block) for 10 consecutive days, during which the platform was left in the same position. The distance taken to locate the escape platform was determined in each trial using the Etho Vision system (Neuroscience Idea Co., Ltd., Osaka, Japan).

#### ***Cued and contextual fear conditioning tests***

For measuring basal levels of freezing response (preconditioning phase), mice were individually placed in the conditioning cage (17 x 27 x 12.5 cm) for 1 min, then in the conditioning cage (25 x 31 x 11 cm) for 2 min. For training (conditioning phase), mice

were placed in the conditioning cage, then a 15 sec tone (80 dB) was delivered as a conditioned stimulus. During the last 5 sec, an unconditioned stimulus was applied through a shock generator (Neuroscience Idea Co., Ltd.). This procedure was repeated 4 times at 15 sec intervals. Cued and contextual tests were carried out 1 day after the fear conditioning. For the cued test, the freezing response was measured in the neutral cage for 1 min in the presence of a continuous-tone stimulus identical to the conditioned stimulus. For the contextual test, mice were placed in the conditioning cage and the freezing response was measured for 2 min in the absence of the conditioned stimulus.

### ***Ethics Statement***

The research protocol was approved by the local animal esthetics committees at Research Institute, National Center for Geriatrics and Gerontology (Animal Care Committee) prior to initiation of the study. The research project was approved by the local ethics committee of Hirosaki University Graduate School of Medicine, and Research Institute, National Center for Geriatrics and Gerontology prior to initiation of the study.

### ***Statistical analyses***

We used factorial design analysis of variance (ANOVA) or Mann-Whitney test to analyze data as appropriate. Significant ANOVA values were subsequently subjected to simple main effects analyses or post hoc comparisons of individual means using the Tukey's or Dannett's method as appropriate. We considered *p* values of 0.05 as significant for all studies. Some of the data obtained from animal experiments were analyzed by two-way ANOVA, with Fisher's test for post hoc analysis.

### **Competing Interests**

YO, TY, and MS are the employees in Immunus Pharma Incorporation. TY and MS hold stock options in Immunus Pharma Incorporation. EM, TY, and MS are co-inventors of two filed provisional patent applications titled "Antibody Specific Binding to a Beta Oligomer and The Use" and "Antibodies That Specifically Bind to A $\beta$  Oligomers and Uses Thereof" that cover the antibodies described in this paper, but this does not alter the adherence to all the Molecular Neurodegeneration policies on sharing data and materials. This study has in some parts been funded by a commercial funder, but that does not alter the author's adherence to all the Molecular Neurodegeneration policies on sharing data and materials.



### **Author's contributions**

Conceived and designed the experiments: MM EM. Performed the experiments: AT YO TK AM TN HS TU NY EM. Analyzed the data: MS MS KY MM. Contributed reagents/ materials/analysis tools: YO TY MS KA. Wrote the paper: AT EM. All authors read and approved the final manuscript.

### **Acknowledgments**

This work was supported in part by a Grant-in-Aid for Advanced Brain Scientific Project-from the Ministry of Education, Culture, Sports, Science and Technology, Japan [15016080 and 16015284 to E.M.]; and the Research Grant for Longevity Sciences from the Ministry of Health, Labour and Welfare [17A-1 to E.M.].

## References

1. Hardy J, Allsop D: **Amyloid deposition as the central event in the aetiology of Alzheimer's disease.** *Trends Pharmacol Sci* 1991, **12**: 383-388.
2. Lue LF, Kuo YM, Roher AE, Brachova L, Shen Y, Sue L, Beach T, Kurth JH, Rydel RE, Rogers J: **Soluble amyloid  $\beta$  peptide concentration as a predictor of synaptic change in Alzheimer's disease.** *Am J Pathol* 1999, **155**: 853-862.
3. McLean CA, Cherny RA, Fraser FW, Fuller SJ, Smith MJ, Beyreuther K, Bush AI, Masters CL: **Soluble pool of Abeta amyloid as a determinant of severity of neurodegeneration in Alzheimer's disease.** *Ann Neurol* 1999, **46**: 860-866.
4. Klein WL, Krafft GA, Finch CE: **Targeting small Abeta oligomers: the solution to an Alzheimer's disease conundrum?** *Trends Neurosci* 2001, **24**: 219-224.
5. Selkoe DJ: **Alzheimer's disease is a synaptic failure.** *Science* 2002, **298**: 789-791.
6. Hass C, Selkoe DJ: **Soluble protein oligomers in neurodegeneration: lessons from the Alzheimer's amyloid  $\beta$ -peptide.** *Nat Rev Mol Cell Biol* 2007, **8**: 101-112.
7. Akiyama H, Barger S, Barnum S, Bradt B, Bauer J, Cole GM, Cooper NR, Eikelenboom P, Emmerling M, Fiebich BL, Finch CE, Frautschy S, Griffin WS,

- Hampel H, Hull M, Landreth G, Lue L, Mrazek R, Mackenzie IR, McGeer PL, O'Banion MK, Pachter J, Pasinetti G, Plata-Salamán C, Rogers J, Rydel R, Shen Y, Streit W, Strohmeier R, Tooyoma I, Van Muiswinkel FL, Veerhuis R, Walker D, Webster S, Wegrzyniak B, Wenk G, Wyss-Coray T: **Inflammation and Alzheimer's disease.** *Neurobiol Aging* 2000, **21**: 383-421.
8. Hardy JA, Higgins GA: **Alzheimer's disease: the amyloid cascade hypothesis.** *Science* 1992, **256**: 184-185.
9. Lambert MP, Barlow AK, Chromy BA, Edwards C, Freed R, Liosatos M, Morgan TE, Rozovsky I, Trommer B, Viola KL, Wals P, Zhang C, Finch CE: **Diffusible, nonfibrillar ligands derived from A $\beta$ <sub>1-42</sub> are potent central nervous system neurotoxins.** *Proc Natl Acad Sci USA.* 1998, **95**: 6448-6453.
10. Dodart JC, Bales KR, Gannon KS, Greene SJ, DeMattos RB, Mathis C, DeLong CA, Wu S, Wu X, Holtzman DM, Paul SM: **Immunization reverses memory deficits without reducing brain A $\beta$  burden in Alzheimer's disease model.** *Nat Neurosci* 2002, **5**:452-457.
11. Chromy BA, Nowak RJ, Lambert MP, Viola KL, Chang L, Velasco PT, Jones BW, Fernandez SJ, Lacor PN, Horowitz P, Finch CE, Krafft GA, Klein WL: **Self-assembly of A $\beta$ <sub>1-42</sub> into globular neurotoxins.** *Biochemistry* 2003, **42**:

12749-12760.

12. Kaye R, Head E, Thompson JL, McIntire TM, Milton SC, Cotman CW, Glabe CG: **Common structure of soluble amyloid oligomers implies common mechanism of pathogenesis.** *Science* 2003, **300**: 486-489.
13. Bayer A, Jones RW, Bullock R, Love S, Neal JW, Zotova E, Nicoll JA: **Long-term effects of Abeta42 immunisation in Alzheimer's disease: follow-up of a randomised, placebo-controlled phase I trial.** *Lancet* 2008, **372**:216-223.
14. Walsh DM, Klyubin I, Fadeeva JV, Cullen WK, Anwyl R, Wolfe MS, Rowan MJ, Selkoe DJ: **Naturally secreted oligomers of amyloid  $\beta$  protein potently inhibit hippocampal long-term potentiation in vivo.** *Nature* 2002, **16**: 535-539.
15. Shankar GM, Li S, Mehta TH, Garcia-Munoz A, Shepardson NE, Smith I, Brett FM, Farrell MA, Rowan MJ, Lemere CA, Regan CM, Walsh DM, Sabatini BL, Selkoe DJ : **Amyloid-beta protein dimers isolated directly from Alzheimer's brains impair synaptic plasticity and memory.** *Nat Med* 2008, **14**:837-842.
16. Lacor PN, Buniel MC, Chang L, Fernandez SJ, Gong Y, Viola KL, Lambert MP, Velasco PT, Bigio EH, Finch CE, Krafft GA, Klein WL: **Synaptic targeting by Alzheimer's-related amyloid  $\beta$  oligomers.** *J Neurosci* 2004, **24**:10191-10200.
17. Cleary JP, Walsh DM, Hofmeister JJ, Shankar GM, Kuskowski MA, Selkoe DJ,

- Ashe KH: **Natural oligomers of the amyloid- $\beta$  protein specifically disrupt cognitive function.** *Nat Neurosci* 2005, **8**:79-84.
18. Lesné S, Koh MT, Kotilinek L, Kaye R, Glabe CG, Yang A, Gallagher M, Ashe KH: **A specific amyloid-beta protein assembly in the brain impairs memory.** *Nature*. 2006, **440**: 352-357.
19. Gong Y, Chang L, Viola KL, Lacor PN, Lambert MP, Finch CE, Krafft GA, Klein WL: **Alzheimer's disease-affected brain: presence of oligomeric A $\beta$  ligands (ADDLs) suggests a molecular basis for reversible memory loss.** *Proc. Natl. Acad. Sci. USA*. 2003, **100**:10417-10422.
20. O'Nuallain B, Wetzel R: **Conformational Abs recognizing a generic amyloid fibril epitope.** *Proc Natl Acad Sci USA* 2002, **99**:1485-1490.
21. Kaye R, Head E, Sarsoza F, Saing T, Cotman CW, Neucula M, Margol L, Wu J, Breydo L, Thompson JL, Rasool S, Gurlo T, Butler P, Glabe CG: **Fibril specific, conformation dependent antibodies recognize a generic epitope common to amyloid fibrils and fibrillar oligomers that is absent in prefibrillar oligomers.** *Mol Neurodegen* 2007, **2**: 18-28.
22. Yamamoto N, Matsubara E, Maeda S, Minagawa H, Takashima A, Maruyama W, Michikawa M, Yanagisawa K: **A ganglioside-induced toxic soluble A $\beta$  assembly.**

- Its enhanced formation from A $\beta$  bearing the Arctic mutation.** *J Biol Chem* 2007, **282**: 2646-2655.
23. Hoshi M, Sato M, Matsumoto S, Noguchi A, Yasutake K, Yoshida N, Sato K:  
**Spherical aggregates of  $\beta$ -amyloid (amylospheroid) show high neurotoxicity and activate tau protein kinase I/glycogen synthase kinase-3 $\beta$ .** *Proc. Natl. Acad. Sci. USA* 2003, **100**:6370-6375.
24. Noguchi A, Matsumura S, Dezawa M, Tada M, Yanazawa M, Ito A, Akioka M, Kikuchi S, Sato M, Ideno S, Noda M, Fukunari A, Muramatsu S, Itokazu Y, Sato K, Takahashi H, Teplow DB, Nabeshima Y, Kakita A, Imahori K, Hoshi M:  
**Isolation and characterization of patient-derived, toxic, high mass amyloid beta-protein (A $\beta$ ) assembly from Alzheimer disease brains.** *J Biol Chem.* 2009, **284**:32895-905.
25. Tero R, Takizawa M, Li Y.J, Yamazaki M, Urisu T: **Lipid membrane formation by vesicle fusion on silicon dioxide surfaces modified with alkyl self-assembled monolayer islands.** *Langmuir* 2004, **20**:7526-7531.
26. Kotilinek LA, Bacskai B, Westerman M, Kawarabayashi T, Younkin L, Hyman BT, Younkin S, Ashe KH: **Reversible memory loss in a mouse model of Alzheimer's disease.** *J Neurosci* 2002, **22**:6331-6335.

27. Lee EB, Leng LZ, Zhang B, Kwong L, Trojanowski JQ, Abel., Lee VM:  
**Targeting amyloid- $\beta$  peptide ( $A\beta$ ) oligomers by passive immunization with a conformation-selective monoclonal antibodies improves learning and memory in  $A\beta$  precursor protein (APP) transgenic mice. *J Biol Chem* 2006, **281**:4292-4299.**
28. Mouri A, Noda Y, Hara H, Mizoguchi H, Tabira T, Nabeshim, T: **Oral vaccination with a viral vector containing  $A\beta$  cDNA attenuates age-related  $A\beta$  accumulation and memory deficits without causing inflammation in a mouse Alzheimer model. *FASEB J* 2007, **21**:2135-2148.**
29. Schmued LC, Hopkins KJ. Fluoro-Jade B: **A high affinity fluorescent marker for the localization of neuronal degeneration. *Brain Res* 2000, **874**: 123-130.**
30. Lambert MP, Viola KL, Chromy BA, Chang L, Morgan TE, Yu J, Venton DL, Krafft GA, Finch CE, Klein WL: **Vaccination with soluble  $A\beta$  oligomers generates toxicity-neutralizing antibodies. *J Neurochem* 2001, **79**:595-605.**
31. Lambert MP, Barlow AK, Chromy BA, Edwards C, Freed R, Liosatos M, Morgan TE, Rozovsky I, Trommer B, Viola KL, Wals P, Zhang C, Finch CE, Lambert MP, Velasco PT, Chang L, Viola KL, Fernandez S, Lacor PN, Khuon D, Gong Y, Bigio EH, Shaw P, De Felice FG, Krafft GA, Klein WL: **Monoclonal antibodies that**

- target pathological assemblies of A $\beta$ .** *J Neurochem* 2007, **100**:23-35.
32. Kawarabayashi T, Shoji M, Younkin LH, Wen-Lang L, Dickson DW, Murakami T, Matsubara E, Abe K, Ashe KH, Younkin SG: **Dimeric amyloid beta protein rapidly accumulates in lipid rafts followed by apolipoprotein E and phosphorylated tau accumulation in the Tg2576 mouse model of Alzheimer's disease.** *J Neurosci.* 2004, **24**:3801-3809.
33. Mori C, Spooner ET, Wisniewsk KE, Wisniewski TM, Yamaguch H, Saido TC, Tolan DR, Selkoe DJ, Lemere CA: **Intraneuronal A $\beta$ 42 accumulation in Down syndrome brain.** *Amyloid* 2002, **9**: 88-102.
34. Hu X, Crick SL, Bu G, Frieden C, Pappu RV, Lee JM. **Amyloid seeds formed by cellular uptake, concentration, and aggregation of the amyloid-beta peptide.** *Proc Natl Acad Sci U S A.* 2009, **106**:20324-20329.
35. Tomiyama T, Matsuyama S, Iso H, Umeda T, Takuma H, Ohnishi K, Ishibashi K, Teraoka R, Sakama N, Yamashita T, Nishitsuji K, Ito K, Shimada H, Lambert MP, Klein WL, Mori H: **A mouse model of amyloid beta oligomers: their contribution to synaptic alteration, abnormal tau phosphorylation, glial activation, and neuronal loss in vivo.** *J Neurosci* 2010, **30**:4845-56.
36. Matsubara E, Frangione B, Ghiso J: **Characterization of apolipoprotein**



**J-Alzheimer's A beta interaction.** *J Biol Chem.* 1995, **270**:7563-7567.

## Figure legends

### Figure 1. Generation and characterization of oligomer-specific antibody. (A)

Electrophoresis of immunogen. SDS-PAGE was performed to isolate the A $\beta$ 1-42 tetramer (red closed arrowhead) alone without any contamination by the A $\beta$ 1-42 trimer (black closed arrowhead) and A $\beta$ 1-42 monomer (opened arrowhead). Lane 1, A $\beta$ 1-42 dissolved in 10 mM phosphate buffer; lane 2, A $\beta$ 1-42 dissolved in distilled deionized water. (B) A $\beta$ 1-42 oligomer formation was observed as a function of time. A $\beta$ 1-42 monomer (25  $\mu$ M) incubated at 37°C for the indicated time (0 – 72 h) were spotted on nitrocellulose membrane and subjected to a dot blot assay using A11 (1:100), 1A9 (1:50), 2C3 (1:50), or 4G8 (1:1000). (C) In this dot blot assay (left half of panel C), 1  $\mu$ g of soluble A $\beta$ 42 oligomers (100,000 g sup for 4-h-incubation at 37°C) and A $\beta$ 42 fibrils (100,000 g pellets for 120-h-incubation at RT) were applied on a nitrocellulose membrane and probed with A11, 1A9, 2C3, or 4G8. EM image of fibrils (right half of panel C). (D) Characterization of A $\beta$ 1-42 oligomers under nondenaturing conditions. A $\beta$ 1-42 monomer (25  $\mu$ M) incubated at 37°C for 4 h was separated on 16% BN-PAGE. (E) Separated peptides under nondenaturing conditions were also subjected to immunoblot analysis using A11, 1A9, 2C3, and 4G8. (F) The 100000 g sup of 4-h-incubated mixture of A $\beta$ 1-42 monomer (25  $\mu$ M) was subjected to two-dimensional

native/SDS-PAGE, followed by 4G8-immunoblot analysis. SDS-stable 15~40-mers are indicated ( ] (red) ). (G) Immunodetection of 4-h-incubated mixture of A $\beta$ 1-42 monomer (25  $\mu$ M) under denaturing conditions probed with A11, 1A9, 2C3, and 4G8. SDS-stable 15~40-mers are indicated ( ] red).

**Figure 2. Biophysical and structural characterization of neurotoxic A $\beta$  assembly.**

(Upper half of panel A) Representative calcein AM/PI stainings of NGF-treated PC12 (PC12N) cells treated at 37 °C for 48 h with: TBS alone; 0-h preincubated A $\beta$ 1-42 (0 h); 2-h preincubated A $\beta$ 1-42 (2 h); 540,000 g supernatant obtained from 2 h (2 h sup); 4-h preincubated A $\beta$ 1-42 (4 h); 540,000 g supernatant obtained from 2 h (4 h sup). Green staining for viable cells versus red staining for dead cells. Resultant cell viability for each treatment is shown in lower half of panel A. Experimental results were analyzed by one-way ANOVA, followed by Tukey's test for posthoc analysis: statistical significance compared with TBS alone (\* $p$ <0.0001). Scale bar = 50  $\mu$ m. (B) The seed-free A $\beta$ 1-42 (25  $\mu$ M) was subjected to a series of membrane ultrafiltration steps with molecular cutoffs at 3, 10, 30, and 100 kDa. The resultant four filtrates and one retentate were designated as Fr. 1 (< 3 kDa), Fr. 2 (3-10 kDa), Fr. 3 (10-30 kDa), and Fr. 4 (30-100 kDa), and final retentate Fr. 5 (>100 kDa). The upper half of panel B shows

the representative calcein AM/PI stainings of NGF-treated PC12 (PC12N) cells incubated at 37 °C for 48 h with seed-free unfractionated A $\beta$ 1-42 or five fractions (25  $\mu$ M each). Resultant cell viability for each treatment is shown in the lower half of panel B. Experimental results were analyzed by one-way ANOVA, followed by Dannett's test for posthoc analysis: statistical significance compared with TBS alone (\* $p$ <0.001). (C) Dot blot analysis of five fractions (Frs. 1-5). The blots were reacted with A11, 1A9, 2C3, and 4G8. (D) Amplitude AFM images (2  $\mu$ m x 2  $\mu$ m) of four fractions (Frs. 2-5). All AFM images were taken on a mica surface.

**Figure 3. Immunolabelling characteristics of 1A9, 2C3, A11, 6E10 and 4G8 in**

**human brains.** Typical A $\beta$ O immunolabelling is observed in diffuse plaques (arrow) and the perikaryon of pyramidal neurons (arrow head) as dense granules (A and C) or diffuse staining (B) in AD brains (400X): 1A9 (1:50, panel A), 2C3 (1:50, panel B), and A11 (1:250, panel C). In control brains (D), 1A9 disclosed small granular intraneuronal staining in isolated clusters of pyramidal neurons (400X). (E) Diffuse intraneuronal staining is a characteristic feature of control brains (A11, 1:250; 400X). (F) 1A9-positive granules were often observed in dendrites ( $\sqcup$ ) in AD brains (400X). (G-H) 4G8 (1:100) and 6E10 (1:100) immunoreactivities were also observed in neurons