

scale in each domain according to the degree of difficulties in life. The anchor-points throughout the domains are as follows: 1: no sign; 2: somewhat but no need to support; 3: special needs by supervisors in groups; 4: special needs by everyone in groups; 5: still difficult even with full-support in groups, and special needs in individual life; and 1.5, 2.5, 3.5 and 4.5 if the conditions are between the adjacent scores. The further criteria for each domain were formulated through repeated meetings among child psychiatrists (22 times in total) and opinions by other professionals. Symptom severity was defined using the concepts of DSM-IV-TR in the domains other than sleep cycle. With respect to learning, we used the concept of learning disorder. We gathered information on behaviors from birth until the present from patients, parents and teachers as much as possible by records and by interview. We evaluated the characteristics of the person assuming an average social environment. If the person was in a very specific social environment, we attempted to exclude the influence of it in our ratings. Similarly, we attempted to exclude the influences of physical diseases or handicaps from our ratings. If the condition was unstable or information was inadequate, we rated the subjects after gathering sufficient information. Most importantly, as we assessed subjects with a range of ages using age-nonspecific measures, the rating was done in reference to normal development. Thus, the knowledge of normal development is an essential requirement for the raters.

Six patients could not remember their language development. Also, we could not score the learning ability in 19 children under school age, especially in the MR group.

2.3. Radar-chart representation

We used a radar-chart representation to visualize the entire profile of 14 domains (Fig. 1). The place of each domain was arranged so that adjacent domains were related. PDD symptoms are at the upper right, domains related to motor are at the lower right, ADHD symptoms are at the lower left and others are at the upper left. With this technique, we could visually grasp the characteristics of subjects at a glance.

2.4. Statistics

To evaluate inter-rater reliability, four trained psychiatrists rated 20 cases independently. The intraclass correlation coefficient (ICC) for each domain was calculated using SPSS 17.0.

Scores of 179 patients on each domain were also analyzed by SPSS 17.0. We compared the scores among the six groups described above using one-way ANOVA, and then we conducted post-hoc tests by the Tukey method.

3. Results

The scoring took about 15 min when adequate information was available.

3.1. Scoring examples

Three examples of the scored charts are shown in Fig. 1a. Case 1 (blue in Fig. 1a) was diagnosed as Asperger's disorder because he did not act as a member of his social group and his interests were very restricted. However, the chief complaint was his going out from the classroom. He also had inattention, that is, ADHD-like symptoms. Patients of this pattern tend to be regarded as ADHD at a glance. The diagnosis of Case 2 (pink in Fig. 1a) was Asperger's disorder, same as Case 1, but the behavioral features were quite different. She was too clumsy to attend physical education class and the chief complaint was repetitive motion. Case 3 (green in Fig. 1a) was diagnosed as PDDNOS and was hypersensitive to noise and touch. He could not stay in the classroom because of noise. Also, he had disabilities in reading and writing.

3.2. Reliability

The inter-rater reliabilities are shown in Table 2 in each domain. The mean and standard deviation of ICC (2,1) was 0.786 ± 0.143 and that of ICC (2,4) was 0.929 ± 0.057 across domains. They were adequately high among the trained psychiatrists participating in the study.

3.3. Feature analysis

We showed mean scores per group in each domain on a radar-chart (Fig. 1b). In general, inattention was present in all the groups. Group variances by one-way ANOVA were significant in the domains other than sleep cycle. We carried out post-hoc tests and show the results in Fig. 2. We also calculated the percentages of patients at clinical level (score of 3 or above) in each group by domains (Table 3).

As a reflection of the diagnostic criteria, all the patients in the two autism and Asperger groups were at clinical level in the domains of sociality, emotion and restricted interests/behavior. In the domain of communication, only one patient with Asperger's disorder had the score of 2, but all other patients in those three groups had the score of 3 or above. In the domains of communication, sociality, and emotion, group differences by post-hoc tests demonstrated a similar pattern: the three groups of autism and Asperger's had equivalent scores in these domains, while the two groups of ADHD scored equivalently.




Fig. 1. Radar-chart representation. (a) Case examples: we showed three cases as examples on a radar-chart. Scores outside red broken lines mean that special supports are required. Cases 1 and 2 had the same diagnosis, Asperger's disorder, but the clinical features were quite different. Case 1 had both symptoms of PDD and ADHD. Case 2 suffered from repetitive motion and clumsiness. Case 3 was diagnosed as PDDNOS and had sensory abnormality and learning disabilities. (b). Mean scores by groups: we showed the mean scores of six diagnostic groups on a radar-chart. Color lines and symbols are explained in the inset. All the groups suffered from inattention. Inattentive-type ADHD had similar but milder features compared with PDDNOS. Combined-type ADHD had a different pattern from other groups. That group had low scores in the domains of restricted interests/behaviors, sensory and motor skills.

Among the three domains of communication, sociality, and emotion, scores in sociality were higher than scores in the other two domains across the groups. In this domain, the average score was around 4 in the groups of autism and Asperger's, indicating the need of considerable assistance from everyone concerned with them. Patients with PDDNOS still needed assistance to some degree at the level of 3, as considered by the leaders or supervisors of their groups. Patients with ADHD had weakness on sociality, but the level of the disturbance was in the range in which they did not need special support.

The degree of restricted interests/behaviors was low in the group with combined-type ADHD, but they still had symptoms to some extent. Half of the patients with inattentive-type ADHD were at clinical level in this domain. The degree in the group with inattentive-type ADHD was slightly higher than that of the group with combined-type ADHD, but this difference was not significant ($p = 0.482$). Also, sensory abnormality and gross and fine motor disabilities were infrequent in the group of combined-type ADHD. None of them were at clinical level in these domains. Variances among individuals on these domains

Table 2
Results of the inter-rater reliability.

Domain	ICC (2,1)	ICC (2,4)
Communication	0.562	0.837
Sociality	0.618	0.866
Emotion	0.563	0.838
Restricted interests/behaviors	0.858	0.960
Sensory	0.855	0.959
Stereotyped/repetitive motion	0.864	0.962
Gross motor	0.909	0.976
Fine motor	0.562	0.837
Inattention	0.800	0.941
Hyperactivity	0.937	0.984
Impulsivity	0.900	0.973
Sleep cycle	0.804	0.942
Learning	0.896	0.972
Language development	0.869	0.964

All the *p*-values by domains were below 0.001 both for ICC (2,1) and ICC (2,4).

were large in all the groups. That is, although some patients suffered much from those features, the mean scores were relatively lower than those in the features such as sociality, communication, and restricted interests/behaviors. Stereotyped/repetitive motion also varied greatly between individuals, but on average was higher in the groups with autism.

Inattention, hyperactivity and impulsivity are known as symptoms of ADHD. However, the other groups also had high scores in those features. The mean scores of inattention were above 3 in all the groups. Also, in the domain of impulsivity, every group had a mean score higher than 2.5 and we did not find group differences in a post-hoc test by the Tukey method, although variation between subgroups by one-way ANOVA was detected ($p = 0.039$). In the domain of hyperactivity, the group of autism with MR had as high a score as the group of combined-type ADHD. Other groups had relatively low scores. However, more than 40% of other PDD patients were hyperactive at the clinical level, whereas most of the patients with inattentive-type ADHD were not hyperactive, as expected from the diagnostic definition. Variances in sleep cycle were large and group differences were not found by one-way ANOVA ($p = 0.362$).

In the domain of learning, only the group of autism without MR had a relatively high score, 2.70 on average. This group showed a significantly higher score than the other groups except for autism with MR. Language development was delayed in the two groups of autism, as expected from the diagnostic definition. Each of these two groups had significantly higher scores than each of the remaining four groups.

4. Discussion

The MSPA was developed with the aim of understanding the entire profile of patients with PDD or ADHD and to facilitate the support that they need more directly. For diagnostic purposes, the assessment tools for these disorders already exist. Many questionnaires have been published and used for each purpose. Questionnaires are useful for gathering information from multiple observers, but there is disagreement between observers especially with psychological and behavioral symptoms as compared with physical symptoms (Marteleto, Lima e Menezes, Tamanaha, Chiari, & Perissinoto, 2008). Thus, questionnaires would be useful for screening, but not adequate for objective assessment of behavioral characteristics.

The currently available objective scales for professional assessment are listed following. The Childhood Autism Rating Scale (CARS) is widely used in clinical practice (Schopler, Reichler, DeVellis, & Daly, 1980) and recently the second edition has been published. The Autism Diagnostic Interview-Revised (ADI-R) (Lord, Rutter, & Le Couteur, 1994), Autism Diagnostic Observation Schedule (ADOS) (Lord et al., 1989) and the Diagnostic Interview for Social and Communication Disorders (DISCO) (Wing, Leekam, Libby, Gould, & Locombe, 2002) are also well known. However, each tool has its own purpose. All the previous four scales are tools for assisting with diagnosis as the names suggest. The advantages of ADI-R, ADOS and DISCO are with high validity for the purpose of diagnosis. ADI-R consists of 93 items and DISCO has 319 items. ADOS is easier to administer than the other two, but still takes about 1 h. Because they are well-structured and take time to administer, they are usually used in research settings rather than in clinical practice (Miles, McCathren, Stichter, & Shinawi, 2010). CARS is as convenient as the present scale and consists of 14 items. However, this is also for the diagnosis of childhood autism, and is formulated so that all the items are correlated and the total score from 14 items indicates the degree of autism (Schopler et al., 1980). That is, the 14 items do not reflect the separate domains. CARS is restricted to children and also CARS, ADI-R and ADOS are intended to assess mainly the core features of autism or PDD. Therefore, they do not always encompass other associated features as independent domains, although each scale includes some of them. The present scale is not only for the assessment of core features of autism or PDD, but also for ADHD, DCD and other associated features with treatment implications.

The most important difference of the present scale from the previous tools is its purpose. The purpose of this scale is not for diagnosis but for understanding behavioral characteristics and difficulties in life. From the standpoint of daily life, even if these patients and their families consult the professionals and receive the diagnosis, their difficulties would continue as long as their living situations do not correspond to their needs by meeting them adequately. When the patients are adolescents or




Fig. 2. Group differences in each domain. Bars indicate the mean scores and the error bars are the standard deviations. * and ** mean that the differences were significant at $0.01 < p < 0.05$, $p \leq 0.01$, respectively. In the domains of communication and sociality, we only showed NS where the group differences were not significant, because most pairs showed significant differences. In other domains, all the significant differences are shown by the above signs.

Table 3
Percentages of patients at clinical level in each group by domains.

	Autism with MR	Autism without MR	Asperger	PDDNOS	ADHD combined	ADHD inattentive
Communication	100	100	97.5	54.1	0	21.4
Sociality	100	100	100	82.5	16.7	28.6
Emotion	100	100	100	58.2	16.7	14.3
Restricted interests/behaviors	100	100	100	83.8	16.7	50
Sensory	57.2	66.7	40	27.1	0	14.2
Stereotyped/repetitive motion	42.9	39	25	9.5	8.3	0
Gross motor	47.6	47.1	23.1	23.3	0	14.2
Fine motor	47.7	29.4	12.8	12.5	0	21.4
Inattention	71.3	77.8	75	64.2	100	100
Hyperactivity	71.3	44.6	40	46	100	0
Impulsivity	71.4	53	62.5	51.4	91.7	42.8
Sleep cycle	28.6	16.7	17.9	18.1	18.2	28.5
Learning	22.2	60	16.2	14.7	9.1	14.3
Language development	100	100	0	13.2	0	7.7

We regarded the clinical level as the score of 3 or above. We calculated each percentage of such patients in each group by domains.

adults, fostering insight or self-understanding would be desirable as well. The diagnosis name with 'disorder' might lower self-esteem (Humphrey & Lewis, 2008), but our chart can be fed back directly to patients and their families at a glance, reflecting needs to be met rather than deficits identified. Since visual cues are known to help recognition in PDD patients (Mesibov & Howley, 2003), the radar-chart format is designed to facilitate insight or self-understanding. Indeed, in our experience, this chart has led to improved insight, to building relationships of mutual trust between patients and psychiatrists, and to patients' acceptance of their diagnosis. Furthermore, it has helped to foster shared understanding of the difficulties between the patients themselves and those in their surroundings.

In addition, the diagnosis alone would be insufficient to represent the entire profile of dysfunction in this group of patients, partly because PDD and ADHD cannot be diagnosed simultaneously in the current diagnostic systems of the International Statistical Classification of Diseases (ICD-10) (WHO, 1992) and DSM-IV-TR, and partly because individual differences still remain even in the same diagnosis. Furthermore, the diagnostic criteria of PDDNOS itself are not fully defined (Myhr, 1998). Not only are the clinical features of these disorders multimodal, but the degree of each feature varies. Therefore, current researchers tend to regard the overall syndrome as a spectrum rather than separate subtypes (Szatmari, 1992). However, the spectrum concept is insufficient to capture the individual features of these patients, and the assessment of the level of each behavioral feature is very important for daily life and social and environmental adaptation. Therefore, we designed this multimodal and nine-stepped representation which is also useful for mild or high-functioning cases. Taken together, we believe that the present scale adds value for treatment beyond the above excellent tools aimed chiefly at diagnosis.

The reliability of the present scale was adequate among trained psychiatrists. However, some training would be needed in raters. Therefore, we are in the process of estimating the amount of training required to increase the number of specialists who can reliably use this assessment. We used the present scale mostly for the outpatients in the department of psychiatry. Therefore, we assessed not only young children but also adolescents and adults. It was very useful for adolescents and adults for understanding themselves. For children, caregivers mainly referred to this scale to support them. The assessment using this scale is based on long-term characteristics, that is, we gathered information from birth throughout development, not limited to the present state. Therefore, the scores are stable and we assume that age does not affect them much when the information is adequate. On the contrary, this scale is not suitable to assess short-term changes such as the effects of intervention. We do not assume that the present scale captures adaptation by intervention or temporary symptoms in exacerbation. However, we are planning a longitudinal study to capture long-term changes of characteristics beyond superficial appearances.

When we consider each domain, sociality is the main domain that divides groups as expected. The outcome that patients with ADHD still had some social difficulty below the clinical level is consistent with a previous report (Nijmeijer et al., 2008). We also showed that inattention was the one domain or feature where all the groups had difficulties at clinical level. Similarly, Strum et al., 2004 reported that 95 out of 101 children with PDD had attention deficit. Thus, we might consider the existence of inattention even in groups other than ADHD, because inattention hinders adaptation in various aspects of daily life. Inattention might be a common feature not only in ADHD but also in PDD. The domain where further consideration is required before a conclusion can be made would be the sleep cycle. Richdale and Schreck (2009) reported that sleep problems are often found in PDD and ADHD, but are multifactorial. They may be an innate feature or may be from stress due to difficulties in life. We will try to detect the innate part and aim to modify the present scale.

Regarding group differences, our results showed that the main difference between the autism and Asperger groups was language development, as defined by the diagnostic criteria. Symptom severity in the Asperger group was milder in the domains of sensory, stereotyped/repetitive motion, and motor skills, but without significant differences (see Fig. 1b). As to the PDDNOS group, symptom severity was even milder than the Asperger group in domains listed as the diagnostic criteria in DSM-IV-TR (communication, sociality, emotion, restricted interests/behaviors, and stereotyped/repetitive motion), but did

not differ in other domains such as motor skills, ADHD symptoms, learning and language development. That is, the difference of features among these three groups is mainly in the domains which constitute the diagnostic definitions of them. Therefore, these three subtypes of PDD are supposed to be a continuum, although they are divided by the diagnostic criteria. Interestingly, features in the group of inattentive-type ADHD were even milder but as a pattern of symptoms, they were similar to those in the group of PDDNOS. This suggests that PDD and ADHD are also related as many previous reports have pointed out (Frazier & Youngstrom, 2006; Jensen et al., 1997; Kochhar et al., 2011; Lee & Ousley, 2006; Nijmeijer et al., 2009; Sinzig et al., 2009; Yoshida & Uchiyama, 2004), indicating the need for a shared assessment tool for PDD and ADHD.

Of interest, only the group of combined-type ADHD showed a different pattern. This group had low scores in the domains of restricted interests/behavior, sensory, and motor skills. Pitcher et al. (2003) also reported that hyperactive/impulsive type had better motor skills than inattentive type. These results identify the possibility that hyperactivity might have positive consequences, such as improved motor skills. These group differences and domain analyses might give novel insights on biological mechanisms of these disorders after further consideration.

5. Conclusions

We developed a multi-dimensional and quantitative assessment chart for PDD and ADHD. The reliability among trained psychiatrists was sufficient. This tool uncovers multi-dimensional clinical features of each patient at a glance, and is useful for fostering understanding of these difficulties by patients themselves and also by those in their surroundings. Furthermore, it discloses the relationships and the differences among subtypes of PDD and ADHD. We hope that it will help patients with these disorders and the clinicians who treat them and also become a useful tool in research settings investigating the biological backgrounds of these disorders.

Acknowledgments

This work was supported by a Health Labour Science Research Grant for Research on Psychiatric and Neurological Diseases and Mental Health (H21-Kokoro-Wakate-021) from the Ministry of Health Labour and Welfare, Japan, and a Grant-in-Aid for Japan Society for the Promotion of Science (JSPS) Fellows (19 45082). We would like to thank all the participants. We also thank Shoshinen Osaka Psychiatric Medical Center and Nagaokakyo City Education Support Center for help in recruiting the participants.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at doi:10.1016/j.ridd.2011.01.052.

References

- American Psychiatric Association. (2000). *Diagnostic and statistical manual of mental disorders-fourth edition-text revision (DSM-IV-TR)*. Washington, DC: American Psychiatric Press.
- Bakker, S. C., van der Meulen, E. M., Buitelaar, J. K., Sandkuijl, L. A., Pauls, D. L., Monsuur, A. J., et al. (2003). A whole-genome scan in 164 Dutch sib pairs with attention-deficit/hyperactivity disorder: Suggestive evidence for linkage on chromosomes 7p and 15q. *American Journal of Human Genetics*, 72, 1251–1260.
- Baron-Cohen, S., Ashwin, E., Ashwin, C., Tavassoli, T., & Chakrabarti, B. (2009). Talent in autism: Hyper-systemizing, hyper-attention to detail and sensory hypersensitivity. *Philosophical Transactions of the Royal Society B*, 364, 1377–1383.
- DeVincent, C. J., & Gadow, K. D. (2009). Relative clinical utility of three Child Symptom Inventory-4 scoring algorithms for differentiating children with autism spectrum disorder vs. attention-deficit hyperactivity disorder. *Autism Research*, 2, 312–321.
- Dewey, D., Cantell, M., & Crawford, S. G. (2007). Motor and gestural performance in children with autism spectrum disorders, developmental coordination disorder, and/or attention deficit hyperactivity disorder. *Journal of International Neuropsychological Society*, 13, 246–256.
- Frazier, T. W., & Youngstrom, E. A. (2006). Evidence-based assessment of attention-deficit/hyperactivity disorder: Using multiple sources of information. *Journal of the American Academy of Child and Adolescent Psychiatry*, 45, 614–620.
- Ghanizadeh, A. (2010). Factor analysis on ADHD and autism spectrum disorder DSM-IV-derived items shows lack of overlap. *European Child and Adolescent Psychiatry*, 19, 797–798.
- Gillberg, I. C., & Gillberg, C. (1988). Children with deficits in attention, motor control and perception (DAMP): Need for specialist treatment. *Acta Paediatrica Scandinavica*, 77, 450–451.
- Harrison, J., & Hare, D. J. (2004). Assessment of sensory abnormalities in people with autistic spectrum disorders. *Journal of Autism and Developmental Disorders*, 34, 727–730.
- Humphrey, N., & Lewis, S. (2008). 'Make me normal': The views and experiences of pupils on the autistic spectrum in mainstream secondary schools. *Autism*, 12, 23–46.
- Ikuzawa, F. M., Matsushita, Y., & Nakase, A. (2001). *Kyoto scale of psychological development 2001*. Kyoto: Kyoto International Social Welfare Exchange Centre.
- Jensen, V. K., Larrieu, J. A., & Mack, K. K. (1997). Differential diagnosis between attention-deficit/hyperactivity disorder and pervasive developmental disorder-not otherwise specified. *Clinical Pediatrics (Philadelphia)*, 36, 555–561.
- Kochhar, P., Batty, M. J., Liddle, E. B., Groom, M. J., Scerif, G., Liddle, P. F., et al. (2011). Autistic spectrum disorder traits in children with attention deficit hyperactivity disorder. *Child: Care, Health and Development*, 37, 103–110.
- Lane, A. E., Young, R. L., Baker, A. E. Z., & Angley, M. T. (2010). Sensory processing subtypes in autism: Association with adaptive behavior. *Journal of Autism and Developmental Disorders*, 40, 112–122.
- Lee, D. O., & Ousley, O. Y. (2006). Attention-deficit hyperactivity disorder symptoms in a clinic sample of children and adolescents with pervasive developmental disorders. *Journal of Child and Adolescent Psychopharmacology*, 16, 737–746.
- Leekam, S. R., Nieto, C., Libby, S. J., Wing, L., & Gould, J. (2007). Describing the sensory abnormalities of children and adults with autism. *Journal of Autism and Developmental Disorders*, 37, 894–910.

- Lichtenstein, P., Carlström, E., Råstam, M., Gillberg, C., & Anckarsäter, H. (2010). The genetics of autism spectrum disorders and related neuropsychiatric disorders in childhood. *American Journal of Psychiatry*, *167*, 1357–1363.
- Lord, C., Rutter, M., Goode, S., Heemsbergen, J., Jordan, H., Mawhood, L., et al. (1989). Autism diagnostic observation schedule: A standardized observation of communicative and social behavior. *Journal of Autism and Developmental Disorders*, *19*, 185–212.
- Lord, C., Rutter, M., & Le Couteur, A. (1994). Autism diagnostic interview-revised: A revised version of a diagnostic interview for caregivers of individuals with possible pervasive developmental disorders. *Journal of Autism and Developmental Disorders*, *24*, 659–685.
- Marteletto, M. R., Lima e Menezes, C. G., Tamanaha, A. C., Chiari, B. M., & Perissinoto, J. (2008). Administration of the Autism Behavior Checklist: Agreement between parents and professionals' observations in two intervention contexts. *Revista brasileira de psiquiatria*, *30*, 203–208.
- Matson, J. L., Boisjoli, J. A., Hess, J. A., & Wilkins, J. (2011). Comorbid psychopathology factor structure on the Baby and Infant Screen for Children with autism traits – Part 2 (BISCUIT-Part 2). *Research in Autism Spectrum Disorders*, *5*, 426–432.
- Matson, J. L., LoVullo, S. V., Rivet, T. T., & Boisjoli, J. A. (2009). Validity of the Autism spectrum disorder-comorbid for children (ASD-CC). *Research in Autism Spectrum Disorders*, *3*, 345–357.
- Matson, J. L., & Nebel-Schwalm, M. S. (2007). Comorbid psychopathology with autism spectrum disorder in children: An overview. *Research in Developmental Disabilities*, *28*, 341–352.
- Mesibov, G., & Howley, M. (2003). *Accessing the curriculum for pupils with autistic spectrum disorders*. London: David Fulton Publishers.
- Miles, J. H., McCathren, R. B., Stichter, J., & Shinawi, M. (2010). Autism spectrum disorders. In R. A. Pagon, T. C. Bird, C. R. Dolan, & K. Stephens (Eds.), *Gene reviews*. Seattle: University of Washington.
- Myhr, G. (1998). Autism and other pervasive developmental disorders: Exploring the dimensional view. *Canadian Journal of Psychiatry*, *43*, 589–595.
- Nijmeijer, J. S., Minderaa, R. B., Buitelaar, J. K., Mulligan, A., Hartman, C. A., & Hoekstra, P. J. (2008). Attention-deficit/hyperactivity disorder and social dysfunctioning. *Clinical Psychology Review*, *28*, 692–708.
- Nijmeijer, J. S., Hoekstra, P. J., Minderaa, R. B., Buitelaar, J. K., Altink, M. E., Buschgens, C. J., et al. (2009). PDD symptoms in ADHD, an independent familial trait? *Journal of Abnormal Child Psychology*, *37*, 443–453.
- Ogdie, M. N., Macphie, I. L., Minassian, S. L., Yang, M., Fisher, S. E., Francks, C., et al. (2003). A genomewide scan for attention-deficit/hyperactivity disorder in an extended sample: Suggestive linkage on 17p11. *American Journal of Human Genetics*, *72*, 1268–1279.
- Pan, C. Y., Tsai, C. L., & Chu, C. H. (2009). Fundamental movement skills in children diagnosed with autism spectrum disorders and attention deficit hyperactivity disorder. *Journal of Autism and Developmental Disorders*, *39*, 1694–1705.
- Pitcher, T. M., Piek, J. P., & Hay, D. A. (2003). Fine and gross motor ability in males with ADHD. *Developmental Medicine and Child Neurology*, *45*, 525–535.
- Richdale, A. L., & Schreck, K. A. (2009). Sleep problems in autism spectrum disorders: Prevalence, nature, and possible biopsychosocial aetiologies. *Sleep Medicine Reviews*, *13*, 403–411.
- Russell, G., Ford, T., Steer, C., & Golding, J. (2010). Identification of children with the same level of impairment as children on the autistic spectrum, and analysis of their service use. *Journal of Child Psychology and Psychiatry*, *51*, 643–651.
- Schopler, E., Reichler, R. J., DeVellis, R. F., & Daly, K. (1980). Toward objective classification of childhood autism: Childhood Autism Rating Scale (CARS). *Journal of Autism and Developmental Disorders*, *10*, 91–103.
- Sinzig, J., Walter, D., & Doepfner, M. (2009). Attention deficit/hyperactivity disorder in children and adolescents with autism spectrum disorder: Symptom or syndrome? *Journal of Attention Disorders*, *13*, 117–126.
- Smalley, S. L., Kustanovich, V., Minassian, S. L., Stone, J. L., Ogdie, M. N., McGough, J. J., et al. (2002). Genetic linkage of attention-deficit/hyperactivity disorder on chromosome 16p13, in a region implicated in autism. *American Journal of Human Genetics*, *71*, 959–963.
- Staples, K. L., & Reid, G. (2010). Fundamental movement skills and autism spectrum disorders. *Journal of Autism and Developmental Disorders*, *40*, 209–217.
- Strum, H., Fernell, E., & Gillberg, C. (2004). Autism spectrum disorders in children with normal intellectual levels: Associated impairments and subgroups. *Developmental Medicine and Child Neurology*, *46*, 444–447.
- Szatmari, P. (1992). The validity of autistic spectrum disorders: A literature review. *Journal of Autism and Developmental Disorders*, *22*, 583–600.
- Wechsler, D. (1991). *Manual for the Wechsler intelligence scale for children* (3rd ed.). San Antonio, Texas: The Psychological Corporation.
- Wechsler, D. (1997). *The Wechsler adult intelligence scale* (3rd ed.). San Antonio, Texas: Psychological Corp (Harcourt).
- Whitman, T. L. (2004). *The development of autism: A self-regulatory perspective*. London: Jessica Kingsley Publishers.
- Wing, L., Leekam, S. R., Libby, S. J., Gould, J., & Larcombe, M. (2002). The diagnostic interview for social and communication disorders: Background, inter-rater reliability and clinical use. *Journal of Child Psychology and Psychiatry*, *43*, 307–325.
- World Health Organization. (1992). *International classification of diseases and related health problems (ICD-10)*. Geneva: WHO.
- Yamagata, T., Aradhya, S., Mori, M., Inoue, K., Momoi, M. Y., & Nelson, D. L. (2002). The human secretin gene: Fine structure in 11p15.5 and sequence variation in patients with autism. *Genomics*, *80*, 185–194.
- Yoshida, Y., & Uchiyama, T. (2004). The clinical necessity for assessing attention deficit/hyperactivity disorder (AD/HD) symptoms in children with high-functioning pervasive developmental disorder (PDD). *European Child and Adolescent Psychiatry*, *13*, 307–314.

