

(c) Co-localization of Rab GTPases and LAMP2. Raw264.7 macrophages transfected with EGFP-fused Rab GTPases were stained with anti-LAMP-2 antibody, followed by observation with laser scanning confocal microscopy. Pearson's correlation (PC) of overlapping fluorescent areas of EGFP-fused Rab GTPases and LAMP-2 was assessed. -, PC < 0, +; 0 < PC < 0.3, ++; PC > 0.3.

References

1. Tisdale EJ, Bourne JR, Khosravi-Far R, Der CJ, Balch WE. GTP-binding mutants of rab1 and rab2 are potent inhibitors of vesicular transport from the endoplasmic reticulum to the Golgi complex. *J Cell Biol* 1992;119:749-761.
2. Abu-Amer Y, Teitelbaum SL, Chappel JC, Schlesinger P, Ross FP. Expression and regulation of RAB3 proteins in osteoclasts and their precursors. *J Bone Miner Res* 1999;14:1855-1860.
3. van der Sluijs P, Hull M, Webster P, Male P, Goud B, Mellman I. The small GTP-binding protein rab4 controls an early sorting event on the endocytic pathway. *Cell* 1992;70:729-740.
4. Bucci C, Parton RG, Mather IH, Stunnenberg H, Simons K, Hoflack B, Zerial M. The small GTPase rab5 functions as a regulatory factor in the early endocytic pathway. *Cell* 1992;70:715-728.
5. Martinez O, Schmidt A, Salamero J, Hoflack B, Roa M, Goud B. The small GTP-binding protein rab6 functions in intra-Golgi transport. *J Cell Biol* 1994;127:1575-1588.
6. Bucci C, Thomsen P, Nicoziani P, McCarthy J, van Deurs B. Rab7: a key to lysosome biogenesis. *Mol Biol Cell* 2000;11:467-480.
7. Yang M, Chen T, Han C, Li N, Wan T, Cao X. Rab7b, a novel lysosome-associated small GTPase, is involved in monocytic differentiation of human acute promyelocytic leukemia cells. *Biochem Biophys Res Commun* 2004;318:792-799.
8. Chen YT, Holcomb C, Moore HP. Expression and localization of two low molecular weight GTP-binding proteins, Rab8 and Rab10, by epitope tag. *Proc Natl Acad Sci U S A* 1993;90:6508-6512.
9. Lombardi D, Soldati T, Riederer MA, Goda Y, Zerial M, Pfeffer SR. Rab9 functions in transport between late endosomes and the trans Golgi network. *EMBO J* 1993;12:677-682.
10. Cardoso CM, Jordao L, Vieira OV. Rab10 regulates phagosome maturation and its overexpression rescues *Mycobacterium*-containing phagosomes maturation. *Traffic* 2010;11:221-235.
11. Cox D, Lee DJ, Dale BM, Calafat J, Greenberg S. A Rab11-containing rapidly recycling compartment in macrophages that promotes phagocytosis. *Proc Natl Acad Sci U S A* 2000;97:680-685.
12. Nokes RL, Fields IC, Collins RN, Folsch H. Rab13 regulates membrane trafficking between TGN and recycling endosomes in polarized epithelial cells. *J Cell Biol* 2008;182:845-853.
13. Junutula JR, De Maziere AM, Peden AA, Ervin KE, Advani RJ, van Dijk SM, Klumperman J, Scheller RH. Rab14 is involved in membrane trafficking between the Golgi complex and endosomes. *Mol Biol Cell* 2004;15:2218-2229.
14. Pavlos NJ, Xu J, Riedel D, Yeoh JS, Teitelbaum SL, Papadimitriou JM, Jahn R, Ross FP, Zheng MH. Rab3D regulates a novel vesicular trafficking pathway that is required for osteoclastic bone resorption. *Mol Cell Biol* 2005;25:5253-5269.
15. Dejgaard SY, Murshid A, Erman A, Kizilay O, Verbich D, Lodge R, Dejgaard K, Ly-Hartig TB, Pepperkok R, Simpson JC, Presley JF. Rab18 and Rab43 have key roles in ER-Golgi trafficking. *J Cell Sci* 2008;121:2768-2781.
16. Das Sarma J, Kaplan BE, Willemsen D, Koval M. Identification of rab20 as a potential regulator of connexin 43 trafficking. *Cell Commun Adhes* 2008;15:65-74.
17. Pellinen T, Arjonen A, Vuoriluoto K, Kallio K, Fransén JA, Ivaska J. Small GTPase Rab21 regulates cell adhesion and controls endosomal traffic of beta1-integrins. *J Cell Biol* 2006;173:767-780.
18. Kauppi M, Simonsen A, Bremnes B, Vieira A, Callaghan J, Stenmark H, Olkkonen VM. The small GTPase Rab22 interacts with EEA1 and controls endosomal membrane trafficking. *J Cell Sci* 2002;115:899-911.
19. Ng EL, Wang Y, Tang BL. Rab22B's role in trans-Golgi network membrane dynamics. *Biochem Biophys Res Commun* 2007;361:751-757.
20. Evans TM, Ferguson C, Wainwright BJ, Parton RG, Wicking C. Rab23, a negative regulator of hedgehog signaling, localizes to the plasma membrane and the endocytic pathway. *Traffic* 2003;4:869-884.
21. Strom M, Hume AN, Tarafder AK, Barkagianni E, Seabra MC. A family of Rab27-binding proteins. Melanophilin links Rab27a and myosin Va function in melanosome transport. *J Biol Chem* 2002;277:25423-25430.
22. Wasmeier C, Romao M, Plowright L, Bennett DC, Raposo G, Seabra MC. Rab38 and Rab32 control post-Golgi trafficking of melanogenic enzymes. *J Cell Biol* 2006;175:271-281.
23. Wang T, Hong W. Interorganellar regulation of lysosome positioning by the Golgi apparatus through Rab34 interaction with Rab-interacting lysosomal protein. *Mol Biol Cell* 2002;13:4317-4332.
24. Kouranti I, Sachse M, Arouche N, Goud B, Echard A. Rab35 regulates an endocytic recycling pathway essential for the terminal steps of cytokinesis. *Curr Biol* 2006;16:1719-1725.
25. Masuda ES, Luo Y, Young C, Shen M, Rossi AB, Huang BC, Yu S, Bennett MK, Payan DG, Scheller RH. Rab37 is a novel mast cell specific GTPase localized to secretory granules. *FEBS Lett* 2000;470:61-64.

Table S2 Primer list for construction of plasmid of EGFP-fused Rab GTPases

Rab GTPase	Primer forward	Primer reverse
Rab1	CAGATCTatgtccagcatgaatcccgaatatg	CGGTACCTtagcagcaacctcccactg
Rab1b	CAGATCTatgaaccccgaatgatgactacctgtttaag	CGAATTCtagcagcagccaccgctagcaggct
Rab2	CAGATCTatggcgtacgcctatctcttcaag	CCGAATTCtaacagcagccgccccca
Rab2b	CAGATCTatgacttatgcttatctcttcaagtatac	CGAATTCtagcagcagccagagttggaccctatgct
Rab3	CCTCGAGCTatggcatccgccacagactcgc	CGAATTCtagcagggcgcagctcctggt
Rab3b	CCTCGAGCTatggcttcagtgacagatggtaaaactgg	CGAATTCtagcatgagcagttctgctgcagcagc
Rab4	CAGATCTatgtcgcagacggccatgtccgaaac	CGAATTCtaacaaccacactcctgagcgttcg
Rab4b	CAGATCTatggctgagacctacgactcctct	CGAATTCagcagccacacggctgaggggccc
Rab5	CAGATCTatggctagtcgagcgcacaacagac	CGAATTCtagttactacaacactgattcctg
Rab5b	CCTCGAGCTatgactagcagaagcacagctaggcc	CGAATTCagttgctacaacactggctctg
Rab6	CCTCGAGCTatgtccacggggcggagactt	CGAATTCtagcaggaacagcctccttcac
Rab6b	CAGATCTatgtccgcagggggagattttggga	CGAATTCtagcagcagcagccgctcctgctggc
Rab7b	CCTCGAGCTatgaatccccggaagaaggt	CGAATTCagcagcatctgctccttactggtctgg
Rab8	CAGATCTatggcgaagacctacgattacctg	CGAATTCcacagaagaacacatcgaaaaagctg
Rab8b	CAGATCTatggcgaagacctacgattatcttcaag	CGAATTCaaagtagcagcaacgaaagaaactggtc
Rab9	CAGATCTatggcaggaataatcatcactttttaaag	CGGTACCTcaacagcaaatgagctaggc
Rab9b	CCTCGAGCTatgagtggaataatcctgctttaaag	CGAATTCtaacagcagcaagacctgctttggagc
Rab10	CAGATCTatggcgaagaagactacgacctg	CGAATTCtagcagcatttctcttccagc
Rab11	CAGATCTatgggcacccgcgacgacgag	CGAATTCtagatgttctgacagcactgcacc
Rab11b	CCTCGAGCTatggggaccgggacgacgagtagc	CGAATTCacaggttctggcagcactgcagctt
Rab13	CAGATCTatggcgaagcctacgaccac	CGAATTCtagccaggaggcactgttg
Rab14	CAGATCTatggcaactgcaccatacaactactc	CGAATTCtagcagccacagcctctc
Rab16	CCTCGAGCTatggcatcagctggagacacc	CGAATTCtagcagctgcagctgctg
Rab18	CAGATCTatggacgaggcagctgctaacca	CGAATTCtataacacagagcaataaccaccacag
Rab20	CCTCGAGCTatgaggaagcccagacagcaag	CGAATTCtaggcacaaccccagatctg
Rab21	CAGATCTccgggaagcagcgggatggc	CGAATTCttaccagaagaacagcaccctcc
Rab22a	CAGATCTatggcgtgaggagctcaa	CGGTACCTcagcagcagctccgcttg
Rab22b	CAGATCTatgatggcgatacgggagctcaaagtg	CGAATTCaacagcaccggcggctggcttg
Rab23	CCTCGAGCTatgttgaggagaagatggaagtgc	CAAGCTtagggtatgctacagctgctaaaagg
Rab24	CAGATCTatgagcgggcagcgcgtgga	CGAATTCtagtgatgacaacagctgtagaag
Rab27	CAGATCTatgtctgatggagattatgattacctc	CGAATTCtaacagccacatgccctttctc
Rab28	CAGATCTatgtcggactctgaggaggag	CGAATTCcactgaactgcacacatagagcttc
Rab29	CAGATCTatggcagccgcgaccacct	CGAATTCtagcagcaggaccagctgg
Rab30	CCTCGAGCTatgagtatggaagattatgattcctgctc	CGGTACCTtagttgaaattacaacaagtcaaatagctg
Rab32	CCAGATCTatggcgggcggaggagccgg	CGAATTCtagcaacactgggatttctctg
Rab34	CCTCGAGCTatgaacattctggcaccctg	CGGTACCTcatgggcaacatgtggcttc
Rab35	CCTCGAGCTatggcccgggactacgacca	CGAATTCtagcagcagcttcttctgttactg
Rab37	CCTCGAGCTatgacgggcacgccaggcgcctgtg	CGAATTCacatgaaggagcagcagctggag
Rab38	CAGATCTatgcaggcccccacaaagga	CGAATTCtaggatttggcacagccagag
Rab39	CAGATCTatggagaccatctggatctaccag	CGAATTCtagcagaagcattcttctctggg
Rab43	CAAGCTTCggcccttggctcttctag	CGAATTCggctctgagggtctggctt

Table S3 Primer list for site-directed mutagenesis

Rab	GTase	dominant negative	constitutive active	dominant negative-F	dominant negative-R	constitutive active-F	constitutive active-R
Rab5	S34N		Q79L	agagctcgcgttggcaaaAA Tggccatgctctgtttt	aaacgaagcaactggctcA Ttttcccaacagggactct	gaatgggatacagctgctc Tgaaacatacattagcctag	ciaagctatgatactgctcA gaaacagcigtatccatattc
Rab7b	T22N		Q67L	gccattggttgggaagaA cctccctccaccabaatg	catattggtggggggag Tcttcccaacaacaaaggc	gactgggcaacagcggctc Tggagcggctcctccatg	catggagcggaaacgctccA gaccgcccgttccagatc
Rab8	T22N		Q67L	gactcgggggtgggaagaA cigtgctcgttccgcttc	gagcgggaacagggcaag Tcttcccccaccocccagtc	gatatgggaacaacccggctc Tgaaacggtttcggacgac	gatactccgaaacccgttccA gaaacccgcttccatatac
Rab8b	T22N		Q67L	ctcgggggtaggcaagaA cigtccctcigtccgcttc	gagaagcggaaacagggggcag Tcttggctacccccggg	gatagggcaacagcggctc Tggaaagattccgaaacaac	gattgtcggaaatcttccA gaaacccgcttccatatac
Rab9	S21N		Q66L	gattggtgggttgggaagaA ttcactatgaaacagatag	catatcigtctataagtgaa Tcttcccaactccaccatc	gggaacagggcaagctc Tggagcgttccgaaaccggagac	gctccagggctcggaaatcgctccA gaaacccgcttccatatac
Rab9b	S21N		Q66L	gattggtgggttgggaagaA ctcgcttatgaaacgcttag	cgtaacggcttcaatagcgaA Tcttcccaactccaccatc	c-tgggacaclycaaggc Tggaaacgttccgaaaccggagac	gctccagggctcggaaatcgctccA gaaacccgcttccatatac
Rab11	S25N		Q70L	gattcctggttgggaagaA taatctcctctctcgtattac	gtaaticggacagggatata Tcttcccaacacagaaatc	gatatgggaacaacagggc Tggaaacgttccgaaaccggagac	gctccagggctcggaaatcgctccA gaaacccgcttccatatac
Rab11b	S25N		Q70L	gactcagcgttgggaagaA caaacctcgtcgtcgcctc	gaaagcgaacagcaggttg Tcttggcccaaccccgcttc	gatacgggaacaacagggc Tggaaacgttccgaaaccggagac	gctccagggctcggaaatcgctccA gaaacccgcttccatatac
Rab13	T22N		Q67L	gactcgggggtgggaagaA cigtctctatcctcgtcttg	caaagcgaatgatccagcaA Tcttcccccaccocccagtc	gatacgggaacaacagggc Tggaaacgttccgaaaccggagac	gctccagggctcggaaatcgctccA gaaacccgcttccatatac
Rab14	S25N		Q70L	gacatggggtaggaaasaA ttgcttctcctcaaattac	gtaaatgatgaaagcaagcaA Ttttctctactcccaatc	gatacgggaacaacagggc Tggaaacgttccgaaaccggagac	gctccagggctcggaaatcgctccA gaaacccgcttccatatac
Rab20	T19N		R59L	catgaaagtggggaagaA Tcgtcgtcagcagcgtatag	calalaccgctcagcagcgaA Tcttcccccactgcaatg	gggaacacagcagggc Tggaaacgttccgaaaccggagac	gctccagggctcggaaatcgctccA gaaacccgcttccatatac
Rab22a	S19N		Q64L	caaggttagttaaA cagattggtggcaggttgg	ccaacaaccgcaacaactactG Ttttaactcaaccg	c-tgggatacagctggac Tcgaacgatttctgcttag	gctccagggctcggaaatcgctccA gaaacccgcttccatatac
Rab20	S19N		Q65L	gacaactggggttgggaasaA Ccagcatcgtcgtcgtattgc	gacaatcggccaccogtagctG Ttttcccaaccccagctc	catctgggaacaacagggc Tggaaacgttccgaaaccggagac	gctccagggctcggaaatcgctccA gaaacccgcttccatatac
Rab23	S20N		Q68L	gaaatggcagttgggaasaA Tgatagtatcagcgalattg	caatatcgtgaatcatacA Ttttcccaactgctccatc	gatacgggaacaacagggc Tggaaacgttccgaaaccggagac	gctccagggctcggaaatcgctccA gaaacccgcttccatatac
Rab27	T23N		Q78L	ctctggttgggaagaA cagatgctcttaccatatac	gtatattgtaagtagcacag Tcttcccaaccccagctc	gatacgggaacaacagggc Tggaaacgttccgaaaccggagac	gctccagggctcggaaatcgctccA gaaacccgcttccatatac
Rab32	T39N		Q85L	gagcttggcgttgggaagaA cagcatcatcaagcgtcag	cgtagccttgtagtagctg Tcttggcccaaccccagctc	gatacgggaacaacagggc Tggaaacgttccgaaaccggagac	gctccagggctcggaaatcgctccA gaaacccgcttccatatac
Rab34	T69N		Q111L	gaccctgggtgggaagaA ttgctctcatatagcttc	gaaactatagtgggcaA Tcttcccccaccocccagctc	c-ttgggatacaccggc Tggaaacgttccgaaaccggagac	gctccagggctcggaaatcgctccA gaaacccgcttccatatac
Rab37	T43N		Q89L	gagacacagcgtcggcaaaaA Cigtctcctgatccaatc	gaaactatagtgggcaA Tcttcccccaccocccagctc	gatacgggaacaacagggc Tggaaacgttccgaaaccggagac	gctccagggctcggaaatcgctccA gaaacccgcttccatatac
Rab38	T23N		Q69L	gactcggggtgggaagaA cagtatcatcaagcgtcag	gaaactatagtgggcaA Tcttcccccaccocccagctc	gatacgggaacaacagggc Tggaaacgttccgaaaccggagac	gctccagggctcggaaatcgctccA gaaacccgcttccatatac
Rab39	S22N		Q72L	gactccacccgttgggaagaA cigtctcctcagcccttc	cgtagccttgtagtagctg Tcttcccccaccocccagctc	c-tgggatacaccggc Tggaaacgttccgaaaccggagac	gctccagggctcggaaatcgctccA gaaacccgcttccatatac
Rab43	T32N		Q77L	cgagcaagcgttgggaagaA Cigtggttggcagccttc	gaaagcgtcagcccagcaA Tcttggcccaaccccagctc	gatacgggaacaacagggc Tggaaacgttccgaaaccggagac	gctccagggctcggaaatcgctccA gaaacccgcttccatatac




Figure S1-1




Figure S1-2




Figure S1-3




Figure S1-4




Figure S1-5




Figure S2A




Figure S2B




Figure S3




Figure S4




Figure S5



RESEARCH ARTICLE

A novel vaccine strategy to induce mycobacterial antigen-specific Th1 responses by utilizing the C-terminal domain of heat shock protein 70

Tomohiro Uto^{1,2}, Kunio Tsujimura¹, Masato Uchijima¹, Shintaro Seto¹, Toshi Nagata^{1,3}, Takafumi Suda², Kingo Chida², Hirotohi Nakamura² & Yukio Koide¹

¹Department of Infectious Diseases, Hamamatsu University School of Medicine, Hamamatsu, Japan; ²Department of Internal Medicine, Hamamatsu University School of Medicine, Hamamatsu, Japan; and ³Department of Health Science, Hamamatsu University School of Medicine, Hamamatsu, Japan

Correspondence: Yukio Koide, Department of Infectious Diseases, Hamamatsu University School of Medicine, 1-20-1 Handa-yama, Higashi-ku, Hamamatsu 431-3192, Japan. Tel.: +81 53 435 2101; fax: +81 53 435 2335; e-mail: koidelb@hama-med.ac.jp

Received 30 June 2010; revised 26 October 2010; accepted 19 November 2010.

DOI:10.1111/j.1574-695X.2010.00762.x

Editor: Willem van Eden

Keywords

Mycobacterium tuberculosis; heat shock protein; DNA vaccine; IFN- γ .

Abstract

Heat shock protein 70 (HSP70) is a member of a highly conserved superfamily of intracellular chaperones called stress proteins that can activate innate and adaptive immune responses. We evaluated the effect of a fusion DNA vaccine that encoded mycobacterial HSP70 and MPT51, a major secreted protein of *Mycobacterium tuberculosis*. Spleen cells from mice immunized with fusion DNA of full-length HSP70 and MPT51 produced a higher amount of interferon- γ (IFN- γ) in response to the CD4+, but not the CD8+ T-cell epitope peptide on MPT51 than those from mice immunized with MPT51 DNA. Furthermore, because HSP70 comprises the N-terminal ATPase domain and the C-terminal peptide-binding domain, we attempted to identify the domain responsible for its enhancing effect. The fusion DNA vaccine that encoded the C-terminal domain of HSP70 and MPT51 induced a higher MPT51-specific IFN- γ production by CD4+ T cells than the vaccine that encoded MPT51 alone, whereas that with the N-terminal domain did not. Similar results were obtained by immunization with the fusion proteins. These results suggest that the DNA vaccine that encodes a chimeric antigen molecule fused with mycobacterial HSP70, especially with its C-terminal domain, can induce a stronger antigen-specific T-helper cell type 1 response than antigen DNA alone.

Introduction

Tuberculosis is a major cause of death worldwide. There were an estimated 9.4 million incident and 11.1 million prevalent cases of tuberculosis, and 1.8 million people died of tuberculosis, in 2008 (World Health Organization, 2009). *Mycobacterium bovis* Bacillus Calmette–Guerin (BCG) is the only available vaccine against tuberculosis, but it has been reported to show variable protective efficacy (Colditz *et al.*, 1994). Although BCG can protect against tuberculosis in childhood, its efficacy in adults varies, especially in endemic countries (Sterne *et al.*, 1998; Kaufmann, 2004; Andersen, 2007). It is also known to represent a risk for immunocompromised patients. In addition, the emergence of multidrug-resistant strains of *Mycobacterium tuberculosis* has lent urgency to the search for novel therapeutic agents and the

development of more effective vaccines capable of protecting adults (Kaufmann, 2000).

MPT51 is a major secreted protein of *M. tuberculosis*, which has a molecular weight of 27 kDa and a primary structure that is similar to the components of the antigen 85 complex (Nagai *et al.*, 1991; Ohara *et al.*, 1995, 1997). We demonstrated previously that MPT51 induces T-cell-mediated immune responses and protective immunity against challenge with *M. tuberculosis* in mice (Miki *et al.*, 2004), and have identified one CD8+ and two CD4+ T-cell epitopes that are presented by H2-D^d and H2-A^b, respectively (Suzuki *et al.*, 2004). MPT51 is recognized by T cells from patients with active tuberculosis and is also considered to be immunogenic in humans (de Araujo-Filho *et al.*, 2008).

Heat shock protein70 (HSP70) is a member of a highly conserved superfamily of intracellular chaperones called

stress proteins (Gething *et al.*, 1995). In addition, HSP70 appears to play important roles in the innate and adaptive immune responses, such as receptor-mediated antigen internalization by sentinel antigen-presenting cells (APCs), stimulation of production of various cytokines, and maturation of dendritic cells (DCs) (Suto & Srivastava, 1995; Basu *et al.*, 2000; Binder *et al.*, 2000; Castellino *et al.*, 2000; Singh-Jasuja *et al.*, 2000). Collectively, the ability of HSP70 to bind antigenic peptides and deliver them to APCs profoundly contributes to the generation of antigen-specific T-cell responses.

In this study, we compared the effect of a DNA vaccine that encoded a fusion protein that consisted of MPT51, a protective antigen against *M. tuberculosis* (Miki *et al.*, 2004), and *M. tuberculosis* HSP70 on the induction of MPT51-specific T-cell responses, with that of MPT51 DNA alone. After showing superior MPT51-specific T-cell induction by the fusion molecule, we attempted to establish the domain of HSP70 that was responsible for this enhancing effect.

Materials and methods

Construction of fusion genes for DNA vaccines

Mycobacterium tuberculosis HSP70 gene was cloned by PCR from genomic DNA of the H37Rv strain. The primers used for PCR were as follows:

HSP70F (full-length HSP70: residues 1–625)

5'-TATGAATTCACCATGGCTCGTGCGGTCGGG-3' (forward)

5'-AATGGTACCCTTGGCCTCCCGGCCGT-3' (reverse)

HSP70N (N-terminal domain of HSP70: residues 1–360)

5'-TATGAATTCACCATGGCTCGTGCGGTCGGG-3' (forward)

5'-AATGGTACCCACCTCGCCCTTGAGGA-3' (reverse)

HSP70C (C-terminal domain of HSP70: residues 354–625)

5'-TATGAATTCACCATGGCGTCCTCAAGGGC-3' (forward)

5'-AATGGTACCCTTGGCCTCCCGGCCGT-3' (reverse)

Forward and reverse primers contained EcoRI and KpnI restriction sites (underlined), respectively. PCR products were purified and cloned into the EcoRI/KpnI sites of the pCI vector (Promega, Madison, WI) that contained MPT51 DNA at the XbaI site (Suzuki *et al.*, 2004). A nucleotide sequence for a 12-amino-acid (GGGSGGGSGGGS) linker was then inserted into the KpnI sites of pCI vectors that contained chimeric MPT51 DNA. The chimeric DNA constructs thus prepared were designated *HSP70F-MPT51*, *HSP70N-MPT51*, and *HSP70C-MPT51*. As a control, *HSP70F* was amplified by PCR using a primer set as follows: 5'-TATGAATTCACCATGGCTCGTGCGGTCGGG-3' (forward, an underline indicates the EcoRI site) and 5'-AATGGTACCCTCAGCTTGGCCTCCCGGCCGT-3' (reverse,

an underline indicates the KpnI site). The PCR product was purified and cloned into the EcoRI/KpnI sites of the pCI vector. The pCI plasmids containing these constructs were propagated in *Escherichia coli* strain XL1-Blue (Stratagene, La Jolla, CA) and purified by EndoFree Plasmid Maxi Kit (Qiagen GmbH, Hilden, Germany). The concentration of endotoxin was tested using the Endospecy ES-24S kit and the Toxicolor DIA kit (both from Seikagaku Biobusiness Corporation, Tokyo, Japan), and all plasmid solutions were found to contain < 0.1 EU endotoxin μg^{-1} DNA.

Preparation of fusion proteins

For protein preparation, *MPT51*, *HSP70F-MPT51*, *HSP70N-MPT51*, and *HSP70C-MPT51* were 6 × histidine-tagged by PCR using the corresponding forward primers listed above and a reverse primer (CAAGCTTTTAATGATGATGATGATGATGGCGGATCGCACCGACGATAT) containing nucleotide sequences for the extreme C-terminal of MPT51, the histidine-tag (double underlined), and the HindIII site (underlined). The PCR products were purified and cloned into the EcoRI/HindIII sites of the pRSET expression vector (Invitrogen, Carlsbad, CA). *Escherichia coli* JM109 competent cells were transformed with the expression vectors, and proteins were induced according to the manufacturer's instructions. Proteins were extracted and purified by Ni²⁺-nitrilotriacetic acid (Ni-NTA) agarose (Qiagen GmbH) in the presence of 7 M urea, according to the manufacturer's instructions. Purified proteins were refolded by decreasing the concentration of urea while immobilized on the Ni-NTA matrix and then eluted by phosphate-buffered saline (PBS) (pH 7.4) containing 250 mM imidazole. The buffer was finally exchanged with PBS (pH 7.4) using a PD-10 desalting column (GE Healthcare, Uppsala, Sweden). The purity of recombinant proteins was checked by sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE). The concentration of endotoxin was tested using the Endospecy ES-24S kit and the Toxicolor DIA kit, and recombinant protein solutions contained < 10 EU endotoxin mg^{-1} protein.

Preparation of anti-MPT51 monoclonal antibody (mAbs)

Anti-MPT51 mAbs were prepared as described previously (Harlow & Lane, 1988). Briefly, purified recombinant MPT51 protein (50 μg per mouse) was mixed with the RIBI Adjuvant System (Corixa, Hamilton, MT) and injected subcutaneously into BALB/c mice three times at 3-week intervals. Spleen cells from immunized mice were fused with SP2/0, and stably hybridized cells were selected using HAT medium (Invitrogen). Hybridomas that produced anti-MPT51 mAbs were screened by enzyme-linked immunosorbent assay (ELISA), and two clones (2B11F5 and 2D9H2) were established by limiting dilution. Both clones produced

anti-MPT51 mAbs of the IgG1 isotype, as determined using a Mouse Immunoglobulin Isotype Kit (BD Biosciences Pharmingen, San Diego, CA).

Western blotting detection of chimeric MPT51 proteins

HEK293T cells were transfected with pCI plasmids that encoded MPT51 DNA of various forms using FuGENE 6 (Roche Diagnostics GmbH, Mannheim, Germany). Cells were collected 24 h after transfection and lysed in the radio-immunoprecipitation assay buffer (25 mM Tris-HCl, pH 7.6, 150 mM NaCl, 1% NP-40, 1% sodium deoxycholate, 0.1% SDS), and Western blotting was performed with mouse anti-MPT51 mAb (from 2B11F5), HRP-conjugated goat anti-mouse immunoglobulins (Zymed Laboratories/Invitrogen), and West Pico Chemiluminescent Substrate (Pierce, Rockford, IL).

Animals and immunization

Female BALB/c and C57BL/6 mice (between 7 and 9 weeks of age) were purchased from Japan SLC (Hamamatsu, Japan) and maintained at the Institute for Experimental Animals, Hamamatsu University School of Medicine. All animal experiments were performed according to the Guidelines for Animal Experimentation, Hamamatsu University School of Medicine.

Mice were epidermally immunized with plasmids (2 µg per mouse per immunization) using the Helios gene gun system (Bio-Rad Laboratories, Hercules, CA) as described previously (Suzuki *et al.*, 2004) or recombinant proteins (50 µg per mouse) emulsified with incomplete Freund's adjuvant (Difco Laboratories/BD Diagnostic Systems, Sparks, MD) by injecting subcutaneously three times at 10-day intervals.

ELISA for interferon-γ (IFN-γ)

Two weeks after the last immunization, spleen cells from immunized BALB/c and C57BL/6 mice were cultured in a 96-well plate ($1 \times 10^6/0.2$ cells mL⁻¹) in the presence of MPT51 peptides (1 µM) for 2 and 4 days, respectively. Antigenic peptides P24-32 and P171-190 (a dominant H2-A^b-restricted epitope) (Suzuki *et al.*, 2004) were used for CD8+ and CD4+ T-cell stimulation, respectively. For protein immunization, spleen cells from immunized or naive C57BL/6 mice were stimulated with an antigenic peptide P171-190 (1 µM) for 4 days as described above. The IFN-γ concentration of the culture supernatants was determined by a sandwich ELISA as described previously (Yoshida *et al.*, 1995).

Preparation of total RNA and semi-quantitative reverse transcriptase (RT)-PCR

Two weeks after the last immunization, spleen cells from immunized C57BL/6 mice were cultured in a 12-well plate (1×10^7 cells mL⁻¹ per well) in the presence of 1 µM MPT51 peptide (P171-190) for 16 h. Total RNA was prepared using ISOGEN (Nippon Gene, Tokyo, Japan), and semi-quantitative RT-PCR was performed as described previously (Uchijima *et al.*, 2005) using primers as follows:

inducible nitric oxide synthase (iNOS)

5'-TACAAAGATGACCCTAAGAGT-3' (forward)

5'-ACATGGCCGAGCGTCAAAGA-3' (reverse)

IFN-γ

5'-TCTGAGACAATAAACGCTAC-3' (forward)

5'-GAATCAGCAGCGACTCCTTT-3' (reverse)

glycerol-3-phosphate dehydrogenase (G3PDH)

5'-ACCACAGTCCAT CCATCAC-3' (forward)

5'-TCCACCACCCTGTTGCTGTA-3' (reverse)

Statistical analysis

Statistical analyses were performed using STATVIEW-J 5.0 statistics program (SAS Institute Inc., Cary, NC). The Mann-Whitney test was used to calculate *P*-values.

Results

Enhancement of MPT51-specific T-cell responses by full-length HSP70

We first examined the enhancing effect of full-length HSP70 (HSP70F) for the induction of MPT51-specific T-cell responses. Because MPT51 induces only CD4+ or CD8+ T-cell response in C57BL/6 or BALB/c, respectively (Suzuki *et al.*, 2004), we used both strains to investigate their corresponding T-cell responses. C57BL/6 and BALB/c mice were immunized with a fusion gene of HSP70F and MPT51 (HSP70F-MPT51) or MPT51 using the Helios gene gun system. Spleen cells from immunized C57BL/6 and BALB/c mice were used for IFN-γ production assays of CD4+ and CD8+ T cells, respectively, in response to corresponding antigenic peptides. CD4+ T cells from C57BL/6 mice immunized with HSP70F-MPT51 produced a significantly higher amount of IFN-γ than those immunized with MPT51 (Fig. 1). The addition of HSP70F to MPT51 did not show a significant enhancing effect on the antigen-specific CD8+ response, although CD8+ T cells from BALB/c mice immunized with HSP70F-MPT51 showed a slightly higher IFN-γ production than those immunized with MPT51. Immunization with HSP70F DNA alone did not induce MPT51-specific immune responses in C57BL/6 and BALB/c mice, and the amounts of IFN-γ production from spleen cells of immunized mice in response to corresponding




Fig. 1. Enhancing effect of HSP70 conjugation on MPT51-specific immune response. BALB/c (for CD8+ T-cell response) and C57BL/6 (for CD4+ T-cell response) mice were immunized with plasmids that encoded *HSP70F-MPT51* or *MPT51* alone, and their spleen cells were stimulated with the corresponding antigenic peptides. The IFN- γ concentration of the culture supernatants was determined by ELISA. The results of four mice per group are presented as mean \pm SD. NS, not significant; * $P < 0.05$. IFN- γ productions from unstimulated spleen cells were nearly undetectable in all groups.

peptides were similar to those of naïve mice (data not shown). In contrast to IFN- γ production, *HSP70F-MPT51* showed no enhancing effect on MPT51-specific interleukin-4 (IL-4) production in both strains, although spleen cells from both strains similarly produced IL-4 in response to anti-CD3 mAb stimulation (data not shown). Taken together, the addition of HSP70 to MPT51 enhanced antigen-specific T-helper cell type 1 (Th1), but not Th2 responses.

Identification of the domain of HSP70 responsible for its enhancing effect on MPT51-specific T-cell response

Because HSP70 comprises two major domains, that is, the N-terminal ATPase domain (44 kDa) and the C-terminal peptide-binding domain (27 kDa), we next attempted to identify the domain that was responsible for the enhancement of the MPT51-specific T-cell response. For this purpose, we prepared additional fusion genes, *HSP70N-MPT51* and *HSP70C-MPT51* (Fig. 2a).

Before immunization, 293 T cells were transfected transiently with plasmids that encoded these chimeric MPT51 DNAs, to confirm the protein expression *in vitro*. Both fusion proteins, as well as HSP70F-MPT51 and MPT51, were detected at around the expected molecular sizes by Western blotting with anti-MPT51 mAb (Fig. 2b), indicating the proper amino acid sequences of expressed proteins.

BALB/c and C57BL/6 mice were then immunized with the plasmids containing *HSP70N-MPT51*, *HSP70C-MPT51*, or *MPT51* by gene-gun and their spleen cells were tested for MPT51-specific T-cell responses. A stronger IFN- γ production from CD4+ T cells was induced by *HSP70C-MPT51* compared with that by *HSP70N-MPT51* and *MPT51* alone (Fig. 3). However, *HSP70C-MPT51* showed no enhancing effect on MPT51-specific IL-4 production (data not shown). Although the CD4+ T-cell response induced by *HSP70N-MPT51* seemed stronger than that induced by *MPT51*, the difference was not statistically significant. In contrast, the

addition of either HSP70N or HSP70C did not have any effect on the induction of CD8+ T-cell responses, which confirmed the results with HSP70F. The enhancing effect of HSP70C was confirmed by semi-quantitative RT-PCR (Fig. 4). The expression of mRNA for IFN- γ was increased considerably, which resulted in the upregulation of mRNA for iNOS, probably in APCs. These results together indicate that the C-terminal domain of *M. tuberculosis* HSP70 contributes to the enhancement of antigen-specific Th1 responses.

Induction of MPT51-specific CD4+ T-cell responses by MPT51-HSP70 fusion proteins

We finally prepared fusion MPT51 proteins and examined their effects on the induction of MPT51-specific CD4+ T-cell responses. Although SDS-PAGE analysis showed all Ni-NTA-purified proteins to have molecular sizes slightly smaller than those algorithmically expected (Fig. 5a), they were confirmed to have proper amino acid sequences by sequencing analyses (data not shown). When mice were immunized with these recombinant proteins, HSP70F-MPT51 and HSP70C-MPT51 again induced stronger immune responses than HSP70N-MPT51 and MPT51, and the effect of HSP70C-MPT51 was superior to that of HSP70F-MPT51 (Fig. 5b). These results confirm the evidence obtained by DNA vaccination that *M. tuberculosis* HSP70, especially its C-terminal domain, facilitates the induction of antigen-specific CD4+ T-cell responses.

Discussion

In DNA vaccines, the plasmid DNA is delivered into somatic cells (such as keratinocytes or myocytes) and professional APCs (such as DCs), but the former cells serve as a predominant reservoir for antigen (Gurunathan *et al.*, 2000; Kutzler & Weiner, 2008). DNA vaccines express a low amount of antigen at the restricted site of inoculation, and the frequency of DC bearing vaccinated DNA is very low




Fig. 2. Chimeric HSP70-MPT51 molecules. (a) Schematic diagram of chimeric molecules. HSP70F, HSP70N, and HSP70C mean full-length, the N-terminal domain, and the C-terminal domain of HSP70, respectively. (b) Expression of chimeric proteins in HEK293T cells. Cells were transfected with plasmids that encoded DNA for various MPT51 molecules, and the protein expression was detected by Western blotting using anti-MPT51 mAb (derived from a clone 2B11F5). 1, MPT51 (expected molecular weight: 27 979.17); 2, HSP70F-MPT51 (95 570.65); 3, HSP70N-MPT51 (67 382.45); and 4, HSP70C-MPT51 (57 774.05). Expected molecular sizes were calculated by an algorithm termed 'COMPUTE PI/MW TOOL' in EXPASY (http://expasy.org/tools/pi_tool.html).




Fig. 3. Comparison of the enhancing effects of the N- and C-domains of HSP70 in DNA vaccination. BALB/c and C57BL/6 mice were immunized with plasmids that encoded *HSP70N-MPT51*, *HSP70C-MPT51*, or *MPT51* alone, and the effects were determined by an IFN- γ production assay using MPT51 peptides for stimulation. The results of four mice per group are presented as mean \pm SD. NS, not significant; * $P < 0.05$. IFN- γ productions from unstimulated spleen cells were nearly undetectable in all the groups.

(Casares *et al.*, 1997; Barouch *et al.*, 2004). Therefore, intercellular antigen spreading seems particularly important to increase the number of APCs. In the present study, we attempted to establish a new fusion DNA vaccine against MPT51 using *M. tuberculosis* HSP70 as a partner to facilitate the receptor-mediated uptake of shed antigens from transfected somatic cells that underwent apoptosis and/or necrosis by professional APCs. With this new vaccine construct, we demonstrated that HSP70, especially its C-terminal domain, had an enhancing effect on the induction of MPT51-specific CD4⁺ (but not CD8⁺) T-cell responses.

Microbial and mammalian HSP70s effectively induce antigen-specific immune responses in various ways (Asea *et al.*, 2000; Kuppner *et al.*, 2001; Somersan *et al.*, 2001), and several receptors have been identified for them, that is, CD91 (Basu *et al.*, 2001), CD40 (Wang *et al.*, 2001; Becker *et al.*, 2002), CCR5 (Floto *et al.*, 2006), TLR-2, and TLR-4 (Asea *et al.*, 2002). Tobian *et al.* (2004a) have reported that exogenous *M. tuberculosis* HSP70 can CD91-dependently enhance the presentation capacity for ovalbumin (OVA)-derived major histocompatibility complex (MHC) class I peptides in macrophages and DCs. This has established

CD91 as a receptor for prokaryotic as well as mammalian HSPs. On the other hand, they have also reported that the enhancing effect of *M. tuberculosis* HSP70 on the presentation of MHC class II peptides is CD91-independent (Tobian *et al.*, 2004b). HSP70 has also been reported to bind to a chemokine receptor CCR5 (Whittall *et al.*, 2006) and transduce various signals into DCs to enhance immune responses (Floto *et al.*, 2006). We reported previously that a fusion protein of MIP-1 α and MPT51 is internalized preferentially into DCs via CCR5, and consequently, induces stronger MPT51-specific CD8⁺ and CD4⁺ T-cell responses than MPT51 alone (Uchijima *et al.*, 2008 and unpublished data). It is possible that *M. tuberculosis* HSP70 exerts its enhancing effect in the same way, although further analysis is necessary. Alternatively, additional signals via other HSP70 receptors, such as CD40 and/or TLRs, may contribute to the enhancing effects. Taken together, these observations suggest that activation signals transduced via HSP70 receptors other than CD91 play pivotal roles in the enhancement of immune responses, although CD91 may contribute to the internalization of MPT51 fusion proteins conjugated with HSP70.

The domain of HSP70 that is responsible for its enhancing effect is still a controversial subject. Some groups have reported that the C-terminal peptide-binding domain might act as a microbial adjuvant (Wang *et al.*, 2002; Qazi *et al.*, 2005). Wang and colleagues showed that the C-, but not the N-terminal domain upregulates the expression of cytokines and their receptors (IL-12, tumor necrosis factor- α , NO, C-C chemokines, and CCR7), costimulatory molecules (CD83, CD86, and CD80), and HLA class II molecules in human hematopoietic cells. Of these upregulated molecules, IL-12 is known to be a very potent cytokine for Th1 polarization (Trinchieri, 1994), but they showed Th1 skewing of immune responses only by an isotype analysis of immunoglobulins specific for HSP70 itself. Qazi and colleagues also described the effectiveness of the C-terminal domain, but they did not compare its effect with that of the N-terminal domain. Together with these observations, our current findings showing the enhancing effect on targeted-antigen-specific IFN- γ production by CD4+ T cells




Fig. 4. Semi-quantitative RT-PCR for iNOS and IFN- γ . Spleen cells from immunized C57BL/6 mice were stimulated with MPT51 peptide (P171–190), and the expression of iNOS and IFN- γ was tested by semi-quantitative RT-PCR. G3PDH was used as a control.

strengthen the rationale for the use of the C-terminal domain of HSP70 as a Th1-polarizing adjuvant. The enhancing effect of C-terminal domain of HSP70 is more prominent when used in protein immunization. As shown in Fig. 5b, the addition of HSP70C (and HSP70F) to MPT51 effectively induced antigen-specific immune responses, even if the antigen alone could not induce the response. In contrast, Udono *et al.* (2001) have identified amino acid residues 280–385 in the N-terminal ATPase domain of HSP70 as the most important region for cytotoxic T lymphocyte (CTL) induction. Huang *et al.* (2000) have also shown that the N-terminal domain of HSP70 (residues 160–370) is sufficient to stimulate the substantial generation of anti-OVA CTL in the absence of an adjuvant. At present, we have no evidence to explain why HSP70N did not show an enhancing effect on CD8+ T-cell responses in our system. Our construct contains the entire structure of the N-terminal domain while other effective constructs lack the extreme N-terminus, and this portion might have an inhibitory effect. Alternatively, the difference(s) in the immunogenicity of antigens (OVA vs. MPT51) and/or strain (C57BL/6 vs. BALB/c) may be accountable for the ineffectiveness. Besides the enhancing effect of the N-terminal domain on immune responses, however, its immunosuppressive role by the production of IL-10 and transforming growth factor- β has also been demonstrated in rats (Kimura *et al.*, 1998; Wendling *et al.*, 2000). In general, the C-terminal peptide-binding domain of HSP70 tends to facilitate CD4+ T-cell responses more effectively than CD8+ T-cell responses, which skews the cytokine milieu toward Th1.

IFN- γ -secreting CD4+ T cells known as Th1 cells are important mediators of tuberculosis protection (Cooper *et al.*, 1993; Flynn *et al.*, 1993), and attempts to induce tuberculosis antigen-specific Th1 cells have been the




Fig. 5. Comparison of the enhancing effects of the N- and C-domains of HSP70 in protein vaccination. (a) SDS-PAGE analysis of recombinant proteins. Proteins were purified with Ni-NTA agarose, subjected to SDS-PAGE, and stained with Coomassie Brilliant Blue R-250. For the expected molecular sizes, see the legend for Fig. 2b. (b) MPT51-specific IFN- γ production. C57BL/6 mice were immunized with recombinant proteins, and their spleen cells were used for an IFN- γ production assay in response to the MPT51 peptide (P171–190). The results of four mice per group are presented as mean \pm SD. * $P < 0.05$. IFN- γ productions from unstimulated spleen cells were nearly undetectable in all the groups.

dominant theme of most tuberculosis-vaccine development (Skeiky & Sadoff, 2006). Although CD8⁺ CTLs have also been reported to contribute to disease resistance (Flynn & Chan, 2001; Kaufmann, 2003), our current findings may pave the way for the establishment of a novel vaccine against *M. tuberculosis*.

Acknowledgements

We thank Ms. Y. Suzuki for her expert assistance. This work was supported in part by Grants-in-Aid for Scientific Research from the Japan Society for the Promotion of Science, the COE Research and the Scientific Research on Priority Areas from the Ministry of Education, Culture, Sports, Science and Technology of Japan, the Health and Labour Science Research Grants for Research into Emerging and Reemerging Infectious Diseases from the Ministry of Health, Labour and Welfare of Japan, and the United States–Japan Cooperative Medical Science Committee.

References

- Andersen P (2007) Vaccine strategies against latent tuberculosis infection. *Trends Microbiol* **15**: 7–13.
- Asea A, Kraeft SK, Kurt-Jones EA, Stevenson MA, Chen LB, Finberg RW, Koo GC & Calderwood SK (2000) HSP70 stimulates cytokine production through a CD14-dependant pathway, demonstrating its dual role as a chaperone and cytokine. *Nat Med* **6**: 435–442.
- Asea A, Rehli M, Kabingu E, Boch JA, Bare O, Auron PE, Stevenson MA & Calderwood SK (2002) Novel signal transduction pathway utilized by extracellular HSP70: role of toll-like receptor (TLR) 2 and TLR4. *J Biol Chem* **277**: 15028–15034.
- Barouch DH, Truitt DM & Letvin NL (2004) Expression kinetics of the interleukin-2/immunoglobulin (IL-2/Ig) plasmid cytokine adjuvant. *Vaccine* **22**: 3092–3097.
- Basu S, Binder RJ, Suto R, Anderson KM & Srivastava PK (2000) Necrotic but not apoptotic cell death releases heat shock proteins, which deliver a partial maturation signal to dendritic cells and activate the NF- κ B pathway. *Int Immunol* **12**: 1539–1546.
- Basu S, Binder RJ, Ramalingam T & Srivastava PK (2001) CD91 is a common receptor for heat shock proteins gp96, hsp90, hsp70, and calreticulin. *Immunity* **14**: 303–313.
- Becker T, Hartl FU & Wieland F (2002) CD40, an extracellular receptor for binding and uptake of Hsp70-peptide complexes. *J Cell Biol* **158**: 1277–1285.
- Binder RJ, Anderson KM, Basu S & Srivastava PK (2000) Cutting edge: heat shock protein gp96 induces maturation and migration of CD11c⁺ cells *in vivo*. *J Immunol* **165**: 6029–6035.
- Casares S, Inaba K, Brumeanu TD, Steinman RM & Bona CA (1997) Antigen presentation by dendritic cells after immunization with DNA encoding a major histocompatibility complex class II-restricted viral epitope. *J Exp Med* **186**: 1481–1486.
- Castellino F, Boucher PE, Eichelberg K, Mayhew M, Rothman JE, Houghton AN & Germain RN (2000) Receptor-mediated uptake of antigen/heat shock protein complexes results in major histocompatibility complex class I antigen presentation via two distinct processing pathways. *J Exp Med* **191**: 1957–1964.
- Colditz GA, Brewer TF, Berkey CS, Wilson ME, Burdick E, Fineberg HV & Mosteller F (1994) Efficacy of BCG vaccine in the prevention of tuberculosis. Meta-analysis of the published literature. *JAMA* **271**: 698–702.
- Cooper AM, Dalton DK, Stewart TA, Griffin JP, Russell DG & Orme IM (1993) Disseminated tuberculosis in interferon gamma gene-disrupted mice. *J Exp Med* **178**: 2243–2247.
- de Araujo-Filho JA, Vasconcelos AC Jr, Martins de Sousa E, Kipnis A, Ribeiro E & Junqueira-Kipnis AP (2008) Cellular responses to MPT-51, GlcB and ESAT-6 among MDR-TB and active tuberculosis patients in Brazil. *Tuberculosis (Edinburgh)* **88**: 474–481.
- Floto RA, MacAry PA, Boname JM *et al.* (2006) Dendritic cell stimulation by mycobacterial Hsp70 is mediated through CCR5. *Science* **314**: 454–458.
- Flynn JL & Chan J (2001) Immunology of tuberculosis. *Annu Rev Immunol* **19**: 93–129.
- Flynn JL, Chan J, Triebold KJ, Dalton DK, Stewart TA & Bloom BR (1993) An essential role for interferon γ in resistance to *Mycobacterium tuberculosis* infection. *J Exp Med* **178**: 2249–2254.
- Gething MJ, Blond-Elguindi S, Buchner J, Fourie A, Knarr G, Modrow S, Nanu L, Segal M & Sambrook J (1995) Binding sites for Hsp70 molecular chaperones in natural proteins. *Cold Spring Harb Symp* **60**: 417–428.
- Gurunathan S, Klinman DM & Seder RA (2000) DNA vaccines: immunology, application, and optimization. *Annu Rev Immunol* **18**: 927–974.
- Harlow E & Lane D (1988) *Monoclonal Antibodies. Antibodies*. Cold Spring Harbor Laboratory Press, New York. pp. 139–243.
- Huang Q, Richmond JF, Suzue K, Eisen HN & Young RA (2000) *In vivo* cytotoxic T lymphocyte elicitation by mycobacterial heat shock protein 70 fusion proteins maps to a discrete domain and is CD4⁺ T cell independent. *J Exp Med* **191**: 403–408.
- Kaufmann SHE (2000) Is the development of a new tuberculosis vaccine possible? *Nat Med* **6**: 955–960.
- Kaufmann SHE (2003) Immunity to intracellular bacteria. *Fundamental Immunology*, 5th edn (Paul WE, ed), pp. 1229–1261. Lippincott Williams & Wilkins Publishers, Philadelphia, PA.
- Kaufmann SHE (2004) New issues in tuberculosis. *Ann Rheum Dis* **63** (suppl 2): ii50–ii56.
- Kimura Y, Yamada K, Sakai T, Mishima K, Nishimura H, Matsumoto Y, Singh M & Yoshikai Y (1998) The regulatory role of heat shock protein 70-reactive CD4⁺ T cells during rat listeriosis. *Int Immunol* **10**: 117–130.