



# Behavioral interventions to reduce the transmission of HIV infection among sex workers...


## 1.4 Mean number of condom use by FSWs at past month



## 1.5 Sexual activity outcomes by FSWs at past month




## 1.6 Mean number of protected sex among FSWs at 6-month




# Behavioral interventions to reduce the transmission of HIV infection among sex workers...


## 1.7 Injected drug use among FSWs at 6-month



## 1.8 Drug use among FSWs at 6-month




## 1.9 Alcohol use among FSWs at 6-month




## 2 Social cognitive theory versus no intervention for promotion of condom use

### 2.1 Consistent condom use by FSWs




### 2.2 HIV knowledge among FSWs at 6-month




# Behavioral interventions to reduce the transmission of HIV infection among sex workers...

## 2.3 Psychosocial barriers to condom use among FSWs at 6-month




## 3 Community empowerment versus standard care for promotion of condom use


### 3.1 STIs prevalence among FSWs at 36-month



### 3.2 Consistent condom use by FSWs with clients




### 3.3 Consistent condom use by FSWs with regular partners




# Behavioral interventions to reduce the transmission of HIV infection among sex workers...


## 3.4 Change in 100% condom use by FSWs




## 3.5 Change in any condom use by FSWs



## 3.6 HIV knowledge among FSWs




## 3.7 HIV testing by FSWs at 36-month




## Behavioral interventions to reduce the transmission of HIV infection among sex workers...


### 3.8 Skills of risk and protective factors among FSWs at 16-month




### 3.9 Type of sexual practices by FSWs at baseline




### 3.10 Drug use by FSWs at baseline



### 3.11 Alcohol use by FSWs at baseline




### 3.12 Verbally abused at 3-month




## 4 Microenterprise plus education intervention versus education alone for reducing the number of sex partners

# Behavioral interventions to reduce the transmission of HIV infection among sex workers...

## 6.1 STIs prevalence among FSWs




## 6.2 Male condom use with clients




# Behavioral interventions to reduce the transmission of HIV infection among sex workers...


## 6.3 Male condom use with non-paying partners




## 6.4 Female condom use by FSWs with clients



## 6.5 Male and female condom use with clients




## 6.6 Male and female condom use in the last sex with non-paying partner




## 7 Peer education plus manager training versus standard care for promoting HIV testing and condom use

# Behavioral interventions to reduce the transmission of HIV infection among sex workers...

## 4.1 Consistent condom use by FSWs at 6-month




## 4.2 Number of FSWs' partners at 6-month




## 5 Peer education versus standard care for promoting HIV testing and condom use


### 5.1 Mean number of consistent condom use by FSWs at 6-month



### 5.2 Mean score of HIV knowledge among FSWs at 6-month




### 5.3 HIV testing by FSWs at 6-month




## Behavioral interventions to reduce the transmission of HIV infection among sex workers...

### 5.4 AIDS perceived control by FSWs at 6-month




### 5.5 AIDS perceived severity by FSWs at 6-month




### 6 Peer education plus clinic based counseling versus peer education only for promotion of condom use

## Behavioral interventions to reduce the transmission of HIV infection among sex workers...

### 6.1 STIs prevalence among FSWs




### 6.2 Male condom use with clients




## Behavioral interventions to reduce the transmission of HIV infection among sex workers...


### 6.3 Male condom use with non-paying partners




### 6.4 Female condom use by FSWs with clients



### 6.5 Male and female condom use with clients




### 6.6 Male and female condom use in the last sex with non-paying partner




### 7 Peer education plus manager training versus standard care for promoting HIV testing and condom use

# Behavioral interventions to reduce the transmission of HIV infection among sex workers...


## 7.1 Mean number of consistent condom use by FSWs at 6-month




## 7.2 Mean score of HIV knowledge among FSWs at 6-month




## 7.3 HIV testing by FSWs at 6-month



## 7.4 AIDS perceived control by FSWs at 6-month




## 7.5 AIDS perceived severity by FSWs at 6-month




## 8 Manager training versus standard care for promoting HIV testing and condom use

# Behavioral interventions to reduce the transmission of HIV infection among sex workers...


## 8.1 Mean number of consistent condom use by FSWs at 6-month




## 8.2 Mean score of HIV knowledge among FSWs at 6-month




## 8.3 HIV testing by FSWs at 6-month



## 8.4 AIDS perceived control by FSWs at 6-month




## 8.5 Mean score of perceived themselves at greater risk (by FSWs at 6-month)




## 9 Promotion of female and male condom versus promotion of male condom

# Behavioral interventions to reduce the transmission of HIV infection among sex workers...


## 9.1 HIV incidence among FSWs at 3-month




## 9.2 STIs incidence among FSWs at 3-month




## 9.3 Consistent male condom use at 3-months



## 9.4 Consistent female condom use by FSWs at 3-month




## 9.5 Consistent female condom use by FSWs at 24-month




# Behavioral interventions to reduce the transmission of HIV infection among sex workers...

## 9.6 Consistent male condom use at 24-month




## 10 Promotion of condom plus health education versus promotion condom alone


### 10.1 Proportion condom use by FSWs and clients due to receive health education at 2-month



### 10.2 Condom provision in motels mainly used for commercial sex at 2-month




### 10.3 Condom provision in motels used mainly for non-commercial sex at 2-month




## 11 Intensive STI screening versus basic STI screening to control STI

# Behavioral interventions to reduce the transmission of HIV infection among sex workers...

## 11.1 STIs prevalence among FSWs at 6-month




## 11.2 Consistent condom use by FSWs at 6-month




## 12 VCT versus standard care of STI for increasing condom use

### 12.1 STIs prevalence among FSWs at 6-month



### 12.2 Consistent condom use by FSWs with clients at 6-month



# Behavioral interventions to reduce the transmission of HIV infection among sex workers...

## 12.3 HIV/STI knowledge among FSWs at 6-month

