

Table 3-1. Representative miRNAs That Were Commonly Repressed In CH-B, CH-C, HCC-B, and HCC-C Compared with Normal Liver (Cluster)

| miRNA | Parametric P Value | Ratio* | No. of Significant Genes/Predicted Target Genes† | Hottelling Test P Value‡ | Differentially Expressed Target Genes§ | Pathway of Regulated Genes¶ |
|-------------|--------------------|--------|--|--------------------------|--|---|
| hsa-miR-219 | 7.3E-05 | 0.28 | 25/109 | 2.59E-04 | Glypican-3, ERP5, PLK2, HIRA, HMG2 ACOX1 NF-X1 | Regulatory T cell differentiation Fatty acid beta-oxidation MHC class II biosynthetic process |
| hsa-miR-320 | 9.8E-05 | 0.50 | 26/88 | 3.50E-06 | Vimentin, ALP (<i>N</i> -acetyltransferase-like), SEC61 beta, G-protein alpha-i2, Filamin A | Protein kinase cascade |
| hsa-miR-154 | 2.7E-04 | 0.15 | 22/70 | 5.40E-06 | Rac1, RhoG Vinexin beta, Profilin I, Ca-ATPase3 OTR, NET1(TSPAN1), NAP1, Vimentin, PDIA3, cytochrome P-450 reductase DLX2 GUAC, ACAT1 | Organelle organization and biogenesis Actin cytoskeleton organization and biogenesis Regulation of apoptosis Morphogenesis Branched chain family amino acid catabolic process |
| hsa-miR-29c | 1.8E-03 | 0.55 | 53/133 | 1.00E-06 | FBX07, ASPP1, HSPA4, Cathepsin O, PDF, COL4A1, HSPA4, TIP30, CXADR NS1-BP, ALP (<i>N</i> -acetyltransferase-like), ACTR10, Beclin 1 SMAD6, LTBR(TNFRSF3), ENPP7 | Cell-substrate adhesion Transcription, DNA-dependent Apoptosis |
| hsa-miR-338 | 5.2E-03 | 0.46 | 30/101 | 3.60E-06 | ID3, GATA-4, NFIA, FR-beta, CREST, HYOU1 G3ST1, CAD, FKBP12, LZIP, PDIA3, Schwannomin (NF2), CREST | Developmental process Immune effector process Immune system process |
| hsa-miR-26a | 6.3E-03 | 0.70 | 37/119 | 2.64E-05 | LIG4, c-FLIP, GADD45 beta, DAPK1, PRDX4, LRP130 Cyclin E, ZDHHC6, Tx1, ATG8 (GATE-16), WASP, C1s COPG1 | Response to stimulus DNA replication initiation Ion transport |
| hsa-miR-126 | 8.1E-03 | 0.65 | 27/101 | 4.04E-03 | ANP32B (april), HSPA4, RLI, LIV-1 (SLC39A6), PTP-MEG2, CD97, DHPR NFKBIA, NMI, MDH1, PDCD2 SMAD6, ATP6AP2, ANP32B (april), NMI, HSPA4 | Regulation of cellular protein metabolic process Response to stress Apoptosis |
| hsa-miR-325 | 8.7E-03 | 0.20 | 18/63 | 2.03E-04 | TRADD, CREST, NEDD8, annexin IV, GPX2, PDF, TNFAIP1 Glypican-3, ID1, PC-TP, SNRNP (Sm-B) | Developmental process Multicellular organismal development RNA splicing |

*Ratio of HCC-B, HCC-C, CH-B, and CH-C to normal.

†The number of significant genes ($P < 0.05$) out of predicted target genes in which expression was evaluated in microarray.

‡Statistical assessment of presence of differentially expressed genes out of predicted target genes of miRNAs.

§Representative differentially expressed genes out of predicted target genes of miRNAs.

¶Representative pathway of differentially expressed genes out of predicted target genes of miRNAs.

the hypergeometrical distribution based on gene ontology terms. Because one gene is frequently involved in multiple pathways, all pathways corresponding to the genes with significance probability were listed.

Verification of Regulation of Candidate Target Genes by miRNAs. Anti-miRNA (Ambion) specific to 13 miRNAs (has-miR-17*, has-miR-20a, has-miR-23a, has-miR-26a, has-miR-27a, has-miR-29c, has-miR-30a, has-miR-92, has-miR-126, has-miR-139, has-miR-187, has-miR-200a, and has-miR-223) showing significant

differences in expression were transfected into Huh7 cells using TransMessenger transfection reagent (QIAGEN, Valencia, CA), and loss of function of each miRNA was evaluated. Similarly, precursor miRNAs of five miRNAs (has-miR-23a, has-miR-26a, has-miR-27a, has-miR-92, and has-miR-200a) were also transfected into Huh7 cells, and gain of function of each miRNA was evaluated. The loss- and gain-of-function of miRNAs were evaluated via RTD-PCR. In addition, different gene expressions regulated by miRNAs were also evaluated via RTD-PCR.

HBV/HCV Infection Model Using Cultured Cells.

The plasmid pHBV 1.2 coding the 1.2-fold length of the HBV genome was transfected into Huh7.5 cells using Fugene6 transfection reagent (Roche Applied Science, Indianapolis, IN). HBeAg production in culture medium was measured using Immunis HBeAg/Ab EIA (Institute of Immunology Co., Ltd., Tokyo, Japan).¹³ The amount of HBV-DNA was measured via RTD-PCR (Supplementary Fig. 1A,B). JFH1-RNA was transfected into Huh7.5 cells using TransMessenger transfection reagent (QIAGEN) and the expression of the core protein was examined via immunofluorescence staining using anti-HCV core antibody (Affinity BioReagent, CO).^{14,15} HCV-RNA amount was also measured via RTD-PCR (Supplementary Fig. 1A,B). JFH1/GND was used as a negative control. miRNA expression was quantitated by RTD-PCR 48 hours after transfection.

Results

Expression of miRNA in Liver Tissue. A panel of miRNA was successfully amplified from liver tissues via RTD-PCR. The representative amplification profile of miRNA as determined with RTD-PCR is shown in Fig. 1. To assess the reliability and reproducibility of this assay system, we first measured RNU6B in duplicate from all samples in different plates. The mean difference in Ct values of RNU6B expression within the same samples was 0.08 ± 0.05 (mean \pm standard deviation), indicating the high reproducibility of this assay. All Ct values from each reaction were collected, and Ct variation obtained by each probe from all patients was calculated. Although RNU6B was frequently used as the internal control, the standard Ct variation was relatively high (Ct, 27 ± 1.94), suggesting that the variances in its value depend on the state of liver disease (N, CH and HCC). Therefore, we selected has-miR-328 as the internal control with the smallest standard deviation (Ct, 30 ± 0.60). The relative expression ratio of individual miRNA to has-miR-328 was calculated and applied to the following analysis using a BRB-array tool.

Hierarchical cluster analysis revealed that the expression profiles of the 188 miRNAs from each patient were roughly classified into normal liver, HBV-infected liver (CH-B+HCC-B; HBV group), and HCV-infected liver (CH-C+HCC-C; HCV group) (Fig. 2A). HCV viremia in two patients with CH-C was persistently cleared by interferon therapy before HCC development. The background liver of one of these patients was clustered in the normal group and those of others in the HCV group. Although these two patients were not clearly differentiated from others, some miRNAs such as miR-194, miR-

211, and miR-340 that were down-regulated in the HCV group were significantly up-regulated in two patients (Fig. 3, cluster 2).

The present CH and HCC expression data were obtained from the same patient; however, each sample clustered irrespective of pairs in all but two patients. miRNA expression profiling was therefore more dependent on the disease condition than on the paired condition, as also confirmed by the Dunnett test.¹² We then attempted to classify the expression profiles into HBV and HCV groups using supervised learning methods (Table 2-1). HBV and HCV groups were significantly differentiated at an 87% accuracy ($P < 0.001$). The normal liver and CH (CH-B + CH-C) and CH and HCC (HCC-B + HCC-C) were also significantly differentiated at a 90% rate of accuracy. These results suggest that different stages of liver disease (normal, CH, and HCC) can be differentiated from each other based on the miRNA expression profile, as well as HBV and HCV infection.

To examine the relationship among five categories of groups, namely, N, CH-B, CH-C, HCC-B and HCC-C, we attempted to differentiate the five groups using a supervised learning algorithm (binary tree classification) used for classifying three or more groups. SVM was used as a prediction method. Expression profiles were first classified into groups N (normal) and non-N (non-normal) (CH-C, CH-B, HCC-C, and HCC-B) (node 1) ($P < 0.01$). The non-N group was then classified into HBV and HCV (node 2) ($P < 0.01$). The HBV group was further classified into CH-B and HCC-B (node 3) ($P < 0.01$), and the HCV group was further classified into CH-C and HCC-C (node 4) ($P < 0.01$) (Fig. 2B, Table 2-2). Thus, the findings support the notion that differences in miRNA expression between HBV and HCV are as distinct as those between CH and HCC.

Out of 20 miRNAs that differentiated node 1 classification (Table 2-2), 12 also differentiated node 3 or node 4 classification. The remaining eight miRNAs specifically differentiated node 1 classification. They were down-regulated in the HBV and HCV groups compared with the normal group (Fig. 3, cluster 1). Nineteen miRNAs differentiated node 2 classification (Table 2-2) and the hierarchical clustering using these miRNAs clearly differentiated the HBV and HCV groups (Fig. 3, cluster 2). There were 15 and 14 miRNAs that differentiated node 3 and 4 classifications, respectively (Table 2-2). Hierarchical clustering using these miRNAs revealed that these miRNAs differentiated CH-B and HCC-B as well as CH-C and HCC-C, respectively; 17 miRNAs were down-regulated in HCC, and six were up-regulated in HCC (Fig. 3, cluster 3).

Table 3-2. Differentially Expressed miRNA Between HCC-B, CH-B, and HCC-C, CH-C, and Their Representative Target Genes (Cluster 2)

| miRNA | Parametric P Value | Ratio* | No. of Significant Genes/Predicted Target Genes† | Hotelling Test P Value‡ | Differentially Expressed Target Genes§ | Pathway of Regulated Genes¶ |
|----------------|--------------------|--------|--|-------------------------|---|--|
| hsa-miR-190 | 1.2E-05 | 2.06 | 21/68 | 4.47E-02 | Chk1, C2orf25, VRK2, USP16, STAF65(gamma) AP1S2, RNASE4 | Regulation of cell cycle Mitotic cell cycle |
| hsa-miR-134 | 2.3E-04 | 5.74 | 11/58 | 3.40E-06 | PPP2R1B, ARHGAP15, UBPY VKDGC, SH2B, MALS-1, DDB2 BCRP1 DDB2 | Negative regulation of apoptosis Multicellular organismal process Regulation of viral reproduction Lipid biosynthetic process |
| hsa-miR-151 | 2.8E-04 | 1.82 | 12/62 | 6.41E-01 | RGS2, UFO, AK2, USP7 eIF4G2, USP7 SLC22A7 | G-protein signaling Regulation of translation Organic anion transport |
| hsa-miR-193 | 5.0E-04 | 1.67 | 23/95 | 9.30E-01 | G-protein alpha-11, p130CAS, VAV-1, PDCD11 Colipase, ACSA DCOR | Cell motility Energy coupled proton transport Intracellular signaling cascade Proteasomal protein catabolic process |
| hsa-miR-133b | 1.7E-03 | 2.42 | 20/97 | 3.69E-02 | DDB2, Bcl-3, Cystatin B Rab-3, RAG1AP1, KCNH2, DCOR AL1B1 | Regulation of biological quality Carbohydrate metabolic process |
| hsa-miR-324-5p | 2.9E-03 | 1.51 | 27/121 | 1.90E-06 | SKAP55, VAV-1, DDB2, E2A, NIP1 MEMO (CGI-27), Rab-3 COPG1, GPX3, OAZ2 | Cellular developmental process Cellular structure morphogenesis Glutathione metabolic process |
| hsa-miR-182* | 3.1E-03 | 2.23 | 28/123 | < 1e-07 | Alpha-endosulfine, HCCR-2, Thioredoxin-like 2, TPT1, USP7 DDB2, TPT1 JIP-1 | Translation initiation in response to stress Cellular developmental process JNK cascade |
| hsa-miR-105 | 4.6E-03 | 4.38 | 18/68 | 4.74E-05 | Beta-2-microglobulin, HLA-B27 PIMT, IL-17RC MHC class I, CDK9, ERG1, Desmocollin 3 PSMD5, SLC26A6 | Antigen processing and presentation Immune response Proteasomal protein catabolic process |
| hsa-miR-211 | 5.3E-03 | 25.61 | 10/56 | 2.00E-04 | Noelin, SC4MOL, Thioredoxin-like 2, CCL5, NALP3 Hic-5/ARA55, USP16, MAP4, Ferroportin 1 | Regulation of apoptosis Positive regulation of cellular process |
| hsa-miR-191 | 6.7E-03 | 1.39 | 25/79 | 7.55E-04 | TOP3A, PLRP1 CDK9, GPS2, CLTA, LXR-alpha ACSA UGCG1, SGPP1 | Oxygen transport Nucleic acid metabolic process Acetyl-CoA biosynthetic process Metal ion transport |
| hsa-miR-340 | 8.5E-03 | 1.48 | 17/81 | 3.73E-03 | FKBP12, DCOR, Gelsolin, VAV-1, ARF6 HXK3 | Calcium ion transport Actin cytoskeleton organization and biogenesis Glucose catabolic process |
| hsa-miR-194 | 8.7E-03 | 1.67 | 13/74 | 5.90E-01 | Cyclin B1, Serglycin PTE2 SLC7A6 | M phase of mitotic cell cycle Acyl-CoA metabolic process Carbohydrate utilization |
| hsa-miR-23a | 1.9E-04 | 0.46 | 14/97 | < 1e-07 | RGL2, MANR, MEK1 (MAP2K1), Caspase-3, AZGP1 FRK, Pyk2(FAK2), CSE1L AZGP1 | Protein kinase cascade Cellular developmental process Defense response |
| hsa-miR-142-5p | 4.9E-04 | 0.40 | 25/89 | 9.10E-06 | Sirtuin4, PAI2, PSAT, RIL, CDC34, SPRY1 E4BP4, DNAJC12, WWP1, PAIP1, PASK, rBAT VCAM1, CaMK I, WWP1, FHL3 | Metabotropic glutamate receptor Regulation of gene expression Cell-matrix adhesion |
| hsa-miR-34c | 5.1E-04 | 0.20 | 31/129 | 7.30E-06 | Diacylglycerol kinase, zeta, PLC-delta 1, ATP2C1, PAI2 MLK3(MAP3K11), MEK1(MAP2K1), CDC25C, MRF-1, XPC GNT-IV | Manganese ion transport Protein kinase cascade Inflammatory cell apoptosis |

Table 3-2. Continued

| miRNA | Parametric P Value | Ratio* | No. of Significant Genes/Predicted Target Genes† | Hotelling Test P Value‡ | Differentially Expressed Target Genes§ | Pathway of Regulated Genes¶ |
|--------------|--------------------|--------|--|-------------------------|---|---|
| hsa-miR-124b | 8.6E-04 | 0.32 | 25/120 | 7.10E-05 | E2F5, Rad51, Jagged1 MLK3(MAP3K11), RGS1 COL16A1 | Muscle development Intracellular signaling cascade MAPKKK cascade |
| hsa-let-7a | 1.0E-03 | 0.45 | 28/136 | 9.35E-04 | RAD51C, CoAA, hASH1, Cockayne syndrome B, Caspase-1, PP5 PLC-delta 1, MANR, ACADVL | Response to DNA damage stimulus Fibroblast proliferation |
| hsa-miR-27a | 3.9E-03 | 0.59 | 18/108 | 1.19E-02 | HGF, NGF COL16A1, RIL, RhoGDI gamma, ANP32B (april) VE-cadherin, NTH1, GATA-2, E4BP4 RAD51C | Cellular developmental process Cytoskeleton organization and biogenesis Response to external stimulus DNA recombination |

*Ratio of HCC-B, CH-B, to HCC-C,CH-C.

†The number of significant genes ($p < 0.05$) out of predicted target genes in which expression was evaluated in microarray.

‡Statistical assesment of presence of differentially expressed genes out of predicted target genes of miRNAs.

§Representative differentially expressed genes out of predicted target genes of miRNAs.

¶Representative pathway of differentially expressed genes out of predicted target genes of miRNAs.

These results indicate that there were two types of miRNAs—one associated with HBV and HCV infection (cluster 2), the other associated with the stages of liver disease (clusters 1 and 2) that were irrelevant to the differences in HBV and HCV infection.

Differential miRNAs and Their Candidate Target Genes and Signaling Pathways. Differentially expressed miRNAs are shown in Table 3. In addition to the expression ratios of miRNAs in each group, the number of genes analyzed on the microarray predicted to be the target genes of miRNAs and that which actually showed significant ($P < 0.05$) differences in expression are also shown. Based on the frequencies and levels of expression of differential genes, the significance of regulation of these gene groups by miRNAs was evaluated using Hotelling T2 test (BRB ArrayTools) (Table 3). The representative candidate target genes and their signaling pathways by each miRNA were shown one by one (Table 3). The signaling pathways regulated by all differential miRNAs in each category of groups are shown in Table 4.

Eight miRNAs were down-regulated in the HBV and HCV groups compared with the normal group (Table 3-1; Fig. 3, cluster 1). These miRNAs were associated with an increased expression of genes related to cell adhesion, cell cycle, protein folding, and apoptosis (Tables 3-1, 4-1), and possibly with the common feature of CH irrespective of the differences in HBV and HCV infection.

Nineteen miRNAs clearly differentiated the HBV and HCV groups (Fig. 3, cluster 2, Table 3-2). Thirteen miRNAs exhibited a decreased expression in the HCV group, and six showed a decreased expression in the HBV group. miRNAs exhibiting a decreased expression in the HCV group regulate genes related to immune response,

antigen presentation, cell cycle, proteasome, and lipid metabolism. On the other hand, those exhibiting a decreased expression in the HBV group regulate genes related to cell death, DNA damage and recombination, and transcription signals. These findings reflected the differences in the gene expression profile between CH-B and CH-C described (Tables 3-2, 4-2).¹⁰ Interestingly, although these miRNAs were HBV and HCV infection-specific, some of them were reported to be tumor-associated miRNAs, suggesting the possible involvement of infection-associated miRNAs in HCC development.

Twenty-three miRNAs clearly differentiated CH and HCC that were irrelevant to the differences in HBV and HCV infection. Seventeen miRNAs were down-regulated in HCC that up-regulated cancer-associated pathways such as cell cycle, adhesion, proteolysis, transcription, translation, and the Wnt signaling pathway (Tables 3-3, 4-3). Six miRNAs were up-regulated in HCC that down-regulated all inflammation-mediated signaling pathways, potentially reflecting impaired antitumor immune response.

Relationship Between Expressions of Infection-Associated miRNA in Liver and Cultured Cells Infected with HBV and HCV. To clarify whether the expression of infection-associated miRNA is regulated by HBV and HCV infection, we investigated the relationship between changes in miRNA in liver tissues and those in miRNA in Huh7.5 cells in which infectious HBV or HCV clones replicated. To evaluate the replication of each clones in Huh7.5 cells, we measured time-course changes in the amounts of HBV-DNA and HCV-RNA in Huh7.5 cells transfected with pHBV1.2 and JFH1-RNA, respectively, by RTD-PCR (Supplementary Fig. 1A). The expression of HBV proteins was examined by measuring the amount

Table 3-3. Differentially Expressed miRNA Between CH and HCC and Their Representative Target Genes (Cluster 3)

| miRNA | Parametric p-value | Ratio* | No. of Significant Genes/Predicted Target Genes† | Hottelling Test P Value‡ | Differentially Expressed Target Genes§ | Pathway of Regulated Genes¶ |
|----------------|--------------------|--------|--|--------------------------|---|---|
| hsa-miR-139 | 4.50E-06 | 0.42 | 19/106 | 2.70E-03 | Cyclin B1, DHX15, MCM5, Histone H2A RBCK1, SYHH | Mitotic cell cycle Protein catabolic process |
| hsa-miR-30a-3p | 2.50E-05 | 0.49 | 26/144 | 1.73E-02 | ILK, IGFBP7, SAFB, CTR9 GGH, Pinn, ZNF207, Annexin VII | Response to external stimulus Regulation of oxidoreductase activity |
| hsa-miR-130a | 7.00E-05 | 0.50 | 22/108 | 1.07E-02 | ILK, LTA4H, ABC50, GNPAT DLC1 | Cell-matrix adhesion Morphogenesis |
| hsa-miR-223 | 3.40E-04 | 0.39 | 14/90 | 6.52E-03 | SPHM, PPP2R5D, RHEB2, SPHM MLK3(MAP3K11), Otubain1, TIMP4 NRBP | Mitotic cell cycle Protein modification process Cell differentiation |
| hsa-miR-187 | 3.55E-04 | 0.12 | 16/66 | 6.76E-04 | Ephrin-A1, Midkine, FDPS K(+) channel, subfamily J HFE2, Otubain1 | Cell morphogenesis Notch signaling pathway Negative regulation of programmed cell death |
| hsa-miR-200a | 6.86E-04 | 0.18 | 20/141 | 2.15E-02 | PRSS11, SUPT5H, RAG1AP1 PLOD3 CDC25B, KAP3, CDK2AP2, CHKA POLD CPSF4 | Developmental process Mitochondrial ornithine transport Cell communication DNA replication RNA splicing |
| hsa-miR-17-3p | 8.42E-04 | 0.58 | 28/108 | 8.98E-04 | MLK3(MAP3K11), Tip60, ACBD6, DOC- 1R, DAX1, RBCK1 WNT5A, 14-3-3 gamma, DHX15 HFE2, MCM5 | Protein kinase cascade BMP signaling pathway DNA recombination Cytokinesis |
| hsa-miR-99a | 1.17E-03 | 0.53 | 33/163 | 9.52E-03 | Calpain small subunit, Thoredoxin-like 2, Survivin IBP2, DNA-PK, KAP3, NFE2L1, PARP-1, HDAC11 | Intracellular signaling cascade Regulatory T cell differentiation |
| hsa-miR-200b | 1.57E-03 | 0.18 | 24/147 | 2.72E-02 | HSP47, HMG2, NRBP SNX17 Ephrin-A1 | Regulation of cell cycle Cell motility Receptor protein signaling pathway |
| hsa-miR-125b | 1.82E-03 | 0.55 | 26/114 | 1.03E-01 | COL4A2, TIP30, HSP47, MSP58 MLK3(MAP3K11), ERK2 (MAPK1), ERK1 (MAPK3), PLOD3 Otubain1, SCN4A(SKM1) | Cell adhesion Nuclear translocation of MAPK Ubiquitin-dependent protein catabolic process |
| hsa-miR-30e | 2.10E-03 | 0.65 | 24/151 | 4.30E-02 | Cyclin B1, XTP3B, GAK, Annexin VII, MIC2, NRBP MSS4 S100A10 | Mitotic cell cycle Protein localization Calcium ion transport |
| hsa-miR-199a* | 4.26E-03 | 0.35 | 11/71 | 7.16E-02 | BUB3, Cyclin B1, LMN8R PRAME | Mitotic cell cycle Cardiac muscle cell differentiation |
| hsa-miR-122a | 6.31E-03 | 0.51 | 11/80 | 1.01E-03 | JAB1, APEX, Clathrin heavy chain PARN DDAH2 | Base-excision repair Translational initiation Regulation of cellular respiration |
| hsa-miR-199a | 8.77E-03 | 0.35 | 18/94 | 3.56E-02 | IL-13, MLK3(MAP3K11), CLK2, ACP33 PAFAH beta, SPA1, CLCN4 | Protein amino acid phosphorylation Small GTPase mediated signal transduction |
| hsa-miR-326 | 9.00E-03 | 0.57 | 29/147 | 2.25E-01 | Midkine, ENT1, IP3KA, PSMC5, ANCO-1 Thy-1, MCM6, Tip60, VILIP3 COMP, Cathepsin A | Regulation of programmed cell death Cell-matrix adhesion Blood vessel development |
| hsa-miR-92 | 9.60E-03 | 0.81 | 28/140 | 2.47E-02 | TUBGCP2, Fibrillin 1, PIPKI gamma, KAP3 SNX15, BCAT2 IGFBP7, FZD6, COPS6 | Rho protein signal transduction LDL receptor and BCAA metabolism Adenosine receptor signaling pathway |
| hsa-miR-221 | 3.40E-06 | 3.34 | 16/67 | 3.59E-01 | Lck, Kallistatin, Neuromodulin, LFA-3, PA24A, AZGP1, MSH2 KYNLU, PMCA3 | Immune response-activating signal transduction DNA repair |

Table 3-3. Continued

| miRNA | Parametric p-value | Ratio* | No. of Significant Genes/Predicted Target Genes† | Hotelling Test P Value‡ | Differentially Expressed Target Genes§ | Pathway of Regulated Genes¶ |
|-------------|--------------------|--------|--|-------------------------|--|--|
| hsa-miR-222 | 6.50E-06 | 2.23 | 18/85 | 1.59E-02 | Thrombospondin 1, Lck, MSH2, ATF-2, CITED2, Kallistatin | Cell motility |
| hsa-miR-301 | 5.22E-05 | 1.96 | 14/71 | 1.16E-01 | PGAR KYN Beta-2-microglobulin, PPCKM, PRC, Fra-1, PPCKM, ACAT2 BMPRI8, ARMER, EHM2, RBBP8 Neuromodulin, LDLR | Triacylglycerol metabolic process DNA replication Antigen processing and presentation |
| hsa-miR-21 | 7.67E-03 | 1.57 | 19/81 | 1.86E-04 | Btk, Fra-1, MSH2, Collectrin, Adipophilin RNASE4, AGXT2L1 SARDH | Meiotic recombination Cell motility Regulation of T cell proliferation |
| hsa-miR-183 | 2.46E-02 | 3.51 | 13/86 | 3.36E-01 | Hdj-2, PEMT, Lck, MKP-5, Chondromodulin-I, ABCA8 IL-16, MTRR, SerRS | Peptidyl-tyrosine phosphorylation Natural killer cell activation during immune response Cell differentiation |
| hsa-miR-98 | 5.22E-02 | 1.32 | 24/130 | 2.95E-04 | ACAA2, LTB4DH, ACADVL, DECR, S14 protein, Rapsyn, Kallistatin, ENPEP, Beta crystallin B1 CYP4F8 | Methionine biosynthetic process Fatty acid metabolic process Multicellular organismal process Prostaglandin metabolic process |

*Ratio of HCC to CH.

†The number of significant genes ($P < 0.05$) out of predicted target genes in which expression was evaluated in microarray.

‡Statistical assessment of presence of differentially expressed genes out of predicted target genes of miRNAs.

§Representative differentially expressed genes out of predicted target genes of miRNAs.

¶Representative pathway of differentially expressed genes out of predicted target genes of miRNAs.

of HBeAg released in culture medium (Supplementary Fig. 1B). HCV protein expression was examined by evaluating the core protein expression after 48 hours by fluorescence immunostaining (Supplementary Fig. 1C). RNA was extracted from the Huh7.5 cells 48 hours after gene transfection, and miRNA expression pattern in the cells was compared with those in liver tissues. We found a strong correlation between differences in miRNA expression between liver tissues of the HBV and HCV groups, and those in miRNA expression between Huh7.5 cells transfected with HBV and HCV clones ($r = 0.73$, $P = 0.0006$) (Fig. 5). These results revealed that differences in the expression of infection-associated miRNA in the liver between the HBV and HCV groups are explained by changes in miRNA expression caused by HBV and HCV infections.

Verification of Regulation of Candidate Target Genes by miRNA. Anti-miRNAs (Ambion) specific to 13 miRNAs (has-miR-17*, has-miR-20a, has-miR-23a, has-miR-26a, has-miR-27a, has-miR-29c, has-miR-30a, has-miR-92, has-miR-126, has-miR-139, has-miR-187, has-miR-200a, and has-miR-223) showing significant differences in expression were transfected into Huh7 cells to examine loss of function of the miRNAs. Five miRNAs (has-miR-23a, has-miR-26a, has-miR-27a, has-miR-92, and has-miR-200a) showed a decreased expression by

more than 50%. Precursor miRNAs of these miRNAs were also transfected into the cells to examine the gain of function of the miRNAs (Supplementary Fig. 2). It was confirmed that the expressions of target genes of the five miRNAs (LIG4 [by has-miR-26a]; RGL2 [by has-miR-23a]; Rad51C [by has-miR-27a]; KAP3, CDC25B, KAP3, CDK2AP2, POLD, and CPSF4 [by has-miR-200a]; and TUBGCP2, SNX15 and BCAT2 [by has-miR-92]) were increased by the suppression of the miRNAs induced by anti-miRNAs and were decreased by the overexpression of precursor miRNAs (Supplementary Fig. 3).

Discussion

miRNA plays an important role in various diseases such as infection and cancer.¹⁻³ In this study, we examined miRNA expression profiles in normal liver and HCC, including nontumor lesions infected with HBV or HCV. Although the expression profiles of miRNAs in HCC have been reported,¹⁶⁻¹⁸ most of the studies were performed using a microarray system. Because we thought that miRNAs could not produce enough detection signals owing to their short length, we applied a highly sensitive and quantitative RTD-PCR method for miRNAs. Moreover, global gene expression in the same tissues was ana-

Table 4-1. Pathway Analysis of Targeted Genes by miRNAs that Were Commonly Repressed in CH-B, CH-C, HCC-B, and HCC-C Compared with Normal Liver (Cluster 1)

| No. | Pathway Name | P Value |
|---|---|----------|
| Down-regulated miRNA in CH-B,HCC-B,CH-C and HCC-C (possibly up-regulating target genes) | | |
| 1 | Cell adhesion_Platelet-endothelium-leukocyte interactions | 1.11E-02 |
| 2 | Cell cycle_S phase | 2.18E-02 |
| 3 | Protein folding_Protein folding nucleus | 2.43E-02 |
| 4 | Cell cycle_G1-S | 3.07E-02 |
| 5 | Development_Cartilage development | 3.89E-02 |
| 6 | Protein folding_Folding in normal condition | 3.89E-02 |
| 7 | Proteolysis_Connective tissue degradation | 3.99E-02 |
| 8 | Proteolysis_Proteolysis in cell cycle and apoptosis | 4.31E-02 |
| 9 | Signal Transduction_BMP and GDF signaling | 5.81E-02 |
| 10 | Immune_Antigen presentation | 6.05E-02 |

lyzed via cDNA microarray to examine whether the differentially expressed miRNAs could regulate their target genes. Because the absolute standard of miRNA is not available at present, and miRNA expression was compared within the samples and genes analyzed in this study, there might be possible errors when a larger number of samples and genes were analyzed.

Using these systems, we found that the expression profile in miRNAs was clearly different according to HBV and HCV infection for the first time. The differences were confirmed by the nonsupervised learning method, hierar-

Table 4-2. Pathway Analysis of Targeted Genes by Differentially Expressed miRNAs Between HBV-Related Liver Disease (CH-B,HCC-B) and HCV Related Liver Disease (CH-C,HCC-C Cluster 2)

| No. | Pathway Name | P Value |
|--|--|----------|
| Down-regulated miRNA in CH-C,HCC-C (possibly up-regulating target genes) | | |
| 1 | Immune_Phagosome in antigen presentation | 5.80E-04 |
| 2 | Muscle contraction | 1.05E-03 |
| 3 | Immune_Antigen presentation | 5.75E-03 |
| 4 | Cell cycle_Meiosis | 1.49E-02 |
| 5 | Reproduction_Male sex differentiation | 2.06E-02 |
| 6 | Cell adhesion_Platelet aggregation | 2.77E-02 |
| 7 | Transport_Synaptic vesicle exocytosis | 3.56E-02 |
| 8 | Inflammation_Kallikrein-kinin system | 3.73E-02 |
| 9 | Inflammation_IgE signaling | 4.10E-02 |
| 10 | Development_Skeletal muscle development | 5.02E-02 |
| Down-regulated miRNA in CH-B,HCC-B (possibly up-regulating target genes) | | |
| 1 | Signal Transduction_Cholecystokinin signaling | 1.15E-04 |
| 2 | Inflammation_NK cell cytotoxicity | 5.29E-03 |
| 3 | Signal transduction_CREM pathway | 5.31E-03 |
| 4 | Reproduction_GnRH signaling pathway | 7.80E-03 |
| 5 | DNA damage_DBS repair | 1.02E-02 |
| 6 | Cell cycle_G2-M | 1.63E-02 |
| 7 | Development_Neuromuscular junction | 2.07E-02 |
| 8 | Apoptosis_Apoptosis mediated by external signals | 2.42E-02 |
| 9 | Reproduction_FSH-beta signaling pathway | 2.92E-02 |
| 10 | Cell adhesion_Amyloid proteins | 3.81E-02 |

Table 4-3. The Pathway Analysis of Targeted Genes by Differentially Expressed miRNAs Between CH and HCC (Cluster 3)

| No. | Pathway Name | P Value |
|---|---|----------|
| Down-regulated miRNA in HCC (possibly up-regulating target genes) | | |
| 1 | Cytoskeleton_Spindle microtubules | 2.15E-03 |
| 2 | Transcription_Chromatin modification | 5.27E-03 |
| 3 | Proteolysis_Ubiquitin-proteasomal proteolysis | 6.43E-03 |
| 4 | Cell adhesion_Cell-matrix interactions | 7.30E-03 |
| 5 | Cell cycle_Meiosis | 7.83E-03 |
| 6 | DNA damage_Checkpoint | 1.69E-02 |
| 7 | Reproduction_Progesterone signaling | 1.94E-02 |
| 8 | Apoptosis_Apoptotic mitochondria | 3.14E-02 |
| 9 | Translation_Regulation of initiation | 4.22E-02 |
| 10 | Signal transduction_WNT signaling | 4.26E-02 |
| Up-regulated miRNA in HCC (possibly down-regulating target genes) | | |
| 1 | Inflammation_IgE signaling | 1.05E-02 |
| 2 | Inflammation_Kallikrein-kinin system | 2.46E-02 |
| 3 | Inflammation_Innate inflammatory response | 2.51E-02 |
| 4 | Inflammation_Histamine signaling | 4.25E-02 |
| 5 | Inflammation_Neutrophil activation | 4.55E-02 |
| 6 | Chemotaxis | 4.68E-02 |
| 7 | Inflammation_IL-12,15,18 signaling | 5.16E-02 |
| 8 | Inflammation_NK cell cytotoxicity | 7.25E-02 |
| 9 | Cell cycle_G0-G1 | 7.53E-02 |
| 10 | Inflammation_Complement system | 7.72E-02 |

chical clustering (Fig. 2A), and supervised learning methods based on SVM at an 87% accuracy ($P < 0.001$) (Table 2-1). As similarly described, the expression profile in miRNAs was significantly different according to the progression of liver disease (normal, CH, and HCC) in this study. The present CH and HCC expression data were derived from the same patient, and some microarray analyses suggested that the noncancerous liver tissue can predict the prognosis of HCC.^{19,20} We examined whether the miRNA expression of paired samples was similar or independent using the Dunnett test¹² (Supplementary Data). Our data indicated that miRNA expression profiling was more dependent on the disease condition than on the paired condition, although the issue of paired samples should be taken into account carefully.

Binary tree prediction analysis and detailed assessment of hierarchical clustering revealed two types of differential miRNAs, one associated with HBV and HCV infection, the other associated with the stages of liver disease that were irrelevant to the differences in HBV and HCV infection. We found that differences in miRNA expression between liver tissues with HBV and HCV (HBV/HCV) were strongly correlated with those in miRNA between cultured cell models of HBV and HCV infection (HBV/HCV) ($r = 0.73$ $P = 0.0006$) (Fig. 5). Thus, there exist HBV- and HCV-infection-specific miRNAs that potentially regulate viral replication and host gene signaling pathways in hepatocytes.

| | HBV/HCV | |
|----------------|---------|--------|
| | Tissue | Huh7.5 |
| hsa-miR-20 | 0.61 | 0.36 |
| hsa-miR-23a | -1.12 | -1.3 |
| hsa-miR-27a | -0.75 | -1.51 |
| hsa-miR-34c | -2.29 | N.D. |
| hsa-miR-105 | 2.13 | N.D. |
| hsa-miR-124b | -1.63 | -10.54 |
| hsa-miR-133b | 1.28 | -3.64 |
| hsa-miR-134 | 2.52 | -0.63 |
| hsa-miR-142-5p | -1.34 | -4.39 |
| hsa-miR-151 | 0.86 | -0.29 |
| hsa-miR-182* | 1.16 | 0.37 |
| hsa-miR-190 | 1.04 | 1.32 |
| hsa-miR-191 | 0.48 | 1.16 |
| hsa-miR-193 | 0.74 | -0.03 |
| hsa-miR-194 | 0.74 | 0.76 |
| hsa-miR-211 | 4.68 | 5.26 |
| hsa-miR-324-5p | 0.59 | 1.16 |
| hsa-miR-340 | 0.57 | 1.68 |
| hsa-let-7a | -1.14 | -4.51 |




Fig. 5. Correlation between differences in miRNA expression between liver tissues infected with HBV and HCV and those in miRNA expression between cultured cell models of HBV and HCV infections. A total of 140 of 188 miRNAs were confirmed to be expressed in Huh7.5 cells. There was a significant correlation of infection-associated miRNA (closed lozenge) in vitro and in vivo ($r = 0.73$, $P = 0.0006$), but none for the other 121 miRNAs (open lozenge) ($r = 0.02$, $P = 0.82$).

The pathway analysis of targeted genes by miRNAs revealed that 13 miRNAs exhibiting a decreased expression in the HCV group regulate genes related to immune response, antigen presentation, cell cycle, proteasome, and lipid metabolism. Six miRNAs showing a decreased expression in the HBV group regulate genes related to cell death, DNA damage and recombination, and transcription signals. These findings reflected differences in the gene expression profile between CH-B and CH-C as described.¹⁰ Many of the miRNAs were down-regulated in the HCV group rather than in the HBV group. It has been reported that human endogenous miRNAs may be involved in defense mechanisms, mainly against RNA viruses.²¹ On the other hand, it is suggested that endogenous miRNAs may be consumed and reduced by defense mechanisms, especially those against RNA viruses.

Although the expressions of these HBV- and HCV-infection-specific miRNAs were irrelevant to the differences in CH and HCC (Fig. 3, cluster 2), some of them have been reported to play pivotal roles in the occurrence of cancer. For example, has-let-7a regulates ras and c-myc genes,²² and has-miR-34 is involved in the p53 tumor suppressor pathway.²³ These miRNAs were down-regulated in the HBV group, possibly participating in a more aggressive and malignant phenotype in HCC-B rather than in HCC-C. High expression of has-miR-191 was shown to be significantly associated with the worse survival in acute myeloid leukemia,²⁴ and has-miR-191 was

overexpressed in the HBV group compared with the HCV group. On the other hand, has-miR-133b, which was reported to be down-regulated in squamous cell carcinoma,²⁵ was repressed in the HCV group compared with the HBV group. Some hematopoietic-specific miRNAs such as has-miR-142-5p were up-regulated in the HCV group. Therefore, these miRNAs were not only HBV and HCV infection-associated but also tumor-associated. These findings indicate different mechanisms of development of HCC infected with HBV and HCV (Fig. 6).

Following HCC development, common changes in miRNA expression between HCC-B and HCC-C appeared (Fig. 3, cluster 3). The 23 miRNAs mentioned above clearly differentiated CH and HCC that were irrelevant to the differences in HBV and HCV infections. Seventeen miRNAs were down-regulated in HCC, which up-regulated cancer-associated pathways. Six miRNAs were up-regulated in HCC that down-regulated all inflammation-mediated signaling pathways, potentially reflecting impaired antitumor immune response in HCC. These results suggest that common signaling pathways are involved in HCC development from CH, and that HBV- and HCV-specific miRNAs participate in generating HCC-specific miRNA expressions (Fig. 6). Therefore, these miRNAs might be good candidates for molecular targeting to prevent HCC occurrence, because they reg-




Fig. 6. Infection-associated and HCC-specific miRNAs and liver disease progression.

ulate a common signaling pathway underlying HCC-B and HCC-C development.

In conclusion, we showed that miRNAs are important mediators of HBV and HCV infections as well as liver disease progression. Further studies are needed to enable more detailed mechanistic analysis of the miRNAs identified here and to evaluate the usefulness of miRNAs as diagnostic/prognostic markers and potential therapeutic target molecules.

Acknowledgement: The authors thank Mikiko Nakamura and Nami Nishiyama for excellent technical assistance.

References

- He L, Thomson JM, Hemann MT, Hernando-Monge E, Mu D, Goodson S, et al. A microRNA polycistron as a potential human oncogene. *Nature* 2005;435:828-833.
- Calin GA, Dumitru CD, Shimizu M, Bichi R, Zupo S, Noch E, et al. Frequent deletions and down-regulation of micro-RNA genes miR15 and miR16 at 13q14 in chronic lymphocytic leukemia. *Proc Natl Acad Sci U S A* 2002;99:15524-15529.
- Cho WC. OncomiRs: the discovery and progress of microRNAs in cancers. *Mol Cancer* 2007;6:60.
- Hutvagner G, Zamore PD. A microRNA in a multiple-turnover RNAi enzyme complex. *Science* 2002;297:2056-2060.
- Ambros V, Bartel B, Bartel DP, Burge CB, Carrington JC, Chen X, et al. A uniform system for microRNA annotation. *RNA* 2003;9:277-279.
- Lee RC, Feinbaum RL, Ambros V. The *C. elegans* heterochronic gene *lin-4* encodes small RNAs with antisense complementarity to *lin-14*. *Cell* 1993;75:843-854.
- Lewis BP, Burge CB, Bartel DP. Conserved seed pairing, often flanked by adenosines, indicates that thousands of human genes are microRNA targets. *Cell* 2005;120:15-20.
- Kiyosawa K, Sodeyama T, Tanaka E, Gibo Y, Yoshizawa K, Nakano Y, et al. Interrelationship of blood transfusion, non-A, non-B hepatitis and hepatocellular carcinoma: analysis by detection of antibody to hepatitis C virus. *HEPATOLOGY* 1990;12:671-675.
- Honda M, Kaneko S, Kawai H, Shirota Y, Kobayashi K. Differential gene expression between chronic hepatitis B and C hepatic lesion. *Gastroenterology* 2001;120:955-966.
- Honda M, Yamashita T, Ueda T, Takatori H, Nishino R, Kaneko S. Different signaling pathways in the livers of patients with chronic hepatitis B or chronic hepatitis C. *HEPATOLOGY* 2006;44:1122-1138.
- Molinari AM, Simon R, Pfeiffer RM. Prediction error estimation: a comparison of resampling methods. *Bioinformatics* 2005;21:3301-3307.
- Dunnnett CW. A multiple comparison procedure for comparing several treatments with a control. *J Am Stat Assoc* 1955;50:1096-1121.
- Weiss L, Kekule AS, Jakubowski U, Burgelt E, Hofschneider PH. The HBV-producing cell line HepG2-4A5: a new in vitro system for studying the regulation of HBV replication and for screening anti-hepatitis B virus drugs. *Virology* 1996;216:214-218.
- Lindenbach BD, Evans MJ, Syder AJ, Wolk B, Tellinghuisen TL, Liu CC, et al. Complete replication of hepatitis C virus in cell culture. *Science* 2005;309:623-626.
- Wakita T, Pietschmann T, Kato T, Date T, Miyamoto M, Zhao Z, et al. Production of infectious hepatitis C virus in tissue culture from a cloned viral genome. *Nat Med* 2005;11:791-796.
- Murakami Y, Yasuda T, Saigo K, Urashima T, Toyoda H, Okanoue T, et al. Comprehensive analysis of microRNA expression patterns in hepatocellular carcinoma and non-tumorous tissues. *Oncogene* 2006;25:2537-2545.
- Varnholt H, Drebber U, Schulze F, Wedemeyer J, Schirmacher P, Dienes HP, et al. MicroRNA gene expression profile of hepatitis C virus-associated hepatocellular carcinoma. *HEPATOLOGY* 2008;47:1223-1232.
- Budhu A, Jia HL, Forgues M, Liu CG, Goldstein D, Lam A, et al. Identification of metastasis-related microRNAs in hepatocellular carcinoma. *HEPATOLOGY* 2008;47:897-907.
- Budhu A, Forgues M, Ye QH, Jia HL, He P, Zanetti KA, et al. Prediction of venous metastases, recurrence, and prognosis in hepatocellular carcinoma based on a unique immune response signature of the liver microenvironment. *Cancer Cell* 2006;10:99-111.
- Hoshida Y, Villanueva A, Kobayashi M, Peix J, Chiang DY, Camargo A, et al. Gene expression in fixed tissues and outcome in hepatocellular carcinoma. *N Engl J Med* 2008;359:1995-2004.
- Jopling CL, Yi M, Lancaster AM, Lemon SM, Sarnow P. Modulation of hepatitis C virus RNA abundance by a liver-specific MicroRNA. *Science* 2005;309:1577-1581.
- Johnson CD, Esquela-Kerscher A, Stefani G, Byrom M, Kelnar K, Ovcharenko D, et al. The let-7 microRNA represses cell proliferation pathways in human cells. *Cancer Res* 2007;67:7713-7722.
- He X, He L, Hannon GJ. The guardian's little helper: microRNAs in the p53 tumor suppressor network. *Cancer Res* 2007;67:11099-11101.
- Garzon R, Volinia S, Liu CG, Fernandez-Cymering C, Palumbo T, Pichiorri F, et al. MicroRNA signatures associated with cytogenetics and prognosis in acute myeloid leukemia. *Blood* 2008;111:3183-3189.
- Wong TS, Liu XB, Chung-Wai Ho A, Po-Wing Yuen A, Wai-Man Ng R, Ignace Wei W. Identification of pyruvate kinase type M2 as potential oncoprotein in squamous cell carcinoma of tongue through microRNA profiling. *Int J Cancer* 2008;123:251-257.

EpCAM-Positive Hepatocellular Carcinoma Cells Are Tumor-Initiating Cells With Stem/Progenitor Cell Features

TARO YAMASHITA,* JUNFANG JI,* ANURADHA BUDHU,* MARSHONNA FORGUES,* WEN YANG,[†] HONG-YANG WANG,[‡] HULIANG JIA,[§] QINGHAI YE,[§] LUN-XIU QIN,[§] ELAINE WAUTHIER,^{||} LOLA M. REID,^{||} HIROSHI MINATO,^{||} MASAO HONDA,^{||} SHUICHI KANEKO,^{||} ZHAO-YOU TANG,[§] and XIN WEI WANG*

*Liver Carcinogenesis Section, Laboratory of Human Carcinogenesis, Center for Cancer Research, National Cancer Institute, Bethesda, Maryland; [†]International Cooperation Laboratory on Signal Transduction, Eastern Hepatobiliary Surgery Institute, Shanghai, China; [‡]Liver Cancer Institute and Zhongshan Hospital, Fudan University, Shanghai, China; [§]Department of Cell and Molecular Physiology, University of North Carolina School of Medicine, Chapel Hill, North Carolina; and the ^{||}Liver Disease Center and Kanazawa University Hospital, Kanazawa University, Kanazawa, Japan

Background & Aims: Cancer progression/metastases and embryonic development share many properties including cellular plasticity, dynamic cell motility, and integral interaction with the microenvironment. We hypothesized that the heterogeneous nature of hepatocellular carcinoma (HCC), in part, may be owing to the presence of hepatic cancer cells with stem/progenitor features. **Methods:** Gene expression profiling and immunohistochemistry analyses were used to analyze 235 tumor specimens derived from 2 recently identified HCC subtypes (EpCAM⁺ α -fetoprotein [AFP]⁺ HCC and EpCAM⁻ AFP⁻ HCC). These subtypes differed in their expression of AFP, a molecule produced in the developing embryo, and EpCAM, a cell surface hepatic stem cell marker. Fluorescence-activated cell sorting was used to isolate EpCAM⁺ HCC cells, which were tested for hepatic stem/progenitor cell properties. **Results:** Gene expression and pathway analyses revealed that the EpCAM⁺ AFP⁺ HCC subtype had features of hepatic stem/progenitor cells. Indeed, the fluorescence-activated cell sorting-isolated EpCAM⁺ HCC cells displayed hepatic cancer stem cell-like traits including the abilities to self-renew and differentiate. Moreover, these cells were capable of initiating highly invasive HCC in nonobese diabetic, severe combined immunodeficient mice. Activation of Wnt/ β -catenin signaling enriched the EpCAM⁺ cell population, whereas RNA interference-based blockage of EpCAM, a Wnt/ β -catenin signaling target, attenuated the activities of these cells. **Conclusions:** Taken together, our results suggest that HCC growth and invasiveness is dictated by a subset of EpCAM⁺ cells, opening a new avenue for HCC cancer cell eradication by targeting Wnt/ β -catenin signaling components such as EpCAM.

Tumors originate from normal cells as a result of accumulated genetic/epigenetic changes. Although considered monoclonal in origin, tumor cells are heterogeneous in their morphology, clinical behavior, and mo-

lecular profiles.^{1,2} Tumor cell heterogeneity has been explained previously by the clonal evolution model³; however, recent evidence has suggested that heterogeneity may be owing to derivation from endogenous stem/progenitor cells⁴ or de-differentiation of a transformed cell.⁵ This hypothesis supports an early proposal that cancers represent “blocked ontogeny”⁶ and a derivative that cancers are transformed stem cells.⁷ This renaissance of stem cells as targets of malignant transformation has led to realizations about the similarities between cancer cells and normal stem cells in their capacity to self-renew, produce heterogeneous progenies, and limitlessly divide.⁸ The cancer stem cell (CSC) (or tumor-initiating cell) concept is that a subset of cancer cells bear stem cell features that are indispensable for a tumor. Accumulating evidence suggests the involvement of CSCs in the perpetuation of various cancers including leukemia, breast cancer, brain cancer, prostate cancer, and colon cancer.⁹⁻¹³ Experimentally, putative CSCs have been isolated using cell surface markers specific for normal stem cells. Stem cell-like features of CSC have been confirmed by functional in vitro clonogenicity and in vivo tumorigenicity assays. For example, leukemia-initiating cells in nonobese diabetic, severe combined immunodeficient (NOD/SCID) mice are CD34⁺CD38⁻.¹¹ Breast cancer CSCs are CD44⁺CD24^{-/low} cells, whereas tumor-initiating cells of the brain, colon, and prostate are CD133⁺.^{10,12,13} CSCs are considered more metastatic and drug-/radiation-resistant than non-CSCs in the tumor, and are responsible for cancer relapse. These findings warrant the development of treatment strategies that can specifically eradicate CSCs.^{14,15}

Abbreviations used in this paper: AFP, α -fetoprotein; BIO, 6-bromoindirubin-3'-oxime; CSC, cancer stem cell; FACS, fluorescence-activated cell sorting; 5-FU, 5-fluorouracil; HpSC, hepatic stem cell; IF, immunofluorescence; IHC, immunohistochemistry; MACS, magnetic-activated cell sorting; MeBIO, 1-methyl-BIO; MH, mature hepatocyte; PCNA, proliferating cell nuclear antigen; siRNA, small interfering RNA.

© 2009 by the AGA Institute

0016-5085/09/\$36.00

doi:10.1053/j.gastro.2008.12.004

Hepatocellular carcinoma (HCC) is the third leading cause of cancer death worldwide.¹⁶ Although the cellular origin of HCC is unclear,^{17,18} HCC has heterogeneous pathologies and genetic/genomic profiles,¹⁹ suggesting that HCC can initiate in different cell lineages.²⁰ The liver is considered as a maturational lineage system similar to that in the bone marrow.²¹ Experimental evidence indicates that certain forms of hepatic stem cells (HpSCs), present in human livers of all donor ages, are multipotent and can give rise to hepatoblasts,^{22,23} which are, in turn, bipotent progenitor cells that can progress either into hepatocytic or biliary lineages.^{22,24} α -fetoprotein (AFP) is one of the earliest markers detected in the liver bud specified from the ventral foregut,^{25,26} but its expression has been found only in hepatoblasts and to a lesser extent in committed hepatocytic progenitors, not in later lineages or in normal human HpSC.²² Recent studies also have indicated that EpCAM is a biomarker for HpSC because it is expressed in HpSCs and hepatoblasts.²²⁻²⁴

We recently identified a novel HCC classification system based on EpCAM and AFP status.²⁷ Gene expression profiles revealed that EpCAM⁺ AFP⁺ HCC (referred to as *HpSC-HCC*) has progenitor features with poor prognosis, whereas EpCAM⁻ AFP⁻ HCC (referred to as *mature hepatocyte-like HCC*; MH-HCC) have adult hepatocyte features with good prognosis. Wnt/ β -catenin signaling, a critical player for maintaining embryonic stem cells,²⁸ is activated in EpCAM⁺ AFP⁺ HCC, and EpCAM is a direct transcriptional target of Wnt/ β -catenin signaling.²⁹ Moreover, EpCAM⁺ AFP⁺ HCC cells are more sensitive to β -catenin inhibitors than EpCAM⁻ HCC cells in vitro.²⁹ Interestingly, a heterogeneous expression of EpCAM and AFP was observed in clinical tissues, a feature that may be attributed to the presence of a subset of CSCs. In this study, we have confirmed that EpCAM⁺ HCC cells are highly invasive and tumorigenic, and have activated Wnt/ β -catenin signaling. We also show a crucial role of EpCAM in the maintenance of hepatic CSCs. Our data shed new light on the pathogenesis of HCC and may open new avenues for therapeutic interventions for targeting hepatic CSCs.

Materials and Methods

Clinical Specimens

HCC samples were obtained with informed consent from patients who underwent radical resection at the Liver Cancer Institute of Fudan University, Eastern Hepatobiliary Surgery Institute, and the Liver Disease Center of Kanazawa University Hospital, and the study was approved by the institutional review boards of the respective institutes. The microarray data from clinical specimens are available publicly (GEO accession number, GSE5975).²⁷ Array data from a total of 156 HCC cases (155 hepatitis B virus [HBV]-positive) corresponding to 2 subtypes of HCC (ie, HpSC-HCC and MH-HCC), were

used to search for HpSC-HCC-associated genes (Supplementary Table 1; see supplementary material online at www.gastrojournal.org). A total of 79 formalin-fixed and paraffin-embedded HCC samples were used for immunohistochemistry (IHC) analyses (Supplementary Table 2; see supplementary material online at www.gastrojournal.org), 56 of which also were used in a recent study.³⁰ The classification of HpSC-HCC and MH-HCC was based on previously described criteria.²⁷

Cell Cultures and Sorting

Human liver cancer cell lines (HuH1 and HuH7) were derived from Health Science Research Resources Bank (JCRB0199 and JCRB0403, respectively) and routinely cultured as previously described.³¹ Normal human MHs, provided by the University of Pittsburgh through Liver Tissue Cell Distribution System, were cultured as previously described.³² Human HpSCs were isolated from fetal livers and cultured in Kubota and Reid's³³ medium as previously described. Wnt10B conditioned medium was prepared as described.³⁴ Embryonic stem cell culture medium was prepared using Knockout Dulbecco's modified Eagle medium supplemented with 18% of Serum Replacement (Invitrogen, Carlsbad, CA). The pTOP-FLASH and pFOP-FLASH luciferase constructs were described previously.²⁹ BIO and MeBIO were generous gifts from Ali Brivanlou (The Rockefeller University, New York, NY). For isolating single cell-derived colonies to determine whether heterogeneity is an intrinsic property of EpCAM⁺ cells, HuH1 and HuH7 cells were resuspended and plated as a single cell per well in 96-well plates. A total of 192 single cells were plated successfully. The clones that grew well were selected 2 weeks after seeding and used for immunofluorescence (IF) analysis. The 5-fluorouracil (5-FU) stock (2 mg/mL; Sigma, St Louis, MO), was prepared in distilled water. Fluorescence-activated cell sorting (FACS) and magnetic-activated cell sorting (MACS) analyses were used to isolate EpCAM⁺ HCC cells (Supplementary materials; see supplementary Materials and Methods online at www.gastrojournal.org).

Clonogenicity, Spheroid Formation, Invasion, Quantitative Reverse Transcription-Polymerase Chain Reaction, and IHC Assays

For colony formation assays, 2000 EpCAM⁺ or EpCAM⁻ cells were seeded in 6-well plates after FACS. After 10 days of culture, cells were fixed by 100% methanol and stained with methylene blue. For spheroid assays, single-cell suspensions of 1000 EpCAM⁺ or EpCAM⁻ cells were seeded in 6-well Ultra-Low Attachment Microplates (Corning, Corning, NY) after FACS. The number of spheroids was measured 14 days after seeding. Invasion assays were performed using BD Bio-Coat Matrigel Matrix Cell Culture Inserts and Control Inserts (BD Biosciences, San Jose, CA) essentially as pre-

viously described.³¹ Reverse transcription-polymerase chain reaction and IHC assays are described in detail in the supplementary materials (see supplementary material online at www.gastrojournal.org).

Tumorigenicity in NOD/SCID Mice

Six-week-old NOD/SCID mice (NOD/NCrCrl-Prkdc^{scid}) were purchased from Charles River (Charles River Laboratories, Inc, Wilmington, MA). The protocol was approved by the National Cancer Institute-Bethesda Animal Care and Use Committee. Cells were suspended in 200 μ L of Dulbecco's modified Eagle medium and Matrigel (1:1), and a subcutaneous injection was performed. The size and incidence of subcutaneous tumors were recorded. For histologic evaluation, tumors were formalin-fixed, paraffin-embedded or embedded directly in OCT compound (Sakura Finetek, Torrance, CA) and stored at -80°C .

RNA Interference

A small interfering RNA (siRNA) specific to *TACSTD1* (SI03019667) and a control siRNA (1022076) were designed and synthesized by Qiagen (Qiagen, Valencia, CA). Transfection was performed using Lipofectamine 2000 (Invitrogen), according to the manufacturer's instructions. A total of 200 nmol/L of siRNA duplex was used for each transfection.

Statistical Analyses

The class comparison and gene clustering analyses were performed as previously described.³⁰ The canonical pathway analysis was performed using Ingenuity Pathways Analysis (v5.5; Ingenuity Systems, Redwood City, CA). The association of HCC subtypes and clinicopathologic characteristics was examined using either the Mann-Whitney *U* test or the chi-square test. Student *t* tests were used to compare various test groups assayed by colony formation, spheroid formation, or invasion assays. The Kaplan-Meier survival analysis was performed to compare patient survival or tumorigenicity.

Results

A Poor Prognostic HCC Subtype With Molecular Features of HpSC


We re-evaluated the gene expression profiles that were uniquely associated with 2 recently identified prognostic subtypes of HCC (ie, HpSC-HCC and MH-HCC), using a publicly available microarray dataset of 156 HCC cases (GEO accession number: GSE5975). Sixty cases were defined as HpSC-HCC with a poor prognosis and 96 cases were defined as MH-HCC with a good prognosis, based on EpCAM and AFP status.²⁷ A class-comparison analysis with univariate *t* tests and a global permutation test (1000 \times) yielded 793 genes that were expressed differentially between HpSC-HCC and MH-HCC ($P < .001$). Hierarchical cluster analyses revealed 2 main gene clusters

that were up-regulated (cluster A; 455 genes) or down-regulated (cluster B; 338 genes) in HpSC-HCC (Figure 1A). Pathway analysis indicated that the enriched genes in cluster A were associated significantly with known stem cell signaling pathways such as transforming growth factor- β , Wnt/ β -catenin, PI3K/Akt, and integrin ($P < .01$) (Figure 1B). In contrast, genes in cluster B were associated significantly with mature hepatocyte functions such as xenobiotic metabolism, complement system, and coagulation system ($P < .01$). Noticeably, known HpSC markers such as *KRT19* (CK19), *TACSTD1* (EpCAM), *AFP*, *DKK1*, *DLK1*, and *PROM1* (CD133) were up-regulated significantly in HpSC-HCC, whereas known liver maturation markers such as *UGT2B7* and *CYP3A4* were expressed more abundantly in MH-HCC (Figure 1C and Supplementary Tables 3 and 4; see supplementary material online at www.gastrojournal.org). Kaplan-Meier survival analysis revealed that HpSC-HCC patients had a significantly shorter survival than MH-HCC patients ($P = .036$) (Figure 1D). Consistently, HpSC-HCC patients had a high frequency of macroscopic and microscopic portal vein invasion (Figure 1E).

However, IHC analyses of an additional 79 HCC cases revealed that among 24 HpSC-HCC cases, EpCAM staining was very heterogeneous with a mixture of EpCAM⁺ and EpCAM⁻ tumor cells in each tumor (Figure 1F, left panel). Noticeably, many of the EpCAM⁺ tumor cells were located at the invasion border zones and often were disseminated at the invasive front (black arrows). IF analysis revealed that HCC cells located at the invasive front co-expressed EpCAM, CK19, and AFP (Figure 1F, right panels). Noticeably, HpSC-HCC patients were significantly younger than MH-HCC patients (Supplementary Tables 1 and 2; see supplementary material online at www.gastrojournal.org). Enrichment of EpCAM⁺ AFP⁺ tumor cells at the tumor-invasive front suggested their involvement in HCC invasion and metastasis.

Isolation and Characterization of EpCAM⁺ Cells in HCC

The results described earlier suggest that HpSC-HCC may be organized in a hierarchical fashion in which EpCAM⁺ tumor cells act as stem-like cells with an ability to differentiate into EpCAM⁻ tumor cells. To test this hypothesis, we first evaluated the expression pattern of 7 hepatic stem/maturation markers (EpCAM, CD133, CD90, CK19, Vimentin, Hep-Par1, and β -catenin) in 6 HCC cell lines (Figure 2A). All 3 AFP⁺ cell lines (HuH1, HuH7, and Hep3B) expressed EpCAM, CD133, and cytoplasmic/nuclear β -catenin, whereas the other 3 AFP⁻ cell lines (SK-Hep-1, HLE, and HLF) did not, consistent with the microarray data. Interestingly, AFP⁻ cell lines had no CD90⁺ cell population, which recently was identified as hepatic CSCs,³⁵ whereas AFP⁻ cell lines had such a population. Consistent with the IF data, FACS analysis showed that AFP⁺ cell lines had a subpopulation of



EpCAM⁺ and CD133⁺, but no CD90⁺ cells, whereas AFP⁻ cell lines had a subpopulation of CD90⁺ cells but no EpCAM⁺ or CD133⁺ cells (Figure 2B). These data indicate that HpSC-HCC and MH-HCC cell lines have distinct stem cell marker expression patterns, and EpCAM as well as CD133 may be hepatic CSC markers specifically in HpSC-HCC.

We selected 2 human HCC cell lines (HuH1 and HuH7) to isolate EpCAM⁺ cells because both lines were heterogeneous in EpCAM, AFP, CK19, and β -catenin expression (Figure 2A and B and Supplementary Figure 1A; see supplementary material online at www.gastrojournal.org).²⁹ We successfully enriched EpCAM⁺ and EpCAM⁻ populations from HuH7 cells by FACS, with more than 80%

purity in EpCAM⁺ cells and more than 90% purity in EpCAM⁻ cells 1 day after sorting (Figure 3A). Similar results were obtained when the purity check was performed immediately after sorting (data not shown). EpCAM⁺ cells also were positive for CK19 and β -catenin (Figure 3B and Supplementary Figure 1B; see supplementary material online at www.gastrojournal.org) and most were AFP⁺ (data not shown). In contrast, EpCAM⁻ cells were negative for these markers but positive for HepPar1, a monoclonal antibody specific to hepatocytes (Figure 3B). Consistent with the microarray data described earlier, the levels of *TACSTD1*, *MYC*, and *bTERT* (known HpSC markers) were increased significantly in EpCAM⁺ HuH7 cells, whereas the levels of *UGT2B7* and *CYP3A4*





Figure 2. Characterization of hepatic stem cell marker expression in HCC cell lines. (A) IF analysis of 6 HCC cell lines (EpCAM⁺ AFP⁺ cell lines: HuH1, HuH7, and Hep3B; EpCAM⁻ AFP⁻ cell lines: SK-Hep-1, HLE, and HLF) stained with anti-EpCAM, anti-CD133, anti-CD90, anti-CK19, anti-Vimentin, anti-Hep-Par1, and anti-β-catenin antibodies. (B) FACS analysis of 6 HCC cell lines stained with anti-EpCAM, anti-CD133, and anti-CD90 antibodies.

(known mature hepatocyte markers) were significantly higher in EpCAM⁻ HuH7 cells (Figure 3C, left upper panel). This expression pattern was reminiscent of human HpSC cells (Figure 3C, left lower panel). Similar results were obtained from HuH1 cells (data not shown). We also compared gene expression patterns of isolated HuH1, HuH7, MH, and HpSC cells using the TaqMan Human Stem Cell Pluripotency Array (Applied Biosystems, Foster City, CA) containing 96 selected human stem cell-related genes. Although a differential expres-

sion pattern of stem cell-related genes was evident among HpSC, EpCAM⁺ HuH1, and EpCAM⁺ HuH7 cells, the EpCAM⁺ HCC cells were related more closely to HpSC cells whereas EpCAM⁻ HCC cells were related more closely to diploid adult mature hepatocytes (Figure 3C, right panel; and Supplementary Figure 1C; see supplementary material online at www.gastrojournal.org). Thus, it appeared that EpCAM⁺ HCC cells had a gene expression pattern that is related more closely to HpSC than EpCAM⁻ HCC cells.

BASIC-LIVER, PANCREAS, AND BILIARY TRACT

Figure 3. Characterization of EpCAM⁺ and EpCAM⁻ cells in HuH7 cells. (A) FACS analysis of EpCAM⁺ and EpCAM⁻ cells on day 1 after cell sorting. (B) IF analysis of cells stained with anti-EpCAM, anti-AFP, anti-CK19, or anti- β -catenin antibodies. (C) Quantitative reverse-transcription polymerase chain reaction analysis of EpCAM⁺ and EpCAM⁻ HuH7 cells (left upper panel) or HpSCs and MHs (left lower panel). Experiments were performed in triplicate. Hierarchical cluster analysis of HpSC, MH, and EpCAM⁺ and EpCAM⁻ HCC cells using a panel of genes expressed in human embryonic stem cells (right panel). Gene expression was measured in quadruplicate. (D) Representative photographs of the plates containing colonies derived from 2000 EpCAM⁺ or EpCAM⁻ HuH7 cells (upper panel). Colony formation experiments were performed in triplicate (mean \pm SD) (middle panel). Cell invasiveness of EpCAM⁺ and EpCAM⁻ cells using the Matrigel invasion assay (lower panel). (E) Flow cytometer analysis of EpCAM⁺ and EpCAM⁻ HuH7 cells stained with anti-EpCAM at days 1 and 14 after cell sorting. (F) Percentage of sorted EpCAM⁺ and EpCAM⁻ cells after culturing for various times as analyzed by IF. Numbers of EpCAM⁺ and EpCAM⁻ cells were counted in 3 independent areas of chamber slides at days 1, 3, 7, and 15 after cell sorting. The average percentages of EpCAM⁺ or EpCAM⁻ cells are depicted as red or blue, respectively.



The isolated EpCAM⁺ HuH7 cells formed colonies efficiently whereas EpCAM⁻ cells failed to do so (Figure 3D, upper and middle panels; and Supplementary Figure 2A for HuH1 cells; see supplementary material online at www.gastrojournal.org). In addition, EpCAM⁺ HuH7 cells were much more invasive than EpCAM⁻ cells ($P < .03$) (Figure 3D, lower panel; and Supplementary Figure 2B for HuH1 cells; see supplementary material online at www.gastrojournal.org). The EpCAM⁺ fraction decreased with time in sorted EpCAM⁺ HuH7 cells from greater than 80% to 50% (Figure 3E). However, a small percentage

of EpCAM⁺ cells remained constant in sorted EpCAM⁻ HuH7 cells. FACS analysis confirmed the results of IF analysis (Figure 3F and Supplementary Figure 2C for HuH7 and HuH1 cells, respectively; see supplementary material online at www.gastrojournal.org), suggesting that EpCAM⁺ cells could differentiate into EpCAM⁻ cells, eventually allowing an enriched EpCAM⁺ fraction to revert back to parental cells after 14 days of culture. In contrast, EpCAM⁻ cells maintained their EpCAM⁻ status. In addition, we successfully isolated 12 HuH1 and 2 HuH7 colonies from 192 single-cell-plated culture wells.

BASIC-LIVER, PANCREAS, AND BILIARY TRACT




Figure 4. Spheroid formation of EpCAM⁺ HuH1 HCC cells. (A) A representative phase-contrast image of an HCC spheroid derived from an EpCAM⁺ cell (scale bar, 100 µm) and (B) total numbers of spheroids from 1000 sorted cells are shown. Experiments were performed in triplicate and data are shown as mean ± SD. (C) Representative confocal images of an HCC spheroid co-stained with anti-EpCAM, anti-AFP, and 4',6-diamidino-2-phenylindole (DAPI) (scale bar, 50 µm). (D) A 3-dimensional image of an HCC spheroid co-stained with anti-EpCAM, anti-AFP, and DAPI (scale bar, 50 µm) reconstructed from confocal images using surface rendering. (E) FACS analysis of EpCAM⁺ cells cultured as spheroid cells (red) or attached cells (blue) for 14 days after cell sorting. (F) Confocal images of an HCC spheroid co-stained with anti-PCNA, anti-AFP, and DAPI (scale bar, 50 µm).

However, all colonies were heterogeneous in EpCAM and AFP expression and no colony was completely EpCAM⁻ (data not shown). Taken together, these results indicate that EpCAM⁺ HCC cells resemble HpSC features. It appears that EpCAM⁺ cells, but not EpCAM⁻ cells, have self-renewal and differentiation capabilities with the ability to form colonies from a single cell, and produce both EpCAM⁺ and EpCAM⁻ cells.

It has been shown previously that stem/progenitor cells and cancer stem/progenitor cells can form spheroids in vitro in a nonattached condition.^{36,37} Consistently, EpCAM⁺ cells could form spheroids efficiently, reaching to about 150 to approximately 200 µm in diameter after 14 days of culture (Figure 4A and B). Interestingly, all cells in a spheroid were EpCAM⁺, whereas AFP expres-

sion was relatively heterogeneous (Figure 4C and D, and Supplementary movie 1; see supplementary material online at www.gastrojournal.org). Rarely, a few spheroids derived from an EpCAM⁻ cell fraction were positive for EpCAM (data not shown), suggesting that these spheroids were derived from contaminated residual EpCAM⁺ cells by FACS sorting. All spheroid cells maintained EpCAM expression while half of the attached cells lost EpCAM expression when the EpCAM⁺ fraction was cultured for 14 days (Figure 4E). Most spheroid cells also abundantly expressed proliferating cell nuclear antigen (PCNA), implying active cell proliferation (Figure 4F and Supplementary movie 2; see supplementary material online at www.gastrojournal.org). Thus, a subset of EpCAM⁺ cells, but not EpCAM⁻ cells, can form spheroids.




Figure 5. Tumorigenic and invasive potential of EpCAM⁺ HCC cells. (A) Representative NOD/SCID mice (upper panel) with subcutaneous tumors (lower panel) from EpCAM⁺ (black arrows) or EpCAM⁻ (white arrows) HuH1 cells. (B) Tumorigenicity of 200 sorted HuH1 cells. (C) Histologic analysis of EpCAM⁺ HuH1-derived xenografts. H&E staining of a subcutaneous tumor (left upper panel) with capsular invasion (left lower panel) and muscular invasion (right lower panel) and IF of the tumor stained with anti-EpCAM, anti-AFP, and 4',6-diamidino-2-phenylindole (DAPI) (right upper panel) (scale bar, 50 μ m). (D) Tumorigenicity of 1000 sorted cells derived from an EpCAM⁺ HuH1 xenograft. Data are generated from 10 mice in each group. (E) Representative NOD/SCID mice (left panel) with subcutaneous tumors from CD133⁻ (black arrows) or CD133⁺ (white arrows) (middle panel) and EpCAM⁺ (black arrows) or EpCAM⁻ (white arrows) (right panel) HuH1 cells. (F) Tumorigenicity of 1000 HuH1 cells sorted by anti-EpCAM (left panel) or anti-CD133 (right panel) antibodies.

EpCAM⁺ HCC Cells as Tumor-Initiating Cells

EpCAM⁺ HCC cells, but not EpCAM⁻ HCC cells, could efficiently initiate invasive tumors in NOD/SCID mice (Figure 5). For example, 10,000 EpCAM⁺ HuH1 cells produced large hypervascular tumors in 100% of mice whereas EpCAM⁻ cell fractions produced only small and pale-looking tumors in 30% of mice 4 weeks after injection (Figure 5A and Supplementary Figure 3A; see supplement-

tary material online at www.gastrojournal.org). Similar results were obtained with HuH7 cells (Supplementary Figure 3B–D; see supplementary material online at www.gastrojournal.org). As little as 200 EpCAM⁺ cells could initiate tumors in 8 of 10 injected mice, whereas 200 EpCAM⁻ cells produced only 1 tumor among 10 injected mice at 6 weeks after transplantation, and the tumor sizes were much larger in the EpCAM⁺ cells than in the EpCAM⁻

BASIC-LIVER, PANCREAS, AND BILIARY TRACT

cells (Figure 5B and Supplementary Figure 3E; see supplementary material online at www.gastrojournal.org). EpCAM⁺ cells produced tumors with a mixture of both EpCAM⁺ and EpCAM⁻ cells in xenografts, and these cells invaded in the capsule and muscles of the leg adjacent to the tumor (Figure 5C). EpCAM⁺ cells derived from tumors again maintained their tumor-initiating capacity, tumor morphology, and invasive ability in an in vivo serial transplantation experiment (Figure 5D). Occasionally, EpCAM⁻ cell fractions produced a few small tumors that always contained a mixture of EpCAM⁺ and EpCAM⁻ cells (data not shown), indicating that the contaminated EpCAM⁺ cells from FACS sorting contribute to the tumor-initiating ability.

To further validate whether EpCAM⁺ HCC cells were tumor-initiating cells, we isolated EpCAM⁺ HCC cells from 2 cases of AFP⁺ (>600 ng/mL serum AFP) HCC clinical specimens using MACS. Consistently, 1×10^4 EpCAM⁺ cells could induce tumors in NOD/SCID mice, but up to 1×10^6 EpCAM⁻ cells failed to do so (Table 1). In addition, similar to HCC cell lines, fresh EpCAM⁺ tumor cells from 2 clinical HCC specimens were more efficient in forming spheroids in vitro than EpCAM⁻ cells (Supplementary Figure 4; see supplementary material online at www.gastrojournal.org).

FACS analysis results indicate that a majority of EpCAM⁺ cells express CD133 in HuH7 cells but not in HuH1 cells (Figure 2B), which prompted us to compare the tumorigenic capacity of EpCAM⁺ and CD133⁺ cells in these cell lines. Noticeably, EpCAM⁺ HuH1 cells showed marked tumor-initiating capacity compared with CD133⁺ HuH1 cells (Figure 5E and F), whereas EpCAM⁻ and CD133⁺ cells had similar tumorigenic ability in HuH7 cells (data not shown).

GSK-3 β Inhibition Augments EpCAM⁺ HCC Cells

To determine the role of Wnt/ β -catenin signaling²⁸ in EpCAM⁺ HCC cells (Figure 1B), we first treated

HuH1, HuH7, and HLF cells with a GSK-3 β inhibitor BIO (Figure 6A), which activates Wnt/ β -catenin signaling (Figure 6B) and maintains undifferentiation of embryonic stem cells.³⁸ 6-bromoindirubin-3'-oxime (BIO) increased the EpCAM⁺ cell population in HuH1 and HuH7 cells when compared with the control methylated BIO (MeBIO) (Figure 6A). In contrast, BIO had no effect on the CD90⁺ cell population, which is more tumorigenic than the CD90⁻ cell population in HLF (Figure 6A and data not shown). Enrichment of EpCAM⁺ cells was provoked further by the treatment of Wnt10B-conditioned media in HuH7 cells (Figure 6C).³⁴ BIO induced morphologic alteration of HuH7 cells because most cells became small and round when compared with MeBIO and suppressed EpCAM⁻ AFP⁻ cell populations (Figure 6D). Moreover, BIO induced *TACSTD1*, *MYC*, and *hTERT* expression and spheroid formation (Figure 6E and F).

EpCAM Blockage by RNA Interference

One of the hallmarks of CSCs is its resistance to conventional chemotherapeutic agents resulting in tumor relapse and thus targeting CSCs is critical to achieve successful tumor remission. Consistently, 5-FU could increase the EpCAM⁺ population and spheroid formation of HuH1 and HuH7 cells (Figure 7A and B) (data not shown), suggesting a differential sensitivity of EpCAM⁺ and EpCAM⁻ HCC cells to 5-FU. In contrast, EpCAM blockage via RNA interference dramatically decreased the population of EpCAM⁺ cells (Figure 7C), and significantly inhibited cellular invasion, spheroid formation, and tumorigenicity of HuH1 cells (Figure 7D-F). Thus, EpCAM may serve as a molecular target to eliminate HCC cells with stem/progenitor cell features.

Discussion

The cellular origin of HCC is currently in debate. In this study, we found that EpCAM can serve as a marker to enrich HCC cells with tumor-initiating ability and with some stem/progenitor cell traits. EpCAM is expressed in many human cancers with an epithelial origin.³⁹ During embryogenesis, EpCAM is expressed in fertilized oocytes, embryonic stem cells, and embryoid bodies, suggesting its role in early stage embryogenesis.⁴⁰ Furthermore, a recent article indicated that EpCAM is expressed in colonic and breast CSCs.⁴¹ Taken together, these data suggest a critical role of EpCAM in CSCs as well as embryonic and somatic stem cells. Consistently, we found that EpCAM expression is regulated by Wnt/ β -catenin signaling²⁹ and tumorigenic and highly invasive HpSC-HCC is orchestrated by a subset of cells expressing EpCAM and AFP with stem cell-like features and self-renewal and differentiation capabilities regulated by Wnt/ β -catenin signaling (this study). Thus, EpCAM may be a common gene expressed in undifferentiated normal cells and HCCs with activated Wnt/ β -catenin signaling. It may act as a downstream molecule

Table 1. The Tumor-Initiating Capacity of EpCAM⁺ Cells From Clinical HCC Specimens

| HCC patients | | | Tumor incidence (mice with tumors/ total no. of mice injected) | | |
|--------------|--------------------------------------|--------------------|---|----------|----------|
| No. | % of EpCAM ⁺ HCC cells | Groups | No. of cells injected | 2 months | 3 months |
| 1 | 5.2 | EpCAM ⁺ | 1×10^3 | 0/3 | 0/3 |
| | | | 1×10^4 | 2/3 | 2/3 |
| | | | 1×10^5 | 2/2 | 2/2 |
| | | EpCAM ⁻ | 1×10^5 | 0/3 | 0/3 |
| | | | 1×10^6 | 0/2 | 0/2 |
| | | | | | |
| 2 | 1.4 | EpCAM ⁺ | 1×10^3 | 0/2 | 0/2 |
| | | | 1×10^4 | 0/1 | 1/1 |
| | | | 1×10^5 | 0/3 | 0/3 |
| | | EpCAM ⁻ | 1×10^4 | 0/3 | 0/3 |
| | | | 1×10^5 | 0/2 | 0/2 |
| | | | | | |




Figure 6. Wnt/ β -catenin signaling augments EpCAM⁺ HCC cells. (A) Flow cytometer analysis of HuH1, HuH7, and HLF cells treated with 2 μ mol/L of BIO (orange) or MeBIO (green) for 10 days and stained with anti-EpCAM, anti-CD133 and anti-CD90 antibodies. (B) TOP-FLASH luciferase assays of HuH7 cells treated with 2 μ mol/L of BIO or MeBIO. (C) Flow cytometer analysis of HuH7 cells cultured in normal media (Dulbecco's modified Eagle medium supplemented with 10% FBS) or Wnt10B conditioned media (details are described in the Materials and Methods section). Cells were cultured in each medium for 2 weeks. (D) Representative phase-contrast images (left panel: scale bar, 100 μ m) or IF images (right panel: scale bar, 50 μ m) of HuH7 cells treated with 2 μ mol/L of BIO or MeBIO for 14 days. (E) Quantitative reverse transcription-polymerase chain reaction analysis of representative HCC-related genes in HuH7 cells treated with 2 μ mol/L of BIO or MeBIO for 14 days. (F) Spheroid formation assay of HuH7 cells treated with 2 μ mol/L of BIO or MeBIO for 14 days (mean \pm SD). FITC, fluorescein isothiocyanate.

to maintain HCC stemness and serve as a good marker for HCC initiating cells.

CD133 or CD90 have been used to identify potential hepatic CSCs.^{35,42} CD133 is expressed in normal and malignant stem cells of the neural, hematopoietic, epithelial, hepatic, and endothelial lineages,^{23,43,44} suggesting that CD133 is also a common marker to detect normal cells and CSCs. Captivatingly, EpCAM expression overlaps with CD133 expression in normal human colon tissues and colorectal cancer tissues, yet CD133⁺ and CD133⁻ cells are equally tumorigenic.⁴⁵ Similarly, we found that EpCAM⁺ and EpCAM⁻ HuH1 cells equally expressed CD133, but only EpCAM⁺ cells de-

veloped large hypervascular tumors. Our data suggest that EpCAM may be a better marker than CD133 to enrich HCC tumor-initiating cells from AFP⁺ tumors. We also found that CD90 expression was limited to HCC cell lines that are EpCAM⁻ AFP⁻, and Wnt/ β -catenin signaling had little effect on CD90⁺ cell enrichment. These results suggest that the expression patterns of various stem cell markers in tumor-initiating cells with stem/progenitor cell features may be different in each HCC subtype, possibly owing to the heterogeneity of activated signaling pathways in normal stem/progenitor cells where these tumor-initiating cells may originate. Therefore, it would be useful to




Figure 7. EpCAM blockage inhibits the tumorigenic and invasive capacity of EpCAM⁺ HCC cells. (A) Enrichment of EpCAM⁺ cells after 5-FU treatment. HuH1 cells refer as control or without treatment (green) or treated with 2 μg/mL of 5-FU (orange) for 3 days and analyzed by FACS using anti-EpCAM and anti-CD133 antibodies. (B) Spheroid formation of HuH1 cells treated with 2 μg/mL of 5-FU for 3 days. (C) FACS analysis of HuH1 cells treated with a control siRNA (orange) or EpCAM-specific siRNA (green) at day 3 after transfection. (D) Spheroid formation or (E) invasive capacity of EpCAM⁺ HuH1 cells transfected with a control siRNA or EpCAM-specific siRNA. Experiments were performed in triplicate and the data are shown as mean ± SD. (D) siRNAs. (F) Inhibition of tumor formation in vivo by EpCAM gene silencing. EpCAM⁺ HuH1 cells were transfected with siRNA oligos and 1000 cells were injected 24 hours after transfection.

comprehensively investigate the expression patterns of stem cell markers to characterize the population of CSCs that may correlate with the activation of their distinct molecular pathways.

CSCs may be more resistant to chemotherapeutic agents than differentiated tumor cells possibly owing to an increased expression of adenosine triphosphate-binding cassette transporters and anti-apoptotic proteins.⁴ Thus, the development of an effective strategy to target CSC pools together with conventional chemotherapies is essential to eradicate a tumor mass.¹⁴ By blocking the programs that activate self-renewal and/or inhibit asymmetric division, CSC features could be destemmed.^{46,47} Consistently, EpCAM blockage could inhibit cellular invasion and tumorigenicity of EpCAM⁺ HCC cells, revealing the feasibility of targeting a CSC marker to destem CSC features. EpCAM may induce c-Myc,⁴⁸ a common molecular node activated in HpSC-HCC.²⁷ c-Myc, together with Oct3/4, Sox2, and Klf4, can induce pluripotent stem cells from adult fibroblasts.⁴⁹ It is possible that EpCAM blockage to inhibit hepatic CSCs may

result in a suppression of c-Myc signaling. Encouragingly, EpCAM-specific antibodies are currently in phase II clinical trials.⁵⁰ Furthermore, a recent study indicated that EpCAM⁺ circulating tumor cells identified by a unique microfluidic platform can be used to monitor outcomes of patients undergoing systemic treatment.⁵¹ Therefore, it may be useful to combine EpCAM antibodies with conventional chemotherapy to target both CSCs and non-CSCs for the treatment of HCC.

Supplementary Data

Note: To access the supplementary material accompanying this article, visit the online version of *Gastroenterology* at www.gastrojournal.org, and at doi: 10.1053/j.gastro.2008.12.004.

References

1. Fialkow PJ. Clonal origin of human tumors. *Biochim Biophys Acta* 1976;458:283-321.