

Research Article




Fig. 5. High doses of MK-0608 and telaprevir combination treatment eliminates virus in HCV-infected mice. HCV-infected mice were treated with 50 mg/kg of MK-0608 and 200 mg/kg of telaprevir for 4 weeks. Mice serum samples were obtained at the indicated times, and HCV RNA titer (upper panel) and human serum albumin concentration (lower panel) were analyzed. Points represent the means \pm SD of five mice.

administration of 7000 IU/g/day of IFN- α treatment. However, the virus rebounded after cessation of the therapy.

Four-week high dose combination therapy of MK-0608 and telaprevir eliminated HCV from mice

We investigated whether combination treatment with high doses of MK-0608 and telaprevir without IFN eliminates viruses from HCV-infected mice. Five HCV-infected mice were treated with high doses of MK-0608 (50 mg/kg) and telaprevir (200 mg/kg) for 4 weeks. Serum HCV RNA titer became undetectable 1 week after commencement of the therapy and remained undetectable in all mice at 30 weeks (18 weeks after cessation of the therapy) (Fig. 5). No apparent toxicity of the drugs was observed as none of the mice showed a decrease in the level of serum HSA.

Discussion

Since we began performing treatment experiments using human hepatocyte chimeric mice with HCV, we have administered many different drugs to analyze the effects on suppression or eradication of the virus. However, until we performed the experiments described in this study, we have never observed long term absence of the virus following cessation of the therapy [12,24]. Strikingly, after only 4 weeks of triple therapy with IFN, telaprevir and MK0608, was long term absence of the virus in mouse serum after cessation of the therapy visible (Fig. 4B). Furthermore, high dose telaprevir and MK-0608 combination therapy resulted in a similar absence of the virus for 16 weeks after cessation of therapy (Fig. 5). In this study, mice were treated with 200 mg/kg of

telaprevir twice a day, and 1 week of the treatment resulted in an approximately 2 log reduction of HCV RNA (Fig. 3A), as has been observed previously in chronic hepatitis C patients treated with 450 mg of telaprevir every 8 hrs [25]. This result suggests that approximately 1/15th of a dose in this mouse model may be equivalent to a dose in humans.

During the observation period, some mice died. We do not think that this is due to the drug regimes because the chimeric mouse is weak, and approximately 50% of mice die spontaneously at week 6 after transplantation [26].

Sustained virological response, the complete elimination of the virus from the human body, is defined as testing negative for HCV RNA in serum for more than 24 weeks after cessation of the therapy. As the chimeric mouse used in this study is a weak animal, we were unable to monitor for absence of the virus beyond 24 weeks following cessation of therapy. However, negative testing for HCV RNA in mouse liver by nested PCR (Fig. 4C) 12 weeks after cessation of the therapy strongly suggests that HCV was completely eliminated from the mouse. Of course the mouse model differs from infection in humans where the virus replicates for years in the livers of infected patients. However, results of this study suggest that we will be able to eliminate the virus in humans by treating patients with regimens similar to those used in this study.

Until recently eradication of the virus with biochemical and histological improvement in chronically infected patients has long been reported only with the use of IFN or PEG-IFN [27,28]. Recently, Suzuki et al. reported for the first time eradication of the virus from chronically infected patients without IFN [29].

Elimination of the virus without IFN is desirable due to the many serious side effects of this drug [3,5–9]. However, emergence of drug resistance is a problem, as demonstrated in this study (Fig. 3) as well as in previous studies using replicon systems and HCV-infected chimpanzees [22,23]. A recent clinical study of NS3-4A and NS5B inhibitor combination therapy has reported that 13 days of this combination treatment achieved robust antiviral suppression in chronic hepatitis C patients [30]. As no study has tested the possibility of development of double drug resistant mutants, we will have to test if long term low dose treatment with any combination of STAT-C compounds might induce emergence of multi-drug resistant strains. Furthermore, as there is no report for emergence of IFN resistant strains, regimens such as combination therapy with multiple STAT-C drugs with a small or standard amount of IFN should be tested to develop the best therapy to eradicate the virus with a minimum of side effects and costs. Our further attempts to test possible combinations in mice to determine the best combination of STAT-C drugs will give us an insight into how to develop more effective therapeutic regimens in humans.

Financial support

This study was supported in part by a Grant-in-Aid for Scientific Research from the Japanese Ministry of Labor, Health and Welfare.

Conflict of interest

The authors who have taken part in this study declared that they do not have anything to disclose regarding conflict of interest with respect to this manuscript.

Acknowledgments

The authors thank Rie Akiyama, Kazuyo Hattori, and Yoko Matsumoto for their expert technical help, and Dr. Naoya Sakamoto for providing Huh7/Rep-Neo cells.

References

- [1] Niederau C, Lange S, Heintges T, Erhardt A, Buschkamp M, Hürter D, et al. Prognosis of chronic hepatitis C: results of a large, prospective cohort study. *Hepatology* 1998;28:1687–1695.
- [2] Kiyosawa K, Sodeyama T, Tanaka E, Gibo Y, Yoshizawa K, Nakano Y, et al. Interrelationship of blood transfusion, non-A, non-B hepatitis and hepatocellular carcinoma: analysis by detection of antibody to hepatitis C virus. *Hepatology* 1990;12:671–675.
- [3] Manns MP, McHutchison JG, Gordon SC, Rustgi VK, Shiffman M, Reindollar R, et al. Peginterferon alfa-2b plus ribavirin compared with interferon alfa-2b plus ribavirin for initial treatment of chronic hepatitis C: a randomised trial. *Lancet* 2001;358:958–965.
- [4] Fried MW, Shiffman ML, Reddy KR, Smith C, Marinos G, Goncales Jr FL, et al. Peginterferon alfa-2a plus ribavirin for patients with chronic hepatitis C virus infection. *N Engl J Med* 2002;347:975–982.
- [5] Hoofnagle JH, Ghany MG, Kleiner DE, Doo E, Heller T, Promart K, et al. Maintenance therapy with ribavirin in patients with chronic hepatitis C who fail to respond to combination therapy with interferon alfa and ribavirin. *Hepatology* 2003;38:66–74.
- [6] Schinazi RF, Bassit L, Gavegnano C. HCV drug discovery aimed at viral eradication. *J Viral Hepat* 2010;17:77–90.
- [7] Lin C, Gates CA, Rao BG, Brennan DL, Fulghum JR, Luong YP, et al. In vitro studies of cross-resistance mutations against two hepatitis C virus serine protease inhibitors, VX-950 and BILN 2061. *J Biol Chem* 2005;280:36784–36791.
- [8] Mo H, Lu L, Pilot-Matias T, Pithawalla R, Mondal R, Masse S, et al. Mutations conferring resistance to a hepatitis C virus (HCV) RNA-dependent RNA polymerase inhibitor alone or in combination with an HCV serine protease inhibitor in vitro. *Antimicrob Agents Chemother* 2005;49:4305–4314.
- [9] Björnsson E, Verbaan H, Oksanen A, Frydén A, Johansson J, Friberg S, et al. Health-related quality of life in patients with different stages of liver disease induced by hepatitis C. *Scand J Gastroenterol* 2009;44:878–887.
- [10] Mercer DF, Schiller DE, Elliott JF, Douglas DN, Hao C, Rinfret A, et al. Hepatitis C virus replication in mice with chimeric human livers. *Nat Med* 2001;7:927–933.
- [11] Kneteman NM, Weiner AJ, O'Connell J, Collett M, Gao T, Aukerman L, et al. Anti-HCV therapies in chimeric scid-Alb/uPA mice parallel outcomes in human clinical application. *Hepatology* 2006;43:1346–1353.
- [12] Hiraga N, Imamura M, Tsuge M, Noguchi C, Takahashi S, Iwao E, et al. Infection of human hepatocyte chimeric mouse with genetically engineered hepatitis C virus and its susceptibility to interferon. *FEBS Lett* 2007;581:1983–1987.
- [13] Kimura T, Imamura M, Hiraga N, Hatakeyama T, Miki D, Noguchi C, et al. Establishment of an infectious genotype 1b hepatitis C virus clone in human hepatocyte chimeric mice. *J Gen Virol* 2008;89:2108–2113.
- [14] Kamiya N, Iwao E, Hiraga N, Tsuge M, Imamura M, Takahashi S, et al. Practical evaluation of a mouse with chimeric human liver model for hepatitis C virus infection using an NS3-4A protease inhibitor. *J Gen Virol* 2010;91:1668–1677.
- [15] Lin C, Kwong AD, Perni RB. Discovery and development of VX-950, a novel, covalent, and reversible inhibitor of hepatitis C virus NS3-4A serine protease. *Infect Disord Drug Targets* 2006;6:3–16.
- [16] Migliaccio G, Tomassini JE, Carroll SS, Tomei L, Altamura S, Bhat B, et al. Characterization of resistance to non-obligate chain-terminating ribonucleoside analogs that inhibit hepatitis C virus replication in vitro. *J Biol Chem* 2003;278:49164–49170.
- [17] Guo JT, Bichko VV, Seeger C. Effect of alpha interferon on hepatitis C virus replicon. *J Virol* 2001;75:8516–8523.
- [18] Tanabe Y, Sakamoto N, Enomoto N, Kurosaki M, Ueda E, Maekawa S, et al. Synergistic inhibition of intracellular hepatitis C virus replication by combination of ribavirin and interferon-alpha. *J Infect Dis* 2004;189:1129–1139.
- [19] Yokota T, Sakamoto N, Enomoto N, Tanabe Y, Miyagishi M, Maekawa S, et al. Inhibition of intracellular hepatitis C virus replication by synthetic and vector-derived small interfering RNAs. *EMBO Rep* 2003;4:602–608.
- [20] Tateno C, Yoshizane Y, Saito N, Kataoka M, Utoh R, Yamasaki C, et al. Near completely humanized liver in mice shows human-type metabolic response to drugs. *Am J Pathol* 2004;165:901–912.
- [21] Tsuge M, Hiraga N, Takaishi H, Noguchi C, Oga H, Imamura M, et al. Infection of human hepatocyte chimeric mouse with genetically engineered hepatitis B virus. *Hepatology* 2005;42:1046–1054.
- [22] Kieffer TL, Kwong AD, Picchio GR. Viral resistance to specifically targeted antiviral therapies for hepatitis C (STAT-Cs). *J Antimicrob Chemother* 2010;65:202–212.
- [23] Carroll SS, Ludmerer S, Handt L, Koeplinger K, Zhang NR, Graham D. Robust antiviral efficacy upon administration of a nucleoside analog to hepatitis C virus-infected chimpanzees. *Antimicrob Agents Chemother* 2009;53:926–934.
- [24] Matsumura T, Hu Z, Kato T, Dreux M, Zhang YY, Imamura M, et al. Amphipathic DNA polymers inhibit hepatitis C virus infection by blocking viral entry. *Gastroenterology* 2009;137:673–681.
- [25] Reesink HW, Zeuzem S, Weegink CJ, Forestier N, van Vliet A, de Rooij J, et al. Rapid decline of viral RNA in hepatitis C patients treated with VX-950: a phase Ib, placebo-controlled, randomized study. *Gastroenterology* 2006;131:997–1002.
- [26] Vanwolleghem T, Libbrecht L, Hansen BE, Desombere I, Roskams T, Meuleman P, et al. Factors determining successful engraftment of hepatocytes and susceptibility to hepatitis B and C virus infection in uPA-SCID mice. *J Hepatol* 2010;53:468–476.
- [27] Chayama K, Saitoh S, Arase Y, Ikeda K, Matsumoto T, Sakai Y, et al. Effect of interferon administration on serum hepatitis C virus RNA in patients with chronic hepatitis C. *Hepatology* 1991;13:1040–1043.
- [28] Poynard T, McHutchison J, Manns M, Trepo C, Lindsay K, Goodman Z, et al. Impact of pegylated interferon alfa-2b and ribavirin on liver fibrosis in patients with chronic hepatitis C. *Gastroenterology* 2002;122:1303–1313.
- [29] Suzuki F, Suzuki Y, Akuta N, Sezaki H, Yatsuji H, Arase Y, et al. Sustained virological response in a patient with chronic hepatitis C treated by monotherapy with the NS3-4A protease inhibitor telaprevir. *J Clin Virol* 2010;47:76–78.
- [30] Gane E, Roberts S, Stedman C, Angus P, Ritchie B, Elston R, et al. 749 Early on-treatment responses during pegylated IFN plus rivavirin are increased following 13 days of combination nucleoside polymerase (RG7128) and protease (RG7227) inhibitor therapy (INFORM-1). *J Hepatol* 2010;51:S291–S292.

MECHANISMS OF GASTROINTESTINAL, PANCREATIC AND LIVER DISEASES

Animal model for study of human hepatitis virusesKazuaki Chayama,^{*,†} C Nelson Hayes,^{*,†} Nobuhiko Hiraga,^{*,†} Hiromi Abe,^{*,†} Masataka Tsuge^{*,†}
and Michio Imamura^{*,†}

*Department of Medicine and Molecular Science, Division of Frontier Medical Science, Programs for Biomedical Research, Graduate School of Biomedical Sciences, Hiroshima University, and [†]Liver Research Project Center, Hiroshima, Japan

Key words

hepatitis B virus, hepatitis C virus, uPA/scid mouse model.

Accepted for publication 26 July 2010.

Correspondence

Professor Kazuaki Chayama, Department of Medical and Molecular Science, Division of Frontier Medical Science, Programs for Biomedical Research, Graduate school of Biomedical Science, Hiroshima University, 1-2-3 Kasumi, Minami-ku, Hiroshima 734-8551, Japan. Email: chayama@hiroshima-u.ac.jp

Abstract

Human hepatitis B virus (HBV) and hepatitis C virus (HCV) infect only chimpanzees and humans. Analysis of both viruses has long been hampered by the absence of a small animal model. The recent development of human hepatocyte chimeric mice has enabled us to carry out studies on viral replication and cellular changes induced by replication of human hepatitis viruses. Various therapeutic agents have also been tested using this model. In the present review, we summarize published studies using chimeric mice and discuss the merits and shortcomings of this model.

Introduction

Hepatitis B virus (HBV) and hepatitis C virus (HCV) are pathogens that cause chronic infection in humans. There are 360 million and 170 million people infected worldwide with HBV or HCV, respectively.^{1,2} Infected individuals develop acute hepatitis, chronic hepatitis and liver cirrhosis. The viruses are also important causative agents of hepatocellular carcinoma, especially in the Asia-Pacific region.³ Study of the biology and development of therapies for each virus has long been hampered by the lack of a small animal model that supports hepatitis virus infection. This is probably as a result of the lack of receptor molecules necessary for viral infection in animal liver cells.

Transgenic mice that express over-length HBV-DNA export viral particles into the serum,⁴ and such animals can be used to evaluate antiviral agents,⁵⁻⁷ as well as HBV-targeted siRNA⁸. However, the virus life cycle is not established in this model, and it is inappropriate for studying drug-resistant HBV strains. Accordingly, researchers attempted to transplant human hepatocytes into mice. The development of the trimera mouse was one such attempt, in which human hepatocytes were transplanted under the kidney capsule of immune-deficient mice after lethal irradiation.^{9,10} However, the number of hepatocytes that could survive on the kidney capsule was small, and normal liver architecture was not present. Although 85% of HBV-inoculated animals developed HBV viremia, the titer was less than 10⁵ virus particles or IU/mL.⁹ Similarly, 85% of HCV-inoculated animals also developed viremia,¹⁰ but the level of the viremia only reached 10²/mL.

Thus, the advent of human hepatocyte transplanted uPA/scid mice has provided the first really useful model for acute and chronic infections of human hepatitis virus.

Human liver cell transplanted uPA/scid mice

Transgenic mice in which the urokinase gene is driven by the human albumin promoter/enhancer were developed and shown to have accelerated hepatocyte death and consequent chronic stimulation of hepatocyte growth.¹¹ Transplanted rat hepatocytes proliferated and repopulated injured livers in immunodeficient uPA mice, which were produced by mating uPA transgenic mice with scid mice.¹² Human hepatocytes were then transplanted into uPA/scid mice; these cells proliferated and replaced the apoptotic mice liver cells (Fig. 1).

Such human hepatocyte chimeric mice have been shown to be susceptible to both HBV¹⁶ and HCV¹⁷ infections. Repopulation levels by human hepatocytes have been estimated by measuring human albumin levels in mouse serum. Replication levels of both HBV¹³ and HCV¹⁷ were higher in mice in which the repopulation index was higher. A unique attempt to remove mouse residual liver cells with the herpes simplex virus type-1 thymidine kinase (HSVtk)/ganciclovir (GCV) system failed to result in a higher repopulation rate as a result of damage to the transplanted human hepatocyte caused by bystander effects.¹⁸ Despite this, mice with livers that have been highly repopulated with human hepatocytes




Figure 1 Generation of human hepatocyte chimeric mice and hepatitis virus infection model. A uPA/scid mouse was created by mating uPA transgenic mouse and scid mouse. Human hepatocytes obtained by surgical resection or donation were transplanted to newborn mice. The chimeric mice can be infected with hepatitis B virus (HBV) or hepatitis C (HCV) virus by injecting human serum containing these viruses. Alternatively, the mice can be infected by HBV¹³ or HCV¹⁴ created in cell culture or by injecting HCV RNA into the mouse liver.¹⁵

are susceptible to infection with both HBV and HCV, and as such comprised the most effective small animal model for chronic hepatitis so far developed.^{19,20} An example of a highly repopulated mouse liver that we are using in experiments is shown in Figure 2.

Highly repopulated mice have been shown to be a valuable model for the study of drug metabolism.^{21–29} Advances in technology for human hepatocyte transplantation have enabled serial passage of human hepatocytes in uPA/scid mice and have been shown to retain infectivity for HBV.³⁰

This mouse model and other animal models for the study of hepatitis viruses have been summarized in reviews by Meuleman and Leroux-Roels,³¹ Dandri *et al.*,^{32,33} Barth *et al.*,³⁴ and Kneteman and Toso.³⁵ The present review will focus on key issues and updated information.

Study of hepatitis B virus infection using human hepatocyte chimeric mice

Since the initial reports of successful transmission of HBV to human hepatocyte chimeric mice in 2001 and 2004,^{16,27} several researchers have reported transmission of HBV into similar

mice.^{13,36,37} In these studies, passage experiments studies show that HBV replicating in mice retain infectivity.^{13,36} Further, the presence of viral proteins has been shown immunohistochemically in human hepatocytes transplanted into mouse livers, but these are not present in mouse hepatocytes.^{13,36,37} Formation of viral particles in infected mouse livers can be shown by electron microscopy.^{36,37} Genetically engineered viruses lacking HBc-antigen have also been shown to infect chimeric mice, proving that c antigen is dispensable for viral infection and replication.¹³ In contrast, HBx protein has been shown to be indispensable for viral replication.³⁸ Transcomplementation of HBx protein with hydrodynamic injection restored HBV infectivity in mice. Interestingly, all revertant viruses show a restored ability to express HBx.³⁸

By infecting chimeric mice with genotype A, B and C, differing proliferative capacity has been shown between HBV genotypes.³⁷ In mice infected for a relatively short time, there are no morphological changes in HBV infected mice livers in studies.^{13,36} In contrast, the occurrence of liver cell damage has been reported after long-term infection of chimeric mice with HBV³⁹ or with specific strains of HBV;⁴⁰ these findings are consistent with direct cytopathic effects of HBV under certain conditions.




Figure 2 Representative uPA/scid mouse livers repopulated by human hepatocytes. (a) Mouse liver almost completely repopulated by human hepatocytes. Only a small portion of mouse hepatocytes are shown by arrows. (b) Microscopic figure of the mouse liver. M and H indicate regions consisting of mouse and human hepatocytes, respectively (Hematoxylin-eosin staining, magnification: $\times 100$). (c) Microscopic figure of the mouse liver stained with antibody directed against human serum albumin.

The biological properties of a newly identified unique strain of HBV, genotype G, which replicates only in the presence of another genotype, were confirmed using the chimeric mouse.⁴¹ Infectivity of another novel HBV strain, identified from a Japanese patient, that is divergent from known human and ape HBV has also been confirmed.⁴² Titration of HBV infectivity, which previously could only be carried out using chimpanzees, can be carried out effectively using chimeric mice.⁴³

Taking advantage of the absence of human immune cells in the chimeric mice, Noguchi *et al.*⁴⁴ showed that hypermutation of HBV increases in human hepatocytes under interferon treatment. Dandri *et al.* measured viral half-life in human and chimeric mice repopulated with woolly monkey hepatocytes.⁴⁵ The results clearly showed that viral half-life is shortened by immunological mechanisms in humans with low viral levels, but not in chimeric mice where functional immunity is absent. Hiraga *et al.*⁴⁶ showed an absence of interference between HBV and HCV.

Evaluation of therapeutic agents is the most important role for this mouse model. Tsuge *et al.*¹³ assessed the effect of interferon and lamivudine using chimeric mice. Similarly, Dandri *et al.*⁴⁷ showed the effects of adefovir using uPA/scid mice repopulated with tupaia hepatocytes, which also support replication of human HBV. Oga *et al.*⁴⁸ identified a novel lamivudine-resistant variant that has an amino acid substitution outside of the YMDD motif. They showed that lamivudine was ineffective against the novel mutant strain. It is thus apparent that this mouse/human liver chimeric model is ideal to study the susceptibility of mutant strains to various drugs, because mutant viruses can easily be made and infected into chimeric mice.¹³ The model has also been utilized to evaluate viral entry inhibitors derived from the large envelope protein.⁴⁹

Study of hepatitis C virus using human hepatocyte chimeric mice

As observed in studies on HBV, HCV infection efficiency was poor and levels of viremia were low in mice where the repopulation rate of the mouse liver with human hepatocyte was low.^{17,50} As shown in Figure 3, human albumin levels in mouse serum were significantly higher in mice in which measurable viremia developed (Hiraga *et al.* unpublished data). Recent studies have therefore been carried out using highly repopulated mice. The usefulness of a newly developed HCV assay,⁵¹ and infectivity of a newly identified intergenotypic recombinant strain,⁵² have been reported using the chimeric mice.

Using the remarkable replication ability of the JFH1 genotype 2a strain,⁵³ infectivity of JFH1 or intergenotypic chimeric viral particles, previously shown in cell culture, has now been shown to be infectious in chimeric mice.⁵⁴⁻⁵⁶ Infectivity of viruses that were replicated in chimeric mice in cell culture has also been shown, and virus fitness has been studied.^{55,56} The role of the HCV core+1 open reading frame and core *cis*-acting RNA elements has also been examined using the chimeric virus.⁵⁷ These elegant studies have the limitation that the non-structural part of the virus is limited to that of JFH1. Hiraga *et al.*¹⁴ have shown that infectious clones of genotype 1a and JFH1 can be infected with direct injection of *in vitro* transcribed RNA into the mouse liver.¹⁴ Similarly, Kimura *et al.*¹⁵ reported the establishment of infectious clones of genotype 1b and ablation of RNA polymerase by site-directed mutagenesis abolish infectivity. These infectious clones will be useful for the study of drug-resistant strains.

The model of HCV infection has also been used to show that infection of the virus can be prevented by antibodies against

Table 1 New therapeutic strategies tested by human hepatocyte chimeric mice

<i>n</i>	Drug or cell	Strategy	Reference
1	Interferon alpha 2b BILN-2061 HCV371	Activation of antiviral genes NS3-4A protease inhibition NS5B polymerase inhibition	Kneteman <i>et al.</i> ⁶⁵
2	Modified BID	Induction of apoptosis	Hsu <i>et al.</i> ⁶⁶
3	Serine palmitoyltransferase inhibitor	Disruption of lipid raft	Umehara <i>et al.</i> ⁶⁷
4	Lymphoblastoid interferon alpha	Activation of antiviral genes	Hiraga <i>et al.</i> ¹⁴
5	Amphipathic DNA polymers	Blocking viral entry	Matsumura <i>et al.</i> ⁶⁸
6	Sec-butyl-analogue of HCV-371	NS5B polymerase inhibition	LaPorte <i>et al.</i> ⁶⁸
7	HCV796	NS5B polymerase inhibition	Kneteman <i>et al.</i> ⁶⁹
8	Liver allograft-derived lymphocyte	Adoptive immunotherapy	Ohira <i>et al.</i> ⁷⁰
9	Telaprevir	NS3-4A protease inhibition	Kamiya <i>et al.</i> ⁷¹




Figure 3 Human albumin levels in mice used in the hepatitis C virus (HCV) infection experiments. A total of 54 mice were injected with HCV positive serum samples containing 5×10^6 virus particles. A total of 24 mice became persistently positive for HCV-RNA, but 30 mice did not. Serum human albumin levels 2 weeks after human hepatocyte transplantation were compared between infected and non-infected mice.

CD81,⁵⁸ polyclonal human immunoglobulin directed to a similar strain,⁵⁹ and amphipathic DNA polymers.⁶⁰ Notably, the presence of broadly neutralizing antibodies to HCV that protect against heterologous viral infection has been reported, suggesting the possibility of a prophylactic vaccine against HCV.⁶¹

With respect to evasion of the virus against the innate immune response, altered intrahepatic expression profiles in the early phase of infection is of particular interest. The chimeric mice model is ideal for such studies; cross-hybridization of mouse and human can be avoided by careful experimental procedures.⁶² Microarray analysis of livers of HCV infected and non-infected mice showed transcriptional activation of genes related to innate immune response, lipid metabolism, endoplasmic reticulum (ER) stress and apoptosis in HCV-infected mice.^{63,64} The HCV infected mouse model is particularly useful for the study of newly developed HCV agents. The effect of recently developed chemicals and a unique therapy using intrahepatic lymphocytes have been shown using

this model (Table 1). However, none of these therapies have yet been able to completely eradicate HCV from mice. It is noteworthy that ultra-rapid cardiotoxicity has been reported with the protease inhibitor BILN 2061 in the uPA/scid mice, but not in scid mice, implicating involvement of the uPA transgene.⁷² Care should therefore be taken in interpreting the results obtained by this model.

Conclusion

Development of a small animal model using human hepatocyte chimeric mice has enabled us to study key aspects of HBV and HCV biology. The characteristic feature of the absence of human immune cells is suitable for studying viral replication and observing changes occurring in liver cells during viral infection, such as the innate immune response and cellular stress and metabolic responses. The model is also useful for studying the effect of drugs without the influence of cytokines and cytotoxic T lymphocytes. Nonetheless, the model is insufficient to study carcinogenesis of hepatitis viruses, because non-parenchymal cells in mouse liver are of mouse origin and do not support inflammation and fibrosis, which are probably closely related to carcinogenesis. The lack of human immune cells also limits the study of inflammation and immunity. Furthermore, the availability of human hepatocytes is limited. Despite these limitations, the current model shows great potential as a mouse model for the study of hepatitis viruses. Development of a small animal model with or without human immunity using stem cells or iPS cells would be an ideal model in the future.

Acknowledgments


This work was supported in part by Grants-in-Aid for scientific research and development from the Ministry of Education, Culture, Sports, Science and Technology, and the Ministry of Health, Labor and Welfare, Government of Japan.

References

- Shepard CW, Simard EP, Finelli L, Flore AE, Bell BP. Hepatitis B virus infection: epidemiology and vaccination. *Epidemiol. Rev.* 2006; **28**: 112–25.

- 2 Sy T, Jamal MM. Epidemiology of Hepatitis C Virus (HCV) infection. *Int. J. Med. Sci.* 2006; **3**: 41–6.
- 3 Yuen MF, Hou JL, Chutaputti A, Prevent APWP. Hepatocellular carcinoma in the Asia pacific region. *J. Gastroenterol. Hepatol.* 2009; **24**: 346–53.
- 4 Guidotti LG, Matzke B, Schaller H, Chisari FV. High-level Hepatitis-B Virus-replication in transgenic mice. *J. Virol.* 1995; **69**: 6158–69.
- 5 Weber O, Schlemmer KH, Hartmann E *et al.* Inhibition of human hepatitis B virus (HBV) by a novel non-nucleosidic compound in a transgenic mouse model. *Antiviral Res.* 2002; **54**: 69–78.
- 6 Julander JG, Sidwell RW, Morrey JD. Characterizing antiviral activity of adefovir dipivoxil in transgenic mice expressing hepatitis B virus. *Antiviral Res.* 2002; **55**: 27–40.
- 7 Julander JG, Colonna RJ, Sidwell RW, Morrey JD. Characterization of antiviral activity of entecavir in transgenic mice expressing hepatitis B virus. *Antiviral Res.* 2003; **59**: 155–61.
- 8 Uprichard SL, Boyd B, Althage A, Chisari FV. Clearance of hepatitis B virus from the liver of transgenic mice by short hairpin RNA. *Proc. Natl. Acad. Sci. U S A* 2005; **102**: 773–8.
- 9 Ilan E, Burakova T, Dagan S *et al.* The hepatitis B virus-trimer mouse: a model for human HBV infection and evaluation of Anti-HBV therapeutic agents. *Hepatology* 1999; **29**: 553–62.
- 10 Ilan E, Arazi J, Nussbaum O *et al.* The hepatitis C virus (HCV)-Trimer mouse: a model for evaluation of agents against HCV. *J. Infect. Dis.* 2002; **185**: 153–61.
- 11 Heckel JL, Sandgren EP, Degen JL, Palmiter RD, Brinster RL. Neonatal bleeding in transgenic mice expressing urokinase-type plasminogen-activator. *Cell* 1990; **62**: 447–56.
- 12 Rhim JA, Sandgren EP, Palmiter RD, Brinster RL. Complete reconstitution of mouse-liver with xenogeneic hepatocytes. *Proc. Natl. Acad. Sci. U S A* 1995; **92**: 4942–6.
- 13 Tsuge M, Hiraga N, Takaishi H *et al.* Infection of human hepatocyte chimeric mouse with genetically engineered hepatitis B virus. *Hepatology* 2005; **42**: 1046–54.
- 14 Hiraga N, Imamura M, Tsuge M *et al.* Infection of human hepatocyte chimeric mouse with genetically engineered Hepatitis C virus and its susceptibility to interferon. *FEBS Lett.* 2007; **581**: 1983–7.
- 15 Kimura T, Imamura M, Hiraga N *et al.* Establishment of an infectious genotype 1b Hepatitis C Virus clone in human hepatocyte chimeric mice. *J. Gen. Virol.* 2008; **89**: 2108–13.
- 16 Dandri M, Burda MR, Torok E *et al.* Repopulation of mouse liver with human hepatocytes and in vivo infection with hepatitis B virus. *Hepatology* 2001; **33**: 981–8.
- 17 Mercer DF, Schiller DE, Elliott JF *et al.* Hepatitis C virus replication in mice with chimeric human livers. *Nat. Med.* 2001; **7**: 927–33.
- 18 Douglas DN, Kawahara T, Sis B *et al.* therapeutic efficacy of human hepatocyte transplantation in a SCID/uPA mouse model with inducible liver disease. *PLoS ONE* 2010; **5**: e9209.
- 19 Tateno C, Yoshizane Y, Saito N *et al.* Near completely humanized liver in mice shows human-type metabolic responses to drugs. *Am. J. Pathol.* 2004; **165**: 901–12.
- 20 Bissig KD, Wieland SF, Tran P *et al.* Human liver chimeric mice provide a model for hepatitis B and C virus infection and treatment. *J. Clin. Invest.* 2010; **120**: 924–30.
- 21 Yu AM, Idle JR, Gonzalez FJ. Polymorphic cytochrome p450 2D6: humanized mouse model and endogenous substrates. *Drug. Metab. Rev.* 2004; **36**: 243–77.
- 22 Katoh M, Sawada T, Soeno Y *et al.* In vivo drug metabolism model for human cytochrome P450 enzyme using chimeric mice with humanized liver. *J. Pharm. Sci.-Us.* 2007; **96**: 428–37.
- 23 Katoh M, Matsui T, Nakajima M *et al.* In vivo induction of human cytochrome P450 enzymes expressed in chimeric mice with humanized liver. *Drug. Metab. Dispos.* 2005; **33**: 754–63.
- 24 Katoh M, Matsui T, Okumura H *et al.* Expression of human phase II enzymes in chimeric mice with humanized liver. *Drug. Metab. Dispos.* 2005; **33**: 1333–40.
- 25 Okumura H, Katoh M, Sawada T *et al.* Humanization of excretory pathway in chimeric mice with humanized liver. *Toxicol. Sci.* 2007; **97**: 533–8.
- 26 Shoda J, Okada K, Inada Y *et al.* Bezafibrate induces multidrug-resistance P-Glycoprotein 3 expression in cultured human hepatocytes and humanized livers of chimeric mice. *Hepatology Res.* 2007; **37**: 548–56.
- 27 Petersen J, Burda MR, Dandri M, Rogler CE. Transplantation of human hepatocytes in immunodeficient UPA mice: a model for the study of hepatitis B virus. *Methods Mol. Med.* 2004; **96**: 253–60.
- 28 Yoshizato K, Tateno C. A human hepatocyte-bearing mouse: an animal model to predict drug metabolism and effectiveness in humans. *PPAR Res.* 2009; **2009**: 476217.
- 29 Yoshizato K, Tateno C, Utoh R. The mechanism of liver size control in mammals: a novel animal study. *Int. J. Design & Nature Ecodynamics* 2009; **4**: 123–42.
- 30 Utoh R, Tateno C, Yamasaki C *et al.* Susceptibility of chimeric mice with livers repopulated by serially subcultured human hepatocytes to hepatitis B virus. *Hepatology* 2008; **47**: 435–46.
- 31 Meuleman P, Leroux-Roels G. The human liver-uPA-SCID mouse: a model for the evaluation of antiviral compounds against HBV and HCV. *Antiviral Res.* 2008; **80**: 231–8.
- 32 Dandri M, Lutgehetmann M, Volz T, Petersen J. Small animal model systems for studying Hepatitis B Virus replication and pathogenesis. *Semin. Liver Dis.* 2006; **26**: 181–91.
- 33 Dandri M, Volz TK, Lutgehetmann M, Petersen J. Animal models for the study of HBV replication and its variants. *J. Clin. Virol.* 2005; **34** (Suppl. 1): S54–62.
- 34 Barth H, Robinet E, Liang TJ, Baumert TF. Mouse models for the study of HCV infection and virus-host interactions. *J. Hepatol.* 2008; **49**: 134–42.
- 35 Kneteman NM, Toso C. In vivo study of HCV in mice with chimeric human livers. *Methods Mol. Biol.* 2009; **510**: 383–99.
- 36 Meuleman P, Libbrecht L, De Vos R *et al.* Morphological and biochemical characterization of a human liver in a uPA-SCID mouse chimera. *Hepatology* 2005; **41**: 847–56.
- 37 Sugiyama M, Tanaka Y, Kato T *et al.* Influence of hepatitis B virus genotypes on the intra- and extracellular expression of viral DNA and antigens. *Hepatology* 2006; **44**: 915–24.
- 38 Tsuge M, Hiraga N, Akiyama R *et al.* HBx protein is indispensable for development of viremia in human hepatocyte chimeric mice. *J. Gen. Virol.* 2010.
- 39 Meuleman P, Libbrecht L, Wieland S *et al.* Immune suppression uncovers endogenous cytopathic effects of the hepatitis B virus. *J. Virol.* 2006; **80**: 2797–807.
- 40 Sugiyama M, Tanaka Y, Kurbanov F *et al.* Direct cytopathic effects of particular hepatitis B virus genotypes in severe combined immunodeficiency transgenic with urokinase-type plasminogen activator mouse with human hepatocytes. *Gastroenterology* 2009; **136**: 652–62.
- 41 Tanaka Y, Sanchez LV, Sugiyama M *et al.* Characteristics of Hepatitis B Virus genotype G coinfecting with genotype H in chimeric mice carrying human hepatocytes. *Virology* 2008; **376**: 408–15.
- 42 Tatematsu K, Tanaka Y, Kurbanov F *et al.* A genetic variant of Hepatitis B Virus divergent from known human and ape genotypes isolated from a Japanese patient and provisionally assigned to new genotype J. *J. Virol.* 2009; **83**: 10538–47.
- 43 Tabuchi A, Tanaka J, Katayama K *et al.* Titration of Hepatitis B Virus infectivity in the sera of pre-acute and late acute phases of HBV infection: transmission experiments to chimeric mice with

- human liver repopulated hepatocytes. *J. Med. Virol.* 2008; **80**: 2064–8.
- 44 Noguchi C, Imamura M, Tsuge M *et al.* G-to-A Hypermutation in Hepatitis B Virus (HBV) and clinical course of patients with chronic HBV infection. *J. Infect. Dis.* 2009; **199**: 1599–607.
 - 45 Dandri M, Murray JM, Lutgehetmann M, Volz T, Lohse AW, Petersen J. Virion half-life in chronic hepatitis B infection is strongly correlated with levels of viremia. *Hepatology* 2008; **48**: 1079–86.
 - 46 Hiraga N, Imamura M, Hatakeyama T *et al.* Absence of viral interference and different susceptibility to interferon between Hepatitis B Virus and Hepatitis C Virus in human hepatocyte chimeric mice. *J. Hepatol.* 2009; **51**: 1046–54.
 - 47 Dandri M, Burda MR, Zuckerman DM *et al.* Chronic infection with Hepatitis B Viruses and antiviral drug evaluation in uPA mice after liver repopulation with tupaia hepatocytes. *J. Hepatol.* 2005; **42**: 54–60.
 - 48 Yatsuji H, Noguchi C, Hiraga N *et al.* Emergence of a novel lamivudine-resistant hepatitis B virus variant with a substitution outside the YMDD motif. *Antimicrob. Agents Chemother.* 2006; **50**: 3867–74.
 - 49 Petersen J, Dandri M, Mier W *et al.* Prevention of Hepatitis B Virus infection in vivo by entry inhibitors derived from the large envelope protein. *Nat. Biotechnol.* 2008; **26**: 335–41.
 - 50 Turrini P, Sasso R, Germoni S *et al.* Development of humanized mice for the study of hepatitis C virus infection. *Transplant. Proc.* 2006; **38**: 1181–4.
 - 51 Cagnon L, Wagaman P, Bartschlagler R *et al.* Application of the trak-C (TM) HCV core assay for monitoring antiviral activity in HCV replication systems. *J. Virol. Methods* 2004; **118**: 23–31.
 - 52 Kurbanov F, Tanaka Y, Chub E *et al.* Molecular epidemiology and interferon susceptibility of the natural recombinant Hepatitis C Virus Strain RF1-2k/1b. *J. Infect. Dis.* 2008; **198**: 1448–56.
 - 53 Wakita T, Pietschmann T, Kato T *et al.* Production of infectious hepatitis C virus in tissue culture from a cloned viral genome. *Nat. Med.* 2005; **11**: 791–6.
 - 54 Grove J, Huby T, Stamataki Z *et al.* Scavenger receptor BI and BII expression levels modulate Hepatitis C Virus infectivity. *J. Virol.* 2007; **81**: 3162–9.
 - 55 Lindenbach BD, Meuleman P, Ploss A *et al.* Cell culture-grown Hepatitis C Virus is infectious in vivo and can be recultured in vitro. *Proc. Natl. Acad. Sci. U S A* 2006; **103**: 3805–9.
 - 56 Kaul A, Woerz I, Meuleman P, Leroux-Roels G, Bartschlagler R. Cell culture adaptation of Hepatitis C Virus and in vivo viability of an adapted variant. *J. Virol.* 2007; **81**: 13168–79.
 - 57 Vassilaki N, Friebe P, Meuleman P *et al.* Role of the Hepatitis C Virus Core+1 open reading frame and core cis-acting RNA Elements in Viral RNA translation and replication. *J. Virol.* 2008; **82**: 11503–15.
 - 58 Meuleman P, Hesselgesser J, Paulson M *et al.* Anti-CD81 antibodies can prevent a Hepatitis C Virus infection in vivo. *Hepatology* 2008; **48**: 1761–8.
 - 59 Vanwolleghem T, Bukh J, Meuleman P *et al.* Polyclonal immunoglobulins from a chronic Hepatitis C Virus patient protect human liver-chimeric mice from infection with a homologous Hepatitis C Virus strain. *Hepatology* 2008; **47**: 1846–55.
 - 60 Matsumura T, Hu ZY, Kato T *et al.* Amphipathic DNA polymers inhibit Hepatitis C Virus infection by blocking viral entry. *Gastroenterology* 2009; **137**: 673–81.
 - 61 Law M, Maruyama T, Lewis J *et al.* Broadly neutralizing antibodies protect against Hepatitis C Virus quasispecies challenge. *Nat. Med.* 2008; **14**: 25–7.
 - 62 Walters KA, Joyce MA, Thompson JC *et al.* Application of functional genomics to the chimeric mouse model of HCV infection: optimization of microarray protocols and genomics analysis. *Virol. J.* 2006; **3**: 37–44.
 - 63 Walters KA, Joyce MA, Thompson JC *et al.* Host-specific response to HCV infection in the chimeric SCID-beige/Alb-uPA mouse model: role of the innate antiviral immune response. *PLoS Pathog.* 2006; **2**: 591–602.
 - 64 Joyce MA, Walters KA, Lamb SE *et al.* HCV Induces Oxidative and ER Stress, and Sensitizes Infected Cells to Apoptosis in SCID/Alb-uPA Mice. *PLoS Pathog.* 2009; **5**: e10000291.
 - 65 Kneteman NM, Weiner AJ, O'Connell J *et al.* Anti-HCV therapies in chimeric scid-Alb/uPA mice parallel outcomes in human clinical application. *Hepatology* 2006; **43**: 1346–53.
 - 66 Hsu EC, Hsi B, Hirota-Tsuchihara M *et al.* Modified apoptotic molecule (BID) reduces hepatitis C virus infection in mice with chimeric human livers. *Nat. Biotechnol.* 2003; **21**: 519–25.
 - 67 Umehara T, Sudoh M, Yasui F *et al.* Serine palmitoyltransferase inhibitor suppresses HCV replication in a mouse model. *Biochem. Biophys. Res. Commun.* 2006; **346**: 67–73.
 - 68 Laporte MG, Jackson RW, Draper TL *et al.* The discovery of pyrano[3,4-b]indole-based allosteric inhibitors of HCV NS5B polymerase with in vivo activity. *Med. Chem.* 2008; **3**: 1508–15.
 - 69 Kneteman NM, Howe AYM, Gao TJ *et al.* HCV796: a selective nonstructural protein 5B polymerase inhibitor with potent Anti-Hepatitis C Virus activity in vitro, in mice with chimeric human livers, and in humans infected with Hepatitis C Virus. *Hepatology* 2009; **49**: 745–52.
 - 70 Ohira M, Ishiyama K, Tanaka Y *et al.* Adoptive immunotherapy with liver allograft-derived lymphocytes induces anti-HCV activity after liver transplantation in humans and humanized mice. *J. Clin. Invest.* 2009; **119**: 3226–35.
 - 71 Kamiya N, Iwao E, Hiraga N *et al.* Practical Evaluation of a Mouse with Chimeric Human Liver Model for Hepatitis C Virus Infection Using an NS3-4A Protease Inhibitor. *J. Gen. Virol.* 2010; **91**: 1668–77.
 - 72 Vanwolleghem T, Meuleman P, Libbrecht L *et al.* Ultra-rapid cardiotoxicity of the hepatitis C virus protease inhibitor BILN 2061 in the urokinase-type plasminogen activator mouse. *Gastroenterology* 2007; **133**: 1144–55.



IL-6-mediated intersubgenotypic variation of interferon sensitivity in hepatitis C virus genotype 2a/2b chimeric clones

Goki Suda ^{a,1}, Naoya Sakamoto ^{a,b,*}, Yasuhiro Itsui ^{a,d}, Mina Nakagawa ^{a,b}, Megumi Tasaka-Fujita ^a, Yusuke Funaoka ^a, Takako Watanabe ^a, Sayuri Nitta ^a, Kei Kiyohashi ^a, Seishin Azuma ^a, Sei Kakinuma ^a, Kiichiro Tsuchiya ^a, Michio Imamura ^c, Nobuhiko Hiraga ^c, Kazuaki Chayama ^c, Mamoru Watanabe ^a

^a Department of Gastroenterology and Hepatology, Tokyo Medical and Dental University, Tokyo, Japan

^b Department for Hepatitis Control, Tokyo Medical and Dental University, Tokyo, Japan

^c Department of Medicine and Molecular science, Division of Frontier Medical Sciences, Programs for Biomedical Research, Graduate School of Biomedical Sciences, Hiroshima University, Hiroshima, Japan

^d Department of Internal Medicine, Soka Municipal Hospital, Saitama, Japan

ARTICLE INFO

Article history:

Received 25 May 2010

Returned to author for revision 19 June 2010

Accepted 26 July 2010

Available online 25 August 2010

Keywords:

HCV-JFH1

Inter-genotypic chimeric virus

SOCS3

Interleukin-6

ABSTRACT

Mechanisms of difference in interferon sensitivity between hepatitis C virus (HCV) strains have yet to be clarified. Here, we constructed an infectious genotype2b clone and analyzed differences in interferon- α sensitivity between HCV-2b and 2a-JFH1 clones using intergenotypic homologous recombination. The HCV-2b/JFH1 chimeric virus able to infect Huh7.5.1 cells and was significantly more sensitive to IFN than JFH1. IFN-induced expression of MxA and 25-OAS was significantly lower in JFH1 than in 2b/JFH1-infected cells. In JFH1-infected cells, expression of SOCS3 and its inducer, IL-6, was significantly higher than in 2b/JFH1-infected cells. The IFN-resistance of JFH1 cells was negated by siRNA-knock down of SOCS3 expression and by pretreatment with anti-IL6 antibody. In conclusion, intergenotypic differences of IFN sensitivity of HCV may be attributable to the sequences of HCV structural proteins and can be determined by SOCS3 and IL-6 expression levels.

© 2010 Elsevier Inc. All rights reserved.

Introduction

Hepatitis C virus (HCV) is one of the most important pathogens causing liver-related morbidity and mortality (Alter, 1997). There is no therapeutic or prophylactic vaccine available for HCV and type I interferons have been the mainstay of HCV therapeutics (Hoofnagle and di Bisceglie, 1997). Antiviral therapeutic options against HCV are limited and yield unsatisfactory responses (Fried et al., 2002). Given these situations, gaining a detailed understanding of the molecular mechanisms of interferon resistance has been a high priority in academia and industry.

Molecular studies of HCV have been hampered by the lack of efficient *in vitro* and *in vivo* models of infection, which has been partly overcome by the development of HCV subgenomic replicons (Blight et al., 2000; Kato et al., 2003; Lohmann et al., 1999) and the HCV-JFH1 cell culture system

(Wakita et al., 2005). HCV-JFH1 is an isolate of HCV genotype 2a that was obtained from a patient with fulminant hepatitis C. The full-length JFH1 genome has been shown to produce infectious particles in cell culture. Simultaneously, a robustly replicating intragenotypic chimera has been reported, which consists of the structural region of a genotype 2a, J6-clone and nonstructural region of JFH-1 (Lindenbach et al., 2005).

HCV isolates are classified into seven major genotypes and multiple subtypes (Gottwein et al., 2009). In infected individuals, HCV exists as quasispecies of closely related genomes (Bukh et al., 1995). A number of studies have suggested that the outcome of HCV infection, as well as the response to interferon treatment, depends on the genotype or quasispecies with which the patient is infected. However, it is not clear how these subtle genetic differences of HCV affect viral replication, infectivity and host responses. Thus, it is important to establish multiple cell culture-permissive strains of different genotypes and isolates of the same genotype for their potential value for characterizing the virus life cycle, drug sensitivity and virus-related cell signaling.

Our present work describes the generation of chimeric viruses with their structural regions from genotype 2b and non-structural genes from the HCV-JFH1 strain. The intergenotypic 2b/JFH1 viruses were compared in terms of intracellular replication, infectious virus production and sensitivity to interferon- α . Here we show that the differences in sensitivity to interferon are attributable to upregulated expression of the cellular interferon signal attenuator, SOCS3, and that this upregulation is caused by overexpression of interleukin-6 (IL-6).

Abbreviations: HCV, hepatitis C virus; TLR, toll-like receptor; FBS, fetal bovine serum; ISG, interferon-stimulated gene; IFN, interferon; SOCS, suppressor of cytokine signaling; IL, interleukin; ALT, alanine aminotransferase; UTR, untranslated region; CLEIA, chemiluminescence enzyme immunoassay; PVDF, polyvinylidene fluoride; STAT, signal transducer and activator of transcription; IFNAR, interferon α / β receptor.

* Corresponding author. Department of Gastroenterology and Hepatology, Tokyo Medical and Dental University, 1-5-45 Yushima, Bunkyo-ku, Tokyo 113-8519, Japan. Fax: +81 3 5803 0268.

E-mail address: nsakamoto.gast@tmd.ac.jp (N. Sakamoto).

¹ G.S. and N.S. contributed equally to this work.

Results

In vitro and *in vivo* infectivity analyses of HCV-2b and 2b/JFH1 intragenotypic chimeras

First, we investigated the infectivity of the full-length genotype 2b clone *in vitro* and *in vivo*. The full-length genotype 2b HCV clone was infectious after direct injection of RNA transcribed *in vitro* into the livers of human hepatocyte engrafted albumin-uPA/SCID mice (see the Supplementary Fig. 1). However, transfection of the HCV RNA into Huh7.5.1 cells did not lead to replication or secretion of virions. Knowing that the full-length genotype 2b HCV was not infectious *in vitro*, we constructed genotype2b/JFH1 intergenotypic recombinants. We constructed three recombinant clones of 2b/JFH1 (Fig. 1A), which were joined between E2 and p7 (JE31F), NS2 and NS3 (JE39F), and within NS2 at nt. 2867 (JEC3F). After transfection of these chimeric HCV RNAs and JFH-1 RNAs into Huh7.5.1 cells, all four clones expressed detectable amounts of HCV core protein in the cells (Fig. 1B) and culture fluid (Fig. 1C). Among the four clones, JEC3F produced the highest level of core protein in the cells and culture fluid. Similarly, in the reinfection assays, JEC3F infected naïve cells most efficiently (Figs. 1D and E). We then compared the infectivity of JEC3F with the other chimeric viruses, genotype2a J6/JFH1 and the JFH1 clone (Supplementary Fig. 2). Transfection of the individual clones into Huh7.5.1 cells showed that JEC3F and the 2b/JFH1 chimera secreted core protein into the medium most efficiently (Fig. 1C). We measured HCV core antigen and HCV-RNA levels in culture supernatant of JEC3F and JFH-1 infected cells. As shown in Fig. 1F, the ratio between supernatant HCV core antigen and HCV-RNA between JEC3F and JFH1 was well correlated each other.

Comparisons of sensitivity to IFN between intragenotypic chimeras and JFH1

Next, we investigated the interferon- α sensitivity of the three 2b/JFH1 chimeric viruses with different junctions, JE31F, JE39F and JEC3F, as well as JFH1. The four viral RNAs were transfected separately into Huh7.5.1 cells and were treated with 0, 1, 3 or 9 IU/mL of interferon- α -2b. Seventy-two hours after addition of interferon, core antigen was measured in the culture fluid. As shown in Fig. 2, all 2b/JFH1 chimeric clones showed significantly higher responses to interferon than JFH1 ($p < 0.01$). These results indicate that the relative interferon sensitivity of 2b/JFH1 clones over JFH1 could be attributable to the sequences of HCV-2b-derived structural proteins, especially core, E1 or E2 protein.

Expression of IFN stimulated genes and STAT1 and 2 phosphorylation in HCV-infected cells

Knowing that the 2b/JFH1 chimeric clones are more sensitive to interferon than JFH1, we next analyzed the effects on cellular interferon signaling. We investigated the expression levels of the interferon-stimulated genes (ISGs), 25OAS and MxA mRNAs that mediate antiviral effects (Itsui et al., 2009; Itsui et al., 2006). Induction of 25OAS and MxA by IFN was significantly suppressed in cells infected with HCV-JFH1 and the JEC3F clones. Of note was that the induction of these ISGs was suppressed substantially in JFH1-infected cells compared to JEC3F-infected cells (Figs. 3A and B). We then detected IFN-induced phosphorylation of STAT1 and STAT2 to pSTAT1 and pSTAT2 in uninfected and JFH1- and JEC3F-infected cells. Phosphorylation of STAT1 and STAT2 occurs within minutes after addition of IFN and substantially decreased at time points later than 8 hours [Itsui, 2006 #1025]. Thus, we detected pSTAT1 and pSTAT2 before and at 15 minutes after IFN treatment. As shown in Figs. 3C and D, production of pSTAT1 and pSTAT2 was decreased substantially in JFH1-infected cells, compared with uninfected and JEC3F-infected

cells. These finding indicated that the differences in sensitivity to interferon of JFH1 and JEC3F were closely associated with attenuation of the cellular IFN signaling pathway.

SOCS 3 is up-regulated in JFH1-infected, IFN-resistant cells

We next investigated the effects of HCV replication on the expression of SOCS1 and SOCS3 that suppress IFN receptor-mediated signaling (Song and Shuai, 1998; Vlotides et al., 2004). While SOCS1 mRNA expression did not differ significantly between uninfected and JFH1- and JEC3F-infected cells, the SOCS3 mRNA expression level was significantly higher in JFH1-infected cells than in uninfected and JEC3F-infected cells (Figs. 4A and B).

Knock down of the SOCS3 gene

To verify that SOCS3 was the key molecule determining the sensitivity to IFN, we performed siRNA knock down of SOCS3 in the virus-infected cells. A SOCS3-directed siRNA was cotransfected with HCV-JFH1 or -JEC3F RNA into Huh7.5.1 cells. Three days after transfection we measured SOCS3 mRNA expression in JFH1 and JEC3F-transfected cells with or without SOCS3-siRNA. Interestingly, SOCS3-knock down in JFH1-transfected cells restored sensitivity of IFN to the same levels as JEC3F-transfected cells (Figs. 5A and B).

Interleukin-6 is involved in SOCS-mediated interferon resistance

It has been reported that SOCS3 is induced principally by phosphorylated STAT3 (pSTAT3) (Hanada et al., 2003) and that interleukin-6 (IL-6) is a strong inducer of pSTAT3 via receptor-mediated Janus kinase activation in the liver (Ramadori and Christ, 1999). This background led us to investigate whether overexpression of SOCS3 is associated with overproduction of IL-6. We investigated Phosphorylated STAT3 (pSTAT3) expression and IL-6 mRNA expression in JFH1- and JEC3F-transfected Huh7.5.1 cells. Phosphorylated STAT3 level was significantly higher in JFH1-transfected cells than JEC3F-transfected cells and naïve Huh7.5.1 cell (Fig. 6A). Moreover IL-6 gene expression level was significantly higher in JFH1-transfected cells than JEC3F-transfected cells (Fig. 6B). Consistent with previous reports, treatment of the Huh7.5.1 cells with IL-6 induced expression of SOCS3 and SOCS1 mRNAs with SOCS3 being much stronger than SOCS1 (Fig. 6C).


Anti-IL-6 antibody restored IFN-resistance to HCV-infected cells

To investigate whether IL-6 is responsible for HCV infection-induced upregulation of SOCS and for resistance to interferon, JFH1 and JEC3F-infected Huh7.5.1 cells were pretreated with antibodies directed against IL-6 and subsequently treated with interferon. Interestingly, anti-IL-6-treated HCV-infected cells became significantly more susceptible to IFN treatment (Fig. 6D) without affecting viral expression levels in the absence of interferon (Fig. 6E). Cellular levels of SOCS3 mRNA were significantly lower in anti-IL-6-treated cells than untreated cells (Fig. 6F). These results strongly suggested that the interferon resistance of HCV-infected cells and the difference between the two viral strains are partly mediated by internal overproduction of IL-6 and subsequent upregulation of SOCS3.


Determination of the HCV structural region that induced SOCS3 and IL6

We studied further which part of HCV structural polyprotein is responsible for the difference in interferon-sensitivity. We constructed two additional chimeric clones between HCV-2b and JFH1. The 2bCoreJFH1 had the 2b-core region followed by the JFH1-structural and nonstructural regions. JCore3F was derived from JEC3F by exchanging the 2b-core with the JFH1-core (Fig. 7A). As


A




B




C




D



E



F



shown in Fig. 7B JFH1 and JCoreC3F, which had a JFH1-derived core region, were significantly more resistant to IFN than JEC3F and 2bCoreJFH1, with a 2b-derived core (Fig. 7B). Consistent with the interferon sensitivity results, JFH1 and JCoreC3F-infected cells expressed SOCS3 and IL6 mRNAs at significantly higher levels than JEC3F and 2bCoreJFH1-infected cells (Figs. 7C and D). These differences in gene expression were inversely associated with the cellular expression levels of each HCV chimeric clone (Fig. 7E). These results indicate that the amino acid sequence of the core protein is responsible for IL-6 and SOCS3-mediated interferon resistance.

Discussion

In this study, we succeeded in establishing a new genotype 2b infectious HCV clone and genotype 2b/JFH1 cell culture-competent intragenotypic chimeric viruses (Fig. 1). Relative interferon sensitivities of 2b/JFH1 chimeras, compared with HCV-JFH1 virus (Fig. 2), led us to conduct a series of assays to investigate the molecular mechanisms of IFN-related response pathways. We found that IFN- α receptor-mediated cellular responses were more attenuated in HCV-JFH1- and 2b/JFH1 chimera-infected than in uninfected Huh7.5.1 cells, but more potently for HCV-JFH1. Precise intragenotypic recombination analyses showed that the amino acid sequence of the HCV core protein is responsible for the differences in interferon sensitivity (Figs. 2, 7). The differences in the interferon-mediated antiviral effects were demonstrated further by the different rates of induction of interferon-inducible MxA and 25-OAS mRNAs (Figs. 3A and B) and IFN induced phosphorylation of STAT1 and STAT2 (Figs. 3D and E). We have demonstrated further that the expression of an interferon signal attenuator, SOCS3, was significantly higher in JFH1 than in 2b/JFH1-infected cells (Song and Shuai, 1998; Vlotides et al., 2004). Indeed, the siRNA-knock down of SOCS3 in JFH1 and 2b/JFH1-infected cells resulted in responsiveness to IFN (Fig. 5). Moreover, cellular expression of IL-6, which increases cytoplasmic phospho-STAT3 (Fig. 6A) and induces SOCS3 expression (Ramadori and Christ, 1999) was significantly higher in JFH1 transfected cells (Fig. 6B). Furthermore, by pre-treatment with anti-IL-6 antibody, JFH1- and 2b/JFH1-infected cells partially recovered elevation of SOCS3 expression and unresponsiveness to IFN (Fig. 6D). Taking all these things together, it is strongly suggested that the differences in IFN sensitivity between genotypes or isolates could be explained by SOCS3-mediated attenuation of interferon responses and, more importantly, IL-6 may constitute a molecular target to reverse such cellular interferon resistance.

Vast numbers of studies have failed to construct infectious HCV clones, other than HCV-JFH1. Murayama, et al. have conducted intragenotypic homologous recombination analyses between HCV-J6 and -JFH1 and have reported that the NS3 protease and NS5B polymerase are essential for replication of the recombinant virus (Murayama et al., 2007). Up to now, several JFH1-based chimeric viruses have been reported, which include genotypes 4a (Scheel et al.,




Fig. 2. Comparison IFN- α sensitivity among 2b/JFH1 chimeric viruses and JFH1. Ten μ g of JE31F, JE39F, JEC3F, JFH1 RNA were transfected into 5×10^6 Huh7.5.1 cells. The transfected cells were divided into 12 wells. Forty eight hours after transfection, cells were washed twice with PBS and treated with 0, 1, 3 and, 9 U/ml IFN- α -2b. Seventy-two hours after IFN- α 2b addition, quantification of HCV core antigen in culture fluids was conducted. The experiments were conducted twice by using Huh 751 cells of different passage, and a representative data was shown. Assays were done in triplicate and the data are shown as mean \pm sd. Asterisks indicate p-values of less than 0.05.

2008), genotype 1a, 1b, 2a (Pietschmann et al., 2006), genotype 3a (Gottwein et al., 2007), genotype 5 a (Jensen et al., 2008) and, genotype 2b, 6a, 7a (Gottwein et al., 2009). Gottwein, et al. constructed intergenotypic chimeric HCV from JFH1 and genotypes 1 through 7 and analyzed differences in sensitivity to antiviral drugs (Gottwein et al., 2009). However, intergenotypic differences in sensitivity to IFN- α and the molecular mechanisms involved have not been well characterized. In this study, we constructed several chimeric virus clones between HCV-2b and HCV-JFH1 (2a), which showed variable sensitivity to IFN and confirmed that the core region is responsible for such IFN sensitivity. This study may support the feasibility of such inter and intragenotypic homologous recombination approaches to characterize differences in viral kinetics and drug responses.

Type I IFNs and their responsive ISGs are the principal mediators of host defense against virus infections, including HCV (Chang et al., 1991; Kalvakolanu, 2003; Ronni et al., 1998). On binding of IFNs to their receptors, IFNAR1 and IFNAR2, Janus kinases-1 and -2 phosphorylate STAT1 and STAT2 to form ISGF-3, which translocates to the nucleus and activates transcription of ISGs (Samuel, 2001; Taniguchi et al., 2001; Taniguchi and Takaoka, 2002). Members of the SOCS family are potent inhibitors of type I and type III IFN-induced activation of the Jak-STAT pathway and subsequent expression of ISGs (Vlotides et al., 2004). In HCV subgenomic replicon-expressing cells, expression levels of SOCS3 were inversely correlated with sensitivity to IFN to suppress viral RNA replication (Zhu et al., 2005).

Fig. 1. Replication and infection competency of HCV-2b/JFH1 chimeric viruses. A. Genomic structures of HCV-JFH1, HCV-2b and 2b/JFH1 chimeric viruses. Intergenotypic homologous recombination was conducted between the HCV-2b and JFH1 (2a) clones and three chimeric clones were constructed that were joined between NS2-NS3 (JE39F), and within E2 at nt2541 (JE31F) and NS2 at nt. 2867 (JEC3F). B. Immunocytochemistry of HCV core. HCV RNA-transfected Huh7.5.1 cells were plated onto 22 mm-round micro cover glasses. Immunocytochemistry was performed 4 days after transfection using mouse-anti-core antibody (green) and DAPI (blue). C. Time courses of 2b/JFH1- and JFH1-transfected cells. *In vitro* transcribed HCV RNAs were transfected into Huh7.5.1 cells by electroporation and HCV core levels of culture fluids were sampled at the time points indicated and core antigen levels were measured. The experiment was done three times with similar results independently. Panel C shows representative data. D. Immunocytochemistry of HCV core. HCV RNA-infected Huh7.5.1 cells using Panel B supernatant that have same amount of HCV core antigen were plated onto 22 mm-round micro cover glasses. Immunocytochemistry was performed 4 days after infection using mouse-anti-core antibody (green) and DAPI (blue). Numbers at the bottom denote percentages of HCV core-positive cells. E. Time courses of 2b/JFH1 infected cells. JE31F, JE39F, JEC3F RNA-transfected cell culture fluids were used to infect naïve Huh7.5.1 cells in 60 mm-diameter plates at density of 3×10^5 cells per plate. Quantification of HCV core antigen in culture supernatants was carried out at 24 hours, 48 hours, 72 hours and 144 hours after inoculation. The experiment was done three times with similar results independently. Panel E shows representative data. F. Comparison between JFH1 and JEC3F supernatant HCV-RNA titer and core antigen. Four days after JFH1 and JEC3F RNA transfection, culture supernatant was harvested and subjected to both HCV core antigen assay and realtime RT-PCR of HCV-RNA. Assays were done in triplicate and the data are shown as mean \pm sd.




Fig. 3. Induction by interferon of the interferon-inducible genes, MxA (panel A), 25-OAS (panel B) and phosphorylated STAT1 (panel C) and STAT2 (panel D). JEC3F and JFH1 10 μ g RNA was transfected into Huh7.5.1 cells. Forty-eight hours after transfection, the cells were treated with 25 U/ml IFN- α 2b. Total cellular RNA was isolated before and 8 and 24 hours after IFN treatment. Relative gene expression levels of MxA (panel A) and 25-OAS (panel B) were determined by real-time PCR at the time points indicated. JEC3F and JFH1 RNA and MOCK was transfected into Huh7.5.1 cells. Forty eight hours after transfection, the cells were treated with 25 U/ml IFN- α 2b. Total cellular protein was isolated before and 15 minutes after IFN treatment. Ten μ g of extracted protein were used for analysis of phosphorylated STAT1, STAT2 protein and STAT1, STAT2 protein as controls. Assays were done in triplicate and the data are shown as mean \pm sd. Asterisks indicate p-values of less than 0.05.

HCV, on the other hand, counteracts such IFN-mediated antiviral pathways. The NS5A and E2 proteins interfere with the action of IFN by inhibiting the activity of PKR (He and Katze, 2002; Taylor et al., 1999). NS5A also induced expression of IL-8 and attenuated expression of ISGs (Polyak et al., 2001). HCV core protein has been reported to bind the STAT1-SH domain (Lin et al., 2006) or destabilize STAT1 (Lin et al., 2005) to block IFN signaling. It also has been reported that overexpression of core protein upregulated SOCS3 expression (Bode et al., 2003). In this study, we used full-length HCV cell culture and found, *for the first time*, that SOCS3 expression is upregulated differently depending on the genetic sequences of HCV strains and that these differences in SOCS3 expression are associated with sensitivity to IFN. Moreover, overexpression and knock down of SOCS3 expression were closely associated with the IFN sensitivity of the HCV-infected cells. These results indicate that interferon-resistance of HCV-infected cells is directed by overexpression of SOCS3, which may be upregulated by HCV proteins as reported (Bode et al., 2003) (Kawaguchi et al., 2004). A sequence comparison of our HCV2b and JFH1 clones has found 16 amino acid differences. These structural differences of the core protein might affect cellular responses to interferon (see the Supplementary Fig. 4).

It has been reported that IL-6 is the principal activator of STAT3 in hepatocytes through binding its receptor (Hanada et al., 2003; Ramadori and Christ, 1999). Furthermore, plasma IL-6 levels are elevated in




Fig. 4. Expression of SOCS1 mRNA (panel A), SOCS3 mRNA (panel B). Forty-eight hours after transfection of JEC3F, JFH1 10 μ g RNA or mock transfection into Huh7.5.1 cells, total RNA and total protein were isolated. Relative gene expression levels of SOCS1 (panel A) and SOCS3 (panel B) and were determined by real time PCR. Values are shown as relative to those of uninfected Huh 751 cells. Assays were done in triplicate and the data are shown as mean \pm sd. Asterisks indicate p-values of less than 0.05.




Fig. 5. Differences in sensitivities to IFN between SOCS3-knock down, HCV transfected cells. JFH1 or JEC3F 10 μ g RNA, and 80 pmol siRNA SOCS3-HSS113312 or MOCK were electroporated into 5×10^6 uninfected Huh7.5.1 cells. A. Expression of SOCS3 mRNA in uninfected and HCV-infected Huh7 cells. Forty-eight hours after transfection, total RNA was isolated. Relative gene expression level of SOCS3 were determined by real time PCR. Values are shown as relative to those of JFH1 infected Huh 751 cells. Assays were done in triplicate and the data are shown as mean \pm sd. Asterisks indicate p-values of less than 0.05. B. Dose-dependent suppression of HCV replication by IFN in SOCS3-knock-down, HCV-infected cells. The above siRNA and HCV RNA-transfected cells were divided into 12 wells. Forty eight hours after transfection, the cells were treated with 0, 1, 5 and 25 U/ml of IFN- α 2b. Seventy two hours after treatment, quantification of HCV core antigen in culture fluids was carried out. Assays were done in triplicate and the data are shown as mean \pm sd. Asterisks indicate p-values of less than 0.05.

chronic hepatitis C patients (Malaguamera et al., 1997). Consistent with those reports, we found that IL-6 strongly induced SOCS3 expression in Huh7.5.1 cells (Fig. 6C). More importantly, cellular IL6 expression levels were in the order of uninfected < JEC3F < JFH1-infected cells, which correlated well with SOCS3 expression (Fig. 4) and with cellular responses to IFN (Fig. 2). In addition, the IFN-resistant JcoreC3F, in which the core region of JEC3F had been re-substituted by the JFH1-core, induced comparatively higher levels of IL-6 and SOCS3 mRNA to JFH1 (Fig. 7). Taken together, our results indicate that the amino acid sequence of the core protein determines IL-6 and SOCS3 expression levels and, as a consequence, resistance to IFNs.

It remains to be clarified what are the inducers of IL-6. There are reports that HCV core protein activates toll-like receptor (TLR)-2 in Huh7 cells and in adult human hepatocytes (Hoffmann et al., 2009; Mozer-Lisewska et al., 2005). TLRs are known to activate downward NF- κ B signaling that upregulates IL-6 expression. Alternatively, IL-6 may be secreted in response to cellular steatosis and insulin resistance. HCV patients with obesity or insulin resistance are refractory to IFN treatments. Such patients have higher levels of hepatic SOCS3 expression than those without obesity or insulin resistance (Miyaaki et al., 2009; Walsh et al., 2006). More recently, Sabio, *et al* have reported that fatty acid-induced secretion of IL-6 from adipocytes upregulates hepatic SOCS3, leading to insulin-resistance (Sabio et al., 2008).

In conclusion, our study demonstrates that HCV intragenotypic and inter-strain differences in IFN sensitivity can be, in most part, attributable to the amino acid sequence of the HCV core protein and that such IFN sensitivities are determined by cellular expression levels of SOCS3 and IL-6. Therapeutic targeting of IL-6 potentially may be a key to targeting IFN-resistance and improving antiviral chemotherapeutics against HCV.

Materials and Methods

Reagents and antibodies

Recombinant human interferon alpha-2b was from Schering-Plough (Kenilworth, NJ). Anti-CD 81 antibody (JS-81) was from BD Biosciences (Franklin Lakes, NJ) (Morikawa et al., 2007), anti-IL6 receptor antibody was from Chugai pharmaceutical Co (Tokyo, Japan), anti-SOCS3 was from Cell Signaling (Beverly, MA), and anti-IL6 antibody was from R&D Systems (Minneapolis, MN).

Cloning of HCV cDNA from patient serum

A serum sample was obtained from a 32-year-old male who developed acute hepatitis after intravenous drug injection. Serum was obtained one week after the onset of symptoms. Total RNA was extracted from 150 μ l of serum using ISOGEN (Nippon Gene, Osaka, Japan). cDNA was synthesized using SuperScript II (Invitrogen, Carlsbad, CA) reverse transcriptase. PCR primers, based on a genotype 2b prototype sequence, HC-J8 (accession number: D10988), were used to amplify 14 fragments of HCV cDNA covering nt. 13–9478 (nucleotide numbers corresponded to HC-J8) by PCR. All amplicons were purified and cloned into the pGEM-T EASY vector (Promega, Madison, WI) and nucleotide sequences were determined using Big Dye Terminator Cycle Sequencing Ready Reaction kits (Applied Biosystems, Foster City, CA) and an automated DNA sequencer (ABI PRISM® 310 Genetic Analyzer; Applied Biosystems). The consensus sequence of five clones was adopted for each region. Each consensus sequence segment of HCV was assembled into pJFH1-full (Wakita et al., 2005) by substituting the insert sequence of pJFH1-full.




Fig. 6. IL-6 expression in HCV infected cells and change in IFN sensitivity by treatment with anti-IL6 antibody. **A.** Expression of cytoplasmic phospho-STAT3 in uninfected and HCV-infected Huh7 cells. JEC3F, JFH1 10 μ g RNA and MOCK was transfected into Huh7.5.1 cells. Forty eight hours total cellular protein was isolated. Ten μ g of extracted protein were used for analysis of phosphorylated STAT3, STAT protein and β -actin as controls. **B.** Expression of Interleukin-6 mRNA in uninfected and HCV-infected Huh7 cells. Forty-eight hours after transfection, total RNA was isolated. Relative gene expression level of IL6 were determined by real time PCR. Values are shown as relative to those of uninfected Huh 751 cells. Assays were done in triplicate and the data are shown as mean \pm sd. **C.** IL-6 induces SOCS3 strongly in uninfected Huh7.5.1 cells. Uninfected Huh7.5.1 cells were treated with 10 ng/ml recombinant human IL6 (PEPRO TEC EC, London, England). Fifteen minutes after treatment, total RNA was isolated. Relative gene expression levels of SOCS1 and SOCS3 were determined by real time PCR. Uninfected Huh7.5.1 cells that were not treated with IL6 were used as a control. Values are shown as relative to those of uninfected Huh 751 cells. Assays were done in triplicate and the data are shown as mean \pm sd. **D.** Dose-dependent suppression of HCV replication by IFN in HCV-infected cells pre-treated with anti-IL-6 antibody. Immediately after electroporation, HCV RNA-transfected cells were divided into 12 wells and pretreated with 1 μ g/ml anti-IL6 antibody. Forty eight hours after transfection, the cells were washed with PBS and treated with 0, 1, 5 and 25 U/ml of IFN-alpha 2b. Seventy two hours after treatment, quantification of HCV core antigen was carried out in culture fluids. Assays were done in triplicate and the data are shown as mean \pm sd. **E.** Core protein secretion levels following treatment of HCV-transfected cells with anti-IL-6 antibody. After treatment with anti-IL-6 antibody, HCV RNA-transfected cells were divided into 12 wells. Five days after transfection, quantification of HCV core antigen was carried out in culture fluids. Assays were done in triplicate and the data are shown as mean \pm sd. **F.** Expression of SOCS3 mRNA in uninfected and HCV-infected Huh7 cells. Forty-eight hours after transfection, total RNA was isolated. Relative SOCS3 gene to beta-actin gene expression were determined by real time PCR. Values are shown as relative to those of uninfected Huh 751 cells. Assays were done in triplicate and the data are shown as mean \pm sd. Asterisks indicate p-values of less than 0.05.




Fig. 6 (continued).

Construction of 2b/JFH-1 based intragenotypic chimeras and transfection

Chimeric HCV constructs of HCV-2b and JFH1 were shown in Figs. 1A and 7A. To construct 2b/JFH1-based intragenotypic chimera, JE31F, the 2b sequence of core through E2 (nt. 342–2541) was fused to the EcoRI-JFH1- 5'-untranslated region (UTR) DNA by fusion PCR. The fused 5'UTR-E2 fragment and JFH1-E2-NS3 (nt2541 through 5324) were assembled by fusion PCR and cloned into pGEM-T EASY. The product was digested by EcoRI and AfeI and insert into pJFH1. Plasmids pJE39F, pJEC3F, pJcoreC3F and p2bcore JFH1 were constructed using a similar procedure. Plasmids pJEC3F and pJE39F were joined between NS2 and NS3, and within NS2 at nt. 2867, respectively. Plasmid pJcoreC3F was made by substitution of the core region of 2b/JFH1 with that of JFH1. The plasmid p2bcoreJFH1 was made by substitution of the core region of JFH1 with that of 2b/JFH1.

Cells and cell culture

Huh7.5.1 cells were maintained in Dulbecco's modified minimal essential medium (Sigma, St. Louis, MO) supplemented with 10% fetal calf serum at 37 °C under 5% CO₂.

HCV cell culture system

Full-length HCV expression plasmids were as follows: pJFH1-full (Wakita et al., 2005), pJE31F, pJE39F, pJEC3F, pJcoreC3F, p2bcoreJFH1, and pFL-H77/JFH1, pFL-J6/JFH1 (Lindenbach et al., 2005). These plasmids were linearized at their 3' ends and used as templates for HCV RNA synthesis using the RiboMax Large Scale RNA Production System (Promega, Madison, WI). After DNase I (RQ-1, RNase-free DNase, Promega) treatment, the HCV RNA was purified using ISOGEN (Nippon Gene, Tokyo, Japan). For the RNA transfection, Huh7.5.1 cells were washed twice with PBS, and 5×10^6 cells were suspended in Opti-MEM I (Invitrogen Carlsbad, CA) containing 10 µg of HCV RNA, transferred into a 4 mm electroporation cuvette and finally subjected to an electric pulse (1,050 µF and 270 V) using the Easy Jet system (EquiBio, Middlesex, UK). After electroporation, the cell suspension

was left for 5 min at room temperature and then incubated under normal culture conditions in a cell culture dish.

Quantification of HCV core antigen in culture supernatants

Culture supernatants of HCV RNA transfected Huh7.5.1 cells were collected on the days indicated, passed through a 0.45 µm filter (MILLEX-HA, Millipore, Bedford, MA) and stored at -80 °C. The concentrations of core antigen in the culture supernatants were measured using a chemiluminescence enzyme immunoassay (CLEIA) according to the manufacturer's protocol (Lumipulse Ortho HCV Antigen, Ortho-Clinical Diagnostics, Tokyo, Japan).

Re-infection analyses

Titer-adjusted supernatants (including 0.03 fmol HCV core antigen) from HCV RNA-transfected cells were inoculated onto naïve Huh7.5.1 cells plated on a 6 cm plate at a density of 3×10^5 cells per plate. Forty-eight hours after inoculation, anti-core immunostaining was carried out with mouse anti-HCV core protein monoclonal antibody and the numbers of infected cells were counted. HCV core antigen in culture supernatants was measured at 24 hours, 48 hours, 72 hours and 144 hours after inoculation.

Real-time RT-PCR analysis

For the detection of HCV RNA in culture supernatant, supernatant was passed through a 0.45 µm filter (MILLEX-HA, Millipore, Bedford, MA) and stored at -80 °C until use. Protocol and primers for the realtime RT-PCR analysis of HCV-RNA has been described previously (Sekine-Osajima et al., 2008). For the detection of endogenous mRNAs, total cellular RNA was isolated using ISOGEN (Nippon Gene). Two micrograms of total cellular RNA were used to generate cDNA from each sample using SuperScript II. Expression of mRNA was quantified using the TaqMan Universal PCR Master Mix and the ABI 7500 Real-Time PCR System (Applied Biosystems, Foster City CA). Some primers have been described (Sekine-Osajima et al., 2008). SOCS3; forward, 5'-CAC ATG GCA CAA GCA CAA GAA G-3' and reverse,




Fig. 7. Replacement of the HCV-2b-core region with JFH1-core causes upregulation of SOCS3 and IL-6 and restores resistance to IFN. **A.** Genome maps of JFH1, JEC3F, J core C3F, 2b core JFH1 recombinant cDNA. J core C3F was made by substitution of the core region of 2b/JFH1 with that of JFH1. The 2b core JFH1 was made by substitution of the core region of JFH1 with that of 2b/JFH1. **B.** Comparison of IFN-alpha sensitivity among JFH1 and JEC3F and core region substitution chimeric viruses. Ten μ g of J core C3F, 2b core JFH1, JEC3F, JFH1 RNA were transfected into 5×10^6 Huh7.5.1 cells and were divided into 12 wells. Forty eight hours after transfection, the cells were treated with 0, 1, 5 and 25 U/ml of IFN-alpha 2b. Seventy two hours after treatment, quantification of HCV core antigen was carried out in culture fluids. Assays were done in triplicate and the data are shown as mean \pm sd. Asterisks indicate p-values of less than 0.05. **C, D.** Core substitution leads to SOCS3 and IL-6 mRNA over-expression. Forty eight hours after transfection into cells, total RNA was isolated. Relative gene expression level SOCS3 (panel C) and IL6 (panel D) were determined by real time PCR. Values are shown as relative to those of uninfected Huh 751 cells. Assays were done in triplicate and the data are shown as mean \pm sd. Asterisks indicate p-values of less than 0.05. **E.** Change of secretion of core protein following core protein substitution. HCV RNA-transfected cells were divided into 12 wells. Five days after transfection, quantification of HCV core antigen was carried out in culture fluids. Assays were done in triplicate and the data are shown as mean \pm sd. Asterisks indicate p-values of less than 0.05.

5'-GGA GAA GCT GGA GACTCA GGT G-3', SOCS1; forward, 5'-CAC TTC CGC ACA TTC CGT TCG-3' and reverse, 5'-GAG GCC ATC TTC ACG CTA AGG-3', IL6; forward, 5'-GGT ACA TCC TCG ACG GCA TCT-3' and reverse, 5'-GTG CCT CTT TGC TGC TTT CAC-3', 25OAS; forward, 5'-CCA CCT TGG AAA GTG CCG ACA ATG CAG ACA-3' and reverse, 5'-CGA GTC TTT AAA AGC GAT TGC CAG ATG ATC -3', MxA; forward, 5'-GCC AGC AGC TTC AGA AGG CCA TGCTGC AGC-3' and reverse, 5'-GGG CAA GCC GGC GCC GAG CCT GCG TCA GCC -3'.

The siRNAs

The siRNAs directed against SOCS3 were designed as follows: SOCS3-HSS113312 stealth (sequence 5'- CCC AGA AGA GCC UAU UAC AUC UAC U-3' and 5'-AGU AGA UGU AAU AGG CUC UUC UGG G-3', Invitrogen) was used. 10 µg in vitro-synthesized HCV-RNA and 80 pmol siRNA SOCS3-HSS113312 or MOCK or control siRNA (negative universal control Med #2, Invitrogen) were electroporated into 5 × 10 naïve Huh7.5.1 cells using the protocol described in *HCV cell culture system*. Forty-eight hours after transfection, expression levels of SOCS3 mRNA were measured by real-time PCR. The difference in IFN sensitivity between SOCS3 knock down HCV infected cells and control HCV infected cells was determined by measuring supernatants HCV core antigen 72 hours after addition of IFN.

Immunohistochemistry for HCV core

HCV-JFH1 transfected or infected Huh7.5.1 cells were cultured on 22 mm-round micro cover glasses (Matsunami, Tokyo, Japan). For detection of HCV core, cells were fixed with cold acetone for 15 min. The cells were incubated with the primary antibodies for 1 hour at 37 °C, and with Alexa Fluor 488 goat anti-mouse IgG antibody (Molecular Probes, Eugene, OR) for 1 hour at room temperature. Cells were mounted with VECTA SHIELD Mounting Medium and DAPI (Vector Laboratories, Burlingame, CA) and visualized by fluorescence microscopy (BZ-8000, KEYENCE, Osaka, Japan).

Western blot analysis

Western blotting was performed as described (Tanabe et al., 2004). Briefly, 10 µg of total cell lysate was separated by SDS-PAGE, and blotted onto a polyvinylidene fluoride (PVDF) membrane. The membrane was incubated with the primary antibodies followed by a peroxidase-labeled anti IgG antibody, and was visualized by chemiluminescence using the ECL Western Blotting Analysis System (Amersham Biosciences, Buckinghamshire, UK).

Statistical analyses

Statistical analyses were performed using Student's *t*-test; *p*-values of less than 0.05 were considered statistically significant.

Acknowledgments

We thank Dr. Frank Chisari for providing Huh7.5.1 cells and Dr. Takaji Wakita for providing pJFH1fill. This study was supported by grants from Ministry of Education, Culture, Sports, Science and Technology-Japan, the Japan Society for the Promotion of Science, Ministry of Health, Labour and Welfare-Japan, Japan Health Sciences Foundation, and National Institute of Biomedical Innovation.

Appendix A. Supplementary data

Supplementary Fig. 1. Infectivity of the full-length 2b HCV RNA and 2b/JFH1 chimeric virus, JEC3F. A. Challenge of human liver-engrafted albumin-uPA/SCID mice with culture fluid from JFH1 and JEC3F cells. Cell culture fluids from the JFH1 clone and JEC3F were injected

intravenously into human liver engrafted albumin-uPA/SCID mice. Serum samples were obtained from the mice every 2 weeks after injection and the HCV RNA titer was determined. B. Fig. 1B Challenge of human liver-engrafted albumin-uPA/SCID mice by intrahepatic injection of in vitro synthesized, full-length 2b HCV RNA. Five hundred µl of RNA solution containing 30 µg of in vitro synthesized full-length 2b HCV RNA was injected into the livers of anesthetized chimeric mice through a small abdominal incision. Serum samples were obtained from the mice every 2 weeks after injection and the HCV RNA titer was determined.

Supplementary Fig. 2. Comparisons of replication efficiency of JFH1 and J6/JFH1, 2b/JFH1 chimeras after transfection into Huh7.5.1-cells. A. Structures of the J6/JFH1 and 2b/JFH1 genomes. J6 is joined between NS2 and NS3 with JFH1. 2b-HCV is joined with JFH1 within NS2 at nt. 2867. B. Measurements of core protein in cell culture fluids. Ten µg of JFH1, J6/JFH1, 2b/JFH1 RNA were transfected into 5 × 10⁶ Huh7.5.1 cells and the cells were cultured in 100 mm-diameter plates. The culture fluids from JFH1, J6/JFH1, H77/JFH1 or 2b/JFH1-transfected Huh7.5.1 cells were collected separately on the days indicated and the levels of core antigen were measured. These experiments were done three times with similar results independently. Panel B shows representative data.

Supplementary Fig. 3. Inhibition of infection by blocking CD81. Huh 7.5.1 cells were plated into a 6 well plate at 1.4 × 10⁵ cells per well. After 48 hours, the cells were incubated with anti-CD81 or isotypematched control antibody at the concentration indicated for 1 hour. Subsequently, cells were infected with 1 ml of JEC3F stock cell culture fluids at day 2 for 4 hours and washed with PBS. 48 hours after inoculation, anti-core immunostaining was performed with mouse anti-HCV core protein monoclonal antibody (Panels B and C). Quantification of HCV core antigen was carried out in culture fluids at 48 hours after infection (Panel A).


Supplementary Fig. 4. Comparison between 2b and JFH-1 core amino acid sequence.

Note: Supplementary materials related to this article can be found online at doi:10.1016/j.virol.2010.07.041.

References

- Alter, M.J., 1997. Epidemiology of hepatitis C. *Hepatology* 26 (3 Suppl 1), 62S–65S.
- Blight, K.J., Kolykhalov, A.A., Rice, C.M., 2000. Efficient initiation of HCV RNA replication in cell culture. *Science* 290 (5498), 1972–1974.
- Bode, J.G., Ludwig, S., Ehrhardt, C., Albrecht, U., Erhardt, A., Schaper, F., Heinrich, P.C., Haussinger, D., 2003. IFN-alpha antagonistic activity of HCV core protein involves induction of suppressor of cytokine signaling-3. *FASEB J.* 17 (3), 488–490.
- Bukh, J., Miller, R.H., Purcell, R.H., 1995. Genetic heterogeneity of hepatitis C virus: quasispecies and genotypes. *Semin. Liver Dis.* 15 (1), 41–63.
- Chang, K.C., Hansen, E., Foroni, L., Lida, J., Goldspink, G., 1991. Molecular and functional analysis of the virus- and interferon-inducible human MxA promoter. *Arch. Virol.* 117 (1–2), 1–15.
- Fried, M.W., Shiffman, M.L., Reddy, K.R., Smith, C., Marinos, G., Goncalves Jr., F.L., Haussinger, D., Diago, M., Carosi, G., Dhumeaux, D., Craxi, A., Lin, A., Hoffman, J., Yu, J., 2002. Peginterferon alfa-2a plus ribavirin for chronic hepatitis C virus infection. *N. Engl. J. Med.* 347 (13), 975–982.
- Gottwein, J.M., Scheel, T.K., Hoegh, A.M., Lademann, J.B., Eugen-Olsen, J., Lisby, G., Bukh, J., 2007. Robust hepatitis C genotype 3a cell culture releasing adapted inter-genotypic 3a/2a (S52/JFH1) viruses. *Gastroenterology* 133 (5), 1614–1626.
- Gottwein, J.M., Scheel, T.K., Jensen, T.B., Lademann, J.B., Prentoe, J.C., Knudsen, M.L., Hoegh, A.M., Bukh, J., 2009. Development and characterization of hepatitis C virus genotype 1-7 cell culture systems: role of CD81 and scavenger receptor class B type I and effect of antiviral drugs. *Hepatology* 49 (2), 364–377.
- Hanada, T., Kinjyo, I., Inagaki-Ohara, K., Yoshimura, A., 2003. Negative regulation of cytokine signaling by CIS/SOCS family proteins and their roles in inflammatory diseases. *Rev. Physiol. Biochem. Pharmacol.* 149, 72–86.
- He, Y., Katze, M.G., 2002. To interfere and to anti-intefere: the interplay between hepatitis C virus and interferon. *Viral Immunol.* 15, 95–119.
- Hoffmann, M., Zeisel, M.B., Jilg, N., Paranhos-Baccala, G., Stoll-Keller, F., Wakita, T., Hafkemeyer, P., Blum, H.E., Barth, H., Henneke, P., Baumert, T.F., 2009. Toll-Like Receptor 2 Senses Hepatitis C Virus Core Protein but Not Infectious Viral Particles. *J. Innate Immun.* 1 (No. 5), 446–454.
- Hoofnagle, J.H., di Bisceglie, A.M., 1997. The treatment of chronic viral hepatitis. *N. Engl. J. Med.* 336 (5), 347–356.
- Itsuji, Y., Sakamoto, N., Kakinuma, S., Nakagawa, M., Sekine-Osajima, Y., Tasaka-Fujita, M., Nishimura-Sakurai, Y., Suda, G., Karakama, Y., Yamamoto, M., Watanabe, T.,

- Ueyama, M., Funaoka, Y., Azuma, S., Watanabe, M., 2009. Antiviral effects of the interferon-induced protein GBP-1 and its interaction with the hepatitis C virus NS5B protein. *Hepatology* 50 (6), 1727–1737.
- Itsui, Y., Sakamoto, N., Kurosaki, M., Kanazawa, N., Tanabe, Y., Koyama, T., Takeda, Y., Nakagawa, M., Kakinuma, S., Sekine, Y., Maekawa, S., Enomoto, N., Watanabe, M., 2006. Expressional screening of interferon-stimulated genes for antiviral activity against hepatitis C virus replication. *J. Viral Hepat.* 13 (10), 690–700.
- Jensen, T.B., Gottwein, J.M., Scheel, T.K., Hoegh, A.M., Eugen-Olsen, J., Bukh, J., 2008. Highly efficient JFH1-based cell-culture system for hepatitis C virus genotype 5a: failure of homologous neutralizing-antibody treatment to control infection. *J. Infect. Dis.* 198 (12), 1756–1765.
- Kalvachlanu, D.V., 2003. Alternate interferon signaling pathways. *Pharmacol. Ther.* 100 (1), 1–29.
- Kato, T., Date, T., Miyamoto, M., Furusaka, A., Tokushige, K., Mizokami, M., Wakita, T., 2003. Efficient replication of the genotype 2a hepatitis C virus subgenomic replicon. *Gastroenterology* 125 (6), 1808–1817.
- Kawaguchi, T., Yoshida, T., Harada, M., Hisamoto, T., Nagao, Y., Ide, T., Taniguchi, E., Kumemura, H., Hanada, S., Maeyama, M., Baba, S., Koga, H., Kumashiro, R., Ueno, T., Ogata, H., Yoshimura, A., Sata, M., 2004. Hepatitis C virus down-regulates insulin receptor substrates 1 and 2 through up-regulation of suppressor of cytokine signaling 3. *Am. J. Pathol.* 165 (5), 1499–1508.
- Lin, W., Choe, W.H., Hiasa, Y., Kamegaya, Y., Blackard, J.T., Schmidt, E.V., Chung, R.T., 2005. Hepatitis C virus expression suppresses interferon signaling by degrading STAT1. *Gastroenterology* 128 (4), 1034–1041.
- Lin, W., Kim, S.S., Yeung, E., Kamegaya, Y., Blackard, J.T., Kim, K.A., Holtzman, M.J., Chung, R.T., 2006. Hepatitis C virus core protein blocks interferon signaling by interaction with the STAT1 SH2 domain. *J. Virol.* 80 (18), 9226–9235.
- Lindenbach, B.D., Evans, M.J., Syder, A.J., Wolk, B., Tellinghuisen, T.L., Liu, C.C., Maruyama, T., Hynes, R.O., Burton, D.R., McKeating, J.A., Rice, C.M., 2005. Complete replication of hepatitis C virus in cell culture. *Science* 309 (5734), 623–626.
- Lohmann, V., Korner, F., Koch, J., Herian, U., Theilmann, L., Bartenschlager, R., 1999. Replication of subgenomic hepatitis C virus RNAs in a hepatoma cell line. *Science* 285 (5424), 110–113.
- Malaguerra, M., Di Fazio, I., Romeo, M.A., Restuccia, S., Laurino, A., Trovato, B.A., 1997. Elevation of interleukin 6 levels in patients with chronic hepatitis due to hepatitis C virus. *J. Gastroenterol.* 32 (2), 211–215.
- Miyaaki, H., Ichikawa, T., Nakao, K., Matsuzaki, T., Muraoka, T., Honda, T., Takeshita, S., Shibata, H., Ozawa, E., Akiyama, M., Miuma, S., Eguchi, K., 2009. Predictive value of suppressor of cytokine signal 3 (SOCS3) in the outcome of interferon therapy in chronic hepatitis C. *Hepatology* 49 (9), 850–855.
- Morikawa, K., Zhao, Z., Date, T., Miyamoto, M., Murayama, A., Akazawa, D., Tanabe, J., Sone, S., Wakita, T., 2007. The roles of CD81 and glycosaminoglycans in the adsorption and uptake of infectious HCV particles. *J. Med. Virol.* 79 (6), 714–723.
- Mozer-Lisewska, I., Sluzewski, W., Kaczmarek, M., Jenek, R., Szczepanski, M., Figlerowicz, M., Kowala-Piaskowska, A., Zeromski, J., 2005. Tissue localization of Toll-like receptors in biopsy specimens of liver from children infected with hepatitis C virus. *Scand. J. Immunol.* 62 (4), 407–412.
- Murayama, A., Date, T., Morikawa, K., Akazawa, D., Miyamoto, M., Kaga, M., Ishii, K., Suzuki, T., Kato, T., Mizokami, M., Wakita, T., 2007. The NS3 helicase and NS5B-to-3'X regions are important for efficient hepatitis C virus strain JFH-1 replication in Huh7 cells. *J. Virol.* 81 (15), 8030–8040.
- Pietschmann, T., Kaul, A., Koutsoudakis, G., Shavinskaya, A., Kallis, S., Steinmann, E., Abid, K., Negro, F., Dreux, M., Cosset, F.L., Bartenschlager, R., 2006. Construction and characterization of infectious intragenotypic and intergenotypic hepatitis C virus chimeras. *Proc. Natl. Acad. Sci. USA* 103 (19), 7408–7413.
- Polyak, S.J., Khabar, K.S., Rezeiq, M., Gretch, D.R., 2001. Elevated levels of interleukin-8 in serum are associated with hepatitis C virus infection and resistance to interferon therapy. *J. Virol.* 75 (13), 6209–6211.
- Ramadori, G., Christ, B., 1999. Cytokines and the hepatic acute-phase response. *Semin. Liver Dis.* 19 (2), 141–155.
- Ronni, T., Matikainen, S., Lehtonen, A., Palvimo, J., Dellis, J., Van Eyllen, F., Goetschy, J.F., Horisberger, M., Content, J., Julkunen, I., 1998. The proximal interferon-stimulated response elements are essential for interferon responsiveness: a promoter analysis of the antiviral MxA gene. *J. Interferon Cytokine Res.* 18 (9), 773–781.
- Sabio, G., Das, M., Mora, A., Zhang, Z., Jun, J.Y., Ko, H.J., Barrett, T., Kim, J.K., Davis, R.J., 2008. A stress signaling pathway in adipose tissue regulates hepatic insulin resistance. *Science* 322 (5907), 1539–1543.
- Samuel, C., 2001. Antiviral actions of interferons. *Clin. Microbiol. Rev.* 14 (4), 778–809.
- Scheel, T.K., Gottwein, J.M., Jensen, T.B., Prentoe, J.C., Hoegh, A.M., Alter, H.J., Eugen-Olsen, J., Bukh, J., 2008. Development of JFH1-based cell culture systems for hepatitis C virus genotype 4a and evidence for cross-genotype neutralization. *Proc. Natl. Acad. Sci. USA* 105 (3), 997–1002.
- Sekine-Osajima, Y., Sakamoto, N., Mishima, K., Nakagawa, M., Itsui, Y., Tasaka, M., Nishimura-Sakurai, Y., Chen, C.H., Kanai, T., Tsuchiya, K., Wakita, T., Enomoto, N., Watanabe, M., 2008. Development of plaque assays for hepatitis C virus-JFH1 strain and isolation of mutants with enhanced cytopathogenicity and replication capacity. *Virology* 371 (1), 71–85.
- Song, M.M., Shuai, K., 1998. The suppressor of cytokine signaling (SOCS) 1 and SOCS3 but not SOCS2 proteins inhibit interferon-mediated antiviral and antiproliferative activities. *J. Biol. Chem.* 273 (52), 35056–35062.
- Tanabe, Y., Sakamoto, N., Enomoto, N., Kurosaki, M., Ueda, E., Maekawa, S., Yamashiro, T., Nakagawa, M., Chen, C.H., Kanazawa, N., Kakinuma, S., Watanabe, M., 2004. Synergistic inhibition of intracellular hepatitis C virus replication by combination of ribavirin and interferon- α . *J. Infect. Dis.* 189 (7), 1129–1139.
- Taniguchi, T., Ogasawara, K., Takaoka, A., Tanaka, N., 2001. IRF family of transcription factors as regulators of host defense. *Annu. Rev. Immunol.* 19, 623–655.
- Taniguchi, T., Takaoka, A., 2002. The interferon- α / β system in antiviral responses: a multimodal machinery of gene regulation by the IRF family of transcription factors. *Curr. Opin. Immunol.* 14, 111–116.
- Taylor, D.R., Shi, S.T., Romano, P.R., Barber, G.N., Lai, M.M., 1999. Inhibition of the interferon-inducible protein kinase PKR by HCV E2 protein. *Science* 285 (5424), 107–110.
- Vlotides, G., Sorensen, A.S., Kopp, F., Zitzmann, K., Cengic, N., Brand, S., Zachoval, R., Auernhammer, C.J., 2004. SOCS-1 and SOCS-3 inhibit IFN- α -induced expression of the antiviral proteins 2, 5-OAS and MxA. *Biochem. Biophys. Res. Commun.* 320 (3), 1007–1014.
- Wakita, T., Pietschmann, T., Kato, T., Date, T., Miyamoto, M., Zhao, Z., Murthy, K., Habermann, A., Krausslich, H.G., Mizokami, M., Bartenschlager, R., Liang, T.J., 2005. Production of infectious hepatitis C virus in tissue culture from a cloned viral genome. *Nat. Med.* 11 (7), 791–796.
- Walsh, M.J., Jonsson, J.R., Richardson, M.M., Lipka, G.M., Purdie, D.M., Clouston, A.D., Powell, E.E., 2006. Non-response to antiviral therapy is associated with obesity and increased hepatic expression of suppressor of cytokine signalling 3 (SOCS-3) in patients with chronic hepatitis C, viral genotype 1. *Gut* 55 (4), 529–535.
- Zhu, H., Nelson, D.R., Crawford, J.M., Liu, C., 2005. Defective Jak-Stat activation in hepatoma cells is associated with hepatitis C viral IFN- α resistance. *J. Interferon Cytokine Res.* 25 (9), 528–539.



Cell culture and in vivo analyses of cytopathic hepatitis C virus mutants^{☆,☆☆}

Kako Mishima^{a,1}, Naoya Sakamoto^{a,b,*}, Yuko Sekine-Osajima^a, Mina Nakagawa^{a,b}, Yasuhiro Itsui^{a,c}, Seishin Azuma^a, Sei Kakinuma^{a,b}, Kei Kiyohashi^a, Akiko Kitazume^a, Kiichiro Tsuchiya^a, Michio Imamura^d, Nobuhiko Hiraga^d, Kazuaki Chayama^d, Takaji Wakita^e, Mamoru Watanabe^a

^a Department of Gastroenterology and Hepatology, Tokyo Medical and Dental University, 1-5-45 Yushima, Bunkyo-ku, Tokyo 113-8519, Japan

^b Department for Hepatitis Control, Tokyo Medical and Dental University, 1-5-45 Yushima, Bunkyo-ku, Tokyo 113-8519, Japan

^c Department of Internal Medicine, Soka Municipal Hospital, 2-21-1 Soka, Soka City, Saitama 340-8560, Japan

^d Department of Medicine and Molecular Science, Division of Frontier Medical Science, Programs for Biomedical Research, Graduate School of Biomedical Sciences, Hiroshima University, 1-2-3 Kasumi, Minami-ku, Hiroshima City, Hiroshima 734-8551, Japan

^e Department of Virology II, National Institute of Infectious Diseases, 1-23-1 Toyama, Shinjuku-ku, Tokyo 162-8640, Japan

ARTICLE INFO

Article history:

Received 11 March 2010

Returned to author for revision 7 April 2010

Accepted 7 June 2010

Available online 6 July 2010

Keywords:

HCV-JFH1 cell culture

Plaque assay

Cytopathic effect

Adaptive mutations

Human hepatocyte chimeric mice

ABSTRACT

HCV-JFH1 yields subclones that develop cytopathic plaques (Sekine-Osajima Y, et al., Virology 2008; 371:71). Here, we investigated viral amino acid substitutions in cytopathic mutant HCV-JFH1 clones and their characteristics in vitro and in vivo. The mutant viruses with individual C2441S, P2938S or R2985P signature substitutions, and with all three substitutions, showed significantly higher intracellular replication efficiencies and greater cytopathic effects than the parental JFH1 in vitro. The mutant HCV-inoculated mice showed significantly higher serum HCV RNA and higher level of expression of ER stress-related proteins in early period of infection. At 8 weeks post inoculation, these signature mutations had reverted to the wild type sequences. HCV-induced cytopathogenicity is associated with the level of intracellular viral replication and is determined by certain amino acid substitutions in HCV-NS5A and NS5B regions. The cytopathic HCV clones exhibit high replication competence in vivo but may be eliminated during the early stages of infection.

© 2010 Elsevier Inc. All rights reserved.

Introduction

Hepatitis C virus (HCV) is one of the most important pathogens causing liver-related morbidity and mortality (Alter, 1997). Antiviral therapeutic options against HCV have been limited to type I interferons and ribavirin and have yielded unsatisfactory responses (Fried et al., 2002). Given this situation, a precise understanding of the molecular mechanisms of interferon resistance has been a high priority of research in academia and industry.

Molecular analyses of the HCV life cycle, virus–host interactions, and mechanisms of liver cell damage by the virus are not understood

completely, mainly because of the lack of cell culture systems. These problems have been overcome to some extent by the development of the HCV subgenomic replicon (Lohmann et al., 1999) and HCV cell culture systems (Lindenbach et al., 2005; Wakita et al., 2005; Zhong et al., 2005). The HCV-JFH1 strain, which is a genotype 2a clone derived from a Japanese fulminant hepatitis patient and can replicate efficiently in Huh7 cells (Kato, 2001; Kato et al., 2003), has contributed to the establishment of the HCV cell culture system. Furthermore, the Huh7-derived cell lines, Huh-7.5 and Huh-7.5.1 cells, allow production of higher viral titers and have a greater permissivity for HCV (Koutsoudakis et al., 2007; Lindenbach et al., 2005; Zhong et al., 2005). The HCV-JFH1 cell culture system now allows us to study the complete HCV life cycle: virus–cell entry, translation, protein processing, RNA replication, virion assembly and virus release.

HCV belongs to the family *Flaviviridae*. One of the characteristics of the *Flaviviridae* is that they cause cytopathic effects (CPE). The viruses have positive strand RNA genomes of ~10 kilo-bases that encode polyproteins of ~3000 amino acids. These proteins are processed post-translationally by cellular and viral proteases into at least 10 mature proteins (Sakamoto and Watanabe, 2009). The viral non-structural proteins accumulate in the ER and direct genomic replication and viral protein synthesis (Bartenschlager and Lohmann, 2000; Jordan et al., 2002; Mottola et al., 2002). It has been recently

Abbreviations: HCV, hepatitis C virus; CPE, cytopathic effect; ER, endoplasmic reticulum; RdRp, RNA dependent RNA polymerase.

[☆] The authors, K.M., N.S., Y.S., M.N., Y.I., S.A., S.K., K.K., A.K., K.T., M.J., N.H., K.C., T.W. and M.W. declare that there is no conflict of interest.

^{☆☆}This study was supported by grants from Ministry of Education, Culture, Sports, Science and Technology–Japan, the Japan Society for the Promotion of Science, Ministry of Health, Labour and Welfare–Japan, Japan Health Sciences Foundation, and National Institute of Biomedical Innovation.

* Corresponding author. Department of Gastroenterology and Hepatology, Tokyo Medical and Dental University, 1-5-45 Yushima, Bunkyo-ku, Tokyo 113-8519, Japan. Fax: +81 3 5803 0268.

E-mail address: nsakamoto.gast@tmd.ac.jp (N. Sakamoto).

¹ K. M. and N. S. contributed equally to this work.