

tee of Mitsui Memorial Hospital and the Faculty of Medicine, University of Tokyo. Between January 2004 and April 2007, 33914 individuals underwent general health screening, among which information on alcohol consumption was available in 26952. Of these 26952 individuals, information on smoking behavior was further available in 24811, of which 15183 were male individuals and were enrolled in the current study. We were unable to identify any specific reasons to explain why some subjects failed to complete the questionnaire about their smoking and drinking status. Among 15183 individuals enrolled in the current study, data on hepatitis C core antigen (HCCAg) and hepatitis B surface antigen (HBsAg) were available in 14829 individuals (98%), of which 71 were positive for HCCAg and 175 were positive for HBsAg. Individuals who were positive for either type of chronic hepatitis virus infection were significantly older (56 ± 10 years) than hepatitis-negative subjects (53 ± 10 years), although GGT levels were not different between hepatitis-positive (52 ± 52 IU/L) and -negative (58 ± 84 IU/L) individuals. We did not exclude individuals who were taking antihypertensive, antidiabetic, or antidyslipidemic drugs, which might have affected insulin resistance and serum GGT levels, from the current study population.

In Japan, regular health check-ups for employees are a legal requirement; all or most of the costs of the screening are paid for either by the employee's company (about two thirds of individuals attending our institute) or by the subject themselves (about one third of individuals attending our institute). Blood pressure was measured after about 10 min of rest by an automated sphygmomanometer. Individuals were judged to be former smokers and/or former drinkers, if they had stopped cigarette smoking and/or alcohol drinking, respectively, more than one month before their attendance.

Laboratory Analysis

Blood samples were taken from the subjects after an overnight fast. Serum levels of total cholesterol (TC), HDL-cholesterol (HDL-C), and triglycerides (TG) were determined enzymatically. Serum GGT levels were measured enzymatically. Hemoglobin A1c was determined by latex agglutination immunoassay. Plasma glucose was measured by the hexokinase method and serum insulin by enzyme immunoassay. Homeostasis model assessment for insulin resistance (HOMA-IR) was calculated according to the following formula: $\text{HOMA-IR} = [\text{fasting immunoreactive insulin } (\mu\text{U/mL}) \times \text{fasting plasma glucose (FPG; mg/dL)}] / 405$.

Statistical Analysis

Data are expressed as the mean \pm SD unless stated otherwise. Analyses of variance with trend analysis, Dunnett's post-hoc analysis and multiple linear regression analysis were appropriate to assess the statistical significance of differences between groups using computer software, StatView ver. 5.0 (SAS Institute, NC) and Dr. SPSS II (SPSS Inc., Chicago, IL). A value of $p < 0.05$ was significant.

Results

Baseline Characteristics

The baseline characteristics of the study subjects are described in **Table 1**. Among 15183 men, 4534 were former smokers and 416 were former drinkers. Individuals who were former smokers and/or drinkers ($n = 4701$) were significantly older than the remaining 10482 individuals.

GGT and HOMA-IR According to Smoking and Drinking Status

Current smokers who smoked 1-9, 10-19, and 20-39 cigarettes per day were significantly younger than never smokers (**Fig. 1A**). The daily amount of alcohol consumption showed a negative graded association with age. The number of cigarettes smoked showed a positive graded association with GGT (**Fig. 1B**) and, as compared with never smokers, individuals who currently smoked 1-9, 10-19, 20-39, and ≥ 40 cigarettes per day had significantly higher GGT levels (by Dunnett's post-hoc analysis). Similarly, the daily amount of alcohol consumption showed a graded association with GGT, and individuals who drank 1-19, 20-39, 40-59, and ≥ 60 g per day had significantly higher GGT levels than never drinkers (by Dunnett's post-hoc analysis). Individuals who smoked 20-39 and ≥ 40 cigarettes per day had significantly higher HOMA-IR than never-smokers (**Fig. 1C**). On the other hand, as compared with never drinkers, individuals who drank 1-19, 20-39, and 40-59 g alcohol per day had significantly lower HOMA-IR levels (by Dunnett's post-hoc analysis), demonstrating a U-shaped association.

GGT and HOMA-IR According to Cross Strata of Number of Cigarettes Smoked and Alcohol Consumption

In the following analysis, we analyzed the data from 10482 individuals after excluding former smokers and/or former drinkers. The mean GGT levels and HOMA-IR values according to the smoking and drinking category are shown in **Table 2**. Current

Table 1. Baseline characteristics

Variables	Whole	Former smokers and/or drinkers [A]	Except former smokers and drinkers [B]	<i>p</i> value ([A] vs. [B])
N	15,183	4,701	10,482	
Age, years	52.9 ± 10.4	55.6 ± 9.9	51.7 ± 10.4	<0.001
Height, cm	169.6 ± 6.0	169.1 ± 5.9	169.7 ± 6.0	<0.001
Weight, kg	68.3 ± 9.5	68.5 ± 8.9	68.2 ± 9.7	0.117
Body mass index, kg/m ²	23.7 ± 2.8	23.9 ± 2.7	23.6 ± 2.9	<0.001
Systolic blood pressure, mmHg	124.7 ± 18.6	127.6 ± 18.5	123.3 ± 18.4	<0.001
Diastolic blood pressure, mmHg	79.0 ± 11.3	81.0 ± 11.0	78.2 ± 11.3	<0.001
Heart rate, bpm	63.3 ± 9.5	63.4 ± 9.6	63.2 ± 9.5	0.373
LDL-cholesterol, mg/dL	126.7 ± 30.5	127.3 ± 30.0	126.5 ± 30.8	0.112
HDL-cholesterol, mg/dL	55.3 ± 13.4	56.9 ± 13.4	54.6 ± 13.3	<0.001
Triglycerides, mg/dL	133.7 ± 94.2	129.8 ± 83.9	135.5 ± 98.4	0.001
AST, IU/L	23.8 ± 12.1	24.0 ± 10.5	23.7 ± 12.7	0.208
ALT, IU/L	27.3 ± 19.4	26.5 ± 18.8	27.6 ± 19.6	0.001
GGT, IU/L	58.2 ± 82.9	58.3 ± 67.0	58.1 ± 89.1	0.926
Fasting glucose, mg/dL	100.3 ± 20.5	101.7 ± 20.8	99.7 ± 20.4	<0.001
Hemoglobin A1c, %	5.38 ± 0.74	5.41 ± 0.72	5.36 ± 0.75	<0.001
HOMA-IR	1.69 ± 1.52	1.74 ± 1.31	1.67 ± 1.60	0.007
Antihypertensive medication, N (%)	1,909 (12.6)	831 (17.7)	1,078 (10.3)	<0.001
Antidiabetic medication, N (%)	474 (3.1)	169 (3.6)	305 (2.9)	0.026
Antidyslipidemic medication, N (%)	674 (4.4)	276 (5.9)	398 (3.8)	<0.001
Smoking and drinking status				
Never smoker				
Never drinker, N (%)	791 (14.1)	0 (0)	791 (14.3)	
Former drinker, N (%)	90 (1.6)	90 (100)	0 (0)	
Current drinker, N (%)	4,744 (84.3)	0 (0)	4,744 (85.7)	
Former smoker				
Never drinker, N (%)	263 (1.7)	263 (1.7)	0 (0)	
Former drinker, N (%)	249 (1.6)	249 (1.6)	0 (0)	
Current drinker, N (%)	4,022 (26.5)	4,022 (26.5)	0 (0)	
Current smoker				
Never drinker, N (%)	416 (8.3)	0 (0)	416 (8.4)	
Former drinker, N (%)	77 (1.5)	77 (100)	0 (0)	
Current drinker, N (%)	4,531 (90.2)	0 (0)	4,531 (91.6)	

BMI, body mass index; LDL, low density lipoprotein; HDL, high density lipoprotein; AST, aspartate aminotransferase; ALT, alanine aminotransferase; GGT, gamma-glutamyl transpeptidase; HOMA-IR, homeostasis model assessment for insulin resistance

drinking showed a graded association with GGT regardless of the smoking status. Cigarette smoking was also positively associated with GGT in some drinking categories: smoking 10–19 ($p < 0.01$), 20–39 ($p < 0.001$) and ≥ 40 ($p < 0.001$) cigarettes per day was associated with greater GGT values than never smoking in individuals who drank 40–59 g/day, and smoking 20–39 ($p < 0.001$) and ≥ 40 ($p < 0.001$) cigarettes per day was associated with greater GGT values than never smoking in individuals who drank ≥ 60 g/day.

Individuals with alcohol consumption of 1–19, 20–39, or 40–59 g/day had lower HOMA-IR value

than never drinkers, showing a U-shaped association between current drinking and HOMA-IR. This U-shaped relationship was absent or not significant in current smoking of 20–39 or ≥ 40 cigarettes per day (Table 2). Individuals who smoked 20–39 ($p < 0.001$) and ≥ 40 ($p < 0.001$) cigarettes per day had higher HOMA-IR than never smokers (Table 2).

Multiple Linear Regression Analysis

Next, multiple linear regression analysis using GGT and HOMA-IR as a dependent variable and age, BMI, amount of smoking, and alcohol consump-




Fig. 1. Age, GGT, and HOMA-IR according to smoking and drinking status. Bar graphs indicate the mean and 95% CI of age (A), GGT (B), and HOMA-IR. *P* values are for ANOVA trend tests. #, †, and * indicate *p* < 0.05, *p* < 0.01, and *p* < 0.001, respectively, versus never smokers or never drinkers by Dunnett's post-hoc analysis.

Table 2. GGT and HOMA-IR according to smoking and drinking status

Smoking category (cig./day)	Drinking category																		
	0 g/day (never drinker)			1-19 g/day			20-39 g/day			40-59 g/day			≥60g/day						
	N	GGT Mean	95%CI	N	GGT Mean	95%CI	N	GGT Mean	95%CI	N	GGT Mean	95%CI	N	GGT Mean	95%CI	[†] p value			
[§] Overall	10,482	58.1	56.4 59.8	1,207	35.5	33.9 37.2	1,872	41.7	39.8 43.6	3,062	50.1	48.0 52.1	2,957	68.7	65.9 71.6	1,384	95.2	85.4 104.9	<0.001
0 (never smoker)	5,535	49.5	48.2 50.9	791	35.6	33.4 37.8	1,222	43.2	40.6 45.8	1,812	47.2	45.3 49.1	1,258	60.7	57.3 64.2	452	69.3	62.0 76.6	<0.001
1-9	771	59.8	54.7 64.9	48	44.0	32.4 55.6	147	39.0	33.8 44.1	207	51.1	45.4 56.8	253	69.5	57.8 81.2	116	87.1	70.3 103.9	<0.001
10-19	2,201	61.9	58.3 65.5	196	32.9	30.0 35.7	322	39.0	35.3 42.7	616	55.4	47.7 63.2	758	72.9	66.3 79.6	309	89.9	79.9 100.0	<0.001
20-39	1,748	75.6	68.1 83.0	151	35.4	32.1 38.7	158	37.0	33.1 40.9	399	55.1	50.3 59.8	623	77.0	71.4 82.6	417	122.1	93.0 151.3	<0.001
≥40	227	91.1	75.3 107.0	21	39.0	27.0 51.0	23	47.4	36.5 58.4	28	40.9	32.1 49.8	65	93.4	62.1 124.7	90	128.5	97.3 159.6	<0.001
																			[‡] p = 0.006

Smoking category (cig./day)	Drinking category																		
	0 g/day (never drinker)			1-19 g/day			20-39 g/day			40-59 g/day			≥60g/day						
	N	HOMA-IR Mean	95%CI	N	HOMA-IR Mean	95%CI	N	HOMA-IR Mean	95%CI	N	HOMA-IR Mean	95%CI	N	HOMA-IR Mean	95%CI	[†] p value			
[§] Overall	10,482	1.67	1.64 1.70	1,207	1.89	1.79 1.99	1,872	1.57	1.51 1.63	3,062	1.5	1.51 1.59	2,957	1.7	1.62 1.77	1,384	1.8	1.78 1.92	<0.001
0 (never smoker)	5,535	1.62	1.58 1.65	791	1.83	1.70 1.96	1,222	1.58	1.50 1.66	1,812	1.54	1.49 1.59	1,258	1.57	1.51 1.64	452	1.80	1.67 1.94	<0.001
1-9	771	1.67	1.58 1.77	48	2.20	1.52 2.89	147	1.50	1.31 1.69	207	1.54	1.41 1.67	253	1.68	1.53 1.84	116	1.90	1.64 2.15	0.002
10-19	2,201	1.64	1.58 1.69	196	1.96	1.77 2.16	322	1.52	1.38 1.66	616	1.43	1.34 1.51	758	1.70	1.60 1.80	309	1.80	1.65 1.96	<0.001
20-39	1,748	1.82	1.70 1.94	151	2.03	1.74 2.32	158	1.49	1.31 1.66	399	1.74	1.61 1.87	623	1.92	1.62 2.22	417	1.80	1.68 1.92	0.274
≥40	227	2.17	1.97 2.37	21	1.91	1.29 2.52	23	2.71	1.98 3.45	28	1.96	1.46 2.46	65	1.84	1.52 2.16	90	2.39	2.02 2.76	0.068
																			[‡] p = 0.385

[†] p values were for ANOVA trend tests. ^{*} p values are versus never drinkers by Dunnett's post-hoc analysis. [‡] Overall indicates all never or current drinkers, and [§]Overall indicates all never or current smokers. GGT, gamma-glutamyl transpeptidase; HOMA-IR, homeostasis model assessment for insulin resistance.

Table 3. Linear regression analysis using GGT and HOMA-IR as dependent variable

	β	95%CI		Standardized β	<i>p</i> value
Dependent variable: GGT					
Age	-0.57	-1.82	0.68	-0.01	0.372
BMI	2.25	1.79	2.71	0.08	<0.001
Smoking	3.18	2.17	4.19	0.05	<0.001
Alcohol consumption	12.34	11.19	13.49	0.17	<0.001
Dependent variable: HOMA-IR					
Age	0.04	0.01	0.06	0.02	0.001
BMI	0.23	0.22	0.24	0.43	<0.001
Smoking	0.04	0.03	0.06	0.04	<0.001
Alcohol consumption	-0.08	-0.10	-0.06	-0.06	<0.001

For the calculation of β values, age was subdivided into 10-year increments. Alcohol consumption (g/day) corresponding to 0 (never drinker), 1-19, 20-39, 40-59, and ≥ 60 was coded as 0, 1, 2, 3, and 4, respectively, and smoking (cigarettes/day) corresponding to 1-9, 10-19, 20-39, and ≥ 40 was coded as 0, 1, 2, 3, and 4, respectively. GGT, gamma-glutamyl transpeptidase; HOMA-IR, homeostasis model assessment for insulin resistance.

tion as independent variables was performed in 10482 individuals (Table 3). In this model, alcohol consumption (g/day) corresponding to 0 (never drinker); 1-19, 20-39, 40-59, and 60 or more was coded as 0, 1, 2, 3, and 4, respectively, and smoking (cigarettes/day) corresponding to 1-9, 10-19, 20-39, and 40 or more were defined as 0, 1, 2, 3, and 4, respectively. Alcohol consumption was associated positively with GGT, but negatively with HOMA-IR. On the other hand, smoking was found to be associated positively with both GGT and HOMA-IR. When an interaction term between alcohol consumption and smoking was used as additional independent variable, the interaction term was found to be significantly associated with GGT ($p < 0.001$), and showed a borderline significant association with HOMA-IR ($p = 0.059$). The variance inflation factor (VIF) scores of all independent variables tested were less than 10 (data not shown).

Association between GGT and HOMA-IR According to Alcohol Consumption

Next, we investigated whether the mode of association between GGT and HOMA-IR differs according to the amount of alcohol consumption. For this purpose, multiple regression analysis was performed in which age, BMI, and GGT were used as independent variables and HOMA-IR was used as a dependent variable after subdividing individuals according to alcohol consumption (Table 4). GGT was found to be a positive predictive value for HOMA-IR in 19 out of the 25 drinking \times smoking categories. In some combi-

nations of drinking and smoking, such as drinking 0 g/day and smoking 1-9 cig./day, GGT was not a statistically significant predictor of HOMA-IR. This may be in part because the number of subjects with specific drinking and smoking conditions was relatively small.

Discussion

In the current study, by analyzing the data of men who underwent general health screening, except former smokers and/or former drinkers, we observed several points: (1) Alcohol consumption showed a graded association with GGT; (2) In individuals who drank 40 g or more per day, smoking 20 cigarettes or more per day further increased GGT levels (Table 2); (3) alcohol consumption showed a U-shaped association with HOMA-IR, when the daily number of cigarettes smoked was less than 20 per day; (4) Individuals who smoked 20-39 and ≥ 40 cigarette per day had higher HOMA-IR than never smokers (Table 2); (5) GGT was found to be a positive predictive value of HOMA-IR in 19 out of the 25 drinking \times smoking categories, and GGT was not a significant negative predictor of HOMA-IR regardless of the drinking or smoking status. These data collectively indicate that, although current drinking may increase GGT and reduce insulin resistance, GGT can be utilized as a marker of insulin resistance regardless of the drinking status.

Many studies have shown that serum GGT is a biomarker of increased alcohol consumption^{1-4, 22}; however, GGT is known to be affected by other con-

Table 4. Linear regression analysis using HOMA-IR as dependent variable

	β	95%CI	Standardized β	p value		β	95%CI	Standardized β	p value	
Current smoking - 0 cig./day (never smoker)										
0 g/day (never drinker)	BMI	0.20	0.16	0.24	<0.001	BMI	0.27	0.18	0.37	<0.001
	GGT	0.07	0.03	0.11	0.001	GGT	0.17	0.04	0.30	0.010
1-19 g/day	BMI	0.22	0.19	0.24	<0.001	BMI	0.12	0.06	0.18	<0.001
	GGT	0.04	0.02	0.05	<0.001	GGT	0.07	0.00	0.14	0.036
20-39 g/day	BMI	0.21	0.19	0.22	<0.001	BMI	0.22	0.18	0.26	<0.001
	GGT	0.03	0.02	0.05	<0.001	GGT	0.04	0.02	0.07	<0.001
40-59 g/day	BMI	0.18	0.16	0.20	<0.001	BMI	0.28	0.18	0.37	<0.001
	GGT	0.04	0.03	0.04	<0.001	GGT	0.02	-0.02	0.06	0.395
≥ 60 g/day	BMI	0.26	0.23	0.30	<0.001	BMI	0.22	0.19	0.26	<0.001
	GGT	0.02	0.01	0.04	0.001	GGT	0.01	0.00	0.01	<0.001
Current smoking - 1-9 cig./day										
0 g/day (never drinker)	BMI	0.44	0.21	0.67	<0.001	BMI	-0.07	-0.29	0.16	0.551
	GGT	0.13	-0.02	0.28	0.084	GGT	0.41	0.18	0.64	0.002
1-19 g/day	BMI	0.24	0.17	0.32	<0.001	BMI	-0.07	-0.29	0.16	0.551
	GGT	0.07	0.01	0.12	0.016	GGT	0.41	0.18	0.64	0.002
20-39 g/day	BMI	0.18	0.13	0.23	<0.001	BMI	0.15	0.02	0.28	0.028
	GGT	0.01	-0.02	0.04	0.396	GGT	0.10	-0.15	0.34	0.425
40-59 g/day	BMI	0.23	0.18	0.27	<0.001	BMI	0.18	0.06	0.30	0.003
	GGT	0.01	0.00	0.03	0.049	GGT	0.01	-0.02	0.03	0.559
≥ 60 g/day	BMI	0.27	0.20	0.34	<0.001	BMI	0.28	0.18	0.38	<0.001
	GGT	0.04	0.02	0.06	<0.001	GGT	0.05	0.03	0.07	<0.001
Current smoking - 10-19 cig./day										
0 g/day (never drinker)	BMI	0.16	0.10	0.22	<0.001	BMI	-0.12	-0.12	0.42	0.551
	GGT	0.27	0.19	0.36	<0.001	GGT	0.41	0.18	0.64	0.002
1-19 g/day	BMI	0.22	0.18	0.26	<0.001	BMI	-0.07	-0.29	0.16	0.551
	GGT	0.08	0.05	0.12	<0.001	GGT	0.41	0.18	0.64	0.002
20-39 g/day	BMI	0.18	0.15	0.21	<0.001	BMI	0.15	0.02	0.28	0.028
	GGT	0.00	-0.01	0.01	0.583	GGT	0.10	-0.15	0.34	0.425
40-59 g/day	BMI	0.25	0.22	0.27	<0.001	BMI	0.18	0.06	0.30	0.003
	GGT	0.02	0.01	0.03	<0.001	GGT	0.01	-0.02	0.03	0.559
≥ 60 g/day	BMI	0.22	0.18	0.26	<0.001	BMI	0.28	0.18	0.38	<0.001
	GGT	0.03	0.01	0.04	<0.001	GGT	0.05	0.03	0.07	<0.001

Standardized β values are estimates resulting from analysis performed on into standardized variables. For the calculation of β values, BMI was subdivided into 1 kg/m² increments, and GGT into 10 IU/L increments. Age, BMI, and GGT were used as independent variables. BMI, body mass index; GGT, gamma-glutamyl transpeptidase.

ditions, such as smoking, obesity, and hepatic steatosis^{23, 24}). Evidence is accumulating that higher serum GGT levels may be associated with an increased incidence of cardiovascular events⁵), metabolic syndrome and diabetes^{8, 25, 26}); therefore, more attention has been paid recently to this liver enzyme. It is possible that the association between GGT and various disorders observed in previous studies may be mediated, in part, by enhanced insulin resistance in subjects with increased GGT levels.

Although mild to moderate alcohol consumption may increase GGT, it may improve insulin sensitivity^{18, 27}), leading to a reduction in the prevalence of metabolic syndrome¹⁷). This finding is in contrast to the observation that cigarette smoking will not improve insulin resistance, even in light smokers¹⁴). As alcohol consumption has opposite effects on GGT and insulin resistance, the mode of association between GGT and HOMA-IR might differ according to the drinking status; however, only a few studies have analyzed the relationship between GGT and insulin resistance in various drinking conditions. Yokoyama and colleagues reported that GGT is associated with increased insulin resistance in non-drinkers²⁸) and light drinkers, but not in heavy drinkers²⁹), a finding that supports the notion that the mode of association between GGT and HOMA-IR differs according to the drinking status. Yamada *et al.* have reported that HOMA-IR rose with increasing serum GGT in both alcohol consumers and non-consumers, and HOMA-IR values corresponding to all serum GGT levels were lower in alcohol consumers than in non-consumers³⁰). A recent study indicated that cigarette smoking may also affect both GGT and insulin resistance independent of the drinking status, and cigarette smoking and alcohol intake may have a synergistic impact on GGT¹³). Smoking status should also be considered when assessing the impact of alcohol intake on the association between GGT and insulin resistance; however, to our knowledge, no previous studies have investigated the relationship between GGT and insulin resistance after stratifying both the drinking status and smoking status, as in the current study.

We found that in 19 of the 25 subgroups divided according to smoking and drinking status, GGT was found to be a positive predictive value of HOMA-IR, which indicates that increased GGT is associated with enhanced insulin resistance regardless of the smoking and drinking status. From this type of cross-sectional study, we cannot conclude whether there is any causal or resultant relationship between GGT and HOMA-IR. A recent study showed that GGT may play a causal role in promoting insulin resistance, pre-

sumably by enhancing oxidative stress^{31, 32}) and hepatic steatosis³³). Whether a change in HOMA-IR would result in a predicted change in GGT should be investigated in future longitudinal studies.

Our study has some limitations. First, we did not take into account coffee intake, which might affect GGT level²). Second, as the prevalence of smokers was low, we did not analyze the data of female subjects. Third, the number of daily cigarettes and alcohol consumption solely reflected the amount that was being consumed at one time, and disregarded the frequency of smoking or drinking consumption. Therefore, this estimation of smoking and drinking quantity was not equal to the mean daily number of cigarettes smoked and the amount of alcohol consumption, except in every-day smokers and drinkers, respectively. We performed such an analysis because the frequency of smoking (or drinking) was reported as a category, two or three times per week, for example; therefore, it was technically difficult to estimate the mean daily number of cigarettes smoked or the alcohol consumption. In the future, however, the frequency of drinking and smoking should also be considered in such an analysis. Fourth, we did not exclude individuals who were taking antihypertensive and/or antidiabetic drugs, which may have affected serum GGT and HOMA-IR values.

In summary, alcohol consumption showed a graded positive association with GGT and a U-shaped negative association with HOMA-IR. Cigarette smoking may further increase GGT levels in individuals who are current drinkers and drink 20 g or more per day. In 19 of the 25 drinking × smoking categories, GGT was found to be a positive predictive value of HOMA-IR, and GGT was not a significant negative predictor of HOMA-IR, regardless of the drinking or smoking status. These data indicate a positive association between GGT and insulin resistance also in current drinkers.

Acknowledgements

The work was supported in part by a grant from the Smoking Research Foundation, Chiyoda Mutual Life Foundation, a St Luke's Grant for the Epidemiological Research, Daiwa Securities Health Foundation, a Gerontology Research Grant from Kowa Life Science Foundation, the Foundation for Total Health Promotion, and the Gout Research Foundation of Japan.

References

- 1) Robinson D, Monk C, Bailey A: The relationship between

- serum gamma-glutamyl transpeptidase level and reported alcohol consumption in healthy men. *J Stud Alcohol*, 1979; 40: 896-901
- 2) Poikolainen K, Vartiainen E: Determinants of gamma-glutamyltransferase: positive interaction with alcohol and body mass index, negative association with coffee. *Am J Epidemiol*, 1997; 146: 1019-1024
 - 3) Stewart SH: Racial and ethnic differences in alcohol-associated aspartate aminotransferase and gamma-glutamyltransferase elevation. *Arch Intern Med*, 2002; 162: 2236-2239
 - 4) Yamada Y, Noborisaka Y, Suzuki H, Ishizaki M, Yamada S: Alcohol consumption, serum gamma-glutamyltransferase levels, and coronary risk factors in a middle-aged occupational population. *J Occup Health*, 2003; 45: 293-299
 - 5) Fraser A, Harris R, Sattar N, Ebrahim S, Smith GD, Lawlor DA: Gamma-glutamyltransferase is associated with incident vascular events independently of alcohol intake: analysis of the British Women's Heart and Health Study and Meta-Analysis. *Arterioscler Thromb Vasc Biol*, 2007; 27: 2729-2735
 - 6) Jousilahti P, Rastenyte D, Tuomilehto J: Serum gamma-glutamyl transferase, self-reported alcohol drinking, and the risk of stroke. *Stroke*, 2000; 31: 1851-1855
 - 7) Hu G, Tuomilehto J, Pukkala E, Hakulinen T, Antikainen R, Vartiainen E, Jousilahti P: Joint effects of coffee consumption and serum gamma-glutamyltransferase on the risk of liver cancer. *Hepatology*, 2008; 48: 129-136
 - 8) Nakanishi N, Suzuki K, Tatara K: Serum gamma-glutamyltransferase and risk of metabolic syndrome and type 2 diabetes in middle-aged Japanese men. *Diabetes Care*, 2004; 27: 1427-1432
 - 9) Lee DH, Gross MD, Steffes MW, Jacobs DR Jr: Is serum gamma-glutamyltransferase a biomarker of xenobiotics, which are conjugated by glutathione? *Arterioscler Thromb Vasc Biol*, 2008; 28: e26-28; author reply e29
 - 10) Kono S, Shinci K, Imanishi K, Todoroki I, Hatsuse K: Coffee and serum gamma-glutamyltransferase: a study of self-defense officials in Japan. *Am J Epidemiol*, 1994; 139: 723-727
 - 11) Nakanishi N, Nakamura K, Nakajima K, Suzuki K, Tatara K: Coffee consumption and decreased serum gamma-glutamyltransferase: a study of middle-aged Japanese men. *Eur J Epidemiol*, 2000; 16: 419-423
 - 12) Li M, Campbell S, McDermott R: gamma-Glutamyltransferase, Obesity, Physical Activity, and the Metabolic Syndrome in Indigenous Australian Adults. *Obesity (Silver Spring)*, 2009; 17: 809-813
 - 13) Breitling LP, Raum E, Muller H, Rothenbacher D, Brenner H: Synergism between smoking and alcohol consumption with respect to serum gamma-glutamyltransferase. *Hepatology*, 2009; 49: 802-808
 - 14) Ishizaka N, Ishizaka Y, Toda E, Hashimoto H, Nagai R, Yamakado M: Association between cigarette smoking, metabolic syndrome, and carotid arteriosclerosis in Japanese individuals. *Atherosclerosis*, 2005; 181: 381-388
 - 15) Oh SW, Yoon YS, Lee ES, Kim WK, Park C, Lee S, Jeong EK, Yoo T: Association between cigarette smoking and metabolic syndrome: the Korea National Health and Nutrition Examination Survey. *Diabetes Care*, 2005; 28: 2064-2066
 - 16) Goude D, Fagerberg B, Hulthe J: Alcohol consumption, the metabolic syndrome and insulin resistance in 58-year-old clinically healthy men (AIR study). *Clin Sci (Lond)*, 2002; 102: 345-352
 - 17) Freiberg MS, Cabral HJ, Heeren TC, Vasani RS, Curtis Ellison R: Alcohol consumption and the prevalence of the Metabolic Syndrome in the US: a cross-sectional analysis of data from the Third National Health and Nutrition Examination Survey. *Diabetes Care*, 2004; 27: 2954-2959
 - 18) Joosten MM, Beulens JW, Kersten S, Hendriks HF: Moderate alcohol consumption increases insulin sensitivity and ADIPOQ expression in postmenopausal women: a randomised, crossover trial. *Diabetologia*, 2008; 51: 1375-1381
 - 19) Yamamoto A, Temba H, Horibe H, Mabuchi H, Saito Y, Matsuzawa Y, Kita T, Nakamura H: Life style and cardiovascular risk factors in the Japanese population--from an epidemiological survey on serum lipid levels in Japan 1990 part 1: influence of life style and excess body weight on HDL-cholesterol and other lipid parameters in men. *J Atheroscler Thromb*, 2003; 10: 165-175
 - 20) Kang YH, Min HK, Son SM, Kim IJ, Kim YK: The association of serum gamma glutamyltransferase with components of the metabolic syndrome in the Korean adults. *Diabetes Res Clin Pract*, 2007; 77: 306-313
 - 21) Shin JY, Chang SJ, Shin YG, Seo KS, Chung CH: Elevated serum gamma-glutamyltransferase levels are independently associated with insulin resistance in non-diabetic subjects. *Diabetes Res Clin Pract*, 2009; 84: 152-157
 - 22) Lamy J, Baglin MC, Weill J, Aron E: Serum gamma-glutamyl-transpeptidase and alcoholism. Diagnosis and control of withdrawal. *Nouv Presse Med*, 1975; 4: 487-490
 - 23) Sato KK, Hayashi T, Nakamura Y, Harita N, Yoneda T, Endo G, Kambe H: Liver enzymes compared with alcohol consumption in predicting the risk of type 2 diabetes: the Kansai Healthcare Study. *Diabetes Care*, 2008; 31: 1230-1236
 - 24) Benini F, Pigozzi MG, Baisini O, Romanini L, Ahmed H, Pozzi A, Ricci C, Lanzini A: Increased serum gamma-glutamyl-transpeptidase concentration is associated with nonalcoholic steatosis and not with cholestasis in patients with chronic hepatitis C. *J Gastroenterol Hepatol*, 2007; 22: 1621-1626
 - 25) Ruttman E, Brant LJ, Concin H, Diem G, Rapp K, Ulmer H: Gamma-glutamyltransferase as a risk factor for cardiovascular disease mortality: an epidemiological investigation in a cohort of 163,944 Austrian adults. *Circulation*, 2005; 112: 2130-2137
 - 26) Shankar A, Li J, Klein BE, Javier Nieto F, Klein R: Serum gamma-glutamyltransferase level and peripheral arterial disease. *Atherosclerosis*, 2008; 199: 102-109
 - 27) Fueki Y, Miida T, Wardaningsih E, Ito M, Nakamura A, Takahashi A, Hanyu O, Tsuda A, Saito H, Hama H, Okada M: Regular alcohol consumption improves insulin resistance in healthy Japanese men independent of obesity. *Clin Chim Acta*, 2007; 382: 71-76
 - 28) Yokoyama H, Hirose H, Moriya S, Saito I: Significant correlation between insulin resistance and serum gamma-glutamyl transpeptidase (gamma-GTP) activity in non-

- drinkers. *Alcohol Clin Exp Res*, 2002; 26: 91S-94S
- 29) Moriya S, Yokoyama H, Hirose H, Ishii H, Saito I: Correlation between insulin resistance and gamma-glutamyl transpeptidase sensitivity in light drinkers. *Alcohol Clin Exp Res*, 2003; 27: 52S-57S
- 30) Yamada Y, Noborisaka Y, Ishizaki M, Tsuritani I, Honda R, Yamada S: Alcohol consumption, homeostasis model assessment indices and blood pressure in middle-aged healthy men. *J Hum Hypertens*, 2004; 18: 343-350
- 31) Whitfield JB: Gamma glutamyl transferase. *Crit Rev Clin Lab Sci*, 2001; 38: 263-355
- 32) Lee DH, Blomhoff R, Jacobs DR Jr: Is serum gamma glutamyltransferase a marker of oxidative stress? *Free Radic Res*, 2004; 38: 535-539
- 33) Nannipieri M, Gonzales C, Baldi S, Posadas R, Williams K, Haffner SM, Stern MP, Ferrannini E: Liver enzymes, the metabolic syndrome, and incident diabetes: the Mexico City diabetes study. *Diabetes Care*, 2005; 28: 1757-1762



ORIGINAL ARTICLE

Effects of the AT₁ receptor blocker losartan and the calcium channel blocker benidipine on the accumulation of lipids in the kidney of a rat model of metabolic syndrome

Nobukazu Ishizaka¹, Makiko Hongo¹, Gen Matsuzaki¹, Kyoko Furuta¹, Kan Saito¹, Ryota Sakurai¹, Aiko Sakamoto¹, Kazuhiko Koike² and Ryozyo Nagai¹

Unfavorable lipid accumulation may occur in the kidneys in the presence of metabolic syndrome and diabetes. The aim of this study was to investigate whether excess lipids would accumulate in the kidneys of Otsuka Long-Evans Tokushima Fatty (OLETF) rats, an animal model of metabolic syndrome. From 34 weeks of age, OLETF rats were treated orally with a calcium channel blocker, benidipine (3 mg kg⁻¹ per day), or an AT₁ receptor blocker, losartan (25 mg kg⁻¹ per day), for 8 weeks. Blood pressure was slightly but significantly higher in the untreated OLETF rats (149 ± 4 mm Hg) than in Long-Evans Tokushima Otsuka (LETO) rats (136 ± 2 mm Hg), and both losartan (135 ± 3 mm Hg) and benidipine (138 ± 3 mm Hg) reduced blood pressure in OLETF rats to a level comparable to that in LETO rats. Tissue content of triglycerides (TG) was greater in OLETF rats than in LETO rats (6.24 ± 3.77 and 2.85 ± 1.32 μg mg⁻¹ · tissue, respectively), and both losartan and benidipine reduced these values. Histological analysis showed lipid droplets in tubular cells in which increased dihydroethidium fluorescence was present. Expression of peroxisome proliferator-activated receptor-α, PGC-1α and uncoupling protein-2 was found to be higher in OLETF rats than in LETO rats; however, the expression of these genes was not altered by treatment with either antihypertensive drug. In contrast, both losartan and benidipine increased the amount of total and phosphorylated forms of AMP kinase and the expression of carnitine palmitoyltransferase-1 (CPT-1). In conclusion, treatment of OLETF rats with losartan and benidipine reduced the tissue content of TG, decreased the production of superoxide and regulated the expression of genes related to fatty acid oxidation such as AMP-activated protein kinase and CPT-1 in the kidneys.

Hypertension Research (2010) 33, 263–268; doi:10.1038/hr.2009.224; published online 8 January 2010

Keywords: calcium channel blocker; diabetes; kidney; lipotoxicity; renin angiotensin system

INTRODUCTION

Unfavorable lipid accumulation in the kidneys may occur in animal models of diabetes,^{1–3} aging,⁴ diet-induced obesity^{5,6} and nephrectomy.⁷ Although the precise mechanism by which lipid content is increased in the kidneys is still not fully elucidated, it may include upregulation of lipogenic gene expression in the kidneys^{1,7,8} and uptake of filtered albumin-bound fatty acids by renal tubular cells when increased urinary excretion of albumin is present.⁹ Transfer of lipogenic genes induced deposition of lipids and upregulation of fibrosis-related gene expression in the kidneys of diabetic animals, whereas lipogenic gene knockdown had the opposite effect,^{1,8} suggesting that the accumulation of excessive lipid in the kidneys is one factor in the pathophysiological process of diabetic nephropathy.⁶

We reported earlier that administration of angiotensin II upregulates the expression of lipogenic genes and increases lipid content in the kidneys,¹⁰ suggesting that activation of the renin angiotensin system may have a role in lipid accumulation in the kidney. It has been reported that lipid content is increased in the liver¹¹ and pancreas¹² of OLETF rats. In this study, therefore, we investigated whether excessive lipid accumulation occurs in the kidneys of Otsuka Long-Evans Tokushima Fatty (OLETF) rats, which exhibit features of metabolic syndrome,¹³ and if present, whether angiotensin II receptor blockers and calcium channel blockers could exert similar effects on renal lipid content in OLETF rats. We used benidipine as the calcium channel blocker because this drug has been reported to reduce the extent of proteinuria in OLETF rats.¹⁴

¹Department of Cardiovascular Medicine, University of Tokyo Graduate School of Medicine, Tokyo, Japan and ²Department of Gastroenterology, University of Tokyo Graduate School of Medicine, Tokyo, Japan

Correspondence: Dr N Ishizaka, Department of Cardiovascular Medicine, University of Tokyo Graduate School of Medicine, Bunkyo-ku, Hongo 7-3-1, Tokyo 113-8655, Japan. E-mail: nobuishizaka-ky@umin.ac.jp

Received 17 August 2009; revised 9 November 2009; accepted 29 November 2009; published online 8 January 2010

METHODS

Animals

The experiments were performed in accordance with the guidelines for animal experimentation approved by the Animal Center for Biomedical Research, Faculty of Medicine, University of Tokyo. Male OLETF and age-matched Long-Evans Tokushima Otsuka (LETO) rats, a genetic control, were obtained from the Tokushima Research Institute (Otsuka Pharmaceutical, Tokushima, Japan) and maintained under constant temperature and lighting conditions with free access to food and water. At 34 weeks of age, the OLETF rats were given 3 mg kg⁻¹ benidipine or 25 mg kg⁻¹ losartan per day orally, which was continued for 8 weeks. One day before sacrifice, the rats were kept in a metabolic cage, and urine was collected for 24 h under fasting conditions. Systolic blood pressure and heart rate were measured in conscious rats by tail-cuff plethysmography (BP-98A, Softron, Tokyo, Japan).

Measurement of lipid content in the serum and kidney

Serum levels of total cholesterol (TC), triglycerides (TG) and non-esterified fatty acid were measured by enzymatic methods (SRL, Tokyo, Japan). Contents of TG and TC in the kidney were measured from homogenate extracts by enzymatic colorimetric determination using the Triglyceride-E Test, the Cholesterol-E Test and the Free cholesterol-E Test, respectively (Wako Pure Chemicals, Osaka, Japan).

Histological analysis

Oil red O staining was performed on sections of unfixed, freshly frozen kidney samples (3 µm in thickness). For semi-quantification of lipid deposition, images of each specimen stained with oil red O were taken with an Olympus BX51 microscope and a DP12 digital camera system (Olympus, Tokyo, Japan). Five images taken in the cortical region of each sample were analyzed. The ratio of the areas of lipid deposition to the total tissue region area was calculated using Adobe Photoshop image analysis software (Adobe Systems, San Jose, CA, USA). *In situ* superoxide production was estimated using the oxidative fluorescent dye dihydroethidium (DHE) in unfixed frozen kidney specimens as described earlier.¹⁰ Images were obtained from at least five fields in each section, and signal intensity was presented as a percentage of that in OLETO rats.

Western blot analysis

Western blot analysis was performed as described earlier.¹⁵ Antibodies against total and phosphorylated forms of AMP-activated protein kinase (AMPK) (Cell Signaling Technology, Danvers, MA, USA), sterol regulatory element-binding protein (SREBP)-1 (Santa Cruz Biotechnology, Santa Cruz, CA, USA) and total and phosphorylated forms of acetyl-CoA carboxylase (ACC) (Cell Signaling Technology) were used at a dilution of 1/1000.

Real time RT-PCR

The mRNA expression of lipid metabolism-related genes was analyzed by real-time quantitative PCR performed using a LightCycler together with Hybri-Probe technology (Roche Diagnostics, Basel, Switzerland). The expression of target genes was normalized to the mRNA expression of the endogenous control, glyceraldehyde-3-phosphate dehydrogenase. The target genes were SREBP-1c, fatty acid synthase (FAS), 3-hydroxy-3-methylglutaryl coenzyme A reductase, peroxisome proliferator-activated receptor (PPAR)-γ, PPAR-α, PPAR-γ coactivator (PGC)-1α, CD36, carnitine palmitoyltransferase (CPT)-1 and uncoupling protein (UCP)-2. The forward and reverse primers used were described earlier.¹⁶

Statistical analysis

Data are expressed as mean ± s.e.m. ANOVA and Kruskal–Wallis analyses followed by a *post hoc* multiple comparison test were performed using the statistical analysis software Dr. SPSS II (SPSS Inc., Chicago, IL, USA). A value of *P* < 0.05 was taken to be statistically significant.

RESULTS

Characteristics of the experimental animals

Body weight, blood pressure, heart rate and blood levels of lipids and glucose in each group have been described elsewhere.¹⁷ Blood pressure was slightly but significantly higher in the untreated OLETF rats (149 ± 4 mm Hg, *n* = 11, *P* = 0.012) than in LETO rats (136 ± 2 mm Hg, *n* = 11), and both losartan (135 ± 3 mm Hg, *n* = 6) and benidipine (138 ± 3 mm Hg, *n* = 11) reduced blood pressure in OLETF rats to a level comparable to that in LETO rats. Treatment of OLETF rats with either antihypertensive drug had no significant effect on circulating levels of triglyceride and glucose, which were higher in OLETF than in LETO rats. Compared with LETO rats, kidney weight was greater in the untreated OLETF rats and the OLETF rats treated with either losartan or benidipine, but no significant difference was observed in creatinine clearance among the groups examined (Figure 1). Urinary protein excretion was greater in OLETF rats than in LETO rats, and both losartan and benidipine reduced proteinuria to a similar extent.

Accumulation of lipids in the kidney

The content of TG in the kidney was significantly greater in untreated OLETF rats than in LETO rats, and both losartan and benidipine treatment reduced renal TG content in the OLETF rats (Figure 2a). The content of TC in the kidney was not significantly different between LETO and untreated OLETF rats; however, both antihyper-




Figure 1 Kidney weight, creatinine clearance and proteinuria in LETO rats and untreated and antihypertensive drug-treated OLETF rats. Absolute values of kidney weight (a) and kidney weight expressed per 100 g body weight (b). Creatinine clearance (c) and daily excretion of urinary protein (d). Summary of data from four to six rats in each group.




Figure 2 Tissue lipid content in the kidneys of LETO rats and untreated and antihypertensive drug-treated OLETF rats. Content of triglycerides (TG) (a) and total cholesterol (TC) (b) in the kidneys of LETO ($n=4$), OLETF ($n=8$), OLETF/Los ($n=8$) and OLETF/Ben ($n=8$) groups is shown.




Figure 3 Accumulation of lipids in the kidney. (a, b, e) Kidney sections from LETO rats. (c, d, f-h) Kidney sections from untreated OLETF rats. (i, j) Kidney sections from OLETF rats treated with losartan (i) and benidipine (j). (a, c, e-g, i, j) Oil red O staining. (b, d, h) Hematoxylin and eosin staining; (a, b), (c, d) and (g, h) are serial sections. Lipid droplets were observed in the tubular (c, g), but not glomerular (f, arrowhead), regions of kidney from untreated OLETF rats. In some hematoxylin and eosin-stained specimens, lipid droplets could be identified by unstained small vesicles (g, h, arrows). The extent of lipid accumulation in the tubular cells was diminished by treatment of OLETF rats with either losartan (i) or benidipine (j). Scale bars indicate 100 μm . (k) Semi-quantification of the oil red O-stained area. Summary of data from five to seven experiments in each group. Kruskal-Wallis analyses followed by a *post hoc* multiple comparison test were performed. * $P<0.05$ and # $P<0.01$ versus untreated OLETF rats.

tensive drugs reduced renal TC content in OLETF rats (Figure 2b). Histological analysis showed that only a trace amount of oil red O-positive lipid droplets was present in the kidneys of LETO rats (Figure 3). By contrast, increased lipid droplets were observed in the tubular and interstitial regions of untreated OLETF rats. Some lipid droplets appeared as small cavities on hematoxylin and eosin-stained frozen specimens (Figures 3g and h, arrows). Areas of oil red O-positive deposits were significantly reduced by the treatment of OLETF rats with losartan or benidipine (Figures 3i-k). The correlation coefficients between tissue TG content and the extent of proteinuria and between the oil red O stained area and the extent of proteinuria were 0.40 ($P<0.05$, $n=33$) and 0.65 ($P<0.001$, $n=36$), respectively.

Localization of superoxide

Fluorescent signals on DHE staining were greater in untreated OLETF rats than in LETO rats and were reduced by treatment with either losartan or benidipine (Figure 4). In untreated OLETF rat kidney,

DHE signals were found to be increased in tubular epithelial (Figures 4e-j, arrowheads) and vascular wall cells (Figures 4e-g, arrows), the former of which contained lipid droplets.

Regulation of genes related to lipid metabolism

The expression of mature SREBP-1 protein did not significantly differ among the four groups examined (Figure 5). Compared with LETO rats, expression of both total and phosphorylated forms of AMPK α was increased in OLETF rats treated with either losartan or benidipine, although it was not increased in untreated OLETF rats. The expression of total ACC protein was unaffected by losartan or benidipine in OLETF rats; however, treatment with either antihypertensive drug significantly increased the amount of the phosphorylated form of ACC.

Among the genes tested, mRNA expression of SREBP-1c, FAS, 3-hydroxy-3-methylglutaryl coenzyme A reductase, PPAR- γ , LDL-r and CD36 did not significantly differ between untreated OLETF and




Figure 4 Localization of lipid droplets and superoxide in kidney sections. (a) Kidney sections from LETO rats. (b, e–j) Kidney sections from untreated OLETF rats. (c, d) Kidney sections from OLETF rats treated with losartan (c) and benidipine (d). (a–d, f, i) Dihydroethidium (DHE) staining. (e, h) Phase contrast (PC) microscopic images. (j) PC microscopic image overlaid with DHE stained images. (g) Oil red O staining; (e, f) and (h–j) are the same section; (e–g) are serial sections. DHE signals were more intense in OLETF kidneys (b) than in LETO kidneys (a), and both losartan (c) and benidipine (d) reduced DHE signal intensity. Granular droplets could be observed in the tubular regions, presumably lipid droplets, by PC imaging (e, arrowheads) of the unstained specimens, and DHE-stained superoxide was increased in these regions and in vascular wall cells (arrow, e, f). Oil red O staining (h) of the serial specimen confirmed that granular materials were lipid droplets. Higher magnification imaging demonstrated that DHE signals were increased in cells with granular droplets. Scale bars indicate 100 μ m. (k) Semi-quantification of the DHE signals. Summary of data from five to seven experiments in each group. Kruskal–Wallis analyses followed by a *post hoc* multiple comparison test were performed. # $P < 0.01$ versus untreated OLETF rats.




Figure 5 Western blot analysis. Western blot of sterol regulatory element-binding protein-1 (SREBP-1), AMP-activated protein kinase α (AMPK), the phosphorylated (activated) form of AMPK α (P-AMPK), acetyl-CoA carboxylase (ACC) and phosphorylated ACC. (a–f) Summary of data from four to six experiments in each group. * $P < 0.01$ and # $P < 0.05$ versus untreated OLETF rats by Dunnett's *post hoc* analysis.

LETO rats (Figure 6). By contrast, mRNA expression of PPAR- α and UCP-2 was greater, whereas that of PGC-1 α was lower, in untreated, losartan-treated and benidipine-treated OLETF rats than in LETO rats. Although the mRNA expression of CPT-1 did not differ between untreated OLETF and LETO rats, treatment of OLETF rats with either losartan or benidipine significantly increased CPT-1 expression.

DISCUSSION

In this study, we found that TG and TC contents were increased in the kidneys of untreated OLETF rats compared with LETO rats. In the kidneys of OLETF rats, oil red O-positive lipid droplets were observed mainly in tubular epithelial cells, in which increased superoxide was present. Treatment of OLETF rats with either losartan or benidipine,




Figure 6 mRNA expression and regulation of lipid metabolism-related genes. (a) Sterol regulatory element-binding protein-1c (SREBP-1c). (b) Fatty acid synthase (FAS). (c) 3-hydroxy-3-methylglutaryl coenzyme A reductase (HMG-CoAR). (d) Peroxisome proliferator-activated receptor (PPAR)-γ. (e) PPAR-α. (f) PPAR-γ coactivator (PGC)-1α. (g) LDL receptor. (h) CD36. (i) Carnitine palmitoyltransferase (CPT)-1. (j) Uncoupling protein (UCP)-2. Summary of data from six to ten experiments in each group * $P < 0.01$ and # $P < 0.05$ versus untreated OLETF rats by Dunnett's *post hoc* analysis.

both of which lowered blood pressure to a similar extent, reduced tissue TG content, oil red O-positive deposits, superoxide signals and urinary protein excretion in the kidneys to a similar extent.

The mechanisms underlying lipid accumulation in the kidney in various animal models and in humans have not been fully elucidated; however, the presumed mechanisms include the regulation of genes related to the uptake, biosynthesis, catabolism and efflux of lipids^{1,3,4,10,18} and the uptake of fatty acids carried on filtered albumin by the renal tubules.^{7,9} What is the possible mechanism underlying the reduction in the renal lipid content of OLETF rats caused by both losartan and benidipine? We reported earlier that long-term administration of angiotensin II upregulates the expression of SREBP-1 and FAS and increases the renal content of lipids, which was suppressed by losartan. This effect of losartan may, at least in part, be independent of its depressor effect.¹⁹ Therefore, it is possible that benidipine may also reduce renal lipid content through a mechanism independent of its depressor effect. To our knowledge, however, no earlier studies have shown that a calcium channel blocker can regulate the expression of certain lipogenic gene and/or reduce tissue lipid content. Toblli *et al.*²⁰ reported that renin angiotensin system inhibition, but not calcium channel blockade, suppressed lipid deposition in the heart²⁰ and liver²¹ in obese Zucker rats.

In this study, the expression of SREBP-1 and FAS was not upregulated in the kidneys of OLETF rats compared with LETO rats, and neither losartan nor benidipine reduced the expression of these genes in the kidneys of OLETF rats. Therefore, it can be stated that the mechanism by which lipid content increased in the kidneys of OLETF rats may be, at least in part, different from that in angiotensin II-infused rats. This could be the reason why losartan was not more effective than benidipine in terms of reducing lipid content in the kidney.

As enhanced urinary albumin excretion may lead to the subsequent tubular absorption of lipid-bound albumin,^{7,9} another possibility is that both losartan and benidipine reduced renal lipid content by their anti-proteinuric effect. Several studies have reported that benidipine may have a greater anti-proteinuric effect than other CCBs such as amlodipine,²² and the renoprotective effect of benidipine may be, in part, mediated by the preservation of an essential cofactor of nitric oxide synthase, (6R)-5,6,7,8-tetrahydrobiopterin.¹⁴ As both losartan and benidipine reduced the extent of proteinuria, both drugs might reduce the transport of the albumin-bound fatty acid to tubular cells. The close relationship between the extent of lipid deposition and proteinuria observed in this study may support this notion. Furthermore, the reduction of blood pressure *per se* might have a role in the modulation of lipid content in the kidney.

In this study, tubular cells that contained lipid droplets were positive for DHE signals (Figure 4). A spatial relationship between lipid droplets and superoxide was also observed in the kidneys and heart of angiotensin II-induced hypertensive animals,^{10,16} indicating that these two phenomena have a relationship²³ under conditions of hypertension and metabolic syndrome, although a causal and resultant relationship has not yet been determined. Taking these observations into account, future studies should examine whether other antihypertensive drugs such as calcium channel blockers of other subclasses, anti-oxidative agents²⁴ and other drugs that may have an anti-proteinuric effect are effective in reducing lipid content in the kidneys of OLETF rats.

In this study, we also found several differences between OLETF and LETO rats in terms of the expression of lipid regulatory genes. The expression of PPAR-α and UCP-2 (mRNA) and ACC (protein) was higher, whereas that of PGC-1α (mRNA) was lower, in untreated OLETF rats than in LETO rats. Several earlier studies have shown

altered expression of these genes under conditions of diabetes or metabolic syndrome in the kidney or other organs. Proctor *et al.* reported that PPAR- α expression is decreased in the kidneys of diabetic animals compared to their non-diabetic counterparts, which results in decreased fatty acid oxidation.² In addition, the expression of UCP-2 was found to be increased in diabetic kidneys.²⁵ The mRNA expression of UCP-2 was also found to be upregulated in the liver, skeletal muscles, heart and aorta in OLETF rats compared to LETO rats.^{26,27} Downregulation of PGC-1 α mRNA expression in skeletal muscles was also reported in diabetes,²⁸ which might have a role in reducing mitochondrial function, leading to muscular lipotoxicity.²⁹

Although regulation of the expression of these genes may have played a role in the increased lipid accumulation in OLETF rat kidneys, treatment with either losartan or benidipine did not significantly alter the mRNA expression of PPAR- α , PGC-1 α or UCP-2 in OLETF rats. Compared with untreated OLETF rats, OLETF rats treated with either losartan or benidipine showed increased phosphorylation of AMPK (activated form) and ACC (inactivated form) and increased expression of CPT-1 mRNA, which may result in increased β -oxidation.¹¹ Therefore, it is possible that changes in the expression of these genes may have a role in the anti-steatotic effects of losartan and benidipine.

In summary, lipid content in the kidney was increased in untreated OLETF rats compared with LETO rats. Oil red O-stainable lipid droplets were primarily found in tubular epithelial cells, which also showed increased superoxide production. Treatment of OLETF rats with either losartan or benidipine, both of which suppressed proteinuria, reduced tissue TG content and modulated the expression of several lipid regulatory genes such as the total and phosphorylated forms of AMPK and CPT-1. These data collectively suggest that losartan and benidipine are both effective in suppressing proteinuria and normalizing lipid homeostasis in the kidneys of a rat model of metabolic syndrome. The underlying mechanisms by which these antihypertensive agents reduced lipid content in the kidney should be investigated in future studies.

ACKNOWLEDGEMENTS

This work was supported by Grants-in-Aid for Scientific Research from the Ministry of Education, Science and Culture of Japan (Grant 19590937); grants from the Takeda Science Foundation, the Sankyo Foundation of Life Science and Okinaka Memorial Institute for Medical Research; and a Grant-in-Aid from the Ministry of Health, Labour and Welfare, Japan.

- 1 Sun L, Halaihel N, Zhang W, Rogers T, Levi M. Role of sterol regulatory element-binding protein 1 in regulation of renal lipid metabolism and glomerulosclerosis in diabetes mellitus. *J Biol Chem* 2002; **277**: 18919–18927.
- 2 Proctor G, Jiang T, Iwahashi M, Wang Z, Li J, Levi M. Regulation of renal fatty acid and cholesterol metabolism, inflammation, and fibrosis in Akita and OVE26 mice with type 1 diabetes. *Diabetes* 2006; **55**: 2502–2509.
- 3 Wang Z, Jiang T, Li J, Proctor G, McManaman JL, Lucia S, Chua S, Levi M. Regulation of renal lipid metabolism, lipid accumulation, and glomerulosclerosis in FVBdb/db mice with type 2 diabetes. *Diabetes* 2005; **54**: 2328–2335.
- 4 Jiang T, Liebman SE, Lucia MS, Li J, Levi M. Role of altered renal lipid metabolism and the sterol regulatory element binding proteins in the pathogenesis of age-related renal disease. *Kidney Int* 2005; **68**: 2608–2620.
- 5 Jiang T, Wang Z, Proctor G, Moskowitz S, Liebman SE, Rogers T, Lucia MS, Li J, Levi M. Diet-induced obesity in C57BL/6J mice causes increased renal lipid accumulation and glomerulosclerosis via a sterol regulatory element-binding protein-1c-dependent pathway. *J Biol Chem* 2005; **280**: 32317–32325.

- 6 Kume S, Uzu T, Araki S, Sugimoto T, Isshiki K, Chin-Kanasaki M, Sakaguchi M, Kubota N, Terachi Y, Kadowaki T, Haneda M, Kashiwagi A, Koya D. Role of altered renal lipid metabolism in the development of renal injury induced by a high-fat diet. *J Am Soc Nephrol* 2007; **18**: 2715–2723.
- 7 Kim HJ, Moradi H, Yuan J, Norris K, Vaziri ND. Renal mass reduction results in accumulation of lipids and dysregulation of lipid regulatory proteins in the remnant kidney. *Am J Physiol Renal Physiol* 2009; **296**: F1297–F1306.
- 8 Jun H, Song Z, Chen W, Zanhua R, Yonghong S, Shuxia L, Huijun D. *In vivo* and *in vitro* effects of SREBP-1 on diabetic renal tubular lipid accumulation and RNAi-mediated gene silencing study. *Histochem Cell Biol* 2009; **131**: 327–345.
- 9 Thomas ME, Morrison AR, Schreiner GF. Metabolic effects of fatty acid-bearing albumin on a proximal tubule cell line. *Am J Physiol* 1995; **268**: F1177–F1184.
- 10 Saito K, Ishizaka N, Hara M, Matsuzaki G, Sata M, Mori I, Ohno M, Nagai R. Lipid accumulation and transforming growth factor-beta upregulation in the kidneys of rats administered angiotensin II. *Hypertension* 2005; **46**: 1180–1185.
- 11 Rector RS, Thyfault JP, Morris RT, Laye MJ, Boregasser SJ, Booth FW, Ibdah JA. Daily exercise increases hepatic fatty acid oxidation and prevents steatosis in Otsuka Long-Evans Tokushima Fatty rats. *Am J Physiol Gastrointest Liver Physiol* 2008; **294**: G619–G626.
- 12 Man ZW, Zhu M, Noma Y, Toide K, Sato T, Asahi Y, Hirashima T, Mori S, Kawano K, Mizuno A, Sano T, Shima K. Impaired beta-cell function and deposition of fat droplets in the pancreas as a consequence of hypertriglyceridemia in OLETF rat, a model of spontaneous NIDDM. *Diabetes* 1997; **46**: 1718–1724.
- 13 Kosegawa I, Katayama S, Kikuchi C, Kashiwabara H, Negishi K, Ishii J, Inukai K, Oka Y. Metformin decreases blood pressure and obesity in OLETF rats via improvement of insulin resistance. *Hypertens Res* 1996; **19**: 37–41.
- 14 Okumura M, Masada M, Yoshida Y, Shintaku H, Hosoi M, Okada N, Konishi Y, Morikawa T, Miura K, Imanishi M. Decrease in tetrahydrobiopterin as a possible cause of nephropathy in type II diabetic rats. *Kidney Int* 2006; **70**: 471–476.
- 15 Aizawa T, Ishizaka N, Taguchi J, Nagai R, Mori I, Tang SS, Ingelfinger JR, Ohno M. Heme oxygenase-1 is upregulated in the kidney of angiotensin II-induced hypertensive rats: possible role in renoprotection. *Hypertension* 2000; **35**: 800–806.
- 16 Hongo M, Ishizaka N, Furuta K, Yahagi N, Saito K, Sakurai R, Matsuzaki G, Koike K, Nagai R. Administration of angiotensin II, but not catecholamines, induces accumulation of lipids in the rat heart. *Eur J Pharmacol* 2009; **604**: 87–92.
- 17 Matsuzaki G, Ishizaka N, Furuta K, Hongo M, Saito K, Sakurai R, Koike K, Nagai R. Comparison of vasculoprotective effects of benidipine and losartan in a rat model of metabolic syndrome. *Eur J Pharmacol* 2008; **587**: 237–242.
- 18 Machado MO, Hirata RD, Sellitti DF, Iotti R, Iotti A, Cusumano AM, Rioridan GP, Coschignano KT, Kopchick JJ, Zuhl I, Nguyen N, Hirata MH, Doi SQ. Growth hormone promotes glomerular lipid accumulation in bGH mice. *Kidney Int* 2005; **68**: 2019–2028.
- 19 Ishizaka N, Matsuzaki G, Saito K, Noiri E, Mori I, Nagai R. Expression and localization of PDGF-B, PDGF-D, and PDGF receptor in the kidney of angiotensin II-infused rat. *Lab Invest* 2006; **86**: 1285–1292.
- 20 Toblli JE, Cao G, Rivas C, DeRosa G, Domecq P. Angiotensin-converting enzyme inhibition reduces lipid deposits in myocardium and improves left ventricular function of obese Zucker rats. *Obesity (Silver Spring)* 2006; **14**: 1586–1595.
- 21 Toblli JE, Munoz MC, Cao G, Mella J, Pereyra L, Mastai R. ACE inhibition and AT1 receptor blockade prevent fatty liver and fibrosis in obese Zucker rats. *Obesity (Silver Spring)* 2008; **16**: 770–776.
- 22 Ohishi M, Takagi T, Ito N, Terai M, Tataru Y, Hayashi N, Shiota A, Katsuya T, Rakugi H, Oghihara T. Renal-protective effect of T-and L-type calcium channel blockers in hypertensive patients: an Amlodipine-to-Benidipine Changeover (ABC) study. *Hypertens Res* 2007; **30**: 797–806.
- 23 Ohtsubo T, Matsumura K, Sakagami K, Fujii K, Tsuruya K, Noguchi H, Rovira II, Finkel T, Iida M. Xanthine oxidoreductase depletion induces renal interstitial fibrosis through aberrant lipid and purine accumulation in renal tubules. *Hypertension* 2009; **54**: 868–876.
- 24 Chung S, Park CW, Shin SJ, Lim JH, Chung HW, Youn DY, Kim HW, Kim BS, Lee JH, Kim GH, Chang YS. Tempol or candesartan prevents high-fat diet-induced hypertension and renal damage in spontaneously hypertensive rats. *Nephrol Dial Transplant* 2009 (e-pub ahead of print 11 September 2009; doi:10.1093/ndt/gfp472).
- 25 Friederich M, Nordquist L, Olerud J, Johansson M, Hansell P, Palm F. Identification and distribution of uncoupling protein isoforms in the normal and diabetic rat kidney. *Adv Exp Med Biol* 2009; **645**: 205–212.
- 26 Mori Y, Tokutate Y, Oana F, Matsuzawa A, Akahane S, Tajima N. Bezafibrate-induced changes over time in the expression of uncoupling protein (UCP) mRNA in the tissues: a study in spontaneously type 2 diabetic rats with visceral obesity. *J Atheroscler Thromb* 2004; **11**: 224–231.
- 27 Minamiyama Y, Bito Y, Takemura S, Takahashi Y, Kodai S, Mizuguchi S, Nishikawa Y, Suehiro S, Okada S. Calorie restriction improves cardiovascular risk factors via reduction of mitochondrial reactive oxygen species in type II diabetic rats. *J Pharmacol Exp Ther* 2007; **320**: 535–543.
- 28 Mensink M, Hesselink MK, Russell AP, Schaart G, Sels JP, Schrauwen P. Improved skeletal muscle oxidative enzyme activity and restoration of PGC-1 alpha and PPAR beta/delta gene expression upon rosiglitazone treatment in obese patients with type 2 diabetes mellitus. *Int J Obes (Lond)* 2007; **31**: 1302–1310.
- 29 Schrauwen P. High-fat diet, muscular lipotoxicity and insulin resistance. *Proc Nutr Soc* 2007; **66**: 33–41.

It's back-to-back for your
ankylosing spondylitis patients.



Abbott
A Promise for Life

The Journal of Rheumatology

The Journal of Rheumatology

Volume 37, no. 2

Changes in Waist Circumference and Body Mass Index in Relation to Changes in Serum Uric Acid in Japanese Individuals

NOBUKAZU ISHIZAKA, YUKO ISHIZAKA, AKIKO TODA, MIZUKI TANI, KAZUHIKO KOIKE, MINORU YAMAKADO and RYOZO NAGAI

J Rheumatol 2010;37:410-416
<http://www.jrheum.org/content/37/2/410>

1. Sign up for our monthly e-table of contents

<http://www.jrheum.org/cgi/alerts/etoc>

2. Information on Subscriptions

<http://jrheum.com/subscribe.html>

3. Have us contact your library about access options

Refer_your_library@jrheum.com

4. Information on permissions/orders of reprints

<http://jrheum.com/reprints.html>

The Journal of Rheumatology is a monthly international serial edited by Duncan A. Gordon featuring research articles on clinical subjects from scientists working in rheumatology and related fields.

Changes in Waist Circumference and Body Mass Index in Relation to Changes in Serum Uric Acid in Japanese Individuals

NOBUKAZU ISHIZAKA, YUKO ISHIZAKA, AKIKO TODA, MIZUKI TANI, KAZUHIKO KOIKE, MINORU YAMAKADO, and RYOZO NAGAI

ABSTRACT. *Objective.* Studies have shown that obesity is associated with an increase in serum uric acid; and few data are available on the relationship between changes in measures of obesity and changes in uric acid concentrations. We investigated the relationship among percentage changes in waist circumference (%dWC), body mass index (%dBMI), and serum uric acid (%dUA).

Methods. The data of 3153 individuals [1968 men, 1185 women (536 premenopausal, 649 postmenopausal)] who underwent general health screening over a 2-year period and were not taking anti-hyperuricemic medication were analyzed.

Results. Stepwise multiple regression analysis showed that %dBMI was associated positively with %dUA in postmenopausal women and men, and the association retained statistical significance after adjustment for changes in blood pressure and in renal function. Association between %dBMI and %dUA was not significant in premenopausal women. In men, %dWC was a predicting factor for %dUA, although it did not remain significant when %dBMI was used as a covariate in the statistical model. Multivariate logistic regression analysis showed that the odds ratio of the association between the lowest %dBMI quartile (%dBMI < -1.86) and the lowest %dUA quartile (%dUA < -7.41) was 2.04 (95% CI 1.35–3.07) in postmenopausal women and 1.46 (95% CI 1.14–1.86) in men.

Conclusion. Weight loss may represent an effective nonmedical strategy for reducing serum UA levels, especially in postmenopausal women and men. (First Release Dec 23 2009; J Rheumatol 2010;37:410–6; doi:10.3899/jrheum.090736)

Key Indexing Terms:

WAIST CIRCUMFERENCE
GLOMERULAR FILTRATION RATE

BODY MASS INDEX

URIC ACID
BLOOD PRESSURE

Obesity and serum uric acid (UA) are both associated with enhanced insulin resistance and incidence of metabolic syndrome¹⁻³. In addition, measures of obesity have been reported to be positively associated with serum levels of UA^{4,5}, an association that may be caused by impaired renal clearance

of UA in the condition of obesity⁶. The finding that a reduction in weight, and thus in body mass index (BMI), may have a significant effect on serum UA⁷ and renal urate excretion⁶ suggests that changes in weight may play a role in the regulation of serum UA levels, although the reverse scenario might also be possible⁸. In our study, by analyzing individuals who underwent general health screening, we examined the influence of changes in waist circumference (WC) and BMI on changes in UA, and the dependency/independency on changes in either blood pressure or renal function, which is the possible critical factor affecting serum UA levels in healthy subjects⁹.

MATERIALS AND METHODS

Study population. The study was approved by The Ethical Committee of Mitsui Memorial Hospital. Between October 2005 and October 2006, 11,558 individuals underwent general health screening at our institute. Of these, 3326 individuals (2113 men, 1213 women) underwent general health screening during this period (first visit) and again the following year (second visit). Among these 3326 individuals, 3179 (1968 men, 1211 women) reported taking no antihyperuricemic drugs at either visit. Among the 1211 women, 1185 (98%) answered the questionnaire concerning whether they still had menstruation, and were enrolled for study. Therefore, we analyzed data of 3153 individuals (1968 men, 1185 women). The mean \pm standard deviation (SD) interval between the 2 visits of the individuals enrolled was

From the Department of Cardiovascular Medicine and Department of Gastroenterology, University of Tokyo Graduate School of Medicine, Tokyo; and Center for Multiphasic Health Testing and Services, Mitsui Memorial Hospital, Tokyo, Japan.

Supported in part by a grant from Chiyoda Mutual Life Foundation, and by the St. Luke's Grant for the Epidemiological Research, Daiwa Securities Health Foundation, Gerontology Research Grant of Kowa Life Science Foundation, Foundation for Total Health Promotion, Gout Research Foundation of Japan, Kurozumi Medical Foundation, and a Grant-in-Aid from the Ministry of Health, Labour, and Welfare, Japan.

N. Ishizaka, MD, Department of Cardiovascular Medicine, University of Tokyo Graduate School of Medicine; Y. Ishizaka, MD; A. Toda, MD; M. Tani, MD, Center for Multiphasic Health Testing and Services, Mitsui Memorial Hospital; K. Koike, MD, Department of Gastroenterology, University of Tokyo Graduate School of Medicine; M. Yamakado, MD, Center for Multiphasic Health Testing and Services, Mitsui Memorial Hospital; R. Nagai, MD, Department of Cardiovascular Medicine, University of Tokyo Graduate School of Medicine.

Address correspondence to Dr. N. Ishizaka, Department of Cardiovascular Medicine, University of Tokyo Graduate School of Medicine, Hongo 7-3-1 Bunkyo-ku, Tokyo 113-8655, Japan. E-mail: nobuishizaka-iky@umin.ac.jp

Accepted for publication September 15, 2009.

Personal non-commercial use only. The Journal of Rheumatology Copyright © 2010. All rights reserved.

356.2 ± 51.7 days. WC, BMI, UA, systolic blood pressure, and estimated glomerular filtration rate (eGFR) at the first visit were designated WC1, BMI1, UA1, BPs1, and eGFR1, respectively, and at the second visit WC2, BMI2, UA2, BPs2, and eGFR2.

The percentage differences between values of WC1 and WC2, BMI1 and BMI2, UA1 and UA2, BPs1 and BPs2, and eGFR1 and eGFR2 were designated %dWC, %dBMI, %dUA, %dBPs, and %deGFR. All participants were seen after an overnight fast. Height and weight were determined, and BMI was expressed as weight (kilograms) divided by the square of height (meters). With the subject standing, waist circumference was measured at the umbilical level to the nearest 1 cm by trained physicians and technicians¹⁰.

Laboratory analysis. Blood samples were taken after an overnight fast. Serum levels of low density lipoprotein (LDL), high density lipoprotein (HDL) cholesterol, and triglycerides were determined enzymatically. Serum UA was measured by the uricase-peroxidase method, and hemoglobin A_{1c} by latex agglutination immunoassay. Creatinine was measured by TBA-200FR (Toshiba Medical Systems, Tochigi, Japan) using a commercial kit. Accuras Auto CRE (Shino-test, Tokyo, Japan) eGFR was calculated by the equation: $eGFR = 194 \times (\text{serum creatinine})^{-1.094} \times (\text{age})^{-0.287} \times 0.739$ if female¹¹. This equation was recently determined by a multicenter study, and differs from the equation¹² that we used in previous studies¹³⁻¹⁵. Blood pressure was measured after about 10 min of rest with an automated sphygmomanometer.

Statistical analysis. Data are expressed as the mean ± SD unless stated otherwise. Analyses of variance with trend analysis and stepwise multiple regression analysis were conducted as appropriate to assess the statistical significance of differences between groups using SPSS II (SPSS Inc., Chicago, IL, USA). A value of $p < 0.05$ was taken to be statistically significant.

RESULTS

Baseline characteristics. We enrolled 536 premenopausal women, 649 postmenopausal women, and 1968 men for study. At the first visit the mean age of premenopausal women was 43.1 ± 5.51 years, postmenopausal women 59.1 ± 6.8 years, and men 53.3 ± 10.2 years. The sex-nonspecific range of the first to the fourth %dUA quartiles (maximum/minimum) was -47.2/-7.5, -7.4/-1.2, 0.0/7.1, and 7.2/77.8 (Table 1). A plot of WC1 and BMI1 compared to UA1 is shown in Figure 1. In both men and women, there was a statistically significant correlation between WC1 or BMI1 and UA1. The correlation coefficient between WC1 and BMI1 was 0.626 ($p < 0.001$) in premenopausal women, 0.563 ($p < 0.001$) in postmenopausal women, and 0.838 ($p < 0.001$) in men.

Relationship between %dWC, %dBMI, and %dUA. The relationship between %dBMI and %dUA, although very weak, was significant in postmenopausal women and men, and the relationship between %dWC and %dUA was significant only in men (Figure 2). In premenopausal women, the relationships between %dWC and %dUA and between %dBMI and %dUA were not statistically significant. The correlation coefficient between %dWC and %dBMI was 0.267 ($p < 0.001$) in premenopausal women, 0.221 ($p < 0.001$) in postmenopausal women, and 0.484 ($p < 0.001$) in men.

We next performed stepwise multiple regression analysis

(Table 2). In a model in which age, UA1, WC1, and %dWC were used as independent variables (model 1), %dWC was found to have independent predictive value for %dUA in men, but not in women. However, after adding BMI1 and %dBMI as independent variables, %dWC in men was no longer a predictor for %dUA (model 2). %dBMI was found to be a predictor for %dUA in postmenopausal women and men even after using either or both %dBPs and %deGFR as independent variables (models 3-5). On the other hand, in premenopausal women, %dBMI was not a significant predictor value for %dUA in any of these models. In addition, in a model in which age, UA1, BMI1, and %dBMI were used as independent variables, %dBMI again was not found to have significant predictive value for %dUA (data not shown). In model 5, variance inflation factor (VIF) scores of all the independent variables were less than 10 (data not shown).

Logistic regression analysis. When the highest %dUA quartile (%dUA ≥ 7.25) was used as a dependent variable, logistic regression analysis showed that the highest %dBMI quartile (%dBMI ≥ 1.47) had a significant positive association in postmenopausal women and in men after adjusting for UA1 and BMI1 (Table 3, model 1). In these groups, statistical significance was retained even after further adjustment for %dBPs and %deGFR (model 2). On the other hand, in premenopausal women, the highest %dBMI quartile was, unexpectedly, negatively associated with the highest %dUA quartile, although statistical significance was lost after further adjustment for %dBPs and %deGFR. In all 3 subgroups tested, %deGFR was negatively associated with the highest quartile of %dUA.

Logistic regression analysis showed that the lowest %dBMI quartile (%dBMI < -1.86) had a significant positive association with the lowest %dUA quartile (%dUA < -7.41) in postmenopausal women and men, and this remained statistically significant even after further adjustment for %dBPs and %deGFR (model 2). But in premenopausal women, association between the lowest %dBMI quartile and lowest %UA was not statistically significant regardless of this further adjustment.

DISCUSSION

Analyzing data of individuals who underwent general health screening and who were taking no antihyperuricemic medication, we found that correlation between percentage changes in BMI (%dBMI) and in UA (%dUA) was statistically significant in postmenopausal women and in men, but not in premenopausal women.

Stepwise multiple regression analysis showed that %dWC is a significant independent variable for %dUA in men, where UA1, WC1, and %dWC was used as possible independent variables (model 1); however, the relationship lost statistical significance after further adjustment for BMI1 and %dBMI (Table 3). %dWC was not found to be a

Table 1. Baseline characteristics at the first visit according to %dUA quartiles.

Variables	Total, n	First (range -47.2/-7.5)	%dUA Quartile			p for Trend
			Second (range -7.4/-1.2)	Third (range 0.0/7.1)	Fourth (range 7.2/77.8)	
Women/men	1185/1968	290/496	252/483	327/514	316/475	
Baseline data at visit 1						
Age, yrs	52.8 ± 10.2	52.6 ± 10.6	53.3 ± 10.1	52.9 ± 10.0	52.3 ± 10.2	0.225
Waist circumference, cm	82.3 ± 9.2	82.8 ± 9.6	82.7 ± 9.1	82.2 ± 8.7	81.7 ± 9.3	0.074
Body mass index, kg/m ²	22.8 ± 3.1	23.0 ± 3.3	22.8 ± 3.0	22.8 ± 3.0	22.6 ± 3.1	0.056
Systolic blood pressure, mmHg	123 ± 19	124 ± 20	123 ± 19	122 ± 19	123 ± 19	0.636
LDL-cholesterol, mg/dl	129.1 ± 31.2	130.2 ± 31.3	129.4 ± 29.5	129.8 ± 31.3	126.9 ± 32.2	0.147
HDL-cholesterol, mg/dl	60.7 ± 15.3	60.7 ± 15.8	60.5 ± 15.5	60.8 ± 14.8	60.9 ± 15.1	0.980
Triglyceride, mg/dl	111 ± 72	112 ± 75	115 ± 81	109 ± 67	109 ± 65	0.249
Uric acid, mg/dl	5.5 ± 1.3	5.8 ± 1.5	5.7 ± 1.2	5.4 ± 1.3	5.1 ± 1.2	< 0.001
Fasting glucose, mg/dl	96 ± 21	97 ± 20	96 ± 19	95 ± 18	96 ± 25	0.486
Hemoglobin A1c, %	5.3 ± 0.7	5.3 ± 0.7	5.3 ± 0.7	5.3 ± 0.7	5.3 ± 0.9	0.941
Blood urea nitrogen, mg/dl	14.1 ± 3.5	14.6 ± 4.0	14.3 ± 3.2	14.1 ± 3.3	13.6 ± 3.5	< 0.001
Serum creatinine, mg/dl	0.77 ± 0.29	0.79 ± 0.42	0.78 ± 0.15	0.76 ± 0.15	0.75 ± 0.34	0.068
Estimated glomerular filtration rate, ml/min/1.73m ²	68.3 ± 11.8	67.3 ± 11.8	67.8 ± 11.5	68.1 ± 11.4	70.2 ± 12.4	< 0.001
Antihypertensive medication, n (%)	306 (9.7)	78 (9.9)	78 (10.6)	76 (9.0)	74 (9.4)	0.736
Antidiabetic medication, n (%)	74 (2.3)	19 (2.4)	14 (1.9)	24 (2.9)	17 (2.1)	0.632
Postmenopause (female), n (%)	649 (54.8)	152 (52.4)	141 (56.0)	194 (59.3)	162 (51.3)	0.165
Current smoker, n (%)	734 (23.3)	180 (22.9)	173 (23.5)	197 (23.4)	184 (23.3)	0.992
Percent change between 2 visits						
%dBMI	-0.28 ± 3.1	-0.70 ± 3.41	-0.46 ± 3.06	0.01 ± 2.88	-0.03 ± 2.98	< 0.001
%dWC	0.17 ± 6.12	-0.02 ± 6.33	-0.35 ± 6.14	0.64 ± 6.14	0.35 ± 5.83	0.008
%deGFR	1.8 ± 10.0	6.3 ± 10.0	2.6 ± 9.0	0.9 ± 9.1	-2.5 ± 10.0	< 0.001

predictive value for %dUA in premenopausal or postmenopausal women. By contrast, %dBMI was found to be a predictor for %dUA, even after adjustment for WC1, %dWC, %dBPs, and %deGFR in postmenopausal women and men, although it was not significant in premenopausal women. In premenopausal women, %dBMI was not a significant predictive value for %dUA in the model in which age, UA1, BMI1, and %dBMI were used as independent variables; therefore, failure of %dBMI as a predictor for %dUA in premenopausal women may not fully be explained by the multicollinearity between %dWC and %dBMI. These findings collectively indicate that mode of association between change in BMI and change in UA differs between premenopausal and postmenopausal women.

There are several previous studies in which changes in obesity measures have been analyzed in relation to the changes in UA over a certain period of time. For example, Heyden, *et al* showed that there was a stepwise progression from decreased UA levels associated with maximum weight loss to increased levels with maximum weight gain¹⁶. In addition, Rathmann, *et al*⁷ analyzed the data of 1249 male and 1362 female subjects aged 17–35 years from the Coronary Artery Risk Development in Young Adults (CARDIA) Study who attended a 10-year followup. They reported that changes in BMI and WC were associated with changes in UA in a statistical model adjusted for age and baseline UA levels⁷. In contrast, we found that %dBMI was,

but %dWC was not, significantly associated with %dUA in multiple linear regression (Table 2, models 2-5). This might be because statistical significance had been weakened after %dWC and %dBMI were simultaneously included into the statistical model; however, %dWC was not significantly associated with %dUA in women even before the adjustment for %dBMI (model 1). From our epidemiological study, we cannot determine what would have caused the different observations between the findings of Rathmann, *et al*⁷ and our own. However, considering that circulating insulin levels may have potential to regulate serum UA levels¹⁷, the difference might derive from the difference in insulin sensitivity^{18,19} and/or difference in the effect of obesity on insulin resistance²⁰ among various ethnicities. This possibility should be investigated in future studies. Choe, *et al* found that mean changes in BMI, but not in WC, were statistically different between subjects who had decreased or had no change in UA and those with increased UA during a 1-year followup in men who underwent health promotion screening⁹. They also found that changes in serum creatinine levels, but not in systolic or diastolic blood pressure, were significantly different between subjects who had decreased or unchanged UA levels and those with increased UA⁹ — a finding that is, in one sense, in agreement with our observations.

What would be the possible underlying mechanisms that explain the difference in the mode of association between




Figure 1. Scatterplot and linear regression between waist circumference at the first visit (WC1) and uric acid at the first visit (UA1) and those between body mass index at the first visit (BMI1) and UA1 in premenopausal and postmenopausal women and in men. Serum uric acid values were not adjusted for age or other possible confounders.

%dBMI and %dUA between premenopausal and postmenopausal women? It has been reported that certain alterations in UA metabolism may occur after the menopause; menopause leads to an increase in serum UA levels^{21,22}, and this may be in part attributed to decreased estrogen production and subsequent reduction of the fractional excretion of UA²³. In addition, a recent study suggested that association between insulin resistance and serum UA levels may be greater in postmenopausal women than premenopausal women²⁴. Whether these phenomena are related to the difference in the mode of association between %dBMI and %dUA of premenopausal and postmenopausal women remains to be investigated.

We previously showed that obesity or overweight was significantly associated with chronic kidney disease¹⁴, and that changes in obesity measures may be associated with changes in eGFR and urinary excretion of albumin²⁵. The strength of the current study was that we demonstrate that change in BMI was positively associated with change in UA in postmenopausal women and men independent of change in eGFR. In addition, we show that mode of association between %dBMI and %dUA was different between pre-

menopausal and postmenopausal women, which may have relation with the fact that menopause causes the elevation of serum UA^{21,22,26}. However, controlling BMI is neither unnecessary nor ineffective in keeping the metabolic measures in optimal ranges in "premenopausal" women, because weight gain may result in the reduced insulin sensitivity and aggravation of cardiovascular risk also in premenopausal women²⁷.

Data for visceral fat volume measured by computed tomography were not available in our study. Recent reports showed that subcutaneous fat accumulation is related to impaired urinary UA excretion⁶, whereas visceral fat accumulation is linked closely to the overproduction of uric acid²⁸, and that serum UA levels are increased both in individuals with subcutaneous fat obesity and in those with visceral fat obesity²⁸. It remains to be determined whether changes of WC will lead to an increase in urinary UA excretion in our population, and whether there is a relationship between %dUA and change in visceral fat volume.

Our study has several potential limitations. First, we had no information on the extent to which modifications of lifestyle and dietary habits affected observed changes in