

Fig. 1. *Continued*

monitored clinically and biochemically after DC infusion (Table 2). A larger proportion (12 of 13) of the patients were complicated with high fever compared to those treated previously with immature DCs (five of 10) [20], due probably to the proinflammatory responses induced by OK432-stimulated DCs. However, there were no grades III or IV

National Cancer Institute Common Toxicity Criteria adverse events, including vomiting, abdominal pain, encephalopathy, myalgia, ascites, gastrointestinal disorders, bleeding, hepatic abscess or autoimmune diseases associated with DC infusion and TAE in this study. There was also no clinical or serological evidence of hepatic failure or autoimmune

Table 2. Adverse events.

Patient no.	Fever (days)	Vomiting	Abdominal pain	Encephalopathy	Others [*]
1	2	No	No	No	No
2	2	No	No	No	No
3	1	No	No	No	No
4	3	No	No	No	No
5	3	No	No	No	No
6	4	No	No	No	No
7	10	No	No	No	No
8	No	No	No	No	No
9	2	No	No	No	No
10	1	No	No	No	No
11	2	No	No	No	No
12	2	No	No	No	No
13	1	No	No	No	No

^{*}Other adverse events include myalgia, ascites, gastrointestinal disorder, bleeding, hepatic abscess and autoimmune diseases.

response in any patients. Thus, concurrent treatment with OK432-stimulated DC infusions can be performed safely at the same time as TAE in patients with cirrhosis and HCC.

Recurrence-free survival following DC infusion

A further objective of this study was to determine clinical response following DC infusion. A group of historical controls treated with TAE without DC administration was reviewed for this study (Table 3). The clinical characteristics including tumour burden and hepatic reserve were comparable between patients treated with TAE and OK432-stimulated DC transfer ($n = 13$) and those historical controls with TAE but without DC administration ($n = 22$). We com-

pared the recurrence-free survival between these patient groups. Kaplan–Meier analysis indicated that patients treated with TAE and OK432-stimulated DC transfer had prolonged recurrence-free survival compared with the historical controls that had been treated with TAE alone (recurrence rates 360 days after the treatments; two of 13 and 12 of 22, respectively; $P = 0.046$, log-rank test) (Fig. 2). The results demonstrated that OK432-stimulated DC transfer during TAE therapy reduces tumour recurrence in HCC patients.

NK cell activity and intracellular cytokine responses in PBMCs

To assess systemic immunomodulatory effects of OK432-stimulated DC transfer, PBMCs were isolated 1 and 3 months after treatment and NK cell cytotoxicity against K562 erythroleukaemia target cells measured using the ⁵¹Cr-release assay (Fig. 3). The level of NK cell was unaltered following treatment. In addition, cytokine production capacity of lymphocyte subsets was quantitated by measuring intracellular IFN- γ and IL-4 using flow cytometry. There were also no significant changes in terms of cytokine production capacity in the CD4⁺, CD8⁺ and CD56⁺ subsets in the patients treated with OK432-stimulated DCs.

Immune responses to peptide epitopes derived from tumour antigens

To assess the effects on T cell responses to tumour antigens, PBMCs were obtained 4 weeks after DC infusion, pulsed with peptides derived from AFP, MRP3, SART2, SART3 and hTERT. IFN- γ production was then quantitated in an

Table 3. Clinical characteristics of patients treated with TAE + OK-DC and TAE alone.

	TAE + OK-DC	TAE	P
No. of patients	13	22	
Age (years)	68.2 \pm 9.1	70.0 \pm 7.6	n.s. [†]
Gender (M/F)	9/4	13/9	n.s. [‡]
White cell count ($\times 10^3/\mu\text{l}$)	34.4 \pm 11.6	41.4 \pm 18.9	n.s. [†]
Lymphocytes ($\times 10^3/\mu\text{l}$)	10.4 \pm 3.6	12.4 \pm 4.7	n.s. [†]
Platelets ($\times 10^3/\mu\text{l}$)	11.5 \pm 10.2	10.3 \pm 5.8	n.s. [†]
Hepaplastin test (%)	64.6 \pm 11.6	75.5 \pm 24.3	n.s. [†]
ALT (IU/l)	56.7 \pm 38.9	67.9 \pm 44.6	n.s. [†]
Total bilirubin (mg/dl)	1.3 \pm 0.7	1.1 \pm 0.6	n.s. [†]
Albumin (g/dl)	3.4 \pm 0.6	3.6 \pm 0.4	n.s. [†]
Non-cancerous liver parenchyma (no.)			
Chronic hepatitis	0	8	
Cirrhosis (Child–Pugh A/B/C)	13 (5/8/0)	14 (6/8/0)	n.s. [‡]
TNM stages (I/II/III/IV-A/IV-B)	0/4/9/0/0	3/8/11/0/0	n.s. [‡]
No. of tumours	2.5 \pm 1.3	1.9 \pm 1.3	n.s. [†]
Largest tumour (mm)	30.2 \pm 9.4	32.6 \pm 15.2	n.s. [†]
AFP	204.8 \pm 404.1	201.8 \pm 544.2	n.s. [†]

Results are expressed as means \pm standard deviation. [†]Mann–Whitney *U*-test. [‡]Fisher's exact test. TAE, transcatheter arterial embolization; OK-DC, OK432-stimulated dendritic cells; ALT, alanine transaminase; TNM, tumour–node–metastasis; AFP, alpha-fetoprotein; Child–Pugh, Child–Pugh classification; n.s., not significant.




Fig. 2. Recurrence-free survival of patients treated with transcatheter hepatic arterial embolization (TAE) with [TAE + OK-stimulated dendritic cells (DC); $n = 13$] and without (TAE: historical controls; $n = 22$) OK432-stimulated DC administration. Time zero is the date of TAE. All patients underwent ultrasound, computed tomography (CT) scan or magnetic resonance imaging (MRI) of the abdomen about 1 month after treatment and at a minimum of once every 3 months thereafter. Kaplan–Meier analysis indicated that TAE + OK-DC treatment prolonged recurrence-free survival compared with the TAE-alone group (recurrence rates 360 days after the treatments; two of 13 and 12 of 22, respectively; $P = 0.046$, log-rank test).

ELISPOT assay. Cells producing IFN- γ in response to stimulation with HLA-A24 [the most common HLA-A antigen (58.1%) in Japanese populations [35]]-restricted peptide epitopes derived from tumour antigens MRP3 and hTERT were induced in three of six HLA-A24-positive patients (numbers 2, 6 and 11) after treatment with TAE and OK432-stimulated DCs (Fig. 4). To understand the immunological and clinical significance of the T lymphocyte responses, PBMCs obtained from the historical control patients who had been treated with TAE without DC administration were also evaluated by ELISPOT. Similarly, positive reactions were observed in four (numbers t8, t19, t20 and t22) of six HLA-A24-positive patients. These data indicate that T lymphocyte responses to HLA-A24 restricted peptide epitopes of tumour antigens were induced following the TAE therapy, but no additional responses were observed as a result of OK432-stimulated DC transfer in the current study.

Serum levels of cytokines, chemokines and arginase activity

To screen for immunobiological responses induced following OK432-stimulated DC transfer, serum levels of cytokines and chemokines were measured simultaneously using the Bio-Plex multiplex suspension array system. The results were compared with the historical control patients treated with TAE without DC administration. Interestingly, serum con-





Fig. 3. Natural killer (NK) cell activity and intracellular cytokine production in peripheral blood mononuclear cells (PBMCs) of patients treated with OK432-stimulated dendritic cells (DCs) during transcatheter hepatic arterial embolization (TAE) therapy ($n = 13$). PBMCs were isolated before and 1 and 3 months after treatment and used for the analyses. Upper panel: NK cell cytotoxicity against K562 erythroleukaemia target cells was evaluated at the effector/target (E/T) cell ratios shown. NK cell activities were not changed following treatment. Middle and lower panels: PBMCs were stimulated with phorbol 12-myristate 13-acetate (PMA) and ionomycin, stained for CD4, CD8 and CD56 expression, permeabilized and stained for intracellular interferon (IFN)- γ and interleukin (IL)-4. Percentages of cytokine-positive cells were quantitated by flow cytometry. There were no significant changes in terms of cytokine production capacity in the CD4+, CD8+ and CD56+ subsets following the treatments. The data are given as means \pm standard deviation of the groups.

Fig. 4. Immune responses to human leucocyte antigen (HLA-DR)-A24-restricted peptide epitopes derived from tumour antigens in HLA-A24-positive patients treated with OK432-stimulated DCs during transcatheter hepatic arterial embolization (TAE) therapy (numbers 2, 5, 6, 7, 10 and 11) and HLA-A24-positive historical controls treated with TAE without dendritic cell (DC) transfer (numbers t8, t14, t15, t19, t20 and t22). Peripheral blood mononuclear cells (PBMCs) were obtained before (open bars) and 1 month after the infusion (solid bars), pulsed with the peptides derived from squamous cell carcinoma antigen recognized by T cells 2 (SART2), SART3, multi-drug resistance protein 3 (MRP3), alpha-fetoprotein (AFP), human telomerase reverse transcriptase (hTERT) and interferon (IFN)- γ production was quantitated by enzyme-linked immunospot (ELISPOT). Negative controls consisted of a human immunodeficiency virus (HIV) envelope-derived peptide (HIVenv₅₈₄). Positive controls consisted of 10 ng/ml phorbol 12-myristate 13-acetate (PMA) or a cytomegalovirus (CMV) pp65-derived peptide (CMVpp65₃₂₈). The number of specific spots was determined by subtracting the number of spots in the absence of antigen from the number of spots in its presence. T lymphocyte responses to the peptide epitopes were induced following TAE therapy, but no additional responses were observed after DC transfer. Numbers denote specific spots beyond the upper limit of y-axis; n.d., not determined.



centrations of IL-9, IL-15 and TNF- α were greatly increased after OK432-stimulated DC infusion, in contrast to their reduction following TAE treatment alone (Fig. 5a). Furthermore, the chemokines eotaxin (CCL11) and MIP-1 β (CCL4) were induced markedly after DC transfer, although they were also decreased after TAE alone. These data indicate that transfer of OK432-stimulated DC during TAE therapy induced unique immune responses that may be mediated by the cytokines IL-9, IL-15 and TNF- α and the chemokines eotaxin and MIP-1 β .

In addition, serum arginase activity was reported to reflect numbers of myeloid-derived suppressor cells (MDSCs) that may inhibit T lymphocyte responses in cancer patients [36]. Therefore, serum arginase activity was measured after OK432-stimulated DC infusion, and it was found that it was

increased six- or sevenfold in patients treated with TAE. However, this increase was independent of the presence or absence of OK432-stimulated DC transfer (Fig. 5b). None the less, serum arginase activity was decreased again 4 weeks after treatment with both TAE and OK432-stimulated DC transfer but tended to be maintained at a high levels in patients treated with TAE without DC transfer. However, these differences did not reach statistical significance ($P > 0.05$). Because arginase activity is known to be relatively high in liver and HCC cells [37], the influence of tissue injury was assessed biochemically by measuring serum levels of ALT and LDH activities. We did not observe ALT or LDH elevation, indicating that the increase of arginase activity was not due to tissue damage following treatment. Collectively, these results demonstrate that infusion of OK432-stimulated




Fig. 5. Cytokine and chemokine profiling and arginase activity in sera of patients treated with OK432-stimulated dendritic cells (DCs) during transcatheter hepatic arterial embolization (TAE) therapy (TAE + OK-DC; $n = 13$) and the historical controls treated with TAE without DC transfer (TAE; $n = 22$). (a) Serum samples were examined for their content of a validated panel of cytokines and chemokines using the Bioplex assay. Percentage changes in serum levels 2 weeks after the treatments were calculated as follows: $[(\text{post-treatment level} - \text{pretreatment level}) / \text{pretreatment level}] \times 100$. The data are means \pm standard error of the mean (s.e.m.) of the groups. $*P < 0.05$ when compared by the Mann-Whitney U -test. (b) Serum samples were tested for arginase activity by conversion of L-arginine to L-ornithine, and for alanine aminotransferase (ALT) and lactic dehydrogenase (LDH) activities. While there was a trend for the arginase activity in the TAE + OK-DC group to decrease 4 weeks after treatment, the difference did not reach statistical significance ($P > 0.05$). Percentage changes in serum levels 2 weeks after the treatments were calculated as follows: $[(\text{post-treatment level} - \text{pretreatment level}) / \text{pretreatment level}] \times 100$. The data indicate means \pm s.e.m. of the groups.

DCs during TAE treatment may reduce the immunosuppressive activities of MDSCs, and assist in developing a favourable environment for the induction of anti-tumour immunity.

Discussion

Although many novel strategies, including immunotherapies, have been developed in an attempt to suppress tumour recurrence after curative treatments for HCC, recurrence rates and survival times have not been improved significantly [38]. In the current study, we first established that OK432-stimulated DC administration during TAE therapy did not cause critical adverse events in patients with cirrhosis and HCC. Most importantly, DC transfer resulted in prolonged recurrence-free survival after combination therapy with TAE and OK432-stimulated DC administration. In terms of the immunomodulatory effects of DC transfer, although NK cell activity, intracellular cytokine production and T lymphocyte-mediated immune responses were not altered in PBMCs from treated patients, serum levels of IL-9, IL-15 and TNF- α and the chemokines eotaxin and MIP-1 β were enhanced markedly after DC transfer. In addition, serum levels of arginase activity were decreased following DC transfer. Collectively, this study demonstrated the feasibility, safety and beneficial anti-tumour effects of OK432-stimulated DC infusion into tumour tissues for patients with cirrhosis and HCC, suggesting the ability of an active immunotherapeutic strategy to reduce tumour recurrence after locoregional treatment of HCC.

DCs were stimulated with OK432 prior to infusion into tumour tissues through an arterial catheter. OK432 was reported to activate DCs through its binding to TLR-2 and -4 [16,39] that can be used for cancer therapy [33]. The current results indicate that OK432 stimulation of immature DCs

from HCC patients promoted their maturation processes while preserving antigen uptake capacity and enhancing tumoricidal activity, consistent with previous observations [16,19] and supporting the current strategy in which OK432-stimulated DCs were infused directly into tumour tissues. Because the tumoricidal activity of unstimulated DCs was not observed in *in vitro* experiments, OK432 stimulation obviously altered the cytotoxic properties of DCs. One of the mechanisms of DC killing was reported to be CD40/CD40 ligand interaction [19]. Further studies are needed to determine the killing mechanisms of DCs derived from HCC patients in a direct [TNF, TNF-related apoptosis inducing ligand (TRAIL), Fas ligand, nitric oxide (NO) and perforin/granzyme] and indirect (MHC-restricted) manner [40–43]. Although the main mechanism by which OK432-stimulated DCs prolonged the recurrence-free survival was not elucidated, the tumoricidal activity of mature DCs was implicated in *in vivo* enhancement of antigen presentation, co-stimulation and inflammatory cytokine production.

Very recent reports document injection of OK432-stimulated DCs into patients with cancer of the gastrointestinal tract or pancreas [44,45], but their anti-tumour effects were not defined clearly. The current study shows for the first time that OK432-stimulated DCs induce beneficial anti-tumour responses when transferred into tumour tissues during TAE therapy. The anti-tumour responses may have been enhanced as a result of optimal activation of the DCs with OK432 or combining infusion of stimulated DCs with TAE therapy. Inappropriately activated DCs may be unable to generate sufficient numbers of properly activated effector T lymphocytes [46]. As shown in Fig. 1, all these alterations could contribute to the further enhancement of anti-tumour effects compared to those in our previous study with immature DCs [20]. Furthermore, the tumour cell death-promoting therapies, e.g. chemotherapy [47] and TAE [48], can be expected to enhance the effects of therapeutic cancer vaccines by redressing the immunosuppressive tumour environment.

NK cell activity and intracellular cytokine responses in CD4⁺ and CD8⁺ T lymphocytes and CD56⁺ NK cell subsets in PBMCs were not changed significantly in patients treated with OK432-stimulated DCs. Furthermore, we did not observe tumour antigen-specific T lymphocyte responses associated clearly with DC administration. The data suggest therefore that the immune responses induced by the therapy applied here were not detectable systemically. Because cytotoxic T lymphocyte responses were enhanced in patients receiving $> 3 \times 10^7$ cells [49,50], the numbers of transferred OK432-stimulated DCs were apparently not sufficient to induce responses detectable in the peripheral blood, but were enough to exert beneficial anti-tumour effects. In addition, many studies have concluded that cytotoxic T lymphocyte responses rarely predict clinical outcomes of DC-based immunotherapies [51,52] and that in many cases, also including our own studies

[28,30], tumour-specific effector T lymphocytes co-exist with the tumours. Consistent with these observations, the current results suggest that cytotoxic T lymphocyte responses in PBMCs are not reliable predictors of beneficial anti-tumour effects in patients treated with the current OK432-stimulated DC strategy.

Serum levels of the cytokines IL-9, IL-15 and TNF- α and the chemokines eotaxin and MIP-1 β were increased following OK432-stimulated DC transfer, but decreased after TAE therapy without DC administration. IL-9 and IL-15 belong to the cytokine receptor common gamma chain (γ ; CD132) family, a member of the type I cytokine receptor family expressed on most lymphocyte populations [53]. IL-9 exerts pleiotropic activities on T and B lymphocytes, mast cells, monocytes and haematopoietic progenitors [54,55]. IL-15 and TNF- α are known to prime T lymphocytes and NK cells when secreted by DCs [56] and to induce anti-tumour immune responses [57]. Eotaxin is known to selectively recruit eosinophils also contributing to anti-tumour effects [58,59], and MIP-1 β is a chemoattractant for NK cells, monocytes and a variety of other immune cells [60]. In addition, serum levels of arginase tended to decrease after DC transfer. Because serum arginase activity reflects the numbers of MDSCs that inhibit T lymphocyte responses in cancer patients [36], the patients treated with OK432-stimulated DCs might have developed lower levels of suppressor cells. Collectively, the results suggest that infusion of OK432-stimulated DCs may orchestrate the immune environment in the whole body that could enhance beneficial anti-tumour effects, although the precise molecular and cellular mechanisms associated with the actions of these cytokines and chemokines were not defined clearly in the current analysis.

Acknowledgements

The authors thank Kazumi Fushimi and Mariko Katsuda for technical assistance. We also thank the patients for participating in this trial. This work was supported in part by research grants from the Ministry of Education, Culture, Sports, Science, and Technology of Japan, the Ministry of Health, Labour and Welfare of Japan and the Japanese Society of Gastroenterology.

Disclosure

The authors have declared that no conflict of interest exists.

References

- 1 Omata M, Tateishi R, Yoshida H, Shiina S. Treatment of hepatocellular carcinoma by percutaneous tumor ablation methods: ethanol injection therapy and radiofrequency ablation. *Gastroenterology* 2004; 127:S159–66.
- 2 Belghiti J. Resection and liver transplantation for HCC. *J Gastroenterol* 2009; 44 (Suppl. 19):132–5.

- 3 Nakamoto Y, Guidotti LG, Kuhlen CV, Fowler P, Chisari FV. Immune pathogenesis of hepatocellular carcinoma. *J Exp Med* 1998; **188**:341–50.
- 4 Ercolani G, Grazi GL, Ravaioli M *et al.* Liver resection for hepatocellular carcinoma on cirrhosis: univariate and multivariate analysis of risk factors for intrahepatic recurrence. *Ann Surg* 2003; **237**:536–43.
- 5 Shankaran V, Ikeda H, Bruce AT *et al.* IFN γ and lymphocytes prevent primary tumour development and shape tumour immunogenicity. *Nature* 2001; **410**:1107–11.
- 6 Vulink A, Radford KJ, Melief C, Hart DN. Dendritic cells in cancer immunotherapy. *Adv Cancer Res* 2008; **99**:363–407.
- 7 Banchereau J, Briere F, Caux C *et al.* Immunobiology of dendritic cells. *Annu Rev Immunol* 2000; **18**:767–811.
- 8 Lemos MP, Esquivel F, Scott P, Laufer TM. MHC class II expression restricted to CD8 α + and CD11b+ dendritic cells is sufficient for control of *Leishmania major*. *J Exp Med* 2004; **199**:725–30.
- 9 Ni K, O'Neill HC. The role of dendritic cells in T cell activation. *Immunol Cell Biol* 1997; **75**:223–30.
- 10 Andrews DM, Andoniou CE, Scalzo AA *et al.* Cross-talk between dendritic cells and natural killer cells in viral infection. *Mol Immunol* 2005; **42**:547–55.
- 11 Heiser A, Coleman D, Dannull J *et al.* Autologous dendritic cells transfected with prostate-specific antigen RNA stimulate CTL responses against metastatic prostate tumors. *J Clin Invest* 2002; **109**:409–17.
- 12 Banchereau J, Steinman RM. Dendritic cells and the control of immunity. *Nature* 1998; **392**:245–52.
- 13 Forster R, Schubel A, Breitfeld D *et al.* CCR7 coordinates the primary immune response by establishing functional microenvironments in secondary lymphoid organs. *Cell* 1999; **99**:23–33.
- 14 MartIn-Fontecha A, Sebastiani S, Hopken UE *et al.* Regulation of dendritic cell migration to the draining lymph node: impact on T lymphocyte traffic and priming. *J Exp Med* 2003; **198**:615–21.
- 15 Ratzinger G, Stoitzner P, Ebner S *et al.* Matrix metalloproteinases 9 and 2 are necessary for the migration of Langerhans cells and dermal dendritic cells from human and murine skin. *J Immunol* 2002; **168**:4361–71.
- 16 Nakahara S, Tsunoda T, Baba T, Asabe S, Tahara H. Dendritic cells stimulated with a bacterial product, OK-432, efficiently induce cytotoxic T lymphocytes specific to tumor rejection peptide. *Cancer Res* 2003; **63**:4112–18.
- 17 Okamoto M, Oshikawa T, Tano T *et al.* Mechanism of anticancer host response induced by OK-432, a streptococcal preparation, mediated by phagocytosis and Toll-like receptor 4 signaling. *J Immunother* 2006; **29**:78–86.
- 18 Pasare C, Medzhitov R. Toll pathway-dependent blockade of CD4+CD25+ T cell-mediated suppression by dendritic cells. *Science* 2003; **299**:1033–6.
- 19 Hill KS, Errington F, Steele LP *et al.* OK432-activated human dendritic cells kill tumor cells via CD40/CD40 ligand interactions. *J Immunol* 2008; **181**:3108–15.
- 20 Nakamoto Y, Mizukoshi E, Tsuji H *et al.* Combined therapy of transcatheter hepatic arterial embolization with intratumoral dendritic cell infusion for hepatocellular carcinoma: clinical safety. *Clin Exp Immunol* 2007; **147**:296–305.
- 21 Steinman RM, Banchereau J. Taking dendritic cells into medicine. *Nature* 2007; **449**:419–26.
- 22 Tacken PJ, de Vries IJ, Torensma R, Figdor CG. Dendritic-cell immunotherapy: from *ex vivo* loading to *in vivo* targeting. *Nat Rev Immunol* 2007; **7**:790–802.
- 23 Makuuchi M. General rules for the clinical and pathological study of primary liver cancer, 2nd edn. Tokyo: Kanehara & Co., Ltd, 2003.
- 24 Veltri A, Moretto P, Doriguzzi A, Pagano E, Carrara G, Gandini G. Radiofrequency thermal ablation (RFA) after transarterial chemoembolization (TACE) as a combined therapy for unresectable non-early hepatocellular carcinoma (HCC). *Eur Radiol* 2006; **16**:661–9.
- 25 Dhodapkar MV, Steinman RM, Sapp M *et al.* Rapid generation of broad T-cell immunity in humans after a single injection of mature dendritic cells. *J Clin Invest* 1999; **104**:173–80.
- 26 Orange JS, Brodeur SR, Jain A *et al.* Deficient natural killer cell cytotoxicity in patients with IKK-gamma/NEMO mutations. *J Clin Invest* 2002; **109**:1501–9.
- 27 Klausner RD, Donaldson JG, Lippincott-Schwartz J, Brefeldin A: insights into the control of membrane traffic and organelle structure. *J Cell Biol* 1992; **116**:1071–80.
- 28 Mizukoshi E, Nakamoto Y, Marukawa Y *et al.* Cytotoxic T cell responses to human telomerase reverse transcriptase in patients with hepatocellular carcinoma. *Hepatology* 2006; **43**:1284–94.
- 29 Mizukoshi E, Nakamoto Y, Tsuji H, Yamashita T, Kaneko S. Identification of alpha-fetoprotein-derived peptides recognized by cytotoxic T lymphocytes in HLA-A24+ patients with hepatocellular carcinoma. *Int J Cancer* 2006; **118**:1194–204.
- 30 Mizukoshi E, Honda M, Arai K, Yamashita T, Nakamoto Y, Kaneko S. Expression of multidrug resistance-associated protein 3 and cytotoxic T cell responses in patients with hepatocellular carcinoma. *J Hepatol* 2008; **49**:946–54.
- 31 Rodriguez PC, Quiceno DG, Zabaleta J *et al.* Arginase I production in the tumor microenvironment by mature myeloid cells inhibits T-cell receptor expression and antigen-specific T-cell responses. *Cancer Res* 2004; **64**:5839–49.
- 32 Itoh T, Ueda Y, Okugawa K *et al.* Streptococcal preparation OK432 promotes functional maturation of human monocyte-derived dendritic cells. *Cancer Immunol Immunother* 2003; **52**:207–14.
- 33 Kuroki H, Morisaki T, Matsumoto K *et al.* Streptococcal preparation OK-432: a new maturation factor of monocyte-derived dendritic cells for clinical use. *Cancer Immunol Immunother* 2003; **52**:561–8.
- 34 Gunn MD, Kyuwa S, Tam C *et al.* Mice lacking expression of secondary lymphoid organ chemokine have defects in lymphocyte homing and dendritic cell localization. *J Exp Med* 1999; **189**:451–60.
- 35 Imanishi T, Akaza T, Kimura A, Tokunaga K, Gojobori T. HLA 1991, Proceedings of the Eleventh International Histocompatibility Workshop and Conference. Tokyo: Oxford University Press, 1992.
- 36 Zea AH, Rodriguez PC, Atkins MB *et al.* Arginase-producing myeloid suppressor cells in renal cell carcinoma patients: a mechanism of tumor evasion. *Cancer Res* 2005; **65**:3044–8.
- 37 Chrzanowska A, Krawczyk M, Baranczyk-Kuzma A. Changes in arginase isoenzymes pattern in human hepatocellular carcinoma. *Biochem Biophys Res Commun* 2008; **377**:337–40.
- 38 Caldwell S, Park SH. The epidemiology of hepatocellular cancer: from the perspectives of public health problem to tumor biology. *J Gastroenterol* 2009; **44** (Suppl. 19):96–101.
- 39 Okamoto M, Oshikawa T, Tano T *et al.* Involvement of Toll-like

- receptor 4 signaling in interferon-gamma production and antitumor effect by streptococcal agent OK-432. *J Natl Cancer Inst* 2003; **95**:316–26.
- 40 Liu S, Yu Y, Zhang M, Wang W, Cao X. The involvement of TNF-alpha-related apoptosis-inducing ligand in the enhanced cytotoxicity of IFN-beta-stimulated human dendritic cells to tumor cells. *J Immunol* 2001; **166**:5407–15.
 - 41 Lu G, Janjic BM, Janjic J, Whiteside TL, Storkus WJ, Vujanovic NL. Innate direct anticancer effector function of human immature dendritic cells. II. Role of TNF, lymphotoxin-alpha(1)beta(2), Fas ligand, and TNF-related apoptosis-inducing ligand. *J Immunol* 2002; **168**:1831–9.
 - 42 Nicolas A, Cathelin D, Larmonier N *et al*. Dendritic cells trigger tumor cell death by a nitric oxide-dependent mechanism. *J Immunol* 2007; **179**:812–18.
 - 43 Stary G, Bangert C, Tauber M, Strohal R, Kopp T, Stingl G. Tumoricidal activity of TLR7/8-activated inflammatory dendritic cells. *J Exp Med* 2007; **204**:1441–51.
 - 44 West E, Morgan R, Scott K *et al*. Clinical grade OK432-activated dendritic cells: *in vitro* characterization and tracking during intralymphatic delivery. *J Immunother* 2009; **32**:66–78.
 - 45 Hirooka Y, Itoh A, Kawashima H *et al*. A combination therapy of gemcitabine with immunotherapy for patients with inoperable locally advanced pancreatic cancer. *Pancreas* 2009; **38**:e69–74.
 - 46 Melief CJ. Cancer immunotherapy by dendritic cells. *Immunity* 2008; **29**:372–83.
 - 47 Zitvogel L, Apetoh L, Ghiringhelli F, Kroemer G. Immunological aspects of cancer chemotherapy. *Nat Rev Immunol* 2008; **8**:59–73.
 - 48 Ayaru L, Pereira SP, Alisa A *et al*. Unmasking of alpha-fetoprotein-specific CD4(+) T cell responses in hepatocellular carcinoma patients undergoing embolization. *J Immunol* 2007; **178**:1914–22.
 - 49 Thurner B, Haendle I, Roder C *et al*. Vaccination with mage-3A1 peptide-pulsed mature, monocyte-derived dendritic cells expands specific cytotoxic T cells and induces regression of some metastases in advanced stage IV melanoma. *J Exp Med* 1999; **190**:1669–78.
 - 50 Banchereau J, Palucka AK, Dhodapkar M *et al*. Immune and clinical responses in patients with metastatic melanoma to CD34(+) progenitor-derived dendritic cell vaccine. *Cancer Res* 2001; **61**:6451–8.
 - 51 Engell-Noerregaard L, Hansen TH, Andersen MH, Thor Straten P, Svane IM. Review of clinical studies on dendritic cell-based vaccination of patients with malignant melanoma: assessment of correlation between clinical response and vaccine parameters. *Cancer Immunol Immunother* 2009; **58**:1–14.
 - 52 Itoh K, Yamada A, Mine T, Noguchi M. Recent advances in cancer vaccines: an overview. *Jpn J Clin Oncol* 2009; **39**:73–80.
 - 53 Sugamura K, Asao H, Kondo M *et al*. The common gamma-chain for multiple cytokine receptors. *Adv Immunol* 1995; **59**:225–77.
 - 54 Temann UA, Geba GP, Rankin JA, Flavell RA. Expression of interleukin 9 in the lungs of transgenic mice causes airway inflammation, mast cell hyperplasia, and bronchial hyperresponsiveness. *J Exp Med* 1998; **188**:1307–20.
 - 55 McMillan SJ, Bishop B, Townsend MJ, McKenzie AN, Lloyd CM. The absence of interleukin 9 does not affect the development of allergen-induced pulmonary inflammation nor airway hyperreactivity. *J Exp Med* 2002; **195**:51–7.
 - 56 de Saint-Vis B, Fugier-Vivier I, Massacrier C *et al*. The cytokine profile expressed by human dendritic cells is dependent on cell subtype and mode of activation. *J Immunol* 1998; **160**:1666–76.
 - 57 Shanmugham LN, Petrarca C, Frydas S *et al*. IL-15 an immunoregulatory and anti-cancer cytokine. Recent advances. *J Exp Clin Cancer Res* 2006; **25**:529–36.
 - 58 Kataoka S, Konishi Y, Nishio Y, Fujikawa-Adachi K, Tomiyama A. Antitumor activity of eosinophils activated by IL-5 and eotaxin against hepatocellular carcinoma. *DNA Cell Biol* 2004; **23**:549–60.
 - 59 Simson L, Ellyard JI, Dent LA *et al*. Regulation of carcinogenesis by IL-5 and CCL11: a potential role for eosinophils in tumor immune surveillance. *J Immunol* 2007; **178**:4222–9.
 - 60 Bystry RS, Aluvihare V, Welch KA, Kallikourdis M, Betz AG. B cells and professional APCs recruit regulatory T cells via CCL4. *Nat Immunol* 2001; **2**:1126–32.

