

the most excellent staging system among the BCLC, Tokyo and JIS staging systems. Therefore, JIS score is currently considered to be the best integrated staging system in Japan. Regarding other integrated staging systems, modified JIS score has been reported<sup>13,18</sup> to be useful for patients undergoing hepatectomy. Biomarker combined JIS score has also been reported to be useful in discrimination in patients with good prognosis.<sup>19</sup> However, the usefulness of these new staging systems will remain unclear until they are assessed in a range of patient sets with HCC.

Regarding the estimation of HCC prognosis, most hepatologists recognize the importance of an integrated staging system rather than applying the TNM stage and hepatic functional reserve scales individually. Furthermore, the JIS score is considered to be the best integrated staging system for current clinical practice. However, it is still difficult to incorporate the integrated staging systems, such as the JIS score, into algorithms for HCC treatment.

*Recommendation 3. Integrated staging system should be used to assess the prognosis of patients with HCC, instead of individually applying scales for TNM stage and liver function stage.*

*Recommendation 4. The JIS score is the best staging system to estimate the prognosis of patients with HCC.*

*Informative Statement 1. Integrated staging systems, such as the JIS score, are not yet suitable for inclusion in algorithms for HCC treatment.*

## SURVEILLANCE AND DIAGNOSIS

### Surveillance programs

IT IS WELL known that HCC mainly occurs in cases with chronic liver disease, particularly cirrhosis. Several cohort studies have shown that the surveillance of high-risk patients with hepatitis B virus (HBV)- or hepatitis C virus (HCV)-related chronic liver disease improves the rate of early detection and the rate of curative treatments.<sup>20–27</sup> For this reason, UK<sup>28</sup>, European<sup>29</sup> and American<sup>3</sup> practice guidelines for HCC recommend routine surveillance of HCC among individuals with viral hepatitis or cirrhosis. Almost all gastroenterologists in Japan conduct surveillance programs using a combination of tumor markers such as AFP, the *lens culinaris* agglutinin-reactive fraction of AFP (AFP-L3%) and DCP, and by ultrasound (US).<sup>30</sup> However, no consensus has been reached in terms of the optimal surveillance strategy. Thompson *et al.* calculated the number of people

who need to be under surveillance to prevent either a single death from HCC or a single premature death (defined as death before the age 75 years) and showed the effectiveness of surveillance programs.<sup>31</sup> In the absence of surveillance, approximately 20% of the mixed etiology cohort died as a result of HCC.

*Recommendation 5. Surveillance with US and three tumor markers including AFP, DCP and AFP-L3 should be performed for early detection of HCC in patients with HBV- and HCV-related chronic liver disease, particularly cirrhosis.*

### Tumor markers

In Japan, AFP, AFP-L3 and DCP are widely and routinely used as serological tumor markers for the surveillance, diagnosis and prognostic estimation of HCC. The Evidence-Based Clinical Practice Guidelines of HCC published in 2005<sup>1</sup> recommended that AFP, AFP-L3 and DCP should be measured at intervals of 3–4 months for very high-risk patients (defined as HBV- or HCV-related liver cirrhosis), and at 6-month intervals for high-risk patients (defined as HBV- or HCV-related chronic liver disease or other causes of liver cirrhosis).<sup>32</sup> Although AFP is the most widely used tumor marker for HCC, the levels of AFP are also increased in patients with liver diseases other than HCC, including viral hepatitis, with a prevalence of 10–42%.<sup>33–35</sup> In contrast, AFP-L3 and DCP are very specific for HCC, compared with AFP alone. The combination assay for AFP, AFP-L3 and DCP should be performed for the early detection of HCC.<sup>36,37</sup> The specificity and sensitivity of the combination assay of AFP and DCP were 83% and 84%, respectively, to detect small HCC of less than 3 cm in diameter.<sup>38</sup> The specificity and sensitivity of the combination assay of DCP and AFP-L3 were 41.7–66.7% and 89.5–89.8%, respectively, to detect small HCC of less than 3 cm in diameter.<sup>39,40</sup>

*Recommendation 6. Periodical measurement of more than two kinds of tumor markers (particularly AFP and DCP) is recommended for the early detection of HCC in high-risk and very high-risk patients.*

*Recommendation 7. The surveillance interval needs to be shorter in very high-risk patients than in high-risk patients.*

### Imaging modalities

Periodic follow-up of chronic liver disease by US, multidetector row computed tomography (MDCT) and magnetic resonance imaging (MRI) allows relatively

easy detection of small HCC.<sup>41–43</sup> However, it is sometimes difficult to characterize small hepatic nodular lesions detected by these imaging modalities. Definitive diagnosis requires invasive methods such as US-guided liver biopsy. Hemodynamic evaluation of the nodule is also important to assess the biological behavior of HCC. The recent advances in MRI and computed tomography (CT) procedures, such as CT during hepatic arteriography (CTHA) and CT during arterial portography (CTAP), have enabled the detailed hemodynamic evaluation of small hepatic nodules.

Recently, liver-specific contrast agents such as superparamagnetic iron oxide particles (SPIO), which are taken up by Kupffer cells, and Gd-EOB-DTPA, which is taken up by hepatocytes, are frequently used in MRI for early diagnosis of HCC. Gd-EOB-DTPA is a superb agent because it provides dynamic and liver-specific MR images.<sup>44–46</sup> This contrast agent is highly liver specific; approximately 50% of the injected dose is taken up by functioning hepatocytes and is excreted in bile, compared with just 3–5% for gadobenate dimeglumine.<sup>46</sup> Early studies comparing Gd-EOB-DTPA-enhanced dynamic MRI with dynamic MDCT showed that Gd-EOB-DTPA-enhanced MRI is significantly more accurate, sensitive and specific than dynamic MDCT for the diagnosis of HCC in patients with cirrhosis.<sup>47,48</sup> In addition, Gd-EOB-DTPA-enhanced MRI has a high detection rate for early stage HCC nodules that are not enhanced in dynamic studies. However, although the differentiation of early HCC from dysplastic nodule by hepatobiliary phase images of Gd-EOB-DTPA MRI is promising, more data are still needed.

**Informative statement 2.** *Gd-EOB-DTPA-enhanced MRI provides dynamic and hepatocyte-specific images and is more accurate than dynamic MDCT or SPIO-MRI for the detection and characterization of small HCC, including early HCC.*

## ABLATION THERAPIES

**I**MAGE-GUIDED PERCUTANEOUS ablation therapies have long played important roles in the treatment of HCC. Percutaneous ethanol injection has been used for unresectable, small HCC since the early 1980s<sup>49–51</sup> and offers us the potential to treat HCC using non-surgical means. Percutaneous microwave coagulation therapy became popular in Japan in the late 1990s.<sup>52</sup> However, since the introduction of radiofrequency ablation (RFA) into clinical practice around 1999, there has been a dramatic shift from ethanol injection or microwave coagulation to RFA.<sup>53</sup> RFA for HCC has been covered by

public health insurance since April 2004 in Japan. Although more than 1700 institutions have experienced RFA in Japan, RFA is estimated to be performed routinely in approximately 1000 institutions throughout Japan at the present.

Radiofrequency ablation often seems to be performed with less than adequate treatment planning or preparation compared with surgical resection. RFA appears to be a very simple procedure. Thus, some physicians may perform RFA without adequate training or experience. In addition, RFA does not require expensive equipment. Thus, several hospitals have introduced RFA into clinical practice without high-performance US and CT.

However, RFA is indicated for malignant tumors and inadequate outcome should be avoided. Thus, only physicians with sufficient experience and appropriate skill should perform the procedure. Furthermore, only well-equipped hospitals should perform RFA because the outcomes of RFA are strongly influenced by the performance of the CT and US equipment available at each institution. It is crucial to offer consistent outcomes for RFA at all institutions and for all operators.

More importantly, before commencing RFA, the tumors should be evaluated by US, contrast-enhanced CT or MRI to determine tumor size, shape, number, presence or absence of extracapsular invasion, presence or absence of satellite lesions, location relative to Glisson's capsule or other critical structures, and to determine the optimal route to approach the tumor.

Within 1–3 days after RFA, contrast-enhanced CT or MRI is essential to objectively assess the treatment response. If the tumor is completely ablated with a sufficient safety margin, the treatment may be considered complete. However, if there is any residual cancer tissue or an insufficient safety margin, RFA should be repeated until complete tumor destruction with a sufficient ablative margin is achieved. The following recommendation was supported by 94% of the experts.

**Recommendation 8.** *Imaging should be performed within 1–3 days after RFA to evaluate treatment response. It is essential that RFA is repeated until entire tumor destruction with a sufficient ablative margin is achieved.*

For accurate tumor evaluation, CT and MRI performed before and after RFA should be done using a thin slice interval. The following recommendation was agreed by 94% of the experts.

**Recommendation 9.** *CT and MRI before and after RFA should be done using a slice thickness and interval of 5 mm or less; slice thickness and interval of 10 mm or more is not adequate.*

A histopathological study has revealed that, in cases with incomplete necrosis, viable cancer tissue remains around the main tumor, in portions isolated by the septa, or along the edge of the tumor after ablation therapies.<sup>54</sup> There may also be extranodular growth, satellite nodules or portal vein invasion, which cannot be detected by imaging modalities.<sup>55,56</sup> The incidence of satellite nodules and portal vein invasion is associated with the gross appearance of the main tumor. The single nodular type with extranodular growth and the confluent multinodular type both show satellite lesions more frequently than early HCC (vaguely nodular-type HCC showing preservation of the preexisting liver structure) and the single nodular type. Thus, it is important to determine the gross appearance of the tumor by imaging. It is also essential to ablate beyond the tumor border to achieve complete tumor necrosis and prevent local tumor progression (ablative margin or safety margin). Sonazoid-enhanced US in the Kupffer phase is useful to determine the gross tumor appearance.<sup>57</sup> The width of the safety margin should be modified based on the gross appearance of the tumor, the number of tumors, the initial tumor or recurrent tumor, the duration of time between the previous treatment and recurrence in recurrent cases, tumor location (particularly in relation to the Glisson's capsule), liver function, comorbid conditions and the patient's age.

Furthermore, the accuracy of contrast-enhanced CT or MRI for evaluating the extent of necrosis is limited because of the partial volume effect.<sup>58</sup> The following recommendation was agreed by 94% of the experts.

**Recommendation 10.** *A safety margin completely surrounding the lesion should be achieved in cases in which RFA is performed as a locally curative treatment (level 6, grade A).*

Ablation therapies, including RFA, are widely accepted as the preferred treatment for unresectable small HCC. On the other hand, it has been strongly debated whether ablation therapies can provide a treatment option for resectable HCC since the introduction of ethanol injection. Although the number of patients treated by RFA has steadily increased, the Clinical Practice Guidelines for Hepatocellular Carcinoma in Japan recommends surgery rather than ablation.<sup>1</sup> Their scientific statement recommends the following: "(i) if only one tumor is present, liver resection is recommended irrespective of the diameter of the tumor. Ablation therapy may also be selected if the severity of liver damage is class B and the diameter of the tumor is no more than 2 cm; (ii) if two to three tumors with diameters of no more than 3 cm are present, liver resection or

local ablation therapy is recommended". This scientific statement is based on a cohort study of patients at clinical stage I (fair liver function), with a solitary tumor of less than 2 cm in diameter, patients across all clinical stages with a solitary tumor greater than 2 cm, and patients of clinical stage II (moderately impaired liver function) with two tumors greater than 2 cm. In that cohort, those who underwent hepatic resection showed higher survival rates than those who received non-surgical interventions.<sup>59</sup>

However, those findings were not based on randomized controlled trials (RCT) and the different survival rates may be subject to bias arising from the background characteristics of the patients. Of note, the hepatic resection group was younger than the ethanol injection group. Furthermore, even among patients at clinical stage I, most patients with normal liver or chronic hepatitis seemed to undergo resection while many with cirrhosis seemed to receive ethanol injection. This might reduce the recurrence rate because of multicentric carcinogenesis and less frequent development of liver failure in the resection group. Moreover, the trend that patients with severe comorbid conditions, such as cardiopulmonary diseases and others, received ethanol injection rather than resection might explain some of the disparity in survival. By contrast, in one RCT the recurrence and survival rates were comparable between surgical resection and ethanol injection.<sup>60</sup> In addition, other non-randomized trials have reported similar or better overall survival after ethanol injection than after resection.<sup>61–63</sup>

In addition, the findings described above only compared resection with ethanol injection. For example, our RCT showed that RFA had higher survival and lower recurrence rates than ethanol injection while the adverse events were similar between the two therapies.<sup>64</sup> Similarly, other RCT have shown that RFA is superior to ethanol injection in terms of treatment outcomes for HCC.<sup>65–67</sup> Another RCT has shown that there was no difference between resection and RFA in terms of overall and disease-free survival, while post-treatment complications occurred more frequently and were more severe after surgery.<sup>68</sup>

Hence, it is inappropriate to generalize the findings for ethanol injection to other percutaneous local ablation therapies such as RFA, and it should not be concluded that hepatectomy is recommended over percutaneous local ablation.

Further trials are needed to determine whether RFA can become a preferred treatment for "resectable HCC". In such trials, the primary end-point should be overall

survival.<sup>69</sup> The AASLD practice guideline clearly states the following: “although a treatment might be less active against the tumor than another treatment and thus result in a higher recurrence rate after initial treatment, the overall survival might not differ or may even be better”.<sup>3</sup>

Recurrence-free survival can be misleading and should not be considered as a surrogate end-point for overall survival. In HCC, unlike other solid tumors, recurrence can still be treated, and the first recurrence does not cause death in most cases. Furthermore, surgery theoretically offers better disease-free survival than RFA because it removes larger liver tissue. However, the better curability associated with hepatectomy could be cancelled out by the surgical invasion and the potential deterioration in liver function. The following recommendation was agreed by 84% of the experts.

**Recommendation 11.** Overall survival should be the end-point to compare results between ablation and hepatectomy.

## SURGICAL TREATMENT: RESECTION AND TRANSPLANTATION

A NATIONWIDE SURVEY by the Japanese Liver Transplantation Society found that a total of 4725 cases of living-donor liver transplantations (LDLT) were reported in Japan as of the end of 2007 since its initiation in 1989. By contrast, during the same period, only 46 cases of deceased-donor liver transplantation (DDLT) were documented. At the end of 2006, 778 patients with HCC had

undergone an LDLT in Japan.<sup>70</sup> Because of the severe shortage of brain-dead donors and the extremely long waiting time for such organs, DDLT is not a realistic treatment option for HCC patients in Japan.

## Algorithm for the treatment of patients with HCC in Japan

Figure 1 shows the treatment algorithm presented in the Japanese evidence-based guideline for the diagnosis and treatment of HCC.<sup>1</sup> Liver transplantation is recommended for HCC patients with liver damage C (similar to Child–Pugh C), but only when the patients meet the Milan criteria proposed by Mazzaferro.<sup>71</sup> In the revised version of the guidelines published at the end of 2009, an age limit of 65 years was added to the criteria for liver transplantation.

## Can the indications for liver transplantation be expanded beyond the Milan criteria?

Until the mid-1990s, HCC was considered a contraindication for liver transplantation because of the extremely poor outcome of early series.<sup>72,73</sup> This pessimistic view was reversed by Mazzaferro *et al.* who conducted a prospective cohort study to identify subgroups of HCC patients who may benefit from DDLT. They presented clear eligibility criteria for transplantation, as follows: the presence of a solitary tumor of 5 cm or less in diameter and no more than three tumor nodules, each 3 cm or less in diameter, in patients with multiple tumors, and the absence of vascular invasion or extrahepatic disease. In their series, the overall and recurrence-free survival rates




Figure 1 Japanese evidence-based treatment algorithm. HCC, hepatocellular carcinoma.

at 4 years for 35 patients who met the above criteria were as high as 85% and 92%, respectively. These criteria were named the “Milan criteria” and became the gold standard for patient selection for liver transplantation. The Milan criteria were also validated for LDLT using data from a nationwide survey in Japan.<sup>74</sup> Since 2004, LDLT for HCC has been covered by social medical insurance in Japan when the preoperative imaging studies indicate that the patient’s condition meets the Milan criteria.

The Milan criteria have encouraged transplant surgeons to increase the number of liver transplantations performed in HCC patients, and the United Network for Organ Sharing (UNOS) has incorporated the Milan criteria as conditions for listing HCC patients. During the extensive application of liver transplantation for HCC, transplant surgeons have noticed that the outcomes of some patients who slightly exceeded the Milan criteria were also favorable. To expand the indications for liver transplantation, several groups from different countries have challenged these restrictive criteria (Table 1).<sup>75–79</sup> Yao *et al.* at the University of California at San Francisco (UCSF) proposed criteria consisting of a single tumor of less than 6.5 cm in diameter or two lesions of less than 4.5 cm in diameter, with a total tumor diameter of less than 8 cm; these criteria are known as the “UCSF criteria”.<sup>76</sup> The utility of the UCSF criteria was subsequently confirmed by the University of California at Los Angeles.<sup>80</sup>

Regarding the indications for LDLT in HCC patients, several proposals from Asian centers have extended the eligibility criteria (Table 1). For example, a group at the University of Tokyo proposed the “5-5 rule”, which allows up to five nodules with a maximum diameter of 5 cm.<sup>77</sup> The 3-year recurrence-free rate of 72 patients who met the Tokyo 5-5 rule was as high as 94%, which was comparable with that of patients within the Milan criteria. A group at the University of Kyoto subsequently proposed a further expansion of the criteria, increasing the upper limit of the number of tumors to 10.<sup>79</sup>

Because LDLT is not governed by an organ-sharing system, some authors have argued that the indications

for LDLT in patients with HCC could be further extended. One might say that “If the patient (recipient) and his/her family (donor) strongly wish to undergo LDLT even in cases of very advanced HCC with full knowledge of potential for poor outcomes, there is no reason for transplant surgeons to reject their wish. The family members may accept the poor outcome after LDLT without doing any harm to the community.” However, we should always remember that, while LDLT does not require a donor from the community, it does require extensive medical resources, including a large workload for surgeons and other hospital staff members, medical supplies, drugs and blood products. Furthermore, the premature death of the recipient is well known to cause severe emotional trauma to the living donors and their family members.

Based on an answer-pad vote at the consensus meeting of 45th JSH congress, 84% of the experts supported keeping the Milan criteria for DDLT, but only 25% supported keeping these criteria for LDLT. Although any expansion of the criteria should be modest, no consensus exists as to the extent to which the criteria can be extended.

*Recommendation 12. For DDLT, the HCC status of the recipients should meet the Milan criteria.*

*Recommendation 13. For LDLT, the HCC status of the recipients does not need to be within the Milan criteria.*

### Which is better, liver resection or transplantation, for HCC patients who are eligible for either treatment?

Because liver transplantation replaces the whole liver, removing the highly carcinogenic background and the cirrhotic liver can avoid multicentric or de novo cancer recurrence.<sup>80</sup> In contrast, liver resection is associated with a very high risk of tumor recurrence. Even after curative liver resection in patients with good liver function, the 5-year recurrence rate is as high as 70–79%.<sup>80</sup> Roughly half of these recurrences are multicentric or de novo recurrences. For this reason, liver transplantation

**Table 1** Summary of proposed criteria for indication of liver transplantation for HCC

Criteria	Conditions	References
Milan criteria	Up to 5 cm for single nodule or up to 3 nodules with a maximum diameter of 3 cm	70
UCSF criteria	Up to 6.5 cm for single nodule or up to 3 nodules with a maximum diameter of 4.5 cm	76
Tokyo 5-5 rule	Up to 5 nodules with a maximum diameter of 5 cm	77
Asan criteria	Up to 6 nodules with a maximum diameter of 5 cm	78
Kyoto criteria	Up to 10 nodules with a maximum diameter of 5 cm and PIVKA-II <400 mAU/mL	79
Up-to-seven criteria	Up to seven as the sum of the size of the largest tumor [in cm] and the number of tumors	75

may be recommended for HCC patients with good liver function who are also eligible for liver resection, as in Western countries.

Another issue is the operative risk of the two treatments. In Japan, the operative mortality rates for LDLT and liver resection are estimated to be 4–10% and 0.8–1.2%, respectively. This striking difference in operative mortality rates might preclude LDLT for patients with good liver function.

Using two databases at the National Cancer Center Hospital in Japan and the University of Pittsburgh Medical Center in the USA, Yamamoto *et al.* compared the long-term outcome of liver resection and transplantation in cirrhotic patients with HCC.<sup>81</sup> The overall survival of Child–Pugh A patients who underwent liver resection was similar to that of the patients without vascular invasion or lymph node metastases who underwent transplantation (most cases with Child–Pugh C). The recurrence rate was significantly lower in the transplantation group. For cases in which either treatment can be performed, the outcome of liver transplantation might be better than that of hepatic resection, particularly in cases with only a few small lesions.<sup>81,82</sup> In cases with large lesions, superior outcomes are achieved with hepatectomy. Because some patients may withdraw from treatment during the pre-transplantation period,<sup>83</sup> the outcomes with resection are better than those for liver transplantation based on intention-to-treat analysis of patients who meet the criteria for resection.

The evidence-based guideline<sup>1</sup> recommends the following: considering the occurrence of dropouts during the pre-transplantation period, the outcome of resection is better than that of liver transplantation among patients who meet the criteria for resection (grade B).

According to a question and answer-analyzer vote at this consensus meeting, 83% of the HCC experts selected LDLT for Child–Pugh C patients meeting the Milan criteria, whereas only 15–19% of the audience selected LDLT for Child–Pugh A or B patients.

*Recommendation 14. LDLT should not be recommended for HCC patients with Child–Pugh A or B liver function.*

## PALLIATIVE TREATMENTS: TRANSARTERIAL CHEMOEMBOLIZATION AND CHEMOTHERAPY

PALLIATIVE TREATMENTS FOR HCC include transarterial chemoembolization (TACE), hepatic arterial infusion chemotherapy (HAIC) and systemic chemotherapy.

## Transarterial embolization/TACE

Transcatheter arterial embolization (TAE)/TACE is one of the treatment options to treat hypervascular HCC. The theoretical basis of embolization is to induce ischemic tumor necrosis by acute arterial occlusion in hypervascular classical HCC. Embolization may be done alone (TAE) or in combination (TACE) with antineoplastic agents such as doxorubicin, epirubicin or cisplatin and a contrast agent, lipiodol. TACE is more effective and, thus, more widely used than embolization alone.

The technique for TACE is well established. The subsegmental artery or a peripheral artery near the target tumor is selected by a micro-catheter technique, followed by selective injection of antineoplastic agents mixed with lipiodol (lipiodol emulsion). The artery is then selectively obstructed with gelatin sponge particles. For bi-lobular multiple HCC with moderately impaired hepatic function (Child–Pugh B), TACE might need to be performed twice with an interval of several weeks to avoid hepatic decompensation.

The survival benefit of TAE/TACE was controversial until the publication of two RCT in 2002, which showed that TACE improved the survival of selected patients (Child–Pugh A with no vascular invasion) compared with conservative treatment.<sup>84,85</sup> A subsequent meta-analysis of seven RCT comparing TAE/TACE as a primary treatment for HCC in comparison with conservative management and/or suboptimal therapies showed a significant improvement in the 2-year survival, favoring TAE/TACE (odds ratio [OR] = 0.53; 95% confidence interval [CI] = 0.32–0.89,  $P = 0.017$ ).<sup>86,87</sup>

According to the Nationwide Follow-up Survey of Primary Liver Cancer in Japan, one-third of all patients with primary HCC were treated by TAE/TACE (Fig. 2). Thus, TAE/TACE, hepatic resection and local ablation therapy are commonly used in Japan. TAE/TACE is the most widely used treatment for unresectable HCC.

In two Japanese treatment guidelines for HCC, evidence-based<sup>1,30,88</sup> and consensus-based guidelines,<sup>89</sup> TACE is recommended for patients with the severity of the liver damage categorized into A or B, in whom there are two or three tumors with a diameter greater than 3 cm, or four or more tumors.

In early stages of HCC, TACE is not indicated as first-line treatment because the outcome review of the Nationwide Follow-up Survey by the LCSGJ reported worse results for TACE than surgery or percutaneous ablation. This survey revealed that the 5-year survival rates for resection, ablation and TACE were 59.2%,




Figure 2 Change of treatment method for hepatocellular carcinoma in Japan. TACE, transcatheter arterial chemoembolization.

48.4% and 29.7%, respectively, for single tumors, and 46.4%, 37.3% and 23.0%, respectively, for two tumors.<sup>90</sup>

In contrast, in a large prospective cohort study of 8510 patients who received TACE for unresectable HCC, according to the LCSGJ, the median survival was 34 months with 1-, 2-, 3-, 5- and 7-year survival rates of 82%, 63%, 47%, 26% and 16%, respectively.<sup>91</sup> In patients with early stage HCC, single tumors of 2 cm or more and preserved liver function (clinical stage I and liver damage A according to the LCSGJ),<sup>92</sup> the median survival was 62 months with 1-, 2-, 3-, 5- and 7-year survival rates of 98%, 92%, 73%, 52% and 38%, respectively.<sup>91</sup> These results for TACE with early stage HCC seem comparable with those for surgery or ablation. Thus, although curative therapies are highly recommended for patients with early stage HCC, TACE can be applied in these patients contraindicated for curative therapies.

Transcatheter arterial chemoembolization can be used in combination with percutaneous ablation, including RFA. A meta-analysis of four RCT comparing combina-

tion therapy (TACE plus percutaneous ethanol injection [PEI]) or RFA versus monotherapy (TACE alone, PEI or RFA alone) showed a significant decrease in mortality favoring combination therapy versus monotherapy in patients with small (<3 cm) or large (>3 cm) HCC (OR = 0.534; 95% CI = 0.288–0.990;  $P = 0.046$ ).<sup>93</sup>

In RFA treatment, as the tumor size increases, the therapeutic response decreases because of the limited volume of coagulation necrosis induced by the electrode. Blood flow also promotes heat loss to result in insufficient necrosis; therefore, reducing blood flow during RFA increases the ablation volume. Therefore, it seems to be reasonable to perform RFA after reducing blood flow by preceding RFA with TACE. Several cohort studies have shown that performing TACE before RFA is feasible and safe, and offers a useful treatment in compensated cirrhosis (Child–Pugh A or B) with relatively small HCC nodules (20–50 mm).<sup>94–97</sup> RFA in combination with preceding TACE is already recommended in the consensus-based treatment algorithm proposed by the JSH.<sup>89</sup>

In the current consensus meeting, for hypervascular HCC of 2 cm in size, 51% of the experts used TACE

before RFA treatment. By contrast, for hypervascular HCC of 3 cm in size, 81% of the experts performed TACE before RFA. This is theoretically reasonable because the possibility of incomplete ablation is greater for tumors of 2–3 cm in size, compared with tumors of less than 2 cm in size, based on the limited volume possible with a single ablation procedure. Additionally, the accumulation of lipiodol in the tumor should facilitate the decision on whether additional RFA treatment is required following the response evaluation by dynamic CT scan. However, the survival benefit of TACE in combination with RFA should be verified by well-designed RCT.

Transcatheter arterial chemoembolization is performed in various stages in the clinical management of HCC, not only for the initially detected HCC, but also for recurrent HCC. TACE has been shown to be valuable for improving the overall survival of HCC patients, although it is difficult to assess its clinical efficacy as second- or third-line therapy.

*Informative Statement 3. TACE performed before RFA is favorable for the curative treatment of hypervascular HCC of 2–3 cm in size.*

*Recommendation 15. TACE performed before RFA is recommended for curative treatment of hypervascular HCC larger than 3 cm in size.*

## Chemotherapy

Chemotherapy for HCC is divided into two types according to the route of administration; the first is systemic chemotherapy and the second is hepatic arterial infusion chemotherapy (HAIC). Systemic chemotherapy can also be divided into two types: intravenous and oral chemotherapy.

According to the Nationwide Follow-up Survey of Primary Liver Cancer by the LCSGJ, chemotherapy is used in 3.4–5.5% of primary HCC patients (Fig. 2). HAIC is theoretically more favorable for HCC than systemic chemotherapy because hepatic arterial infusion of anticancer drugs enables the delivery of high doses of drugs directly to the hypervascular HCC. In addition, HAIC provides a lower systemic level of the drugs than systemic administration, because the first-pass effect in the liver, and thus reduces toxicity and side-effects. Because of these advantages, HAIC is frequently used in Japan for intrahepatic advanced HCC with portal vein tumor thrombosis and/or intrahepatic multiple HCC. A recent report from the Japanese Nationwide Survey revealed that almost 90% of the chemotherapeutic regimens for HCC are done by hepatic arterial infusion. Thus, HAIC has become widely used in Japan, despite

there being no solid evidence for a survival benefit of HAIC compared with systemic chemotherapy or best supportive care (Fig. 3).


*Recommendation 16. HAIC is recommended for advanced HCC with major portal vein tumor thrombi with preserved liver function.*

Various anticancer drugs and treatment regimens are used for HAIC in Japan. Two regimens in particular are widely used for HAIC. The first is interferon (IFN) in combination with 5-fluorouracil (5-FU); the second is low-dose cisplatin (CDDP) in combination with 5-FU. For IFN plus 5-FU, the response rate was reported to be 52.6%, with 16.4% achieving complete response (CR) and 36.2% achieving partial response (PR) among 116 patients with tumor thrombosis of the major portal vein or first branches of the portal vein. The survival rates at 6, 12 and 24 months were 53%, 34% and 18%, respectively, with a median survival of 6.9 months, compared with survival rates of 40%, 15% and 5%, respectively, in the historical control group.<sup>98</sup> The survival was significantly different between the two groups ( $P < 0.01$ ). For low-dose CDDP plus 5-FU, the response rate was 48%, including 8% with CR and 40% with PR among 48 patients with portal vein tumor thrombosis. The 1-, 2-, 3- and 5-year cumulative survival rates were 45%, 31%, 25% and 11%, respectively, with a median survival of 10.2 months.<sup>99</sup>

In a review of previously reported small-size phase II studies of HAIC for advanced HCC,<sup>10,17,98–108</sup> the response rate varied from 14% to 71%. The mean survival duration also varied from 2.6 months to 32.4 months. However, few reports have compared systemic chemotherapy or HAIC using cytotoxic agents with placebo or best supportive care (Table 2).

The results of a randomized placebo-controlled double-blind phase III study with the multikinase inhibitor sorafenib were recently reported, representing a breakthrough in the chemotherapy for advanced HCC. Sorafenib is an oral drug that inhibits the platelet-derived growth factor (PDGF)-R, vascular endothelial growth factor (VEGF)-R, c-Kit-R and raf signaling pathways in tumor cells and in surrounding endothelial cells. In that study, 602 patients with advanced HCC, who were not indicated for other loco-regional treatments such as hepatic resection, who had not received prior systemic treatment and who had good liver functional reserve (Child–Pugh A) were randomized to sorafenib (400 mg b.i.d.) or placebo. Sorafenib was well tolerated and yielded a statistically significant improvement (44%) in overall survival. The median survival increased from 7.9 to 10.7 months (hazard ratio, 0.69;





**Figure 3** Consensus-based treatment algorithm for hepatocellular carcinoma proposed by the Japan Society of Hepatology (JSH) revised in 2010. (1) Treatment should be performed as if extrahepatic spread is negative, when extrahepatic spread is not regarded as a prognostic factor. (2) Sorafenib is the first choice of treatment in this setting as a standard of care. (3) Intensive follow-up observation is recommended for hypovascular nodules by the Japanese Evidence-Based Clinical Practice Guidelines. However, local ablation therapy is frequently performed in the following cases: (i) when the nodule is diagnosed pathologically as early hepatocellular carcinoma (HCC); (ii) when the nodules show decreased uptake on gadolinium ethoxybenzyl magnetic resonance imaging (Gd-EOB-MRI); or (iii) when the nodules show decreased portal flow by computed tomography during arterial portography (CTAP), because these nodules are known to frequently progress to the typical advanced HCC. (4) Even for HCC nodules exceeding 3 cm in diameter, combination therapy of transcatheter arterial chemoembolization (TACE) and ablation is frequently performed when resection is not indicated. (5) TACE is the first choice of treatment in this setting. Hepatic arterial infusion chemotherapy (HAIC) using an implanted port is also recommended for TACE refractory patients. The regimen for this treatment is usually low-dose FP (5-fluorouracil [5-FU] + cisplatin [CDDP]) or intra-arterial 5-FU in fusion combined with systemic interferon therapy. Sorafenib is also a treatment of choice for TACE/HAIC refractory patients with Child–Pugh A liver function. (6) Resection is sometimes performed even when numbers of nodules are over 4. Furthermore, ablation is sometimes performed in combination with TACE. (7) Milan criteria: tumor size  $\leq 3$  cm and tumor numbers  $\leq 3$ ; or solitary tumor  $\leq 5$  cm. Even when liver function is good (Child–Pugh A/B), transplantation is sometimes considered for relatively younger patients with frequently or early recurring HCC after curative treatments. (8) HAIC or sorafenib is recommended for HCC patients with Vp3 (portal invasion at the 1<sup>st</sup> portal branch) or Vp4 (portal invasion at the main portal branch). Sorafenib is only recommended for HCC patients with Child–Pugh A liver function. (9) Resection and TACE is frequently performed when portal invasion is minimal such as Vp1 (portal invasion at the 3<sup>rd</sup> or more peripheral portal branch) or Vp2 (portal invasion at the 2<sup>nd</sup> portal branch). (10) Local ablation therapy or subsegmental TACE is performed even for Child–Pugh C patients when transplantation is not indicated when there is no hepatic encephalopathy, no uncontrollable ascites and a low bilirubin level ( $<3.0$  mg/dL). However, it is regarded as an experimental treatment since there is no evidence of its survival benefit in Child–Pugh C patients. A prospective study is necessary to clarify this issue. Even in Child–Pugh A/B patients, transplantation is sometimes performed for relatively younger patients with frequently or early recurring HCC after curative treatments.

Table 2 Response rates and survival periods in studies of intrahepatic arterial infusion chemotherapy for advanced hepatocellular carcinoma

	Drugs	No. of Patients	Response rate (CR + PR, %)	Median survival time (months)	References
Single	Doxorubicin (IHAC)	72	60	7.0	Tzorcoleftherakis <i>et al.</i> <sup>102</sup>
	Doxorubicin (systemic)		44.1	6.5	
	CDDP	67	37	10.7	Court <i>et al.</i> <sup>103</sup>
Multiple	CDDP, 5-FU (low FP)	52	71	ND	Okuda <i>et al.</i> <sup>104</sup>
	CDDP, 5-FU (low FP)	48	48	10.2	Ando <i>et al.</i> <sup>99</sup>
	CDDP, 5-FU (low FP)	37	56.3	32.4	Sumie <i>et al.</i> <sup>101</sup>
	CDDP, 5-FU (low FP)	38	47	6.2	Tanioka <i>et al.</i> <sup>105</sup>
	CDDP, 5-FU	41	22	12.0	Park <i>et al.</i> <sup>106</sup>
	CDDP, Mitomycin C, 5-FU, LV	53	28.3	13.2	Lin <i>et al.</i> <sup>100</sup>
	IFN, CDDP	68	33	4.4	Chung <i>et al.</i> <sup>107</sup>
	CDDP		14	2.6	
	BSC			1.2	
	IFN, CDDP, 5-FU, MTX, LV	34	45	ND	Kaneko <i>et al.</i> <sup>108</sup>
	IFN, 5-FU	116	52	6.9	Obi <i>et al.</i> <sup>98</sup>

IHAC, intrahepatic arterial chemotherapy; CDDP, cisplatin; 5-FU, 5-fluorouracil; low FP, 5-fluorouracil + cisplatin. BSC, best support care; IFN, interferon; LV, leucovorin; MTX, methotrexate.

95% CI = 0.55–0.87). Side-effects included hand–foot skin reaction, diarrhea and fatigue, but sorafenib was not found to be toxic to the liver.<sup>109</sup> Similar findings were reported in a subsequent Asia–Pacific RCT.<sup>110</sup>

Based on the results of these RCT, sorafenib has become the first-line therapy for advanced HCC worldwide. Some Japanese experts for HCC are claiming low response rates, although the survival was significantly prolonged compared with placebo. This phenomenon could be explained by a longer period with stable disease with sorafenib than with placebo, or the necrotic change in the tumor is present without size reduction.

In Japan, sorafenib was approved for the treatment of HCC on 20 May 2009. In the consensus meeting held in June, 35% of the Japanese experts agreed that sorafenib should be selected as the first-line therapy for advanced HCC considered unsuitable for resection, RFA or TACE. A further 36% of the experts were undecided because they did not have enough experience with using sorafenib.

*Informative Statement 4. Sorafenib is the first-line therapy for advanced HCC with major vascular invasion and/or extrahepatic spread and good liver function. However, further studies are needed to compare the overall efficacy of HAIC and sorafenib.*

## TREATMENT ALGORITHM

TO TREAT HCC, the most appropriate therapeutic option needs to be selected among the available treatment modalities, including resection, percutaneous ablation, TACE and transplantation, but few evidence-based guidelines have been developed to aid decision-making.<sup>1,28,29,88,89,111</sup> Recently, two treatment algorithms for HCC have been proposed in the Japanese guidelines. The profile of these algorithms is briefly described here, in addition to the results of two questions and answers at the JSH Consensus Meeting for HCC at Kobe.

### Evidence-based treatment algorithm

The Clinical Practice Guidelines for HCC was established in 2005 based on evidence-based methodology, and covers six topics including prevention, diagnosis, surgery, chemotherapy, TACE and percutaneous ablation. To develop these guidelines, a systematic review of the English medical published work was performed and a total of 7118 articles on HCC were identified, mainly from MEDLINE (1966–2002), of which 334 were selected based on the evidence level to form 58 pairs of clinical questions and recommendations.<sup>1,88</sup> For convenience in clinical use, two algorithms were created for

the surveillance and treatment of HCC. A full English version was uploaded to the website of the JSH ([www.jsh.or.jp/](http://www.jsh.or.jp/)) in 2006.

The treatment algorithm for HCC was made on the basis of three independent factors: degree of liver damage, tumor number and tumor size. For the resulting six patients' subgroups, the first- and second-line therapies were recommended as objectively as possible (Fig. 1). The degree of liver damage is a modified system based on the Child–Pugh classification: "encephalopathy" was replaced by ICGR<sub>15</sub>, to provide an accurate evaluation of liver functional reserve, particularly in surgical candidates.

Patients with mild (class A) or moderate (class B) liver damage are subject to the following recommendations: (i) in patients with a single tumor, liver resection is recommended, irrespective of the tumor size (percutaneous ablation may be performed if liver damage is of class B and the tumor is no more than 2 cm in size); (ii) for patients with two or three tumors smaller than 3 cm, resection or ablation are recommended; (iii) for patients with two or three tumors larger than 3 cm, resection or TACE are recommended; and (iv) for patients with more than four tumors, TACE or HAIC is recommended. The recommendations for patients with severe (class C) liver damage are as follows: (v) in patients with tumor(s) meeting the Milan criteria, liver transplantation is recommended; and (vi) for patients with more than four tumors, palliative treatment is recommended. For patients with extrahepatic metastasis, chemotherapy may be performed.

The rationale for selecting resection or ablation in patients with class A or B liver damage is based on the outcome of the largest multicenter study involving 12 888 patients in Japan.<sup>59</sup> The recommendation for TACE is based on the findings of two RCT showing a significant improvement in the survival of patients with multiple tumors and class A or B liver damage.<sup>84,85</sup> The indication for liver transplantation is derived from a prospective cohort study using the Milan criteria,<sup>71</sup> and a nationwide survey of Japan justifying the criteria in living donor transplantation.<sup>74</sup>

### Consensus-based treatment algorithm

An expert panel of the JSH established a consensus-based treatment algorithm based on the therapeutic policies that are widely used in Japan.<sup>89,111</sup> This algorithm categorizes the patients on five clinical variables (extrahepatic spread, liver function, vascular invasion, tumor number and tumor size), and it divides the treatment options into resection, ablation, TACE, HAIC, liver

transplantation and palliative treatment (Fig. 3).<sup>89,111</sup> Because of the recent introduction of sorafenib in Japan, this consensus-based treatment algorithm was further revised and approved by the experts at the consensus meeting.<sup>111,112</sup>

Essentially, the consensus-based algorithm follows the evidence-based algorithm, but the treatments widely used in Japan were included by consensus, even though the evidence may be weak. The major differences in the consensus-based algorithm include: (i) ablation is sometimes performed in patients with a single, hypovascular early HCC; (ii) sorafenib is recommended for use in Child–Pugh A patients with vascular invasion, TACE failure or extrahepatic spread of HCC,<sup>109,112</sup> and (iii) liver transplantation is recommended, even for Child–Pugh A/B patients, if the Milan criteria are met.

The consensus-based algorithm based on the consensus of a large number of specialists, and a treatment strategy for management of HCC in Japan is important, and should be revised based on prospective trials for aspects of the algorithm lacking sufficient evidence.<sup>111,112</sup>

**Informative statement 5.** RFA might be recommended as a first-line treatment option in patients with a single, hypervascular HCC of less than 2 cm in size and with preserved liver function (Child–Pugh A or Liver Damage Class A). However, there was a discrepancy between surgeons and non-surgeons for this statement. This statement is strongly supported by non-surgeons (68%), whereas 80% of the surgeons favor resection rather than RFA. **Recommendation 17.** Resection should be considered as the first-line treatment option for patients with a single, hypervascular HCC of 3 cm or more in size and with preserved liver function (Child–Pugh A or Liver Damage Class A).

The revised version of the consensus-based treatment algorithm for HCC proposed by the JSH (Fig. 3) should aid decision-making at every stage in clinical practice. By sharing the information contained within the treatment algorithm chart, the physicians can offer recommended treatment options to the patient who can then choose one based on their preference (Fig. 3).

## CONCLUSIONS

THIS CONSENSUS STATEMENT is a conclusion of the consensus meeting of HCC, which was held at the 45th JSH meeting, Kobe, Japan on 4–5 June 2009 (Congress President: Professor Masatoshi Kudo). This manuscript and recommendations largely reflect the daily practice in the real world carried out throughout

Japan. The biggest difference of Japan's HCC practice from Western countries are pathological assessment issue, prognostic staging system, surveillance and diagnostic strategy, treatment strategy including role of HAIC, and method of RFA procedure, and treatment algorithm shown in Figure 3.

We believe every reader of this manuscript will well understand the real Japanese HCC practice much better than the other already published arterial articles. It is needless to say that consensus statements like this article should be regularly revised every 3–4 years because solid evidence or new diagnostic and treatment tool/drug or concept will be published and then established in clinical practice every year.

## REFERENCES

- 1 Makuuchi M, Kokudo N, Arai S *et al.* Development of evidence-based clinical guidelines for the diagnosis and treatment of hepatocellular carcinoma in Japan. *Hepatology Res* 2008; 38 (1): 37–51.
- 2 Liver Cancer Study Group of Japan. *General Rules for the Clinical and Pathological Study of Primary Liver Cancer*. English 2nd edn. Tokyo: Kanehara; 2003.
- 3 Bruix J, Sherman M. Management of hepatocellular carcinoma. *Hepatology* 2005; 42 (5): 1208–36.
- 4 Rockey DC, Caldwell SH, Goodman ZD, Nelson RC, Smith AD. Liver biopsy. *Hepatology* 2009; 49 (3): 1017–44.
- 5 Liu YW, Chen CL, Chen YS, Wang CC, Wang SH, Lin CC. Needle tract implantation of hepatocellular carcinoma after fine needle biopsy. *Dig Dis Sci* 2007; 52 (1): 228–31.
- 6 Chang S, Kim SH, Lim HK, Lee WJ, Choi D, Lim JH. Needle tract implantation after sonographically guided percutaneous biopsy of hepatocellular carcinoma: evaluation of doubling time, frequency, and features on CT. *AJR Am J Roentgenol* 2005; 185: 400–5.
- 7 Durand F, Regimbeau JM, Belghiti J *et al.* Assessment of the benefits and risks of percutaneous biopsy before surgical resection of hepatocellular carcinoma. *J Hepatol* 2001; 35: 254–8.
- 8 Takamori R, Wong LL, Dang C, Wong L. Needle-tract implantation from hepatocellular cancer: is needle biopsy of the liver always necessary? *Liver Transpl* 2000; 6: 67–72.
- 9 Sakamoto M, Hirohashi S. Natural history and prognosis of adenomatous hyperplasia and early hepatocellular carcinoma: multi-institutional analysis of 53 nodules followed up for more than 6 months and 141 patients with single early hepatocellular carcinoma treated by surgical resection or percutaneous ethanol injection. *Jpn J Clin Oncol* 1998; 28: 604–8.
- 10 Okuda K, Ohtsuki T, Obata H *et al.* Natural history of hepatocellular carcinoma and prognosis in relation to

- treatment. Study of 850 patients. *Cancer* 1985; 56 (4): 918–28.
- 11 The Cancer of the Liver Italian Program (CLIP) investigators. A new prognostic system for hepatocellular carcinoma: a retrospective study of 435 patients. *Hepatology* 1998; 28 (3): 751–5.
  - 12 Llovet JM, Burroughs A, Bruix J. Hepatocellular carcinoma. *Lancet* 2003; 362 (9399): 1907–17.
  - 13 Kudo M, Chung H, Osaki Y. Prognostic staging system for hepatocellular carcinoma (CLIP score): its value and limitations, and a proposal for a new staging system, the Japan Integrated Staging Score (JIS score). *J Gastroenterol* 2003; 38: 207–15.
  - 14 Cillo U, Bassanello M, Vitale A et al. The critical issue of hepatocellular carcinoma prognostic classification: which is the best tool available? *J Hepatol* 2004; 40: 124–31.
  - 15 Tateishi R, Yoshida H, Shiina S et al. Proposal of a new prognostic model for hepatocellular carcinoma: an analysis of 403 patients. *Gut* 2005; 54: 419–25.
  - 16 Kudo M, Chung H, Haji S et al. Validation of a new prognostic staging system for hepatocellular carcinoma: the JIS score compared with the CLIP score. *Hepatology* 2004; 40 (6): 1396–405.
  - 17 Chung H, Kudo M, Takahashi S et al. Comparison of three current staging systems for hepatocellular carcinoma: Japan integrated staging score, new Barcelona Clinic Liver Cancer staging classification, and Tokyo score. *J Gastroenterol Hepatol* 2008; 23: 445–52.
  - 18 Ikai I, Takayasu K, Omata M et al. A modified Japan Integrated Stage score for prognostic assessment in patients with hepatocellular carcinoma. *J Gastroenterol* 2006; 41: 884–92.
  - 19 Kitai S, Kudo M, Minami Y et al. Validation of a new prognostic staging system for hepatocellular carcinoma: a comparison of the biomarker-combined Japan integrated staging score, the conventional Japan integrated staging score and the BALAD score. *Oncology* 2008; 75 (Suppl 1): 83–90.
  - 20 Oka H, Kurioka N, Kim K et al. Prospective study of early detection of hepatocellular carcinoma in patients with cirrhosis. *Hepatology* 1990; 12 (4 Pt 1): 680–7.
  - 21 Colombo M, de Franchis R, Del Ninno E et al. Hepatocellular carcinoma in Italian patients with cirrhosis. *N Engl J Med* 1991; 325: 675–80.
  - 22 Pateron D, Ganne N, Trinchet JC et al. Prospective study of screening for hepatocellular carcinoma in Caucasian patients with cirrhosis. *J Hepatol* 1994; 20: 65–71.
  - 23 Zoli M, Magalotti D, Bianchi G, Gueli C, Marchesini G, Pisi E. Efficacy of a surveillance program for early detection of hepatocellular carcinoma. *Cancer* 1996; 78: 977–85.
  - 24 Yuen MF, Cheng CC, Laufer IJ, Lam SK, Ooi CG, Lai CL. Early detection of hepatocellular carcinoma increases the chance of treatment: Hong Kong experience. *Hepatology* 2000; 31: 330–5.
  - 25 Bolondi L, Sofia S, Siringo S et al. Surveillance programme of cirrhotic patients for early diagnosis and treatment of hepatocellular carcinoma: a cost effectiveness analysis. *Gut* 2001; 48: 251–9.
  - 26 Chen TH, Chen CJ, Yen MF et al. Ultrasound screening and risk factors for death from hepatocellular carcinoma in a high risk group in Taiwan. *Int J Cancer* 2002; 98: 257–61.
  - 27 Danta M, Barnes E, Dusheiko G. The surveillance and diagnosis of hepatocellular carcinoma. *Eur J Gastroenterol Hepatol* 2005; 17: 491–6.
  - 28 Ryder SD. Guidelines for the diagnosis and treatment of hepatocellular carcinoma (HCC) in adults. *Gut* 2003; 52 (Suppl 3): iii1–8.
  - 29 Bruix J, Sherman M, Llovet JM et al. Clinical management of hepatocellular carcinoma. Conclusions of the Barcelona-2000 EASL conference. European Association for the Study of the Liver. *J Hepatol* 2001; 35: 421–30.
  - 30 Kokudo N, Makuuchi M. Evidence-based clinical practice guidelines for hepatocellular carcinoma in Japan: the J-HCC guidelines. *J Gastroenterol* 2009; 44 (Suppl 19): 119–21.
  - 31 Thompson Coon J, Rogers G, Hewson P et al. Surveillance of cirrhosis for hepatocellular carcinoma: systematic review and economic analysis. *Health Technol Assess* 2007; 11 (34): 1–206.
  - 32 Makuuchi M for the group formed to establish Guidelines for evidence-based clinical practice for the treatment of liver cancer. Clinical practice guidelines for hepatocellular carcinoma: Tokyo, Kanehara; 2005 (in Japanese).
  - 33 Kew MC, Purves LR, Bersohn I. Serum alpha-fetoprotein levels in acute viral hepatitis. *Gut* 1973; 14: 939–42.
  - 34 Alpert E, Feller ER. Alpha-fetoprotein (AFP) in benign liver disease. Evidence that normal liver regeneration does not induce AFP synthesis. *Gastroenterology* 1978; 74 (5 Pt 1): 856–8.
  - 35 Eleftheriou N, Heathcote J, Thomas HC, Sherlock S. Serum alpha-fetoprotein levels in patients with acute and chronic liver disease. Relation to hepatocellular regeneration and development of primary liver cell carcinoma. *J Clin Pathol* 1977; 30: 704–8.
  - 36 Taketa K. Alpha-fetoprotein: reevaluation in hepatology. *Hepatology* 1990; 12 (6): 1420–32.
  - 37 Mita Y, Aoyagi Y, Yanagi M, Suda T, Suzuki Y, Asakura H. The usefulness of determining des-gamma-carboxy prothrombin by sensitive enzyme immunoassay in the early diagnosis of patients with hepatocellular carcinoma. *Cancer* 1998; 82: 1643–8.
  - 38 Tsai SL, Huang GT, Yang PM, Sheu JC, Sung JL, Chen DS. Plasma des-gamma-carboxyprothrombin in the early stage of hepatocellular carcinoma. *Hepatology* 1990; 11: 481–8.
  - 39 Shimauchi Y, Tanaka M, Kuromatsu R et al. A simultaneous monitoring of Lens culinaris agglutinin A-reactive alpha-fetoprotein and des-gamma-carboxy prothrombin

- as an early diagnosis of hepatocellular carcinoma in the follow-up of cirrhotic patients. *Oncol Rep* 2000; 7: 249–56.
- 40 Nomura F, Ishijima M, Kuwa K, Tanaka N, Nakai T, Ohnishi K. Serum des-gamma-carboxy prothrombin levels determined by a new generation of sensitive immunoassays in patients with small-sized hepatocellular carcinoma. *Am J Gastroenterol* 1999; 94 (3): 650–4.
  - 41 Kawata S, Murakami T, Kim T *et al.* Multidetector CT: diagnostic impact of slice thickness on detection of hypervascular hepatocellular carcinoma. *AJR Am J Roentgenol* 2002; 179: 61–6.
  - 42 Ichikawa T, Erturk SM, Araki T. Multiphasic contrast-enhanced multidetector-row CT of liver: contrast-enhancement theory and practical scan protocol with a combination of fixed injection duration and patients' body-weight-tailored dose of contrast material. *Eur J Radiol* 2006; 58: 165–76.
  - 43 Noguchi Y, Murakami T, Kim T *et al.* Detection of hepatocellular carcinoma: comparison of dynamic MR imaging with dynamic double arterial phase helical CT. *AJR Am J Roentgenol* 2003; 180 (2): 455–60.
  - 44 Kim SH, Lee J, Kim MJ *et al.* Gadoteric acid-enhanced MRI versus triple-phase MDCT for the preoperative detection of hepatocellular carcinoma. *AJR Am J Roentgenol* 2009; 192 (6): 1675–81.
  - 45 Dohr O, Hofmeister R, Treher M, Schweinfurth H. Pre-clinical safety evaluation of Gd-EOB-DTPA (Primovist). *Invest Radiol* 2007; 42: 830–41.
  - 46 Bartolozzi C, Crocetti L, Lencioni R, Cioni D, Della Pina C, Campani D. Biliary and reticuloendothelial impairment in hepatocarcinogenesis: the diagnostic role of tissue-specific MR contrast media. *Eur Radiol* 2007; 17 (10): 2519–30.
  - 47 Di Martino M, Marin D, Guerri A *et al.* Intraindividual comparison of gadoteric acid (Gd-EOB-DTPA) enhanced MR imaging and multiphasic 64-slice CT for the detection of hepatocellular carcinoma (HCC) in patients with cirrhosis. *B-096, RNSA* 2008.
  - 48 Luca A, Grazioli L, Caruso S *et al.* A two-centre study for the comparison of Gd-EOB-DTPA (PRIMOVIST)-enhanced MRI versus triple-phase MDCT for the detection of hepatocellular carcinoma in cirrhosis. *B-097, RNSA* 2008.
  - 49 Sugiura NTK, Ohto M *et al.* Ultrasound image-guided percutaneous intratumor ethanol injection for small hepatocellular carcinoma. *Kanzo* 1983; 24: 920.
  - 50 Livraghi T, Festi D, Monti F, Salmi A, Vettori C. US-guided percutaneous alcohol injection of small hepatic and abdominal tumors. *Radiology* 1986; 161: 309–12.
  - 51 Shiina S, Yasuda H, Muto H *et al.* Percutaneous ethanol injection in the treatment of liver neoplasms. *AJR Am J Roentgenol* 1987; 149: 949–52.
  - 52 Seki T, Wakabayashi M, Nakagawa T *et al.* Percutaneous microwave coagulation therapy for patients with small hepatocellular carcinoma: comparison with percutaneous ethanol injection therapy. *Cancer* 1999; 85 (8): 1694–702.
  - 53 Shiina S, Teratani T, Obi S, Hamamura K, Koike Y, Omata M. Nonsurgical treatment of hepatocellular carcinoma: from percutaneous ethanol injection therapy and percutaneous microwave coagulation therapy to radiofrequency ablation. *Oncology* 2002; 62 (Suppl 1): 64–8.
  - 54 Shiina S, Tagawa K, Unuma T *et al.* Percutaneous ethanol injection therapy for hepatocellular carcinoma. A histopathologic study. *Cancer* 1991; 68: 1524–30.
  - 55 Nakashima Y, Nakashima O, Tanaka M, Okuda K, Nakashima M, Kojiro M. Portal vein invasion and intrahepatic micrometastasis in small hepatocellular carcinoma by gross type. *Hepatol Res* 2003; 26: 142–7.
  - 56 Okusaka T, Okada S, Ueno H *et al.* Satellite lesions in patients with small hepatocellular carcinoma with reference to clinicopathologic features. *Cancer* 2002; 95 (9): 1931–7.
  - 57 Hatanaka K, Chung H, Kudo M *et al.* Usefulness of the post-vascular phase of contrast-enhanced ultrasonography with Sonazoid in the evaluation of gross types of hepatocellular carcinoma. *Oncology* 2010 (in press).
  - 58 Burgener FA, Hamlin DJ. Contrast enhancement of focal hepatic lesions in CT: effect of size and histology. *AJR Am J Roentgenol* 1983; 140: 297–301.
  - 59 Arii S, Yamaoka Y, Futagawa S *et al.* Results of surgical and nonsurgical treatment for small-sized hepatocellular carcinomas: a retrospective and nationwide survey in Japan. The Liver Cancer Study Group of Japan. *Hepatology* 2000; 32 (6): 1224–9.
  - 60 Huang GT, Lee PH, Tsang YM *et al.* Percutaneous ethanol injection versus surgical resection for the treatment of small hepatocellular carcinoma: a prospective study. *Ann Surg* 2005; 242 (1): 36–42.
  - 61 Ryu M, Shimamura Y, Kinoshita T *et al.* Therapeutic results of resection, transcatheter arterial embolization and percutaneous transhepatic ethanol injection in 3225 patients with hepatocellular carcinoma: a retrospective multicenter study. *Jpn J Clin Oncol* 1997; 27: 251–7.
  - 62 Livraghi T, Bolondi L, Buscarini L *et al.* No treatment, resection and ethanol injection in hepatocellular carcinoma: a retrospective analysis of survival in 391 patients with cirrhosis. Italian Cooperative HCC Study Group. *J Hepatol* 1995; 22: 522–6.
  - 63 Castells A, Bruix J, Bru C *et al.* Treatment of small hepatocellular carcinoma in cirrhotic patients: a cohort study comparing surgical resection and percutaneous ethanol injection. *Hepatology* 1993; 18 (5): 1121–6.
  - 64 Shiina S, Teratani T, Obi S *et al.* A randomized controlled trial of radiofrequency ablation with ethanol injection for small hepatocellular carcinoma. *Gastroenterology* 2005; 129: 122–30.
  - 65 Lencioni RA, Allgaier HP, Cioni D *et al.* Small hepatocellular carcinoma in cirrhosis: randomized comparison of

- radio-frequency thermal ablation versus percutaneous ethanol injection. *Radiology* 2003; 228 (1): 235–40.
- 66 Lin SM, Lin CJ, Lin CC, Hsu CW, Chen YC. Radiofrequency ablation improves prognosis compared with ethanol injection for hepatocellular carcinoma < or =4 cm. *Gastroenterology* 2004; 127 (6): 1714–23.
  - 67 Lin SM, Lin CJ, Lin CC, Hsu CW, Chen YC. Randomised controlled trial comparing percutaneous radiofrequency thermal ablation, percutaneous ethanol injection, and percutaneous acetic acid injection to treat hepatocellular carcinoma of 3 cm or less. *Gut* 2005; 54: 1151–6.
  - 68 Chen MS, Li JQ, Zheng Y *et al.* A prospective randomized trial comparing percutaneous local ablative therapy and partial hepatectomy for small hepatocellular carcinoma. *Ann Surg* 2006; 243: 321–8.
  - 69 Llovet JM, Di Bisceglie AM, Bruix J *et al.* Design and endpoints of clinical trials in hepatocellular carcinoma. *J Natl Cancer Inst* 2008; 100: 698–711.
  - 70 The Japanese Liver Transplantation Society. Liver transplantation in Japan in 2006 (part 2)-Registry by the Japanese Liver Transplantation Society. *Ishoku* 2008; 43: 45–55, (in Japanese).
  - 71 Mazzaferro V, Regalia E, Doci R *et al.* Liver transplantation for the treatment of small hepatocellular carcinomas in patients with cirrhosis. *N Engl J Med* 1996; 334: 693–9.
  - 72 Iwatsuki S, Starzl TE, Sheahan DG *et al.* Hepatic resection versus transplantation for hepatocellular carcinoma. *Ann Surg* 1991; 214: 221–8, discussion 228–229.
  - 73 Penn I. Hepatic transplantation for primary and metastatic cancers of the liver. *Surgery* 1991; 110: 726–34, discussion 734–725.
  - 74 Todo S, Furukawa H, Tada M. Extending indication: role of living donor liver transplantation for hepatocellular carcinoma. *Liver Transpl* 2007; 13 (11 Suppl 2): S48–54.
  - 75 Mazzaferro V, Llovet JM, Miceli R *et al.* Predicting survival after liver transplantation in patients with hepatocellular carcinoma beyond the Milan criteria: a retrospective, exploratory analysis. *Lancet Oncol* 2009; 10 (1): 35–43.
  - 76 Yao FY, Ferrell L, Bass NM *et al.* Liver transplantation for hepatocellular carcinoma: expansion of the tumor size limits does not adversely impact survival. *Hepatology* 2001; 33 (6): 1394–403.
  - 77 Sugawara Y, Tamura S, Makuuchi M. Living donor liver transplantation for hepatocellular carcinoma: Tokyo University series. *Dig Dis* 2007; 25: 310–12.
  - 78 Lee SG, Hwang S, Moon DB *et al.* Expanded indication criteria of living donor liver transplantation for hepatocellular carcinoma at one large-volume center. *Liver Transpl* 2008; 14: 935–45.
  - 79 Ito T, Takada Y, Ueda M *et al.* Expansion of selection criteria for patients with hepatocellular carcinoma in living donor liver transplantation. *Liver Transpl* 2007; 13: 1637–44.
  - 80 Duffy JP, Vardanian A, Benjamin E *et al.* Liver transplantation criteria for hepatocellular carcinoma should be expanded: a 22-year experience with 467 patients at UCLA. *Ann Surg* 2007; 246: 502–9, discussion 509–511.
  - 81 Yamamoto J, Iwatsuki S, Kosuge T *et al.* Should hepatomas be treated with hepatic resection or transplantation? *Cancer* 1999; 86: 1151–8.
  - 82 Figueras J, Jaurrieta E, Valls C *et al.* Resection or transplantation for hepatocellular carcinoma in cirrhotic patients: outcomes based on indicated treatment strategy. *J Am Coll Surg* 2000; 190: 580–7.
  - 83 Llovet JM, Fuster J, Bruix J. Intention-to-treat analysis of surgical treatment for early hepatocellular carcinoma: resection versus transplantation. *Hepatology* 1999; 30 (6): 1434–40.
  - 84 Llovet JM, Real MI, Montana X *et al.* Arterial embolisation or chemoembolisation versus symptomatic treatment in patients with unresectable hepatocellular carcinoma: a randomised controlled trial. *Lancet* 2002; 359 (9319): 1734–9.
  - 85 Lo CM, Ngan H, Tso WK *et al.* Randomized controlled trial of transarterial lipiodol chemoembolization for unresectable hepatocellular carcinoma. *Hepatology* 2002; 35 (5): 1164–71.
  - 86 Llovet JM, Bruix J. Systematic review of randomized trials for unresectable hepatocellular carcinoma: chemoembolization improves survival. *Hepatology* 2003; 37 (2): 429–42.
  - 87 Bruix J, Sala M, Llovet JM. Chemoembolization for hepatocellular carcinoma. *Gastroenterology* 2004; 127 (5 Suppl 1): S179–88.
  - 88 Makuuchi M, Kokudo N. Clinical practice guidelines for hepatocellular carcinoma: the first evidence based guidelines from Japan. *World J Gastroenterol* 2006; 12: 828–9.
  - 89 Kudo M, Okano T. Management of hepatocellular carcinoma in Japan: consensus-based clinical practice manual proposed by the Japan Society of Hepatology. *Oncology* 2007; 72: S2–15.
  - 90 Ikai I, Aii S, Okazaki M *et al.* Report of the 17th Nationwide Follow-up Survey of Primary Liver Cancer in Japan. *Hepatol Res* 2007; 37: 676–91.
  - 91 Takayasu K, Aii S, Ikai I *et al.* Prospective cohort study of transarterial chemoembolization for unresectable hepatocellular carcinoma in 8510 patients. *Gastroenterology* 2006; 131 (2): 461–9.
  - 92 Ueno S, Tanabe G, Nuruki K *et al.* Prognostic performance of the new classification of primary liver cancer of Japan (4th edition) for patients with hepatocellular carcinoma: a validation analysis. *Hepatol Res* 2002; 24: 395–403.
  - 93 Marelli L, Stigliano R, Triantos C *et al.* Treatment outcomes for hepatocellular carcinoma using chemoembolization in combination with other therapies. *Cancer Treat Rev* 2006; 32: 594–606.
  - 94 Veltri A, Moretto P, Doriguzzi A, Pagano E, Carrara G, Gandini G. Radiofrequency thermal ablation (RFA) after

- transarterial chemoembolization (TACE) as a combined therapy for unresectable non-early hepatocellular carcinoma (HCC). *Eur Radiol* 2006; 16 (3): 661–9.
- 95 Buscarini L, Buscarini E, Di Stasi M, Quaretti P, Zangrandi A. Percutaneous radiofrequency thermal ablation combined with transcatheter arterial embolization in the treatment of large hepatocellular carcinoma. *Ultraschall Med* 1999; 20 (2): 47–53.
  - 96 Yamakado K, Nakatsuka A, Akeboshi M, Shiraki K, Nakano T, Takeda K. Combination therapy with radiofrequency ablation and transcatheter chemoembolization for the treatment of hepatocellular carcinoma: short-term recurrences and survival. *Oncol Rep* 2004; 11: 105–9.
  - 97 Koda M, Ueki M, Maeda Y *et al.* The influence on liver parenchymal function and complications of radiofrequency ablation or the combination with transcatheter arterial embolization for hepatocellular carcinoma. *Hepatol Res* 2004; 29 (1): 18–23.
  - 98 Obi S, Yoshida H, Toune R *et al.* Combination therapy of intraarterial 5-fluorouracil and systemic interferon-alpha for advanced hepatocellular carcinoma with portal venous invasion. *Cancer* 2006; 106 (9): 1990–7.
  - 99 Ando E, Tanaka M, Yamashita F *et al.* Hepatic arterial infusion chemotherapy for advanced hepatocellular carcinoma with portal vein tumor thrombosis: analysis of 48 cases. *Cancer* 2002; 95: 588–95.
  - 100 Lin CP, Yu HC, Cheng JS *et al.* Clinical effects of intra-arterial infusion chemotherapy with cisplatin, mitomycin C, leucovorin and 5-fluorouracil for unresectable advanced hepatocellular carcinoma. *J Chin Med Assoc* 2004; 67: 602–10.
  - 101 Sumie S, Yamashita F, Ando E *et al.* Interventional radiology for advanced hepatocellular carcinoma: comparison of hepatic artery infusion chemotherapy and transcatheter arterial lipiodol chemoembolization. *AJR Am J Roentgenol* 2003; 181 (5): 1327–34.
  - 102 Tzoracoleftherakis EE, Spiliotis JD, Kyriakopoulou T, Kakkos SK. Intra-arterial versus systemic chemotherapy for non-operable hepatocellular carcinoma. *Hepatogastroenterology* 1999; 46 (26): 1122–5.
  - 103 Court WS, Order SE, Siegel JA *et al.* Remission and survival following monthly intraarterial cisplatin in non-resectable hepatoma. *Cancer Invest* 2002; 20 (5–6): 613–25.
  - 104 Okuda K, Tanaka M, Shibata J *et al.* Hepatic arterial infusion chemotherapy with continuous low dose administration of cisplatin and 5-fluorouracil for multiple recurrence of hepatocellular carcinoma after surgical treatment. *Oncol Rep* 1999; 6: 587–91.
  - 105 Tanioka H, Tsuji A, Morita S *et al.* Combination chemotherapy with continuous 5-fluorouracil and low-dose cisplatin infusion for advanced hepatocellular carcinoma. *Anticancer Res* 2003; 23 (2C): 1891–7.
  - 106 Park JY, Ahn SH, Yoon YJ *et al.* Repetitive short-course hepatic arterial infusion chemotherapy with high-dose 5-fluorouracil and cisplatin in patients with advanced hepatocellular carcinoma. *Cancer* 2007; 110: 129–37.
  - 107 Chung YH, Song IH, Song BC *et al.* Combined therapy consisting of intraarterial cisplatin infusion and systemic interferon-alpha for hepatocellular carcinoma patients with major portal vein thrombosis or distant metastasis. *Cancer* 2000; 88 (9): 1986–91.
  - 108 Kaneko S, Urabe T, Kobayashi K. Combination chemotherapy for advanced hepatocellular carcinoma complicated by major portal vein thrombosis. *Oncology* 2002; 62 (Suppl 1): 69–73.
  - 109 Llovet JM, Ricci S, Mazzaferro V *et al.* Sorafenib in advanced hepatocellular carcinoma. *N Engl J Med* 2008; 359: 378–90.
  - 110 Cheng AL, Kang YK, Chen Z *et al.* Efficacy and safety of sorafenib in patients in the Asia-Pacific region with advanced hepatocellular carcinoma: a phase III randomised, double-blind, placebo-controlled trial. *Lancet Oncol* 2009; 10 (1): 25–34.
  - 111 Kudo M. Hepatocellular carcinoma 2009 and beyond: from the surveillance to molecular targeted therapy. *Oncology* 2008; 75: S1–12.
  - 112 Kudo M. The 2008 Okuda lecture: management of hepatocellular carcinoma: from surveillance to molecular targeted therapy. *J Gastroenterol Hepatol* 2010; 25 (3): 439–52.



## Survival Benefit of Hepatopancreatoduodenectomy for Cholangiocarcinoma in Comparison to Hepatectomy or Pancreatoduodenectomy

Yuji Kaneoka · Akihiro Yamaguchi ·  
Masatoshi Isogai · Takashi Kumada

Published online: 7 July 2010  
© Société Internationale de Chirurgie 2010

### Abstract

**Background** Perihilar and distal cholangiocarcinoma remain difficult to treat, and long-term survival is poor. We conducted a retrospective study of patients with cholangiocarcinoma to examine whether hepatopancreatoduodenectomy, in comparison to standard surgeries, provides a survival benefit.

**Methods** Subjects were 75 patients with perihilar or distal cholangiocarcinoma who, between April 1997 and May 2007, underwent hepatectomy with bile duct resection (Hx,  $n = 29$ ), pancreatoduodenectomy (PD,  $n = 32$ ), or hepatopancreatoduodenectomy (HPD,  $n = 14$ ) at our hospital. We compared surgical outcomes and survival between groups and identified factors negatively influencing survival.

**Results** Morbidity and in-hospital mortality did not differ significantly between groups (Hx group, 34% and 10%, respectively; PD group, 44% and 3%; and HPD, 57% and 0%). The overall median survival time was 39 months, and overall 5-year survival (including in-hospital mortality) was 42%. Respective group values were as follows: Hx, 24 months and 31%; PD, 51 months and 49%, and HPD, 63 months and 50%. Although the number of patients was small, survival in the HPD was not influenced by the type of invasion whether widespread intramural invasion ( $n = 8$ ), superficial spread ( $n = 4$ ), or hepatoduodenal ligament invasion ( $n = 2$ ). Multivariate analysis (Cox proportional hazards model) showed only perineural invasion ( $p = .007$ )

and decreased curability (R1/2 resection) ( $p = .017$ ) to be independent risk factors influencing survival.

**Conclusions** In cases of perihilar or distal cholangiocarcinoma, aggressive surgery must be aimed at overcoming perineural invasion. Our findings indicate that HPD improves survival of patients undergoing surgery for widespread cholangiocarcinoma in comparison to standard surgeries.

### Introduction

Typically, perihilar cholangiocarcinoma is treated surgically by means of hepatectomy with bile duct resection (Hx). This treatment has been reported by some institutions to yield satisfactory survival rates with high curative resection rates [1–4]. In contrast, distal cholangiocarcinoma, i.e., carcinoma of the middle and lower extrahepatic bile duct, has long been treated by pancreatoduodenectomy (PD) [5, 6]. Only in some institutions has hepatopancreatoduodenectomy (HPD) been offered as a surgical option for selected patients with cholangiocarcinoma showing widespread infiltration [7–9]. In 1998 we began performing HPD at our institution for bile duct carcinoma characterized by widespread intramural invasion. To obtain curative resection in cases of perihilar and distal cholangiocarcinoma, we now choose among the following three procedures as standard surgical therapy: Hx, PD, or HPD. Use of HPD, a challenging operation introduced in 1980 [7], has increased in Japan over the past two decades, but worldwide it has not been established as a standard operation for cholangiocarcinoma. Safety of the procedure, the type of disease for which it is indicated, and long-term survival have not been adequately analyzed. Therefore we conducted a retrospective study at a single institution to compare surgical outcomes and survival between HPD and

Y. Kaneoka (✉) · A. Yamaguchi · M. Isogai  
Department of Surgery, Ogaki Municipal Hospital,  
4-86 Minaminokawa-cho, Ogaki 503-8502, Japan  
e-mail: y-kaneoka@omh.ogaki.gifu.jp

T. Kumada  
Department of Gastroenterology, Ogaki Municipal Hospital,  
4-86 Minaminokawa-cho, Ogaki 503-8502, Japan

the other two surgical resection procedures for perihilar and distal cholangiocarcinoma.

### Patients and methods

Between April 1997 and May 2007, 124 patients diagnosed with perihilar or distal cholangiocarcinoma were admitted to Ogaki Municipal Hospital. Forty-nine of these patients were considered poor surgical candidates because of advanced age, distant metastases, or impaired liver function and were thus not treated surgically; 4 of the 49 patients underwent exploratory laparotomy alone, and 3 underwent non-curative bile duct resection alone after intraoperative identification of non-resectable distant metastases. The remaining 75 patients, who comprised our study group, underwent radical surgery (resection rate: 60%). The presenting symptoms were obstructive jaundice ( $n = 57$ ; 76%), abdominal pain ( $n = 10$ ; 13.3%), abnormal laboratory values ( $n = 7$ ; 9.3%), and weight loss ( $n = 1$ ; 1.3%). Twenty-nine of the 75 patients underwent Hx, 32 underwent PD, and 14 underwent HPD. Clinical results of these three surgical procedures, including survival and prognostic factors, were analyzed.

### Selection between types of surgery

Tumor extension into the biliary tree was assessed by percutaneous transhepatic cholangiography, endoscopic retrograde cholangiography, or magnetic resonance cholangiopancreatography when necessary. In early cases of perihilar cholangiocarcinoma with obstructive jaundice, percutaneous transhepatic biliary drainage (PTBD) was performed to facilitate precise evaluation of the intrahepatic biliary system. More recently, endoscopic nasobiliary drainage has often been used to prevent spillage of the cancer cell-containing bile. We now consider evaluation of periductal infiltration, vascular involvement, and regional lymph node enlargement by abdominal multidetector-row computed tomography (CT) to be the most reliable means of deciding on the appropriate operative procedure and the extent of resection.

The Hx procedure was selected for patients with perihilar cholangiocarcinoma (extrahepatic cholangiocarcinoma located in the hepatic hilum—so-called Klatskin tumor,  $n = 17$ ; and intrahepatic cholangiocarcinoma invading the hepatic hilum,  $n = 12$ ). Depending on the predominant location of the tumor, right or left hepatectomy with total caudate lobectomy was routinely performed. However, 2 patients underwent central bisegmentectomy with total caudate lobectomy for impaired liver function, whereas 1 patient underwent medial (segment 4) segmentectomy with total caudate lobectomy for a small (limited) Klatskin tumor.

Pancreatoduodenectomy was performed for distal cholangiocarcinoma that did not extend into the hepatic hilum and did not give the appearance of superficial spread, either by preoperative cholangiography or cholangioscopy.

The HPD procedure was performed for tumor that could not be radically resected by Hx or PD alone, especially when the proximal extent of the residual hepatic duct was restricted by the tumor within the first order of hepatic duct (right or left hepatic trunk). Therefore, right or left trisectionectomy was not carried out in patients with HPD. This included 12 cases of extrahepatic cholangiocarcinoma widely invading the proximal or distal bile duct and 2 cases of intrahepatic cholangiocarcinoma invading the hepatic hilum. The types of invasion were: *widespread intramural invasion*, i.e., longitudinal stricture of the extrahepatic bile duct reaching both the hepatic hilum and the lower (pancreatic head) bile duct ( $n = 8$ ); *superficial spread*, i.e., papillary carcinoma arising from a bile duct and spreading to the mucosa ( $n = 4$ ); and *hepatoduodenal ligament invasion*, i.e., advanced cholangiocarcinoma invading the hepatic hilum and main vessels of the hepatoduodenal ligament, requiring hepato-ligamentopancreatoduodenectomy (HLPD) [10] ( $n = 2$ ) to remove microscopic cancer cell infiltration.

Preoperative portal vein embolization was performed according to the ipsilateral percutaneous transhepatic method described by Nagino et al. [11] to enhance remnant liver function for 8 of 13 patients who were scheduled to undergo right (extended) hepatectomy by Hx and for all 6 patients who were scheduled to undergo right hepatectomy by HPD. Our institutional criteria for portal vein embolization were right-sided hepatectomy with the future remnant liver volume less than 40% in Hx and without regard for potential remnant liver volume in HPD. The primary surgeries were performed more than 2 weeks after embolization.

### HPD procedure


After making a midline incision with a right subcostal extension, we first looked for the absence of macroscopic para-aortic lymph node metastases by the Kocher maneuver. Pancreatoduodenectomy was begun upon dissection of the duodenum (pylorus-preserving PD) or stomach; the pancreas was divided at the left side of the portal vein. The jejunum was then divided 10 cm distally from the ligament of Treitz. The uncinate process of the pancreas was dissected along the superior mesenteric vein, and the right-sided superior mesenteric plexus was resected simultaneously. After skeletonization of the hepatoduodenal ligament, hepatic transection was performed with an ultrasonic surgical aspirator under intermittent total hepatic inflow vascular occlusion (15-min clamp time, 3-min declamp time, and so on). The alimentary tract was

reconstructed after stent placement for hepaticojejunostomy and pancreaticojejunostomy. A drain was placed in the right subphrenic space, beneath the hepaticojejunostomy, and in the cranial portion of the pancreaticojejunostomy. These drains were routinely removed within 1 week after surgery, but all stents were left in place for 3 weeks.


Segmental portal vein resection and reconstruction (PVR) was performed immediately after tumor removal. In cases of right hepatectomy, however, PVR followed division of the left hepatic ducts with a small part of the medial hepatic segment because PVR hastened residual hepatic transection by facilitating a good view of the surgical field (Fig. 1). When graft reconstruction of the portal vein was required, the right external iliac vein was harvested extraperitoneally through a right groin incision. For the 2 cases requiring HLPD, left hepatectomy with total caudate lobectomy was performed. In these 2 cases, skeletonization of the hepatoduodenal ligament was unnecessary; the common hepatic and right hepatic arteries were divided, and the right hepatic artery reconstruction (HAR) was performed by using the middle colic artery, which was mobilized with the middle colic vein. The portal vein was divided between the confluence of the splenic vein and the right portal branch and then reconstructed with an external iliac vein graft (Fig. 2).

#### Study definitions

Because of the typically invasive spread of cholangiocarcinoma, tumor location (perihilar versus distal) was determined according to the depth of invasion and the degree of bile duct stricture. The International Union against Cancer (UICC) staging system [12] for extrahepatic bile duct carcinoma was applied to all patients, allowing us to treat intrahepatic cholangiocarcinoma invading the hepatic hilum (14 cases) as perihilar cholangiocarcinoma [13].



**Fig. 1** Right hepatectomy with total caudate lobectomy, pylorus-preserving pancreatoduodenectomy, and portal vein resection and reconstruction (PVR) were performed (*right oblique view*). Arrow suture line of the portal vein anastomosis, B bile duct stump, P pancreatic stump



**Fig. 2** Left hepatectomy with total caudate lobectomy, pancreatoduodenectomy, and portal vein grafting (PVG; 4 cm in length) were performed (*left oblique view*). Arrow anastomosis of the right hepatic artery and the mobilized middle colic artery, MHV middle hepatic vein, G graft, PT pancreatic stent tube

Liver failure was defined as a maximum postoperative serum total bilirubin level  $>5$  mg/dl. Pancreatic fistula was defined as pancreatic drainage that persisted for more than 3 weeks after resection, i.e., grade B pancreatic fistula (according to the International Study Group of Pancreatic Fistula criteria [14]). Delayed gastric emptying (DGE) was defined as a patient's inability to tolerate oral intake by postoperative day (POD) 7, i.e., grade A DGE (according to the International Study Group of Pancreatic Surgery criteria [15]). Perineural invasion was defined as microscopic evidence of cancer cell infiltration into the perineural space to an intermediate degree or more (according to the *Japanese General Rules for Surgical and Pathological Studies on Cancer of the Biliary Tract* [16]). Arterial invasion was defined as the presence of a macroscopic mass around the artery and/or microscopic observation of crowded cancer cells in the perineural space beside the artery. Portal vein invasion was defined as microscopic invasion into the adventitia or intima of the portal vein.

#### Preoperative and postoperative therapy

Adjuvant chemoradiation therapy was not performed in any of the study patients. Recurrence was identified in 6 patients who were subsequently treated with 5-fluorouracil or gemcitabine. The hepatic hilum was irradiated in 2 patients. Because the number of patients was small, data pertaining specifically to chemotherapy or radiation therapy were not entered into the analysis.

#### Data acquisition

By accessing patient records, we were able to obtain the following clinical data: patient age, sex, tumor location, whether right or left hepatectomy was performed, whether

**Table 1** Profiles of the patients

	Hx ( <i>n</i> = 29)	PD ( <i>n</i> = 32)	HPD ( <i>n</i> = 14)	<i>p</i> value*
Age (range)	65 ± 2 (43–81)	67 ± 2 (48–81)	62 ± 2 (50–78)	.18
Sex, M/F	18/11	23/9	10/4	.68
Predominant location of tumor				
Perihilar	29	0	9	
Distal	0	32	5	
Operative procedure				
RH	13 (2)		6	
LH	13 (2)		7	
CBS	2			
Seg	1		1	
PPPD		25	11	
PD		7	3	
PVR	13 (4)	4 (1)	5 (2)	
HAR	6		2	
Operative time, min (range)	429 ± 21 (247–655)	275 ± 22 (131–585)	550 ± 30 (370–1,025)	<.0001
Blood loss, ml (range)	1,096 ± 107 (390–2,497)	863 ± 115 (230–2,470)	1,354 ± 153 (754–2,900)	.041

The figures in parentheses indicate the number of graft reconstructions

*RH* right hepatectomy (two right trisectionectomy), *LH* left hepatectomy (two left trisectionectomy), *CBS* central bisegmentectomy, *Seg* segmentectomy, *PPPD* pylorus-preserving pancreatoduodenectomy, *PD* pancreatoduodenectomy, *PVR* (segmental) portal vein reconstruction, *HAR* hepatic artery reconstruction

\* *p* values compare variables among the three surgical methods

PVR or HAR was performed, HAR method, operative time, total blood loss, histopathologic features of the resected tumor, morbidity, follow-up time, site of recurrence, and survival time.

#### Statistics

Continuous variables are expressed as mean ± standard error of the mean. Differences between groups were analyzed by chi-square or Mann–Whitney test, as appropriate. Survival was plotted by the Kaplan–Meier method and analyzed by log-rank test (univariate analysis) to identify potential prognostic factors. Multivariate analysis was carried out by means of a Cox proportional hazards model to identify independent risk factors influencing survival. Statistical significance was accepted at *p* < .05. Data analysis was performed with SPSS, version 17.0 (SPSS Japan Inc., Tokyo, Japan).

## Results

#### Clinical variables

Mean age of the 75 study patients was 65 ± 1 years (range: 43–81 years); the male/female ratio was 51/24 (Table 1). Both perihilar (*n* = 9) and widespread distal (*n* = 5) cholangiocarcinomas were treated by HPD. The numbers of right and left hepatectomies were evenly distributed in the Hx and HPD groups, whereas for pancreatic resection, pylorus-preserving pancreaticoduodenectomy

(PPPD) was most prevalent procedure, according to our institutional policy. Portal vein reconstruction was performed for macroscopic cancer invasion in 13 patients during Hx, in 4 patients during PD, and in 5 patients during HPD, for an overall PVR rate of 29%. HAR was required for 6 patients who underwent Hx and 2 patients who underwent HPD, for a total HAR rate of 11%. Six patients underwent both PVR and HAR. Regarding arterial reconstruction, simple end-to-end arterial reconstruction was performed for 1 patient during Hx, whereas others required mobilization of another artery (either the gastroduodenal or middle colic artery) for anastomosis to the future remnant hepatic artery.

As expected, operative time and blood loss differed between groups, with PD resulting in the shortest operative time and least blood loss.

#### Follow-up time

Mean follow-up for all patients was 46 ± 4 months. There were 4 in-hospital deaths (in-hospital mortality, 5.3%). Thirty-nine patients died from cancer relapse, and 3 died from other diseases. The remaining 29 patients were alive as of December 31, 2009. The minimum follow-up period for any patient was 31 months.

#### Postoperative complications

Morbidity and mortality did not differ statistically between groups (Table 2). The overall incidence of morbidity was lower in the Hx group than in the other two groups, but