

- 31 Yu J, Qiao L, Zimmermann L *et al.* Troglitazone inhibits tumor growth in hepatocellular carcinoma *in vitro* and *in vivo*. *Hepatology* 2006; 43: 134–43.
- 32 Marchesini G, Bianchi G, Merli M *et al.* Nutritional supplementation with branched-chain amino acids in advanced cirrhosis: a double-blind, randomized trial. *Gastroenterology* 2003; 124: 1792–801.
- 33 Muto Y, Sato S, Watanabe A *et al.* Effects of oral branched-chain amino acid granules on event-free survival in patients with liver cirrhosis. *Clin Gastroenterol Hepatol* 2005; 3: 705–13.
- 34 Nishitani S, Takehana K, Fujitani S, Sonaka I. Branched-chain amino acids improve glucose metabolism in rats with liver cirrhosis. *Am J Physiol Gastrointest Liver Physiol* 2005; 288: G1292–300.
- 35 Kawaguchi T, Nagao Y, Matsuoka H, Ide T, Sata M. Branched-chain amino acid-enriched supplementation improves insulin resistance in patients with chronic liver disease. *Int J Mol Med* 2008; 22: 105–12.
- 36 Urata Y, Okita K, Korenaga K, Uchida K, Yamasaki T, Sakaida I. The effect of supplementation with branched-chain amino acids in patients with liver cirrhosis. *Hepatology Res* 2007; 37: 510–16.
- 37 Muto Y, Sato S, Watanabe A *et al.* Overweight and obesity increase the risk for liver cancer in patients with liver cirrhosis and long-term oral supplementation with branched-chain amino acid granules inhibits liver carcinogenesis in heavier patients with liver cirrhosis. *Hepatology Res* 2006; 35: 204–14.
- 38 Yoshiji H, Noguchi R, Kitade M *et al.* Branched-chain amino acids suppress insulin-resistance-based hepatocarcinogenesis in obese diabetic rats. *J Gastroenterol* 2009; 44: 483–91.
- 39 Shimizu M, Shirakami Y, Iwasa J *et al.* Supplementation with branched-chain amino acids inhibits azoxymethane-induced colonic preneoplastic lesions in male C57BL/KsJ-db/db mice. *Clin Cancer Res* 2009; 15: 3068–75.

