

such as occludin and claudin, and among them, claudin is a key structural and functional component of TJ seals (Furuse and Tsukita, 2006). Claudin, a tetra-transmembrane protein, comprises a family consisting of more than 20 members. The expression profiles and sealing functions of claudins differ among tissues. Claudin expression is often deregulated in human cancers such as breast, prostate, ovarian, gastric, and pancreatic cancers (Morin, 2005; Kominsky, 2006). There is a relationship between dysregulated claudin and metastasis (Agarwal et al., 2005; Dhawan et al., 2005). These findings indicate that claudin can be a potent target for cancer therapy.

Clostridium perfringens enterotoxin (CPE), a 35-kDa polypeptide, causes food poisoning in humans. CPE binds to its receptor, and then causes changes in the membrane permeability by complex formation on the plasma membrane followed by the induction of oncosis and apoptosis (McClane and Chakrabarti, 2004). The local administration of CPE suppresses solid tumor growth (Michl et al., 2001; Kominsky et al., 2004; Santin et al., 2005); however, whole CPE had never been applied into a ligand for claudin CPE receptors because of its strong cytotoxicity. The receptor-binding region of CPE (C-CPE) can be used for claudin-targeted cancer therapy (Saeki et al., 2009). Immunotoxins, consisting of a protein toxin connected to a binding ligand, such as an antibody or growth factor, have been developed and used for clinical therapy. Protein synthesis inhibitory factor (PSIF) derived from *Pseudomonas* exotoxin is a widely used protein toxin (Kreitman and Pastan, 2006), and intratumoral administration of the claudin-4-targeting PSIF has been shown to attenuate solid tumor growth (Saeki et al., 2009). In the present study, we investigated whether claudin CPE receptors can be a target for tumor metastasis by using the claudin-4-targeting PSIF as a claudin-targeting agent.

Materials and Methods

Cell Culture. Mouse melanoma cell line B16-BL6 and mouse breast cancer cell line 4T1 were cultured in modified Eagle's medium supplemented with 10% fetal bovine serum (FBS) and Dulbecco's modified Eagle's medium (DMEM) containing 10% FBS and 10 mmol/ml HEPES, respectively. The cells were maintained in a 5% CO₂ atmosphere at 37°C.

Preparation of B16 Cells Stably Expressing Claudin-4. Mouse claudin-4 cDNA was subcloned into pcDNA3.1 plasmid coding a neomycin resistance gene. The claudin-4 expression vector was transfected into B16 cells, and B16 cells stably expressing claudin-4 (CL4-B16 cells) were isolated by genetic selection.

Immunoblot Analysis. Cells were lysed in lysis buffer [50 mM Tris (pH 7.4), 8.25 mg/ml NaCl, 1% NP-40, 2 mM SDS, and protease inhibitor cocktail (Sigma-Aldrich, St. Louis, MO)]. The cell lysates were subjected to SDS-polyacrylamide gel electrophoresis. The separated proteins were transferred onto a polyvinylidene difluoride membrane, followed by immunoblotting with anti-claudin-4 (Zymed Laboratories, South San Francisco, CA) or anti- β -actin Ab (Sigma-Aldrich). After incubation with a peroxidase-labeled secondary antibody (Millipore Bioscience Research Reagents, Temecula, CA), the immunoreactive band was visualized by chemiluminescence reagents (GE Healthcare, Little Chalfont, Buckinghamshire, UK).

Cell Proliferation Assay. B16 or CL4-B16-expressing cells (CL4-B16 cells) (2×10^4 cells) were seeded into a 24-well plate. At the indicated time points, the cells were stained with trypan blue, and the number of viable cells was counted by using a hemocytometer under a microscope.

In Vitro Metastasis Analysis. A cell culture insert with an 8- μ m pore size membrane (BD Biosciences Discovery Labware, Bedford,

MA) was used for the invasion assay. The upper surface of the chamber was coated with 50 μ l of 0.1 mg/ml Matrigel (BD Biosciences, San Jose, CA), and the lower chamber was filled with DMEM containing 10% FBS. Cells were cultured to approximately 80% confluence and serum-starved overnight (0.5% FBS). Then, cells (1×10^5 cells) suspended in DMEM with 0.5% FBS were placed into the upper chamber and incubated at 37°C for 24 h. After incubation, noninvading cells were removed, and the membrane was stained with Diff-Quick reagent. The stained cells in five randomly selected fields were counted under a microscope.

Preparation of C-CPE-PSIF. The C-terminal fragment of CPE (C-CPE)-fused PSIF (C-CPE-PSIF) was prepared as described previously (Saeki et al., 2009). In brief, pET-C-CPE-PSIF was transfected into *Escherichia coli* BL21 (DE3) (Novagen, Darmstadt, Germany), and the production of C-CPE-PSIF was induced by the addition of 0.25 mM isopropyl-D-thiogalactopyranoside. The cells were harvested and then lysed in buffer A [10 mM Tris-HCl (pH 8.0), 400 mM NaCl, 5 mM MgCl₂, 0.1 mM phenylmethylsulfonyl fluoride, 1 mM 2-mercaptoethanol, and 10% glycerol]. The lysates were centrifuged, and the resultant supernatant was applied to HiTrap Cheating HP (GE Healthcare). The proteins were eluted by imidazole in buffer A. The buffer was exchanged with phosphate-buffered saline (PBS) by using a PD-10 column (GE Healthcare), and the purified protein was stored at -80°C until use. Protein was quantified by using a BCA protein assay kit (Pierce Chemical, Rockford, IL) with BSA as a standard.

In Vitro Cytotoxic Analysis. In the cytotoxic assay, B16 and CL4-B16 cells were seeded onto a 96-well culture dish at 1×10^4 cells per well. After 24 h, the cells were treated with C-CPE-PSIF for 24 h at the indicated concentrations. The cytotoxicity was determined by a WST-8 kit, according to the manufacturer's instructions (Nacalai Tesque, Kyoto, Japan).

In Vivo Antitumor Activity. Female BALB/c mice (7–8 weeks old) and C57/BL6 mice (7–8 weeks old) were obtained from Shimizu Laboratory Supplies Co., Ltd. (Kyoto, Japan). The mice were housed in an environmentally controlled room at $23 \pm 1.5^\circ\text{C}$ with a 12-h light/dark cycle. The animal experiments were performed according to the guidelines of Osaka University.

For an experimental metastasis model, B16 or CL4-B16 cells (5×10^5 or 1×10^6 cells) in 100 μ l of PBS were injected intravenously into the tails of C57/BL6 mice. Then, mice received intravenously administered PBS or C-CPE-PSIF three times per week. Two weeks after the inoculation of the cells, the mice were sacrificed, and the number of lung metastasis colonies was counted under a microscope. For antitumor activity in B16 cells, 1×10^5 B16 or CL4-B16 cells were injected subcutaneously into the right flanks of C57BL/6 mice. Vehicle or C-CPE-PSIF was intravenously administered three times per week. Calipers were used to measure the minimal and maximal tumor diameters, and the tumor volume was calculated as $a \times b \times b/2$, where a represents the minimal tumor diameter, and b represents the maximal tumor diameter. The body weights of mice were also monitored.

For antitumor activity in 4T1 cells, 4T1 cells (1×10^5 cells) in 50 μ l of PBS were injected subcutaneously into the right flanks of BALB/c mice. PBS, C-CPE-PSIF, or adriamycin (ADR) was intravenously administered, and the tumor size and body weight of mice were monitored. Mice were sacrificed 35 days after tumor inoculation, and lung metastasis was evaluated by staining with India ink.

Statistical Analysis. Data were analyzed by using Dunnett's method. The statistical significance for all comparisons was set at $p < 0.05$.

Results

Preparation of Claudin-4-Expressing B16 BL6 Cells.

To investigate the effects of C-CPE-PSIF on tumor metastasis, we selected murine B16 cells, which have a high propensity

to metastasize to the lung (Saiki, 1997). Western blotting analysis revealed no expression of claudin-4 in B16 cells (Fig. 1A). B16 cells were not sensitive to C-CPE-PSIF (Fig. 2A). We transfected claudin-4 cDNA into B16 cells and established stable CL4-B16 cells (Fig. 1A). The proliferation rate was not affected by exogenously expressed claudin-4 (Fig. 1B). Metastasis has multiple processes, including motility and invasion (Steeg, 2006). To investigate the invasion of CL4-B16 cells, we performed a Boyden chamber migration assay. Cells were seeded onto the cell culture insert of Matrigel-coated membrane with an 8- μ m pore size, and the cells that invaded the apical membrane and reached the basal membrane were counted. As shown in Fig. 1C, the invasion activity was increased 17-fold in CL4-B16 cells compared with that in parental B16 cells. Lung metastasis of CL4-B16 cells was observed when intravenously injected into mice; however, the number of lung metastasis colonies of CL4-B16 cells was smaller than that of the parental B16 cells (Fig. 1D). These findings indicate that CL4-B16 cells can be used as a metastasis model of claudin-4-expressing cancer cells. We discuss the elevation of migration activity and lower lung metastasis in CL4-B16 cells under *Discussion*.

Antitumor Activity of C-CPE-PSIF in CL4-B16 Cells.

Before *in vivo* experiments, we investigated the *in vitro* cytotoxicity of C-CPE-PSIF in CL4-B16 cells. As shown in Fig. 2A, C-CPE-PSIF showed dose-dependent cytotoxicity in CL4-B16 cells, decreasing their viability to 35% at 100 ng/ml. In contrast, parental B16 cells were not sensitive to C-CPE-PSIF even at 1 μ g/ml, indicating that C-CPE-PSIF may target claudin-4. Claudin-4 is expressed in the intestines, liver, and kidney (Morita et al., 1999). To determine a safe dose of C-CPE-PSIF, we checked serum biochemical markers of liver (alanine aminotransferase) and kidney (blood urea nitrogen) injury in mice injected with C-CPE-PSIF. After intravenous administration of C-CPE-PSIF (5 μ g/kg), the mice showed no signs of injury (data not shown). In the following *in vivo* experiments, the doses of C-CPE-PSIF were less than or equal to 5 μ g/kg. B16 or CL4-B16 cells were intravenously injected into mice, and then C-CPE-PSIF was intravenously administered every 2 days. Two weeks after the tumor cell injection, the number of lung metastasis colonies was counted. As shown in Fig. 2B, C-CPE-PSIF treatment decreased the number of lung metastasis colonies from 39 ± 17 to 10 ± 4 at 5 μ g/kg. In contrast, C-CPE-PSIF treatment did not affect the lung metastasis of B16 cells (Fig. 2B). C-CPE-PSIF suppressed the growth of CL4-B16 cells but not B16 cells inoculated into the right flank of mice (Fig. 2C). These data suggest that claudin-4 targeting may be a potent strategy for suppressing tumor metastasis and growth.

Suppression of the Primary Tumors and Metastasis of 4T1 Cells. To clarify the potency of a claudin-4-targeting strategy, we investigated the effect of C-CPE-PSIF on a spontaneous metastasis model. 4T1 cells are murine cancer cells that spontaneously metastasize to the lung after being subcutaneously inoculated (Wong et al., 2002). We investigated whether C-CPE-PSIF suppresses the spontaneous lung metastasis of 4T1 cells. On day 33, the tumor volume was 1801.2 ± 848.5 mm³ in the vehicle-treated group and 740.5 ± 94.6 mm³ in the group treated with 5 μ g/kg C-CPE-PSIF (Fig. 3A). The number of lung metastasis colonies was decreased to 2 ± 1 colonies at 5 μ g/kg C-CPE-PSIF (Fig. 3B). A dose of 2 μ g/kg C-CPE-PSIF did not suppress tumor growth




Fig. 1. Claudin-4-expressing B16 melanoma (CL4-B16) cells. A, preparation of CL4-B16 cells. Cell lysates from B16 and CL4-B16 cells were subjected to SDS-polyacrylamide gel electrophoresis, followed by Western blotting with claudin-4 and β -actin. β -Actin is a control for an endogenous protein. B, the effect of claudin-4 on cellular proliferation in B16 cells. B16 or CL4-B16 cells (2×10^4 cells) were seeded onto a 24-well plate. Then, the cell numbers were counted by trypan blue dye exclusion assay at the indicated periods. Data are shown as means \pm S.D. ($n = 4$). C, effect of claudin-4 on invasion in B16 cells. B16 or CL4-B16 cells (1×10^5 cells) were seeded into the upper well of the cell culture insert coated with Matrigel. After 24 h, the cells that invaded the bottom membrane of the insert were stained with DiffQuick reagent and counted under a microscope. Data are shown as means \pm S.D. ($n = 4$). *, significantly different from B16 cells ($p < 0.01$). D, lung metastasis of CL4-B16 cells. B16 or CL4-B16 cells (5×10^5 or 1×10^6 cells) were injected into the tail veins of mice. After 14 days, the mice were sacrificed, the lungs were fixed, and the colonies on the lung surface were counted. Data are shown as means \pm S.D. ($n = 5$). *, significantly different from B16 cells ($p < 0.01$).




Fig. 2. Antitumor activity of C-CPE-PSIF for CL4-B16 cells. A, cytotoxicity of C-CPE-PSIF in CL4-B16 cells. B16 cells (top) and CL4-B16 cells (bottom) were treated with PSIF or C-CPE-PSIF at the indicated concentrations for 24 h. Cell viability (%) was measured by a WST-8 kit, according to the manufacturer's instructions (Nacalai Tesque). Data represent the mean \pm S.D. ($n = 3$). *, significantly different from the vehicle-treated group ($p < 0.05$). B, anti-metastatic activity of C-CPE-PSIF on lung metastasis of B16 (top) or CL4-B16 (bottom) cells. B16 or CL4-B16 cells (1×10^6 cells) were injected into the tail veins of mice on day 0, and vehicle or C-CPE-PSIF (2 or 5 $\mu\text{g}/\text{kg}$) was intravenously injected on days 0, 2, 4, 7, 9, 11, and 13. On day 14, the mice were sacrificed, their lungs were fixed, and the colonies on the lung surface were counted. Data are shown as means \pm S.D. ($n = 5$). *, significantly different from the vehicle-treated group ($p < 0.05$). C, antitumor activity of C-CPE-PSIF on CL4-B16 subcutaneously inoculated allograft. B16 (top) or CL4-B16 (bottom) cells (1×10^5 cells) were intradermally inoculated into the right flank of mice on day 0, and PBS or C-CPE-PSIF (2 or 5 $\mu\text{g}/\text{kg}$) was intravenously injected three times a week. Tumor volume was monitored. Each point is the mean \pm S.D. ($n = 5$). The data are representative of two independent experiments.

but did prevent lung metastasis. The circulating tumor cells might be more sensitive to C-CPE-PSIF than tumor cells in the solid tumor tissue. C-CPE-PSIF treatments did not cause a decrease in body weight (Fig. 3C), and there were no apparent biochemical side effects (Supplemental Fig. 1). ADR, which is frequently used in clinical chemotherapy, suppressed the tumor growth from 970.3 ± 278.4 to $458.6 \pm 51.4 \text{ mm}^3$ at 4 mg/kg (Fig. 3D). As shown in Fig. 3E, 4 mg/kg ADR decreased the number of lung metastasis colonies (24 ± 13 colonies in the vehicle-treated group; 6 ± 4 colonies in the ADR-treated group). However, the ADR-treated mice experienced a 26% loss of body weight, which is a sign of side effects (Fig. 3F). Thus, the antitumor activity of C-CPE-PSIF might

be more potent than that of ADR. These results indicate that claudin-4-targeting therapy might be a potent strategy for tumor therapy with a low level of side effects and a high level of antitumor activity.

Discussion

Most malignant tumors are derived from the epithelium, and metastasis is the major cause of death from cancers. In the present study, we found that systemic administration of a claudin-targeting molecule suppressed cancer metastasis, indicating that claudin targeting might be an effective therapy against cancer metastasis.




Fig. 3. Antitumor activity of C-CPE-PSIF for murine breast cancer 4T1 cells. 4T1 cells (1×10^5 cells) were intradermally inoculated into the right flanks of mice on day 0, and C-CPE-PSIF (A–C) or ADR (D–F) was intravenously injected three or two times a week at the indicated doses, respectively. Tumor volume (A and D) and body weight (B and E) were monitored. On day 35, the mice were sacrificed, their lungs were stained with India ink, and the number of spontaneous metastases was determined (C and F). Data are shown as means \pm S.D. ($n = 5$). *, significantly different from the vehicle-treated group ($p < 0.01$). The data are representative of two independent experiments.

Although the *in vitro* metastasis activity of CL4-B16 cells was higher than that of parental B16 cells, the *in vivo* lung metastasis of CL4-B16 cells was lower than that of B16 cells. As shown in Supplemental Fig. 2, claudin-4 increased the invasiveness and migration activity of B16 cells *in vitro* and decreased lung metastasis *in vivo*. A possible explanation for this discrepancy might be the experimental model, which evaluates a different stage of metastasis. The migration and invasion activity involved in the early stage of metastasis was investigated in the *in vitro* analysis, whereas extravasation and colonization to an organ involved in the late stage of metastasis was evaluated by the *in vivo* experiment. The altered expression of claudin-4 changed the metastasis of 4T1 cells to the heart and liver, suggesting that claudin affects organ-specific metastasis (Erin et al., 2009). Claudin-4 might suppress the lung metastasis of B16 cells.

Claudin is a structural and functional component of TJs (Furuse and Tsukita, 2006). What is the role of claudin in metastasis? Metastasis is composed of three steps: leaving the primary site, entering the blood flow, and invading the distant site. In the first step, the combination of claudin members in the TJ strands may be important. The claudin family contains at least 24 members. Claudin is believed to form homo- and hetero-type claudin strands, and the pattern

of the strands differs among tissues and determines the properties of TJ seals (Furuse and Tsukita, 2006). For example, rigid TJ seals were formed when claudin-11 or claudin-15 was exogenously expressed in Madin-Darby canine kidney cells, whereas the expression of claudin-11 or claudin-15 reduced the TJ integrity in LLC-PK1 cells by its dominant negative effects on TJ sealing (Van Itallie et al., 2003). Dominant negative effects of claudin-4 on the TJ barrier might contribute to an acceleration in the detachment of cancer cells from the primary tumor tissue. In the second and third steps of metastasis, cancer cells must move through the extracellular matrix at the primary site and the distant site. Cancer cells must degrade the extracellular matrix by the expression of matrix metalloproteinase and increase their motility. Claudin expression enhanced invasion with increased matrix metalloproteinase activity (Agarwal et al., 2005). There is a relationship between the levels of claudin-1/claudin-4 and the metastasis of human cancers, including hepatic, colonic, ovarian, and gastric cancers (Miwa et al., 2001; Agarwal et al., 2005; Resnick et al., 2005; Halder et al., 2008; Lee et al., 2008). The overexpression of claudin suppressed cancer metastasis in human pancreatic and gastric cancers (Michl et al., 2003; Mima et al., 2005; Ohtani et al., 2009). Claudin-4 suppressed or accelerated *in vitro* and *in*

AQ: D

AQ: E

vivo metastasis of human cancer cells (Agarwal et al., 2005; Ohtani et al., 2009). Cell-cell interaction through TJs regulates cell growth signaling (Matter et al., 2005). Taken together, these findings indicate that claudin family members might control several steps of cancer metastasis. The precise molecular mechanism and role of claudin in cancer metastasis remain to be determined.

Whether a claudin-4-targeting method causes severe side effects is critical for its clinical application in cancer therapy. Claudins play pivotal roles in TJ barrier and fence functions by maintaining cellular polarity in normal epithelium (Furuse and Tsukita, 2006). Claudins are believed to be more accessible in tumors than in normal epithelium. Claudins form TJ seals in lateral membranes between adjacent cells in normal epithelium, whereas claudins are exposed on the cell surface during tumorigenesis (Soler et al., 1999; Kominsky, 2006). Indeed, no local or systemic side effects have been observed after the intratumoral administration of CPE (Kominsky et al., 2007; Santin et al., 2007). Here, we also found that the systemic administration of C-CPE-PSIF causes no significant increase in biochemical markers (aspartate aminotransferase, alanine aminotransferase, and blood urea nitrogen) for toxicity at a therapeutic dose of 5 $\mu\text{g}/\text{kg}$ (Supplemental Fig. 2). Thus, a claudin-targeting strategy might have weak side effects.

It is difficult to prepare recombinant claudin protein because of its hydrophobic property, and claudin has low antigenicity. Until recently, an antibody against the extracellular loop domain of claudin had never been successfully prepared, and C-CPE was the only known claudin binder. Recently, Romani et al. (2009) prepared a single-chain antibody fragment against claudin-3 by using phage display technology. They found that the antibody fragment binds to ovarian and uterine carcinoma cells in vitro. More importantly, a therapeutic monoclonal antibody against claudin-4 was developed. Suzuki et al. (2009) successfully prepared anti-claudin-4 antibody by immunizing claudin-4-expressing tumor cells into a mouse with autoimmune disease. The antibody mediates antibody-dependent cellular cytotoxicity and both in vitro and in vivo antitumor activity. Although the preparation of anti-claudin antibody may lead to a breakthrough in cancer therapy, the immunogenicity associated with immunotoxin clinical therapies is a future problem (Kreitman and Pastan, 2006). The C-terminal 30 amino acids are the minimum functional domain of C-CPE to bind to claudin-4 (Hanna et al., 1991). The C-terminal 30-amino-acid fragment was used to deliver a cytokine to claudin-4-expressing cells by genetic fusion (Yuan et al., 2009). Humanized antibody and the claudin-4-targeting peptide may be useful for cancer therapy in the near future.

In summary, this is the first report to indicate that systemic injection of a claudin-targeting molecule suppresses tumor growth and metastasis. Hematologic cells do not develop TJs; therefore, a claudin-targeting therapy may have no hematologic toxicity. We anticipate that claudin targeting will be a potent strategy for cancer therapy.

Acknowledgments

We thank members of our laboratory for useful comments and discussion and Drs. Y. Horiguchi (Osaka University, Osaka, Japan), S. Tsunoda (National Institute of Biomedical Innovation, Osaka, Japan), and M. Furuse (Kobe University, Hyogo, Japan) for providing C-CPE cDNA, PSIF cDNA, and claudin cDNA, respectively.

References

- Agarwal R, D'Souza T, and Morin PJ (2005) Claudin-3 and claudin-4 expression in ovarian epithelial cells enhances invasion and is associated with increased matrix metalloproteinase-2 activity. *Cancer Res* **65**:7378–7385.
- Dhawani P, Singh AB, Deane NG, No Y, Shiou SR, Schmidt C, Neff J, Washington MK, and Beauchamp RD (2005) Claudin-1 regulates cellular transformation and metastatic behavior in colon cancer. *J Clin Invest* **115**:1765–1776.
- Erin N, Wang N, Xin P, Bui V, Weisz J, Barkan GA, Zhao W, Shearer D, and Clawson GA (2009) Altered gene expression in breast cancer liver metastases. *Int J Cancer* **124**:1503–1516.
- Furuse M and Tsukita S (2006) Claudins in occluding junctions of humans and flies. *Trends Cell Biol* **16**:181–188.
- Gupta GP and Massagué J (2006) Cancer metastasis: building a framework. *Cell* **127**:679–695.
- Halder SK, Rachakonda G, Deane NG, and Datta PK (2008) Smad7 induces hepatic metastasis in colorectal cancer. *Br J Cancer* **99**:957–965.
- Hanna PC, Mietzner TA, Schoolnik GK, and McClane BA (1991) Localization of the receptor-binding region of *Clostridium perfringens* enterotoxin using cloned toxin fragments and synthetic peptides. The 30 C-terminal amino acids define a functional binding region. *J Biol Chem* **266**:11037–11043.
- Jemal A, Siegel R, Ward E, Hao Y, Xu J, Murray T, and Thun MJ (2008) Cancer statistics, 2008. *CA Cancer J Clin* **58**:71–96.
- Kominsky SL (2006) Claudins: emerging targets for cancer therapy. *Expert Rev Mol Med* **8**:1–11.
- Kominsky SL, Tyler B, Sosnowski J, Brady K, Doucet M, Nell D, Smedley JG 3rd, McClane B, Brem H, and Sukumar S (2007) *Clostridium perfringens* enterotoxin as a novel-targeted therapeutic for brain metastasis. *Cancer Res* **67**:7977–7982.
- Kominsky SL, Vali M, Korz D, Gabig TG, Weitzman SA, Argani P, and Sukumar S (2004) *Clostridium perfringens* enterotoxin elicits rapid and specific cytotoxicity of breast carcinoma cells mediated through tight junction proteins claudin 3 and 4. *Am J Pathol* **164**:1627–1633.
- Kreitman RJ and Pastan I (2006) Immunotoxins in the treatment of hematologic malignancies. *Curr Drug Targets* **7**:1301–1311.
- Lee LY, Wu CM, Wang CC, Yu JS, Liang Y, Huang KH, Lo CH, and Hwang TL (2008) Expression of matrix metalloproteinases MMP-2 and MMP-9 in gastric cancer and their relation to claudin-4 expression. *Histol Histopathol* **23**:515–521.
- Martin TA and Jiang WG (2001) Tight junctions and their role in cancer metastasis. *Histol Histopathol* **16**:1183–1195.
- Matter K, Aijaz S, Tsapara A, and Balda MS (2005) Mammalian tight junctions in the regulation of epithelial differentiation and proliferation. *Curr Opin Cell Biol* **17**:453–458.
- McClane BA and Chakrabarti G (2004) New insights into the cytotoxic mechanisms of *Clostridium perfringens* enterotoxin. *Anaerobe* **10**:107–114.
- Michl P, Barth C, Buchholz M, Lerch MM, Rolke M, Holzmann KH, Menke A, Fensterer H, Giehl K, Lohr M, et al. (2003) Claudin-4 expression decreases invasiveness and metastatic potential of pancreatic cancer. *Cancer Res* **63**:6265–6271.
- Michl P, Buchholz M, Rolke M, Kunsch S, Lohr M, McClane B, Tsukita S, Leder G, Adler G, and Gress TM (2001) Claudin-4: a new target for pancreatic cancer treatment using *Clostridium perfringens* enterotoxin. *Gastroenterology* **121**:678–684.
- Mima S, Tsutsumi S, Ushijima H, Takeda M, Fukuda I, Yokomizo K, Suzuki K, Sano K, Nakanishi T, Tomisato W, et al. (2005) Induction of claudin-4 by nonsteroidal anti-inflammatory drugs and its contribution to their chemopreventive effect. *Cancer Res* **65**:1868–1876.
- Mitic LL and Anderson JM (1998) Molecular architecture of tight junctions. *Annu Rev Physiol* **60**:121–142.
- Morin PJ (2005) Claudin proteins in human cancer: promising new targets for diagnosis and therapy. *Cancer Res* **65**:9603–9606.
- Morita K, Furuse M, Fujimoto K, and Tsukita S (1999) Claudin multigene family encoding four-transmembrane domain protein components of tight junction strands. *Proc Natl Acad Sci USA* **96**:511–516.
- Mullin JM (1997) Potential interplay between luminal growth factors and increased tight junction permeability in epithelial carcinogenesis. *J Exp Zool* **279**:484–489.
- Ohtani S, Terashima M, Satoh J, Soeta N, Saze Z, Kashimura S, Ohsuka F, Hoshino Y, Kogure M, and Gotoh M (2009) Expression of tight junction-associated proteins in human gastric cancer: down-regulation of claudin-4 correlates with tumor aggressiveness and survival. *Gastric Cancer* **12**:43–51.
- Resnick MB, Konkin T, Routhier J, Sabo E, and Pricolo VE (2005) Claudin-1 is a strong prognostic indicator in stage II colonic cancer: a tissue microarray study. *Mod Pathol* **18**:511–518.
- Romani C, Comper F, Bandiera E, Ravaggi A, Bignotti E, Tassi RA, Pecorelli S, and Santin AD (2009) Development and characterization of a human single-chain antibody fragment against claudin-3: a novel therapeutic target in ovarian and uterine carcinomas. *Am J Obstet Gynecol* **201**:70,e71–e79.
- Saeki R, Kondoh M, Kakutani H, Tsunoda S, Mochizuki Y, Hamakubo T, Tsutsumi Y, Horiguchi Y, and Yagi K (2009) A novel tumor-targeted therapy using a claudin-4-targeting molecule. *Mol Pharmacol* **76**:918–926.
- Saiki I (1997) Cell adhesion molecules and cancer metastasis. *Jpn J Pharmacol* **75**:215–242.
- Santin AD, Bellone S, Marizzoni M, Palmieri M, Siegel ER, McKenney JK, Hennings L, Comper F, Bandiera E, and Pecorelli S (2007) Overexpression of claudin-3 and claudin-4 receptors in uterine serous papillary carcinoma: novel targets for a type-specific therapy using *Clostridium perfringens* enterotoxin (CPE). *Cancer* **109**:1312–1322.
- Santin AD, Cané S, Bellone S, Palmieri M, Siegel ER, Thomas M, Roman JJ, Burnett A, Cannon MJ, and Pecorelli S (2005) Treatment of chemotherapy-resistant human ovarian cancer xenografts in C.B-17/SCID mice by intraperitoneal administration of *Clostridium perfringens* enterotoxin. *Cancer Res* **65**:4334–4342.
- Soler AP, Miller RD, Laughlin KV, Carp NZ, Klurfeld DM, and Mullin JM (1999) Increased tight junctional permeability is associated with the development of colon cancer. *Carcinogenesis* **20**:1425–1431.

- Q: G** Sonoda N, Furuse M, Sasaki H, Yonemura S, Katahira J, Horiguchi Y, and Tsukita S (1999) *Clostridium perfringens* enterotoxin fragment removes specific claudins from tight junction strands: Evidence for direct involvement of claudins in tight junction barrier. *J Cell Biol* **147**:195–204.
- Stahelin LA (1973) Further observations on the fine structure of freeze-cleaved tight junctions. *J Cell Sci* **13**:763–786.
- Steeg PS (2006) Tumor metastasis: mechanistic insights and clinical challenges. *Nat Med* **12**:895–904.
- Suzuki M, Kato-Nakano M, Kawamoto S, Furuya A, Abe Y, Misaka H, Kimoto N, Nakamura K, Ohta S, and Ando H (2009) Therapeutic antitumor efficacy of monoclonal antibody against claudin-4 for pancreatic and ovarian cancers. *Cancer Sci* **100**:1623–1630.
- Van Itallie CM, Fanning AS, and Anderson JM (2003) Reversal of charge selectivity in cation or anion-selective epithelial lines by expression of different claudins. *Am J Physiol Renal Physiol* **285**:F1078–F1084.
- Vermeer PD, Einwalter LA, Moninger TO, Rokhlina T, Kern JA, Zabner J, and Welsh MJ (2003) Segregation of receptor and ligand regulates activation of epithelial growth factor receptor. *Nature* **422**:322–326.
- Wodarz A and Näthke I (2007) Cell polarity in development and cancer. *Nat Cell Biol* **9**:1016–1024.
- Wong CW, Song C, Grimes MM, Fu W, Dewhirst MW, Muschel RJ, and Al-Mehdi AB (2002) Intravascular location of breast cancer cells after spontaneous metastasis to the lung. *Am J Pathol* **161**:749–753.
- Yuan X, Lin X, Manorek G, Kanatani I, Cheung LH, Rosenblum MG, and Howell SB (2009) Recombinant CPE fused to tumor necrosis factor targets human ovarian cancer cells expressing the claudin-3 and claudin-4 receptors. *Mol Cancer Ther* **8**:1906–1915.

Address correspondence to: Dr. Masuo Kondoh, Laboratory of Bio-Functional Molecular Chemistry, Graduate School of Pharmaceutical Sciences, Osaka University, Suita, Osaka 565-0871, Japan. E-mail: masuo@phs.osaka-u.ac.jp

AQ: H

NOT FOR



DISTRIBUTION

A Novel Screening System for Claudin Binder Using Baculoviral Display

Hideki Kakutani¹*, Azusa Takahashi¹*, Masuo Kondoh^{1*}, Yumiko Saito¹, Toshiaki Yamaura¹, Toshiko Sakihama², Takao Hamakubo², Kiyohito Yagi^{1*}

¹ Laboratory of Bio-Functional Molecular Chemistry, Graduate School of Pharmaceutical Sciences, Osaka University, Suita, Osaka, Japan, ² Department of Molecular Biology and Medicine, Research Center for Advanced Science and Technology, The University of Tokyo, Meguro, Tokyo, Japan

Abstract

Recent progress in cell biology has provided new insight into the claudin (CL) family of integral membrane proteins, which contains more than 20 members, as a target for pharmaceutical therapy. Few ligands for CL have been identified because it is difficult to prepare CL in an intact form. In the present study, we developed a method to screen for CL binders by using the budded baculovirus (BV) display system. CL4-displaying BV interacted with a CL4 binder, the C-terminal fragment of *Clostridium perfringens* enterotoxin (C-CPE), but it did not interact with C-CPE that was mutated in its CL4-binding region. C-CPE did not interact with BV and CL1-displaying BV. We used CL4-displaying BV to select CL4-binding phage in a mixture of a scFv-phage and C-CPE-phage. The percentage of C-CPE-phage in the phage mixture increased from 16.7% before selection to 92% after selection, indicating that CL-displaying BV may be useful for the selection of CL binders. We prepared a C-CPE phage library by mutating the functional amino acids. We screened the library for CL4 binders by affinity to CL4-displaying BV, and we found that the novel CL4 binders modulated the tight-junction barrier. These findings indicate that the CL-displaying BV system may be a promising method to produce a novel CL binder and modulator.

Citation: Kakutani H, Takahashi A, Kondoh M, Saito Y, Yamaura T, et al. (2011) A Novel Screening System for Claudin Binder Using Baculoviral Display. *PLoS ONE* 6(2): e16611. doi:10.1371/journal.pone.0016611

Editor: Vladimir Uversky, University of South Florida College of Medicine, United States of America

Received: November 22, 2010; **Accepted:** December 24, 2010; **Published:** February 14, 2011

Copyright: © 2011 Kakutani et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by a Grant-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology, Japan (21689006), by a Health and Labor Sciences Research Grant from the Ministry of Health, Labor and Welfare of Japan, by Takeda Science Foundation, by a Suzuken Memorial Foundation, by a grant from Kansai Biomedical Cluster project in Saito, which is promoted by the Knowledge Cluster Initiative of the Ministry of Education, Culture, Sports, Science and Technology, Japan and by a Research Grant for Promoting Technological Seeds from Japan Science and Technology Agency. A.T. is supported by Research Fellowships of the Japan Society for the Promotion of Science for Young Scientists. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: masuo@phs.osaka-u.ac.jp (MK); yagi@phs.osaka-u.ac.jp (KY)

☯ These authors contributed equally to this work.

Introduction

Tight junctions (TJ) are intercellular adhesion complexes in epithelial and endothelial cells; TJs are located in the most apical part of the complexes [1]. TJs have a barrier function and a fence function [2–4]. TJs contribute to epithelial and endothelial barrier functions by restricting the diffusion of solutes through the paracellular pathway. TJs maintain cellular polarity by preventing the free movement of membrane proteins between the apical and basal membranes [5]. Loss of cell-cell adhesion and cellular polarity commonly occurs in the early stages of cancer [6]. Modulation of the TJ barrier function can be a method to enhance drug absorption, and TJ components exposed on the surface of cancer cells can be a target for cancer therapy.

Biochemical analyses of TJs have identified TJ components, such as occludin, claudins (CLs) and junction adhesion molecule [7]. The CL family contains more than 20 integral tetra-transmembrane proteins that play pivotal roles in the TJ barrier and fence functions. CL1-deficient mice lack the epidermal barrier, while CL5-deficient mice lack the blood-brain barrier [8,9], indicating that the regulation of the TJ barrier by modulation of CLs may be a promising method for drug delivery. *Clostridium perfringens* enterotoxin (CPE) causes food poisoning in

humans [10]. An interaction between the C-terminal domain of CPE (C-CPE) with CL4 deregulates the TJ barrier [11,12]. We previously found that C-CPE enhances jejunal absorption through its interaction with CL4, indicating that a CL binder is a potent drug-delivery system [13].

The majority of lethal cancers are derived from epithelial tissues [14]. Malignant tumor cells frequently exhibit abnormal TJ function, followed by the deregulation of cellular polarity and intercellular contact, which is commonly observed in both advanced tumors and the early stages of carcinogenesis [6]. Some CLs are overexpressed in various types of cancers. For example, CL3 and CL4 are overexpressed in breast, prostate, ovarian, pancreatic and gastric cancers. CL1, CL7, CL10 and CL16 are overexpressed in colon, gastric, thyroid and ovarian cancers, respectively [15,16]. These findings indicate that the CLs may be a target molecule for cancer therapy. A receptor for CPE is CL4 [11,12]. CPE has anti-tumor activity against human pancreatic and ovarian cancers without side effects [17,18]. The CLs binders will be useful for cancer-targeting therapy.

As above, recent investigations of CLs provide new insight into their use as pharmaceutical agents; for example, a CL binder may be used in drug delivery and anti-tumor therapy. Selection of a CL binder by using a recombinant CL protein is a putative method to

prepare a CL binder. However, CLs are four-transmembrane proteins with high hydrophobicity; there has been little success in the preparation of intact CL protein. Recently, a novel type of protein expression system that uses baculovirus has been developed. Membrane proteins are displayed on the budded baculovirus (BV) in their active form [19–21], indicating that the BV system may be useful for the preparation of a CL binder. In the present study, we investigated whether a CL binder was screened by using a CL-displaying BV.

Results

Preparation of CL4-displaying BV

C-CPE is the only known CL binder and modulator [12,13,22]. C-CPE has affinity to CL4 in a nanomolar range [23]. We chose C-CPE and CL4 as models of the CL binder and CL, respectively. Several reports indicate that membrane proteins expressed on the surface of BV are in an intact form [19–21]. To check the expression of CL4 on the BV, we performed immunoblot analysis of the lysate of CL4-BV against CL4. As shown in Fig. 1A, CL4 was detected in the virus lysates. To determine if the CL4 expressed on the virus has an intact form, we performed enzyme-linked immunosorbent assay (ELISA) with CL4-BV-coated immunoplates. C-CPE binds to the extracellular loop domain of CL4 [23]. After the addition of C-CPE to the CL4-BV-coated plate, the C-CPE bound to the CL4-BV-coated plate was detected by anti-his-tag antibody, followed by incubation with horseradish peroxidase-labeled antibody. C-CPE was dose-dependently bound to CL4-BV, whereas C-CPE did not interact with wild-BV (Fig. 1B). Deletion of the CL4-binding region (C-CPE303) attenuated the interaction of C-CPE with CL4-BV (Fig. 1C). Together, these results indicate that the CL4 displayed on BV may have an intact extracellular loop region.

Selection of C-CPE-phage by using CL4-BV

We next examined the interaction between C-CPE-phage and CL4-BV. As shown in Fig. 2A, C-CPE-phage bound to CL4-BV but not to wild-BV, and a scFv-phage did not bind to CL4-BV. To determine if CL-BV can be used to select CL binders, we prepared a mixture of C-CPE-phage and scFv-phage at a ratio of 2:10 and used CL4-BV to select CL4-binding phage in the mixtures. The amount of C-CPE-phage was increased to 11 of 12 clones in the mixture (Fig. 2B), indicating that CL-BV may be useful in the preparation of CL binders.

We previously found that each substitution of S304, S305, S307, N309, S313 and K318 with alanine increased the binding of C-CPE to CL4 [24]. Here, we prepared a phage library for C-CPE by randomly changing the functional 6 amino acids to any of the 20 amino acids. To confirm the diversity of the library, we checked the sequences of 17 randomly isolated clones. Each of the 17 clones had a different sequence, indicating that the library has a diverse population of C-CPE mutants (Table 1).

Then, we screened the CL4-binding phage by their affinity to CL4-BV. After addition of the C-CPE library to CL4-BV-adsorbed tubes, the CL4-BV-bound phages were recovered (1st screening). We repeated this screening process two more times (2nd screening and 3rd screening). If the number of CL4-bound phage is increased during the screening, the ratio of the incubated phage titers to the recovered phage titers will increase. As shown in Fig. 3A, the ratio was increased during screening from 4.5×10^{-7} to 5.5×10^{-5} , indicating that the screening system for CL4 binders may work. Indeed, the number of monoclonal phage clones with high affinity to CL4-BV was increased after the 3rd screening compared with that after the 2nd screening (Fig. 3B).




Figure 1. Preparation of CL4-displaying BV. A) Immunoblot analysis. Wild-BV and CL4-BV (0.1 µg/lane) were subjected to SDS-PAGE, followed by immunoblot analysis with anti-CL4 antibody. The lysate of CL4-expressing L (CL4/L) cells was used as a positive control. B, C) Interaction of a CL4 binder with CL4-BV. Immunotubes were coated with the wild-BV or CL4-BV, and C-CPE (B) or mutated C-CPE (C) was added to the BV-coated immunotubes at the indicated concentration. C-CPE bound to the BV-coated tubes was detected by ELISA with an anti-his-tag antibody. doi:10.1371/journal.pone.0016611.g001

We analyzed the sequences of the CL4-BV-bound phages and got novel CL4-binder candidates with amino acid sequences that differed from the wild-type sequence (Table 2). To investigate their CL4-binding, we prepared the recombinant proteins of the binders and investigated their interaction with CL4 by ELISA with CL-BVs. As shown in Fig. 4A, the novel C-CPE derivatives had affinity to CL4 but not CL1. Next, we investigated whether the novel CL4 binders modulate TJ barrier in Caco-2 monolayer cell sheets, a popular model for the evaluation of TJ barriers [25].




Figure 2. Selection of C-CPE-displaying phage by using the CL4-BV system. A) Interaction of C-CPE-displaying phage with CL4-BV. Wild-BV or CL4-BV was coated on an immunoplate, and then scFv-displaying phage or C-CPE-displaying phage was added to the BV-coated immunoplate at the indicated concentrations. The BV-bound phages were detected by ELISA with anti-M13 antibody as described in Materials and methods. Data are representative of two independent experiments. Data are means \pm SD ($n=3$). B) Enrichment of C-CPE-displaying phage by the BV system. A mixture of scFv-phage and C-CPE-phage (mixing ratio of scFv-phage to C-CPE-phage=2:10) was incubated with a CL4-BV-coated immunotube, and the bound phages were recovered. Each phage clone was identified by PCR amplification, followed by agarose gel electrophoresis. Upper and lower pictures are before and after the selection, respectively. The putative sizes of the PCR products are 856 and 523 bp in scFv and C-CPE, respectively. The data are representative of two independent experiments. doi:10.1371/journal.pone.0016611.g002

Treatment of the cells with C-CPE resulted in decreased transepithelial electrical resistance (TEER) values, a marker of TJ integrity, and the TEER values increased after removal of C-CPE. The C-CPE derivatives (clones 1–5) had TJ-modulating activity similar to that of C-CPE (Fig. 4B).

Discussion

CL is a promising target for pharmaceutical therapy. However, CL has low antigenicity, and there has been little success in the preparation of monoclonal antibody against the extracellular loop region of CL. The three-dimensional structure of CL has never been determined, so it is impossible to perform a theoretical design

Table 1. C-CPE phage library.

	304	305	307	309	313	318
C-CPE	S	S	S	N	S	K
Clone 1	V	T	C	V	N	K
2	C	P	A	H	L	T
3	A	G	G	V	P	P
4	R	G	H	L	E	H
5	A	A	P	S	R	Q
6	P	A	P	D	P	A
7	C	T	T	T	N	K
8	H	P	S	P	G	H
9	R	G	G	R	N	R
10	A	P	S	T	Q	P
11	V	L	G	N	M	R
12	P	P	A	T	F	R
13	G	D	C	S	N	L
14	F	R	V	F	R	N
15	S	Q	Q	W	T	T
16	S	R	L	E	W	Q
17	K	R	E	R	Q	S

Phage clones were randomly picked up from the C-CPE phage library, and the amino acids sequences of C-CPE mutant were analyzed. doi:10.1371/journal.pone.0016611.t001

of a CL binder based on the structural information. In the present study, we developed a novel screening system for CL binders by using a BV system and a C-CPE phage display library, and we used this system to identify novel CL4 binders.

In ligand screening, the preparation of a receptor for the ligand is very critical. Membrane proteins are especially difficult to prepare as recombinant protein with an intact structure. Functional membrane proteins such as cell-surface proteins are heterologously expressed on BV in their native forms [19–21]. Interactions between membrane proteins can be detected by using receptor-displaying and ligand-displaying BV [21]. In the present report, we found that CL4-BV interacts with a CL4 binder, C-CPE, but it does not interact with C-CPE303 that lacks the CL4-binding residues of C-CPE. The CL4-binding site of C-CPE corresponds to that of CPE; so, the second extracellular loop of CL appears to be the C-CPE-binding site [23,26]. These findings indicate that CL4 displayed on BV may have native form. We anticipate that CL-BV will be useful for the preparation of CL binders, such as peptides and antibodies.

To the best of our knowledge, the preparation of CL binder has been performed by only four groups. Offner et al. prepared polyclonal antibodies against extracellular domains of CL3 and CL4 [27], Ling et al. screened peptide types of CL4 binder by using a 12-mer peptide phage display library and CL4-expressing cells [28], Suzuki et al. generated a monoclonal antibody against the second extracellular loop of CL4 from mice immunized with a human pancreatic cancer cell line [29] and Romani et al. screened scFv against CL3 by using a human antibody phage display library [30]. However, the CL modulators have never been developed; thus, C-CPE is the only known CL4 modulator [12]. In the present study, we prepared a C-CPE phage library containing C-CPE mutants in which each of the 6 functional amino acids was randomly replaced with an amino acid, and we isolated CL4 binders by using CL4-BV as a screening ligand. Interestingly, all of




Figure 3. Screening of a novel CL4 binder. A) Enrichment of phages with affinity to CL4-BV. CL4-BVs coated on immunotubes were incubated with the C-CPE-derivative phage library at 1.6×10^{12} CFU titer (1st input phage). The phages bound to CL4-BV were recovered (1st output phage). The CL4-BV-binding phages were subjected to two additional cycles of the incubation and wash step, resulting in 2nd, 3rd output phage. The ratio of output phage to input phage titers was calculated. B) Monoclonal analysis of C-CPE-derivative phage. CL4-BV-bound phage clones were isolated from the 2nd and 3rd output phages, and the interaction of the monoclonal phage with CL4-BV was examined by ELISA with anti-M13 antibody as described in Materials and methods. Data are expressed as relative binding to that of C-CPE-phage indicated by the most right column. doi:10.1371/journal.pone.0016611.g003

the CL4 binders modulated TJ barriers. We are investigating why the substitution of the amino acids with the other amino acids modulated CL4. These findings indicate that a BV screening system with a C-CPE library may be a powerful method to develop CL modulators.

The CL family forms various types of TJ barriers through combinations of its more than 20 members in homophilic/heterophilic CL strands [31,32]. Intercellular proteins ZO-1 and ZO-2 determine the localization of CL strands [33]. If a screening system to reconstitute heterogeneous CL strands with ZO-1 and/

Table 2. CL4-binding phages.

	304	305	307	309	313	318
C-CPE	S	S	S	N	S	K
Clone 1	R	V	S	A	R	R
2	R	S	V	A	R	K
3	G	D	G	R	T	R
4	S	A	P	R	S	A
5	R	S	L	K	S	K

The sequences of C-CPE mutant in the CL4-binding phages were analyzed. doi:10.1371/journal.pone.0016611.t002

or ZO-2 is developed, then useful and effective CL modulators can be identified. In this point, the BV system has extremely superior features. G protein and G protein-coupled receptors have been




Figure 4. Isolation of a novel CL4 modulator. A) Interaction of the C-CPE derivatives with CL4. C-CPE derivatives were prepared as his-tagged recombinant proteins. The C-CPE derivatives (0.02 μ g) were added to CL-BV-coated immunoplates, followed by detection of the C-CPE derivatives bound to CL-BV. Data are means \pm SD (n=4). B) Modulation of tight junction-barriers. Caco-2 cells were cultured on TranswellTM chambers. When TEER values reach a plateau, the cells were treated with C-CPE or C-CPE derivatives at the indicated concentrations. After 18 h of exposure to the C-CPEs, the cells were washed with medium to remove C-CPEs, and then the cells were cultured for an additional 24 h. Changes in TEER values were monitored during the C-CPEs treatment. Relative TEER values were calculated as the ratio of TEER values at 0 h. Data are representative of two independent experiments. The data are means \pm SD (n=4). doi:10.1371/journal.pone.0016611.g004

functionally reconstituted in BV [20,34], and functional γ -secretase complexes have also been reconstituted on BV [35]. In the near future, the reconstituted CL system on BV will be developed and used for the screening of CL binders and modulators, hopefully leading to breakthroughs in pharmaceutical therapies that target CLs.

Materials and Methods

Recombinant BV construction and Sf9 cell culture

Recombinant BV was prepared by using the Bac-to-Bac expression system, according to the manufacturer's instructions (Invitrogen, Gaithersburg, MD). Briefly, mouse CL1 and CL4 cDNA (kind gifts from Dr. M Furuse, Kobe University, Japan) were inserted into pFastBac1, and the resulting plasmids were transduced into DH10Bac *E. Coli* cells. Recombinant bacmid DNA was extracted from the cells. Sf9 cells were transduced with the bacmid coding CL, and the recombinant BV was recovered by centrifugation of the conditioned medium [36].

Preparation of the BV fractions

Sf9 cells (2×10^6 cells) were infected with recombinant BV at a multiplicity of infection of 5. Seventy-two hours after infection, the BV fraction was recovered from the culture supernatant of infected Sf9 cells by centrifugation. The pellets of the BV fraction were resuspended in Tris-buffered saline (TBS) containing 1% protease inhibitor cocktail (Sigma-Aldrich, St. Louis, MO) and then stored at 4°C until use. The expression of CL1 and CL4 in the BV was confirmed by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) and immunoblot analysis with anti-CL antibodies (Zymed Laboratory, South San Francisco, CA).

Preparation of mutant C-CPE library

C-CPE fragments in which the functional amino acids (S304, S305, S307, N309, S313 and K318) [24] were randomly mutated were prepared by polymerase chain reaction (PCR) with pET-H₁₀PER as a template, a forward primer (5'-catgcatgcccgatagaaaaagaaatccttgattagctgctg-3', Nco I site is underlined) and a reverse primer (5'-ttttcitttgcggcgcgcaaa^{smnt}gaaataat^{smnata}agg^{gta}smntcc^{smnata}smn^{smnat}-tagcttt-3', Not I site is underlined, and the randomly mutated amino acids are in italics). The PCR fragments were inserted into a pY03 phagemid at the NcoI/NotI sites [22]. The resultant phagemid containing the C-CPE mutant library was transduced into *E. coli* TG1 cells, and then the cells were stored at -80°C.

Preparation of phage

TG1 cells containing phagemid coding a scFv, C-CPE, C-CPE mutant or C-CPE mutant library were culture in 2YT medium containing 2% glucose and ampicillin. When the cells grew to a growing phase, M13K07 helper phages (Invitrogen) were added, and the medium was changed into 2YT medium containing ampicillin and kanamycin. After an additional 6 h of culture, the phages in the conditioned medium were precipitated with polyethylene glycol. The phages were suspended in phosphate-buffered saline (PBS) and stored at 4°C until use.

ELISA

Wild-BVs or CL-BVs (0.5 μ g/well) were adsorbed onto an immunoplate (Greiner Bio-One, Frickenhausen, Germany). The wells were washed with PBS and blocked with TBS containing 1.6% BlockAce (Dainippon Sumitomo Pharma, Osaka, Japan). C-CPEs or phages were incubated in the immunoplate, and the BV-bound C-CPEs or phages were detected by using anti-his-tag

antibody (Novagen, Darmstadt, Germany) or anti-M13 antibody (Amersham-Pharmacia Biotech, Uppsala, Sweden), respectively, horseradish peroxidase-labelled secondary antibody and TMB peroxidase substrate (Nacalai Tesque, Kyoto, Japan). The immunoreactive C-CPEs or phages were quantified by the measurement of absorbance at 450 nm. In the screening of phages, the data were normalized by the amounts of phages, which were quantified by ELISA for the FLAG-tag contained in the coat protein.

Selection of phage by using BV

A total of 0.5 μ g of BV was adsorbed onto an immunotube (Nunc, Roskilde, Denmark). The tube was washed with PBS and blocked with TBS containing 4.0% BlockAce. The BV-coated tubes were incubated with mixture of phages, and then the tubes were washed 15 times with PBS and 15 times with PBS containing 0.05% Tween 20. The phages bound to the tube were eluted with 100 mM HCl. TG1 cells were infected with the eluted phages, and phages were prepared as described above. The resulting phages were subjected to repeated selection by using the BV-coated immunotubes.

Identification of a phage clone

To identify an isolated phage clone, we performed PCR or sequencing analysis. We amplified the inserted fragment into the phagemid by PCR using forward primer 5'-caggaacacgctatgac-3' and reverse primer 5'-gtaaataaatttctgtatgagg-3'. The resultant PCR products were subjected to agarose gel electrophoresis followed by staining with ethidium bromide. We performed a sequence analysis with primer 5'-gtaaataaatttctgtatgagg-3'.

Measurement of phage titer

To quantify the concentration of phages, we measured the titer (colony formation unit (CFU)/ml) of the phage solution. Briefly, the phage solution was diluted to 10^{-5} - 10^{-10} with PBS. The diluted solution was seeded onto PetrifilmTM (Tech-Jam, Osaka, Japan). After 24 h of incubation, the colonies were counted, and the titer was calculated.

Purification of C-CPE mutants

C-CPE and C-CPE303, in which the CL-4 binding region of C-CPE was deleted, were prepared as described previously [13]. To prepare plasmid containing C-CPE mutants, the C-CPE mutant fragment was PCR-amplified by using phagemids coding C-CPE mutants as a template. The resulting PCR fragment was inserted into pET16b, and the sequence was confirmed. The plasmids were transduced into *E. coli* strain BL21 (DE3), and production of mutant C-CPEs was induced by the addition of isopropyl-D-thiogalactopyranoside. The harvested cells were lysed in buffer A (10 mM Tris-HCl, pH 8.0, 400 mM NaCl, 5 mM MgCl₂, 0.1 mM phenylmethanesulfonyl fluoride, 1 mM 2-mercaptoethanol, and 10% glycerol) that was supplemented with 8 M urea when necessary. The lysates were applied to HiTrapTM Chelating HP (GE Healthcare, Buckinghamshire, UK), and mutant C-CPEs were eluted with buffer A containing 100-400 mM imidazole. The buffer was exchanged with PBS by using a PD-10 column (GE Healthcare), and the purified protein was stored at -80°C until use. Purification of the mutant C-CPEs was confirmed by SDS-PAGE, followed by staining with Coomassie Brilliant Blue and by immunoblotting with anti-his-tag antibody (Novagen). Protein was quantified by using a BCA protein assay kit with bovine serum albumin as a standard (Pierce Chemical, Rockford, IL).

TEER assay

Caco-2 cells were seeded in Transwell™ chambers (Corning, NY) at a subconfluent density. The TEER of the Caco-2 monolayer cell sheets on the chamber was monitored by using a Millicell-ERS epithelial volt-ohmmeter (Millipore, Billerica, MA). When TEER values reached a plateau, indicating that TJs were well-developed in the cell sheets, the Caco-2 monolayers were treated with C-CPE or C-CPE mutants on the basal side of the chamber. Changes in TEER values were monitored. The TEER values were normalized by the area of the Caco-2 monolayer, and the TEER value of a blank Transwell™ chamber (background) was subtracted.

References

- Farquhar MG, Palade GE (1963) Junctional complexes in various epithelia. *J Cell Biol* 17: 375–412.
- Anderson JM, Van Itallie CM, Fanning AS (2004) Setting up a selective barrier at the apical junction complex. *Curr Opin Cell Biol* 16: 140–145.
- Balda MS, Matter K (1998) Tight junctions. *J Cell Sci* 111(Pt 5): 541–547.
- Tsukita S, Furuse M, Itoh M (2001) Multifunctional strands in tight junctions. *Nat Rev Mol Cell Biol* 2: 285–293.
- Mitic LL, Anderson JM (1998) Molecular architecture of tight junctions. *Annu Rev Physiol* 60: 121–142.
- Wodarz A, Nathke I (2007) Cell polarity in development and cancer. *Nat Cell Biol* 9: 1016–1024.
- Schneeberger EE, Lynch RD (2004) The tight junction: a multifunctional complex. *Am J Physiol* 286: C1213–C1228.
- Furuse M, Hata M, Furuse K, Yoshida Y, Haratake A, et al. (2002) Claudin-based tight junctions are crucial for the mammalian epidermal barrier: a lesson from claudin-1-deficient mice. *J Cell Biol* 156: 1099–1111.
- Nitta T, Hata M, Gotoh S, Seo Y, Sasaki H, et al. (2003) Size-selective loosening of the blood-brain barrier in claudin-5-deficient mice. *J Cell Biol* 161: 653–660.
- McClane BA (1994) *Clostridium perfringens* enterotoxin acts by producing small molecule permeability alterations in plasma membranes. *Toxicology* 87: 43–67.
- Katahira J, Inoue N, Horiguchi Y, Matsuda M, Sugimoto N (1997) Molecular cloning and functional characterization of the receptor for *Clostridium perfringens* enterotoxin. *J Cell Biol* 136: 1239–1247.
- Sonoda N, Furuse M, Sasaki H, Yonemura S, Katahira J, et al. (1999) *Clostridium perfringens* enterotoxin fragment removes specific claudins from tight junction strands: Evidence for direct involvement of claudins in tight junction barrier. *J Cell Biol* 147: 195–204.
- Kondoh M, Masuyama A, Takahashi A, Asano N, Mizuguchi H, et al. (2005) A novel strategy for the enhancement of drug absorption using a claudin modulator. *Mol Pharmacol* 67: 749–756.
- Jemal A, Siegel R, Ward E, Hao Y, Xu J, et al. (2008) Cancer statistics, 2008. *CA Cancer J Clin* 58: 71–96.
- Kominsky SL (2006) Claudins: emerging targets for cancer therapy. *Expert Rev Mol Med* 8: 1–11.
- Morin PJ (2005) Claudin proteins in human cancer: promising new targets for diagnosis and therapy. *Cancer Res* 65: 9603–9606.
- Michl P, Buchholz M, Rolke M, Kunsch S, Lohr M, et al. (2001) Claudin-4: a new target for pancreatic cancer treatment using *Clostridium perfringens* enterotoxin. *Gastroenterology* 121: 678–684.
- Santin AD, Cane S, Bellone S, Palmieri M, Siegel ER, et al. (2005) Treatment of chemotherapy-resistant human ovarian cancer xenografts in C.B-17/SCID mice by intraperitoneal administration of *Clostridium perfringens* enterotoxin. *Cancer Res* 65: 4334–4342.
- Loisel TP, Ansanay H, St-Onge S, Gay B, Boulanger P, et al. (1997) Recovery of homogeneous and functional beta 2-adrenergic receptors from extracellular baculovirus particles. *Nat Biotechnol* 15: 1300–1304.
- Sakihama T, Masuda K, Sato T, Doi T, Kodama T, et al. (2008) Functional reconstitution of G-protein-coupled receptor-mediated adenylyl cyclase activation by a baculoviral co-display system. *J Biotechnol* 135: 28–33.
- Sakihama T, Sato T, Iwanari H, Kitamura T, Sakaguchi S, et al. (2008) A simple detection method for low-affinity membrane protein interactions by baculoviral display. *PLoS ONE* 3: e4024.
- Ebihara C, Kondoh M, Hasuike N, Harada M, Mizuguchi H, et al. (2006) Preparation of a claudin-targeting molecule using a C-terminal fragment of *Clostridium perfringens* enterotoxin. *J Pharmacol Exp Ther* 316: 255–260.
- Fujita K, Katahira J, Horiguchi Y, Sonoda N, Furuse M, et al. (2000) *Clostridium perfringens* enterotoxin binds to the second extracellular loop of claudin-3, a tight junction integral membrane protein. *FEBS Lett* 476: 258–261.
- Takahashi A, Komiya E, Kakutani H, Yoshida T, Fujii M, et al. (2008) Domain mapping of a claudin-4 modulator, the C-terminal region of C-terminal fragment of *Clostridium perfringens* enterotoxin, by site-directed mutagenesis. *Biochem Pharmacol* 75: 1639–1648.
- Meunier V, Bourrie M, Berger Y, Fabre G (1995) The human intestinal epithelial cell line Caco-2; pharmacological and pharmacokinetic applications. *Cell Biol Toxicol* 11: 187–194.
- Hanna PC, Mietzner TA, Schoolnik GK, McClane BA (1991) Localization of the receptor-binding region of *Clostridium perfringens* enterotoxin utilizing cloned toxin fragments and synthetic peptides. *J Biol Chem* 266: 11037–11043.
- Ofner S, Hekele A, Teichmann U, Weinberger S, Gross S, et al. (2005) Epithelial tight junction proteins as potential antibody targets for pancreatic cancer therapy. *Cancer Immunol Immunother* 54: 431–445.
- Ling J, Liao H, Clark R, Wong MS, Lo DD (2008) Structural constraints for the binding of short peptides to claudin-4 revealed by surface plasmon resonance. *J Biol Chem* 283: 30585–30595.
- Suzuki M, Kato-Nakano M, Kawamoto S, Furuya A, Abe Y, et al. (2009) Therapeutic antitumor efficacy of monoclonal antibody against Claudin-4 for pancreatic and ovarian cancers. *Cancer Sci* 100: 1623–1630.
- Romani C, Comper F, Bandiera E, Ravaggi A, Bignotti E, et al. (2009) Development and characterization of a human single-chain antibody fragment against claudin-3: a novel therapeutic target in ovarian and uterine carcinomas. *Am J Obstet Gynecol* 201: 70 e71–79.
- Furuse M, Furuse K, Sasaki H, Tsukita S (2001) Conversion of zonulae occludentes from tight to leaky strand type by introducing claudin-2 into Madin-Darby canine kidney I cells. *J Cell Biol* 153: 263–272.
- Furuse M, Sasaki H, Tsukita S (1999) Manner of interaction of heterogeneous claudin species within and between tight junction strands. *J Cell Biol* 147: 891–903.
- Umeda K, Ikenouchi J, Katahira-Tayama S, Furuse K, Sasaki H, et al. (2006) ZO-1 and ZO-2 independently determine where claudins are polymerized in tight-junction strand formation. *Cell* 126: 741–754.
- Masuda K, Itoh H, Sakihama T, Akiyama C, Takahashi K, et al. (2003) A combinatorial G protein-coupled receptor reconstitution system on budded baculovirus. *J Biol Chem* 278: 24552–24562.
- Hayashi I, Urano Y, Fukuda R, Isoo N, Kodama T, et al. (2004) Selective reconstitution and recovery of functional gamma-secretase complex on budded baculovirus particles. *J Biol Chem* 279: 38040–38046.
- Saeki R, Kondoh M, Kakutani H, Tsunoda S, Mochizuki Y, et al. (2009) A novel tumor-targeted therapy using a claudin-4-targeting molecule. *Mol Pharmacol* 76: 918–926.

Acknowledgments

We thank Drs. S. Tsunoda (National Institute of Biomedical Innovation, Japan), Y. Tsutsumi, Y. Mukai (Osaka University, Japan) for their kind instruction of phage display technology. We also thank Drs. Y. Horiguchi (Osaka University, Japan), S. Tsukita (Kyoto University, Japan) and members of our laboratory for providing us C-CPE cDNA, CL-expressing cells and their useful comments and discussion, respectively.

Author Contributions

Conceived and designed the experiments: MK TS TH KY. Performed the experiments: HK AT MK YS TY TS. Analyzed the data: HK AT MK KY. Contributed reagents/materials/analysis tools: HK AK TS TH. Wrote the manuscript: HK MK TY.



Claudin as a Target for Drug Development

A. Takahashi, M. Kondoh*, H. Suzuki and K. Yagi*

Graduate School of Pharmaceutical Sciences, Osaka University, Suita, Osaka 565-0871, Japan

Abstract: Tight junctions (TJs) play pivotal roles in the fence and barrier functions of epithelial and endothelial cell sheets. Since the 1980s, the modulation of the TJ barrier has been utilized as a method for drug absorption. Over the last decade, the structural and functional biochemical components of TJs, such as occludin and claudin, have been determined, providing new insights into TJ-based pharmaceutical therapy. For example, the modulation of the claudin barrier enhances the jejunal absorption of drugs, and claudin expression is deregulated in cancer cells. Claudin is a co-receptor for the hepatitis C virus. Moreover, claudin is modulated during inflammatory conditions. These findings indicate that claudins are promising drug targets. In this review, we discuss the seeds of claudin-based drug development, which may provide potential pharmaceutical breakthroughs in the future.

Keywords: Tight junction, claudin, cancer, inflammation, infection.

INTRODUCTION

Tight junctions (TJs) limit the movement of molecules through the intercellular space in epithelial and endothelial sheets, and they are located on the most apical part of cells [1, 2]. Electron microscopy has revealed that TJs appear as a series of continuous, anastomotic and intramembranous particle strands. Tsukita's group performed a series of biochemical analyses that clearly showed that the tetra-transmembrane proteins occludin and claudin are components of the TJ [3-5]. The claudin family contains more than 20 members. Interestingly, the expression profiles and the TJ-barrier function of the claudin family members are tissue-specific. For example, claudin-1 is involved in the epidermal barrier, and claudin-5 is involved in the blood-brain barrier [2, 6, 7]. It appears that claudin forms heteromeric and/or homomeric strands in TJs and that the combination and mixing ratios of different claudins determines the tissue-specific barrier properties of TJs [5, 8]. Epithelial cell sheets have bicellular TJs between adjacent cells and tricellular TJs at which three adjacent cells join together. Occludin and claudins are components of bicellular TJs. The occludin-related protein tricellulin has been recently identified to be a component of tricellular TJs [9]. Tricellulin is ubiquitously expressed in epithelial junctions of tissues and organs throughout the body. Down regulation of tricellulin mRNA by RNA interference resulted in disruption of epithelial barrier in an epithelial cell line [9]. However, human tricellulin mutations had no effect on epidermal, respiratory, renal or intestinal barrier [10]. Whether tricellulin can be a target for drug development is unclear.

Functions of TJs are classified as fence- and barrier- functions. Modulation of the TJ barrier has been a popular strategy used to promote drug absorption since the 1980s (See reviews [11, 12]). Sodium caprate is clinically used as an absorption enhancer of drug. Disturbance of either the TJ-fence function or the TJ-barrier function causes human diseases. Disturbance of the TJ-fence function followed by a loss of cellular polarity often occurs in tumorigenesis (See reviews [13-16]). TJs regulate the paracellular passage of ions, molecules, pathogens and inflammatory cells in epithelial and endothelial cell sheets [17-19]. The TJ-barrier becomes deregulated in various human diseases, including infections, inflammation and hereditary diseases (See reviews [20, 21]). Based on these findings, novel therapeutic strategies for TJ-related diseases have been proposed. In the present review, we discuss the seeds of claudin-based pharmaceutical therapies for human diseases relevant to TJs.

CANCER AND CLAUDIN

Malignant tumors are a major cause of death. Approximately 7.6 million people worldwide died from cancer in 2007, and 90% of tumors are derived from epithelial tissue [22]. Normal epithelial tissues develop cellular polarity, whereas the epithelial polarity is often deregulated during tumorigenesis [23]. TJs are localized between adjacent epithelial cells and separate the apical and basolateral membrane domains, which vary in protein and lipid content, resulting in the maintenance of the cell polarity. Claudins are deregulated in various cancers [13-16]. Claudin may regulate cancer metastasis by modulating activation of matrix metalloproteinases [11]. In this section, we discuss recent breakthroughs in claudin-targeted cancer therapy.

Claudin as a Diagnostic Marker

Claudin proteins are frequently overexpressed in ovarian cancers. In ovarian cancer cells with a high level of claudin-4, the critical claudin-4 promoter region exhibits a low level of DNA methylation and a high level of histone H3 acetylation [24]. Claudin-4 was detected in the 32 of 63 plasma samples of patients with ovarian cancers. Among 50 patients without ovarian cancer, only one had claudin-4-positive plasma. Thus, claudin-4 has a high specificity for the detection of ovarian cancers *via* a blood test, indicating that claudin-4 may be a diagnostic marker for ovarian cancer [25]. Because of the high specificity of claudin expression patterns in cancers, claudin might be a novel non-invasive diagnostic marker for cancer therapy.

Anti-Claudin Antibody

One of the most popular strategies for claudin-targeted cancer therapy is the preparation of antibody against the extracellular region of claudin. However, attempts to prepare anti-claudin antibodies have had little success because claudin has low antigenicity and is highly conserved in various species. A strain of autoimmune mice, BXS^B, was immunized with a human pancreatic cancer cell line, resulting in the successful preparation of anti-claudin-4 monoclonal antibody that recognizes the extracellular region of claudin-4 [26, 27]. Moreover, the antibody mediated antibody-dependent cell cytotoxicity (ADCC) and *in vivo* anti-tumor activity. ScFv against the extracellular region of claudin-3 was isolated by using the ETH-2 Gold phage display library, which is a synthetic human recombinant antibody library that contains >10⁹ possible antibody combinations in an scFv format [28, 29]. Immunization with DNA encoding the first extracellular loop of claudin-18 made success on preparation of anti-claudin-18 monoclonal antibody [30]. These successes in the preparation of anti-claudin antibody are likely to lead to a breakthrough in the development of claudin-targeted cancer therapy.

*Address correspondence to these authors at the Graduate School of Pharmaceutical Sciences, Osaka University, Suita, Osaka 565-0871, Japan; Tel: +81-6-6879-8196; Fax: +81-6-6879-8199; E-mail: masuo@phs.osaka-u.ac.jp
Tel: +81-6-6879-8195; Fax: +81-6-6879-8195; E-mail: yagi@phs.osaka-u.ac.jp

Clostridium Perfringens Enterotoxin

Another approach to targeting claudin in cancer therapy is the use of *Clostridium perfringens* enterotoxin (CPE). CPE is a single-chain polypeptide of 35 kDa that causes food poisoning in humans. The functional domains of CPE consist of the N-terminal cytotoxic region and the C-terminal receptor-binding region [31]. Claudin-3 and -4 serve as the receptors for CPE. CPE binds to the second extracellular loop of claudin-3 and -4 [32] (Fig. 1). We previously prepared a claudin-targeting molecule (C-CPE-PSIF) by fusion of the C-terminal fragment of CPE (C-CPE) with the protein synthesis inhibitory factor (PSIF) derived from *Pseudomonas aeruginosa* exotoxin. C-CPE-PSIF, but not PSIF, is cytotoxic to claudin-4 expressing cells. TJ-undeveloped cells are more sensitive to C-CPE-PSIF than TJ-developed cells. Polarized epithelial cells are sensitive to the basolaterally applied C-CPE-PSIF, but they are less sensitive to the apically applied C-CPE-PSIF. A claudin-targeting molecule may recognize the cellular polarity. Intratumoral injection of C-CPE-PSIF reduced tumor growth. These findings indicate that C-CPE may be a novel molecule for drug delivery and cancer therapy [33]. The receptor-binding region of C-CPE fused to TNF was cytotoxic in human ovarian cancer cells [34]. Thus, CPE fragments might be a tool for claudin-targeting therapy. Treatment of mice with claudin-3 siRNA suppressed ovarian tumor growth and metastasis [35]. Claudin gene silencing with siRNA is also potent anti-tumor agents.




Fig. (1). Schematic illustration of interaction of C-CPE and claudin. Claudin is a tetra-transmembrane protein. C-CPE interacted with the 2nd loop region of claudin *via* its C-terminal domain [32, 76].

INFECTION AND CLAUDINS

Twenty million people die from infectious diseases each year. Most pathogens enter the body through nasal, pulmonary, intestinal and genital mucosa, and the mucosal epithelial cell sheets play a pivotal role as the first line of defense against the pathogens. Invading pathogens are distributed throughout the organ *via* endothelial cells of the blood vessels. TJs seal intercellular spaces between adjacent cells, preventing entry of the pathogens into the body and into the organ across the paracellular spaces. Disruption of mucosal TJ seals allows pathogens to enter into the body and the organ. In this section, we review the recent findings on the relationship between infections and claudins.

West Nile Virus (WNV) and Claudin

WNV, a neurotropic flavivirus, is a human pathogen that targets neurons and causes potentially lethal encephalitis in 1% to 2% of WNV-infected febrile patients [36]. No therapeutic agents or vaccines have been approved for use against WNV infection. Langerhans cells in the skin become infected with WNV by the bite of a

carrier mosquito. WNV replicates in the regional tissues and lymph nodes, which results in the dissemination of the virus into the bloodstream. The following second replication proceeds at several sites in the host, including epithelial cells in the skin, kidney, intestine and testis, and then WNV may ultimately invade the brain [37]. The infection of the nervous system is characteristic of the most severe cases of WNV disease, and it often results in death or long-term neurologic sequelae [38]. Understanding the mechanism of the second infection and the viral entry into the brain is critical for the development of therapies against WNV. In WNV-infected epithelial cells, claudin-1, -2, -3 and -4 are degraded, followed by a disruption in the TJ barrier without cell death. The capsid of the WNV was responsible for the modulation of the TJ barrier [39]. These findings suggest that an inducer of claudin may be a promising candidate for pharmaceutical agents to inhibit the dissemination of WNV. Whether or not the WNV modulates the blood-brain barrier *via* the modulation of claudin-5 is an unsettled question.

Human Immunodeficiency Virus (HIV) and Claudin

HIV encephalitis (HIVE), including behavioral, motor, and cognitive impairments, is a common condition in the late stage of HIV-associated dementia [40]. Invasion of HIV into the brain and the transmigration of HIV-infected lymphocytes into the brain are the major causes of HIVE [41]. The blood-brain barrier (BBB), which is responsible for the regulation of solutes and cells between the peripheral circulation and the central nervous system, is comprised of the brain microvascular endothelial cells. Adjacent brain microvascular endothelial cells are connected by TJs that limit paracellular flux and restrict permeability [42]. The BBB frequently breaks down in patients with HIVE [41]. Claudin-5 plays a pivotal role in the BBB [7]. Treatment of human brain microvascular endothelial cells with HIV Gp120 envelope glycoprotein decreased the claudin-5 levels, followed by a disruption of the TJ barrier [43]. Claudin-5 levels were lower in brain microvessels from HIV patients with HIVE compared with brain microvessels from HIV patients without HIVE [44]. The deregulation of the claudin-5 barrier by HIV may be responsible for the breakdown of the BBB in HIV patients. Cannabinoids, the active ingredients in marijuana, reduce pain and improve the quality of life in HIV patients [45]. HIV activates signal transducers and activators of transcription-1 (STAT-1) [46]. Cannabinoids and an inhibitor of STAT-1 prevented the down-regulation of claudin-5 in the HIV Gp120- and HIV-treated human brain microvascular endothelial cells, respectively [43, 44]. These findings indicate that an inducer of claudin-5 may be a pharmaceutical agent for HIVE.

Hepatitis C Virus (HCV) and Claudin

Approximately 170 million people worldwide are infected with HCV. More than 80% of acute infections become persistent, resulting in liver fibrosis, cirrhosis, and hepatocellular carcinoma [47]. HCV infects human hepatocytes but not murine hepatocytes, and the detailed mechanism responsible for this difference has remained obscure. There is no pharmaceutical agent that prevents HCV infection. HCV attaches to tetraspanin CD81 and scavenger receptor class B type I (SR-BI) on host cells through its envelop glycoprotein [48, 49]. However, when CD81 and SR-BI were expressed in non-primate cell lines, the cells were still resistant to HCV entry [50, 51]. Recent studies to identify the additional factors that are needed to render non-human cells susceptible to HCV entry revealed that claudin-1 and occludin are co-receptors for HCV entry [51, 52]. HCV envelop proteins interact with the first extracellular loop region of claudin-1 and the second extracellular loop region of occludin [51, 52]. Binders to CD81, SR-BI, claudin-1 or/and occludin are expected to inhibit HCV entry. The HCV genome is frequently mutated; thus, pharmaceutical agents that recognize host molecules, such as the receptors, may be promising candidates for the prevention of HCV infection.

INFLAMMATORY BOWEL DISEASE (IBD) AND CLAUDIN

Inflammatory bowel disease (IBD), including ulcerative colitis and Crohn's disease, is characterized by an activated mucosal immune system that leads to impaired epithelial barrier function and tissue destruction with relapsing diarrhea [53, 54]. Ulcerative colitis is characterized by chronic inflammation and ulcers in the colon, while Crohn's disease causes ulcers and swelling of the mucosa on all areas of the digestive tract from the mouth to the anus. A common feature of IBD is enhanced permeability of the intestinal epithelium and disruption of the epithelial barrier. In this section, we summarize the recent findings on the relationship between IBD and claudins.

Changes of Claudins in IBD

The epithelial barrier function is impaired in ulcerative colitis, and ulcerative colitis is associated with decreased numbers of TJ strands in the epithelial barrier [55]. Biochemical analysis of TJ components in rectal biopsy specimens from patients with active ulcerative colitis revealed that the protein and mRNA levels of claudin-4 and -7 were decreased, whereas the protein and mRNA levels of claudin-2 were increased, as compared with control patients [56]. Overexpression of claudin-2 led to a decrease in the TJ barrier in an epithelial cell line, whereas claudin-4 or -7 transfection elevated the epithelial barrier function [57, 58]. Thus, the down-regulation of claudin-4/7 and the up-regulation of claudin-2 can lead to altered TJ structure, resulting in impaired epithelial function in active ulcerative colitis. However, claudin-deficient mice or claudin-overexpressing mice did not reproduce the pathology of IBD. Whether change in claudins is cause of IBD or result from IBD remains to be proved.

Although the precise etiology of IBD remains unknown, it is well accepted that IBD results from a deregulated mucosal immune response to environmental factors in genetically susceptible hosts. In IBD patients, the primary defect may be due to an abnormal intestinal epithelial barrier function [59]. The SAMP1/YitFc (SAMP) mouse strain is a spontaneous model of IBD that closely resembles Crohn's disease due to its histological features and localization to the terminal ileum [60]. The deregulated epithelial barrier function in SAMP mice is accompanied by an increase in claudin-2 and a decrease in occludin [61, 62].

FoxO4 is a member of the forkhead box transcription factor O (FoxO) subfamily, which has unique cell type-specific functions that regulate target genes and are involved in the regulation of immune responses [63, 64]. FoxO4-null mice were more susceptible to trinitrobenzene sulfonic acid-induced colitis [65]. FoxO4 deficiency increased the intestinal epithelial permeability and down-regulated the TJ proteins ZO-1 and claudin-1. Immunohistochemical analysis revealed that epithelial expression of FoxO4 was significantly down-regulated in patients with active ulcerative colitis as compared to patients with inactive ulcerative colitis [66]. Thus, FoxO4 might be a target for ulcerative colitis therapy.

A Potent Pharmaceutical Agent for IBD

Pro-inflammatory cytokines, such as tumor necrosis factor- α (TNF- α) and interferon- γ , are key mediators for the disruption of the epithelial barrier associated with Crohn's disease [55, 66, 67]. Expression of claudin-2 was increased by TNF- α in epithelial cells [68]. Experimental colitis model mice showed the down-regulation of occludin and up-regulation of claudin-2. Deletion of TNF- α receptor attenuated these changes of occludin and claudin-2 in the experimental colitis model. Importantly, anti-TNF treatment infliximab, which is currently used in Crohn's disease and ulcerative colitis, suppressed the reduction of occludin and elevation of claudin-2 in the experimental colitis model [69].

n-3 polyunsaturated fatty acids (PUFAs), which are abundant in fish oil and include eicosapentaenoic acid and docosahexaenoic acid, have beneficial effects on IBD [70-72]. In an experimental IBD model induced by treatment with trinitrobenzene sulfonic acid, the distribution of TJ proteins, including occludin and claudin-1, was affected; however, the administration of n-3 PUFAs prevented this redistribution of TJ proteins [73].

Probiotics are living bacteria that, when ingested in sufficient quantity, improve the health of the host beyond their inherent basic nutrition [74]. Probiotics have anti-inflammatory effects in IBD. VSL#3, a mixture of 8 probiotic bacterial strains, provided protection against intestinal inflammation in an experimental colitis model. Probiotics also attenuated the enhancement of epithelial permeability and the reduction of TJ components, including occludin, claudin-1 and -4 in the experimental model [75]. Therefore, compounds that enhance the TJ barrier function are candidates for IBD therapy.

CONCLUSIONS

Epithelium and endothelium are located between the outer and inner components of the body or tissues. Most malignant tumors are derived from epithelium. Moreover, epithelium and endothelium are also barriers that prevent invading pathogens and inflammatory cells from entering into the body and tissues. Therefore, the epithelium and endothelium are excellent targets for drug delivery systems, anti-tumor agents, anti-infection agents and anti-inflammatory agents.

Recent studies have revealed the involvement of claudin in some human diseases relevant to TJs (Table 1). Claudin is often overexpressed in human cancers [13-16]. Therefore, a cancer therapy approach that uses claudin ligands is sought. Suzuki *et al.* used autoimmune mice to successfully prepare an anti-claudin-3 monoclonal antibody that mediated ADCC [26]. We anticipate that a novel claudin-targeted cancer therapy will be forthcoming. TJ components are also associated with infections. Claudin-1 and occludin are co-receptors for HCV [51, 52]. The claudin-5 level was reduced in brain microvessels of patients with HIV [44], and cannabinoids, a clinically used agent for HIV patients, prevented the down-regulation of claudin-5 [43]. These findings indicate that a

Table 1. Perspective on Claudin-Targeted Therapies

Applications	Claudins	References
A diagnostic marker for ovarian cancers	Claudin-4	[25]
Inhibitor of WNV dissemination	Claudin-1-4	[39]
Inhibitor of HIV encephalitis	Claudin-5	[43-45]
Inhibitor of HCV infection	Claudin-1	[51]
Inhibitor of intestinal inflammation in IBD	Claudin-1-4	[69, 73, 75]

WNV, west Nile virus; HIV, human immunodeficiency virus; HCV, hepatitis C virus; IBD, inflammatory bowel disease.

claudin/occludin binder and an inducer of claudin-5 may be an inhibitor of HCV infection and a therapeutic agent for HIVE patients. Disruption of the intestinal epithelial barrier is a common feature in patients with IBD. A chemical compound that strengthens the claudin barrier function will be a promising drug for IBD.

Biochemical and functional information regarding TJs has accumulated since the identification of occludin in 1993, and the deregulation of claudins has been observed in several human diseases [16, 20, 21]. The potential of TJ-based therapies is promising. We believe that TJ-targeted therapies might provide a breakthrough in pharmaceutical therapy in the future.

ACKNOWLEDGEMENTS

We also thank the members of our laboratory for their useful comments and discussion. Our work referred in this review was partly supported by MEXT KAKENHI (21689006), by a Health and Labor Sciences Research Grants from the Ministry of Health, Labor and Welfare of Japan, by Takeda Science Foundation, by a Suzuken Memorial Foundation and by a grant from Kansai Biomedical Cluster project in Saito, which is promoted by the Knowledge Cluster Initiative of the Ministry of Education, Culture, Sports, Science and Technology, Japan. A.T. is supported by Research Fellowships of the Japan Society for the Promotion of Science for Young Scientists.

ABBREVIATIONS

TJ	=	Tight junction
ADCC	=	antibody-dependent cell cytotoxicity
CPE	=	<i>Clostridium perfringens</i> enterotoxin
C-CPE	=	the C-terminal fragment of CPE
PSIF	=	protein synthesis inhibitory factor
WNV	=	West Nile virus
HIV	=	human immunodeficiency virus
HIVE	=	HIV encephalitis
BBB	=	blood-brain barrier
STAT-1	=	signal transducers and activators of transcription-1
HCV	=	hepatitis C virus
SR-BI	=	scavenger receptor class B type I
IBD	=	inflammatory bowel disease
FoxO	=	forkhead box transcription factor O
TNF	=	tumor necrosis factor
PUFAs	=	n-3 polyunsaturated fatty acids

REFERENCES

- Anderson, J. Molecular structure of tight junctions and their role in epithelial transport. *News Physiol. Sci.*, **2001**, *16*, 126-130.
- Tsukita, S.; Furuse, M.; Itoh, M. Multifunctional strands in tight junctions. *Nat. Rev. Mol. Cell Biol.*, **2001**, *2*, 285-293.
- Furuse, M.; Fujita, K.; Hiiragi, T.; Fujimoto, K.; Tsukita, S. Claudin-1 and -2: novel integral membrane proteins localizing at tight junctions with no sequence similarity to occludin. *J. Cell Biol.*, **1998**, *141*, 1539-1550.
- Furuse, M.; Hirase, T.; Itoh, M.; Nagafuchi, A.; Yonemura, S.; Tsukita, S.; Tsukita, S. Occludin: a novel integral membrane protein localizing at tight junctions. *J. Cell Biol.*, **1993**, *123*, 1777-1788.
- Furuse, M.; Tsukita, S. Claudins in occluding junctions of humans and flies. *Trends Cell Biol.*, **2006**, *16*, 181-188.
- Furuse, M.; Hata, M.; Furuse, K.; Yoshida, Y.; Haratake, A.; Sugitani, Y.; Noda, T.; Kubo, A.; Tsukita, S. Claudin-based tight junctions are crucial for the mammalian epidermal barrier: a lesson from claudin-1-deficient mice. *J. Cell Biol.*, **2002**, *156*, 1099-1111.
- Nitta, T.; Hata, M.; Gotoh, S.; Seo, Y.; Sasaki, H.; Hashimoto, N.; Furuse, M.; Tsukita, S. Size-selective loosening of the blood-brain barrier in claudin-5-deficient mice. *J. Cell Biol.*, **2003**, *161*, 653-660.
- Furuse, M.; Sasaki, H.; Tsukita, S. Manner of interaction of heterogeneous claudin species within and between tight junction strands. *J. Cell Biol.*, **1999**, *147*, 891-903.
- Ikenouchi, J.; Furuse, M.; Furuse, K.; Sasaki, H.; Tsukita, S.; Tsukida, S. Tricellulin constitutes a novel barrier at tricellular contacts of epithelial cells. *J. Cell Biol.*, **2005**, *171*, 939-945.
- Riazuddin, S.; Ahmed, Z.M.; Fanning, A.S.; Lagziel, A.; Kitajiri, S.; Ramzan, K.; Khan, S.N.; Chatteraj, P.; Friedman, P.L.; Anderson, J.M.; Belyantseva, I.A.; Forge, A.; Riazuddin, S.; Friedman, T.B. Tricellulin is a tight-junction protein necessary for hearing. *Am. J. Hum. Genet.*, **2006**, *79*, 1040-1051.
- Kondoh, M.; Yoshida, T.; Kakutani, H.; Yagi, K. Targeting tight junction proteins-significance for drug development. *Drug Discov. Today*, **2008**, *13*, 180-186.
- Matsuhisa, K.; Kondoh, M.; Takahashi, A.; Yagi, K. Tight junction modulator and drug delivery. *Expert Opin. Drug Deliv.*, **2009**, *6*, 509-515.
- Kominsky, S.L. Claudins: emerging targets for cancer therapy. *Expert Rev. Mol. Med.*, **2006**, *8*, 1-11.
- Morin, P.J. Claudin proteins in human cancer: promising new targets for diagnosis and therapy. *Cancer Res.*, **2005**, *65*, 9603-9606.
- Swissel, K.; Macek, R.; Kubbies, M. Role of claudins in tumorigenesis. *Adv. Drug Deliv. Rev.*, **2005**, *57*, 919-928.
- Tsukita, S.; Yamazaki, Y.; Katsuno, T.; Tamura, A.; Tsukita, S. Tight junction-based epithelial microenvironment and cell proliferation. *Oncogene*, **2008**, *27*, 6930-6938.
- Schulzke, D.; Ploeger, S.; Amasheh, M.; Fromm, A.; Zeissig, S.; Troeger, H.; Richter, J.; Bojarski, C.; Schumann, M.; Fromm, M. Epithelial tight junctions in intestinal inflammation. *Ann. N.Y. Acad. Sci.*, **2009**, *1165*, 294-300.
- Van Itallie, C.M.; Betts, L.; Smedley, J.G. 3rd; McClane, B.A.; Anderson, J.M. Structure of the claudin-binding domain of *Clostridium perfringens* enterotoxin. *J. Biol. Chem.*, **2008**, *283*, 268-274.
- Argaw, A.T.; Gurfein, B.T.; Zhang, Y.; Zameer, A.; John, G.R. VEGF-mediated disruption of endothelial CLN-5 promotes blood-brain barrier breakdown. *Proc. Natl. Acad. Sci. USA*, **2009**, *106*, 1977-1982.
- Furuse, M. Knockout animals and natural mutations as experimental and diagnostic tool for studying tight junction functions *in vivo*. *Biochim. Biophys. Acta*, **2009**, *1788*, 813-819.
- Sawada, N.; Murata, M.; Kikuchi, K.; Osanai, M.; Tobioka, H.; Kojima, T.; Chiba, H. Tight junctions and human diseases. *Med. Electron. Microsc.*, **2003**, *36*, 147-156.
- Jemal, A.; Siegel, R.; Ward, E.; Hao, Y.; Xu, J.; Murray, T.; Thun, M.J. Cancer statistics, 2008. *CA Cancer J. Clin.*, **2008**, *58*, 71-96.
- Wodarz, A.; Nathke, I. Cell polarity in development and cancer. *Nat. Cell Biol.*, **2007**, *9*, 1016-1024.
- Honda, H.; Pazin, M.J.; Ji, H.; Wernyj, R.P.; Morin, P.J. Crucial roles of Sp1 and epigenetic modifications in the regulation of the CLDN4 promoter in ovarian cancer cells. *J. Biol. Chem.*, **2006**, *281*, 21433-21444.
- Li, J.; Sherman-Baust, C.A.; Tsai-Turton, M.; Bristow, R.E.; Roden, R.B.; Morin, P.J. Claudin-containing exosomes in the peripheral circulation of women with ovarian cancer. *BMC Cancer*, **2009**, *9*, 244.
- Suzuki, M.; Kato-Nakano, M.; Kawamoto, S.; Furuya, A.; Abe, Y.; Misaka, H.; Kimoto, N.; Nakamura, K.; Ohta, S.; Ando, H. Therapeutic antitumor efficacy of monoclonal antibody against Claudin-4 for pancreatic and ovarian cancers. *Cancer Sci.*, **2009**, *100*, 1623-1630.
- Yin, B.W.; Wong, G.Y.; Lloyd, K.O.; Oettgen, H.F.; Welt, S. Increased yields of IgG2a- and IgG3-secreting hybridomas after fusion of B cells from mice with autoimmune diseases. *J. Immunol. Methods*, **1991**, *144*, 165-173.
- Pini, A.; Viti, F.; Santucci, A.; Carnemolla, B.; Zardi, L.; Neri, P.; Neri, D. Design and use of a phage display library. Human antibodies with subnanomolar affinity against a marker of angiogenesis eluted from a two-dimensional gel. *J. Biol. Chem.*, **1998**, *273*, 21769-21776.
- Romani, C.; Comper, F.; Bandiera, E.; Ravaggi, A.; Bignotti, E.; Tassi, R.A.; Pecorelli, S.; Santin, A.D. Development and characterization of a human single-chain antibody fragment against claudin-3: a novel therapeutic target in ovarian and uterine carcinomas. *Am. J. Obstet. Gynecol.*, **2009**, *201*, 70. e71-79.
- Sahin, U.; Koslowski, M.; Dhaene, K.; Usener, D.; Brandenburg, G.; Seitz, G.; Huber, C.; Tureci, O. Claudin-18 splice variant 2 is a pan-cancer target suitable for therapeutic antibody development. *Clin. Cancer Res.*, **2008**, *14*, 7624-7634.
- Hanna, P.C.; Wieckowski, E.U.; Mietzner, T.A.; McClane, B.A. Mapping of functional regions of *Clostridium perfringens* type A enterotoxin. *Infect. Immun.*, **1992**, *60*, 2110-2114.
- Fujita, K.; Kathaira, J.; Horiguchi, Y.; Sonoda, N.; Furuse, M.; Tsukita, S. *Clostridium perfringens* enterotoxin binds to the second extracellular loop of claudin-3, a tight junction integral membrane protein. *FEBS Lett.*, **2000**, *476*, 258-261.
- Saeki, R.; Kondoh, M.; Kakutani, H.; Tsunoda, S.; Mochizuki, Y.; Hamakubo, T.; Tsutsumi, Y.; Horiguchi, Y.; Yagi, K. A novel tumor-targeted therapy using a claudin-4-targeting molecule. *Mol. Pharmacol.*, **2009**, *76*, 918-926.
- Yuan, X.; Lin, X.; Manorek, G.; Kanatani, I.; Cheung, L.H.; Rosenblum, M.G.; Howell, S.B. Recombinant CPE fused to tumor necrosis factor targets human ovarian cancer cells expressing the claudin-3 and claudin-4 receptors. *Mol. Cancer Ther.*, **2009**, *8*, 1906-1915.

- [35] Huang, Y.H.; Bao, Y.; Peng, W.; Goldberg, M.; Love, K.; Bumcrot, D.A.; Cole, G.; Langer, R.; Anderson, D.G.; Swicki, J.A. Claudin-3 gene silencing with siRNA suppresses ovarian tumor growth and metastasis. *Proc. Natl. Acad. Sci. USA*, **2009**, *106*, 3426-3430.
- [36] Hayes, E.B.; Gubler, D.J. West Nile virus: epidemiology and clinical features of an emerging epidemic in the United States. *Annu. Rev. Med.*, **2006**, *57*, 181-194.
- [37] Samuel, M.A.; Diamond, M.S. Pathogenesis of West Nile Virus infection: a balance between virulence, innate and adaptive immunity, and viral evasion. *J. Virol.*, **2006**, *80*, 9349-9360.
- [38] Klee, A.L.; Maidin, B.; Edwin, B.; Poshni, I.; Mostashari, F.; Fine, A.; Layton, M.; Nash, D. Long-term prognosis for clinical West Nile virus infection. *Emerg. Infect. Dis.*, **2004**, *10*, 1405-1411.
- [39] Medigeshi, G.R.; Hirsch, A.J.; Brien, J.D.; Uhrlaub, J.L.; Mason, P.W.; Wiley, C.; Nikolich-Zugich, J.; Nelson, J.A. West Nile virus capsid degradation of claudin proteins disrupts epithelial barrier function. *J. Virol.*, **2009**, *83*, 6125-6134.
- [40] McArthur, J.C. HIV dementia: an evolving disease. *J. Neuroimmunol.*, **2004**, *157*, 3-10.
- [41] Banks, W.A.; Ercal, N.; Price, T.O. The blood-brain barrier in neuroAIDS. *Curr. HIV Res.*, **2006**, *4*, 259-266.
- [42] Hawkins, B.T.; Davis, T.P. The blood-brain barrier/neurovascular unit in health and disease. *Pharmacol. Rev.*, **2005**, *57*, 173-185.
- [43] Lu, T.S.; Avraham, H.K.; Seng, S.; Tachado, S.D.; Kozel, H.; Makriyannis, A.; Avraham, S. Cannabinoids inhibit HIV-1 Gp120-mediated insults in brain microvascular endothelial cells. *J. Immunol.*, **2008**, *181*, 6406-6416.
- [44] Chaudhuri, A.; Yang, B.; Gendelman, H.E.; Persidsky, Y.; Kanmogne, G.D. STAT1 signaling modulates HIV-1-induced inflammatory responses and leukocyte transmigration across the blood-brain barrier. *Blood*, **2008**, *111*, 2062-2072.
- [45] Pacher, P.; Batkai, S.; Kunos, G. The endocannabinoid system as an emerging target of pharmacotherapy. *Pharmacol. Rev.*, **2006**, *58*, 389-462.
- [46] Bovolenta, C.; Camorali, L.; Lorini, A.L.; Ghezzi, S.; Vicenzi, E.; Lazzarin, A.; Poli, G. Constitutive activation of STATs upon *in vivo* human immunodeficiency virus infection. *Blood*, **1999**, *94*, 4202-4209.
- [47] Poynard, T.; Yuen, M.F.; Ratziu, V.; Lai, C.L. Viral hepatitis C. *Lancet*, **2003**, *362*, 2095-2100.
- [48] Pileri, P.; Uematsu, Y.; Campagnoli, S.; Galli, G.; Falugi, F.; Petracca, R.; Weiner, A.L.; Houghton, M.; Rosa, D.; Grandi, G.; Abrignani, S. Binding of hepatitis C virus to CD81. *Science*, **1998**, *282*, 938-941.
- [49] Scarselli, E.; Ansuini, H.; Cerino, R.; Roccasecca, R.M.; Acali, S.; Filocamo, G.; Traboni, C.; Nicosia, A.; Cortese, R.; Vitelli, A. The human scavenger receptor class B type I is a novel candidate receptor for the hepatitis C virus. *Embo. J.*, **2002**, *21*, 5017-5025.
- [50] Bartosch, B.; Dubuisson, J.; Cosset, F.L. Infectious hepatitis C virus pseudoparticles containing functional E1-E2 envelope protein complexes. *J. Exp. Med.*, **2003**, *197*, 633-642.
- [51] Evans, M.J.; von Hahn, T.; Tschernig, D.M.; Syder, A.J.; Panis, M.; Wolk, B.; Hatzioannou, T.; McKeating, J.A.; Bieniasz, P.D.; Rice, C.M. Claudin-1 is a hepatitis C virus co-receptor required for a late step in entry. *Nature*, **2007**, *446*, 801-805.
- [52] Ploss, A.; Evans, M.J.; Gaysinskaya, V.A.; Panis, M.; You, H.; de Jong, Y.P.; Rice, C.M. Human occludin is a hepatitis C virus entry factor required for infection of mouse cells. *Nature*, **2009**, *457*, 882-886.
- [53] Gitter, A.H.; Wullstein, F.; Fromm, M.; Schulzke, J.D. Epithelial barrier defects in ulcerative colitis: characterization and quantification by electrophysiological imaging. *Gastroenterology*, **2001**, *121*, 1320-1328.
- [54] Marin, M.L.; Greenstein, A.J.; Geller, S.A.; Gordon, R.E.; Aufses, A.H., Jr. A freeze fracture study of Crohn's disease of the terminal ileum: changes in epithelial tight junction organization. *Am. J. Gastroenterol.*, **1983**, *78*, 537-547.
- [55] Schmitz, H.; Barmeyer, C.; Fromm, M.; Runkel, N.; Foss, H.D.; Bentzel, C.J.; Riecken, E.O.; Schulzke, J.D. Altered tight junction structure contributes to the impaired epithelial barrier function in ulcerative colitis. *Gastroenterology*, **1999**, *116*, 301-309.
- [56] Oshima, T.; Miwa, H.; Joh, T. Changes in the expression of claudins in active ulcerative colitis. *J. Gastroenterol. Hepatol.*, **2008**, *23 Suppl 2*, S146-150.
- [57] Alexandre, M.D.; Lu, Q.; Chen, Y.H. Overexpression of claudin-7 decreases the paracellular Cl⁻ conductance and increases the paracellular Na⁺ conductance in LLC-PK1 cells. *J. Cell Sci.*, **2005**, *118*, 2683-2693.
- [58] Van Itallie, C.; Rahner, C.; Anderson, J.M. Regulated expression of claudin-4 decreases paracellular conductance through a selective decrease in sodium permeability. *J. Clin. Invest.*, **2001**, *107*, 1319-1327.
- [59] Shorter, R.G.; Huizenga, K.A.; Spencer, R.J.; Guy, S.K. Inflammatory bowel disease. The role of lymphotoxin in the cytotoxicity of lymphocytes for colonic epithelial cells. *Am. J. Dig. Dis.*, **1972**, *17*, 689-696.
- [60] Kosiewicz, M.M.; Nast, C.C.; Krishnan, A.; Rivera-Nieves, J.; Moskaluk, C.A.; Matsumoto, S.; Kozaiwa, K.; Cominelli, F. Th1-type responses mediate spontaneous ileitis in a novel murine model of Crohn's disease. *J. Clin. Invest.*, **2001**, *107*, 695-702.
- [61] Reuter, B.K.; Pizarro, T.T. Mechanisms of tight junction dysregulation in the SAMPI/YitFc model of Crohn's disease-like ileitis. *Ann. N.Y. Acad. Sci.*, **2009**, *1165*, 301-307.
- [62] Vidrich, A.; Buzan, J.M.; Barnes, S.; Reuter, B.K.; Skaar, K.; Ilo, C.; Cominelli, F.; Pizarro, T.; Cohn, S.M. Altered epithelial cell lineage allocation and global expansion of the crypt epithelial stem cell population are associated with ileitis in SAMPI/YitFc mice. *Am. J. Pathol.*, **2005**, *166*, 1055-1067.
- [63] Paik, J.H.; Kollipara, R.; Chu, G.; Ji, H.; Xiao, Y.; Ding, Z.; Miao, L.; Tothova, Z.; Horner, J.W.; Carrasco, D.R.; Jiang, S.; Gilliland, D.G.; Chin, L.; Wong, W.H.; Castrillon, D.H.; DePinho, R.A. FoxOs are lineage-restricted redundant tumor suppressors and regulate endothelial cell homeostasis. *Cell*, **2007**, *128*, 309-323.
- [64] Tothova, Z.; Kollipara, R.; Huntly, B.J.; Lee, B.H.; Castrillon, D.H.; Cullen, D.E.; McDowell, E.P.; Lazo-Kallanian, S.; Williams, I.R.; Sears, C.; Armstrong, S.A.; Passegue, E.; DePinho, R.A.; Gilliland, D.G. FoxOs are critical mediators of hematopoietic stem cell resistance to physiologic oxidative stress. *Cell*, **2007**, *128*, 325-339.
- [65] Zhou, W.; Cao, Q.; Peng, Y.; Zhang, Q.J.; Castrillon, D.H.; DePinho, R.A.; Liu, Z.P. FoxO4 inhibits NF- κ B and protects mice against colonic injury and inflammation. *Gastroenterology*, **2009**, *137*, 1403-1414.
- [66] Gitter, A.H.; Bendfeldt, K.; Schulzke, J.D.; Fromm, M. Leaks in the epithelial barrier caused by spontaneous and TNF- α -induced single-cell apoptosis. *Faseb. J.*, **2000**, *14*, 1749-1753.
- [67] Mankertz, J.; Tavalali, S.; Schmitz, H.; Mankertz, A.; Riecken, E.O.; Fromm, M.; Schulzke, J.D. Expression from the human occludin promoter is affected by tumor necrosis factor alpha and interferon gamma. *J. Cell. Sci.*, **2000**, *113*, 2085-2090.
- [68] Mankertz, J.; Amasheh, M.; Krug, S.M.; Fromm, A.; Amasheh, S.; Hillenbrand, B.; Tavalali, S.; Fromm, M.; Schulzke, J.D. TNF α up-regulates claudin-2 expression in epithelial HT-29/B6 cells via phosphatidylinositol-3-kinase signaling. *Cell Tissue Res.*, **2009**, *336*, 67-77.
- [69] Fries, W.; Muja, C.; Crisafulli, C.; Cuzzocrea, S.; Mazon, E. Dynamics of enterocyte tight junctions: effect of experimental colitis and two different anti-TNF strategies. *Am. J. Physiol.*, **2008**, *294*, G938-947.
- [70] Aslan, A.; Triadafilopoulos, G. Fish oil fatty acid supplementation in active ulcerative colitis: a double-blind, placebo-controlled, crossover study. *Am. J. Gastroenterol.*, **1992**, *87*, 432-437.
- [71] Hawthorne, A.B.; Daneshmand, T.K.; Hawkey, C.J.; Belluzzi, A.; Everitt, S.J.; Holmes, G.K.; Malkinson, C.; Shaheen, M.Z.; Willars, J.E. Treatment of ulcerative colitis with fish oil supplementation: a prospective 12 month randomised controlled trial. *Gut*, **1992**, *33*, 922-928.
- [72] Kitsukawa, Y.; Saito, H.; Suzuki, Y.; Kasanuki, J.; Tamura, Y.; Yoshida, S. Effect of ingestion of eicosapentaenoic acid ethyl ester on carrageenan-induced colitis in guinea pigs. *Gastroenterology*, **1992**, *102*, 1859-1866.
- [73] Li, Q.; Zhang, Q.; Zhang, M.; Wang, C.; Zhu, Z.; Li, N.; Li, J. Effect of n-3 polyunsaturated fatty acids on membrane microdomain localization of tight junction proteins in experimental colitis. *FEBS J.*, **2008**, *275*, 411-420.
- [74] Fuller, R. Probiotics in man and animals. *J. Appl. Bacteriol.*, **1989**, *66*, 365-378.
- [75] Mennigen, R.; Nolte, K.; Rijcken, E.; Utech, M.; Loeffler, B.; Senninger, N.; Bruewer, M. Probiotic mixture VSL#3 protects the epithelial barrier by maintaining tight junction protein expression and preventing apoptosis in a murine model of colitis. *Am. J. Physiol.*, **2009**, *296*, G1140-1149.
- [76] Takahashi, A.; Kondoh, M.; Masuyama, A.; Fujii, M.; Mizuguchi, H.; Horiguchi, Y.; Watanabe, Y. Role of C-terminal regions of the C-terminal fragment of *Clostridium perfringens* enterotoxin in its interaction with claudin-4. *J. Control. Release*, **2005**, *108*, 56-62.

