




Contents lists available at ScienceDirect

Molecular Genetics and Metabolism

journal homepage: [www.elsevier.com/locate/ymgme](http://www.elsevier.com/locate/ymgme)



# Functional analysis of genetic variations in the 5'-flanking region of the human MDR1 gene

Mayumi Saeki, Kouichi Kurose, Ryuichi Hasegawa, Masahiro Tohkin \*

Division of Medicinal Safety Science, National Institute of Health Sciences, 1-18-1 Kamiyoga, Setagaya-ku, Tokyo 158-8501, Japan

## ARTICLE INFO

### Article history:

Received 26 August 2010

Accepted 26 August 2010

Available online 19 September 2010

### Keywords:

5'-flanking region

Multidrug resistance 1

Nuclear receptors

P-glycoprotein

Single nucleotide polymorphism

## ABSTRACT

P-glycoprotein (P-gp), the product of the *MDR1* gene, shows large interindividual variations in expression, which leads to differences in the pharmacokinetics of the substrate drugs. The functions of single nucleotide polymorphisms located in the nuclear receptor-responsive element of the 5'-flanking region in the human *MDR1* gene were analyzed in order to clarify the mechanism underlying the interindividual variation in P-gp expression. Electrophoretic mobility shift assays revealed that the –7833C>T substitution in the nuclear receptor-responsive region of *MDR1* decreases the binding affinities of four nuclear receptors to their responsive elements: vitamin D receptor (VDR), thyroid hormone receptor (TR), constitutive androstane receptor (CAR), and pregnane X receptor (PXR). A reporter gene assay revealed that the C-to-T substitution at –7833 also reduces the transcriptional activation of *MDR1* by VDR, TR $\beta$ , CAR, and PXR. However, another SNP (–1211T>C substitution), which results in the formation of a xenobiotic responsive element-like sequence and a hypoxia responsive element-like sequence, failed to affect the aryl hydrocarbon receptor-dependent and hypoxia-induced transcriptional activation of *MDR1*. Although the frequency of the –7833C>T substitution in *MDR1* is relatively low, the SNP is crucial because it may alter the pharmacokinetics of P-gp substrates in a small subset of the population.

© 2010 Elsevier Inc. All rights reserved.

## 1. Introduction

P-glycoprotein (P-gp) transports a wide variety of compounds, including foreign xenobiotics and endogenous substrates, to the outside of cells [1,2]. P-gp is encoded by the human *MDR1* gene and is expressed in various physiological barriers such as intestinal epithelium [3,4], kidney tubules cells [3], the liver [3], and the capillary endothelium of the central nervous system [5]. It also plays an important role in pharmacokinetic processes such as drug absorption [1,6–8], renal secretion [9], biliary excretion [10], and brain distribution [11–13]. The large interindividual variations in P-gp expression and activity levels [14–17] suggest that the systemic exposure level and tissue concentrations of drugs that are P-gp substrates vary depending on the subject [15,18]. The pharmacokinetic differences ultimately lead to interindividual variation in drug efficacy and adverse reactions. Therefore, P-gp expression levels and activity are crucial factors in drug efficacy and safety.

The results of several studies suggest that the basis of interindividual variation in P-gp expression and activity resides in the presence of single nucleotide polymorphisms (SNPs) in the coding region of *MDR1* [2,19–22]. Hoffmeyer et al. showed that 3435C>T, a synonymous SNP in exon 26, is associated with reduced P-gp

expression in the duodenum and increased plasma levels of digoxin, a typical P-gp substrate, following its oral administration to healthy volunteers [23]. Wang et al. also reported that 3435C>T appears to affect the allelic variation of *MDR1* expression in the liver via a change in mRNA stability [24]. In addition to digoxin, other studies have reported that the T allele of 3435C>T is related to higher plasma concentrations of cyclosporine A. While these reports suggest the importance of *MDR1* exonic SNPs in the regulation of P-gp expression, conflicting results have been reported [18,23,25–27].

In addition to SNPs in the coding region, there are variations in the 5'-flanking region that could affect *MDR1* gene transcription and mRNA expression levels. Haplotypes in the *MDR1* transcriptional regulatory region suggest the existence of functional haplotypes that could alter P-gp expression [17,28,29]. In these studies, the –1789G>A haplotype, alone or in combination with –145C>G, was associated with decreased P-gp expression. However, the reported effects on P-gp expression of haplotypes carrying –129T>C and two other linked SNPs were contradictory, showing reduction and enhancement [27,30,31]. These reports suggest that functional SNPs in the 5'-flanking region have not been fixed, and the relationship between these SNPs and the transcriptional factors that regulate the *MDR1* gene transcription is unknown. Therefore, the mechanism underlying interindividual variations in the expression levels of P-gp remains unclear.


Geick et al. found that the induction of *MDR1* mRNA by rifampicin is mediated by pregnane X receptor (PXR), which binds to direct

\* Corresponding author. Fax: +81 3 3700 9788.

E-mail address: [tohkin@nihs.go.jp](mailto:tohkin@nihs.go.jp) (M. Tohkin).

repeat sequences separated by four bases (DR4) located  $-7.9$  to  $-7.8$  kbp upstream of the transcription start site [32]. This region contains several DR sequences and functions as the enhancer region in *MDR1* [32]. The constitutive androstane receptor (CAR) also induces *MDR1* mRNA expression by binding to several DR4s located in the

same region [33]. Recently, we reported that the thyroid hormone receptor  $\beta$  (TR $\beta$ ) and vitamin D receptor (VDR) regulate the expression of *MDR1* by binding to several DRs located in this region [34,35]. A SNP,  $-7833\text{C}>\text{T}$ , has been reported within one of the half-sites (Hs), a pair of which composes a DR or ER (AGTTCA $\rightarrow$ AGTTTA,



**Fig. 1.** The C-to-T substitution at  $-7833$  affects the binding affinity of nuclear receptors to ER8 and/or DR4(II) in the *MDR1* gene. (A) Oligonucleotide sequences used for the electrophoretic mobility shift assay (EMSA). Several half-sites (designated as Hs1 to 8 in this study) are boxed, and arrows indicate the direction of the half-site. The C-to-T substitution at  $-7833$  is located within Hs6. The numbers are in reference to the transcriptional start site at +1. Schematic representations of probes containing the  $-7833\text{C}$  (ER8-WT and DR4(II)-WT) or  $-7833\text{T}$  (ER8-VT and DR4(II)-VT) allele are shown. Only nucleotides that differ from the wild-type are shown as letters; asterisks represent unchanged nucleotides. (B) EMSA was performed using *in vitro* translated VDR and RXR $\alpha$ . The FITC-labeled probe was incubated with VDR and RXR $\alpha$  as described in the Materials and methods. The complexes were resolved by electrophoresis on a 6% Long Ranger gel. This protocol was also used for the EMSAs shown in C–G. (C) EMSA was performed using *in vitro* translated TR $\beta$  and RXR $\alpha$ . (D) EMSA was performed using *in vitro* translated CAR and RXR $\alpha$ . (E) EMSA was performed using *in vitro* translated CAR and RXR $\alpha$ . (F) EMSA was performed using *in vitro* translated PXR and RXR $\alpha$ . (G) We carried out the translation reaction using the empty vector, and performed EMSA using the reaction product instead of *in vitro* translated nuclear receptors, as the negative control.

Fig. 1A) [9,36]. Although the frequency of this substitution is very low (0.002), there is the possibility the  $-7833\text{C}>\text{T}$  variant may influence the transcription of the *MDR1* gene.

*MDR1* mRNA expression was also induced by 2,3,7,8-tetrachlorodibenzo-p-dioxin (aryl hydrocarbon receptor (AhR) ligand) in three of seven primary human hepatocytes [37], an indication of the interindividual variation in AhR-induced expression, although the AhR responsive element (xenobiotics responsive element; XRE) in *MDR1* has not been identified. The  $-1211\text{T}>\text{C}$  (rs28746504,  $-1910$  in Taniguchi et al., [17] and  $-1017$  in Takane et al. [29]) SNP has been found in Japanese and Caucasian populations at allelic frequencies of 0.08–0.1 (Japanese) and 0.016 (Caucasian). The T-to-C substitution at nucleotide  $-1211$  results in an XRE-like sequence (possible XRE, TGGTGTG>TGGCGTG, Fig. 3A) that we hypothesize may cause the interindividual variability in the response to TCDD. In addition to AhR-induced *MDR1* mRNA expression, hypoxia-inducible factor-1 (HIF-1) regulates *MDR1* gene transcription [38]. The  $-1211\text{T}>\text{C}$  substitution also results in an HIF-1-responsive element (hypoxia responsive elements; HRE)-like sequence (possible HRE, GTGTG>GCGTG, Fig. 3A). However, it was unknown if HIF-1 could bind the candidate HRE in addition to the known HRE located between  $-49$  and  $-45$  ( $-53$  and  $-49$  in this study) in the promoter region of *MDR1*.

In order to clarify the functional significance of variants in the transcriptional regulatory region of *MDR1* gene, the effect of two SNPs ( $-7833\text{C}>\text{T}$  and  $-1211\text{T}>\text{C}$ ) on the binding properties of nuclear receptors and on the transcriptional activity of *MDR1* was evaluated using electrophoretic mobility shift assay (EMSA) and a luciferase-reporter gene assay, respectively. The  $-7833\text{C}>\text{T}$  substitution resulted in a decrease in nuclear receptor binding and transcriptional activation *in vitro*, while the  $-1211\text{T}>\text{C}$  substitution did not produce an effect. These results suggest that the  $-7833\text{C}>\text{T}$  substitution influences the regulation of *MDR1* mRNA levels through PXR, CAR, TR $\beta$ , and VDR.

## 2. Materials and methods

### 2.1. Chemicals

Rifampicin and 3-methylcholanthrene (3MC) were purchased from Wako Pure Chemicals (Osaka, Japan).  $\text{CoCl}_2$ , 3,3',5'-triiodo-L-thyronine (T3), and 1 $\alpha$ ,25-dihydroxyvitamin D $_3$  (1,25-(OH) $_2$ D $_3$ ) were purchased from Sigma-Aldrich (St. Louis, MO, USA). All chemicals, except T3 and  $\text{CoCl}_2$ , were dissolved in dimethyl sulfoxide (DMSO). T3 and  $\text{CoCl}_2$  were dissolved in 0.2 M NaOH and water, respectively.

### 2.2. Plasmid constructs

The expression plasmids, pDEST12.2-hPXR, pEF6/V5-hVDR, pCMVTNT-hVDR, and pcDNA3.1-TR $\beta$ , were previously constructed in our laboratory [34,35,39]. The expression plasmid encoding human RXR $\alpha$  cDNA (pcDNA3.1-hRXR $\alpha$ ) was a generous gift from Dr. Shuichi Koizumi (Yamanashi University, Japan). pCMVTNT-hTR $\beta$  was constructed by ligating the *EcoRI*/*NotI* fragment from pcDNA3.1-TR $\beta$  into the pCMVTNT expression plasmid (Promega, Madison, WI, USA), which was digested with *EcoRI* and *NotI*. The pEGFP-hCAR was a generous gift from Dr. Hideto Jinno (National Institute of Health Sciences, Japan) [40]. The nucleotide at the position 540 of pEGFP-hCAR differed from the reference sequence (NM\_005122) (540C>T, synonymous substitution) [40]. Thus, wild-type sequence was introduced into the pEGFP-hCAR using a QuikChange Site-Directed Mutagenesis Kit (Stratagene, La Jolla, CA, USA). pEGFP-hCAR was then digested with *XhoI* and *EcoRI*, and the resulting fragment was ligated into the pcDNA3.1/Zeo expression plasmid (Invitrogen, Carlsbad, CA, USA), which was digested with *XhoI* and *EcoRI*. This expression plasmid (pcDNA3.1-hCAR) was digested with *XhoI* and *PmeI*, and the resulting fragment was ligated into the pCMVTNT expression plasmid,

which was digested with *XhoI* and *SmaI* (pCMVTNT-hCAR). The pDEST12.2-hPXR, pEF6/V5-hVDR, pcDNA3.1-TR $\beta$ , and pcDNA3.1-hCAR plasmids were used for transfections. The pDEST12.2-hPXR, pCMVTNT-hVDR, pCMVTNT-hTR $\beta$ , pCMVTNT-hCAR, and pcDNA3.1-hRXR $\alpha$  plasmids were used for *in vitro* synthesis.

The luciferase reporter gene plasmids, pMD10082L, pMD\*824 $\Delta$ 90L, and pMD457L were previously constructed in our laboratory [34]. The pMD2912L plasmid was constructed by deleting the *NsiI* fragment from pMD10082L. Mutations were introduced at positions  $-7833$  and  $-1211$  of pMD\*824 $\Delta$ 90L-WT and pMD2912L-WT to create pMD\*824 $\Delta$ 90L-VT and pMD2912L-VT, respectively, using a QuikChange Multi Site-Directed Mutagenesis Kit (Stratagene), according to the manufacturer's instructions, with the following oligonucleotides:

$-7833\text{C}>\text{T}$ -SNP: 5'-GCTCCTGGGAGAGAGTTTATTGAGATTAACAAG-3'

$-1211\text{T}>\text{C}$ -SNP: 5'-CAGGAGAATGGCGTGAACCCGGGAG-3'.

The pGL2A8-2504 [41] plasmid, which contains the XRE of the Syrian hamster CYP2A8, was digested with *KpnI* and *HindIII*. The resulting fragment was ligated to the *KpnI*/*HindIII* site of the firefly luciferase rapid response reporter vector pGL4.12 (Promega). This plasmid (pGL4.12-2A8) was used as a positive control.

### 2.3. Electrophoretic mobility shift assay (EMSA)

TNT T7 Quick Coupled Transcription/Translation Systems (Promega) were used for *in vitro* synthesis of human RXR $\alpha$  protein from pcDNA3.1-hRXR $\alpha$ , according to the manufacturer's instructions. TNT SP6 Quick Coupled Transcription/Translation Systems (Promega) were used for *in vitro* synthesis of human VDR, TR $\beta$ , CAR, and PXR proteins from pCMVTNT-hVDR, pCMVTNT-hTR $\beta$ , pCMVTNT-hCAR, and pDEST12.2-hPXR, respectively, according to the manufacturer's instructions. The plus strand sequences of probes used in the EMSAs are shown in Fig. 1A. The oligonucleotides were purchased from Sigma Genosys (Hokkaido, Japan) and equal amounts of complementary strands were annealed. The reaction mixture was prepared as follows: a 2.5  $\mu\text{L}$  aliquot of the *in vitro* translated proteins (nuclear receptor alone, or mixed at a ratio of 1:1) or unprogrammed reticulocyte lysate was incubated for 20 min at room temperature with 1  $\mu\text{L}$  of 5 $\times$  binding buffer (15 mM MgCl $_2$ , 0.5 mM EDTA, 2.5 mM dithiothreitol (DTT), 50% glycerol and 100 mM HEPES, pH 7.75), 0.5  $\mu\text{L}$  of 1 mg/mL poly(dI-dC) (GE Healthcare, Little Chalfont, UK), and 0.5  $\mu\text{L}$  of 0.33  $\mu\text{M}$  5'-fluorescein isothiocyanate (FITC)-labeled double stranded oligonucleotide probe. The protein-DNA complexes were resolved by electrophoresis on 6% non-denaturing Long Ranger gels (Lonza, Rockland, ME) run in 0.5 $\times$ TBE (44.5 mM Tris, 44.5 mM boric acid, and 1.25 mM EDTA) at 500 V constant voltage, and visualized and quantified on a slab gel DNA sequencer (DSQ-2000L; Shimadzu Co., Kyoto, Japan).

### 2.4. Cell culture

Caco-2 cells, a human colon adenocarcinoma cell line, were obtained from the American Type Culture Collection (Manassas, VA, USA). Caco-2 cells were cultured in low glucose Dulbecco's modified Eagle's medium (DMEM, Sigma-Aldrich) supplemented with 10% heat-inactivated fetal bovine serum (FBS), 100 U/mL penicillin G/100  $\mu\text{g}$ /mL streptomycin (Gibco-Invitrogen, Carlsbad, CA, USA), and 1 $\times$  MEM non-essential amino acids solution (Gibco-Invitrogen) at 37  $^{\circ}\text{C}$  under 5% CO $_2$ -95% air.

### 2.5. Transfection and luciferase reporter gene assays

Caco-2 cells were seeded into 96-well plates ( $1.7 \times 10^4$  cells/well), grown overnight, and transiently transfected using HilyMax (at a ratio of DNA to HilyMax of 1:5; Dojindo Laboratories, Kumamoto, Japan)

according to the manufacturer's instructions with 10 ng/well of the indicated expression plasmid, 100 ng/well of the indicated luciferase reporter plasmid, and 10 ng/well of the *Renilla* luciferase reporter plasmid, pGL4.74 [hRluc/TK] (Promega) to normalize the transfection efficiency. After 24 h, the medium was replaced by phenol red-free DMEM (Gibco-Invitrogen) supplemented with 10% dextran-coated charcoal-stripped FBS (Hyclone Laboratories, Logan, UT, USA) containing 1  $\mu$ M 3MC, 150  $\mu$ M CoCl<sub>2</sub>, 25 nM 1,25-(OH)<sub>2</sub>D<sub>3</sub>, 50 nM T<sub>3</sub>, 10  $\mu$ M rifampicin, or vehicle (DMSO, water, or 0.2 mM NaOH) for 3.5 h except for CoCl<sub>2</sub>. CoCl<sub>2</sub> treatment was performed for 6 h. Firefly and *Renilla* luciferase activities were measured using the Dual-Glo Luciferase Assay System (Promega) according to the manufacturer's instructions and a luminometer (Wallac 1420 ARVO sx Multilabel Counter, Perkin-Elmer Life Sciences, Boston, MA, USA). Firefly luciferase activity was normalized to *Renilla* luciferase activity (Luc activity). The inducibility (fold induction) was calculated as the ratio of luciferase activity of ligand-treated cells to that of control cells. As for CAR, fold transactivation was calculated as the ratio of luciferase activity of CAR-transfected cells to that of pcDNA3.1 transfected cells. The results were presented as the mean  $\pm$  standard deviation (S.D.) of at least four independent experiments. The statistical analysis was performed using two-tailed, unpaired t-tests with a significance level of  $P < 0.05$ .

### 3. Results

#### 3.1. C-to-T substitution at –7833 of MDR1 decreases binding affinity of nuclear receptors to ER8 and/or DR4(II)

Nuclear receptors, such as VDR, TR $\beta$ , CAR, and PXR reportedly up-regulate *MDR1* expression through the response region located 7880 to 7810 bp upstream of the *MDR1* gene. This region contains several half-sites, a pair of which composes a DR or ER (Hs1–8 in Fig. 1A). The previously reported variation –7833C>T is located within the Hs6 half-site (Fig. 1A). The effect of the C-to-T substitution at –7833 on the binding affinity of nuclear receptors to the response region was examined with EMSA using *in vitro* translated VDR, TR $\beta$ , PXR, CAR and RXR $\alpha$ . ER8 and DR4(II) oligonucleotide probes containing either –7833C (WT) or –7833T (VT) were used because Hs6 is involved in both ER8 and DR4(II) (Fig. 1A). The probes used for the EMSA are summarized in Fig. 1A. As described in previous reports, VDR/RXR $\alpha$  forms DNA–protein complexes with ER8-WT and DR4(II)-WT, with a weaker binding affinity for DR4(II) than for ER8. Introduction of the variant into the probes (DR4(II)-VT and ER8-VT) decreased the formation of VDR/RXR $\alpha$  and probe complexes (Fig. 1B). As described previously, the DNA–TR $\beta$ /RXR $\alpha$  complex was formed using DR4(II)-WT, but not with ER8-WT (Fig. 1C). The DR4(II)-TR $\beta$ /RXR $\alpha$  complex significantly decreased when the variant type probe (DR4(II)-VT) was used. It has been reported that CAR binds to Hs6 as a monomer [33].

Consistent with this report, the DNA–CAR monomer complex was formed with either ER8-WT or DR4(II)-WT, and the C-to-T substitution at –7833 (ER8-VT and DR4(II)-VT) caused a decrease in band formation (Fig. 1D). The CAR/RXR $\alpha$  DNA–protein complex was also formed when either ER8-WT or DR4(II)-WT was used as a probe, although the binding affinity of ER8-CAR/RXR $\alpha$  was weak (Fig. 1E). Complex formation between CAR/RXR $\alpha$  and the probes decreased when the WT probes were replaced with ER8-VT and DR4(II)-VT (Fig. 1E). PXR/RXR $\alpha$  DNA–protein complexes formed using both ER8-WT and DR4(II)-WT produced weak bands, which disappeared when ER8-VT and DR4(II)-VT were used as probes (Fig. 1F). These results indicate that the C-to-T substitution at –7833 leads to decreased binding affinity of ER8 and/or DR4(II) for VDR/RXR $\alpha$ , TR $\beta$ /RXR $\alpha$ , CAR, CAR/RXR $\alpha$ , and PXR/RXR $\alpha$ .

#### 3.2. C-to-T substitution at –7833 of MDR1 affects the inducibility by 1,25-(OH)<sub>2</sub>D<sub>3</sub>, T<sub>3</sub>, CAR, and PXR, but does not affect the inducibility by rifampicin

The EMSA experiments revealed that the C-to-T substitution at –7833 results in decreased binding affinity of VDR/RXR $\alpha$ , TR $\beta$ /RXR $\alpha$ , CAR, CAR/RXR $\alpha$ , and PXR/RXR $\alpha$  for ER8 and/or DR4(II). To investigate the effect of the substitution on VDR-, TR $\beta$ -, CAR-, and PXR-induced *MDR1* transcription, –7833C>T was introduced into pMD\*824 $\Delta$ 90L reporter plasmid, which contains the nuclear receptor-response region (Fig. 2A), and a luciferase reporter assay was performed using the Caco-2 intestinal epithelial cell line. In order to exclude the interaction of other nuclear receptors, we used the Caco-2 because Caco-2 cells express the nuclear receptors at relatively lower levels. As described in previous reports [34,35], 1,25-(OH)<sub>2</sub>D<sub>3</sub> (VDR ligand) or triiodothyronine (T<sub>3</sub>, TR $\beta$  ligand) increased the luciferase activity in Caco-2 cells that were co-transfected with VDR or TR $\beta$  expression plasmid (Figs. 2B and C) although VDR or TR $\beta$  alone could not activate the transcriptional activity without the ligand (data not shown). The C-to-T substitution resulted in significantly reduced transcriptional activation by 1,25-(OH)<sub>2</sub>D<sub>3</sub> (Fig. 2B,  $P = 0.0015$ , unpaired t-test, two-tailed). The substitution also led to a slight decrease in T<sub>3</sub>-induced transcriptional activation, but this decrease was not significant (Fig. 2C,  $P = 0.0688$ , unpaired t-test, two-tailed). Co-transfection of a CAR expression plasmid led to increased transcriptional activity of the reporter gene compared with the mock transfection control group (fold transactivation in Fig. 2D) due to the constitutive activity of CAR. The C-to-T substitution also partially reduced constitutive CAR transactivity (Fig. 2D,  $P = 0.0326$ , unpaired t-test, two-tailed). Rifampicin increased transactivation of the reporter gene following transfection with the PXR expression plasmid. The –7833C>T substitution did not affect the fold induction produced by rifampicin (Fig. 2E). However, the substitution reduced both the rifampicin-dependent and -independent transcriptional activation by PXR (Fig. 2F). These results indicated that –7833C>T in the *MDR1* gene reduced the transcriptional activation which is induced by 1,25-(OH)<sub>2</sub>D<sub>3</sub>.

**Fig. 2.** Effect of –7833C>T on the transactivation of *MDR1* by VDR, TR $\beta$ , CAR, and PXR. (A) The C-to-T mutation at –7833 was introduced into a reporter gene plasmid containing the 5' upstream region from –7880 to –7810 bp of *MDR1* (pMD\*824 $\Delta$ 90L). Several half-sites (designated as Hs1 to Hs8) are boxed and arrows indicate the direction of the half-site. The C-to-T substitution at –7833 is located within Hs6. The numbers are in reference to the transcriptional start site at +1. In the variant construct (pMD\*824 $\Delta$ 90L-VT), only the nucleotide that differs from the wild-type (pMD\*824 $\Delta$ 90L-WT) is shown as a letter; asterisks represent unchanged nucleotides. (B) Caco-2 cells were co-transfected with the indicated luciferase reporter plasmid, VDR expression plasmid, and the *Renilla* luciferase reporter plasmid, and were subsequently treated with DMSO or 1,25-(OH)<sub>2</sub>D<sub>3</sub>. Firefly luciferase activity was normalized to *Renilla* luciferase activity, and the fold induction was calculated as the ratio of luciferase activity in 1,25-(OH)<sub>2</sub>D<sub>3</sub>-treated cells to that of DMSO-treated cells. Each value represents the mean  $\pm$  S.D. of independent seven experiments. Statistical analysis was performed using a two-tailed, unpaired t-test, and a statistically significant difference, as compared with the wild-type, is indicated by an asterisk (\*\* $P < 0.005$ ). (C) The luciferase assay was performed as described in (B) except that the TR $\beta$  expression plasmid, T<sub>3</sub>, and 0.2 mM NaOH were used instead of the VDR expression plasmid, 1,25-(OH)<sub>2</sub>D<sub>3</sub>, and DMSO, respectively. The fold induction was calculated as the ratio of luciferase activity in T<sub>3</sub>-treated cells to that in 0.2 mM NaOH-treated cells. Each value represents the mean  $\pm$  S.D. of independent six experiments. Statistical analysis was performed using the two-tailed, unpaired t-test, ( $P = 0.069$ ). (D) Luciferase activity was analyzed as described in the Materials and methods. Firefly luciferase activity was normalized to *Renilla* luciferase activity, and the fold transactivation was calculated as the ratio of luciferase activity of CAR-transfected cells to that of control plasmid-transfected cells. Each value represents the mean  $\pm$  S.D. of independent four experiments. Statistical analysis was performed using the two-tailed, unpaired t-test, and a statistically significant difference, as compared with the wild-type, is indicated by asterisk (\* $P < 0.05$ ). (E) The luciferase assay was performed as described in (B) except that PXR expression plasmid and rifampicin were used instead of VDR expression plasmid and 1,25-(OH)<sub>2</sub>D<sub>3</sub>, respectively. The fold induction was calculated as the ratio of luciferase activity in rifampicin-treated cells to that in DMSO-treated cells. Each value represents the mean  $\pm$  S.D. of independent five experiments. (F) These data are from the experiments described in (E), but are expressed in each treatment group. The firefly luciferase activity was normalized for transfection efficiency, using the activity of co-transfected *Renilla* luciferase, and represented as Luc activity.





and T3. The –7833C>T also reduced CAR and PXR-dependent transcriptional activities.

3.3. XRE- and HRE-like sequences, resulting from a –1211T>C substitution in MDR1, are not functional

The T-to-C substitution at –1211 from the transcriptional start site of the human *MDR1* gene (–1211T>C, rs28746504) results in the formation of a XRE-like sequence (TGGTGTG>TGGCGTG, Fig. 3A). To examine the functionality of the XRE-like sequence, –1211T>C was

introduced into a reporter plasmid containing the 5' upstream region from –2912 to +117 bp of the *MDR1* gene and transfected into Caco-2 cells. The pGL4.12-2A8 construct (2A8), which contains the functional XRE, was used as the positive control [41]. As shown in Fig. 3B, 3MC (AhR ligand) did not affect transcriptional activity of Caco-2 cells transfected with either the wild-type (pMD2912L-WT) or mutated (pMD2912L-VT) construct, although the luciferase activity of the positive control reporter construct (2A8) was increased approximately 5-fold by 3MC, indicating that the XRE-like sequence containing –1211T>C is not functional.





**Fig. 3.** Effect of  $-1211\text{T}>\text{C}$  on the transactivation of the *MDR1* gene by AhR and HIF. (A) Schematic representation of reporter plasmids containing the  $-1211\text{T}$  (pMD2912L-WT) or  $-1211\text{C}$  (pMD2912L-VT) allele. The pMD2912L plasmid was constructed by deleting the *NsiI* fragment in pMD10082L. The numbers are in reference to the transcriptional start site at +1. Xenobiotic response element (XRE)-like and hypoxia response element (HRE)-like sequences (TGCGGTG and GCGTG, respectively) resulting from the T-to-C substitution at  $-1211$  are boxed. pMD457L was constructed by deleting the *SphI* fragment from pMD10082L. (B) Caco-2 cells were co-transfected with the indicated luciferase reporter plasmid and the *Renilla* luciferase reporter plasmid, and were subsequently treated with DMSO or 3MC as described in Materials and methods. Firefly luciferase activity was normalized to *Renilla* luciferase activity, and the fold induction was calculated as the ratio of luciferase activity in 3MC-treated cells to that in DMSO-treated cells. As a positive control, pGL4.12-2A8 (2A8), which contains the functional XRE of Syrian hamster CYP2A8, was used. Each value represents the mean  $\pm$  S.D. of four independent experiments. (C) Caco-2 cells were co-transfected with the indicated luciferase reporter plasmid and the *Renilla* luciferase reporter plasmid, and then replaced by media with or without  $\text{CoCl}_2$  as described in Materials and methods. Firefly luciferase activity was normalized to *Renilla* luciferase activity, and the fold induction was calculated as the ratio of luciferase activity in  $\text{CoCl}_2$ -treated cells to that of untreated cells. The pMD457L, which contains the functional HRE but does not contain the HRE-like sequence, was used as a control. Each value represents the mean  $\pm$  S.D. of four independent experiments.

The T-to-C substitution at  $-1211$  also results in the formation of an HRE-like sequence (GTGTG>GCGTG, Fig. 3A). The effect of the  $-1211\text{T}>\text{C}$  mutation on hypoxia responsiveness was measured by transfecting Caco-2 cells with the pMD2912L-WT or pMD2912L-VT. The cells were cultured with or without  $\text{CoCl}_2$ , a hypoxia-mimetic agent [38]. As shown in Fig. 3C,  $\text{CoCl}_2$ -dependent inducibility of the wild-type construct (pMD2912L-WT) was similar to that of the variant construct (pMD2912L-VT). Moreover, the inducibility of the wild-type construct, which includes the HRE-like sequence and the known HRE between  $-53$  and  $-49$ , was also similar to that of pMD457L, which contains the reported functional HRE alone [38]. Therefore, the  $-1211\text{T}>\text{C}$  substitution did not affect the hypoxic response, and the HRE-like sequence is also non-functional.

#### 4. Discussion

Many studies have tried to clarify the relationship between P-gp expression levels and SNPs located in the coding and transcriptional regulatory regions of *MDR1* [2,19,21–23,28,42]. However, most of these studies have not explored the functionality of these SNPs based on the nuclear receptor-responsive elements that play important roles in the

expression of *MDR1* mRNA. Therefore, the proposed functions of SNPs in the 5'-flanking region, such as  $-129\text{T}>\text{C}$ , remain controversial [27,30,31]. In this study, we analyzed the SNP located in the nuclear receptor-responsive region of *MDR1* and the SNP that results in conversion to a potential nuclear receptor-responsive element. The  $-7833\text{C}>\text{T}$  substitution, located in the nuclear receptor-responsive region, was found to significantly decrease the binding affinity of VDR/RXR $\alpha$ , TR $\beta$ /RXR $\alpha$ , CAR, CAR/RXR $\alpha$ , and PXR/RXR $\alpha$  (Fig. 1). In addition, the  $-7833\text{C}>\text{T}$  substitution reduced the transcriptional activation induced by  $1,25\text{-(OH)}_2\text{D}_3$  and T3, and suppressed CAR and PXR-dependent transcriptional activities *in vitro* (Fig. 2).

Recently, we reported that the relative binding ability of VDR/RXR $\alpha$  to the elements located in this region is: DR4(I)>DR3>ER8 (Mdc3)>DR4(III)>DR4(II), and DR4(I), with ER8(Mdc3), and DR4 (III) the main players in VDR-mediated *MDR1* induction [35]. Since  $-7833\text{C}>\text{T}$  is involved in ER8(Mdc3) (Fig. 1A), and ER8(Mdc3) contributes to VDR-mediated *MDR1* induction [35], it is possible that the  $-7833\text{C}>\text{T}$  substitution decreases VDR-mediated *MDR1* mRNA induction through ER8(Mdc3). We also previously indicated that TR $\beta$ /RXR $\alpha$  bound to several DRs in the order: DR4(I)>DR4(II)>DR3 $\approx$ DR4(III), and that every direct repeat contributes to TR $\beta$ -

mediated MDR1 induction [34]. Since  $-7833\text{C}>\text{T}$  is located in DR4(II) (Fig. 1A), and TR $\beta$ /RXR $\alpha$  can bind to DR4(II) at a relative high affinity [34], it is reasonable that the  $-7833\text{C}>\text{T}$  substitution could affect TR $\beta$ -mediated MDR1 mRNA induction through DR4(II). Burk et al. reported that CAR binds DR4(I) and DR4(III) as a heterodimer with RXR $\alpha$ , and to the 5' half-site of DR4(II) (designated as Hs6 in Fig. 1A) as a CAR monomer, and suggested that DR4(I) and the 5' half-site of DR4(II) (Hs6) were important elements for the CAR-mediated MDR1 induction [33]. In this study, the CAR monomer formed a complex with Hs6 both in ER8(MdC3) and DR4(II), and CAR/RXR $\alpha$  formed a complex with DR4(II) (Figs. 1D and E). These results and those of Burk indicate that Hs6 contributes to the CAR-mediated MDR1 mRNA induction. Since  $-7833\text{C}>\text{T}$  is found in Hs6 and DR4(II) (Fig. 1A), the  $-7833\text{C}>\text{T}$  substitution could decrease CAR-mediated MDR1 mRNA induction (Fig. 2D) even if the CAR monomer or CAR/RXR $\alpha$  contributes to MDR1 mRNA induction. Geick et al. reported that PXR/RXR $\alpha$  bind to DR4(I), DR4(II), and DR4(III) in the nuclear receptor-responsive elements of MDR1, with the highest affinity for DR4(III). DR4(I) is involved in rifampicin-mediated induction by PXR [32]. They suggest that DR4(II) does not contribute to the rifampicin-mediated induction by PXR even if the PXR/RXR $\alpha$  can bind DR4(II). In our study, PXR/RXR $\alpha$  formed a weak complex with DR4(II), and the  $-7833\text{C}>\text{T}$  substitution decreased the binding affinity of PXR/RXR $\alpha$  to DR4(II) (Fig. 1F). The  $-7833\text{C}>\text{T}$  substitution also decreased the transcriptional activation by PXR both in the presence and absence of rifampicin (Fig. 2F), but fold induction of MDR1 mRNA by rifampicin was not affected (Fig. 2E). These results suggest that DR4(II) does not contribute to rifampicin-activated MDR1 mRNA induction by PXR, which was consistent with previous observations [32]. PXR-dependent MDR1 gene transcription without rifampicin was observed not only in this study, but also in other reports [32,43,44], and PXR-dependent CYP3A4 gene transcription without rifampicin also has been reported [32,43,44]. Since the  $-7833\text{C}>\text{T}$  substitution decreased PXR-dependent basal activation without rifampicin (Fig. 2F), DR4(II), which contains  $-7833\text{C}>\text{T}$ , may be involved in PXR-dependent basal MDR1 expression, not ligand-dependent expression. Although  $-7833\text{C}>\text{T}$  affected nuclear receptor (VDR, TR $\beta$ , CAR, and PXR)-mediated MDR1 gene transactivation, it suppressed only about one third of total transactivity in each nuclear receptor (Fig. 2). We propose that this partial inhibition may be due to the multiple responsive elements that mediate MDR1 mRNA induction. Since xenobiotics and endogenous substrates-activated P-gp inductions play key roles in physiological functions [1,4,11,45], nuclear receptors might stimulate MDR1 transcription through the multiple responsive elements.

Since the T-to-C substitution at nucleotide  $-1211$  forms an XRE-like sequence (Fig. 3A) and AhR-induced MDR1 expression shows interindividual variation in human hepatocytes [37,46], we hypothesized that the  $-1211\text{T}>\text{C}$  substitution could affect the interindividual variation in MDR1 mRNA expression. However, AhR-dependent MDR1 activation was not observed with either  $-1211\text{T}$  or  $-1211\text{C}$  (Fig. 3B), indicating that the  $-1211\text{T}>\text{C}$  substitution does not form a functional XRE. The  $-1211\text{T}>\text{C}$  substitution also forms an HRE-like sequence (Fig. 3A), and the  $-129\text{T}>\text{C}$  substitution, which is completely linkage disequilibrium with  $-1211\text{T}>\text{C}$  in Japanese [27,31], reportedly affects the expression level of MDR1 mRNA in intestine and colon cancer cells. Since the concentrations of oxygen in the intestine and cancer cells are relatively low compared to other tissues [30], it is possible that the  $-1211\text{T}>\text{C}$  substitution affects the HIF-induced MDR1 expression in a synergistic or inhibitory manner. However, the  $-1211\text{T}>\text{C}$  substitution could not affect the MDR1 expression, which was induced by a chemically-induced hypoxic condition, indicating that the  $-1211\text{T}>\text{C}$  substitution could not form the functional HRE (Fig. 3C).

In summary, we have demonstrated that the  $-7833\text{C}>\text{T}$  substitution in the MDR1 gene decreases the binding affinities of nuclear

receptors, VDR/RXR $\alpha$ , TR $\beta$ /RXR $\alpha$ , CAR, CAR/RXR $\alpha$ , and PXR/RXR $\alpha$ , to their responsive elements located around  $-7833$ . We also showed that the C-to-T substitution at  $-7833$  reduces transcriptional activation of MDR1 by VDR, TR $\beta$ , CAR, and PXR. However, another SNP at  $-1211$  (T>C), which forms XRE-like and HRE-like sequences, failed to affect the AhR-dependent and hypoxia-induced MDR1 transcriptional activation. Although the frequency of  $-7833\text{C}>\text{T}$  substitution in the MDR1 gene is relatively low, knowledge of the  $-7833\text{C}>\text{T}$  substitution in MDR1 is crucial for subjects who hold the  $-7833\text{T}$  allele because the pharmacokinetics of P-gp substrates may differ from wild-type profile. Further study, especially clinical studies, is necessary to confirm the functional significance of the  $-7833\text{C}>\text{T}$  substitution in the interindividual differences in P-gp expression level.

## Acknowledgments

We thank Prof. Shuichi Koizumi (Yamanashi University, Japan) and Dr. Hideo Jinno (National Institute of Health Sciences, Japan) for providing the human RXR $\alpha$  cDNA and human CAR cDNA, respectively. This work was supported in part by grants from the Ministry of Health, Labor and Welfare of Japan (to M.T.) and the Japan Health Sciences Foundation (Research on Publicly Essential Drugs and Medical Devices) (to K.K.).

## References

- [1] V.J. Wachter, J.A. Silverman, Y. Zhang, L.Z. Benet, Role of P-glycoprotein and cytochrome P450 3A in limiting oral absorption of peptides and peptidomimetics, *J. Pharm. Sci.* 87 (1998) 1322–1330.
- [2] T. Sakaeda, T. Nakamura, K. Okumura, MDR1 genotype-related pharmacokinetics and pharmacodynamics, *Biol. Pharm. Bull.* 25 (2002) 1391–1400.
- [3] F. Thiebaut, T. Tsuruo, H. Hamada, M.M. Gottesman, I. Pastan, M.C. Willingham, Cellular localization of the multidrug-resistance gene product P-glycoprotein in normal human tissues, *Proc. Natl. Acad. Sci. USA* 84 (1987) 7735–7738.
- [4] M. Thorn, N. Finnstrom, S. Lundgren, A. Rane, L. Loof, Cytochromes P450 and MDR1 mRNA expression along the human gastrointestinal tract, *Br. J. Clin. Pharmacol.* 60 (2005) 54–60.
- [5] C. Cordon-Cardo, J.P. O'Brien, D. Casals, L. Rittman-Grauer, J.L. Biedler, M.R. Melamed, J.R. Bertino, Multidrug-resistance gene (P-glycoprotein) is expressed by endothelial cells at blood–brain barrier sites, *Proc. Natl. Acad. Sci. USA* 86 (1989) 695–698.
- [6] P.B. Watkins, The barrier function of CYP3A4 and P-glycoprotein in the small bowel, *Adv. Drug Deliv. Rev.* 27 (1997) 161–170.
- [7] J. Taipaleensuu, H. Tornblom, G. Lindberg, C. Einarsson, F. Sjoqvist, H. Melhus, P. Garberg, B. Sjöström, B. Lundgren, P. Artursson, Correlation of gene expression of ten drug efflux proteins of the ATP-binding cassette transporter family in normal human jejunum and in human intestinal epithelial Caco-2 cell monolayers, *J. Pharmacol. Exp. Ther.* 299 (2001) 164–170.
- [8] C. Zimmermann, H. Gutmann, P. Hruz, J.P. Gutzwiller, C. Beglinger, J. Drewe, Mapping of multidrug resistance gene 1 and multidrug resistance-associated protein isoform 1 to 5 mRNA expression along the human intestinal tract, *Drug Metab. Dispos.* 33 (2005) 219–224.
- [9] K. Sai, N. Kaniwa, M. Itoda, Y. Saito, R. Hasegawa, K. Komamura, K. Ueno, S. Kamakura, M. Kitakaze, K. Shirao, H. Minami, A. Ohtsu, T. Yoshida, N. Saijo, Y. Kitamura, N. Kamatani, S. Ozawa, J. Sawada, Haplotype analysis of ABCB1/MDR1 blocks in a Japanese population reveals genotype-dependent renal clearance of irinotecan, *Pharmacogenetics* 13 (2003) 741–757.
- [10] K.N. Faber, M. Muller, P.L. Jansen, Drug transport proteins in the liver, *Adv. Drug Deliv. Rev.* 55 (2003) 107–124.
- [11] A. Doran, R.S. Obach, B.J. Smith, N.A. Hosea, S. Becker, E. Callegari, C. Chen, X. Chen, E. Choo, J. Cianfroga, L.M. Cox, J.P. Gibbs, M.A. Gibbs, H. Hatch, C.E. Hop, I.N. Kasman, J. Laperle, J. Liu, X. Liu, M. Logman, D. MacLain, F.M. Nedza, F. Nelson, E. Olson, S. Rahematpura, D. Raunig, S. Rogers, K. Schmidt, D.K. Spracklin, M. Szeew, M. Troutman, E. Tseng, M. Tu, J.W. Van Deusen, K. Venkatakrishnan, G. Walens, E.Q. Wang, D. Wong, A.S. Yaggar, C. Zhang, The impact of P-glycoprotein on the disposition of drugs targeted for indications of the central nervous system: evaluation using the MDR1A/1B knockout mouse model, *Drug Metab. Dispos.* 33 (2005) 165–174.
- [12] N.H. Hendrikse, A.H. Schinkel, E.G. de Vries, E. Fluks, W.T. Van der Graaf, A.T. Willemsen, W. Vaalburg, E.J. Franssen, Complete in vivo reversal of P-glycoprotein pump function in the blood–brain barrier visualized with positron emission tomography, *Br. J. Pharmacol.* 124 (1998) 1413–1418.
- [13] M.B. Muller, M.E. Keck, E.B. Binder, A.E. Kresse, T.P. Hagemeyer, R. Landgraf, F. Holsboer, M. Uhr, ABCB1 (MDR1)-type P-glycoproteins at the blood–brain barrier modulate the activity of the hypothalamic–pituitary–adrenocortical system: implications for affective disorder, *Neuropsychopharmacology* 28 (2003) 1991–1999.

- [14] E.G. Schuetz, K.N. Furuya, J.D. Schuetz, Interindividual variation in expression of P-glycoprotein in normal human liver and secondary hepatic neoplasms, *J. Pharmacol. Exp. Ther.* 275 (1995) 1011–1018.
- [15] K.S. Lown, R.R. Mayo, A.B. Leichtman, H.L. Hsiao, D.K. Turgeon, P. Schmiedlin-Ren, M.B. Brown, W. Guo, S.J. Rossi, L.Z. Benet, P.B. Watkins, Role of intestinal P-glycoprotein (mdr1) in interpatient variation in the oral bioavailability of cyclosporine, *Clin. Pharmacol. Ther.* 62 (1997) 248–260.
- [16] A. Asghar, J.C. Gorski, B. Haehner-Daniels, S.D. Hall, Induction of multidrug resistance-1 and cytochrome P450 mRNAs in human mononuclear cells by rifampin, *Drug Metab. Dispos.* 30 (2002) 20–26.
- [17] S. Taniguchi, Y. Mochida, T. Uchiyama, T. Tahira, K. Hayashi, K. Takagi, M. Shimada, Y. Maehara, H. Kuwano, S. Kono, H. Nakano, M. Kuwano, M. Wada, Genetic polymorphism at the 5' regulatory region of multidrug resistance 1 (MDR1) and its association with interindividual variation of expression level in the colon, *Mol. Cancer Ther.* 2 (2003) 1351–1359.
- [18] R. Kerb, A.S. Aynacioglu, J. Brockmoller, R. Schlagenhauser, S. Bauer, T. Szekeres, A. Hamwi, M. Fritzer-Szekeres, C. Baumgartner, H.Z. Ongen, P. Guzelbey, I. Roots, U. Brinkmann, The predictive value of MDR1, CYP2C9, and CYP2C19 polymorphisms for phenytoin plasma levels, *Pharmacogenomics J.* 1 (2001) 204–210.
- [19] U. Brinkmann, M. Eichelbaum, Polymorphisms in the ABC drug transporter gene MDR1, *Pharmacogenomics J.* 1 (2001) 59–64.
- [20] S. Saito, A. Iida, A. Sekine, Y. Miura, C. Ogawa, S. Kawauchi, S. Higuchi, Y. Nakamura, Three hundred twenty-six genetic variations in genes encoding nine members of ATP-binding cassette, subfamily B (ABCB/MDR/TAP), in the Japanese population, *J. Hum. Genet.* 47 (2002) 38–50.
- [21] R.B. Kim, B.F. Leake, E.F. Choo, G.K. Dresser, S.V. Kubba, U.I. Schwarz, A. Taylor, H.G. Xie, J. McKinsey, S. Zhou, L.B. Lan, J.D. Schuetz, E.G. Schuetz, G.R. Wilkinson, Identification of functionally variant MDR1 alleles among European Americans and African Americans, *Clin. Pharmacol. Ther.* 70 (2001) 189–199.
- [22] C. Kimchi-Sarfaty, J.M. Oh, I.W. Kim, Z.E. Sauna, A.M. Calcagno, S.V. Ambudkar, M.M. Gottesman, A "silent" polymorphism in the MDR1 gene changes substrate specificity, *Science* 315 (2007) 525–528.
- [23] S. Hoffmeyer, O. Burk, O. von Richter, H.P. Arnold, J. Brockmoller, A. John, I. Cascorbi, T. Gerloff, I. Roots, M. Eichelbaum, U. Brinkmann, Functional polymorphisms of the human multidrug-resistance gene: multiple sequence variations and correlation of one allele with P-glycoprotein expression and activity in vivo, *Proc. Natl Acad. Sci. USA* 97 (2000) 3473–3478.
- [24] D. Wang, A.D. Johnson, A.C. Papp, D.L. Kroetz, W. Saade, Multidrug resistance polypeptide 1 (MDR1, ABCB1) variant 3435C>T affects mRNA stability, *Pharmacogenet. Genomics* 15 (2005) 693–704.
- [25] T. Sakaeda, T. Nakamura, M. Horinouchi, M. Kakimoto, N. Ohmoto, T. Sakai, Y. Morita, T. Tamura, N. Aoyama, M. Hirai, M. Kasuga, K. Okumura, MDR1 genotype-related pharmacokinetics of digoxin after single oral administration in healthy Japanese subjects, *Pharm. Res.* 18 (2001) 1400–1404.
- [26] N. Morita, T. Yasumori, K. Nakayama, Human MDR1 polymorphism: G2677T/A and C3435T have no effect on MDR1 transport activities, *Biochem. Pharmacol.* 65 (2003) 1843–1852.
- [27] Y. Moriya, T. Nakamura, M. Horinouchi, T. Sakaeda, T. Tamura, N. Aoyama, T. Shirakawa, A. Gotoh, S. Fujimoto, M. Matsuo, M. Kasuga, K. Okumura, Effects of polymorphisms of MDR1, MRP1, and MRP2 genes on their mRNA expression levels in duodenal enterocytes of healthy Japanese subjects, *Biol. Pharm. Bull.* 25 (2002) 1356–1359.
- [28] B. Wang, S. Ngoi, J. Wang, S.S. Chong, C.G. Lee, The promoter region of the MDR1 gene is largely invariant, but different single nucleotide polymorphism haplotypes affect MDR1 promoter activity differently in different cell lines, *Mol. Pharmacol.* 70 (2006) 267–276.
- [29] H. Takane, D. Kobayashi, T. Hirota, J. Kigawa, N. Terakawa, K. Otsubo, I. Ieiri, Haplotype-oriented genetic analysis and functional assessment of promoter variants in the MDR1 (ABCB1) gene, *J. Pharmacol. Exp. Ther.* 311 (2004) 1179–1187.
- [30] T. Koyama, T. Nakamura, C. Komoto, T. Sakaeda, M. Taniguchi, N. Okamura, T. Tamura, N. Aoyama, T. Kamigaki, Y. Kuroda, M. Kasuga, K. Kadoyama, K. Okumura, MDR1 T-129C polymorphism can be predictive of differentiation, and thereby prognosis of colorectal adenocarcinomas in Japanese, *Biol. Pharm. Bull.* 29 (2006) 1449–1453.
- [31] W. Qian, M. Homma, F. Itagaki, H. Tachikawa, Y. Kawanishi, K. Mizukami, T. Asada, S. Inomata, K. Honda, N. Ohkohchi, Y. Kohda, MDR1 gene polymorphism in Japanese patients with schizophrenia and mood disorders including depression, *Biol. Pharm. Bull.* 29 (2006) 2446–2450.
- [32] A. Geick, M. Eichelbaum, O. Burk, Nuclear receptor response elements mediate induction of intestinal MDR1 by rifampin, *J. Biol. Chem.* 276 (2001) 14581–14587.
- [33] O. Burk, K.A. Arnold, A. Geick, H. Tegude, M. Eichelbaum, A role for constitutive androstane receptor in the regulation of human intestinal MDR1 expression, *Biol. Chem.* 386 (2005) 503–513.
- [34] K. Kurose, M. Saeki, M. Tohkin, R. Hasegawa, Thyroid hormone receptor mediates human MDR1 gene expression—identification of the response region essential for gene expression, *Arch. Biochem. Biophys.* 474 (2008) 82–90.
- [35] M. Saeki, K. Kurose, M. Tohkin, R. Hasegawa, Identification of the functional vitamin D response elements in the human MDR1 gene, *Biochem. Pharmacol.* 76 (2008) 531–542.
- [36] K. Sai, M. Itoda, Y. Saito, K. Kurose, N. Katori, N. Kaniwa, K. Komamura, T. Kotake, H. Morishita, H. Tomoike, S. Kamakura, M. Kitakaze, T. Tamura, N. Yamamoto, H. Kunitoh, Y. Yamada, Y. Ohe, Y. Shimada, K. Shirao, H. Minami, A. Ohtsu, T. Yoshida, N. Saijo, N. Kamatani, S. Ozawa, J. Sawada, Genetic variations and haplotype structures of the ABCB1 gene in a Japanese population: an expanded haplotype block covering the distal promoter region, and associated ethnic differences, *Ann. Hum. Genet.* 70 (2006) 605–622.
- [37] E. Jigorel, M. Le Vee, C. Boursier-Neyret, Y. Parmentier, O. Fardel, Differential regulation of sinusoidal and canalicular hepatic drug transporter expression by xenobiotics activating drug-sensing receptors in primary human hepatocytes, *Drug Metab. Dispos.* 34 (2006) 1756–1763.
- [38] K.M. Comerford, T.J. Wallace, J. Karhausen, N.A. Louis, M.C. Montalto, S.P. Colgan, Hypoxia-inducible factor-1-dependent regulation of the multidrug resistance (MDR1) gene, *Cancer Res.* 62 (2002) 3387–3394.
- [39] S. Koyano, Y. Saito, H. Fukushima-Uesaka, S. Ishida, S. Ozawa, N. Kamatani, H. Minami, A. Ohtsu, T. Hamaguchi, K. Shirao, T. Yoshida, N. Saijo, H. Jinno, J. Sawada, Functional analysis of six human aryl hydrocarbon receptor variants in a Japanese population, *Drug Metab. Dispos.* 33 (2005) 1254–1260.
- [40] H. Jinno, T. Tanaka-Kagawa, N. Hanioka, S. Ishida, M. Saeki, A. Soyama, M. Itoda, T. Nishimura, Y. Saito, S. Ozawa, M. Ando, J. Sawada, Identification of novel alternative splice variants of human constitutive androstane receptor and characterization of their expression in the liver, *Mol. Pharmacol.* 65 (2004) 496–502.
- [41] K. Kurose, M. Tohkin, M. Fukuhara, A novel positive regulatory element that enhances hamster CYP2A8 gene expression mediated by xenobiotic responsive element, *Mol. Pharmacol.* 55 (1999) 279–287.
- [42] J.M. Gow, L.W. Chinn, D.L. Kroetz, The effects of ABCB1 3'-untranslated region variants on mRNA stability, *Drug Metab. Dispos.* 36 (2008) 10–15.
- [43] A. Pfunder, H. Gutmann, C. Beglinger, J. Drewe, Gene expression of CYP3A4, ABC-transporters (MDR1 and MRP1-MRP5) and hPXR in three different human colon carcinoma cell lines, *J. Pharm. Pharmacol.* 55 (2003) 59–66.
- [44] L. Cerveny, L. Svecova, E. Anzenbacherova, R. Vrzal, F. Staud, Z. Dvorak, J. Ulrichova, P. Anzenbacher, P. Pavek, Valproic acid induces CYP3A4 and MDR1 gene expression by activation of constitutive androstane receptor and pregnane X receptor pathways, *Drug Metab. Dispos.* 35 (2007) 1032–1041.
- [45] S. Marchetti, R. Mazzanti, J.H. Beijnen, J.H. Schellens, Concise review: clinical relevance of drug–drug and herb–drug interactions mediated by the ABC transporter ABCB1 (MDR1, P-glycoprotein), *Oncologist* 12 (2007) 927–941.
- [46] P. Olinga, M.G. Elferink, A.L. Draaisma, M.T. Merema, J.V. Castell, G. Perez, G.M. Groothuis, Coordinated induction of drug transporters and phase I and II metabolism in human liver slices, *Eur. J. Pharm. Sci.* 33 (2008) 380–389.



## Review

### Prediction of Severe Adverse Drug Reactions Using Pharmacogenetic Biomarkers

Masahiro TOHKIN<sup>1,\*</sup>, Akihiro ISHIGURO<sup>1,2</sup>, Nahoko KANIWA<sup>1</sup>,  
Yoshiro SAITO<sup>1</sup>, Kouichi KUROSE<sup>1</sup> and Ryuichi HASEGAWA<sup>1</sup>

<sup>1</sup>Division of Medicinal Safety Science, National Institute of Health Sciences, Tokyo, Japan

<sup>2</sup>Office of Safety II, Pharmaceuticals and Medical Devices Agency, Tokyo, Japan

Full text of this paper is available at <http://www.jstage.jst.go.jp/browse/dmpk>

**Summary:** Severe adverse drug reactions (ADRs) are a major issue for drug therapy because they can cause serious disorders and be life-threatening. Many severe ADRs appear to be idiosyncratic and unpredictable. Genetic factors may underlie susceptibility to severe ADRs, and identification of predisposing genotypes may improve drug therapy by facilitating prescreening of carriers for specific genetic biomarkers. In this review, we clarify the current status of ADRs in Japan from open ADR data sources. Then, we introduce recent progress in the field of pharmacogenetic biomarkers for severe cutaneous ADRs, liver injury, and statin-induced myopathy. Key challenges for discovery of predictable risk alleles for these severe ADRs are also discussed.

**Keywords:** drug-induced liver injury; drug-induced myopathy; human lymphocyte antigen; Stevens-Johnson syndrome; toxic epidermal necrolysis

#### Introduction

Severe adverse drug reaction (ADR) is a major reason for failure of new drug development and withdrawal of approved drugs from the market. The classical pharmacological classification of ADRs by Rawlins and Thompson distinguished two types of severe ADRs.<sup>1)</sup> Type A reactions are dose-dependent and predictable on the basis of the drug's known pharmacological actions. Type A reactions are relatively common and include hypoglycemia induced by diabetic drugs and bleeding induced by warfarin, an oral anti-coagulant. By contrast, type B reactions are idiosyncratic, unpredictable from the pharmacological action of the drug, and are not necessarily dose-dependent. These type B reactions make up approximately 10–15% of all ADRs and include severe cutaneous disorders, such as Stevens-Johnson syndrome (SJS) and toxic epidermal necrolysis (TEN), and drug-induced liver injury (DILI) caused by various drugs.

The aim of this review is to provide an update on the current understanding of pharmacogenetic analysis related to severe ADRs, especially severe cutaneous ADRs,

DILI, and statin-induced myopathy; great progress has been recently observed in pharmacogenetic biomarkers of these ADRs, and this should facilitate early-stage detection of severe ADRs. Therefore, pharmacogenetic biomarkers of ADRs hold promise for reducing severe ADRs and pave the way for creating more affordable pharmaceuticals.

#### Domestic case reports for severe adverse drug reactions in Japan

Domestic cases of severe ADRs are reported to the Pharmaceuticals and Medical Devices Agency (PMDA) by pharmaceutical companies based on the Pharmaceutical Affairs Law in Japan. The ADR report in Japan includes information on suspicious drugs; ADR diagnoses, which are expressed using the Medical Dictionary for Regulatory Activities preferred terms (MedDRA-PT); and patient background such as gender, age, and concomitant use of other drugs. Because the quotation frequency of MedDRA-PTs in ADR reports reflects the number of ADR events, the quotation frequency provides basic information for estimating the event frequency of each ADR. Ac-

Received; November 9, 2009, Accepted; December 31, 2009

\*To whom correspondence should be addressed: Masahiro TOHKIN, Ph.D., Division of Medicinal Safety Science, National Institute of Health Sciences, 1-18-1 Kamiyoga, Setagaya-ku, Tokyo 158-8501, Japan. Tel. +81-3-3700-9789, Fax. +81-3-3700-9788, E-mail: [tohkin@nihs.go.jp](mailto:tohkin@nihs.go.jp)

The URL of the Drug Safety Information website of the Pharmaceuticals and Medical Devices Agency is "[http://www.info.pmda.go.jp/fukusayou/menu\\_fukusayou\\_attention.html](http://www.info.pmda.go.jp/fukusayou/menu_fukusayou_attention.html)" and the URL of material from the Committee on Safety of Drugs of the Ministry of Health, Labour and Welfare is "<http://www.mhlw.go.jp/shingi/yakuji.html#anzen>".

**Table 1.** Accumulated number of each adverse drug reaction term in cases reported from April 2004 to February 2009

Classification of accumulated frequency of each ADR term	Number of ADR terms in each classification	Representative ADR terms concerning SJS/TEN, hepatotoxicity, and rhabdomyolysis*
> 1,000	30	Hepatic function abnormal (4,866), Liver disorder (3,936), Rhabdomyolysis (1,648), Stevens-Johnson syndrome (1,202)
500–1,000	26	Toxic epidermal necrolysis (751), Jaundice (715)
100–499	221	Hepatitis fulminant (406), Hepatitis acute (388), Oculomucocutaneous syndrome (324), Hepatitis (310), Cholestasis (184)
< 100	4,491	
Total	4,768	

Based on the open data source MHLW website: <http://www.mhlw.go.jp/shingi/yakuji.html#anzen>

\*Accumulated frequency is listed in parentheses.

**Table 2.** Number of case reports concerning SJS/TEN for fiscal years 2005 to 2008

Year (fiscal) Molecular entities	2005			2006			2007			2008		
	SJS	TEN	Sum	SJS	TEN	Sum	SJS	TEN	Sum	SJS	TEN	Sum
Allopurinol	14	12	26	22	8	30	15	11	27	33	18	51
Carbamazepine	17	5	22	17	4	21	24	6	30	26	10	36
Diclofenac	9	5	14	8	2	10	6	1	7	7	9	16
Loxoprofen	9	2	11	11	5	16	7	7	14	12	9	21
Phenobarbital	5	5	10	2	4	6	6	0	6	8	4	12
Non-pyridines	6	4	10	2	4	6	1	9	10	2	4	6
Zonisamide	9	1	10	3	2	5	1	4	5	9	6	15
Acetaminophen	1	4	5	1	5	6	1	12	13	3	10	13
Mortality (rate, %)	13 (6)	41 (27)		14 (5)	41 (30)		17 (7)	36 (23)		12 (4)	50 (26)	
Total	223	151	374	271	136	407	260	156	416	289	189	478

Based on the open data source PMDA drug safety information website: [http://www.info.pmda.go.jp/fukusayou/menu\\_fukusayou\\_attention.html](http://www.info.pmda.go.jp/fukusayou/menu_fukusayou_attention.html)

cording to material from the Committee on Safety of Drugs of the Ministry of Health, Labour and Welfare (MHLW, see footnote for URL address), the total number of MedDRA-PTs was 4,768 terms cited among 168,045 ADR events from domestic cases reported to PMDA from April 2004 to February 2009. As shown in **Table 1**, the number of MedDRA-PTs cited more than 100 times among 168,045 ADR events was 277. SJS/TEN, hepatotoxicity, and rhabdomyolysis are considered to be major ADRs in Japan because the frequently cited top 277 MedDRA-PTs included most ADR terms concerning SJS/TEN, hepatotoxicity, and rhabdomyolysis. In particular, accumulated quotation frequency of abnormal hepatic function (4,866) and liver disorder (3,936) ranked second and third, followed by interstitial pneumonia (5,190).

Major severe cutaneous ADRs, SJS and TEN, are life-threatening skin disorders which are often accompanied by high fever and systemic complications.<sup>2,3)</sup> SJS/TEN incidence is generally very low and more than 100 different causative drugs have been reported.<sup>4–6)</sup> We counted the event number for major suspected drugs in domestic

cases concerning SJS and TEN from April 2005 to March 2009 based on the open data source of PMDA, and the results are shown in **Table 2**. Allopurinol, an anti-hyperuricemia drug, is the most frequently reported drug for SJS and TEN (**Table 2**). Many cases were reported with anticonvulsant drugs including carbamazepine (CBZ), phenobarbital, and zonisamide and non-steroidal anti-inflammation drugs (NSAID), which include diclofenac, loxoprofen, non-pyridines, and acetaminophen (**Table 2**). Although it is difficult to calculate the exact incidence of SJS and TEN in Japan, the Japanese mortality rates of SJS and TEN based on domestic cases were about 4–7% and 23–30%, respectively, which were in accordance with rates reported in other populations<sup>7,8)</sup> (**Table 2**). Therefore, the mortality rate calculated by ADR reports can be a useful reference to estimate the mortality rate of severe ADRs such as SJS and TEN.

The liver is a common target for drug toxicity due to its pivotal role in drug metabolism. Moreover, any drug has the potential to cause liver injury.<sup>9)</sup> Therefore, hepatotoxicity is the most common ADR, causing drug withdrawals and post-marketing regulatory decisions and

Table 3. Case reports concerning hepatotoxicity in fiscal year 2008

Molecular entities	Hepatic function abnormal	Liver disorder	Jaundice	Hepatitis fulminant	Hepatitis acute	Hepatitis	Cholestasis
Terbinafine HCl	52	47	3	0	5	11	2
Fluvastatin Na	33	21	0	0	0	1	0
Itraconazole	17	10	0	0	0	0	0
Loxoprofen Na	12	15	0	0	0	2	0
Carbamazepine	11	13	0	1	1	2	0
Rosuvastatin Ca	15	7	0	1	0	3	0
Tegafur/Uracil	12	10	1	3	1	0	0
Ticlopidine HCl	9	12	4	0	0	1	15
Gefitinib	16	2	0	1	0	0	0
Cyclosporine	11	3	3	0	0	0	1
Atorvastatin Ca	7	8	0	0	0	0	1
Fenofibrate	7	5	0	0	0	4	0
Acarbose	2	8	0	0	0	0	0
Tranilast	6	4	1	0	0	1	1
Aspirin	1	8	1	0	0	1	0
Non-pyrines (4)	3	5	0	0	0	0	0
Voglibose	3	4	1	0	1	3	0
Temozolomide	6	1	0	0	0	0	0
Cefcapene pivoxil HCl	3	3	2	2	0	0	0
Total	1,063	771	90	82	68	58	55

Based on the open data source PMDA drug safety information website: [http://www.info.pmda.go.jp/fukusayou/menu\\_fukusayou\\_attention.html](http://www.info.pmda.go.jp/fukusayou/menu_fukusayou_attention.html)

is associated with significant mortality.<sup>10)</sup> We counted the event number of major suspected drugs in domestic cases concerning various hepatotoxicities based on the open data source from PMDA from April 2008 to March 2009, and the results are shown in **Table 3**. Highly reported hepatotoxicities were abnormal hepatic function, liver disorder, jaundice, fulminant hepatitis, acute hepatitis, hepatitis, and cholestasis (**Table 3**). Although the difference between abnormal hepatic function and hepatic disorder is unclear, these MedDRA-PTs are exclusively used. The total number of case reports for hepatotoxicity was 2,509. Other hepatotoxicity related reports (more than 10 cases) not included in **Table 3** were hepatic failure (44), hepatocellular injury (41), cholelithiasis (21), acute hepatic failure (19), autoimmune hepatitis (19), hyperbilirubinemia (17), veno-occlusive liver disease (15), cholecystitis (14), mixed liver injury (12), cholecystitis acute (11), jaundice cholestatic (10), and cholangitis (10). There were 89 other hepatotoxicity-related reports (less than 9 cases) not included in **Table 3**. The estimated frequency of reported hepatotoxicity ranges from 1 per 10,000 to 1 per 10 million patient-years of exposure<sup>11)</sup> (**Table 3**).

Rhabdomyolysis, one of the most serious myopathies, is characterized by the leakage of muscle cell content, including electrolytes, myoglobin, and other sarcoplasmic proteins [e.g., creatine kinase, aldolase, lactate dehydrogenase, alanine aminotransferase (AST), and aspar-

Table 4. Case reports concerning rhabdomyolysis for fiscal years 2004 to 2008

Molecular entities	Fiscal year				
	2004	2005	2006	2007	2008
Atorvastatin	51	41	48	31	28
Bezafibrate	16	22	17	16	11
Pravastatin	21	24	19	9	11
Simvastatin	21	15	5	8	2
Levofloxacin	9	10	10	9	4
Fluvastatin	13	8	7	4	6
Omeprazole	10	8	10	3	4
Propofol	8	14	2	5	8
Rosuvastatin	—	0	4	15	14
Risperidone	7	5	5	5	5
Fenofibrate	10	9	3	1	5
Pitavastatin	4	3	10	3	9
Total	389	351	359	291	332

Based on the open data source PMDA drug safety information website: [http://www.info.pmda.go.jp/fukusayou/menu\\_fukusayou\\_attention.html](http://www.info.pmda.go.jp/fukusayou/menu_fukusayou_attention.html)

tate aminotransferase (AST)] into the circulation.<sup>12)</sup> Lipid-lowering drugs (e.g., statins and fibrates) are well known causes of rhabdomyolysis, and reports produced by the US Food and Drug Administration (FDA) showed that the rate of fatal rhabdomyolysis was 0.15 per 1 million statin

prescriptions dispensed.<sup>13,14</sup> **Table 4** shows suspicious drugs in Japan, which are similar to US data, and the reported range of case numbers is 300 to 400 per year for the last 5 years (April 2004–March 2009).

### Genomic analysis of severe adverse drug reactions

Severe ADRs affect only a minority of patients taking drugs. However, hereditary forms of severe ADRs and cases occurring in identical twins have been reported, implying involvement of certain genetic factors in predisposing individuals to such severe ADRs.<sup>15,16</sup> The genetic basis of ADRs can be categorized into two broad groups. The first group involves genes that drive pharmacological mechanisms (drug targets, drug metabolizing enzymes, and drug transporters).<sup>17</sup> Common mechanisms underlying these severe ADRs are unusual drug accumulation in the target organ due to polymorphisms in drug metabolizing enzyme and drug transporter genes, and unusual sensitivity in the target organ due to changes in drug target genes.<sup>18</sup> The second category involves the immune system in a drug-induced allergic reaction. One important molecule for ADRs associated with immune reactions is the human lymphocyte antigen (HLA), which plays a key role in initiation of immune responses and killing target cells by presenting antigens to the T-cell receptor.<sup>19</sup> The *HLA* gene region codes for three classical class I (HLA-A, HLA-B, and HLA-C) and three class II (HLA-DR, HLA-DP, and HLA-DQ) antigens. Class I antigens are recognized by cytotoxic CD8<sup>+</sup> T cells and class IIs by CD4<sup>+</sup> T cells. Of the highly polymorphic *HLA* genes, *HLA-B* is the most polymorphic with over 800 variants reported in the human genome.<sup>20</sup> *HLA* genes within each class encode structurally similar but distinct *HLA* proteins that bind and present *HLA*-type-specific antigenic peptides to T-cell receptors.<sup>19,20</sup> *HLA* disease associations that are related to genes with immunological and inflammatory functions have been identified in many autoimmune and inflammatory conditions.

In addition to these susceptible genes, recent advances in molecular biology have led to analysis of the association of whole genome polymorphisms with ADRs.<sup>21,22</sup> For example, recent high-density DNA microarrays can analyze more than one million genomic biomarkers at the same time. Therefore, association analysis has been conducted by both candidate gene and genome-wide analysis.

### Severe cutaneous adverse drug reactions

Carbamazepine (CBZ) is one of the most widely used aromatic anticonvulsants and is often used as a pain-relief drug for prosopalgia. CBZ is metabolized by mainly hepatic CYP3A4, CYP2B6, and CYP2C8, which generate various potentially reactive metabolites, such as CBZ-10,11-epoxide; 3-hydroxy-CBZ; 2-hydroxy-CBZ; and CBZ-2,3-epoxide.<sup>23,24</sup> CBZ is generally well tolerated but

also is associated with idiosyncratic adverse reactions such as SJS/TEN. A high frequency of CBZ-induced SJS/TEN was reported in Han Chinese (0.25% in new exposures to CBZ) compared to Caucasians (0.014% in new exposures to CBZ).<sup>25–29</sup> Furthermore, CBZ-induced SJS/TEN has been reported in identical twins.<sup>15</sup> These studies suggest that susceptibility to such reactions may be genetically determined. Since most reactive CBZ metabolites are detoxified to non-toxic dihydrodiols by liver microsomal epoxide hydrolase 1 (EPHX1) or to glutathione conjugates by glutathione S-transferase  $\mu$  1 (GSTM1),<sup>30,31</sup> some researchers have attempted to find the defective alleles of *EPHX* and *GST* genes in patients with SJS/TEN; however, these attempts have failed to find associations, indicating that reactive metabolite generation from CBZ is not sufficient to cause SJS/TEN.<sup>32,33</sup> Recently, Chung *et al.* reported a tight association between CBZ-induced SJS/TEN and *HLA-B\*1502* allele in Han Chinese<sup>25</sup> (**Table 5**). They showed that all 44 Han Chinese patients with CBZ-induced SJS/TEN carried the *HLA-B\*1502* allele and its odds ratio was 2,505 (95% confidence interval, 195 to 27,483,  $P_c = 2.02 \times 10^{-32}$ ). The finding was further confirmed by the same group in

**Table 5.** Association between severe cutaneous adverse drug reaction and *HLA* type

Drug	HLA	Population	OR or (N) <sup>a</sup>	P <sup>b</sup>	Ref
Carbamazepine	B*1502	Han Chinese	2505	$2.0 \times 10^{-32}$	25)
Carbamazepine	B*1502	Asian ancestry	(4/4)	—	37)
Carbamazepine	B*1502	European	(0/8)	—	37)
Carbamazepine	B*1502	Thai	25.5	0.0005	106)
Carbamazepine	B*1502	Japanese	(0/7)	—	36)
Carbamazepine	B*1502	Han Chinese	1357	$1.6 \times 10^{-41}$	34)
Phenytoin	B*1502	Thai	18.5	0.005	106)
Antiepileptic	B*1502	Han Chinese	17.6	0.001	35)
Allopurinol	B*5801	Han Chinese	580	$4.7 \times 10^{-24}$	44)
Allopurinol	B*5801	European	80	$< 10^{-6}$	5)
Allopurinol	B*5801	Japanese	40.8	$< 10^{-4}$	36)
Abacavir	B*5701	Australian	117	$< 10^{-4}$	48)
Abacavir	B*5701	British	24	$< 10^{-4}$	47)
Abacavir	B*5701	Australian	960	$< 10^{-4}$	50)
Abacavir	B*5701	White	(36/65)	—	49)
		Black	(0/9)	—	
		Other	(1/10)	—	
Abacavir	B*5701	Japanese	(0/7)	—	40)
Nevirapine	DRB1*0101 and high CD4	Caucasian Australian	18	0.0006	59)
Nevirapine	Cw*0802-B*1402	Sardinian	15	0.05	60)
Nevirapine	B*3505	Thai	18.96	$4.6 \times 10^{-6}$	62)
Nevirapine	Cw8	Japanese	6.2	0.03	61)

<sup>a</sup> OR is odds ratio and (N) is sensitivity (carrier cases/all cases).

<sup>b</sup> P<sub>c</sub> indicates corrected P value.



an additional study that included patients who were Han Chinese or Chinese descendants from Taiwan, Hong Kong, China, and the USA<sup>34)</sup> (Table 5). The involvement of *HLA-B\*1502* was also detected in SJS/TEN caused by other aromatic anti-epileptic drugs, such as phenytoin in Han Chinese<sup>35)</sup> (Table 5). However, such strong association between *HLA-B\*1502* and CBZ-induced SJS/TEN has not been detected in Caucasian and Japanese SJS/TEN patients<sup>36,37)</sup> (Table 5). *HLA-B\*1502* is present at a higher allele frequency in South-east Asian populations than in Caucasian and Japanese populations.<sup>38)</sup> *HLA-B\*1502* was not detected in 486 healthy Japanese subjects<sup>39)</sup> and in 935 USA Caucasians (<http://www.allelefrequencies.net/>), while the allele frequency in Han Chinese is 8.6%.<sup>25)</sup> The low frequency of *HLA-B\*1502* in Caucasian and Japanese populations may account for the fact that no association between *HLA-B\*1502* and CBZ-induced SJS/TEN was observed in Caucasians and Japanese.<sup>36,40)</sup> Alternatively, these results suggest that *HLA-B\*1502* is involved in the mechanism, but is not sufficient for CBZ-induced SJS/TEN. There could be other co-factors, such as virus infection or other variants of genes, for example, *CYP3A4*, *CYP2B6*, *CYP2C8*, *EPHX1*, and *GSTM1*; T cell receptors; genes related to apoptosis; or genes for co-stimulatory molecules involved in the interaction between antigen-presenting cells and T cells.

Allopurinol is a xanthine oxidase inhibitor that prevents the production of uric acid and is commonly used for hyperuricemia and gout.<sup>41)</sup> Allopurinol is metabolized by xanthine oxidase to oxipurinol, which forms ribonucleotide adduct and ribonucleoside adduct.<sup>42)</sup> Allopurinol has been reported to be a causative drug of a variety of delayed cutaneous adverse reactions, such as SJS/TEN.<sup>43)</sup> Recently, a strong association of *HLA-B\*5801* with allopurinol-induced severe cutaneous adverse reactions (drug-induced hypersensitivity syndrome and SJS/TEN) was found in Han Chinese in Taiwan<sup>44)</sup> (Table 5). They showed that the *HLA-B\*5801* allele was present in all patients (51/51) with allopurinol-induced severe cutaneous adverse reactions, but only in 15% of tolerant patients (20/135). The odds ratio was 580 (95% confidence interval, 34 to 9781,  $P_c = 4.7 \times 10^{-24}$ ). Although the association was confirmed in Caucasians<sup>5)</sup> and Japanese,<sup>36)</sup> the odds ratio in Han Chinese (580) was much higher than that in Caucasians (80) and Japanese (40) (Table 5). Approximately 9 to 11% of Han Chinese are carriers of the allele, and its prevalence is generally lower in Caucasian (1 to 6%), Japanese (0.68%), and African (2 to 4%) populations.<sup>43,44)</sup> These reports suggest that *HLA-B\*5801* might be a genetic biomarker for allopurinol-induced severe cutaneous adverse reactions; however, the extent of the association showed ethnic differences.

Abacavir is a potent nucleoside analog reverse transcriptase inhibitor that is used in combination with other

drugs to treat human immunodeficiency virus infection. The most serious adverse reaction of abacavir that limits its use in therapy is a hypersensitivity reaction which includes the combination of fever, skin rash, constitutional symptoms, gastrointestinal tract symptoms, and respiratory symptoms.<sup>45)</sup> Hypersensitivity to abacavir occurs in approximately 5 to 8% of patients treated with abacavir, typically within 1 to 6 weeks of the initial dose.<sup>45)</sup> Abacavir is metabolized by type 1 alcohol dehydrogenase to an aldehyde-reactive metabolite.<sup>46)</sup> The initial association between *HLA-B\*5701* and abacavir-induced hypersensitivity reaction was elucidated by two independent research groups in 2002<sup>47,48)</sup> (Table 5). The association was reported only in Caucasians and not in Africans or Japanese<sup>40,49)</sup> (Table 5) because the allelic frequency of *HLA-B\*5701* in Caucasians is approximately 8%, but low in individuals of African or Asian descent.<sup>47,48)</sup> Fine recombinant genetic mapping has identified a significant linkage disequilibrium of the haplotypic M493T polymorphism of heat shock protein-Hom (Hsp70-Hom M493T) and *HLA-B\*5701* in abacavir-induced hypersensitivity reaction cases in Western Australians, which enhanced the discrimination of hypersensitive subjects from the tolerant control (odds ratio, 3,893;  $P_c < 0.00001$ ) when compared to *HLA-B\*5701* only (odds ratio, 960;  $P_c < 0.00001$ ).<sup>50)</sup> The Hsp70-Hom M493T polymorphism may facilitate the loading of abacavir- or its metabolite-hapten endogenous peptide onto *HLA-B\*5701*.<sup>51)</sup> A large randomized, controlled clinical trial assessing the clinical effectiveness of *HLA-B\*5701* screening in Caucasians (PREDICT-1 study),<sup>52)</sup> and a case-control study of *HLA-B\*5701* in both Caucasians and African Americans (SHAPE study),<sup>53)</sup> were highly supportive of the use of *HLA-B\*5701* screening in clinical practice to exclude *HLA-B\*5701* carriers from patients treated with abacavir. However, 7 Japanese patients, who were all *HLA-B\*5701*-negative, had abacavir-induced hypersensitivity reactions.<sup>40)</sup> Thus, the genetic screening of *HLA-B\*5701* is unlikely to be effective for the Japanese.

Nevirapine is a non-nucleoside reverse transcriptase inhibitor that is used in combination with antiretroviral therapy.<sup>54)</sup> The major treatment-limiting toxicity associated with nevirapine use is skin rash and hypersensitivity, which emerge in 5% of patients who have initiated nevirapine therapy.<sup>55)</sup> Nevirapine is metabolized by *CYP3A4* predominantly, and to a lesser degree by *CYP2B6* and other *CYP* isoforms, to several hydroxylated metabolites: 12-hydroxynevirapine has been implicated as a putative nevirapine metabolite causing hypersensitivity reactions.<sup>56)</sup>  $CD4^+$  T cells have been shown to be involved in the nevirapine-induced hypersensitivity reaction.<sup>57,58)</sup> Thus, a high level of  $CD4^+$  T cells (more than 25% above normal) is one risk factor for nevirapine-induced hypersensitivity reaction.<sup>55)</sup> In addition to  $CD4^+$  T cell levels, associations between several types of HLA and

nevirapine-induced hypersensitivity reactions have been reported in different countries or populations. For example, *HLA-DRB1\*0101* (Western Australian),<sup>59)</sup> *HLA-Cw\*0802-B\*1402* haplotype (Sardinian patients, people in Italian autonomous regions),<sup>60)</sup> *HLA-Cw8* (Japanese),<sup>61)</sup> and *HLA-B\*3505* (Thai)<sup>62)</sup> have been reported (Table 5). These results imply that the primary determining *HLA* allele may be different among populations in the nevirapine-induced hypersensitivity reaction.

### Drug-induced liver injury

Several patterns of DILI exist, with the most useful classification being hepatocellular, cholestatic, or a combination of both.<sup>63)</sup> Hepatocellular injury involves marked elevation of serum ALT and AST levels, usually preceding an increase in total bilirubin and no increase or only modest increases in alkaline phosphatase (ALP) levels.<sup>9)</sup> In cholestatic injury, increases in ALP levels predominate and precede increases in ALT and AST.<sup>9)</sup> The patterns in which a drug causes liver injury are regarded as being either predictable (dose-dependent) or unpredictable (idiosyncratic).<sup>64)</sup> Acetaminophen-induced hepatotoxicity has been considered the classic example of a dose-related hepatotoxin,<sup>65)</sup> although few other drugs fit this pattern.<sup>64)</sup> Rather, the majority of drugs that are capable of producing liver injury do so in an unpredictable fashion with variable latency periods.<sup>9,66)</sup>

Single nucleotide polymorphisms (SNPs) in drug metabolizing enzymes and drug transporters, which regulate the metabolism and disposition of drugs, represent the best studied set of pharmaceutically important genetic markers of DILI. *N*-Acetyltransferase (NAT) functions by acetylating drugs, therefore causing active drug metabolites to be detoxified. Deficient alleles of *NAT* (such as *NAT1\*14* and *\*15* and *NAT2\*5*, *\*6*, and *\*7*), which reduce detoxification activity, increase the toxicity of drugs including isoniazid, sulfonamides, and procainamides.<sup>67,68)</sup> The allele distribution of the Caucasian population differs from that reported in the Japanese population.<sup>67,69)</sup> Another enzyme pathway of importance is that of glutathione in the detoxification of reactive metabolites. Genetically determined deficiencies in glutathione synthetase and GST have been associated with increased hepatotoxicity of certain drugs, including acetaminophen, metronidazole, and nitrofurantoin.<sup>70,71)</sup> Frequencies of GST-deficient alleles show ethnic differences. For example, the homozygous deletion genotype frequency in *GSTM1* ranges from 0.38 to 0.67 in Caucasians, from 0.33 to 0.63 in East Asians, and from 0.22 to 0.35 in Africans and African-Americans.<sup>72)</sup> Pacific Islanders have the highest reported frequency of homozygous deletion genotypes (0.64–1.0) of any group studied.<sup>73,74)</sup> Troglitazone is a 2,4-thiazolidinedione anti-diabetic drug with insulin-sensitizing activities.<sup>75,76)</sup> Troglitazone-associated idiosyncratic hepatic dysfunction and hepatic

**Table 6.** Association between drug-induced liver injury and genetic polymorphisms

Drug	Gene variant	Population	OR <sup>a</sup>	Pc <sup>b</sup>	Ref
Troglitazone	GSTM1/T1	Japanese	3.69	0.008	81)
Ticlopidine	HLA-A*3303	Japanese	36.5	$7.32 \times 10^{-7}$	86)
Diclofenac	UGT2B7*2	Unknown	8.5	0.03	82)
	ABCC2(C-24T)	Unknown	5.0	0.005	
	CYP2C8 haplotypes	Unknown	—	0.04	
Flucloxacillin	HLA-B*5701	European	45.0	$8.7 \times 10^{-33}$	21)

<sup>a</sup> OR is the odds ratio.

<sup>b</sup> Pc indicates corrected P value.

failure were reported after introduction of the drug into the market.<sup>77–79)</sup> Yamamoto *et al.* reported that CYP3A4 catalyzed troglitazone into an epoxide of a quinone metabolite which may be eliminated by GSTs and EPHX.<sup>80)</sup> To address the susceptible genetic factors responsible for the hepatotoxicity associated with the drug, Watanabe *et al.* performed a genetic polymorphic analysis by a target gene approach in troglitazone-treated Japanese patients with type 2 diabetes mellitus<sup>81)</sup> (Table 6). They observed a correlation between hepatic failure and both *GSTT1* and *GSTM1* null genotypes. They reported that the odds ratio was 3.692 and its 95% confidence interval was 1.354 to 10.066 (Pc = 0.008). A more recent example of drug hepatotoxicity resulting from genetic polymorphisms of drug metabolizing enzymes and drug transporters is that of diclofenac, a non-steroidal anti-inflammatory drug that is among the most common drugs to cause idiosyncratic hepatotoxicity. Diclofenac-induced hepatotoxicity occurs at a rate of 6 per 100,000 users, and 8 to 20% of the patients who develop jaundice die of liver failure. It has been concluded that diclofenac hepatotoxicity is associated with the possession of variant UDP-glucuronosyltransferase 2B7 (*UGT2B7\*2*), ATP-binding cassette transporter C2 (*ABCC2*, –24C>T), and *CYP2C8* haplotypes<sup>82)</sup> (Table 6).

Immune-mediated mechanisms via the reactive metabolite binding to macromolecules are believed to be associated with idiosyncratic DILI. *HLA* has been considered to be involved in T-cell mediated cytotoxic reactions and drug-induced allergic reactions. Therefore, *HLA* might be another type of candidate genetic biomarker of DILI.<sup>64,66)</sup> Ticlopidine, an anti-platelet agent, which has been widely used for the secondary prevention of atherothrombosis,<sup>83)</sup> has shown severe hepatotoxicity, mainly of the cholestatic type,<sup>84)</sup> and there appears to be an increased rate of hepatic adverse reactions in Japanese compared with Caucasian patients.<sup>85)</sup> Hirata *et al.* explored genetic risk factors for ticlopidine-induced hepatotoxicity using 22 Japanese patients with ticlopidine-induced hepatotoxicity and 85 Japanese patients who tolerated ticlopidine therapy without ex-

periencing adverse reactions and they found a significant correlation between ticlopidine-induced hepatotoxicity and *HLA-A\*3303*<sup>86)</sup> (Table 6). Allelic frequency of *HLA-A\*3303* is 7.54% in Japanese, 0.6% in Caucasians, and 4.5% in African-Americans (<http://www.allelefrequencies.net/>). They reported that 12 patients (86%) among 14 patients who showed ticlopidine-induced cholestatic hepatotoxicity had *HLA-A\*3303* and the odds ratio was 36.50 (95% confidence interval, 7.25 to 183.82). Another example of the correlation between *HLA* and DILI is that of flucloxacillin, which is widely used in many European countries and Australia for treatment of staphylococcal infection. Its use has been associated with a characteristic cholestatic hepatitis that is more common in females, the elderly, and patients with prolonged treatment courses.<sup>87-89)</sup> In the United Kingdom, the incidence of flucloxacillin-related DILI has been estimated at 8.5 in every 100,000 new users in days 1 to 45 after starting treatment.<sup>89)</sup> Daly *et al.* conducted a genome-wide association study using 51 cases (white European ancestry) of flucloxacillin-related DILI and 282 controls matched for sex and ancestry and found the strongest correlation between flucloxacillin-related DILI and a genetic marker (rs2395029) in complete linkage disequilibrium with *HLA-B\*5701*<sup>21)</sup> (Table 6). They reported that the odds ratio was 45 (95% confidence interval, 19.4 to 105). Among 51 cases, 43 patients (84%) carried the risk allele (G), which has a frequency of approximately 5% in the population controls and in European populations generally.

### Statin-induced myopathy

In rare cases, statins can cause muscle pain or weakness in association with elevated creatine kinase levels (*i.e.*, myopathy), and occasionally this leads to muscle breakdown and myoglobin release (*i.e.*, rhabdomyolysis), with a risk of kidney failure and death.<sup>90)</sup> The mechanisms by which statins cause myopathy remain unknown but appear to be related to statin concentrations in blood and muscle.<sup>90)</sup>

Morimoto *et al.* studied genetic factors contributing to the risk of statin-induced myopathy and showed that the frequencies of *OATP-C\*15* [tagged by 388A>G (N130D) and 521T>C, V174A], a mutant allele of *OATP-C* (*OATP1B1*, *SLC21A6/SLC01B1*) was significantly higher in Japanese patients with myopathy who were receiving pravastatin or atorvastatin than in patients without myopathy<sup>91)</sup> (Table 7). They also found another *OATP-C* mutant allele, 1628T>G (L543W), which is located in exon 12 of *SLC21A6/SLC01B1* in a Japanese patient with pravastatin-induced myopathy<sup>92)</sup> (Table 7). They examined the transporting activity for pravastatin and other substrates and found that the activity decreased significantly in HEK293 cells expressing mutant proteins with V174A and L543W compared to those in cells ex-

**Table 7.** Candidates of *OATP-C* genomic biomarkers for statin-induced myopathy

Variant	<i>In vitro</i> activity	Statin	Ref
521T>C	Decrease	Pravastatin or atorvastatin	91)
521T>C	Decrease	Simvastatin	22)
1628T>G	Decrease	Pravastatin	91, 94)

pressing *OATP-C\*1a*, the reference allele of *OATP-C*.<sup>93,94)</sup> *OATP-C* has been shown to mediate the hepatic uptake of statins.<sup>95)</sup> From these results, they speculated that patients who are carrying these defective *OATP-C* mutant alleles have increased plasma concentrations of these statins and are thus more susceptible to the myotoxic effects of these statins compared to non-carrier patients treated with pravastatin and atorvastatin.<sup>92)</sup> In fact, Ide *et al.* recently reported that *OATP-C\*15* significantly influenced the relative bioavailability [F(rel)] of pravastatin; F(rel) was increased 1.50- and 1.95-fold in heterozygous and homozygous participants, respectively, for the *OATP-C\*15* allele in comparison with participants without the allele from a covariate analysis of population pharmacokinetic analysis.<sup>96)</sup>

The SEARCH Collaborative Group, which aims to determine whether a daily dose of 80 mg of simvastatin safely produces greater benefit than a daily dose of 20 mg, found 98 definite or incipient cases of myopathy among 6,031 participants who were assigned to receive 80 mg of simvastatin.<sup>22)</sup> All participants were from the United Kingdom, but their ethnicity was not specified. They performed a genome-wide association study using approximately 300,000 genomic markers in 85 subjects with definite or incipient myopathy and 90 controls, and found a single strong association of myopathy with the rs4149056 (521T>C) SNP located within *SLC01B1* (Table 7). They reported that the odds ratio for myopathy was 4.5 (95% confidence interval, 2.6 to 7.7) per copy of the C allele, and 16.9 (95% confidence interval, 4.7 to 61.1) in CC as compared with TT homozygotes. They concluded that this variant of *SLC01B1* is strongly associated with an increased risk of statin-induced myopathy.

These studies suggest that variant *OATP-C* decreased the hepatic uptake of statin and increased blood and muscle concentrations of statin. The increase of the blood and muscle concentrations of statin may cause myopathy or rhabdomyolysis. Genotyping these *SLC01B1* variants might help to achieve the benefits of statin therapy more safely.

### Use of pharmacogenetic biomarkers in clinical practice

In principle, identifying genetic risk factors for severe ADRs, particularly type B reactions, could significantly

decrease the incidence rate of ADRs and improve the process of drug development.<sup>97)</sup> Among these type B ADRs which we consider in this review, the usefulness of abacavir *HLA*-genetic biomarker (*HLA-B\*5701*) has been confirmed in Caucasians from several prospective studies, such as the PREDICT-1 study.<sup>52)</sup> The association of CBZ-induced SJS/TEN and an *HLA*-genetic biomarker (*HLA-B\*1502*) in Han Chinese is extremely high compared with other drugs.<sup>25)</sup> Therefore, *HLA-B\*1502* screening is recommended for CBZ in clinical practice by the US FDA, and *HLA-B\*5701* screening is recommended for abacavir by the US FDA and European Medical Agency. Before treatment with CBZ or abacavir, *HLA* analysis should be performed to exclude *HLA-B\*1502* or *HLA-B\*5701* unless the patient is from a population who shows extremely low frequency of these *HLA* types. Such exclusion of patients from treatment with causative drugs would markedly reduce the possibility of severe ADRs and prevent overestimation of severe cutaneous ADRs that could otherwise result in excessive discontinuation of treatment.<sup>34,98,99)</sup> In Japan, package inserts of CBZ and abacavir describe these research results.

#### Perspective on pharmacogenetic biomarkers

An unresolved issue for genetic biomarkers is ethnic differences, since these *HLA* markers show ethnic specificity. For example, an *HLA* marker of abacavir (*B\*5701*) or CBZ (*B\*1502*) is present only in Caucasians or Han Chinese (and South East Asians), respectively, and its usefulness has not been shown in other populations such as Japanese.<sup>37,40)</sup> On the other hand, the association between allopurinol treatment and *HLA-B\*5801* was observed not only in Han Chinese but also in Caucasians and Japanese, although the odds ratios were lower than that of CBZ.<sup>5,36,44)</sup> Therefore, it is absolutely necessary to explore the *HLA* marker for each population against each drug and also to find the universal genetic biomarker (if one exists) of severe ADRs for clinical practice. As shown for nevirapine, the association between the rash with hepatitis and *HLA-DRB1\*0101* was observed in Western Australian patients with CD4<sup>+</sup> T-cell levels greater than 25% above normal levels.<sup>59)</sup> This case suggests that the combination of *HLA* genetic biomarkers and other biomarkers might be useful to predict ADRs for some drugs. A prospective study or comparative study with other populations is necessary for *HLA* biomarkers of ticlopidine- and flucloxacillin-related DILI and *SLCO1B1* biomarkers of statin-induced myopathy.

In most cases of allergic reactions, such as SJS/TEN and DILI, *HLA*-drug toxicity associations are thought to arise as a result of the interaction of a specific *HLA* allele with the drug or its metabolite, causing an immune reaction to be triggered.<sup>38,100,101)</sup> As shown in the abacavir-induced hypersensitivity reaction, the drug metabolite may play an important role in the allergic reaction process,<sup>102)</sup>

suggesting that sequential reactions from drug metabolism to the immune mechanism can exist in the allergic process. Thus, drug toxicities that are driven primarily by the immune response may require bioactivation of the drug to a specific metabolite to evoke the specific immune response that will lead to the generation of an adverse reaction.<sup>38,100,101)</sup> These complex mechanisms may be involved in most cases of allergic reactions because reactive metabolites have been detected not only in abacavir but also in nevirapine, CBZ, and allopurinol.

Most of the currently available genetic biomarkers are limited in relation to *HLA*, drug metabolizing enzymes, and drug transporters.<sup>17)</sup> Considering that the technology to identify genetic variants across the whole genome is advancing rapidly, many more significant genetic factors for ADRs are likely to be identified in the future. In such whole-genome case-control analysis, there might be critical points to resolve. The first problem is the size of case and control groups. Accrual of large numbers of cases is necessary for genome-wide association study of genetic factors underlying severe ADRs, even though the number of patients with specific types of ADRs is small.<sup>103)</sup> Control subjects for such studies should be matched for drug exposure, concomitant use of other drugs that could affect the pharmacokinetics and pharmacodynamics of the drug in question, and subject background such as age, gender, and ethnicity.<sup>103)</sup> The second problem is objective diagnosis of ADRs. Because one drug could induce many ADR phenotypes in which the mechanism may be different, standardization of diagnosis is necessary.<sup>40,103)</sup> These critical points affect the sufficient statistical power to detect the genetic biomarker. In order to resolve these problems, several regional networks to study severe ADRs have been established, including our research group in Japan (for SJS/TEN, DILI, and myopathy in Japanese),<sup>36)</sup> European collaboration for studying the genetic basis of adverse drug reactions (EUDRAGENE for six severe ADRs in multiple European populations),<sup>104)</sup> the United States Drug Induced Liver Injury Network (DILIN, for DILI),<sup>105)</sup> and the International Serious Adverse Event Consortium (SAEC, for SJS/TEN and DILI in global populations).<sup>21)</sup> These networks involve scientists in regulatory agencies, healthcare systems, and pharmaceutical industries as well as academia. Moreover, for the goal of standardizing phenotypes and comparing ethnicity regarding genetic risk factors for severe ADRs, these networks may form a global consortium together with new networks from other communities in the future.

#### Conclusions

Specific types of *HLA* which showed strong association with severe cutaneous ADRs and DILI have been found as candidate pharmacogenetic biomarkers for each ADR. The *HLA* type was different for different causative drugs,



and the allelic frequency of *HLA* genetic polymorphisms showed ethnic differences. The genetic polymorphism of drug transporter gene *SLCO1B1* has been shown to be associated with statin-induced myopathy. It is necessary to conduct prospective studies to establish valid pharmacogenetic biomarkers for severe ADRs. A large research network for the collection of DNA samples from patients with ADRs is also necessary to explore a variety of pharmacogenetic biomarkers for ADRs.

### References

- 1) Rawlins, M. D. and Thompson, J. W.: Pathogenesis of adverse drug reactions. In Davis, D. M. (ed.): *Text book of adverse drug reactions*, Oxford, Oxford University Press, 1977, pp.10.
- 2) Bastuji-Garin, S., Rzany, B., Stern, R. S., Shear, N. H., Naldi, L. and Roujeau, J. C.: Clinical classification of cases of toxic epidermal necrolysis, Stevens-Johnson syndrome, and erythema multiforme. *Arch. Dermatol.*, **129**: 92–96 (1993).
- 3) Roujeau, J. C. and Stern, R. S.: Severe adverse cutaneous reactions to drugs. *N. Engl. J. Med.*, **331**: 1272–1285 (1994).
- 4) Chave, T. A., Mortimer, N. J., Sladden, M. J., Hall, A. P. and Hutchinson, P. E.: Toxic epidermal necrolysis: current evidence, practical management and future directions. *Br. J. Dermatol.*, **153**: 241–253 (2005).
- 5) Lonjou, C., Borot, N., Sekula, P., Ledger, N., Thomas, L., Halevy, S., Naldi, L., Bouwes-Bavinck, J. N., Sidoroff, A., de Toma, C., Schumacher, M., Roujeau, J. C., Hovnanian, A. and Mockenhaupt, M.: A European study of HLA-B in Stevens-Johnson syndrome and toxic epidermal necrolysis related to five high-risk drugs. *Pharmacogenet. Genomics*, **18**: 99–107 (2008).
- 6) Roujeau, J. C.: Stevens-Johnson syndrome and toxic epidermal necrolysis are severity variants of the same disease which differs from erythema multiforme. *J. Dermatol.*, **24**: 726–729 (1997).
- 7) Chung, W. H., Hung, S. I., Yang, J. Y., Su, S. C., Huang, S. P., Wei, C. Y., Chin, S. W., Chiou, C. C., Chu, S. C., Ho, H. C., Yang, C. H., Lu, C. F., Wu, J. Y., Liao, Y. D. and Chen, Y. T.: Granulysin is a key mediator for disseminated keratinocyte death in Stevens-Johnson syndrome and toxic epidermal necrolysis. *Nat. Med.*, **14**: 1343–1350 (2008).
- 8) Murata, J. and Abe, R.: Soluble Fas ligand: is it a critical mediator of toxic epidermal necrolysis and Stevens-Johnson syndrome? *J. Invest. Dermatol.*, **127**: 744–745 (2007).
- 9) Haleboua-De Marzio, D. and Navarro, V. J.: Drug-induced hepatotoxicity in humans. *Curr. Opin. Drug Discov. Devel.*, **11**: 53–59 (2008).
- 10) Lucena, M. I., Andrade, R. J., Kaplowitz, N., Garcia-Cortes, M., Fernandez, M. C., Romero-Gomez, M., Bruguera, M., Hallal, H., Robles-Diaz, M., Rodriguez-Gonzalez, J. F., Navarro, J. M., Salmeron, J., Martinez-Odrizola, P., Perez-Alvarez, R., Borraz, Y. and Hidalgo, R.: Phenotypic characterization of idiosyncratic drug-induced liver injury: the influence of age and sex. *Hepatology*, **49**: 2001–2009 (2009).
- 11) Fontana, R. J., Watkins, P. B., Bonkovsky, H. L., Chalasani, N., Davern, T., Serrano, J. and Rochon, J.: Drug-Induced Liver Injury Network (DILIN) prospective study: rationale, design and conduct. *Drug Saf.*, **32**: 55–68 (2009).
- 12) Bosch, X., Poch, E. and Grau, J. M.: Rhabdomyolysis and acute kidney injury. *N. Engl. J. Med.*, **361**: 62–72 (2009).
- 13) Psaty, B. M., Furberg, C. D., Ray, W. A. and Weiss, N. S.: Potential for conflict of interest in the evaluation of suspected adverse drug reactions: use of cerivastatin and risk of rhabdomyolysis. *JAMA*, **292**: 2622–2631 (2004).
- 14) Staffa, J. A., Chang, J. and Green, L.: Cerivastatin and reports of fatal rhabdomyolysis. *N. Engl. J. Med.*, **346**: 539–540 (2002).
- 15) Edwards, S. G., Hubbard, V., Aylett, S. and Wren, D.: Concordance of primary generalised epilepsy and carbamazepine hypersensitivity in monozygotic twins. *Postgrad. Med. J.*, **75**: 680–681 (1999).
- 16) Gennis, M. A., Vemuri, R., Burns, E. A., Hill, J. V., Miller, M. A. and Spielberg, S. P.: Familial occurrence of hypersensitivity to phenytoin. *Am. J. Med.*, **91**: 631–634 (1991).
- 17) Nakamura, Y.: Pharmacogenomics and drug toxicity. *N. Engl. J. Med.*, **359**: 856–858 (2008).
- 18) Pirmohamed, M. and Park, B. K.: Genetic susceptibility to adverse drug reactions. *Trends Pharmacol. Sci.*, **22**: 298–305 (2001).
- 19) Rudolph, M. G., Stanfield, R. L. and Wilson, I. A.: How TCRs bind MHCs, peptides, and coreceptors. *Annu. Rev. Immunol.*, **24**: 419–466 (2006).
- 20) Horton, R., Wilming, L., Rand, V., Lovering, R. C., Bruford, E. A., Khodiyar, V. K., Lush, M. J., Povey, S., Talbot, C. C., Jr., Wright, M. W., Wain, H. M., Trowsdale, J., Ziegler, A. and Beck, S.: Gene map of the extended human MHC. *Nat. Rev. Genet.*, **5**: 889–899 (2004).
- 21) Daly, A. K., Donaldson, P. T., Bhatnagar, P., Shen, Y., Pe'er, I., Floratos, A., Daly, M. J., Goldstein, D. B., John, S., Nelson, M. R., Graham, J., Park, B. K., Dillon, J. F., Bernal, W., Cordell, H. J., Pirmohamed, M., Aithal, G. P. and Day, C. P.: HLA-B\*5701 genotype is a major determinant of drug-induced liver injury due to flucloxacillin. *Nat. Genet.*, **41**: 816–819 (2009).
- 22) Link, E., Parish, S., Armitage, J., Bowman, L., Heath, S., Matsuda, F., Gut, I., Lathrop, M. and Collins, R.: *SLCO1B1* variants and statin-induced myopathy—a genomewide study. *N. Engl. J. Med.*, **359**: 789–799 (2008).
- 23) Kerr, B. M., Thummel, K. E., Wurden, C. J., Klein, S. M., Kroetz, D. L., Gonzalez, F. J. and Levy, R. H.: Human liver carbamazepine metabolism. Role of CYP3A4 and CYP2C8 in 10,11-epoxide formation. *Biochem. Pharmacol.*, **47**: 1969–1979 (1994).
- 24) Madden, S., Maggs, J. L. and Park, B. K.: Bioactivation of carbamazepine in the rat in vivo. Evidence for the formation of reactive arene oxide(s). *Drug Metab. Dispos.*, **24**: 469–479 (1996).
- 25) Chung, W. H., Hung, S. I., Hong, H. S., Hsieh, M. S., Yang, L. C., Ho, H. C., Wu, J. Y. and Chen, Y. T.: Medical genetics: a marker for Stevens-Johnson syndrome. *Nature*, **428**: 486 (2004).
- 26) Kamaliah, M. D., Zainal, D., Mokhtar, N. and Nazmi, N.: Erythema multiforme, Stevens-Johnson syndrome and toxic epidermal necrolysis in northeastern Malaysia. *Int. J. Dermatol.*, **37**: 520–523 (1998).
- 27) Khoo, A. K. and Foo, C. L.: Toxic epidermal necrolysis in a burns centre: a 6-year review. *Burns*, **22**: 275–278 (1996).
- 28) Tennis, P. and Stern, R. S.: Risk of serious cutaneous disorders after initiation of use of phenytoin, carbamazepine, or sodium

- valproate: a record linkage study. *Neurology*, **49**: 542–546 (1997).
- 29) Vittorio, C. C. and Muglia, J. J.: Anticonvulsant hypersensitivity syndrome. *Arch. Intern. Med.*, **155**: 2285–2290 (1995).
  - 30) Bellucci, G., Berti, G., Chiappe, C., Lippi, A. and Marioni, F.: The metabolism of carbamazepine in humans: steric course of the enzymatic hydrolysis of the 10,11-epoxide. *J. Med. Chem.*, **30**: 768–773 (1987).
  - 31) Lillibridge, J. H., Amore, B. M., Slattery, J. T., Kalhorn, T. F., Nelson, S. D., Finnell, R. H. and Bennett, G. D.: Protein-reactive metabolites of carbamazepine in mouse liver microsomes. *Drug Metab. Dispos.*, **24**: 509–514 (1996).
  - 32) Gaedigk, A., Spielberg, S. P. and Grant, D. M.: Characterization of the microsomal epoxide hydrolase gene in patients with anticonvulsant adverse drug reactions. *Pharmacogenetics*, **4**: 142–153 (1994).
  - 33) Green, V. J., Pirmohamed, M., Kitteringham, N. R., Gaedigk, A., Grant, D. M., Boxer, M., Burchell, B. and Park, B. K.: Genetic analysis of microsomal epoxide hydrolase in patients with carbamazepine hypersensitivity. *Biochem. Pharmacol.*, **50**: 1353–1359 (1995).
  - 34) Hung, S. I., Chung, W. H., Jee, S. H., Chen, W. C., Chang, Y. T., Lee, W. R., Hu, S. L., Wu, M. T., Chen, G. S., Wong, T. W., Hsiao, P. F., Chen, W. H., Shih, H. Y., Fang, W. H., Wei, C. Y., Lou, Y. H., Huang, Y. L., Lin, J. J. and Chen, Y. T.: Genetic susceptibility to carbamazepine-induced cutaneous adverse drug reactions. *Pharmacogenet. Genomics*, **16**: 297–306 (2006).
  - 35) Man, C. B., Kwan, P., Baum, L., Yu, E., Lau, K. M., Cheng, A. S. and Ng, M. H.: Association between HLA-B\*1502 allele and antiepileptic drug-induced cutaneous reactions in Han Chinese. *Epilepsia*, **48**: 1015–1018 (2007).
  - 36) Kaniwa, N., Saito, Y., Aihara, M., Matsunaga, K., Tohkin, M., Kurose, K., Sawada, J., Furuya, H., Takahashi, Y., Muramatsu, M., Kinoshita, S., Abe, M., Ikeda, H., Kashiwagi, M., Song, Y., Ueta, M., Sotozono, C., Ikezawa, Z. and Hasegawa, R.: HLA-B locus in Japanese patients with anti-epileptics and allopurinol-related Stevens-Johnson syndrome and toxic epidermal necrolysis. *Pharmacogenomics*, **9**: 1617–1622 (2008).
  - 37) Lonjou, C., Thomas, L., Borot, N., Ledger, N., de Toma, C., Krouget, H., Graf, E., Schumacher, M., Hovnanian, A., Mockenhaupt, M. and Roujeau, J. C.: A marker for Stevens-Johnson syndrome: ethnicity matters. *Pharmacogenomics J.*, **6**: 265–268 (2006).
  - 38) Phillips, E. J. and Mallal, S. A.: HLA and drug-induced toxicity. *Curr. Opin. Mol. Ther.*, **11**: 231–242 (2009).
  - 39) Tanaka, H., Akaza, T. and Juji, T.: Report of the Japanese Central Bone Marrow Data Center. *Clin. Transpl.*, 139–144 (1996).
  - 40) Gatanaga, H., Honda, H. and Oka, S.: Pharmacogenetic information derived from analysis of HLA alleles. *Pharmacogenomics*, **9**: 207–214 (2008).
  - 41) Wortmann, R. L.: Gout and hyperuricemia. *Curr. Opin. Rheumatol.*, **14**: 281–286 (2002).
  - 42) Marr, J. J., Berens, R. L., Nelson, D. J., Krenitsky, T. A., Spector, T., LaFon, S. W. and Elion, G. B.: Antileishmanial action of 4-thiopyrazolo (3,4-d) pyrimidine and its ribonucleoside. Biological effects and metabolism. *Biochem. Pharmacol.*, **31**: 143–148 (1982).
  - 43) Chung, W. H., Hung, S. I. and Chen, Y. T.: Human leukocyte antigens and drug hypersensitivity. *Curr. Opin. Allergy Clin. Immunol.*, **7**: 317–323 (2007).
  - 44) Hung, S. I., Chung, W. H., Liou, L. B., Chu, C. C., Lin, M., Huang, H. P., Lin, Y. L., Lan, J. L., Yang, L. C., Hong, H. S., Chen, M. J., Lai, P. C., Wu, M. S., Chu, C. Y., Wang, K. H., Chen, C. H., Fann, C. S., Wu, J. Y. and Chen, Y. T.: HLA-B\*5801 allele as a genetic marker for severe cutaneous adverse reactions caused by allopurinol. *Proc. Natl. Acad. Sci. USA*, **102**: 4134–4139 (2005).
  - 45) Hetherington, S., McGuirk, S., Powell, G., Cutrell, A., Naderer, O., Spreen, B., Lafon, S., Pearce, G. and Steel, H.: Hypersensitivity reactions during therapy with the nucleoside reverse transcriptase inhibitor abacavir. *Clin. Ther.*, **23**: 1603–1614 (2001).
  - 46) Walsh, J. S., Reese, M. J. and Thurmond, L. M.: The metabolic activation of abacavir by human liver cytosol and expressed human alcohol dehydrogenase isozymes. *Chem. Biol. Interact.*, **142**: 135–154 (2002).
  - 47) Hetherington, S., Hughes, A. R., Mosteller, M., Shortino, D., Baker, K. L., Spreen, W., Lai, E., Davies, K., Handley, A., Dow, D. J., Fling, M. E., Stocum, M., Bowman, C., Thurmond, L. M. and Roses, A. D.: Genetic variations in HLA-B region and hypersensitivity reactions to abacavir. *Lancet*, **359**: 1121–1122 (2002).
  - 48) Mallal, S., Nolan, D., Witt, C., Masel, G., Martin, A. M., Moore, C., Sayer, D., Castley, A., Mamotte, C., Maxwell, D., James, I. and Christiansen, F. T.: Association between presence of HLA-B\*5701, HLA-DR7, and HLA-DQ3 and hypersensitivity to HIV-1 reverse-transcriptase inhibitor abacavir. *Lancet*, **359**: 727–732 (2002).
  - 49) Hughes, A. R., Mosteller, M., Bansal, A. T., Davies, K., Haneline, S. A., Lai, E. H., Nangle, K., Scott, T., Spreen, W. R., Warren, L. L. and Roses, A. D.: Association of genetic variations in HLA-B region with hypersensitivity to abacavir in some, but not all, populations. *Pharmacogenomics*, **5**: 203–211 (2004).
  - 50) Martin, A. M., Nolan, D., Gaudieri, S., Almeida, C. A., Nolan, R., James, I., Carvalho, F., Phillips, E., Christiansen, F. T., Purcell, A. W., McCluskey, J. and Mallal, S.: Predisposition to abacavir hypersensitivity conferred by HLA-B\*5701 and a haplotypic Hsp70-Hom variant. *Proc. Natl. Acad. Sci. USA*, **101**: 4180–4185 (2004).
  - 51) Martin, A. M., Almeida, C. A., Cameron, P., Purcell, A. W., Nolan, D., James, I., McCluskey, J., Phillips, E., Landay, A. and Mallal, S.: Immune responses to abacavir in antigen-presenting cells from hypersensitive patients. *AIDS*, **21**: 1233–1244 (2007).
  - 52) Mallal, S., Phillips, E., Carosi, G., Molina, J. M., Workman, C., Tomazic, J., Jagel-Guedes, E., Rugina, S., Kozyrev, O., Cid, J. F., Hay, P., Nolan, D., Hughes, S., Hughes, A., Ryan, S., Fitch, N., Thorborn, D. and Benbow, A.: HLA-B\*5701 screening for hypersensitivity to abacavir. *N. Engl. J. Med.*, **358**: 568–579 (2008).
  - 53) Saag, M., Balu, R., Phillips, E., Brachman, P., Martorell, C., Burman, W., Stancil, B., Mosteller, M., Brothers, C., Wanamaker, P., Hughes, A., Sutherland-Phillips, D., Mallal, S. and Shaefer, M.: High sensitivity of human leukocyte antigen-b\*5701 as a marker for immunologically confirmed abacavir hypersensitivity in white and black patients. *Clin. Infect. Dis.*, **46**: 1111–1118 (2008).
  - 54) van der Valk, M., Kastelein, J. J., Murphy, R. L., van Leth, F.,

- Katlama, C., Horban, A., Glesby, M., Behrens, G., Clotet, B., Stellato, R. K., Molhuizen, H. O. and Reiss, P.: Nevirapine-containing antiretroviral therapy in HIV-1 infected patients results in an anti-atherogenic lipid profile. *AIDS*, **15**: 2407–2414 (2001).
- 55) Phillips, E. and Mallal, S.: Drug hypersensitivity in HIV. *Curr. Opin. Allergy Clin. Immunol.*, **7**: 324–330 (2007).
  - 56) Erickson, D. A., Mather, G., Trager, W. F., Levy, R. H. and Keirns, J. J.: Characterization of the in vitro biotransformation of the HIV-1 reverse transcriptase inhibitor nevirapine by human hepatic cytochromes P-450. *Drug Metab. Dispos.*, **27**: 1488–1495 (1999).
  - 57) Kiertiburanakul, S., Sungkanuparph, S., Charoenyingwattana, A., Mahasirimongkol, S., Sura, T. and Chantratita, W.: Risk factors for nevirapine-associated rash among HIV-infected patients with low CD4 cell counts in resource-limited settings. *Curr. HIV Res.*, **6**: 65–69 (2008).
  - 58) van Leth, F., Andrews, S., Grinsztejn, B., Wilkins, E., Lazanas, M. K., Lange, J. M. and Montaner, J.: The effect of baseline CD4 cell count and HIV-1 viral load on the efficacy and safety of nevirapine or efavirenz-based first-line HAART. *AIDS*, **19**: 463–471 (2005).
  - 59) Martin, A. M., Nolan, D., James, I., Cameron, P., Keller, J., Moore, C., Phillips, E., Christiansen, F. T. and Mallal, S.: Predisposition to nevirapine hypersensitivity associated with HLA-DRB1\*0101 and abrogated by low CD4 T-cell counts. *AIDS*, **19**: 97–99 (2005).
  - 60) Littera, R., Carcassi, C., Masala, A., Piano, P., Serra, P., Ortu, F., Corso, N., Casula, B., La Nasa, G., Contu, L. and Manconi, P. E.: HLA-dependent hypersensitivity to nevirapine in Sardinian HIV patients. *AIDS*, **20**: 1621–1626 (2006).
  - 61) Gatanaga, H., Yazaki, H., Tanuma, J., Honda, M., Genka, I., Teruya, K., Tachikawa, N., Kikuchi, Y. and Oka, S.: HLA-Cw8 primarily associated with hypersensitivity to nevirapine. *AIDS*, **21**: 264–265 (2007).
  - 62) Chantarangsu, S., Mushiroda, T., Mahasirimongkol, S., Kiertiburanakul, S., Sungkanuparph, S., Manosuthi, W., Tantisiriwat, W., Charoenyingwattana, A., Sura, T., Chantratita, W. and Nakamura, Y.: HLA-B\*3505 allele is a strong predictor for nevirapine-induced skin adverse drug reactions in HIV-infected Thai patients. *Pharmacogenet. Genomics*, **19**: 139–146 (2009).
  - 63) Navarro, V. J. and Senior, J. R.: Drug-related hepatotoxicity. *N. Engl. J. Med.*, **354**: 731–739 (2006).
  - 64) Pachkoria, K., Lucena, M. I., Molokhia, M., Cueto, R., Carballo, A. S., Carvajal, A. and Andrade, R. J.: Genetic and molecular factors in drug-induced liver injury: a review. *Curr. Drug Saf.*, **2**: 97–112 (2007).
  - 65) Laine, J. E., Auriola, S., Pasanen, M. and Juvonen, R. O.: Acetaminophen bioactivation by human cytochrome P450 enzymes and animal microsomes. *Xenobiotica*, **39**: 11–21 (2009).
  - 66) Andrade, R. J., Lucena, M. I., Alonso, A., Garcia-Cortes, M., Garcia-Ruiz, E., Benitez, R., Fernandez, M. C., Pelaez, G., Romero, M., Corpas, R., Duran, J. A., Jimenez, M., Rodrigo, L., Nogueras, F., Martin-Vivaldi, R., Navarro, J. M., Salmeron, J., de la Cuesta, F. S. and Hidalgo, R.: HLA class II genotype influences the type of liver injury in drug-induced idiosyncratic liver disease. *Hepatology*, **39**: 1603–1612 (2004).
  - 67) Hickman, D. and Sim, E.: N-acetyltransferase polymorphism. Comparison of phenotype and genotype in humans. *Biochem Pharmacol.*, **42**: 1007–1014 (1991).
  - 68) Huang, Y. S., Chern, H. D., Su, W. J., Wu, J. C., Lai, S. L., Yang, S. Y., Chang, F. Y. and Lee, S. D.: Polymorphism of the N-acetyltransferase 2 gene as a susceptibility risk factor for antituberculosis drug-induced hepatitis. *Hepatology*, **35**: 883–889 (2002).
  - 69) Deguchi, T., Mashimo, M. and Suzuki, T.: Correlation between acetylator phenotypes and genotypes of polymorphic arylamine N-acetyltransferase in human liver. *J. Biol. Chem.*, **265**: 12757–12760 (1990).
  - 70) Huang, Y. S., Su, W. J., Huang, Y. H., Chen, C. Y., Chang, F. Y., Lin, H. C. and Lee, S. D.: Genetic polymorphisms of manganese superoxide dismutase, NAD(P)H:quinone oxidoreductase, glutathione S-transferase M1 and T1, and the susceptibility to drug-induced liver injury. *J. Hepatol.*, **47**: 128–134 (2007).
  - 71) Spielberg, S. P.: In vitro assessment of pharmacogenetic susceptibility to toxic drug metabolites in humans. *Fed. Proc.*, **43**: 2308–2313 (1984).
  - 72) Rebbeck, T. R.: Molecular epidemiology of the human glutathione S-transferase genotypes GSTM1 and GSTT1 in cancer susceptibility. *Cancer Epidemiol. Biomarkers Prev.*, **6**: 733–743 (1997).
  - 73) Board, P., Coggan, M., Johnston, P., Ross, V., Suzuki, T. and Webb, G.: Genetic heterogeneity of the human glutathione transferases: a complex of gene families. *Pharmacol. Ther.*, **48**: 357–369 (1990).
  - 74) Lin, H. J., Han, C. Y., Bernstein, D. A., Hsiao, W., Lin, B. K. and Hardy, S.: Ethnic distribution of the glutathione transferase Mu 1-1 (GSTM1) null genotype in 1473 individuals and application to bladder cancer susceptibility. *Carcinogenesis*, **15**: 1077–1081 (1994).
  - 75) Iwamoto, Y., Kuzuya, T., Matsuda, A., Awata, T., Kumakura, S., Inooka, G. and Shiraishi, I.: Effect of new oral antidiabetic agent CS-045 on glucose tolerance and insulin secretion in patients with NIDDM. *Diabetes Care*, **14**: 1083–1086 (1991).
  - 76) Yoshioka, T., Fujita, T., Kanai, T., Aizawa, Y., Kurumada, T., Hasegawa, K. and Horikoshi, H.: Studies on hindered phenols and analogues. I. Hypolipidemic and hypoglycemic agents with ability to inhibit lipid peroxidation. *J. Med. Chem.*, **32**: 421–428 (1989).
  - 77) Herrine, S. K. and Choudhary, C.: Severe hepatotoxicity associated with troglitazone. *Ann. Intern. Med.*, **130**: 163–164 (1999).
  - 78) Shibuya, A., Watanabe, M., Fujita, Y., Saigenji, K., Kuwano, S., Takahashi, H. and Takeuchi, H.: An autopsy case of troglitazone-induced fulminant hepatitis. *Diabetes Care*, **21**: 2140–2143 (1998).
  - 79) Watkins, P. B. and Whitcomb, R. W.: Hepatic dysfunction associated with troglitazone. *N. Engl. J. Med.*, **338**: 916–917 (1998).
  - 80) Yamamoto, Y., Yamazaki, H., Ikeda, T., Watanabe, T., Iwabuchi, H., Nakajima, M. and Yokoi, T.: Formation of a novel quinone epoxide metabolite of troglitazone with cytotoxicity to HepG2 cells. *Drug Metab. Dispos.*, **30**: 155–160 (2002).
  - 81) Watanabe, I., Tomita, A., Shimizu, M., Sugawara, M., Yasuno, H., Koishi, R., Takahashi, T., Miyoshi, K., Nakamura, K., Izumi, T., Matsushita, Y., Furukawa, H., Haruyama, H. and Koga,

- T.: A study to survey susceptible genetic factors responsible for troglitazone-associated hepatotoxicity in Japanese patients with type 2 diabetes mellitus. *Clin. Pharmacol. Ther.*, **73**: 435–455 (2003).
- 82) Daly, A. K., Aithal, G. P., Leathart, J. B., Swainsbury, R. A., Dang, T. S. and Day, C. P.: Genetic susceptibility to diclofenac-induced hepatotoxicity: contribution of UGT2B7, CYP2C8, and ABCC2 genotypes. *Gastroenterology*, **132**: 272–281 (2007).
  - 83) Jacobson, A. K.: Platelet ADP receptor antagonists: ticlopidine and clopidogrel. *Best Pract. Res. Clin. Haematol.*, **17**: 55–64 (2004).
  - 84) Takikawa, H.: Lessons from ticlopidine-induced liver injury. *Hepatol. Res.*, **33**: 193–194 (2005).
  - 85) Mizushima, M., Iwata, N., Fujimoto, T. T., Ishikawa, K. and Fujimura, K.: Patient characteristics in ticlopidine hydrochloride-induced liver injury: Case-control study. *Hepatol. Res.*, **33**: 234–240 (2005).
  - 86) Hirata, K., Takagi, H., Yamamoto, M., Matsumoto, T., Nishiya, T., Mori, K., Shimizu, S., Masumoto, H. and Okutani, Y.: Ticlopidine-induced hepatotoxicity is associated with specific human leukocyte antigen genomic subtypes in Japanese patients: a preliminary case-control study. *Pharmacogenomics J.*, **8**: 29–33 (2008).
  - 87) Fairley, C. K., McNeil, J. J., Desmond, P., Smallwood, R., Young, H., Forbes, A., Purcell, P. and Boyd, I.: Risk factors for development of flucloxacillin associated jaundice. *BMJ*, **306**: 233–235 (1993).
  - 88) Olsson, R., Wiholm, B. E., Sand, C., Zettergren, L., Hultcrantz, R. and Myrhed, M.: Liver damage from flucloxacillin, cloxacillin and dicloxacillin. *J. Hepatol.*, **15**: 154–161 (1992).
  - 89) Russmann, S., Kaye, J. A., Jick, S. S. and Jick, H.: Risk of cholestatic liver disease associated with flucloxacillin and flucloxacillin prescribing habits in the UK: cohort study using data from the UK General Practice Research Database. *Br. J. Clin. Pharmacol.*, **60**: 76–82 (2005).
  - 90) Thompson, P. D., Clarkson, P. and Karas, R. H.: Statin-associated myopathy. *JAMA*, **289**: 1681–1690 (2003).
  - 91) Morimoto, K., Ueda, S., Seki, N., Igawa, Y., Kameyama, Y., Shimizu, A., Oishi, T., Hosokawa, M., Iesato, K., Mori, S., Saito, Y. and Chiba, K.: OATP-C(OATP-1B1)\*15 is associated with statin-induced myopathy in hypercholesterolemic patients. *Clin. Pharmacol. Ther.*, **77**: P21 (2005).
  - 92) Morimoto, K., Oishi, T., Ueda, S., Ueda, M., Hosokawa, M. and Chiba, K.: A novel variant allele of OATP-C (SLCO1B1) found in a Japanese patient with pravastatin-induced myopathy. *Drug Metab. Pharmacokinet.*, **19**: 453–455 (2004).
  - 93) Furihata, T., Satoh, N., Ohishi, T., Ugajin, M., Kameyama, Y., Morimoto, K., Matsumoto, S., Yamashita, K., Kobayashi, K. and Chiba, K.: Functional analysis of a mutation in the SLCO1B1 gene (c.1628T>G) identified in a Japanese patient with pravastatin-induced myopathy. *Pharmacogenomics J.*, **9**: 185–193 (2009).
  - 94) Kameyama, Y., Yamashita, K., Kobayashi, K., Hosokawa, M. and Chiba, K.: Functional characterization of SLCO1B1 (OATP-C) variants, SLCO1B1\*5, SLCO1B1\*15 and SLCO1B1\*15+C1007G, by using transient expression systems of HeLa and HEK293 cells. *Pharmacogenet. Genomics*, **15**: 513–522 (2005).
  - 95) Nishizato, Y., Ieiri, I., Suzuki, H., Kimura, M., Kawabata, K., Hirota, T., Takane, H., Irie, S., Kusuhaara, H., Urasaki, Y., Urae, A., Higuchi, S., Otsubo, K. and Sugiyama, Y.: Polymorphisms of OATP-C (SLC21A6) and OAT3 (SLC22A8) genes: consequences for pravastatin pharmacokinetics. *Clin. Pharmacol. Ther.*, **73**: 554–565 (2003).
  - 96) Ide, T., Sasaki, T., Maeda, K., Higuchi, S., Sugiyama, Y. and Ieiri, I.: Quantitative population pharmacokinetic analysis of pravastatin using an enterohepatic circulation model combined with pharmacogenomic information on SLCO1B1 and ABCC2 polymorphisms. *J. Clin. Pharmacol.*, **49**: 1309–1317 (2009).
  - 97) Giacomini, K. M., Krauss, R. M., Roden, D. M., Eichelbaum, M., Hayden, M. R. and Nakamura, Y.: When good drugs go bad. *Nature*, **446**: 975–977 (2007).
  - 98) Easterbrook, P. J., Waters, A., Murad, S., Ives, N., Taylor, C., King, D., Vyakarnam, A. and Thorburn, D.: Epidemiological risk factors for hypersensitivity reactions to abacavir. *HIV Med.*, **4**: 321–324 (2003).
  - 99) Rauch, A., Nolan, D., Martin, A., McKinnon, E., Almeida, C. and Mallal, S.: Prospective genetic screening decreases the incidence of abacavir hypersensitivity reactions in the Western Australian HIV cohort study. *Clin. Infect. Dis.*, **43**: 99–102 (2006).
  - 100) Lavergne, S. N., Park, B. K. and Naisbitt, D. J.: The roles of drug metabolism in the pathogenesis of T-cell-mediated drug hypersensitivity. *Curr. Opin. Allergy Clin. Immunol.*, **8**: 299–307 (2008).
  - 101) Park, B. K., Naisbitt, D. J., Gordon, S. F., Kitteringham, N. R. and Pirmohamed, M.: Metabolic activation in drug allergies. *Toxicology*, **158**: 11–23 (2001).
  - 102) Chessman, D., Kostenko, L., Lethborg, T., Purcell, A. W., Williamson, N. A., Chen, Z., Kjer-Nielsen, L., Mifsud, N. A., Tait, B. D., Holdsworth, R., Almeida, C. A., Nolan, D., Macdonald, W. A., Archbold, J. K., Kellerher, A. D., Marriott, D., Mallal, S., Bharadwaj, M., Rossjohn, J. and McCluskey, J.: Human leukocyte antigen class I-restricted activation of CD8+ T cells provides the immunogenetic basis of a systemic drug hypersensitivity. *Immunity*, **28**: 822–832 (2008).
  - 103) Wilke, R. A., Lin, D. W., Roden, D. M., Watkins, P. B., Flockhart, D., Zineh, I., Giacomini, K. M. and Krauss, R. M.: Identifying genetic risk factors for serious adverse drug reactions: current progress and challenges. *Nat. Rev. Drug Discov.*, **6**: 904–916 (2007).
  - 104) Molokhia, M. and McKeigue, P.: EUDRAGENE: European collaboration to establish a case-control DNA collection for studying the genetic basis of adverse drug reactions. *Pharmacogenomics*, **7**: 633–638 (2006).
  - 105) Liss, G. and Lewis, J. H.: Drug-induced liver injury: what was new in 2008? *Expert Opin. Drug Metab. Toxicol.*, **5**: 843–860 (2009).
  - 106) Locharnkul, C., Loplumlert, J., Limotai, C., Korkij, W., Desudchit, T., Tongkobpetch, S., Kangwanthirata, O., Hirankarn, N., Suphaphetiporn, K. and Shotelersuk, V.: Carbamazepine and phenytoin induced Stevens-Johnson syndrome is associated with HLA-B\*1502 allele in Thai population. *Epilepsia*, **49**: 2087–2091 (2008).