

the existence of thousands of T-DMRs that are hypo- and hypermethylated in ESCs (ES hypo hyper T-DMRs) [24]. Thus, an adequate DNA methylation profile during self-renewal and differentiation is critical for pluripotency. Recently, enrichment of short interspersed elements has been observed as a genomic property of regions harboring hypomethylated T-DMRs in ESCs [28]. Therefore, DNA methylation and heterochromatin signals will be useful for epigenetic assessment.

Chemicals that cause alterations in epigenetic systems are called epimutagens [29]. A DNA demethylating agent, 5-azacytidine, which is a cytidine analog, is a well-established epimutagen that causes reactivation of silenced genes [30]. It is known that Trichostatin A (TSA), an inhibitor of histone deacetylases, causes alteration in epigenetic regulation and nuclear structure of the cells [31]. Common chemicals in the environment also cause epigenetic alteration. We previously demonstrated that dimethyl sulfoxide (DMSO), which has been used as a cryopreservant for various cell lines, altered DNA methylation status in both gene areas and condensed repetitive sequences during the differentiation of ESCs into the embryoid body (EB) [32]. Carcinogenic nickel compounds have also been reported as an epigenetic modifier that induces histone methylation and changes in gene expression and chromatin structure [33]. Thus, environmental chemicals may have the potential to affect epigenetic states in the embryonic tissue or cells.

We here report the epigenetic effects of 25 environmental chemicals at low concentration, which were detected in maternal peripheral blood and cord blood, on heterochromatin signals and DNA methylation status of T-DMRs in mouse ESCs.

MATERIALS AND METHODS

Maternal peripheral blood and cord blood samples.

Maternal peripheral blood and cord blood samples at term were collected at Yamaguchi University during 2008–2010 for analysis of 3-phenoxybenzoic acid (3-PBA), 3,5,6-trichloro-2-pyridinol (TCP), dimethylphosphate (DMP), diethylphosphate (DEP), dimethylthiophosphate (DMTP), diethylthiophosphate (DETP), perfluorooctanoate (PFOA), and PFOS (Table 1). The blood samples were collected at Tokai University during 2005–2007 for other chemicals. This study was approved by the Ethics Committee of Yamaguchi Grand Medical Center, the Yamaguchi University School of Medicine, and the Human Subjects Committee of Tokai University Hospital. All samples were taken with written informed consent.

Sample preparation and chromatography conditions.

Determination of the serum concentrations of chemicals except for organophosphate and pyrethroid insecticides (Group A) and heavy metals (Group E) and volatile organic compounds (VOCs) (Group F) was as previously described [3, 4, 34, 35]. For octachlorodipropyl ether (S-421) (Group A), polybrominated diphenyl ethers (PBDEs) (Group D) and tin (Sn) (Group E), the exposure level to cells was determined by concentrations of nonserum sample, breast milk (S-421), and adult urine (penta BDE [36] and Sn). Concentration of dimethyldithiophosphate (DMDTP), diethyldithiophosphate (DEDTP), and deca BDE was based on the data from previous reports [37, 38].

The determination of 3-PBA and TCP (Group A) was by a modified previous method [39, 40] using liquid chromatography–tandem mass spectrometry (LC/MS/MS) (Alliance2695 and Quattro micro; Waters, Milford, MA) equipped with an electrospray ionization probe. After addition of beta-glucuronidase for derivatization, samples were purified with an Oasis HLB extraction column (Waters), and then fractionated by reverse-phase chromatography on a symmetric C18 column (50 mm × 2.1 mm i.d., 3.0 μm film thickness; Waters) with 35% acetonitrile and 0.1% acetic acid. Following sample load, MS/MS was performed in the negative-ion mode.

For S-421 (Group A), sample preparation from human milk was performed according to the

method described in Kakimoto et al. [41] with slight modifications, and the concentration of S-421 was determined by gas chromatography–mass spectrometry (GC/MS).

DMP, DEP, DMTP, and DETP (Group A) were determined using the method described in Ueyama et al. [42] with modifications. Samples were purified on a C18 cartridge column (Aisti Science, Wakayama, Japan), and then deproteinized with acetonitrile. After derivatization with pentafluorobenzylbromide, samples were purified using a three-layer column of Florisil, PSA, and anhydrous sodium sulfate. The concentrations of DMP, DEP, DMTP, and DETP were determined using gas chromatography with flame-photometric detector (GC/FPD) (Agilent Technologies, Tokyo, Japan) equipped with an LVI-S200 injection port (Aisti Science) on a DB-1701 or DB-5MS column (30 m × 0.25 mm i.d., 0.25 μm film thickness; Agilent Technologies).

The determination of heavy metals (Group E) were by a modified previous method [43]. Serum and urine samples were mixed with HNO₃ in a Nano-Band MV (GL Science, Tokyo, Japan) overnight. After microwave digestion, samples were mixed with internal standard (scandium (Sc), yttrium (Y), iridium (Ir)) and subjected to analysis by inductively coupled plasma–mass spectrometry (ICP/MS) (Agilent Technologies 7500i).

VOCs (Group F) in serum samples were determined using the previously described procedure [44]. Samples were diluted with saturated saline solution and spiked with internal standard (toluene-d₈) in a CV-22 vial (Chromacol, Trumbull, CT). VOCs were determined by AUTO MASS SYSTEM II (JEOL, Tokyo, Japan) coupled with Tekmar 7000 headspace sampler (Tekmar Co., Cincinnati, OH). Separation was accomplished with a Vocol column (60 mm × 0.25 mm i.d., 1.5 μm film thickness; Supelco, Bellefonte, PA). Analysis was performed in a selected ion monitoring mode and quantification masses for each of the analytes.

ESC culture and exposure of chemicals.

ESCs (J1) derived from 129 strain mice were cultured on 0.1% gelatin-coated dishes with ES medium, containing 15% fetal bovine serum (FBS; Biowest, Nuaille, France), 2 mM of L-glutamine (Invitrogen, Tokyo, Japan), 100 mM of nonessential amino acid (Invitrogen), 100 mM of

2-mercaptomethanol (Invitrogen), 1 mM of Na-pyruvate (Invitrogen), 50 U/mL of penicillin, 50 µg/mL of streptomycin (Invitrogen), and 2000 U/mL of leukemia inhibitory factor (LIF, ESGRO; Chemicon, Temecula, CA) in Dulbecco's modified Eagle's medium (DMEM; Wako Pure Chemicals, Osaka, Japan). To induce formation of EBs, ESCs were transferred to bacterial Petri dishes after removal of feeder cells. EBs were cultured in DMEM containing 10% FBS (JRH, Lenexa, KS), 50 U/mL penicillin, and 50 µg/mL streptomycin.

ESCs or EBs from J1 ESCs were cultured with chemicals at the serum level concentrations or tenfold higher level (Table 1). Selenium (Se), Cd, Hg, and Pb were dissolved in 0.0025% nitric acid (HNO₃). Sn was diluted with 0.007% hydrochloric acid (HCl). The other chemicals including TSA (Sigma-Aldrich, Tokyo, Japan) were dissolved in 0.1% ethanol (EtOH).

Slide preparation for heterochromatin analysis.

In the heterochromatin and DNA methylation analyses, all reagents, unless otherwise stated, were purchased from Wako Pure Chemicals. ESCs exposed to chemicals or their solvents were fixed with fix solution, consisting of EtOH (A) and acetic acid (B) (A/B:3/1, v/v). After centrifuging and removal of the supernatant liquid, the cells were suspended with 20–50 µL of fix solution, and then the interphase nuclei were spread on the glass slide. Prepared glass slides were stored at –20 °C until DAPI staining or DNA fluorescence *in situ* hybridization (DNA-FISH) analysis. For heterochromatin analysis with DAPI staining, the nuclei were stained with 0.5 µg/mL of DAPI (Dojindo, Kumamoto, Japan) for 5 min at room temperature, followed by washing in phosphate-buffered saline (PBS). The samples were mounted with antifade medium, PermaFluor Aqueous Mounting Medium (Thermo Scientific, Inc., Wilmington, DE).

DNA-FISH.

DNA-FISH using the major satellite repeat sequence, according to the previous reports [45, 46]. Probes specific to major satellite sequences in pericentric regions were labeled with biotin using Biotin-Nick Translation Mix (Roche Diagnostics, Tokyo, Japan). The interphase nuclei on a glass

slide were immersed in 2x SSC (0.3 M NaCl, 30 mM sodium citrate, pH 7.0) for 30 min at 37 °C, and then dehydrated with 70%, 80%, and 95% EtOH each for 2 min at room temperature. After denaturation with 70% formamide in 2x SSC (pH 7.0) for 2 min at 70 °C, the samples were dehydrated again with cold 70%, 80%, and 95% EtOH each for 2 min at -20 °C. The probes were diluted to 10 ng/mL with hybridization solution (50% formamide, 10% dextran sulfate, 50 mM phosphate buffer in 2x SSC, pH 7.0), followed by denaturation for 5 min at 95 °C. Denatured probe was added to the nuclear samples and incubated for 16 hr at 37 °C. The samples were then immersed in wash buffer (100 mM Tris-Cl, 150 mM NaCl, 0.05% Tween-20, pH 7.0) for 5 min at room temperature. Following incubation in blocking solution (100 mM Tris-Cl, 150 mM NaCl, 0.5% blocking reagent (Roche Diagnostics), pH 7.5) for 30 min at 37 °C, the samples were incubated with 10 µg/mL of biotinylated antistreptavidin (Vector Laboratories, Burlingame, CA) for 60 min at 37 °C. After three sets of washing in wash buffer, the samples were incubated with 5 µg/mL of Texas red-conjugated streptavidin (Vector Laboratories) for 30 min at 37 °C, followed by three sets of washing. Finally, 10 µg/mL of biotinylated antistreptavidin was added to the samples again for 30 min at 37 °C, and then the nuclei were counterstained with DAPI at room temperature.

Immunofluorescence.

ESCs were fixed in 4% paraformaldehyde for 10 min. After permeabilization with 0.2% Triton X-100 for 5 min, samples were treated with blocking buffer (5% BSA, 0.1% Tween-20 in PBS) for 30 min. The samples were incubated with anti-mouse heterochromatin protein 1a (HP1a, Chemicon, Temecula, CA) dissolved in blocking buffer (1:500) for 45 min, followed by three sets of washing in PBS containing 0.05% Tween-20. After incubation with fluorescent secondary antibody (anti-mouse IgG-FITC conjugated, KPL, Gaithersburg, MD) dissolved in blocking buffer (1:20) for 60 min, the samples were washed again. Finally, the samples were mounted with PermaFluor Aqueous Mounting Medium, containing 0.2 µg/mL of DAPI on the glass slide. All reactions were performed at room temperature.

2-mercaptomethanol (Invitrogen), 1 mM of Na-pyruvate (Invitrogen), 50 U/mL of penicillin, 50 µg/mL of streptomycin (Invitrogen), and 2000 U/mL of leukemia inhibitory factor (LIF, ESGRO; Chemicon, Temecula, CA) in Dulbecco's modified Eagle's medium (DMEM; Wako Pure Chemicals, Osaka, Japan). To induce formation of EBs, ESCs were transferred to bacterial Petri dishes after removal of feeder cells. EBs were cultured in DMEM containing 10% FBS (JRH, Lenexa, KS), 50 U/mL penicillin, and 50 µg/mL streptomycin.

ESCs or EBs from J1 ESCs were cultured with chemicals at the serum level concentrations or tenfold higher level (Table 1). Selenium (Se), Cd, Hg, and Pb were dissolved in 0.0025% nitric acid (HNO₃). Sn was diluted with 0.007% hydrochloric acid (HCl). The other chemicals including TSA (Sigma-Aldrich, Tokyo, Japan) were dissolved in 0.1% ethanol (EtOH).

Slide preparation for heterochromatin analysis.

In the heterochromatin and DNA methylation analyses, all reagents, unless otherwise stated, were purchased from Wako Pure Chemicals. ESCs exposed to chemicals or their solvents were fixed with fix solution, consisting of EtOH (A) and acetic acid (B) (A/B:3/1, v/v). After centrifuging and removal of the supernatant liquid, the cells were suspended with 20–50 µL of fix solution, and then the interphase nuclei were spread on the glass slide. Prepared glass slides were stored at –20 °C until DAPI staining or DNA fluorescence *in situ* hybridization (DNA-FISH) analysis. For heterochromatin analysis with DAPI staining, the nuclei were stained with 0.5 µg/mL of DAPI (Dojindo, Kumamoto, Japan) for 5 min at room temperature, followed by washing in phosphate-buffered saline (PBS). The samples were mounted with antifade medium, PermaFluor Aqueous Mounting Medium (Thermo Scientific, Inc., Wilmington, DE).

DNA-FISH.

DNA-FISH using the major satellite repeat sequence, according to the previous reports [45, 46]. Probes specific to major satellite sequences in pericentric regions were labeled with biotin using Biotin-Nick Translation Mix (Roche Diagnostics, Tokyo, Japan). The interphase nuclei on a glass

Image acquisition and counting heterochromatin foci.

Image acquisition of DAPI staining and DNA-FISH was performed using a fluorescent microscope (BZ-8000; KEYENCE, Osaka, Japan), and 5–10 visual fields were analyzed for individual samples. The exposure time of fluorescence by DAPI staining and major satellite DNA-FISH was fixed to 1/1.5 sec and 1.0 sec, respectively. After deconvolution of the fluorescent image by BZ-8000 software, the number of DAPI and major satellite signals in the nuclei was determined using ImageJ software provided by the National Institute of Health (<http://rsb.info.nih.gov/ij/>). Briefly, RGB images were first converted to 8-bit grayscale by linearly scaling min–max to 0–255. Next, the thresholds of intensity in DAPI and DNA-FISH images were decided within 79–84 and 15–20, respectively. Finally, the number of signals per nucleus ($n = 30–80$) was determined by ImageJ. Statistical analysis between solvent- and chemical-exposed cells was performed using the Wilcoxon test.

Combined Bisulfite Restriction Analysis (COBRA) assay.

Genomic DNA extraction and bisulfite conversion were carried out as previously described [26]. DNA methylation analysis was performed based on the previous finding [24] focusing on 48 T-DMRs by COBRA assay [47].

Genomic DNA was extracted from ESCs in lysis buffer (100 mM Tris-Cl (pH 8.0), 5 mM EDTA, 0.2% SDS, 200 mM NaCl, and 200 μ g/mL Proteinase K) at 55 °C for 30 min. After treatment with PCI, consisting of phenol (A), chloroform (B), and isoamyl alcohol (C) (A/B/C:50/49/1, v/v) for removal of proteins, RNaseA (Roche Diagnostics) was added and genomic DNA was purified by EtOH precipitation. Genomic DNA was digested with a restriction enzyme, *Hind* III (TaKaRa, Kyoto, Japan), and purified by EtOH precipitation. After denaturation of digested genomic DNA with 0.3 M NaOH, sodium metabisulfite (pH 5.0) and hydroquinone were added to final concentrations of 2.0 M and 0.5 mM, respectively. The bisulfite reaction was performed on a PCR machine as follows: 20 cycles of 95 °C for 30 sec and 55 °C for 15 min, followed by 55 °C for 10 hr. Bisulfite-treated genomic DNA was purified with QIAquick Gel Extraction Kit (Qiagen GmbH, Hilden, Germany), and desulfonated with 0.3 M NaOH at 37 °C for 15 min, followed by EtOH precipitation. Purified

bisulfite-treated DNA was amplified with BioTaq (BIOLINE, London, UK) using specific primers for gene loci (Supplemental Table 1). Amplified PCR products were digested with *Hpy* CH4 IV (New England BioLabs, Inc., Beverly, MA) at 37 °C for 3 hr, and then analyzed by microchip electrophoresis using MCE-202 (MultiNA; Shimadzu, Kyoto, Japan). The DNA methylation level by COBRA assay was calculated using the formula: estimated methylation degree (%) = $100 \times I^C / (I^C + I^{CU})$, where I^C and I^{CU} represent the sum of the intensities of digested bands and undigested bands, respectively.

Statistical Analysis.

Statistical comparisons of the number of DAPI signals were performed using the Wilcoxon test, and those of DNA methylation and area of EBs were performed using the Student's *t* test.

RESULTS

Chemical screening by heterochromatin marks.

Outline of the chemical exposure experiments was illustrated in Fig. 1. We first examined the effect of the 25 chemicals detected in maternal serum of pregnant mothers and cord blood (Table 1, Groups A–G) by counting the number of DAPI signals colocalized with signals of HP1a and major satellite DNA (Fig 2A). Reduction of numbers was observed in ESCs treated with TSA (1.0 nM) for 48 hr, but not with TSA at a lower concentration (0.2 nM) or control solvent (Fig. 2B). The other solvents, EtOH (0.1%), HCl (0.007%), or HNO₃ (0.0025%) for other chemicals, also had no effect on the DAPI signals in the treated ESCs (Fig. 2C).

Based on the levels of the 25 chemicals in the specimens (Table 1, Groups A–G), we assessed the effects of the chemicals on ESCs for 48 hr culture with each chemical at low (serum level, ×1, Table 1) and high (10 times, ×10, Table 1) concentrations (Exp. 1 in Fig. 1). DEP at serum concentration (0.1 ppb) significantly decreased the number of DAPI signals (Figure 2D). Cotinine (Group B) and Hg (Group E) also caused a decrease in DAPI signals at the serum concentration. In contrast, S-421 (Group A) and Se (Group E) increased the number of signals. Chemicals such as TCP, DMP, DETP, and DMDTP in Group A, deca BDE in Group D and mono(2-ethylhexyl)phthalate (MEHP) in group G exhibited effects on the number of DAPI signals only at high concentrations ($p < 0.01$) (Fig. 2D). It is noteworthy that five chemicals (DEP, Hg, cotinine, Se, and S-421) could affect the epigenetic status of ESCs at the concentration in the fetal environment.

Effects of DEP, Hg, cotinine, Se, and S-421 on the size of heterochromatin signals.

We further evaluated the effects of the five chemicals (DEP, Hg, cotinine, Se, and S-421) by examining the size of signals with both DAPI staining and DNA-FISH with major satellite repeats, in which signals were merged closely together in all preparations (Fig. 2E).

The size of the heterochromatin signals by DAPI and DNA-FISH were decreased by treatments with DEP and Se, but not in the Hg, cotinine, and S-421 treatments (Fig. 2F), although DEP, Hg, and cotinine decreased the number of signals and Se and S-421 increased the number as observed in the

screening using DAPI staining. These data indicate that the effects of DEP are, at least, distinct from those of Hg and cotinine, and the effect of S-421 is distinguished from that of Se. These distinct reactions by chemicals suggested that the size of heterochromatin signals should provide another dimension of heterochromatin status in addition to the number of DAPI signals, and that DEP, Hg, cotinine, Se, and S-421 have distinct effects on the epigenetic status of ESCs.

Effects of DEP, Hg, cotinine, Se, and S-421 on DNA methylation status at T-DMRs in ESCs.

To explore the effects of five chemicals (DEP, Hg, cotinine, Se, and S-421) on the epigenetic status of the genome at serum level concentrations, we investigated the DNA methylation status of 48 T-DMRs of genic regions, in which the methylation degree (%) varied from 0 to 80 in controls by COBRA assay using *Hpy* CH4 IV (Figure 3A, Exp. 2 in Fig. 1, and Supplemental Table 2).

DNA methylation levels in the control and Hg-exposed cells were $62.2\% \pm 0.79\%$ and $72.3\% \pm 2.38\%$ at *Rnd2* T-DMR, respectively, indicating that Hg caused hypermethylation (Fig. 3A and 3B). DNA methylation status at the *Aebp2* and *Prickle2* T-DMRs was also changed, resulting in hypomethylation in Se-exposed cells. DNA methylation levels in the control and Se-exposed cells were $28.0\% \pm 1.04\%$ and $18.9\% \pm 0.64\%$ at the *Aebp2* T-DMR, and $21.5\% \pm 0.70\%$ and $16.0\% \pm 0.73\%$ at the *Prickle2* T-DMR, respectively (Fig. 3B). These results indicate that Hg and Se affect the epigenetic status of DNA methylation as well as higher-ordered levels (heterochromatin), and that simple DAPI staining of ESCs enables us to screen for chemicals that affect epigenetic status.

Studies on prolonged effects of DEP, Hg, cotinine, Se, and S-421.

Changes in DNA methylation, which has been regarded as a stable epigenetic mark, prompted us to assess the prolonged effects of chemicals. We examined heterochromatin in ESCs exposed to the chemicals DEP, Hg, cotinine, Se, and S-421 at the serum level concentrations for 48 hr followed by 48 hr culture after their removal (Exp. 3 in Fig. 1). In the pretreatment of ESCs with Hg, cotinine, Se, and S-421, the number of DAPI signals in ESCs was reversed to almost the same levels as their controls (Fig. 4A, upper panel), suggesting that altered heterochromatin was reversible in the cells exposed to

these four chemicals. In contrast, the reduced number of DAPI signals in DEP-exposed cells did not recover to the control level after removal of DEP (Fig. 4A, upper panel). In addition, the area occupied by major satellite signals also remained at affected levels even after removal of DEP (Fig. 4A, lower panel). Thus, DEP-induced heterochromatin change was irreversible, even after 48 hr.

Effects of DEP, Hg, cotinine, Se, and S-421 on differentiation potentials in formation of EBs.

To test the effects of the chemicals on the developmental potential, we investigated the effects of chemicals on EB formation (Exp. 4 in Fig. 1). Although exposure to cotinine and Se had no effect on EB formation judged by the sizes of EBs between chemical-exposed and control cells, in the presence of DEP, Hg, and S-421 at low concentration, the size of EBs was significantly smaller than that of the controls (Fig. 4B).

DISCUSSION

The assessment of environmental chemicals has usually been performed with high doses of chemicals resulting in growth arrest in the previous studies. However, many chemicals have been detected in human samples with much lower concentrations at which severe growth arrest and cell death are not observed. In the present study, using mouse ESCs as an *in vitro* model for early embryos, we demonstrated that at least five chemicals (DEP, cotinine, Hg, Se, and S-421) have epimutagenic effects at the serum concentration of maternal peripheral blood and cord blood. The majority of the chemicals (20/25) examined had no effect on heterochromatin formation at the serum level, whereas the 10 times or more higher pharmacological dose may have some effects. In addition, exposure to Hg and Se caused alteration of DNA methylation status at unique gene loci harboring T-DMRs, DEP shows a long-term effect, and DEP, Hg, and S421 affect the developmental potentials. Thus, certain chemicals that are detected in human samples affect the epigenetic levels. A summary of the results is shown in Fig. 4C.

Of the five epimutagens, DEP-exposed ESCs did not recover from the altered heterochromatin status even after removal of the chemical, whereas other chemicals (Hg, cotinine, Se, and S-421) exhibited reversal of the number of DAPI signals after removal of the chemicals. This implies that DEP can cause long-lasting heterochromatin changes even at concentrations that have been detected in the serum of cord blood. DEP is a metabolite of chlorpyrifos, which is a component of pesticides. Prenatal exposure to pesticides caused lasting developmental disorders such as neural dysfunction in children even after birth [48, 49]. The EB is an *in vitro* model for differentiating early embryos [50], and EB formation was impaired in the presence of serum levels of DEP, Hg, and S-421, which caused altered heterochromatin formation in ESCs. In particular, DEP-induced epigenetic changes should be long lasting even at serum level concentrations. Collectively, certain chemicals such as DEP have the potential to cause epigenetic dysfunction in developing early embryos.

The genes showing disturbed DNA methylation status upon Hg or Se exposure have important roles in fetal development. *Aebp2* is a component of the epigenetic regulator, Polycomb Repression Complex 2 [51]. *Prickle2* and *Rnd2* are related to neural differentiation [52, 53]. This implies that the

genes with various functions regulated by DNA methylation are affected in ESCs that are thought of as an *in vitro* model of early embryonic cells.

The cytotoxic effect of chemicals is more severe in early stages of development than in adulthood [54, 55]. Heavy metals such as Se and Cd have been reported to cause epigenetic alteration in human cancer cell lines or fibroblasts [56, 57], although the cells were treated with high doses, more than 10 times those of serum level. On the other hand, human prostate cancer cell line LNCsP, exposed to Se for four days, did not exhibit altered DNA methylation [58]. In the present study, Hg and Se at serum levels caused altered heterochromatin marks and DNA methylation changes in ESCs.

In conclusion, the method using mouse ESCs to assess the epimutagens was simple and sensitive. We have demonstrated that DEP, Hg, cotinine, Se, and S-421 altered epigenetic systems at serum level concentrations detected in human maternal peripheral blood and cord blood.

ACKNOWLEDGMENTS

This work was supported by a Health Science Research Grant from the Ministry of Health, Labor and Welfare, Japan (to TM). The authors claim no competing financial interests.

REFERENCES

1. **Al-Saleh I, Shinwari N, Mashhour A, Mohamed GE, Rabah A.** Heavy metals (lead, cadmium and mercury) in maternal, cord blood and placenta of healthy women. *Int J Hyg Environ Health* in press
2. **Inoue K, Okada F, Ito R, Kato S, Sasaki S, Nakajima S, Uno A, Saijo Y, Sata F, Yoshimura Y, Kishi R, Nakazawa H.** Perfluorooctane sulfonate (PFOS) and related perfluorinated compounds in human maternal and cord blood samples: assessment of PFOS exposure in a susceptible population during pregnancy. *Environ Health Perspect* 2004; 112: 1204–1207.
3. **Okano K, Hinohara M, Iwasaki Y, Ito R, Saito K, Izumi S, Makino T, Nakazawa H.** Determination of nicotine and cotinine in human serum for evaluation of tobacco smoke exposure by hydrophilic interaction chromatography/mass spectrometry. *Bunseki Kagaku* 2007; 56: 785–790 (In Japanese).
4. **Takatori S, Kitagawa Y, Kitagawa M, Nakazawa H, Hori S.** Determination of di(2-ethylhexyl)phthalate and mono(2-ethylhexyl)phthalate in human serum using liquid chromatography-tandem mass spectrometry. *J. Chromatogr. B Analyt. Technol. Biomed. Life Sci.* 2004; 804: 397–401.
5. **Whyatt RM, Camann DE, Kinney PL, Reyes A, Ramirez J, Dietrich J, Diaz D, Holmes D, Perera FP.** Residential pesticide use during pregnancy among a cohort of urban minority women. *Environ Health Perspect* 2002; 110: 507–514.
6. **Zhang H, Feng X, Larssen T, Qiu G, Vogt RD.** In inland China, rice, rather than fish, is the major pathway for methylmercury exposure. *Environ Health Perspect* 2010; 118: 1183–1188.
7. **Jaddoe VW, Troe EJ, Hofman A, Mackenbach JP, Moll HA, Steegers EA, Witteman JC.** Active and passive maternal smoking during pregnancy and the risks of low birthweight and preterm birth: the Generation R Study. *Paediatr Perinat Epidemiol* 2008; 22: 162–171.
8. **Washino N, Saijo Y, Sasaki S, Kato S, Ban S, Konishi K, Ito R, Nakata A, Iwasaki Y,**

- Saito K, Nakazawa H, Kishi R.** Correlations between prenatal exposure to perfluorinated chemicals and reduced fetal growth. *Environ Health Perspect* 2009; 117: 660–667.
9. **Whyatt RM, Rauh V, Barr DB, Camann DE, Andrews HF, Garfinkel R, Hoepner LA, Diaz D, Dietrich J, Reyes A, Tang D, Kinney PL, Perera FP.** Prenatal insecticide exposures and birth weight and length among an urban minority cohort. *Environ Health Perspect* 2004; 112: 1125–1132.
10. **Zhu M, Fitzgerald EF, Gelberg KH, Lin S, Druschel CM.** Maternal low-level lead exposure and fetal growth. *Environ Health Perspect* 2010; 118: 1471–1475
11. **Ananth CV, Smulian JC, Vintzileos AM.** Incidence of placental abruption in relation to cigarette smoking and hypertensive disorders during pregnancy: a meta-analysis of observational studies. *Obstet Gynecol* 1999; 93: 622–628.
12. **Kippler M, Hoque AM, Raquib R, Ohrvik H, Ekström EC, Vahter M.** Accumulation of cadmium in human placenta interacts with the transport of micronutrients to the fetus. *Toxicol Lett* 2010; 192: 162–168.
13. **Llanos MN, Ronco AM.** Fetal growth restriction is related to placental levels of cadmium, lead and arsenic but not with antioxidant activities. *Reprod Toxicol* 2009; 27: 88–92.
14. **Todaka E, Sakurai K, Fukata H, Miyagawa H, Uzuki M, Omori M, Osada H, Ikezuki Y, Tsutsumi O, Iguchi T, Mori C.** Fetal exposure to phytoestrogens--the difference in phytoestrogen status between mother and fetus. *Environ Res* 2005; 99: 195–203.
15. **Edwards TM, Myers JP.** Environmental exposures and gene regulation in disease etiology. *Environ Health Perspect* 2007; 115: 1264–1270.
16. **Wade PA, Archer TK.** Epigenetics: environmental instructions for the genome. *Environ Health Perspect* 2006; 114: A140–A141.
17. **Golob JL, Paige SL, Muskheli V, Pabon L, Murry CE.** Chromatin remodeling during mouse and human embryonic stem cell differentiation. *Dev Dyn* 2008; 237: 1389–1398.
18. **Lieb JD, Beck S, Bulyk ML, Farnham P, Hattori N, Henikoff S, Liu XS, Okumura K, Shiota K, Ushijima T, Gready JM.** Applying whole-genome studies of epigenetic

- regulation to study human disease. *Cytogenet Genome Res* 2006; 114: 1–15.
19. **Shiota K.** DNA methylation profiles of CpG islands for cellular differentiation and development in mammals. *Cytogenet Genome Res* 2004; 105: 325–334.
 20. **Ikegami K, Ohgane J, Tanaka S, Yagi S, Shiota K.** Interplay between DNA methylation, histone modification and chromatin remodeling in stem cells and during development. *Int J Dev Biol* 2009; 53: 203–214.
 21. **Lachner M, O’Sullivan RJ, Jenuwein T.** An epigenetic road map for histone lysine methylation. *J Cell Sci* 2003; 116: 2117–2124.
 22. **Ng HH, Bird A.** DNA methylation and chromatin modification. *Curr Opin Genet Dev* 1999; 9: 158–163.
 23. **Vissel B, Choo KH.** Mouse major (gamma) satellite DNA is highly conserved and organized into extremely long tandem arrays: implications for recombination between nonhomologous chromosomes. *Genomics* 1989; 5: 407–414.
 24. **Sato S, Yagi S, Arai Y, Hirabayashi K, Hattori N, Iwatani M, Okita K, Ohgane J, Tanaka S, Wakayama T, Yamanaka S, Shiota K.** Genome-wide DNA methylation profile of tissue-dependent and differentially methylated regions (T-DMRs) residing in mouse pluripotent stem cells. *Genes Cells* 2010; 15: 607–618.
 25. **Shiota K, Kogo Y, Ohgane J, Imamura T, Urano A, Nishino K, Tanaka S, Hattori N.** Epigenetic marks by DNA methylation specific to stem, germ and somatic cells in mice. *Genes Cells* 2002; 7: 961–969.
 26. **Yagi S, Hirabayashi K, Sato S, Li W, Takahashi Y, Hirakawa T, Wu G, Hattori N, Hattori N, Ohgane J, Tanaka S, Liu XS, Shiota K.** DNA methylation profile of tissue-dependent and differentially methylated regions (T-DMRs) in mouse promoter regions demonstrating tissue-specific gene expression. *Genome Res* 2008; 18: 1969–1978.
 27. **Sakamoto H, Suzuki M, Abe T, Hosoyama T, Himeno E, Tanaka S, Greally JM, Hattori N, Yagi S, Shiota K.** Cell type-specific methylation profiles occurring disproportionately in CpG-less regions that delineate developmental similarity. *Genes Cells*

- 2007; 12: 1123–1132.
28. **Muramoto H, Yagi S, Hirabayashi K, Sato S, Ohgane J, Tanaka S, Shiota K.** Enrichment of short interspersed transposable elements to embryonic stem cell-specific hypomethylated gene regions. *Genes Cells* 2010; 15: 855–865.
 29. **Holliday R.** The inheritance of epigenetic defects. *Science* 1987; 238: 163–170.
 30. **Holliday R, Ho T.** DNA methylation and epigenetic inheritance. *Methods* 2002; 27: 179–183.
 31. **Taddei A, Maison C, Roche D, Almouzni G.** Reversible disruption of pericentric heterochromatin and centromere function by inhibiting deacetylases. *Nat Cell Biol* 2001; 3: 114–120.
 32. **Iwatani M, Ikegami K, Kremenska Y, Hattori N, Tanaka S, Yagi S, Shiota K.** Dimethyl sulfoxide has an impact on epigenetic profile in mouse embryoid body. *Stem Cells* 2006; 24: 2549–2556.
 33. **Chen H, Ke Q, Kluz T, Yan Y, Costa M.** Nickel ions increase histone H3 lysine 9 dimethylation and induce transgene silencing. *Mol Cell Biol* 2006; 26: 3728–3737.
 34. **Iwasaki Y, Goto M, Mochizuki K, Terayama E, Ito R, Saito K, Sugino N, Makino T, Nakazawa H.** Development and validation of a hydrophilic interaction chromatography-tandem mass spectrometry for quantification of nicotine and its metabolites in human maternal and cord sera. *Biomed Chromatogr* in press.
 35. **Nakata H, Nakata A, Okada F, Ito R, Inoue K, Saito K, Nakazawa H.** Development of online solid-phase extraction-HPLC/MS/MS method for the determination of perfluorochemicals in human plasma. *Bunseki Kagaku* 2005; 54: 877–884 (In Japanese).
 36. **Akutsu K, Takatori S, Nozawa S, Yoshiike M, Nakazawa H, Hayakawa K, Makino T, Iwamoto T.** Polybrominated diphenyl ethers in human serum and sperm quality. *Bull Environ Contam Toxicol* 2008; 80: 345–350.
 37. **Berman T, Hochner-Celnikier D, Barr DB, Needham LL, Amitai Y, Wormser U, Richter E.** Pesticide exposure among pregnant women in Jerusalem, Israel: results of a pilot

- study. *Environ Int* 2011; 37: 198–203.
38. **Takasuga T, Senthilkumar K, Takemori H, Ohi E, Tsuji H, Nagayama J.** Impact of fermented brown rice with *Aspergillus oryzae* (FEBRA) intake and concentrations of polybrominated diphenyl ethers (PBDEs) in blood of humans from Japan. *Chemosphere* 2004; 57: 795–811.
 39. **Baker SE, Olsson AO, Barr OD.** Isotope dilution high-performance liquid chromatography–tandem mass spectrometry method for quantifying urinary metabolites of synthetic pyrethroid insecticides. *Arch Environ Contam Toxicol* 2004; 46: 281–288.
 40. **Olsson AO, Baker SE, Nguyen JV, Romanoff LC, Udunka SO, Walker RD, Flemmen KL, Barr DB.** A liquid chromatography–tandem mass spectrometry multiresidue method for quantification of specific metabolites of organophosphorus pesticides, synthetic pyrethroids, selected herbicides, and deet in human urine. *Anal Chem* 2004; 76: 2453–2461.
 41. **Kakimoto K, Akutsu K, Konishi Y, Tanaka Y.** Time trend of hexabromocyclododecane in the breast milk of Japanese women. *Chemosphere* 2007; 71: 1110–1114.
 42. **Ueyama J, Saito I, Kamijima M, Nakajima T, Gotoh M, Suzuki T, Shibata E, Kondo T, Takagi K, Miyamoto K, Takamatsu J, Hasegawa T, Takagi K.** Simultaneous determination of urinary dialkylphosphate metabolites of organophosphorus pesticides using gas chromatography–mass spectrometry. *J Chromatogr B Analyt Technol Biomed Life Sci* 2006; 832: 58–66.
 43. **Razagui IB, Ghribi I.** Maternal and neonatal scalp hair concentrations of zinc, copper, cadmium, and lead: relationship to some lifestyle factors. *Biol Trace Elem Res* 2005; 106: 1–28.
 44. **Kondo F, Ikai Y, Goto T, Ito Y, Oka H, Nakazawa H, Odajima Y, Kamijima M, Shibata E, Torii S, Miyazaki Y.** Serum levels of volatile organic compounds in patients with sick building syndrome. *Bull Environ Contam Toxicol* 2006; 77: 331–337.
 45. **Lehnertz B, Ueda Y, Derijck AA, Braunschweig U, Perez-Burgos L, Kubicek S, Chen T, Li E, Jenuwein T.** Peters, A.H., Suv39h-mediated histone H3 lysine 9 methylation

- directs DNA methylation to major satellite repeats at pericentric heterochromatin. *Curr Biol* 2003; 13: 1192–1200.
46. **Senda S, Blanche S, Costagliola D, Cibert C, Nigon F, Firtion G, Floch C, Parat S, Viegas-Péquignot E.** Altered heterochromatin organization after perinatal exposure to zidovudine. *Antivir Ther* 2007; 12: 179–187.
47. **Xiong Z, Laird PW.** COBRA: a sensitive and quantitative DNA methylation assay. *Nucleic Acids Res* 1997; 25: 2532–2534.
48. **Eskenazi B, Marks AR, Bradman A, Harley K, Barr DB, Johnson C, Morga N, Jewell NP.** Organophosphate pesticide exposure and neurodevelopment in young Mexican–American children. *Environ Health Perspect* 2007; 115: 792–798.
49. **Harari R, Julvez J, Murata K, Barr D, Bellinger DC, Debes F, Grandjean P.** Neurobehavioral deficits and increased blood pressure in school-age children prenatally exposed to pesticides. *Environ Health Perspect* 2010; 118: 890–896.
50. **O’Shea KS.** Embryonic stem cell models of development. *Anat Rec* 1999; 257: 32–41.
51. **Kim H, Kang K, Kim J.** AEBP2 as a potential targeting protein for Polycomb Repression Complex PRC2. *Nucleic Acids Res* 2009; 37: 2940–2950.
52. **Heng JI, Nguyen L, Castro DS, Zimmer C, Wildner H, Armant O, Skowronska-Krawczyk D, Bedogni F, Matter JM, Hevner R, Guillemot F.** Neurogenin 2 controls cortical neuron migration through regulation of Rnd2. *Nature* 2008; 455: 114–118.
53. **Okuda H, Miyata S, Mori Y, Tohyama M.** Mouse Prickle1 and Prickle2 are expressed in postmitotic neurons and promote neurite outgrowth. *FEBS Lett* 2007; 581: 4754–4760.
54. **Benke GM, Murphy SD.** The influence of age on the toxicity and metabolism of methyl parathion and parathion in male and female rats. *Toxicol Appl Pharmacol* 1975; 31: 254–269.
55. **Chakraborti TK, Farrar JD, Pope CN.** Comparative neurochemical and neurobehavioral effects of repeated chlorpyrifos exposures in young and adult rats. *Pharmacol Biochem Behav* 1993; 46: 219–224.

56. **Jiang G, Xu L, Song S, Zhu C, Wu Q, Zhang L, Wu L.** Effects of long-term low-dose cadmium exposure on genomic DNA methylation in human embryo lung fibroblast cells. *Toxicology* 2008; 244: 49–55.
57. **Xiang N, Zhao R, Song G, Zhong W.** Selenite reactivates silenced genes by modifying DNA methylation and histones in prostate cancer cells. *Carcinogenesis* 2008; 29: 2175–2181.
58. **Ramachandran K, Navarro L, Gordian E, Das PM, Singal R.** Methylation-mediated silencing of genes is not altered by selenium treatment of prostate cancer cells. *Anticancer Res* 2007; 27: 921–926.