

特に無し

2. 実用新案取得

特に無し

3. その他

特に無し

別紙 4

研究成果の刊行に関する一覧表

雑誌

発表者氏名	論文タイトル名	発表誌名	巻号	ページ	出版年
Lei, K., Cusing, B.S., Musatov, S., <u>Ogawa, S.</u> , and Kramer, K.M.	Estrogen receptor-alpha in the bed nucleus of the stria terminalis regulates social affiliation in male prairie voles (<i>Microtus ochrogaster</i>).	PLoS One	5	e8931	2010
Furuta M, Mitsushima D, Shinohara K, Kimura F, <u>Funabashi T.</u>	Food availability affects orexin a/hypocretin-1-induced inhibition of pulsatile luteinizing hormone secretion in female rats.	Neuroendocrinology	91	41-47	2010
<u>Funabashi T.</u> , Furuta M, Fukushima A, Kimura F.	Age- and sex-specific changes in naloxone-induced luteinizing hormone secretion and Fos expression in gonadotropin-releasing hormone neurons of gonadectomized rats.	Neurosci Lett.	471	157-161	2010
Hagiwara H, Kimura F, Mitsushima D, <u>Funabashi T.</u>	Formalin-induced nociceptive behavior and c-Fos expression in middle-aged female rats.	Physiol Behav.	100	101-104	2010
<u>Bito, H.</u>	The chemical biology of synapses and neuronal circuits.	Nature Chem.Biol.	6	560-563	2010
Win-Shwe TT, <u>Tsukahara S.</u> , Yamamoto S, Fukushima A, Kunugita N, Arashidani K, Fujimaki H.	Up-regulation of neurotrophin-related gene expression in mouse hippocampus following low-level toluene exposure.	Neurotoxicology	31	85-93	2010

発表者氏名	論文タイトル名	発表誌名	巻号	ページ	出版年
Koike-Kuroda Y, <u>Kakeyama M</u> , Fujimaki H, <u>Tsukahara S</u> .	Use of live imaging analysis for evaluation of cytotoxic chemicals that induce apoptotic cell death.	Toxicol In Vitro.	24	2012-2020	2010
<u>Tsukahara S</u> , Tsuda MC., Kurihara R, Kato Y, Kuroda Y, Nakata M, Xiao K, Nagata K, Toda K, <u>Ogawa S</u> .	Effects of aromatase or estrogen receptor gene deletion on masculinization of the principal nucleus of the bed nucleus of	Neuroendocrinology		Epub ahead of print	2011
Tsuda MC, Yamaguchi N, <u>Ogawa S</u> .	Early life stress disrupts peripubertal development of aggression in male mice.	Neuroreport	22	259-263	2011
Endo T, <u>Maekawa E</u> , Vöikar V, Haijima A, <u>Uemura Y</u> , Zhang Y, Miyazaki W, Suyama S, Shimazaki K, Wolfer DP, Yada T, Tohyama C, Lipp HP, <u>Kakeyama M</u> .	Automated test of behavioral flexibility in mice using a behavioral sequencing task in IntelliCage.	Behav Brain Res	221	172-181	2011

研究成果の刊行物・別刷

*現在の資源・エネルギー事情を踏まえて、本研究課題の集大成というべき一報(Endo *et al.* 2011、本冊子の末尾に掲載)を除いて、別刷は2ページを1項に縮小、モノクロ印刷とさせていただきました。

Estrogen Receptor- α in the Bed Nucleus of the Stria Terminalis Regulates Social Affiliation in Male Prairie Voles (*Microtus ochrogaster*)

Kelly Lei^{1*}, Bruce S. Cushing^{2*}, Sergei Musatov³, Sonoko Ogawa⁴, Kristin M. Kramer¹

1 Department of Biology, University of Memphis, Memphis, Tennessee, United States of America, **2** Department of Biology and Integrated Bioscience Program, The University of Akron, Akron, Ohio, United States of America, **3** Neurologi, Inc. Laboratory of Neurobiology and Behavior, The Rockefeller University, New York, New York, United States of America, **4** Kansei, Behavioral and Brain Sciences Graduate School of Comprehensive Human Sciences, University of Tsukuba, Tsukuba, Ibaraki, Japan

Abstract

Estrogen receptor alpha (Er α) typically masculinizes male behavior, while low levels of Er α in the medial amygdala (MeA) and the bed nucleus of the stria terminalis (BST) are associated with high levels of male prosocial behavior. In the males of the highly social prairie vole (*Microtus ochrogaster*), increasing Er α in the MeA inhibited the expression of spontaneous alloparental behavior and produced a preference for novel females. To test for the effects of increased Er α in the BST, a viral vector was used to enhance Er α expression in the BST of adult male prairie voles. Following treatment, adult males were tested for alloparental behavior with 1–3 day-old pups, and for heterosexual social preference and affiliation. Treatment did not affect alloparental behavior or 73% of Er α -BST males and 62.5% of control males were alloparental. Increasing Er α in the BST affected heterosexual affiliation, with Er α -BST males spending significantly less total time in side-by-side contact with females relative to time spent with control males. Er α -BST males did not show a preference for either the familiar or novel female. These findings differed significantly from those reported in Er α -MeA enhanced males, where Er α inhibited alloparental behavior and produced a preference for a novel female. The findings from this study suggest two things: first, that increased Er α in the BST decreases social affiliation and second, that altering Er α in different regions of the social neural circuit differentially impacts the expression of social behavior.

Citation: Lei K, Cushing BS, Musatov S, Ogawa S, Kramer KM (2010) Estrogen Receptor- α in the Bed Nucleus of the Stria Terminalis Regulates Social Affiliation in Male Prairie Voles (*Microtus ochrogaster*). PLoS ONE 5(1): e8951. doi:10.1371/journal.pone.008951

Editor: Wim E. Crusio, Centre National de la Recherche Scientifique, France

Received: November 18, 2009; **Accepted:** December 23, 2009; **Published:** January 27, 2010

Copyright: © 2010 Lei et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This research was supported by funds provided by the University of Memphis (KMK) and a grant from the National Institute of Mental Health (MH-041992) (BSC). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* Email: kcruc@uakron.edu

Current address: Department of Psychology, Program in Neuroscience, Florida State University, Tallahassee, Florida, United States of America

Introduction

Estrogen acting via estrogen receptor alpha (ER α) masculinizes male behavior [1–4], which is typically associated with low levels of prosocial (positive affiliative) behavior and high levels of aggression. In male rats, treatment with a selective ER α agonist increased male aggression and anxiety [5] and masculinized serotonergic (5-HT) projections in female rats [6]. Conversely, data suggest that low levels of ER α are associated with high levels of male prosocial behavior. Highly social males, such as prairie voles (*Microtus ochrogaster*), pine voles (*M. pinetorum*), and Djungarian hamsters (*Phodopus campbelli*), display low levels of ER α in two regions of the brain that play a critical role in the expression of social and sociosexual behavior, the bed nucleus of the stria terminalis (BST) and medial amygdala (MeA) [7–9]. In male prairie voles, increased ER α is associated with decreased prosocial behavior. Neonatal castration inhibits the expression of prosocial behavior in male prairie voles [10,11], and significantly increases ER α expression in the BST and MeA [12]. Enhancing ER α in the MeA of male prairie voles decreased spontaneous alloparental behavior and resulted in a preference for a novel female over a familiar female [13]. These findings are significant for a couple of reasons. First, spontaneous

alloparental behavior is extremely difficult to disrupt in male prairie voles [reviewed in 12]. Second, of the numerous studies testing partner preference in male prairie voles only one other study, in that case adrenalectomized males, has even reported a trend for a preference for the novel female [14].

Based on extensive overlap in function and interconnections between nuclei in the limbic system, the BST and MeA have been classified as part of a social neural network that also includes the medial preoptic area, lateral septum, ventromedial hypothalamus, and the anterior hypothalamus [15]. While social behavior requires the interplay of a number of regions of the brain, the BST and MeA may play a particularly critical role in regulating social interactions as these areas receive direct input from the accessory olfactory bulb, have bi-directional communication, and are among the first regions to show neuronal activation during social contact [16–19]. Not surprisingly, the BST and MeA have been implicated in regulating a variety of social and sociosexual behaviors, including social preferences, affiliation, aggression [15,20,21], and mating [22]. The MeA, in particular, is necessary for social recognition [21]. While many of these studies have been conducted in rats and mice, the MeA and BST are involved in regulating the same social behaviors in prairie voles [23–25], suggesting that these regions

have a similar function in both highly social and less social species. Many of the studies that have examined the role of the BST in regulating social behavior have either found responses in both regions, which is not unexpected given the intimate relationships and interconnection, or have examined the response in only one region. One of the difficulties of interpreting the effects on behavior through the manipulation of a single region is that it is often unclear whether the change is due to the direct effect of the region or a response of the neural circuit to the manipulation. Therefore, one of the goals of this study is to use the same manipulation in a different region within the same neural circuit to determine if the effects are the same of if they vary by region.

Given that the BST and MeA have efferent and afferent neural connections and are part of the social neural network [15], ER α expression is low in both areas in male prairie voles [8], and ER α expression in the MeA decreased social behavior [13], we sought to determine the role of ER α in the BST. It also has been suggested that studying the role of ER α in the BST in regulating male social behavior is an essential next step [26]. Therefore the objectives of this study were to test the prediction that increasing ER α in the BST would reduce the expression of male prosocial behavior and to determine if the enhancing ER α in the BST had the same or different effects on male prosocial behavior as it did in the MeA. To accomplish this, a viral vector was used to enhance ER α expression in the BST of adult male prairie voles. This is a powerful technique for teasing apart the roles of specific brain regions as viral vectors can be delivered to specific sites using stereotaxic injection and they have limited spread, compared to the chronic and wide spread effect in knock out models. We replicated the design of Cushing et al. [13]. Treated males, along with the appropriate controls, were then tested for the expression of spontaneous alloparental behavior and heterosexual social preference and affiliation.

Methods

Subjects

Animals used in this study were laboratory-reared prairie voles that originated from wild stock trapped near Urbana, Illinois. Animals were housed under a 14:10 light/dark cycle and provided Purina high fiber rabbit chow (cat # 5326) and water *ad libitum*. Litters were weaned at 21 days of age and housed in same-sex pairs until treatment in polycarbonate cages (28.3 cm \times 17.5 cm \times 12.5 cm) with wood shavings for bedding. At the time of testing all subjects were sexually naive adults, 60–90 days of age. Animals were maintained in accordance with USDA and NIH guidelines and all procedures were approved by the University of Memphis Institutional Animal Care and Use Committee prior to conducting any study.

Viral Vector

Adeno-associated viral (AAV) vectors were used to enhance the expression of ER α in the BST. The following is a brief description of the vector; for complete details see Mustafaev et al. [27]. The vectors express shRNA containing human ER α (AAV-ER α) or luciferase (AAV-luciferase) target sequences. Vectors also express enhanced green fluorescent protein (GFP) as a reporter that allows for visual detection of transduced neurons. These vectors have been previously demonstrated to permit effective transfection in prairie voles [13].

ER α Adenoviral Vector Transfection

To achieve site-specific over-expression of ER α in the BST, adult males (60–70 days of age) were stereotactically injected,

bilaterally, with AAV-ER α . A site-specific control was generated by transfecting the BST of males with a vector encoding firefly luciferase cDNA, and an ER α control was generated by transfecting the caudate putamen with ER α , as the caudate does not express ER α . At approximately 60 days of age, a stereotaxic apparatus was used for site-specific injections of the AAV vector into the brain of experimental males. Males were randomly assigned to one of three treatment groups: 1) AAV-ER α into the BST ($n = 17$), 2) AAV-ER α into the caudate putamen (ER α transfection control) ($n = 9$), or 3) AAV-luciferase into the BST (injection control) ($n = 17$). Prior to the procedure, animals were deeply anesthetized with a combination of Ketamine (67.7 mg/kg) and Xylazine (13.3 mg/kg). The viral vector (1 μ l) was infused over a 5-min period with a micropump injector, and the infusion needle was left in place for an additional 3 min. The injection coordinates were determined to be AP -0.17 mm, ML ± 1.75 mm, 4.6 ventral from Bregma and, for the caudate, AP 1.3 mm anterior, ML ± 2.0 mm lateral, and 4.0 mm ventral. These coordinates specifically target the medial division of the BST. Animals were given 2–3 wks to recover and allow for expression of the vector prior to behavioral testing. Based upon a pilot study with prairie voles and other published results this amount of time is sufficient for the expression of transcribed ER α and expression lasts at least 13 weeks [27,28]. Following completion of testing, animals were euthanized and their brains collected to verify the accuracy of the injection and successful transfection. Only animals with bilateral expression of transcribed ER α were included in the analysis.

Verification of Transfection

Upon completion of the social preference test, brains from experimental animals were fixed using immersion fixation, sectioned at 30 μ m on a freezing-sliding microtome, and then the free-floating sections were stained for ER α using standard AB immunocytochemistry (for complete details see [8]). The human-specific primary antibody RA9101-s (Neomarkers, Fremont CA, 1:1000 dilution) was used to label transcribed ER α and then visualized using DAB. RA9101-s does not label prairie vole ER α . Therefore, all ER α observed with this antibody were the product of transfection. Successful transfection was determined qualitatively by visually examining ER α expression using a Nikon E-800 microscope. It should be noted that there were no animals that displayed only a few transcribed-ER α immunoreactive cells; transcribed males either displayed no or a significant amount of transcribed-ER α cells. Fig 1 shows a typical level of transcribed ER α expression. Control BST transfection was determined by staining for GFP expression, as described in [13].

Alloparental Behavior Test

Subjects were removed from the home cage and placed into a testing apparatus for 30–45 min to acclimate; food and water were provided during the acclimation period. The testing apparatus consisted of two standard size mouse cages (29 cm \times 19 cm \times 13 cm) connected by an 8-cm long clear acrylic tube. After the acclimation period, during which time all subjects investigated both mouse cages, an unlabeled vole pup 1–3 days of age was introduced. The vole pup was placed in the cage without the subject so that latency to approach the pup could be assessed. Recording began as soon as the stimulus pup was introduced. The test continued for 10 min after the subject entered the cage with the pup. If an attack occurred, the test was ended immediately so that, if necessary, the pup could be treated and to prevent further injury. Subjects were given 30 min to make an approach before ending the test. All tests were recorded with a digital video camera and scored by the same






Figure 1. Shows photomicrograph of AAV-mediated ERα expression at 200x in the medial division of the BST. ERα was labeled using the human-specific antibody RM9101's (Neomarkers, Fremont CA), which does not bind to endogenous prairie vole ERα. Immunoreactivity was visualized with DAB. doi:10.1371/journal.pone.0089931.g001

experimentally-blind observer using the JWatcher program (UCLA). Behaviors scored and analyzed included latency to enter the pup cage, duration of pup-directed licking and huddling, and pup-directed attacks. Animals that displayed pup-directed aggression were classified as non-alloparental.

Heterosexual Social Preference and Affiliation

One week after the alloparental test, treated males were tested for heterosexual social preference. This was done using the same apparatus as that used in the standardized vole partner preference test [11,29]. Each subject was placed in a clean cage with a sexually naive female for a 1-hr period of cohabitation. This female was designated as the "familiar." Immediately after the cohabitation period, social preferences of the subject were assessed. The social preference arena consists of three polycarbonate standard size mouse cages in a modified Y-shape. The two cages housing stimulus animals are in parallel and a third cage (neutral) is attached separately to each stimulus cage by Plexiglas tubes. The familiar female is gently tethered in one of the stimulus cages while an age- and size-matched sexually naive female, classified as "novel," that is unrelated to both the familiar female and the subject is tethered in the other stimulus cage. At the start of the test, immediately after the cohabitation period, the experimental animal is placed in the neutral cage and is free to move among the three cages. The social preference test lasted 3 hrs and was recorded using a digital video camera and then scored using the JWatcher program at a 10:1 temporal reduction. Behaviors scored included: time in each cage, time investigating each stimulus animal, and time spent in side-by-side contact with each stimulus animal. It should be noted that without hormonal manipulations, 1 hr of cohabitation is insufficient for prairie voles to form a preference for the familiar partner [14,30] and so control males were expected to spend equal amounts of time with the familiar and novel females. All stimulus females were sexually naive adults, 60–90 days of age, and mating is not a concern as female prairie voles do not undergo a spontaneous estrous cycle, requiring 24 or more hours of contact with a novel male before becoming sexually receptive [31].

Statistical Analyses

For all data sets, the two control groups were compared using a *t*-test. In no case were there differences between the two control groups; these were combined into a single control group for all subsequent analyses. A Fisher's exact test was used to analyze whether or not there was a difference between the proportion of control and ERα-BST males that expressed alloparental behavior. A one-way ANOVA was used to compare alloparental behaviors, licking and huddling, by treatment. A one-way ANOVA was used to analyze between treatment effects on social preference, while a paired *t*-test was used to analyze within treatments effects. For all statistical tests, the criterion for significance was $P \leq 0.05$.

Results

Eleven males were successfully transferred with ERα in the BST. Figure 1 shows a representation of successful transfection.

Alloparental

There was no significant difference in number of males that express alloparental behavior; 8 of 11 ERα-BST males were alloparental compared to 10 of 16 control males (Fisher's Exact $P = 0.69$). Non-alloparental behavior was comprised of, for ERα-BST males: 2 pup attacks and 1 with no contact; for controls: 2 pup attacks and 4 that did not enter the pup cage. In animals that expressed alloparental behavior, there was no effect of treatment on time spent licking or huddling with the pup (Fig. 2).

Heterosexual Social Preference & Affiliation

There was a significant effect of treatment on affiliation (Fig. 3). ERα-BST males (48 ± 7.3 s.e. min) spent significantly less total time in side-by-side with both familiar and novel females than did control males (81.9 ± 5.6 min; ANOVA $F_{1,25} = 13.4$, $P < 0.005$; $t_0 = 3.43$, $P = 0.004$). There was no significant difference between or within treatment groups for time spent in side-by-side contact with the familiar versus the novel female. Although a preference for the familiar animal is not expected after only 1 hr of cohabitation, control males did display a trend toward a preference for the familiar female (Control paired- $t_0 = -2.01$; $P = 0.06$; ERα-BST paired- $t_0 = -1.64$; ns; Fig. 3).




Figure 2. Show the mean (± s.e.) time spent licking and huddling pups during the alloparental tests by treatment. There was no significant difference for either licking or huddling between control and ERα-BST males. doi:10.1371/journal.pone.0089931.g002




Figure 3. Shows the mean (± s.e.) time spent in the cage and in physical (side-by-side) contact with the familiar and novel stimulus female by treatment. While there was no significant difference either within or between treatments for time spent in the cage or in contact with the familiar versus the novel female ERα-BST treated males spent significantly less total time (combined familiar and novel) than control males (see text). doi:10.1371/journal.pone.0089931.g003

Discussion

The results from this study support the hypothesis that low levels of ERα in the BST play a role in the expression of male prosocial behavior. Increasing ERα in the BST significantly reduced heterosexual social affiliation, but did not impact the expression of alloparental behavior. These findings are significant on several levels. First, they suggest that mechanisms underlying different types of affiliation differ, as increasing ERα in the BST decreased physical contact with females, but did not alter the amount of time spent licking or huddling with pups. Second, the results differed significantly from the effects of enhancing ERα expression in the MeA, where enhanced ERα inhibited the expression of alloparental behavior and resulted in the formation of a preference of the novel female [13]. Although the BST and MeA are both part of the extended amygdala, these results indicate that they regulate different aspects of social behavior and that the MeA is more important in regulating the expression of alloparental behavior than is the BST. Finally, the differential effects of increasing ERα in the BST versus in the MeA suggest the possibility that the suite of prosocial behaviors seen in male prairie voles is, at least in part, a product of the interaction between these two critical regions of the social neural circuit.

Enhancing ERα in the BST did not affect the expression of alloparental behavior. Male prairie voles typically display high levels of spontaneous alloparental behavior, ranging from 70 to 100% of males displaying alloparental behavior [13,32]. While at the lower end of the range, 73% of ERα-BST males were alloparental. This is compared with 33% of males in which ERα was enhanced in the MeA [13]. The expression of male parental behavior has been associated with a number of regions in the brain, including the BST and the MeA. In prairie voles, male parental behavior was associated with increased fos expression in both the MeA and BST [16]. Also in male prairie voles, adult castration reduced the expression of male parental behavior and the expression of arginine vasopressin, which has been shown to play a significant role in male social behavior [33,34], in the BST and MeA; treatment with testosterone restored both [23]. While Wang and De Vries [23] hypothesized that this resulted from the effect of testosterone on vasopressin expression, there is evidence that estrogen may play a direct role, with estrogen receptors

increasing in the BST and MeA in male mice in response to interactions with pups [35]. While these studies did show changes in several regions of the brain associated with parental behavior, they did not differentiate the role of the individual regions. The findings from the current study, along with the previous examination of ERα manipulation in the MeA of male prairie voles, suggest that the MeA, and not the BST, is critical for the expression of paternal behavior. A significant role for the MeA in paternal behavior is supported by the finding that axon sparing lesions of the MeA disrupted male parental behavior [36]. It is, however, still possible that the BST plays a significant role in male parental behavior through the actions of other compounds such as arginine vasopressin [23,37].

Enhancing ERα in the BST significantly reduced affiliation with females. This is in contrast to the effect of enhanced ERα in the MeA which did not affect total time spent in contact with females, but instead produced a preference for the novel female. Increased ERα expression resulted in a shift of affiliation, with male ERα-MeA males spending almost no time in contact with the familiar female [13]. In male prairie voles, neonatal castration not only inhibited the ability of central vasopressin to stimulate the formation of partner preferences [11], but also produced results similar to the current study, markedly reducing total side-by-side contact. Neonatal castration also resulted in a significant increase in ERα in both the MeA and the BST in adult males [12]. It is possible the results observed in the castration studies are the product of changes in both the BST and MeA, with increased ERα disrupting the formation of a partner preference, while changes in the BST reduced affiliation. This suggests that modification of prosocial behavior could result from subtle changes within one region of the brain, while changes in multiple regions may produce markedly different changes.

In conclusion, the ability to manipulate the expression of receptors in a site-specific manner can be a powerful tool. Studies that have looked at responses within the social neural circuit have been unable to differentiate the roles of the BST and the MeA. Although, it has been shown that there is a strong correlation between low levels of ERα expression levels in both the BST and MeA and the display of high levels of male prosocial levels [7–9], our study indicates that the role of ERα is not redundant in these two brain areas. In terms of prosocial behavior a lack of ERα in

Food Availability Affects Orexin A/ Hypocretin-1-Induced Inhibition of Pulsatile Luteinizing Hormone Secretion in Female Rats

Miyako Furuta^{a,d}, Dai Mitsushima^a, Kazuyuki Shinohara^b,
Toshiya Funabashi^a, Fukuko Kimura^{a,c}

^aDepartment of Physiology, Yokohama City University Graduate School of Medicine, Yokohama,
^bDepartment of Neurobiology and Behavior, Nagasaki University Graduate School of Biomedical Sciences,
Nagasaki, ^cInternational University of Health and Welfare, Tokyo, and ^dDepartment of Physiology,
St. Marianna University School of Medicine, Kawasaki, Japan

Key Words

Orexin A/hypocretin-1 • Luteinizing hormone pulse •
Glucose • Lactic acid • Stress • Fasting

Abstract

Orexin A/hypocretin-1 inhibits pulsatile luteinizing hormone (LH) secretion in female rats. In this study, we investigated whether this inhibition was tied to the fasting state, as suggested by our previous study. We first examined whether orexin A inhibited pulsatile LH secretion when food was available ad libitum during blood sampling. Next, we investigated the effect of intravenous administration of glucose (400 mg/kg) or lactic acid (negative control; 400 mg/kg) on orexin A-induced inhibition of pulsatile LH secretion. We found that orexin A did not affect pulsatile LH secretion in the presence of food, although it increased feeding behavior. Injection of orexin A significantly inhibited pulsatile LH secretion when food was withheld during blood sampling ($p < 0.05$); this inhibitory effect was rapidly reversed by intravenous injection of glucose but not lactic acid. Because orexin A did not seem to affect pulsatile LH secretion when food was available ad libitum, we speculate that orexin A has an effect on LH secretion when orexin A-induced hunger is ac-

companied by stress, such as the absence of food. Furthermore, glucose as well as food may act as a satiety factor in gonadotropin-releasing hormone pulse generation.

Copyright © 2009 S. Karger AG, Basel

Nutrient availability plays an important role in the maintenance of reproductive functions [1]. For example, fasting and insulin-induced hypoglycemia inhibit pulsatile luteinizing hormone (LH) secretion [2, 3] by inhibiting the electrical activity of the gonadotropin-releasing hormone (GnRH) pulse generator in the hypothalamus [4, 5]. Thus, the site of action of fasting and insulin-induced hypoglycemia is most likely the hypothalamus. Cellular depletion of ATP due to cellular glycopenia could result in inhibition of the GnRH pulse generator activity. However, this may not be the case, since glucose availability, rather than a simple measure of metabolism, such as ATP level, is essential for maintaining the electrical activity of the GnRH pulse generator in insulin-induced hypoglycemia [6]. Therefore, hypoglycemia-sensing mechanisms in the brain play an important role in conveying information

Acknowledgments

We wish to thank N. Cushing for her comments and suggestions on this manuscript and the animal care staff at the University of Memphis.

Author Contributions

Conceived and designed the experiments: BC. Performed the experiments: KL. Analyzed the data: KL, KK. Contributed reagents/materials/analysis tools: SM, SO. Wrote the paper: KL, BC, KK.

References

- Wessinger SR, Samson K, Villalba C, Lubahn DB, Rissman EF, et al. (1997) Masculine sexual behavior is disrupted in male and female mice lacking a functional estrogen receptor α gene. *Horm Behav* 32: 176–183.
- Ogawa S, Washburn TF, Lubahn DB, Korach KS, Pfaff DW (1996) Modifications of testosterone-dependent behaviors by estrogen receptor- α gene in mice. *Endocrinology* 138: 1099–1106.
- Kissman EF, Wessinger SR, Fink TH, Fink TC (1999) Sex with knockout: modulate behavioral studies of estrogen receptor alpha. *Brain Res* 833: 80–90.
- Cooke BM, Breedlove SM, Jordan C (2003) Both estrogen receptors and androgen receptors contribute to testosterone-induced changes in the morphology of the medial amygdala and sexual arousal in male rats. *Horm Behav* 43: 335–346.
- Pulsatile LH, Bateman HL (2008) Neonatal exposure to endocrine active compounds or an ER agonist increases adult anxiety and aggression in gonadally intact female mice. *Endocrinology* 153: 1099–1106.
- Pulsatile LH, Alesci H, Meuwissen JA (2009) Neonatal agonism of ER alpha modulates serotonergic (5-HT) projections to the female rat ventromedial nucleus of the hypothalamus (VMN) but does not impair lordosis. *Behav Brain Res* 196: 317–322.
- Hnaczk OC, Laciato CA, DonCarlos LL, Carter CS, Morrel JI (1994) Estrogen receptor immunoreactivity in specific brain areas of the prairie vole (*Mitovis adogenus*) is altered by sexual receptivity and genetic sex. *J Neuroendocrinol* 6: 81–90.
- Pulsatile LH, Morrel JI, Murphy AZ, Epperson PD, Le WW, et al. (2004) Luteinizing hormone secretion in the adult prairie vole is regulated by the same sociosexual behavior in two populations of prairie voles. *Brain Res* 1016: 247–254.
- Cushing BS, Wynne-Edwards KE (2006) Estrogen receptor- α distribution in male rodents is associated with social organization. *J Comp Neurol* 494: 595–605.
- Loewen JS, Rood BD, De Vries GJ (2002) Parental responsiveness is feminized after neonatal castration in virgin male prairie voles, but is not masculinized by neonatal testosterone. *Endocrinology* 151: 40–47.
- Cushing BS, Okere U, Wynne-Edwards KE (2003) The administered agnate vasopressin and the expression of V_1 receptors in adult male prairie voles. *J Neuroendocrinol* 15: 1021–1026.
- Cushing BS, Kruger KM (2005) Mechanisms underlying pituitary effects of early social experience: the role of neurotrophils and steroids. *Neurosci Biobehav Rev* 29: 1089–1105.
- Cushing BS, Perry A, Munoz S, Ogawa S, Papadimitrakou F (2008) Estrogen receptors in the brain inhibit the expression of male prosocial behaviors in prairie voles. *J Neurosci* 28: 10396–10407.
- DeVries AG, DeVries MB, Tayanian SE, Carter CS (1996) The effects of stress on social preferences are sexually dimorphic in prairie voles. *Proc Natl Acad Sci U S A* 93: 11980–11984.
- Newman SW (1999) The medial extended amygdala in male reproductive behavior: a node in the mammalian social behavior network. *Ann NY Acad Sci* 877: 242–257.
- Kirkpatrick B, Kim JW, Insel TR (1994a) Lumbic sex expression in prairie voles (*Mitovis adogenus*). *Brain Res* 638: 112–118.
- Philo HG, Hsueh MM (1996) The effects of gonadal steroid hormones on the brain following sexual stimulation of male and female rodents. *Brain Res Bull* 44: 397–407.
- Curtis JT, Wang ZX (2003) Forebrain c-fos expression under conditions conducive to pair bonding in female prairie voles (*Mitovis adogenus*). *Physiol Behav* 80: 95–101.
- Cushing BS, Mookewala N, Le WW, Hoffman GE, Carter CS (2003) Colocalization of Fos immunoreactivity in the monogamous prairie vole. *Brain Res* 965: 205–211.
- Raaij-Fillio AA, Lendiero RG, Achaval M (2000) Functional activities of the amygdala: an overview. *J Psych Neurosci* 25: 14–23.
- Ferguson JN, Aldag JM, Insel TR, Young LJ (2001) Oxytocin in the medial amygdala is essential for social recognition in the mouse. *J Neurosci* 21: 8278–8285.
- Vale P, Fim PD, Hoffman NW, Sogay N (1991) Sexually dimorphic cell groups in the medial preoptic area that are affected by male sex behavior and by the medial amygdala. *Endocrinology* 129: 463–470.
- Wang ZX, DeVries GJ (1998) Testosterone effects on parental behavior and vasopressin immunoreactive projections in prairie voles (*Mitovis adogenus*). *Brain Res* 631: 156–160.
- De Vries GJ, Villalba C (1997) Brain sexual dimorphism and sex differences in parental and other social behaviors. *Ann N Y Acad Sci* 807: 273–286.
- Wang Z, Hulthén TJ, Insel TR (1997) Sexual and social experience is associated with different patterns of behavior and neural activation in male prairie voles. *Brain Res* 767: 321–332.
- Trezza V, Campolongo P (2009) Toward understanding the neurobiology of social attachment: role of estrogen receptors in the medial amygdala. *J Neurosci* 29: 1519–1526.
- Munoz S, Chen W, Pfaff DW, Kaplan MG, Owen S (2006) RNA-mediated silencing of estrogen receptor α in the ventromedial nucleus of the hypothalamus abolishes female sexual behaviors. *Proc Natl Acad Sci U S A* 103: 10456–10460.
- Schultz KN, von Esenwein SA, Hu M, Bennett AL, Kennedy RT, et al. (2009) Viral vector-mediated overexpression of estrogen receptor α in striatum enhances the estradiol-induced motor activity in female rats and estradiol-modulated GABA release. *J Neurosci* 29: 1897–1903.
- Liu Y, Curtis JT, Wang ZX (2001) Vasopressin in the lateral septum regulates pair bonding and formation in male prairie voles (*Mitovis adogenus*). *Behav Neurosci* 115: 919–926.
- Che MM, DeVries AG, Williams JR, Carter CS (1999) The effects of oxytocin and vasopressin on partner preferences in male and female prairie voles (*Mitovis adogenus*). *Behav Neurosci* 113: 1071–1079.
- Carter CS, DeVries AG, Getz LL (1995) Physiological substrates of mammalian monogamy—the prairie vole model. *Neurosci Biobehav Rev* 19: 309–314.
- Roberts RL, Williams JR, Wang AK, Carter CS (1998) Cooperative breeding and monogamy in prairie voles: influence of the sex and geographical variation. *Anim Behav* 55: 1331–1340.
- Revenne EB, Curtis JT (2004) Vasopressin, oxytocin and social behaviour. *Curr Opin Neurobiol* 14: 772–783.
- Yan Y, Wang ZX (2004) The neurobiology of pair bonding. *Nat Neurosci* 7: 1048–1054.
- Elert G, Jurgens A, Kiehl M (1993) Endogenous-receptor occurrence in the mouse brain: modulation by paternal experience. *Neuroreport* 11: 1247–1250.
- Kirkpatrick B, Carter CS, Newman SW, Insel TR (1994b) Aton-sparing lesions of the medial nucleus of the amygdala decrease affiliative behaviors in the prairie vole (*Mitovis adogenus*): behavioral and anatomical specificity. *Behav Neurosci* 108: 501–513.
- Frazier CRM, Trainor BC, Cravens CJ, Whitney TK, Marder CA (2006) Paternal behavior influences development of aggression and vasopressin expression in male California mouse offspring. *Horm Behav* 50: 609–620.
- Cushing BS, Kim JW, Insel TR, Wang ZX (2004) Species differences in paternal behavior and aggression in prairie voles (*Mitovis adogenus*) with vasopressin immunoreactivity and receptors. *Horm Behav* 46: 25–38.
- Bester-Meredith JK, Marder CA (2001) Vasopressin and aggression in the rodent *Peromyscus leucopus*. *Physiol Behav* 74: 51–64.
- Gadaghe KL, Liu Y, Jia XX, Wang ZX (2007) Anterior hypothalamic neural activation and neurochemical associations with aggression in pair-bonded male prairie voles. *J Comp Neurol* 502: 1109–1122.

the BST appears to facilitate heterosexual affiliation. Finally, future studies will examine the role of the BST in regulating male/female and heterosexual aggression. Reduced aggression is the other side of increasing prosocial behavior, and there are a number of studies that suggest that the BST may play an important role in regulating aggression [38–40].

ICMARTING

© 2009 S. Karger AG, Basel
Fax +41 61 306 12 34
E-Mail karger@karger.ch
www.karger.com/nen

Toshiya Funabashi
Department of Physiology, Yokohama City University Graduate School of Medicine
3-9 Fukuura, K. Yokohama
Yokohama 236-0004 (Japan)
Tel: +81 45 787 2578, Fax: +81 45 787 2578, E-Mail toshiya@med.yokohama-cu.ac.jp

Materials and Methods

Animals

Seven-week-old female Wistar rats (Charles River, Yokohama, Japan) were maintained under controlled temperature (24–26°C) and lighting conditions (light on from 05:00 to 19:00 h) with food (Oriental Yeast Co., Ltd., Tokyo, Japan) and water available ad libitum. The rats were OVX at 8 weeks of age. All animal housing conditions and surgical procedures were in line with the guidelines provided by the Institutional Animal Care and Use Committee of the Yokohama City University School of Medicine.

Two weeks after ovariectomy, a stainless steel guide cannula (outer diameter, 0.7 mm; length, 13 mm) was stereotaxically placed into the third ventricle of each rat (stereotaxic coordinates: A = 6.0, V = 2.0 and L = 0.0 [19]) under sodium pentobarbital anesthesia (31.5 mg/kg body weight). The animals were allowed to recover for more than 10 days.

Estrogen treatment was performed under ether anesthesia around noon on the day before the sampling experiment. A silicone tube (inner diameter, 1.5 mm; outer diameter, 2.5 mm; length, 25 mm) containing 17 β -estradiol (E₂; Sigma Chemical Co., St. Louis, Mo., USA) dissolved in sesame oil at a concentration of 20 μ g/ml was implanted subcutaneously. We previously showed that this procedure results in serum estradiol levels that are equivalent to those observed on the day of diestrus in intact female rats [15]. An intra-atrial cannula was implanted through the jugular vein at the same time that the silicone tube was implanted.

Experimental Protocols

We first examined the effect of orexin A on pulsatile LH when food was available ad libitum during blood sampling. Blood samples were collected in a sound-attenuated room. Approximately 120 μ l of blood was collected under free-moving conditions at 6-min intervals from 09:00 to 12:00 h. After each blood sample was taken, the blood was replaced by an equal volume of heparinized saline (2 IU/ml). One hour after the first blood sample was collected, 0.3 nmol of orexin A (Peptide Institute Inc., Osaka, Japan) dissolved in 3 μ l artificial cerebrospinal fluid (aCSF) was injected into the third ventricle of each rat. The same volume of aCSF was injected into control animals.

In the next experiments, food was removed from individual cages just before the start of blood collection; there was no longer any food in the room with these rats, ensuring that the rats could neither see nor smell food. Blood samples were collected as described above. Orexin A (0.3 nmol) was injected i.c.v. after 1 h had passed, and glucose or lactic acid (as a negative control) were injected i.v. at a dose of 400 mg/kg [6], dissolved in 0.5 ml saline, through the same intracardiac cannula 2 h after the experiment started. We used lactic acid as a negative control because its effect on pulsatile LH secretion is identical to that of saline [6]; however, it maintains an ATP level similar to that maintained by glucose [20].

LH Assay and Pulse Analysis

Serum concentrations of LH were measured by double antibody radioimmunoassay with materials supplied by the National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK). The reference standard was NIDDK rat LH-RP-3, but the LH levels were expressed in terms of NIH LH-SI.

about the metabolic state to the GnRH pulse generating system, and the underlying key factor may be more complex than ATP level alone. Fasting may cause stress in rats [7]; however, in monkeys, it has been suggested that the suppression of pulsatile LH secretion by fasting is not caused by the psychological stress of food deprivation [5].

The lateral hypothalamic area of the brain is known as a hunger center [8] that contains orexin/hypocretin neurons [9, 10]. Orexins/hypocretins were first identified as neuropeptides that regulated feeding behavior [9, 10]. Subsequently, these neuropeptides were found to play a critical role in regulating the sleep/wake cycle [11], and it was suggested that they also mediated sleep and metabolic imbalances [12]. Regarding neuroendocrine functions, orexin inhibits LH secretion in ovariectomized (OVX) rats [13, 14], and this inhibitory effect is markedly enhanced by a low dose of estrogen that does not itself induce an LH surge [15]. However, orexin also stimulates LH secretion as part of a positive feedback cycle in which estrogen induces an LH surge [13]. Thus, orexin can either inhibit or suppress LH secretion, depending on other physiological conditions. Orexin also acts in a stimulatory manner in the rostral preoptic area and in an inhibitory manner in the medial preoptic area or arcuate/median eminence region, suggesting that it acts in a site-specific manner [16]. Although the precise mechanisms underlying orexin's distinct roles in LH secretion are unknown, orexins may act directly on GnRH neurons [17].

Preliminary experiments suggested that orexin A did not inhibit pulsatile LH secretion when the blood samples were collected under conditions where the rats had access to food ad libitum. In our previous report of the inhibitory effect of orexin A on LH secretion, the rats were under conditions where food was not available during the sampling [15]. We hypothesized that the inhibitory effect of orexin A was due to the absence of food, since orexin-induced hunger might be accompanied by stress stemming from the absence of food. We reasoned that if orexins acted as physiological feeding signals, orexin neurons would be activated at night, since rats eat more at night than in the daytime. It would follow that LH would thus be secreted less frequently at night. However, the LH pulse frequency increased rather than decreased at night [18], even though the rats should have been hungrier then.

The aim of the present study was to systematically determine whether orexin induces inhibition of pulsatile LH secretion when food is available ad libitum. We also examined whether a single satiety factor, such as glucose, can abolish the inhibitory effect of orexins in female rats.




Fig. 1. Representative examples of the effects of i.c.v. injection of aCSF alone (left panels) or aCSF plus orexin A (right panels) on the serum concentrations of LH in rats when food was available ad libitum. Data are shown as mean \pm SEM in the bottom two panels. Arrowheads indicate statistically defined LH pulses. Arrows indicate injections (i.c.v.) of aCSF or aCSF plus orexin A. * $p < 0.05$ versus the LH value during the 1st hour.

Results

Effect of Orexin on Pulsatile LH Secretion in Rats in the Presence of Food

Blood samples were collected every 6 min for 3 h from female rats that had free access to food. One group of rats was injected with 0.3 nmol of orexin A after 1 h, while another group was injected with vehicle (aCSF) alone (Fig. 1). In the aCSF group, each rat exhibited pulsatile LH

LH pulse detection was based on a method reported previously [6]. The analysis of LH pulsatility included the determination of pulse frequency (number of LH pulses per hour) and pulse amplitude (the difference between the peak and the pre-peak nadir). Group data of serum LH concentrations over a 3-hour period were also obtained by averaging values at each time point (Fig. 1). Analysis of variance (ANOVA) followed by Fisher's PLSD post-hoc comparison was used to test for statistical significance, and significance was accepted at $p < 0.05$. The mean LH concentration was analyzed by a paired *t*-test (Fig. 1, Fig. 3).




Fig. 2. Effects of i.c.v. injection of aCSF alone (a; n = 6) or aCSF plus orexin A (b; n = 8) on LH pulse frequency and pulse amplitude with food available ad libitum. Data are expressed as mean \pm SEM for the 1st hour, 2nd hour, and 3rd hour. Orexin A was injected after the 1st hour. No statistically significant differences were observed (ANOVA, $p > 0.05$).

secretion, which is normal for OVX rats (fig. 1). We did not observe orexin-induced suppression of pulsatile LH secretion in these conditions, i.e., when food was available ad libitum (fig. 1). However, orexin injection induced feeding behavior (data not shown), indicating that the orexin A was injected correctly and had a biological effect. LH pulse frequency in the aCSF group was not significantly different before versus after injection of aCSF (ANOVA, $p > 0.5$, fig. 2). In the orexin group, the LH pulse frequency decreased slightly but not significantly (ANOVA, $p > 0.1$, fig. 2). Thus, orexin A did not affect pulsatile LH secretion when food was available ad libitum during blood sampling. Neither the aCSF nor the orexin group showed a significant change in the amplitude of the LH pulses before and after injection (ANOVA, $p > 0.5$, fig. 2). As shown in the lower panels in figure 1, injection of orexin A resulted in small but statistically significant

cant decreases in the mean LH concentration at two time points (paired t test, $p < 0.05$).

Effect of Lactic Acid or Glucose Injection on Orexin-Induced Suppression of Pulsatile LH Secretion in the Absence of Food

Next, blood samples were collected every 6 min for 3 h from rats without access to food (fig. 3). Rats were injected with 0.3 nmol of orexin A after 1 h. After 2 h, glucose or lactic acid (as a negative control) was injected i.v. In both groups, each rat exhibited a spontaneous pulsatile LH secretion during the 1st hour of blood sampling (fig. 3), as in figure 1. This indicated that pulsatile LH secretion was observed regardless of whether food was available or not; this makes sense because rats normally eat very little food during this time of day (09:00–12:00 h). However, in both the lactic acid and the glucose groups, pulsatile LH secretion was markedly inhibited by i.c.v. injection of orexin A during the 2nd hour of blood samplings because food was withheld (fig. 3). Thus, orexin A did not affect LH pulse frequency (fig. 1, 2, the 2nd hour) but did decrease it (fig. 3, 4, the 2nd hour). The difference between these experiments was food availability: it was either available ad libitum (fig. 1) or was withheld during blood sampling (fig. 3). During the 3rd hour of blood sampling, after lactic acid or glucose injection, some LH pulses were observed (fig. 3). Analysis showed that LH pulse frequency decreased in both groups after orexin injection but immediately recovered in the 3rd hour after glucose injection (compared with the 1st hour, $p > 0.5$), but not after lactic acid injection (fig. 4, compared with the 1st hour, $p < 0.001$). LH pulse amplitude in the groups was not significantly changed during blood sampling ($p > 0.5$, fig. 4).

Discussion

In the present study, we confirmed that orexin A inhibited pulsatile LH secretion in estrogen-primed OVX rats when food was not available during blood sampling. We further showed that i.c.v. injection of orexin A did not inhibit pulsatile LH secretion when food was available ad libitum. When food was not available, the inhibitory effect of orexin A disappeared rapidly after i.v. injection of glucose. These data lead us to speculate that when orexin A is accompanied by stress due to the absence of food, LH secretion is affected. In general, if food is not available when rats want to eat, the result can be death. The stress due to the combination of absence of food plus injection




Fig. 3. Representative examples of the effects of i.v. injection of lactic acid (left panels) or glucose (right panels) on the serum concentration of LH after i.c.v. injection of orexin A. Data are shown as mean \pm SEM in the bottom two panels. Arrowheads indicate statistically defined LH pulses. Solid arrows indicate orexin A injection (i.c.v.). Open arrows indicate lactic acid or glucose injection (i.v.). * $p < 0.05$ versus the LH value during the 1st hour.

of orexin A can be eliminated by glucose as well as by food.

In contrast to the present results showing no effect of orexin A on pulsatile LH secretion in the presence of food, treatment with either neuropeptide Y (NPY) [21] or 2-deoxyglucose [3] inhibits pulsatile LH secretion even if neurons are thought to mediate the inhibitory effect of

food is available ad libitum. The effect of 2-deoxyglucose is probably mediated by NPY [22]. Thus, both NPY and orexin A generate hunger and drive a feeding response, but only NPY [21] inhibits pulsatile LH secretion in the presence of food. This discrepancy is puzzling since NPY neurons are thought to mediate the inhibitory effect of




Fig. 4. Effects of i.v. injection of lactic acid (a; n = 15) or glucose (b; n = 17) on LH pulse frequency and pulse amplitude after i.c.v. orexin A injection. Data are expressed as mean \pm SEM for the 1st hour, 2nd hour, and 3rd hour. Orexin A was injected after the 1st hour, and lactic acid or glucose was injected after the 2nd hour. * p < 0.05 versus the value during the 1st hour.

orexin A on pulsatile LH secretion and the stimulatory role of orexin A in feeding [23].

Consequently, we speculate that orexin A inhibits LH secretion in the absence of food, such as in the fasting state, but does not affect LH secretion in the presence of food. This raises the question of when orexin A acts to inhibit LH secretion. At present, it is not known whether orexin neurons are activated under physiological conditions, such as during spontaneous feeding when food is available ad libitum. We have preliminary data which suggest that orexin neurons are activated by fasting but not during free-feeding conditions (data not shown). Furthermore, orexin neurons are activated by subnormal glucose levels when food is absent (as in 48-hour fasting), but the activation of orexin neurons was not simply due to increased appetite [24]. Thus, we hypothesize that orexin A is activated by stressful conditions, such as fasting. This stress can be relieved not only by food but also by glucose. In line with this hypothesis, it has been suggested that in addition to affecting feeding behavior and wakefulness [25], orexin neurons are part of the circuitry that mediates the hypothalamic response to acute stress [26], although fasting-induced inhibition of LH secretion cannot be attributed solely to stress [27].

Acknowledgments

We are grateful to the NIDDK for providing radioimmunoassay materials. This work was supported by the Umehara Prize, which was awarded by the Yokohama Foundation for the Advancement of Medical Science.

References

- 1 Wade GN, Schneider JE. Metabolic fuels and reproduction in female mammals. *Neurosci Biobehav Rev* 1992;16:235-272.
- 2 Cagampang FRA, Maeda KI, Tsukamura H, Okkura S, Ota K. Involvement of ovarian steroids and endogenous opioids in the fasting-induced suppression of pulsatile LH release in ovariectomized rats. *J Endocrinol* 1991;129:321-328.
- 3 Nagatani S, Tsukamura H, Murahashi K, Maeda KE. A rapid suppressive effect of estrogen in the paraventricular nucleus on pulsatile LH release in fasting-ovariectomized rats. *J Neuroendocrinol* 1996;8:267-273.
- 4 Chen MD, O'Byrne KT, Chiappini SE, Hochkiss J, Knobil E. Hypoglycemic stress and gonadotropin-releasing hormone pulse generator activity in the rhesus monkey: role of the ovary. *Neuroendocrinology* 1992;56:666-673.
- 5 Goubillon ML, Thalabard JC. Insulin-induced hypoglycemia decreases luteinizing hormone secretion in the castrated male rat: involvement of opiate peptides. *Neuroendocrinology* 1996;64:49-56.
- 6 He D, Funabashi T, Sano A, Uemura T, Minaguchi H, Kimura F. Effects of glucose and related substrates on the recovery of the electrical activity of gonadotropin-releasing hormone pulse generator which is decreased by insulin-induced hypoglycemia in the estrogen-primed ovariectomized rat. *Brain Res* 1999;820:71-76.
- 7 Maeda K, Nagatani S, Estacio MA, Tsukamura H. Novel estrogen feedback sites associated with stress-induced suppression of luteinizing hormone secretion in female rats. *Cell Mol Neurobiol* 1996;16:311-324.
- 8 Bernardis LL, Bellinger LL. The lateral hypothalamic area revisited: ingestive behavior. *Neurosci Biobehav Rev* 1996;20:189-287.
- 9 de Lecea L, Kilduff TS, Peyron C, Gao X, Foye PE, Danielson PE, Fukuhara C, Bartberger EL, Gautvik VT, Bartlett ES, 2nd, Frankel WN, van den Pol AN, Bloom FE, Gautvik KM, Sutcliffe JG. The hypocretins: hypothalamus-specific peptides with neuroexcitatory activity. *Proc Natl Acad Sci USA* 1998;95:322-327.
- 10 Sakurai T, Amemiya A, Ishii M, Matsuzaki I, Chemel RM, Tanaka H, Williams SC, Richardson JA, Kozlowski GP, Wilson S, Arch JR, Buckingham RE, Haynes AC, Carr SA, Annan RS, McNulty DE, Liu WS, Terrett JA, Elshourbagy NA, Bergsma DJ, Yanagisawa M. Orexins and orexin receptors: a family of hypothalamic neuropeptides and G protein-coupled receptors that regulate feeding behavior. *Cell* 1998;92:1 page following 696.

- 11 Ohno K, Sakurai T. Orexin neuronal circuitry: role in the regulation of sleep and wakefulness. *Front Neuroendocrinol* 2008;29:70-87.
- 12 Adamantidis A, de Lecea L. The hypocretins as sensors for metabolism and arousal. *J Physiol* 2009;587:33-40.
- 13 Pu S, Jain MR, Kalra PS, Kalra SP. Orexins, a novel family of hypothalamic neuropeptides, modulate pituitary luteinizing hormone secretion in an ovarian steroid-dependent manner. *Regul Pept* 1998;78:133-136.
- 14 Tamura T, Irahara M, Tezuka M, Kiyokawa M, Aono T. Orexins, orexigenic hypothalamic neuropeptides, suppress the pulsatile secretion of luteinizing hormone in ovariectomized female rats. *Biochem Biophys Res Commun* 1999;264:759-762.
- 15 Furuta M, Funabashi T, Kimura F. Suppressive action of orexin A on pulsatile luteinizing hormone secretion is potentiated by a low dose of estrogen in ovariectomized rats. *Neuroendocrinology* 2002;75:151-157.
- 16 Small CJ, Goubillon ML, Murray JF, Siddiqui A, Grimshaw SE, Young H, Swanson V, Kalamatianos T, Kennedy AR, Coen CW, Bloom SR, Wilson CA. Central orexin A has site-specific effects on luteinizing hormone release in female rats. *Endocrinology* 2003;144:3225-3236.
- 17 Campbell RE, Grove KL, Smith MS. Gonadotropin-releasing hormone neurons coexpress orexin 1 receptor immunoreactivity and receive direct contacts by orexin fibers. *Endocrinology* 2003;144:1542-1548.
- 18 Hiruma H, Nishihara M, Kimura F. Hypothalamic electrical activity that relates to the pulsatile release of luteinizing hormone exhibits diurnal variation in ovariectomized rats. *Brain Res* 1992;582:119-122.
- 19 Albe-Fessard D, Sturmsky E, Liboutan S, Alblas stereotaxique du diencéphale du rat blanc. Paris: Editions du Centre National de la Recherche Scientifique; 1966.
- 20 Saitoh M, Okada Y, Nabetani M. Effect of mannose, fructose and lactate on the preservation of synaptic potentials in hippocampal slices. *Neurosci Lett* 1994;171:125-128.
- 21 Kalra SP, Kalra PS. Nutritional infertility: the role of the interconnected hypothalamic neuropeptide Y-galanin-opioid network. *Front Neuroendocrinol* 1996;17:371-401.
- 22 He B, White BD, Edwards GL, Martin RJ. Neuropeptide Y antibody attenuates 2-deoxy-D-glucose-induced feeding in rats. *Brain Res* 1998;781:348-350.
- 23 Yamamata A, Kunita K, Nambu T, Tajino N, Sakai A, Matsuzaki I, Miwa Y, Goto K, Sakurai T. Orexin-induced food intake involves neuropeptide Y pathway. *Brain Res* 2000;859:404-409.
- 24 Cai XI, Widdowson PS, Harrold J, Wilson S, Buckingham RE, Arch JR, Tadayyon M, Clapham JC, Wilding J, Williams G. Hypothalamic orexin expression: modulation by blood glucose and feeding. *Diabetes* 1999;48:2132-2137.
- 25 Kote CM. Integration of feeding and spontaneous physical activity: role for orexin. *Physiol Behav* 2006;88:294-301.
- 26 Winsky-Sommerer R, Boutrel B, de Lecea L. Stress and arousal: the corticotropin-releasing factor/hypocretin circuitry. *Mol Neurobiol* 2005;33:285-294.
- 27 Cameron JL, Helmreich DL, Schreihöfer DA. Modulation of reproductive hormone secretion by nutritional intake: stress signals versus metabolic signals. *Hum Reprod* 1993;18(suppl 2):162-167.



Contents lists available at ScienceDirect

Neuroscience Letters

journal homepage: www.elsevier.com/locate/neulet

Age- and sex-specific changes in naloxone-induced luteinizing hormone secretion and Fos expression in gonadotropin-releasing hormone neurons of gonadectomized rats

Toahiya Funabashi^{a,*}, Miyako Furuta^{a,b}, Atsushi Fukushima^a, Fukuko Kimura^{a,c}^a Department of Physiology, Yokohama City University Graduate School of Medicine, 3-9 Fikura, Kanazawa-ku, Yokohama 226-0004, Japan^b Department of Physiology, St. Marianna University School of Medicine, 2-16-1 Sugao, Miyamae-ku, Kawasaki 216-8511, Japan^c International University of Health and Welfare, Amity-Neigazaka-Bldg 1-21-1, Minamioyama, Minato-ku, Tokyo 107-0062, Japan

ARTICLE INFO

Article history:

Received 14 November 2009

Received in revised form

20 December 2009

Accepted 15 January 2010

Keywords:

Opioid

Pulse

Luteinizing hormone

Gonadotropin-releasing hormone

Immediate early genes

ABSTRACT

In the present study, we examined sex-specific changes in luteinizing hormone (LH) secretion and Fos expression in gonadotropin-releasing hormone (GnRH) neurons in response to naloxone in young (3 months old) and old (24 months old), gonadectomized male and female rats. We revealed by immunocytochemistry that, regardless of age and sex, naloxone significantly increased the number of GnRH neurons expressing Fos, which was associated with increased LH secretion. Additionally, although the magnitude of the increase in Fos-expressing GnRH neurons did not change in old males compared to young males, it was attenuated by almost half in old females compared to young females. LH levels decreased 60% in old males compared to young males and 15% in old females compared to young females. These results suggest LH secretion is impaired with age, but the ability of GnRH neurons to be stimulated by naloxone is preserved. However, the opioid-controlling mechanism is more fragile in females than males during aging.

© 2010 Elsevier Ireland Ltd. All rights reserved.

The gonadotropin-releasing hormone (GnRH) pulse generator includes a neuronal circuit and components for controlling pulsatile GnRH secretion. It is manifested by a characteristic increase in electrical neuronal activities in the hypothalamus accompanied by the initiation of each luteinizing hormone (LH) pulse. This event is called a multi-unit activity (MUA) volley [20,30,36]. Although the nature of the electrical activity and anatomically detailed evidence of the GnRH pulse generator are still uncertain.

The frequency of GnRH pulses is an important determinant of LH secretion from the anterior pituitary [21]. Estrogen can act in a negative feedback loop by decreasing the frequency of MUA volleys [19,20,25]. In rats, this effect of estrogen is at least partly mediated by opioid neurons [16]. The effects of the opioid receptor antagonist, naloxone, are also observable without estrogen [20]; therefore, changes in the frequency of MUA volleys in response to naloxone may reflect an important function of the GnRH pulse generator related to opioid neurons.

On the other hand, anatomical evidence has suggested that β-endorphin neurons make direct contact with GnRH neurons located




Fig. 1. Naloxone-induced changes in LH secretion during aging. Naloxone (filled circle with solid line for young and filled square with dashed line for old rats) was injected into males (left panel) and females (right panel). Control rats were injected with saline (open circle with solid line for young and open square with dashed line for old rats). Each point and vertical line indicate mean and SEM, respectively. Numbers in parentheses indicate numbers of rats, and * indicate $p < 0.05$ vs. saline control and old rats, respectively.

the influence of estrogen on pulse frequency in females, because estrogen decreases drastically with age. The aim of the present study was to determine whether the responsiveness of GnRH neurons to naloxone changed with age when negative feedback by gonadal steroid hormones was interrupted. GnRH responsiveness was measured by counting the number of GnRH neurons that expressed Fos in male and female Wistar rats ($n = 8$) were obtained (Charles River, Yokohama, Japan) at 7–8 weeks of age and were maintained under controlled lighting conditions (7:00–19:00 lights on) with food and water available ad libitum. Two weeks after castration and the day before the experiment, young (3 months old) and old (24 months old) rats were implanted with an intraventricular cannula. Naloxone (5 mg/kg) was intravenously injected after the first blood sampling in half of the rats, and thereafter blood samples were obtained through the cannula 15, 30, 45, and 60 min after injection. The remaining rats were controls and received saline injections. All animal housing and surgical procedures were in accordance with the guidelines of the Institutional Animal Care and Use Committee of the Yokohama City University School of Medicine.

Serum concentrations of LH were measured by double-antibody radioimmunoassay with materials supplied by the National Institute of Diabetes and Digestive and Kidney Disease (NIDDK). The reference standard was NIDDK rat LH-RP-3, but the amounts of LH are expressed in terms of NIH LH-S1. Changes in LH concentrations were analyzed by analyses of variance (ANOVA) for repeated measures, followed by the Fisher's protected LSD *post hoc* test and significance was attained at $p < 0.05$.

After blood sampling, the rats were killed by intravenous injection of an overdose of pentobarbital (70 mg/kg, bw) and prepared for the dual-immunocytochemical study, according to a previously reported method [8]. Briefly, the rats were perfused through the cardiac ventricle with ice-cold 4% paraformaldehyde in phosphate buffer (pH 7.5). Frozen coronal sections were cut 30 μ m thick and incubated overnight with rabbit polyclonal antibody to Fos (Oncogene Science, PC-05 Lot#2940101, 0.05 μ g/ml). The next day, sections were incubated with biotinylated anti-rabbit IgG (diluted 1:200) and then incubated with an avidin-biotin-peroxidase complex (Vectastain Elite ABC Kit, Vector Laboratories, Burlingame, USA). The sections were then incubated for 5 min in 0.05% 3,3'-diaminobenzidine with H_2O_2 . Following the detection of Fos immunoreactivity, the sections were rinsed well, incubated overnight with mouse monoclonal antibody to

GnRH (LRH13 [26]) (diluted 1:4000), and further incubated for 1 h with biotinylated anti-mouse IgG (diluted 1:200) and for 2 h with streptavidin-FITC (30 μ g/ml). The sections were rinsed and covered with aqueous medium and then examined under a light microscope equipped with appropriate filters for the visualization of the fluorescent signal. Omission of primary antibodies eliminated all staining. Secondary anti-mouse IgG and anti-rabbit IgG antibodies did not bind to the rabbit polyclonal antibody and mouse monoclonal antibody, respectively.

The counting was performed by an investigator unaware of the experimental conditions and expectations. The GnRH-immunoreactive (GnRH-ir) cells, in which a nucleus was visible as a clear halo surrounded by cytoplasmic fluorescent staining, were counted. GnRH-ir cells were defined as double-labeled for both Fos-ir and GnRH-ir when a blue-black nucleus signaling a Fos-ir cell was surrounded by cytoplasmic fluorescent staining. The forebrain was arbitrarily divided as shown in Fig. 2a: the rostral part of the forebrain was defined as the region including the diagonal band of Broca (DBB), the organum vasculosum of the lamina terminalis (OVLT), the anterior preoptic area (aPOA), the POA, the supraoptic nucleus-para-ventricular nucleus (SON-PVN), and the caudal part of the medial basal hypothalamic nucleus (MBH). The mean number of sections for each area was approximately 5.8, 3.9, 2.8, 8.6, 15.9, or 18.3 for the DBB, the OVLT, the aPOA, the POA, the SON-PVN, or the MBH, respectively. The number of immunoreactive cells was summed in each rat and the mean numbers were then calculated for each treatment group. Statistical comparisons were carried out by ANOVA followed by the Fisher's protected LSD *post hoc* test, and considered significant at $p < 0.05$.

Regardless of sex and treatment, LH secretion appeared to decrease with age; yet, naloxone seemed to induce LH secretion (Fig. 1). In male rats, repeated ANOVA showed a significant effect of treatment (saline vs. naloxone, $p < 0.001$) and age (young vs. old, $p < 0.0001$) without interaction ($p > 0.4$). Therefore, LH secretion significantly decreased with age and naloxone significantly induced LH secretion regardless of age (Fig. 1, left panel). In saline-injected rats, mean LH levels in old males were 62% of the LH levels in young males.

In female rats, repeated ANOVA showed a significant effect of treatment (saline vs. naloxone, $p < 0.0001$) and age (young vs. old, $p < 0.0001$) with interaction ($p < 0.02$). Since the interaction was significant, we further analyzed the data by one-way ANOVA ($p < 0.0001$). Naloxone significantly increased LH secretion in both young and old female rats (young $p < 0.0001$, old $p < 0.05$). Regard-

* Corresponding author. Tel.: +81 45 787 2578; fax: +81 45 787 2578.

E-mail address: toshiya@med.yokohama-cu.ac.jp (T. Funabashi).

¹ Present address: Department of Physiology, St. Marianna University School of Medicine, 2-16-1 Sugao, Miyamae-ku, Kawasaki 216-8511, Japan.





Fig. 2. Naloxone-induced Fos expression in GnRH-ir cells in young and old rats. (a) Schematic illustration of a coronal section showing the brain area where GnRH-ir cells expressing Fos-ir were quantified (modified from Abe-Fessard et al., 1996 [1]). aPOA, anterior preoptic area; CA, anterior commissure; DBB, diagonal band of the Broca; FX, fornix; IV, lateral ventricle; ME, median eminence; MBH, mediobasal hypothalamus; OT, optic tract; OX, optic chiasm; OVLT, organum vasculosum of the lamina terminalis; POA, preoptic area; SCN, suprachiasmatic nucleus; Sep, septum; SOM, supraoptic nucleus; SV, third ventricle. (b) The number of GnRH-ir cells (upper panels) and GnRH-ir cells that also express Fos-ir (lower panels) in each brain area in males (left panels) and females (right panels). Values are the mean \pm SEM and the numbers of rats are indicated in parentheses. * and ** indicate $p < 0.05$ vs. saline-injected rats at each age and saline-injected rats regardless of age, respectively.

females and does not change with age in male rats. Furthermore, LH levels in old rats dropped 40% in males and 85% in females, compared to young rats. The LH secretion data suggest that reproductive aging is due primarily to aging of the anterior pituitary function.

There are approximately 20% fewer GnRH-ir cells in old rats compared to young rats, irrespective of sex, suggesting age-related loss of GnRH neurons. This result is in good accord with our previous report [8], but contradictory to others [11,22]. Differences in the ages of animals used in the studies may be one reason for the variation; we used approximately 2-year-old rats, while other studies used rats approximately 10–12 months old. A more likely factor may be differences in the hormonal milieu of the animals; we used castrated rats, while other studies used intact animals.

We previously reported the percentage of GnRH-ir cells that express Fos in response to naloxone is 2.4% in male rats [24] and 6.8% in female rats [7]. In the present study, we found approximately 1.9% and 3.4% of GnRH-ir cells in young and old male rats, respectively, and 16.2% and 7.6% of GnRH-ir cells in young and old female rats, respectively, express Fos in response to naloxone. Small differences in the number of Fos-expressing GnRH-ir cells between the previous and present study are probably due to differences in the methods of naloxone administration (bulk intravenous injection in the present study and intravenous infusion in the previous study) and the time after naloxone treatment. We consider our overall results in the present study to be in accord with our previous results [7,24]. We conclude the population of GnRH neurons that express Fos is larger in young female rats than in young male rats, and there is no sex-specific difference in GnRH populations among old rats. Of note, both the percentage and distribution pattern of GnRH-ir cells that are also Fos-ir in the present study are in good accord with our previous studies; GnRH neurons located in the caudal region (i.e. the MBH) are more likely to express Fos in response to naloxone. This distribution pattern may suggest an important difference between GABAergic neurons and opioid neurons, which control GnRH secretion [7].

Opioid peptides are critically involved in the regulation of LH secretion via control of GnRH secretion [13]. Naloxone induces LH secretion by increasing GnRH secretion from the hypothalamus in rats [5]. Opioid peptides at least partly mediate the negative feedback of gonadal steroid hormones in both sexes of rats [13,23], as well as in human men [33] and women [34]. This suggests that the response to naloxone is due to the side of the hypothalamo-pituitary-gonadal axis related to the negative feedback of gonadal steroid hormones. Furthermore, in rats, opioid neurons play a distinct role in the control of GnRH secretion; naloxone stimulates the electrical activity of the GnRH pulse generator both in the presence and absence of gonadal hormones [12,20]. This phenomenon is also reported in goats [12], but not in monkeys [18]. Although the nature of the electrical activity of the GnRH pulse generator is not known, naloxone-induced Fos expression in GnRH neurons suggests activation of GnRH neurons. In support of this speculation, it has been reported that naloxone induces GnRH secretion in ovariectomized rats [14]. Thus, naloxone induces GnRH secretion [14], likely by increasing the GnRH pulse frequency, which is reflected as an increase in MUA volleys [20] and the resultant induction of Fos expression in GnRH neurons [7]. However, whether naloxone treatment induces Fos expression by directly activating GnRH neurons or by modulating other neurons is unclear from the present study, since naloxone was administered by an intravenous injection. Anatomical evidence [4] suggests direct action but we cannot deny the possibility that excitatory neurons activate GnRH neurons due to disinhibition by opioid neurons.

What is the physiological meaning of this sex-specific difference in Fos expression of GnRH neurons in young rats? Why is the GnRH pulse generator more active in females than males, as the present study in young rats demonstrates? We do not have

less of treatment, LH secretion in old females was significantly less than in young females ($p < 0.0001$). In saline-injected rats, mean LH levels in old females were only 15% of the LH levels in young females.

In Fig. 2b, the number of GnRH-ir cells (upper panels) is expressed as the mean number per section for each area, and the number of GnRH-ir cells also expressing Fos-ir (lower panels) are presented for each area (as schematized in Fig. 2a). The overall distribution of GnRH-ir cells in the forebrain was similar to that reported previously [38]. As we noted previously, a small number of GnRH-ir cells existed in the MBH. It was obvious that the number of GnRH-ir cells decreased with age, regardless of sex. In both the male and female rats, ANOVA revealed a significant effect of age (young vs. old males, $p < 0.0001$; young vs. old females, $p < 0.02$) but not of treatment (males: saline vs. naloxone, $p > 0.3$; females: saline vs. naloxone, $p > 0.7$) without interaction (males: $p > 0.3$; females: $p > 0.9$). These results suggested that aging caused a loss of GnRH neurons in male rats. Similar results have previously been reported in female rats lacking gonadal hormones [8].

There were Fos-ir cells throughout the forebrain in saline-injected rats, but few GnRH-ir cells were also Fos-ir, regardless of age or sex. However, the number of Fos-ir cells seemed to be increased in all subdivisions, extending from the rostral to the caudal part of the forebrain, in naloxone-injected rats (data not shown). For the total number of GnRH-ir cells that were also Fos-ir in male rats, two-way ANOVA revealed a significant effect of treatment (saline vs. naloxone, $p < 0.05$) but not of age (young vs. old, $p > 0.5$) without interaction ($p > 0.9$). These results suggested that naloxone-induced Fos expression in GnRH neurons regardless of age. There was no significant difference between young and old rats in the number of naloxone-induced Fos-ir cells in male rats. For the total number of GnRH-ir cells that were also Fos-ir in female rats, ANOVA revealed a significant effect of treatment (saline vs. naloxone, $p < 0.0005$) and age (young vs. old, $p < 0.04$) with significant interaction ($p < 0.04$). Further analysis by one-way ANOVA ($p < 0.0009$) revealed that the number of GnRH-ir cells expressing Fos-ir was significantly larger in naloxone-injected young females than in saline-injected young females ($p < 0.0003$) or naloxone-injected old females ($p < 0.007$). There were significantly more GnRH-ir cells that were also Fos-ir in naloxone-injected old females than in saline-injected old females ($p < 0.05$). These results suggested that naloxone treatment induced Fos expression in GnRH neurons, but the effect was larger in young females than in old females.

Finally, we analyzed the total number of double-labeled cells in each age to see a possible sex difference. In young rats, ANOVA revealed a significant effect of treatment (saline vs. naloxone, $p < 0.0005$) and sex (male vs. female, $p < 0.003$) with significant interaction ($p < 0.003$). Further analysis by one-way ANOVA ($p < 0.0005$) revealed the number of GnRH-ir cells that were also Fos-ir was significantly larger in naloxone-injected females than in saline-injected females or naloxone-injected males. This result suggested that naloxone treatment in young rats induced Fos expression in GnRH neurons, but the effect was larger in females than in males. In old rats, two-way ANOVA revealed a significant effect of treatment (saline vs. naloxone, $p < 0.002$) but not of sex (male vs. female, $p > 0.1$) without interaction ($p > 0.06$). This result suggested that naloxone-induced Fos expression in old rats regardless of sex.

The present study demonstrated, in females, the number of GnRH-ir cells that expressed Fos in response to naloxone was approximately two times larger in young rats than old rats; while, no difference was observed between young and old male rats. However, there was a sex-specific difference in the response of GnRH neurons to naloxone in young rats that disappeared in old rats. We concluded, in the absence of gonadal steroid hormones, the GnRH pulse generator does not differ between older males and older

answers to these questions yet, but one possible explanation is that GnRH neurons in females are active to maintain estrous cyclicity. The LH pulse frequency is influenced by the milieu of gonadal steroid hormones. Without testosterone in males and estrogen in females, the frequency of the GnRH pulse generator could be theoret-

ically be the same. In keeping with this hypothesis, the MUA volley intervals are identical in ovariectomized and orchidectomized rats [20]. Kato and Kimura personal communications). Therefore, basal pulse generator activity might be essentially the same in both sexes. However, the frequency is obviously higher in the follicular phase

compared to the luteal phase in female rats. This leads us to speculate that the GnRH pulse generator has the capacity to generate a higher frequency in females than in males, and this ability is lost during reproductive senescence. In female rats, the GnRH pulse generator is for the ovary, which shows phasic changes in function. After loss of ovary function due to reproductive senescence, it is quite reasonable there would be no sex-specific difference in the GnRH pulse generator in old rats, as found in the present study.

A simple interpretation of the present study may be that the pulse generator is not aged in both sexes, but rather GnRH neurons involved in surge generation are impaired with age. It has been suggested that opioid neurons are involved in the surge of LH secretion [2,13]. The mechanism that governs aging of the GnRH pulse generator in females is unknown at this time. GnRH neurons may be aged by definition [35]. An important point is that the effects of aging on the hypothalamus, in regard to GnRH neurons or other neural components of the GnRH pulse generator, are mild compared to age-related effects on the anterior pituitary.

Acknowledgement

This study was supported in part by a Grant-in-Aid for Scientific Research (C: 17590209 to TF) from the Ministry of Education, Culture, Sports, Science and Technology, Japan.

References

- Albe-Fessard, F., Stutinsky, S., Libouan, A., Atlas Stérolaxique du Diewéphale du Rat Blanc, Centre National de la Recherche Scientifique, Paris, 1966.
- Allen, E., Kahn, D., Caton, S.P., Kalia, D., Evidence that a decrease in opioid tone on proestrus changes the episodic pattern of luteinizing hormone (LH) secretion: implications in the precovulatory LH hypersecretion. *Endocrinology* 122 (1988) 1004–1013.
- R.J. Bicknell, Endogenous opioid peptides and hypothalamic neuroendocrine neurons. *J. Endocrinol.* 107 (1985) 437–446.
- W.-P. Chien, J.W. Witten, A.J. Silverman, Sexual dimorphism in the synaptic localization of gonadotropin-releasing hormone neurons. *Endocrinology* 126 (1990) 695–702.
- T.J. Cicero, P.F. Schmoekel, E.R. Meyer, R.T. Miller, Luteinizing hormone-releasing hormone mediates naloxone's effects on serum luteinizing hormone levels in normal and morphine-sensitized male rats. *Life Sci.* 37 (1985) 467–474.
- A. Coquelin, C. Desjardins, Luteinizing hormone and testosterone secretion in young and old male mice. *Am. J. Physiol.* 243 (1982) E257–E263.
- T. Funabashi, K. Jinma, F. Kimura, Fos expression by naloxone in LH neurons of the mediobasal hypothalamus and effects of pentobarbital sodium in the preovulatory rat. *J. Neuroendocrinol.* 9 (1997) 87–92.
- T. Funabashi, F. Kimura, The number of luteinizing hormone-releasing hormone immunoreactive neurons is significantly decreased in the forebrain of old-aged female rats. *Endocrinology* 136 (1995) 35–38.
- A.C. Gore, B.M. Windrose-Figueroa, E. Teague, Menopausal increases in pulsatile gonadotropin-releasing hormone release in a nonhuman primate (*Macaca mulatta*). *Endocrinology* 145 (2004) 4653–4659.
- R.A. Gorski, Gonadal hormones and the perinatal development of neuroendocrine function. *Front. Neuroendocrinol.* 2 (1971) 237–290.
- G.E. Hoffman, C.E. Finch, LH/RH neurons in the female (C57BL/6J) mouse brain during reproductive aging: no loss up to middle age. *Neurobiol. Aging* 7 (1986) 45–48.
- K. Ito, T. Tanaka, Y. Mori, Opioid peptide/gene control of gonadotropin-releasing hormone pulse generator activity in the ovariectomized goat. *Neuroendocrinology* 57 (1993) 694–699.
- S.P. Keenan, J.D. Veldhuis, The control of LH/RH secretion: a model for estrous cycle regulation. *Soc. Biol. Behav.* 23 (1985) 733–740.
- D.G. Karalunas, J.E. Levine, Naloxone stimulation of in vivo LH/RH release is not diminished following ovariectomy. *Neuroendocrinology* 47 (1988) 504–510.
- A.E. Krapav, W.J. Bremner, D.K. Clifton, R.A. Steiner, D.M. Dorsa, Diminished luteinizing hormone pulse frequency and amplitude with aging in the male rat. *Endocrinology* 112 (1983) 788–792.
- A. Kato, H. Hiruma, F. Kimura, Acute estradiol modulation of electrical activity of the LH/RH pulse generator in the ovariectomized rat: restoration by naloxone. *Neuroendocrinology* 59 (1994) 426–431.
- D.M. Keenan, J.D. Veldhuis, Disruption of the hypothalamic luteinizing hormone pulsing mechanism in aging men. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 281 (2001) R1917–R1924.
- J.S. Kessler, J.M. Kaufman, K.C. Wilson, G. Kuroda, E. Knobil, The effect of morphology on the electrophysiological activity of the hypothalamic neuroendocrine pulse generator in the rhesus monkey. *Neuroendocrinology* 43 (1986) 686–688.
- J.S. Kessler, K.C. Wilson, J.M. Kaufman, J. Haccibass, Y. Chen, H. Yamamoto, R.R. Pardo, E. Knobil, Unexpected responses of the hypothalamic gonadotropin-releasing hormone "pulse generator" to physiological estradiol inputs in the absence of the ovary. *Proc. Natl. Acad. Sci. U.S.A.* 84 (1987) 8745–8749.
- F. Kimura, M. Nishihara, H. Hiruma, T. Funabashi, Naloxone increases the frequency of the electrical activity of luteinizing hormone-releasing hormone pulse generator in long-term ovariectomized rats. *Neuroendocrinology* 53 (1991) 97–102.
- E. Knobil, The neuroendocrine control of the menstrual cycle. *Recent Prog. Horm. Res.* 36 (1980) 33–88.
- B.H. Koo, J.S. Kessler, The role of the μ -opioid receptor subunit expression in GnRH neurons changes during reproductive senescence in the female rat. *Endocrinology* 143 (2002) 3568–3574.
- M.A. Miller, W.J. Bremner, D.K. Clifton, D.M. Dorsa, R.A. Steiner, Opioid regulation of luteinizing hormone secretion in the male rat. *Biol. Reprod.* 35 (1986) 17–26.
- D. Mitsuhashi, T. Funabashi, F. Kimura, Fos expression in gonadotropin-releasing hormone neurons by naloxone or butorphanol in intact male rats. *Biol. Res.* 839 (1999) 209–212.
- Y. Mori, M. Nishihara, T. Tanaka, T. Shimizu, M. Yamaguchi, Y. Takouchi, K. Hoshino, Chronic recording of electrophysiological manifestation in the hypothalamic gonadotropin-releasing hormone pulse generator activity in the ovariectomized rat. *Neuroendocrinology* 43 (1986) 29–35.
- M.K. Park, K. Wakabayashi, Preparation of a monoclonal antibody to a common amino acid sequence of LH/RH and its application. *Endocrinol. Jpn.* 33 (1986) 257–272.
- W.G. Rossmanith, C. Reichert, W.A. Scherbaum, Neuroendocrinology of aging in humans: attenuated sensitivity to sex steroid feedback in elderly postmenopausal women. *Neuroendocrinology* 59 (1994) 355–362.
- A. Sano, F. Kimura, Electrical activity of the pulse generator of gonadotropin-releasing hormone in 26-month-old female rats. *Neuroendocrinology* 72 (2000) 199–207.
- K. Scarborough, P.M. Wise, Age-related changes in pulsatile luteinizing hormone release: precocious transition to continuous acyclicity and depend upon estrogen cycle block. *Endocrinology* 126 (1990) 1436–1443.
- T. Tanaka, Y. Mori, K. Hoshino, Hypothalamic GnRH pulse generator activity during the estradiol-induced LH surge in ovariectomized goats. *Neuroendocrinology* 56 (1992) 641–645.
- J.D. Veldhuis, A. Iranmanesh, E. Samojlik, R.J. Urban, Differential sex steroid negative feedback regulation of pulsatile follicle-stimulating hormone secretion in healthy older men: deconvolution analysis and steady-state sex-steroid hormone infusions in frequently sampled healthy older individuals. *J. Clin. Endocrinol. Metab.* 82 (1997) 1248–1254.
- J.D. Veldhuis, D.M. Keenan, P.Y. Liu, A. Iranmanesh, P.Y. Takahashi, A.X. Nehra, The aging male hypothalamic-pituitary-gonadal axis: pulsatility and feedback. *Endocrinology* 136 (1995) 2569–2576.
- J.D. Veldhuis, D.M. Keenan, E. Samojlik, N.H. Firth, Role of endogenous opiates in the expression of negative feedback actions of androgens on gonadotropin pulsatile properties of luteinizing hormone secretion in man. *J. Clin. Invest.* 74 (1984) 47–55.
- M. Vermeir, P.D. Silva, R.A. Lobo, Endogenous opioids modulate the inhibitory effects of androgen on the hypothalamic-pituitary axis of normal women. *J. Clin. Endocrinol. Metab.* 65 (1987) 1183–1186.
- Y. Wang, M. Garro, H.A. Dantzer, J.A. Taylor, D.D. Klim, M.C. Kuehl-Kovarik, Age affects spontaneous activity and depolarizing after potentials in isolated gonadotropin-releasing hormone neurons. *Endocrinology* 149 (2008) 1947–1954.
- R.C. Wilcox, J.S. Kessler, J.M. Kaufman, T. Uemura, T. Akema, E. Knobil, Central rhesus monkey. *Neuroendocrinology* 39 (1984) 256–260.
- J.J. Winters, R.J. Sherins, P. Treen, The gonadotropin-suppressive activity of androgen is increased in elderly men. *Metabolism* 33 (1984) 1052–1059.
- J.W. Wirtkin, C.M. Paden, A.J. Silverman, The luteinizing hormone-releasing hormone (LHRH) systems in the rat brain. *Neuroendocrinology* 35 (1982) 429–438.
- The impact of the estrous cycle on the nociceptive response in middle-aged female rats was assessed using the formalin test and c-Fos immunoreactivity as a marker of neural activation. Young (2-month-old) and middle-aged (11-month-old) rats were examined, dividing the middle-aged rats into two groups based on their estrous cycle: regular 4-day estrous cycle and irregular estrous cycle. The right hind paw was subcutaneously injected with 50 μ l of 2% formalin or saline as a control. Behavioral changes were observed for 1 h. Cycling rats were used during proestrus. Middle-aged female rats had a significantly higher score for nociceptive behavior compared to young rats, irrespective of estrous cyclicity, which suggests that aging, not the ability to maintain estrous cyclicity, causes hypersensitivity to the formalin injection. Immunohistochemical analysis found that the brain response to formalin injection was also more sensitive in middle-aged rats than young rats; a significant increase in the number of c-Fos immunoreactive cells was found in the ventral portion of the lateral septum of middle-aged rats injected with formalin compared to young and middle-aged rats injected with saline, irrespective of estrous cyclicity. Based on these results, we conclude that the sensitivity to painful stimuli in middle-aged female rats, which are in a neuroendocrine state similar to pre- and peri-menopausal women, is associated with age and not affected by reproductive ability. © 2010 Elsevier Inc. All rights reserved.
- Women often report pain-related symptoms during menopause [1,2], and pain is a significant health problem for middle-aged women [3]. However, the factors that affect the pain response in menopausal females are not clear. A lack of understanding regarding these factors and mechanisms is mostly due to limited studies on age-related changes in nociception and pain behavior, which have only been performed in males [4]; no studies have focused on females during the arrest of spontaneous ovulation. For example, an age-associated change has been demonstrated in the sensitivity to formalin-induced tonic pain, which peaks at mid-life [5], but the study only used male rats. Reproductive aging in female rodents is characterized by an arrest of spontaneous ovulation, with a transition from regular to irregular cycling in middle age [6,7]. Because the menopausal transition in women and female rats share significant characteristics, including patterns of luteinizing hormone secretion and responsiveness to estradiol, middle-aged female rats are an appropriate model for studying reproductive aging [7]. Importantly, changes in the blood estrogen levels of pre- and peri-menopausal women during aging [8] are similar to those of middle-aged rats [9–11]; the responsiveness to estrogen in the brain plays an essential role in the initiation of reproductive senescence before any changes in circulating estrogens [7]. In females, estrogens are not only essential for reproduction [12], but they are also an important factor in the modulation of pain responses. Interestingly, although the formalin test is a model of behavioral responses to tissue injury (phase 1) and inflammation (phase 2) [13,14], an estrogen effect is evident during interphase 1 [15–18]. Thus, estrogens, the responsiveness to estrogens, and/or aging per se may alter pain responses in aged female rats, and the formalin test is suitable for measuring these alterations. If 17 β -estradiol is an important factor for pain responses, a change in pain-related behavior would not be observed unless serum 17 β -estradiol is altered. If the responsiveness to 17 β -estradiol is important, pain-related behavior would be different between regular-cycling and irregular-cycling groups of female rats. If aging itself is important, female rats would exhibit altered pain responses regardless of reproductive state. Therefore, the present study examined changes in the response to formalin-induced nociceptive stimuli using a behavioral test and focusing on the effect of the estrous cycle in middle-aged rats with unchanged serum 17 β -estradiol levels [9–11]. In addition to behavioral examinations, we analyzed c-Fos expression

Contents lists available at ScienceDirect

Physiology & Behavior

journal homepage: www.elsevier.com/locate/physbeh



Formalin-induced nociceptive behavior and c-Fos expression in middle-aged female rats

Hiroko Hagiwara^a, Fukuko Kimura^{a,b}, Dai Mitsushima^a, Toshiya Funabashi^{a,*}

^a Department of Physiology, Yokohama City University Graduate School of Medicine, 3-9 Fukuura, Kanazawa-Ku, Yokohama 236-0004, Japan
^b International University of Health and Welfare, Amity-Nogizaka-Bldg. 1-2-1, Minamioyama, Miho-Ari-Ku, Tokyo 107-0062, Japan, Japan

ARTICLE INFO

ABSTRACT

The impact of the estrous cycle on the nociceptive response in middle-aged female rats was assessed using the formalin test and c-Fos immunoreactivity as a marker of neural activation. Young (2-month-old) and middle-aged (11-month-old) rats were examined, dividing the middle-aged rats into two groups based on their estrous cycle: regular 4-day estrous cycle and irregular estrous cycle. The right hind paw was subcutaneously injected with 50 μ l of 2% formalin or saline as a control. Behavioral changes were observed for 1 h. Cycling rats were used during proestrus. Middle-aged female rats had a significantly higher score for nociceptive behavior compared to young rats, irrespective of estrous cyclicity, which suggests that aging, not the ability to maintain estrous cyclicity, causes hypersensitivity to the formalin injection. Immunohistochemical analysis found that the brain response to formalin injection was also more sensitive in middle-aged rats than young rats; a significant increase in the number of c-Fos immunoreactive cells was found in the ventral portion of the lateral septum of middle-aged rats injected with formalin compared to young and middle-aged rats injected with saline, irrespective of estrous cyclicity. Based on these results, we conclude that the sensitivity to painful stimuli in middle-aged female rats, which are in a neuroendocrine state similar to pre- and peri-menopausal women, is associated with age and not affected by reproductive ability. © 2010 Elsevier Inc. All rights reserved.

1. Introduction

Women often report pain-related symptoms during menopause [1,2], and pain is a significant health problem for middle-aged women [3]. However, the factors that affect the pain response in menopausal females are not clear. A lack of understanding regarding these factors and mechanisms is mostly due to limited studies on age-related changes in nociception and pain behavior, which have only been performed in males [4]; no studies have focused on females during the arrest of spontaneous ovulation. For example, an age-associated change has been demonstrated in the sensitivity to formalin-induced tonic pain, which peaks at mid-life [5], but the study only used male rats.

Reproductive aging in female rodents is characterized by an arrest of spontaneous ovulation, with a transition from regular to irregular cycling in middle age [6,7]. Because the menopausal transition in women and female rats share significant characteristics, including patterns of luteinizing hormone secretion and responsiveness to estradiol, middle-aged female rats are an appropriate model for

* Corresponding author. Department of Physiology, Yokohama City University Graduate School of Medicine, 3-9 Fukuura, Kanazawa-Ku, Yokohama 236-0004, Japan. Tel./fax: +81 45 787 2578.
E-mail address: toshiya@med.yokohama-cu.ac.jp (T. Funabashi).

2.3. Immunohistochemistry

Frozen coronal sections (30 μ m) were cut using a Bright cryostat and washed with 0.1 M phosphate buffered saline (PBS). Samples were incubated overnight with rabbit polyclonal c-Fos antibody diluted 1:40,000 (PC38, Ab-5, Calbiochem) in PBS containing 1.5% normal goat serum and 0.1% Triton X-100. The next day, sections were incubated with biotinylated anti-rabbit IgG (1:200), followed by incubation with streptavidin–biotin–peroxidase complex (Vectastain Elite ABC Kit, Vectorstain Lab).

Bound peroxidase was visualized by incubating the sections for 8 min in 0.05% 3,3'-diaminobenzidine with H₂O₂. Samples were then mounted on glass slides, dehydrated in graded alcohol, cleared in xylene, and coverslipped with Permount.

The lateral septal nucleus (LS) was subdivided into the dorsal (LSD), intermediate (LSI), and ventral (LSV) regions [20] (Fig. 1), and the number of c-Fos immunoreactive (c-Fos-ir) cells in each region was determined by an investigator blinded to the experimental conditions. Two sections per rat were selected and matched across all animals in all experimental groups. Microscopic images (5 \times 10 magnification) were imported into a computer with a Penguin 600CL digital camera (Plexera Corporation, Los Gatos) and analyzed using Image-Pro plus version 5.1 (MediakCybernetics, Inc.). The cut-off value was defined so that the number of nuclei obtained by this method was consistent with the number obtained by visual inspection.




Fig. 1. Behavioral responses of young and middle-aged female rats during the formalin test. R, regular 4-day estrous cycle; IR, irregular estrous cycle. Each data point in the upper panel represents the amount of time the animals raised the injected paw (elevation duration) every 5 min during 1 h of observation. For statistical purposes, the data were separated into three phases in the lower panel: phase I (0–5 min after formalin injection), interphase (5–15 min after injection), and phase II (15–60 min after injection). The data are presented as mean \pm SEM. a, $P < 0.05$ vs. young rats.

as a marker of neural activity in order to determine the cerebral region associated with middle-age-related changes in the pain response.

2. Methods

2.1. Animals

Female 2-month-old (young) and 11-month-old (middle-aged) Wistar rats (Charles River, Yokohama, Japan) were maintained at a constant temperature of 24–26 °C under controlled lighting conditions (lights on 05:00–19:00) with food and water available ad libitum. Daily vaginal smears were obtained, and rats exhibiting three or more consecutive 4-day estrous cycles were defined as having a regular cycle. For young rats, only those exhibiting a regular cycle were used. Middle age was defined as 11 months of age because it is when rats transition from a regular estrous cycle to an irregular cycle [7]. The middle-aged rats were divided into two groups: rats exhibiting a regular 4-day estrous cycle (R group) and those that had ceased cycling (IR group). Regular-cycling rats were used on the day of proestrus. Irregular-cycling rats exhibiting persistent estrus or persistent diestrus were examined on these days. Twenty-four hours before formalin injection, a silicone cannula for intravenous anesthetic injection was implanted into the right atrium of all rats under ether anesthesia. This procedure did not interfere with the estrous cycle. The right hindpaws of some young (173 \pm 3.3 g, $n = 8$) and middle-aged rats (R group 376.3 \pm 3.4 g, $n = 8$; IR group 374.3 \pm 4.5 g, $n = 11$) were injected with 50 μ l of 2% formalin and behavioral changes observed for 1 h. The remaining young (172.1 \pm 4.1 g, $n = 7$) and regular middle-aged rats (375 \pm 7.6 g, $n = 6$) were injected with saline on the day of proestrus as a control. All rats were sacrificed by an overdose of intravenous pentobarbital (100 mg/kg) and subjected to immunohistochemical analysis.

Heparinized phosphate buffer (PB; pH 7.5) at approximately 4 °C was perfused through the cardiac ventricle, followed by paraformaldehyde (4%) in PB. After perfusion, the brain was removed from the cranium, fixed at 4 °C overnight in PB containing 4% paraformaldehyde, and incubated at 4 °C overnight in 25% sucrose in PB. The brains were then frozen with powdered dry ice and stored at –70 °C until processed for immunohistochemistry.

Blood samples were taken before the injection and the serum 17 β -estradiol concentrations determined using an EIA kit (Cayman Chemical Co., Ann Arbor, MI, USA). Serum samples were extracted once with diethyl ether and reconstituted with assay buffer. All animal housing and surgical procedures were in accordance with the guidelines specified by the institutional animal care and use committee of Yokohama City University School of Medicine. The experiments followed the ethical guidelines of the International Association for the Study of Pain [10].

2.2. Behavioral test

Immediately after the formalin injection, the rat was placed in a transparent Plexiglas box (30 \times 30 \times 30 cm) with a transparent floor positioned over a mirror angled at 45° to allow for observations of nociceptive behavior. The nociceptive behavior assessment [13,14] was performed on a single parameter by scoring the time of elevation for the injected paw. This simple method successfully detects sex differences in the formalin test [18] and was chosen because changes in single behavioral responses may be overlooked when using the weight-scores method [19]. We determined the length of time that the paw was elevated from the floor every 5 min for 1 h, and a mean response was calculated for each phase. For statistical purposes, the data were separated into three phases: phase I (0–5 min after formalin injection), interphase (5–15 min after injection), and phase II (15–60 min after injection).




Fig. 2. Representative photomicrographs of the lateral septum 1 h after saline or formalin injection. The c-Fos-immunoreactive cells are shown in a saline-injected young rat (Young-R-Saline), formalin-injected young rat (Young-R-Formalin), saline-injected middle-aged rat with a regular estrous cycle (Middle-R-Saline), formalin-injected middle-aged rat with a regular estrous cycle (Middle-R-Formalin), and a formalin-injected middle-aged rat with an irregular estrous cycle (Middle-IR-Formalin). Scale bar = 100 μ m.

All counting parameters, including the cut-off value and microscopic illumination, were unchanged throughout the counting procedure. The number of c-Fos-ir cells in a square region (600 mm \times 600 mm) in the LSI and LSV and a rectangular region (300 mm \times 600 mm) on the LSD was counted bilaterally. The average number of c-Fos-ir cells per area in each rat was used for statistical analysis. Data were analyzed by repeated or one-way analysis of variance (ANOVA), followed by Fisher's protected least significant difference post-hoc test. Differences were considered significant if $P < 0.05$.

3. Results

The serum concentrations of 17 β -estradiol in the young-R-formalin ($n = 8$), middle-R-formalin ($n = 6$), and middle-IR-formalin ($n = 6$) groups were 21.6 \pm 3.0 pg/ml, 23.3 \pm 4.2 pg/ml, and 26.4 \pm 4.1 pg/ml, respectively. No significant difference in the serum concentrations was found (ANOVA, $P > 0.6$).

3.1. Behavioral test

Formalin-injected rats exhibited two typical phases of nociceptive behavior (Fig. 1), which is in agreement with what was previously reported [13,14,18]. This pattern was similar in young and middle-aged rats, but we observed a different length of nociceptive behavior during interphase and phase II. Repeated measures ANOVA indicated a significant difference among the groups ($F_{2, 24} = 4.950$, $P < 0.05$) and phases ($F_{2, 48} = 69.297$, $P < 0.0001$) with a significant interaction ($F_{4, 48} = 2.895$, $P = 0.0317$). Because the interaction was significant, the groups were compared at each phase. We did not detect any significant differences among the groups during phase I ($F_{2, 24} = 0.691$, $P > 0.5$). In interphase ($F_{2, 24} = 4.273$, $P < 0.05$) and phase II ($F_{2, 24} = 5.880$, $P < 0.01$), ANOVA revealed significant differences among the groups. In both phases, the post-hoc comparison indicated that the duration of elevation in middle-aged rats was significantly longer than the duration in young rats, irrespective of the estrous cycle (interphase $P < 0.05$, phase II $P < 0.01$).

3.2. Number of c-Fos-ir cells

Based on visual inspection, the number of c-Fos-ir cells in the LS, but not other brain areas, including the paraventricular nucleus, the pre-optic area, and the bed nucleus stria terminalis, was associated with age. Formalin injection appeared to increase the expression of

c-Fos in the cells of the lateral septum (Fig. 2), particularly in the LSV. ANOVA indicated a significant difference in the number of c-Fos cells in the LSV among the groups ($F_{4, 35} = 6.326$, $P < 0.001$; Fig. 3). The post-hoc comparison revealed that the number of c-Fos-ir cells in the middle-aged rats was significantly greater than in the control groups of young and middle-aged rats, irrespective of estrous cycle ($P < 0.02$). We did not detect any significant differences in the number of c-Fos-ir cells between young saline- and formalin-injected rats




Fig. 3. Quantification of c-Fos immunoreactive cells in the lateral septum. Brain sections were from saline-injected young rats (Young-R-Saline), formalin-injected young rats (Young-R-Formalin), saline-injected middle-aged rats with a regular estrous cycle (Middle-R-Saline), formalin-injected middle-aged rats with a regular estrous cycle (Middle-R-Formalin), and formalin-injected middle-aged rats with an irregular estrous cycle (Middle-IR-Formalin). An illustration of the coronal region of the rat brain is shown in the upper right panel. The areas with bold and hatched lines indicate the regions in which c-Fos immunoreactive cells were counted and those taken for photomicrographs in Fig. 2, respectively. Ac, anterior commissure; cc, corpus callosum; LSD, lateral septum dorsal; LSI, lateral septum intermediate; LSV, lateral septum ventral; n, optic nerve. Data are presented as mean \pm SEM. a, $P < 0.05$ vs. young rats injected with saline; b, $P < 0.05$ vs. middle-aged rats injected with saline.

($P < 0.05$). In the LSI, changes in the number of c-Fos-ir cells were similar to the changes in the LSV, but ANOVA found no significant difference ($F_{4, 36} = 2.411, P > 0.05$). ANOVA also found no significant difference in the number of c-Fos-ir cells in the LSD among the groups ($F_{4, 36} = 0.758, P > 0.5$; Fig. 3).

4. Discussion

We used the formalin test and immunohistochemical analysis to demonstrate that middle-aged female rats are more sensitive to formalin-induced nociceptive stimuli, irrespective of estrous cyclicity. We speculate that the brain response of the middle-aged rats, whose neuroendocrine status is similar to that of pre- and peri-menopausal women, exhibit hypersensitivity to nociceptive stimuli due to age, and this response is unrelated to reproductive ability.

Studies have suggested that females are sensitive to formalin-induced pain due to the influence of ovarian steroid hormones [15]. For example, normal female rats exhibit more nociceptive responses to formalin than ovariectomized rats, suggesting modulatory roles of ovarian steroid hormones [16]. Ovarian steroid hormone replacement in ovariectomized rats restores the ovariectomy-induced decline in nociceptive response [21], though contradictory results have been reported [22]. Thus, age-related changes in blood 17 β -estradiol levels and/or responses to 17 β -estradiol might determine pain hypersensitivity in middle-aged rats. However, 17 β -estradiol levels of middle-aged rats, not only on the day of proestrus during regular-cycling, but also on another day during irregular cycling, are similar to the levels in young rats on the day of proestrus. These results are consistent with previous reports [9–11]. Therefore, we speculate that 17 β -estradiol levels are not an important factor for hyperalgesia in middle-aged, and the pain response is elevated in association with age not only in males [5], but also in females. If this situation is the case, we would like to note that the aging effects are also related to phase in the formalin test; middle-aged rats were hyperalgesic in interphase and phase 2 but not in phase 1. The absence of age effects in phase 1 of the formalin test suggests that the tissue injury induced by formalin injection in middle-aged rats is not different from the injury induced in young rats. Middle age may be too early to see changes in phase 1, but previous results in old male rats may eliminate this possibility [5]. Currently, the mechanism for the phase-specific effects of age is not known, but the results of the present study support the idea that age differences in the pain responses are not linear in models of tissue injury and inflammation [4].

Another finding from the present study is that the activation of the LSV after formalin injection is stronger in middle-aged female rats than young female rats. This observation is in accordance with previous findings in male rats [23]. However, in young female rats, we observed a tendency of increased c-Fos expression after formalin injection, but it was not significantly different from the expression in saline-injected rats. This finding is contradictory to previous reports, which found increased c-Fos expression in the LS of male rats [24]. This discrepancy may be due to the assessment time after injection; we examined expression 1 h after injection, whereas the other report analyzed expression 2 h after injection [24]. In addition, Aloisi et al. [25] did not find a significant difference 90 min after injection. Nevertheless, our results suggest that pain responses are elevated in middle age. This speculation is supported not only by the results from our behavioral examination, but also the morphological study.

How the LSV is associated with the pain response in the brain after formalin injection is not yet known. Because pain is an emotional experience, and the lateral septum plays an important role in regulating emotional responses [18,26–28], we speculate that middle-aged females are hypersensitive to a pain-induced emotional

process. Future studies should address the role of the LSV and the underlying mechanisms in the pain response in middle-aged females.

Acknowledgment

This work was supported by a Grant-in-Aid for Scientific Research (C: 17500208 to FK) from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

References

- Peat G, Thomas E, Handy J, Croft P. Social networks and pain interference with daily activities in middle and old age. *Pain* 2004;112:397–405.
- Barnabei VM, Cochran BB, Aragaki AK, Nygaard I, Williams RS, McGovern JG, et al. Menopausal symptoms and treatment-related effects of estrogen and progesterin in the Women's Health Initiative. *Obstet Gynecol* 2005;105:1063–73.
- Roussouni W, Cortiub S, Pabst A, midliff J. Midlife Women's Health. 2004;49:38–38.
- Geisler L, Mézack R. Age differences in nociception and pain behaviours in the rat. *Neurosci Biobehav Rev* 2000;24:843–54.
- Gaigle L, Mézack R. Age differences in the response to the formalin test in rats. *Neurobiol Aging* 1999;20:699–707.
- Lu JK, Anzalone CR, LaRoit PS. Relation of neuroendocrine function to reproductive decline during aging in the female rat. *Neurobiol Aging* 1994;15:541–4.
- Downs JL, Wise PM. The role of the brain in female reproductive aging. *Mol Cell Endocrinol* 2009;299:32–8.
- Santoro N, Brown JR, Adeli T, Skurnick JH. Characterization of reproductive hormonal dynamics in the perimenopause. *J Clin Endocrinol Metab* 1996;81:1495–501.
- Geisler L, Mézack R, Thayer WV, Butcher RL. Age-related alterations in follicular development and hormonal profiles in rats with 4-day estrous cycles. *Biol Reprod* 1990;42:6133–8.
- Robin BS, Mitchell S, Lee CE, King JC. Reconstructions of populations of luteinizing hormone releasing hormone neurons in young and middle-aged rats reveal progressive increases in subgroups expressing Fos protein on proestrus and age-related deficits. *Endocrinology* 1995;136:3823–30.
- Gore AC, Yeung G, Morrison JH, Ong T. Neuroendocrine aging in the female rat: the changing relationship of hypothalamic gonadotropin-releasing hormone neurons and N-methyl-D-aspartate receptors. *Endocrinology* 2000;141:4757–67.
- Morrison JH, Limon R, Schmidt PJ, Gore AC. Estrogen, menopause, and the aging brain: how basic neuroscience can inform hormone therapy in women. *J Neurosci* 2006;26:11933–40.
- Honey JL, Yazdani K, Pritchard GM, Cedars TL. Physiological evidence that the menopause is a neuroendocrine event. *Endocrinology* 1998;143:1637–43.
- Tjolsen A, Berge OG, Hunskaar S, Rosland JH, Hole K. The formalin test: an evaluation of the method. *Pain* 1992;51:1–17.
- Aloisi AM, Bonifazi M. Sex hormones, central nervous system and pain. *Horm Behav* 2006;50:11–7.
- Gaumond I, Arsenault P, Marchand S. The role of sex hormones on formalin-induced nociceptive responses. *Brain Res* 2002;958:139–45.
- Spencer MF, Robichaud P, Carrier JC, Marchand S. Endogenous pain modulation during the formalin test in estrogen receptor beta knockout mice. *Neuroscience* 2007;150:975–80.
- Yoshida T, Uehashi T, Mizushima D, Kimura F. Effects of neonatal testosterone treatment on sex differences in formalin-induced nociceptive behavior in rats. *Neurosci Lett* 2007;412:264–7.
- Capone F, Aloisi AM. Refinement of pain evaluation techniques. *The formalin test. Ann Ist Super Sanita* 2004;40:223–9.
- Aloisi JR, Fritscher M. Organization of the septal region in the rat brain: a Golgi EM study of lateral septal neurons. *J Comp Neurol* 1989;286:472–87.
- Gaumond I, Arsenault P, Marchand S. Specificity of female and male sex hormones on excitatory and inhibitory phases of formalin-induced nociceptive responses. *Brain Res* 2005;1052:105–11.
- Kuba T, Quimones-Jenab V. The role of female gonadal hormones in behavioral sex differences in persistent and chronic pain: clinical versus preclinical studies. *Brain Res Bull* 2005;68:65–70.
- Lei LC, Zhang YQ, Zhao ZQ. Pain-related aversion and Fos expression in the central nervous system of rats. *NeuroReport* 2004;15:67–71.
- Ohtori S, Takahashi K, Chiba T, Takahashi Y, Yamagata M, Sameda H, et al. Fos expression in the rat brain and spinal cord evoked by noxious stimulation to low back muscle and skin. *Spine* 2000;25:2425–30.
- Aloisi AM, Zimmermann M, Herdigen T. Sex-dependent effects of formalin and restraint on c-Fos expression in the septum and hippocampus of the rat. *Neuroscience* 1997;81:951–8.
- Mongeau K, Miller GA, Chang E, Anderson DJ. Neural correlates of competing fear behaviors evoked by an imminently aversive stimulus. *J Neurosci* 2003;23:3655–68.
- Keefe TJ, Rumble MC, Szpilo CD, Giordano JA, Ferreri DM. Psychological aspects of rheumatoid arthritis. *Psychiatr Clin North Am* 2003;26:33–45.
- Sheehan TP, Chambers RA, Russell DS. Reevaluation of the lateral septum: implications for neuropsychiatry. *Brain Res Brain Res Rev* 2004;46:71–117.

commentary

The chemical biology of synapses and neuronal circuits

Haruhiko Bito

Excitatory synapses are located in confined chemical spaces called the dendritic spines. These are atypical femtoliter-order microdomains where the behavior of even single molecules may have important biological consequences. Powerful chemical biological techniques have now been developed to decipher the dynamic stability of the synapses and to further interrogate the complex properties of neuronal circuits.

More than a century ago, the visionary anatomist (and later Nobel laureate) Santiago Ramon y Cajal uncovered the immense potential that the central nervous system acquires during development when innumerable connections, rather than fusions, are formed between diverse neuronal cells. These connections are called synapses, and their anatomical, physiological, biochemical and pharmacological characteristics have been studied extensively for more than 100 years. It is estimated that there are about 10^{11} synapses formed between 10^{11} neurons in each and every human brain. Through the explosion of knowledge we have gained over the past few decades, we are now able to pin down the locations of most synaptic proteins and even identify the favorite binding partners of many. Intense efforts are currently being devoted to cataloging the physical arrays of synaptic connections in the cerebral cortex and to extracting the organizing principles that govern these connectivities.

Synapses are subcellular microcompartments that are essential for a digital form of transneuronal communication, mediated by chemical mediators termed neurotransmitters. At excitatory synapses in the cerebral cortex, the most numerous synapses in our brain, the presynaptic nerve ending contains several tens to a few hundreds of synaptic vesicles filled with the neurotransmitter glutamate. Each time a terminal receives an electrical impulse from the cell soma, within one to a few milliseconds several vesicles fuse to presynaptic membrane zones and release a large amount of glutamate into the intersynaptic space. Released glutamate traverses this space by passive diffusion, and because its target, glutamate receptors, in the postsynaptic neuron are heavily enriched

in specialized membrane structures that are directly apposed to the glutamate release sites, it will take no more than a few milliseconds for the postsynaptic ligand-gated channels to open and trigger a cascade of electrochemical signaling pathways.

This postsynaptic reception of neurotransmitter signals is carried out at the surface of very tiny (<1 μ m; l.f.l. = (1 μ m)) specialized protrusions emanating from dendritic branches (Fig. 1a), which are termed dendritic 'spines'. Since Cajal's time, what we have learned about the spatial and structural regularity of these postsynaptic specializations strongly suggests that this synaptic apparatus (Fig. 1b) could serve as an elementary signal-generating (and

perhaps also signal-processing) unit of a neuron. Here, I will discuss the growing impact of chemical biology in improving our understanding of the biology of the dendritic spines, with a view toward modeling, measuring and manipulating the dynamic functions of the synapses. Indeed, we have recently witnessed a tremendous amount of excitement on this subject, fueled by the ever-growing availability of new chemical and molecular tools that allow the thought experiments of the past to become reality today.

Spines are odd chemical reaction spaces
Information transmitted by synaptic activity is first received at the dendritic




Figure 1 Dendritic spines are highly atypical chemical reaction spaces at the boundary between mass and stochastic reaction schemes. (a) Golgi staining of dendritic branches in a typical neuron (courtesy of H. Okuno). (b) On the basis of the standard size of three-dimensionally reconstructed dendritic spines, we can estimate that in each spine, most signaling molecules are likely to be present only in small numbers (1 to about 100). The image in panel b is reproduced from ref. 33, with permission from Elsevier.

Box 1 | Back-of-the-envelope calculations

Estimating the number of molecules in a standard spine (~0.16 fl), given that concentration \times volume \times Avogadro constant = number of molecules:
 At 1 μM , $1 \times 10^{-6} \text{ (mol l}^{-1}\text{)} \times 0.16 \times 10^{-15} \text{ (l)} \times 6.02 \times 10^{23} = 1 \times 10^2$ molecules
 At 10 nM, $10 \times 10^{-9} \text{ (mol l}^{-1}\text{)} \times 0.16 \times 10^{-15} \text{ (l)} \times 6.02 \times 10^{23} = 1$ molecule

spines. But despite the fundamental importance of these structures, neuroscientists have yet to answer some of the simplest questions about the spines. For example, let us consider the physical scale of the dendritic spines (Fig. 2a), where the synapses are. A rough estimate of the mean volume of the dendritic spines, obtained through serial electron microscopic reconstructions of excitatory synapses, shows them to be smaller than 1 fl (ref. 4) (Fig. 1b). A back-of-the-envelope calculation (Box 1) will then reveal that a densely enriched synaptic protein, such as glutamate receptor, at about 1 μM concentration will represent about 100 molecules per synapse. Strikingly, this also means that the presence of one single molecule in an average spine will represent a 10 nM concentration.

A dissociation constant of this order is frequently observed in *in vitro* reactions between a pair of high-affinity interacting proteins or between an enzyme and a substrate. Thus, one might imagine that in a femtoliter-order space such as the spines, the addition or loss of each single molecule may perhaps be biologically important. This is assuming that a chemical mass reaction based on free reaction and diffusion can occur in spines at the nano-to-meso scale. However, we do not actually know how diffuse synaptic proteins enriched in the spines are, as many of them are bound to either membrane-spanning proteins (such as channels and receptors) or cytoskeletal proteins (such as actin and actin-interacting) or both (Fig. 2b). It has been speculated that stochastic processes might be important in such tiny biological spaces, and indeed, models of stochastic reactions have been shown to reliably predict some of the signal transduction outcomes within a spine¹. Conversely, experimental data suggest that the opening of a single presynaptic Ca^{2+} channel, for instance, can trigger neurotransmitter release², whereas even at the peak of a synaptic response, less than one NMDA receptor channel may be open on average³, thus lending support to a possible stochastic scenario of synaptic transmission-triggered Ca^{2+} signaling. On the other hand, classical models of actin remodeling still remain useful for describing actin polymerization states within a spine⁴, and a simple linear inverse correlation was found between spine length and PDZ domain-based interactions of PSD-95 and SynGAP, two dominantly expressed proteins at maturing synapses⁵.

Thus, the truth may be a mixture of these two possibilities, to say the least, and dendritic spines appear to represent highly atypical chemical reaction spaces at the

boundary between mass and stochastic reaction schemes. As these tiny synapses may behave quite distinctly from continuous fluids or ideal gas, one may need to go beyond the classical Boltzmann equation and adopt other frameworks, such as the Langevin⁶ or Fokker-Planck equations⁷, in order to more accurately describe and model the chemistry of synapses.

The dynamic stability of the synapses

For many decades, *in vivo* and *in vitro* recordings of synaptic transmission in many brain areas have shown remarkable stability in the magnitude of the electrical responses that can be elicited at synapses. Recent *in vivo* live imaging studies of dendritic spines in superficial layers of the cerebral cortex confirms that a large majority of synapses are indeed stable over a long period of time^{8,9}. Similarly, the molecular make-up of the synapses appears to be quite stable and uniform across synapses, both presynaptically and postsynaptically, when measured in fixed brain tissues and cells that have already reached adulthood and maturity. However, given that most neurons in our brain lose their ability to divide during early development, how can such regularity of structural and molecular components persist over many months, if not years or a whole lifetime, in a majority of synapses?

Clearly, the replacement of both structural (membrane lipids and cytoskeleton) and proteinous components of the synapses has to take place under rigorous control, as most neuronal proteins degrade within hours to a few days once translated. In such a self-organized system, regulation of protein trafficking, as well as mRNA transcriptions and translation, may be truly vital.

Even more surprising, however, are recent measurements taken within the dendritic spines, both *in vitro* and *in vivo*, indicating that the turnover of synaptic proteins can be very fast, on the order of seconds to minutes^{10,11}. Proteins may even travel across—and be shared between—neighboring spines within the same dendritic tree, without losing their activity¹². Furthermore, during synaptic plasticity, which is a sudden change in the efficacy of the synaptic




Figure 2 | Going inside the spines and exploring synaptic proteins that support the function of individual synapses. (a) Golgi stain of individual spines (Courtesy of H. Okuno). (b) Analyzing one synapse at a time. Proteins expressed in dendritic spines are highly organized and form clusters; scaffolding proteins are directly and indirectly associated with membrane-spanning proteins such as channels and receptors, while also interacting with F-actin or F-actin-interacting proteins. (c) Localized distribution of activated caged fluorescein isothiocyanate (FITC) near the synaptic sites where focused UV light was shined.

fluorophore-labeled antibodies has revealed that lateral diffusion of synaptic receptors can be unexpectedly fast. Quantitative imaging using fluorescence resonance energy transfer (and fluorescence lifetime measurement) has also proven to be very effective in measuring signal transduction at single-synapse resolution.

Although not all the mysteries have been solved, these experiments collectively provide an increasingly quantitative picture of the protein machines that are present at synapses. Through engineering brighter dyes and tags with improved cell penetration, enhancing protein labeling techniques and

Clicking synapses 'on' and 'off'

As discussed above, the dendritic spines represent one of the most interesting challenges for scientists elucidating the elementary chemical interface at the intersection of neural circuitry and




Figure 3 | Deconstructing and reconstructing cognition. Manipulating and interrogating synapses and neuronal circuits, by triggering and altering synaptic activity, has simultaneous effects at several distinct hierarchical levels: behavioral, systems, microcircuits, cellular and molecular. Intuitively, these levels can be compared to various complexities of electronic computation.

intracellular signal transduction cascades. There are, however, many practical issues that need to be overcome to expand our knowledge in this area. We still do not even know the basic anatomical rules governing interaction and interference between physically proximal and distal synapses. Are there any mechanisms to amplify chemical signaling across distance? Do topologically organized chemical gradients exist inside dendrites, and do they play a significant role in modulating synaptic information processing?

Furthermore, we currently have little understanding of the functional states of synapses on a cell-assembly basis. What is the overall signaling impact of synaptic inputs at physically non-adjacent, yet temporally correlated, connections between neurons in an active cell assembly? What are the exact numbers (few or many?) and the locations (random or topologically organized?) of the synapses that are usually engaged in performing any given neuronal or cognitive task?

To address these questions, we clearly need to push the current limits of data acquisition and promote massively parallel imaging approaches using a battery of sensors of synaptic activity and of specific signal transduction pathways (obtained through careful chemical biological designs or encoded genetically), with both synaptic and cellular resolutions.

Our current ability to deconstruct the chemistry inside the synapses in depth gives hope that we will soon be able to reconstruct the function of a synapse and causally connect the networks of chemical reactions inside synapses with the encoding and representation of information within the neuronal circuit. The most direct way to do this is to choose a synapse of interest, identify unitary events and artificially manipulate them, preferably mimicking them so as to replace endogenous information inputs, and test how one has managed to alter (compromise or ameliorate) the neuronal circuitry's outputs.

Pioneering chemists have previously designed 'caged' molecules, in which photosensitive reaction moieties such as carboxy-2-nitrobenzyl (CNB), 4-methoxy-7-nitroindolyl (MNI), carboxymethoxy-5,7-dinitroindolyl (CDNI) or ruthenium-bipyridine-trimethylphosphine (RUBI) groups mask the functional groups of the molecule^{13,14}. Exposure to light of a specific wavelength triggers photocatalysis of these moieties, thereby revealing the protected groups and 'activating' the molecules instantaneously (Fig. 2c). Shining light on synapses bathed with caged neurotransmitters has provided a powerful

© 2010 Nature America, Inc. All rights reserved.

means to demonstrate the role of glutamate as a neurotransmitter, to map functional circuit connectivities *in vitro*²⁵ and *in vivo*²⁶ and to uncover mechanisms of synaptic plasticity at single-spine resolution^{10,18,26} at millisecond to second temporal resolution. Additionally, chemical genetic strategies such chemical dimerization²⁷, inducible protein stabilization²⁸ or molecular re-engineering²⁹ are actively sought after to dissect the role of specific chemical signaling pathways.

Systematically manipulating a specific set of synapses *in vivo* will unavoidably modify neural function at all levels: molecular, cellular, circuits, systems, and behavioral (Fig. 3). Spectacular achievements using optogenetic technologies³⁰ demonstrate the critical importance of precisely targeting specific sets of synapses within a neuronal circuit. Clearly, finding the most astute combination of remotely activatable chemicals, circuit-oriented genetic manipulation and massively parallel high-throughput readouts at multiple levels, *ex vivo* or *in vivo*, will pave the way for a successful chemical biology exploration of the synapses and neuronal circuits.

What lies ahead?

More and more, our efforts to tease apart the principles of chemistry and how they pertain to communication inside and across synapses are met with success. Through these continuing efforts, we will better resolve the molecular states and kinetics of the most basic chemical events, events that are central to the computation carried out in each of the 10^{14} synapses. This level of deconstruction of the chemical nature of the synapses now provides us with an unprecedented window of opportunity to address a fundamental

desire of a synaptologist: can we sneak into the synapses' design and even perhaps reprogram them?

Although constructing and designing synapses may have sounded preposterous five years ago, it does not sound so far-fetched in 2010. In fact, a tremendous amount of translational research money is now being devoted to the study of the malfunction of so-called "induced synapses" formed *in vitro* between neuron-like cells massively derived from human induced pluripotent stem cells, which were obtained from living neuropsychiatric patients. If we could identify the faulty information processing event involved at induced synapses in such diverse neurons, we might think of using drugs to correct synaptic dysregulation there. The possibility of *en masse* reprogramming of embryonic stem cells into "induced neurons"³¹ and of building layered neuronal structures among them³² in fact, provides hope that drug screening and pre-clinical trials performed to study neuronal activity in human induced neuronal circuits *in vitro* may become a reality in the not-too-distant future.

Haruhiko Bito is in the Department of Neurochemistry at University of Tokyo Graduate School of Medicine, Tokyo, Japan.
e-mail: hito@nu.u-tokyo.ac.jp

References

1. Ramón y Cajal, S. *Textura del sistema nervioso del hombre y de los vertebrados* (Librería de Nicolás Moya, Madrid, 1899).
2. Kandel, E.R., Schwartz, J.H. & Jessell, T.M. *Principles of Neural Science*, 4th ed. (McGraw-Hill, New York, 2000).
3. Markram, H. *Nat. Rev. Neurosci.* **7**, 153–160 (2006).
4. Harris, K.M. & Stevens, J.K. *J. Neurosci.* **9**, 2882–2897 (1989).
5. Bhalla, U.S. *Biophys. J.* **87**, 745–753 (2004).
6. Stanley, E.F. *Neuron* **11**, 1007–1011 (1993).
7. Nimchinsky, E.A., Yasuda, R., Oertner, T.G. & Svoboda, K. *J. Neurosci.* **24**, 2054–2064 (2004).
8. Honbara, N., Mizusawa, M., Noguchi, J., Ellis-Davies, G.C. & Kasai, H. *Neuron* **57**, 719–729 (2008).

9. Novak, M., Dai, T., Fujyoshi, Y., Takemoto-Kimura, S. & Bito, H. *J. Neurosci.* **26**, 774–777 (2006).
10. Kasai, H., Fukuda, M., Watanabe, S., Hayashi-Tokaji, A. & Noguchi, J. *Trends Neurosci.* **33**, 121–129 (2010).
11. Harrison, L.M., David, O. & Friston, K.J. *Phil. Trans. R. Soc. Lond. B* **360**, 1075–1091 (2005).
12. Jing, G., Pan, F. & Com, M.L. *Nature* **463**, 93–95 (2009).
13. Honbara, N. & Svoboda, K. *Nat. Rev. Neurosci.* **10**, 647–658 (2009).
14. Kratochvil, T., Inoue, A., Bito, H., Sobue, K. & Ohabe, S. *J. Neurosci.* **26**, 7693–7706 (2006).
15. Gross, N.W., Wémeur, R.M., Bhirani, L. & Svoboda, K. *PLoS Biol.* **4**, e370 (2006).
16. Tsunashima, S. *et al. PLoS Biol.* **4**, e271 (2006).
17. Kowalek, H.W. & Malhotra, R. *Neuron* **61**, 340–350 (2009).
18. Matsuzaka, M., Honkura, N., Ellis-Davies, G.C. & Kasai, H. *Nature* **429**, 791–796 (2004).
19. Gheysmans, B.N., Adams, S.R., Ellsman, M.H. & Tsien, R.Y. *Science* **312**, 217–224 (2006).
20. Tsunashima, S. *et al. Nature* **48**, 189–190 (2005).
21. Ciampini, M., Nelson, L., Papeghoian, G., Corrie, J.E.T. & Ogden, D.J. *Neurosci. Methods* **112**, 29–42 (2001).
22. Ellis-Davies, G.C. *Nat. Methods* **4**, 619–628 (2007).
23. Inno, E. *et al. Front. Neural Circuits* **3**, 2 (2009).
24. Yoshimura, T., Dauter, J.L. & Callaway, E.M. *Nature* **433**, 868–873 (2005).
25. Honkura, N., Ellis-Davies, G.C. & Kasai, H. *J. Neurophysiol.* **99**, 1535–1541 (2008).
26. Tanaka, K. & Augustine, G.J. *Neuron* **59**, 608–620 (2008).
27. Prinsak, M.N. *et al. Chem. Biol.* **13**, 172 (1994).
28. Bonasirski, L.A., Chen, L.C., Maynard-Smith, L.A., Ooi, A.G. & Waudesch, T.J. *Cell* **126**, 995–1004 (2006).
29. Bishop, A.C. *et al. Nature* **407**, 395–401 (2000).
30. Gradinaru, V. *et al. Cell* **141**, 154–165 (2010).
31. Vierbuchen, T. *et al. Nature* **463**, 1035–1041 (2010).
32. Ehrlich, M. *et al. Cell Stem Cell* **3**, 519–532 (2008).
33. Bourne, J.R. & Harris, K.M. *Curr. Opin. Neurobiol.* **17**, 381–386 (2007).

© 2010 Nature America, Inc. All rights reserved.

Acknowledgments

This commentary is based on past and ongoing works that received support from grants-in-aid from the Japan Society for Promotion of Science, the Ministry of Education, Culture, Sports, Science and Technology of Japan, the Ministry of Health, Labor and Welfare of Japan, a Global Center of Excellence program "Comprehensive Center for Education and Research of the Chemical Biology of Disease" and awards from the Astellas Foundation for Research on Metabolic Disorders, the Cell Science Foundation, the Takeda Science Foundation and the Toray Science Foundation. The author thanks H. Okuno, S. Takemoto-Kimura and H. Fujii for critical comments on an earlier version of the manuscript.

Competing interests statement

The author declares no competing financial interests.



Contents lists available at ScienceDirect

NeuroToxicology

Up-regulation of neurotrophin-related gene expression in mouse hippocampus following low-level toluene exposure

Tin-Tin Win-Shwe^a, Shinji Tsukahara^a, Shoji Yamamoto^a, Atsushi Fukushima^a, Naoki Kunugita^b, Keiichi Arashidani^b, Hidekazu Fujimaki^{a,b,*}

^aResearch Center for Environmental Risk, National Institute for Environmental Studies, 16-2 Onogawa, Tsukuba, Ibaraki 305-8506, Japan
^bUniversity of Occupational and Environmental Health, Japan 1-1 Isegaki, Yahatanishi-ku, Kitakyushu, Fukuoka 807-8555, Japan

ARTICLE INFO

Article history:

Received 8 May 2009
Accepted 17 November 2009
Available online 22 November 2009

Keywords:

Neurotrophins
Toluene
Brain
Allergy
Strain difference

ABSTRACT

To investigate the role of strain differences in sensitivity to low-level toluene exposure on neurotrophins and their receptor levels in the mouse hippocampus, 8-week-old male C3H/HeN, BALB/c and C57BL/6J mice were exposed to 0, 5, 50, or 500 ppm toluene for 6 h per day, 5 days per week for 6 weeks in an inhalation chamber. We examined the expressions of neurotrophin-related genes and receptors in the mouse hippocampus using real-time reverse transcription polymerase chain reaction (RT-PCR). The expression of nerve growth factor (NGF), brain-derived neurotrophic factor (BDNF), tyrosine kinase (Trk) A, and TrkB mRNAs in the C3H/HeN mice hippocampus was significantly higher in the mice exposed to 500 ppm toluene. Among the three strains of mice, the C3H/HeN mice seemed to be sensitive to toluene exposure. To examine the combined effect of toluene exposure and allergic challenge, the C3H/HeN mice stimulated with ovalbumin were exposed to toluene. The allergy group of C3H/HeN mice showed significantly elevated level of NGF mRNA in the hippocampus following exposure to 50 ppm toluene. Then, we also examined the expression of transcription factor, dopamine markers and oxidative stress marker in the hippocampus of sensitive strain C3H/HeN mice and found that the expression of CREB1 mRNA was significantly increased at 50 ppm toluene. In immunohistochemical analysis, the density of the NGF-immunoreactive signal was significantly stronger in the hippocampal CA3 region of the C3H/HeN mice exposed to 500 ppm toluene in non-allergy group and 50 ppm in allergy group. Our results indicate that low-level toluene exposure may induce up-regulation of neurotrophin-related gene expression in the mouse hippocampus depending on the mouse strain and an allergic stimulation in sensitive strain may decrease the threshold for sensitivity at lower exposure level.

© 2009 Elsevier Inc. All rights reserved.

1. Introduction

Toluene is a well-known neurotoxicant and the central nervous system is a primary target in both acute and chronic exposure of toluene-toxicity (US EPA, 1994). Previous data from *in vivo* and *in vitro* animal studies strongly support that the hippocampus is target for toluene (Korbo et al., 1996; Terashi et al., 1997; Gelazonia et al., 2006). In addition, we had measured brain toluene levels by solid-phase microextraction technique and reported the hippocampal toluene levels following inhalation exposure (Nakajima et al., 2006) and intraperitoneal administration (Win-Shwe et al., 2007a). Little is known about the role of hippocampus in allergic sensitivity. Toluene exposure may cause alterations in the hippocampal functions of individuals with underlying allergic or stressful conditions (Lindvall et al., 1992). The number of corticotrophin-releasing hormone (CRH)-immunoreactive (ir) neurons in the hypothalamus, the

number of adrenocorticotrophic hormone (ACTH)-ir cells, and ACTH-mRNA expression in the anterior pituitary of allergy (AG) group mice were significantly higher than those of non-allergy (NAG) group mice (Sari et al., 2004, 2006). Recently, we have reported that the ovalbumin (OVA) immunization may enhance the sensitivity to toluene exposure by modulating the NMDA receptor subunit expressions in the olfactory bulb of allergic mouse model (Win-Shwe et al., *in press*). Therefore, in the present study, we used OVA as a stressor to investigate whether exposure to low levels of toluene, in combination with OVA, might also affect hippocampal neurotrophin-related gene expression.

Neurotrophins are a group of structurally related polypeptides that support the survival, differentiation, and maintenance of neuronal populations expressing appropriate high-affinity neurotrophin receptors. Neurons in the hippocampus are maintained by neurotrophins, such as nerve growth factor (NGF), brain-derived neurotrophic factor (BDNF), and the tyrosine kinase (Trk) family of neurotrophin receptors. Neurotrophins and their related receptors have been identified as targets for neurotoxicants and are known to play a role in bidirectional signaling between cells of the

* Corresponding author. Tel.: +81 29 850 2518; fax: +81 29 850 2518.
E-mail address: fujimaki@nies.go.jp (H. Fujimaki).

0161-813X/\$ - see front matter © 2009 Elsevier Inc. All rights reserved.
doi:10.1016/j.neuro.2009.11.004

immune and nervous systems. Recently, many researchers have examined the functions of neurotrophic factors and their receptors in the brain, because these factors and receptors might be possible targets for neurotoxicants like lead (Kidd et al., 2008) and chlorpyrifos (Betancourt et al., 2006).

While the neurobehavioral and neurotoxic effects of toluene have been thoroughly examined, the mechanisms whereby toluene exerts its effects in the brain are not fully understood. The occupational exposure limit for toluene is 50 ppm in Japan (Japan Society for Occupational Health, 1994) and recently updated threshold limit value is 20 ppm in United States (ACGIH, 2006). Exposure to low concentrations of toluene leads to persistent deficits in spatial learning and memory function in a rat model (von Euler et al., 1993, 2000). From a dosimetric analysis of behavioral effects of acute toluene exposure in rat and humans, behavioral outcomes depend on internal dose rather than external dose and duration of exposure (Benignus et al., 1998). Recently, a study using a mouse model showed that mice exposed to a low level of toluene showed little change in the number of rewards obtained in a waiting-for-reward task, whereas mice exposed to a high concentration of toluene exhibited a significantly poorer performance, suggesting that exposure to high concentrations of toluene had a significant impact on cognitive and/or psychomotor function (Bowen and McDonald, 2009). Toluene also affects the neurotransmitter systems in the hippocampus. Our previous study indicates that the extracellular levels of neurotransmitter glutamate and taurine in the mouse hippocampus were rapidly and reversibly increased within 30 min after the toluene administration in a dose-dependent manner and returned to the basal level by 1 h, however, the glycine and GABA were stable and no significant changes were observed (Win-Shwe et al., 2007a).

We hypothesized that toluene exposure may affect the function of the hippocampus via the modulation of neurotrophin-related genes and signaling pathway in strain-dependent manner and allergic stimulation may influence the threshold for sensitivity. Therefore, the present study was designed to investigate the intra-species variation in sensitivity to the expression of toluene-mediated neurotrophins and related receptors in the mouse hippocampus and examine the combined effect of toluene

exposure and allergic challenge on neurogenesis-related markers in the hippocampus of sensitive mouse strain.

2. Materials and methods

2.1. Animals

Seven-week-old male C3H/HeN, BALB/c and C57BL/10 mice were obtained from Japan SLC Inc. (Shizuoka, Japan) and were used at an age of 8 weeks. Food and water were given *ad libitum*. The mice were housed in plastic cages under controlled environmental conditions (temperature, 23 ± 0.5 °C; humidity, 50 ± 5%; lights between 07:00 and 19:00 h). This study was approved by the Ethics Committee for Experimental Animals of the University of Occupational and Environmental Health, Japan.

2.2. Experimental design

To detect the strain differences in response to toluene exposure, C3H/HeN, BALB/c and C57BL/10 mice ($n = 9$ each) were exposed to either a filtered air control (0 ppm) or 5, 50, or 500 ppm of toluene for 6 h (from 10:00 to 16:00 h) per day, 5 days per week for 6 weeks. One day following the final toluene inhalation, the hippocampi from nine mice (six for mRNA analysis and three for immunohistochemical analysis) from each strain were collected. Among the three strains of mice, C3H/HeN mice seemed to be most sensitive to toluene exposure and this mouse strain was used to detect the combined effect of toluene exposure and OVA immunization. C3H/HeN mice ($n = 9$ each, six for mRNA analysis and three for immunohistochemical analysis) were exposed to toluene as same schedule and the AC groups were treated with OVA on days 0, 7, 21 and 42 approximately 1 h before toluene exposure. These mice were injected with 10 µg of OVA plus 2 mg of alum intraperitoneally on day 0 and 10 µg of OVA only on day 7. Each of these mice was then challenged with nebulized OVA as a booster once every 3 weeks (days 21 and 42) during the exposure period, as described previously (Fujimaki et al., 2007). Fig. 1 shows the detail experimental schedule for the present study.




Fig. 1. (A) Experimental schedule for detecting different sensitivity in three strains of mice and (B) experimental schedule for investigating the combined effect of toluene and OVA immunization in C3H/HeN mice. Mice were exposed to a filtered air control (0 ppm), 5, 50, or 500 ppm of toluene for 6 h (from 10:00 to 16:00 h) per day, 5 days per week for 6 weeks. The allergy (AC) groups were treated with OVA on days 0, 7, 21 and 42 approximately 1 h before toluene exposure.

2.3. Generation of toluene

Toluene vapor was generated using an organic solvent gas generator (Shibata Scientific Technology, Ltd.) and was diluted with clean, filtered air to achieve the desired gas concentrations; the vapor was then introduced into a stainless steel and glass chamber, as described previously (Hori et al., 1999). The average toluene levels (mean ± SD) in the 5-, 50- and 500-ppm exposure groups were 4.9 ± 0.6, 49.5 ± 4.4, and 496.7 ± 17.9 ppm, respectively.

2.4. Quantitative real-time PCR

One day following the final toluene inhalation, the mice were sacrificed under deep pentobarbital anesthesia and the hippocampus was collected from each group of three strains of mice, frozen quickly in liquid nitrogen, and then stored at -80 °C until the total RNA was extracted. The expression levels of NGF, BDNF, TrkA, TrkB, calcium/calmodulin-dependent protein kinase (CaMK)-IV, cyclic AMP responsive element binding protein (CREB)-1, CREB2, D1, D2, tyrosine hydroxylase (TH), heme oxygenase (HO)-1, Bax, Bbc3 and 18S mRNA in the hippocampus were estimated using quantitative real-time PCR analysis with the ABI Prism 7000 Sequence Detection System (Applied Biosystems Inc., USA), as described previously (Win-Shwe et al., 2007b). The tissue 18S level was examined as an internal control. The primer sequences used in the present study were shown in Table 1. Some primers (BDNF, ID_12064; TrkA, ID_18211; TrkB, ID_18212; D1, ID_13488; D2, ID_13489; TH, ID_21823; Bbc3, ID_170770; Bax, ID_12028; HO-1, ID_15368) were purchased from Qiagen, Sample & Assay Technologies (Qiagen GmbH, Germany). Data were analyzed using the comparative threshold cycle method. The relative expression levels of each mRNA were normalized individually according to the 18S content in the respective samples and were expressed as mRNA signals per 18S.

2.5. Immunohistochemistry

On the day after the final toluene exposure, the brains were removed from three C3H/HeN mice from each group of control and toluene-exposed non-allergic and AC groups after the animals had been deeply anesthetized with sodium pentobarbital; the brains were then fixed with 10% formalin. The fixed brains were dehydrated using a graded series of ethanol, cleared with xylene, and embedded in paraffin. Coronal paraffin sections were cut at a thickness of 10 µm using a cryostat and were mounted on 3-aminopropyltriethoxysilane-coated glass slides. Cells containing NGF polypeptide in the hippocampus were detected immunohistochemically.

To determine the expression levels of NGF polypeptide in the hippocampus, series of NGF-immunostained sections taken from 3 mice in each group (4 sections for each mouse) were used to analyze the density of NGF-immunoreactive signals in the

Table 1
Primer sequences used in real-time PCR.

Gene	Primer	Sequence
18S rRNA	Forward	5'-TACCACTCCACGAAGGAG-3'
	Reverse	5'-TCCCTCATGATGATCC-3'
NGF	Forward	5'-TGGCCCTCAGGAGGAGATC-3'
	Reverse	5'-CAGCTTCTTACTGCGCCAG-3'
CREB1	Forward	5'-ATACGCCATCGCAGAAC-3'
	Reverse	5'-AATGCCATCGCAGAAC-3'
CREB2	Forward	5'-CAATCTCTGGATCGGAGC-3'
	Reverse	5'-AATCAGCCCTGCTCTTGA-3'
CaMKIV	Forward	5'-TCTGTTTTCAGTCACTG-3'
	Reverse	5'-TCTGTTTTCAGTCACTG-3'

hippocampus. Photomicrographic digital images (150 dpi, 256 scales) of the hippocampal regions were taken using a CCD camera connected to a light microscope. In the pyramidal cell layer of the CA1 and CA3 regions, the granular cell layer of the DC, and all of these regions combined, the densities of the NGF-immunoreactive signals were determined with the aid of a computer running image-analysis software (ImageJ 1.38X; National Institutes of Health, Bethesda, MD, USA). The density of the NGF-immunoreactive signal was expressed as a percentage of the NGF-immunoreactive area darker than 70 of the 256 units in the total area of interest.

2.6. Statistical analysis

All data were expressed as the mean ± standard error (S.E.). The statistical analysis was performed using the StatMate II statistical analysis system for Microsoft Excel, Version 5.0 (Nankaido Inc., Tokyo, Japan). Dose-response data were analyzed using a one-way analysis of variance with a Bonferroni/Dunn post-hoc analysis. A value of $P < 0.05$ was considered statistically significant.

3. Results

3.1. Detection of intra-species variation of sensitivity to toluene exposure

To explore the sensitive strain of mice to toluene exposure, we compared the expression of various marker mRNA in the hippocampus of C3H/HeN, BALB/c and C57BL/10 mice.

3.1.1. Neurotrophins and related receptors

NGF mRNA expression in the hippocampus of the C3H/HeN mice was significantly increased in mice exposed to 500 ppm of toluene ($P < 0.01$, Fig. 2a). The mRNA expression of TrkA, a receptor for NGF, was also significantly increased in mice exposed to 500 ppm of toluene ($P < 0.01$, Fig. 2d). In BALB/c mice, however, no significant change in NGF or TrkA receptor expression was observed after toluene exposure (Fig. 2b and e). Moreover, no significant difference in NGF or TrkA receptor expression was observed in C57BL/10 mice after toluene exposure (Fig. 2c and f).

BDNF mRNA expression in the hippocampus of C3H/HeN mice was significantly increased in mice exposed to 500 ppm of toluene ($P < 0.01$, Fig. 3a). The mRNA expression of TrkB, the receptor for BDNF, was also significantly increased in mice exposed to 500 ppm of toluene ($P < 0.01$, Fig. 3d). In BALB/c mice, BDNF mRNA expression was significantly increased in the mice exposed to 500 ppm of toluene, compared with expression in the control group ($P < 0.01$, Fig. 3b). No significant change in TrkB receptor expression was observed in BALB/c mice after toluene exposure (Fig. 3e). Furthermore, no significant difference in BDNF or TrkB receptor expression was observed in C57BL/10 mice after toluene exposure (Fig. 3c and f).

3.1.2. Transcription factor

In NGF signal transduction pathways, multiple protein kinases translocate to the nucleus to activate transcriptional activator CREB. One of these kinases is CaMKIV, which phosphorylates CREB. We examined the expression of CaMKIV, and did not observe any difference of CaMKIV mRNA expression between the mice exposed to toluene and filtered air (data not shown). We also detected the expression of NGF-related transcription factors, CREB1 mRNA in the hippocampus of C3H/HeN mice after toluene exposure. An increased CREB1 mRNA was found in mice exposed to 50 ppm of toluene compared to that of mice exposed to filtered air ($P < 0.05$,




Fig. 2. Expression of NGF and TrkA in the hippocampus of three strains of mice. (a) NGF and (b) TrkA in C3H/HeN mice; (c) NGF and (d) TrkA in BALB/c mice; and (e) NGF and (f) TrkA in C57BL/10 mice following exposure to 0, 5, 50 or 500 ppm of toluene for 6 weeks. Each bar represents the mean \pm S.E. ($n = 6$) ($*P < 0.05$).




Fig. 4. Expression of CREB1 and CREB2 in the hippocampus of C3H/HeN mice. (a) CREB1 and (b) CREB2 mRNAs in mice following exposure to 0, 5, 50 or 500 ppm of toluene for 6 weeks. Each bar represents the mean \pm S.E. ($n = 6$) ($*P < 0.05$).

In addition, we examined the expression of CREB2 mRNA, which functions to repress CREB1 activity. We did not observe any significant change of CREB2 mRNA (Fig. 4b).

3.1.3. Dopamine markers

Then, we postulated that the signaling pathway mediating the effect of toluene may be common or specific in the hippocampus for this C3H/HeN of mice. It is possible that the dopamine receptor signaling pathway may cross-talk with neurotrophin signaling pathways following exposure to toluene. We did not find any difference of D1 mRNA expression between mice exposed to toluene and filtered air (Fig. 5a). Although statistically insignificant, the results indicate a trend of increased D2 mRNA expression in mice exposed to 50 ppm of toluene (Fig. 5b). Tyrosine hydroxylase (TH) has a link with

dopamine receptor activity. Therefore, we examined the expression of TH mRNA in the hippocampus of mice exposed to toluene. There was a tendency of higher TH mRNA expression in mice exposed to toluene as compared to that of mice exposed to filtered air (Fig. 5c).

3.1.4. Oxidative stress marker and apoptotic markers

NGF served as a growth factor as well as a regulator for cell survival. Balance of NGF-TrkA-MEK1-ERK1/2-CREB and NGF-TrkA-MEK3/6-p38 MAP pathway could contribute to NGF-induced cell death (Yan et al., 2002). To investigate the effect of toluene exposure on oxidative stress in the hippocampus of C3H/HeN mice, we examined the expression of oxidative stress marker, HO-1 mRNA. We found a significantly increased HO-1 mRNA expression in mice exposed to 500 ppm toluene as compared to that of mice




Fig. 3. Expression of BDNF and TrkB in the hippocampus of three strains of mice. (a) BDNF and (b) TrkB in C3H/HeN mice; (c) BDNF and (d) TrkB in BALB/c mice; and (e) BDNF and (f) TrkB in C57BL/10 mice following exposure to 0, 5, 50 or 500 ppm of toluene for 6 weeks. Each bar represents the mean \pm S.E. ($n = 6$) ($*P < 0.01$).




Fig. 5. Expression of D1, D2 and TH in the hippocampus of C3H/HeN mice. (a) D1, (b) D2 and (c) TH mRNAs in mice following exposure to 0, 5, 50 or 500 ppm of toluene for 6 weeks. Each bar represents the mean \pm S.E. ($n = 6$).




Fig. 6. Expression of HO-1, Bax and Bcl-2 in the hippocampus of C3H/HeN mice. (a) HO-1, (b) Bax and (c) Bcl-2 mRNAs in mice following exposure to 0, 5, 50 or 500 ppm of toluene for 6 weeks. Each bar represents the mean \pm S.E. ($n = 6$) ($*P < 0.05$).




Fig. 7. Representative digital photomicrographs of NGF-immunostained sections taken from the hippocampal CA3 region in C3H/HeN mice following exposure to toluene (a) 0 ppm and (b) 500 ppm for 6 weeks (Scale bars = 50 μm). (c) Density of NGF-immunoreactivity in the hippocampal CA3 region of C3H/HeN mice following exposure to 0 and 500 ppm of toluene for 6 weeks. Each bar represents the mean ± S.E. (n = 3).

exposed to filtered air ($P < 0.05$, Fig. 6a). In addition, to assess the apoptotic activity, we measured the expression of apoptotic related genes (Bax for pro-apoptosis and Bcl-2 binding component) for anti-apoptosis). There was a tendency of higher Bax mRNA expression in mice exposed to toluene, but not statistically significant (Fig. 6b). We did not observe any significant difference of Bcl-2 mRNA expression between mice exposed to toluene and filtered air (Fig. 6c).

3.1.5. NGF-immunoreactivity in the hippocampus of C3H/HeN mice

To confirm the effect of toluene on the expression levels of NGF-polypeptide in C3H/HeN mice, we examined the density of NGF-immunoreactive signals in the CA1–3 and DG hippocampal regions. In the hippocampus of 500 ppm toluene-exposed C3H/HeN mice, intense immunoreactive signals for NGF were found in the pyramidal cell layers of the CA1–3 regions and in the granular cell layer of the DG. The representative digital photomicrograph of the CA3 of mice exposed to 0 ppm (Fig. 7a) and 500 ppm (Fig. 7b) and, the percentage of NGF-immunoreactivity (Fig. 7c) were shown.

Taken together, our findings indicate that C3H/HeN mice seemed to be sensitive to toluene exposure than the other two strains of mice.

3.2. Detection of combined effect of toluene exposure and allergic stimulation on NGF expression in the hippocampus of sensitive mouse strain

To examine whether the allergic condition affect NGF expression in the hippocampus modulated by exposure to toluene, C3H/HeN mice were stimulated with OVA during toluene exposure period. A significant increase in NGF mRNA




Fig. 9. Representative digital photomicrographs of NGF-immunostained sections taken from the hippocampal CA3 region in AG group of C3H/HeN mice following exposure to toluene (a) 0 ppm and (b) 500 ppm for 6 weeks (Scale bars = 50 μm). (c) Density of NGF-immunoreactivity in the hippocampal CA3 region of AG group of C3H/HeN mice following exposure to 0 and 500 ppm of toluene for 6 weeks. Each bar represents the mean ± S.E. (n = 3). (** $P < 0.01$, AG = allergy).

4. Discussion

The major findings of our present study are that low levels of toluene exposure can up-regulate the expressions of neurotrophins and their related receptors in the mouse hippocampus in strain-dependent manner, and allergic stimulation lower the threshold for the sensitivity to toluene in the most sensitive strain C3H/HeN mice. Little is known about differences in sensitivity to volatile organic compounds (VOCs) among mouse strains. Immunological studies have indicated that C3H mice are more resistant to infection with *Salmonella*, *Leishmania*, or *Mycobacterium avium* than C57BL/6 mice (Kamath et al., 2004). Acute restraint stress has also been reported to decrease the infiltration of inflammatory cells in BALB/c mice, but not in C57BL/6 mice (Okuyama et al., 2007). Neurological studies have also demonstrated that different sensory abilities may influence strain differences in learning and memory performances (Brown and Wong, 2007). In toxicological aspect, our research group showed that the strain differences influence the extracellular amino acid neurotransmitter levels in the hippocampus of major histocompatibility complex (MHC) congenic mice (Win-Shwe et al., 2009) and NMDA receptor subunit expression in olfactory bulb of an allergic mouse model (Win-Shwe et al., in press) following toluene exposure. In view of these previous results, the three above-mentioned mice strains were used to compare the effect of low-level toluene on neurotrophin-related gene expression in the hippocampus.

Our present data clearly showed that toluene exposure alone significantly increased the mRNA expressions of NGF, TrkA, BDNF and TrkB in the hippocampus of C3H/HeN mice exposed to 500 ppm of toluene, but not in the BALB/c and C57BL mice. It may be due to different sensitivities of three strains of mice to brain neurotrophins in response to low-level toluene exposure. However, it is not known the exact mechanism for different sensitivity to toluene-mediated neurotrophin expressions in the hippocampus of three mice strains in the present study. It has been reported that MHC class I protein perform crucial roles outside the immune system, specifically in activity-dependent development and plasticity of the vertebrate nervous system (Huh et al., 2000; Goddard et al., 2007). Currently, our research group showed the strain differences in extracellular amino acid neurotransmitter levels in the hippocampus of H-2 haplotype congenic mice [C57BL/10 (H-2^b) and B10.BR/sg (H-2^k)] in response to toluene exposure (Win-Shwe et al., 2009). Therefore, it may be suggested that H-2 haplotype plays a role in different sensitivity to environmental chemical exposure.

In our present study, using sensitive model of mice, we examined brain NGF level in the hippocampus and found that up-regulation of NGF level in non-allergic mice exposed to 500 ppm toluene whereas in AG mice exposed to 500 ppm toluene. Our findings indicate that toluene exposure at lower level up-regulates the expression of neurotrophins in an allergic mouse model. However, a dose-dependency in adjuvant activity was not found. Some discrepancies were found between high and low dose effect of toluene on NGF expression in mouse hippocampus. No mechanism has been proposed to explain the above stimulatory effect of low-level toluene exposure on NGF expression in the hippocampus. One possibility is that toluene inhalation contributes to neural dysfunction related to sensory stimulation, and this neurotrophin expression and homeostatic mechanisms compensate for toxicant-induced changes. Neurotrophins also take part in neurogenic inflammation by modulating the activity of sensory neurons and enhancing the synthesis and release of neuropeptides (Lindsay and Harmar, 1989). Another possibility is that in OVA-immunized mice, immune system may be activated and may be sensitive to low-level toluene. Furthermore, we

previously reported that OVA immunization acts as a stressor to activate hypothalamo-pituitary adrenal axis (Sari et al., 2004) and then that signal further activates some of immune cells and neural cells in the hippocampus and these neuroimmune interactions promote to up-regulate the neurotrophin expressions. Toluene exposure alone induces NGF synthesis at higher exposure level (500 ppm) and co-exposure of toluene and OVA, like double stressors may induce NGF synthesis at lower exposure level (50 ppm). This finding was consistent with our previous study that showed increased sensitivity of CRH-ir neurons in the hypothalamus and ACTH-ir neurons and mRNA expression in AG mice were higher than those of non-allergic mice in response to toluene (Sari et al., 2006).

We also suggested that the NGF endogenously released after toluene and/or OVA stimulation possibly participates in promoting cell repair and the remodeling of damaged tissue due to acute and chronic stressful events (Aloe et al., 2002). Taken together, brain NGF may represent a link between nervous, endocrine and immune systems and may translate environmental messages into pathophysiological responses. In our previous study, we demonstrated an increased NGF protein level in the hippocampus of C3H/HeN mice following exposure to low levels of formaldehyde (400 ppb) after OVA stimulation (Fujimaki et al., 2004). NGF and BDNF play a key role in the fine tuning of learning and memory performances and in some behavioral processes associated with stress situations (Aloe et al., 2002; Chao et al., 2006). In the immunohistochemical analysis, we observed significantly denser NGF-immunoreactivities in the CA1 and CA3 regions of the hippocampus of C3H/HeN mice in the AG group after exposure to 50 ppm of toluene. This finding is consistent with the expression of NGF mRNA levels in the hippocampus in AG group of C3H/HeN mice. Taken together, our findings strongly suggest that low levels of toluene exposure up-regulated the expression of NGF in the hippocampus at both the mRNA and protein levels in AG C3H/HeN mice.

Currently, neurotrophins and their related receptors have been identified as targets for neurotoxicant. NGF can regulate allergic airway inflammation as well as neural hyper-responsiveness (Bonini et al., 1996; Sanico et al., 1999), indicating that NGF is involved in both neural and immune sensitization. Under physiologic conditions, the local production of neurotrophins is low in both human and animal studies (Nockher and Renz, 2006). Among the neurotrophins, NGF and BDNF play a crucial role in the survival and development of specific peripheral and brain neurons (Chao et al., 2006; Allen and Dawbarn, 2006). They are produced and released by a variety of cells localized in the central and peripheral nervous systems and by cells of the immune and endocrine systems (Levi-Montalcini et al., 1990; Barde, 1994). As mentioned above, the exact mechanism behind the alterations in neurotrophins and their receptors in the mouse hippocampus following toluene exposure is not clear. Using PC12 cells, Bedogni and colleagues demonstrated the dual role of reactive oxygen species (ROS) in survival signaling and in cell death (Bedogni et al., 2003). They reported that ROS rapidly generated in the cell cytosol in response to NGF and Rac-1 have a role in survival signaling to Akt and CREB, whereas harmful mitochondrial oxidants produced at the onset of apoptotic death are reduced by the neurotrophin, likely through the induction of the mitochondrial scavenger manganese-dependent superoxide dismutase. In the present study, oxidative stress marker, the expression of HO-1 mRNA was significantly higher in the hippocampus of C3H/HeN mice exposed to 500 ppm of toluene. There was a tendency of higher proapoptotic Bax mRNA expression in mice exposed to toluene, but not statistically significant. At least in part, our present study showed that the up-regulation of neurotrophins and related receptors in the C3H/HeN mice hippocampus may be occurred to