

REFERENCES

- (1) Cho, A. K., Stefano, E. D., You, Y., Rodriguez, C. E., Schmitz, D. A., Kumagai, Y., Miguel, A. H., Eiguren-Fernandez, A., Kobayashi, T., Avol, E., and Froines, J. R. (2004) Determination of four quinones in diesel exhaust particles, SRM 1649a, and atmospheric PM_{2.5}. *Aerosol Sci. Technol.* 38, 1–14.
- (2) Chen, J., Ehrenhauser, F. S., Valsaraj, K. T., and Wornat, M. J. (2006) Uptake and UV-photooxidation of gas-phase PAHs on the surface of atmospheric water films. 1. Naphthalene. *J. Phys. Chem. A* 110, 9161–9168.
- (3) Sugiyama, K., Wang, T. C., Simpson, J. T., Rodriguez, L., Kador, P. F., and Sato, S. (1999) Aldose reductase catalyzes the oxidation of naphthalene-1, 2-dihydrodiol for the formation of ortho-naphthoquinone. *Drug Metab. Dispos.* 27, 60–67.
- (4) Wilson, A. S., Davis, C. D., Williams, D. P., Buckpitt, A. R., Pirmohamed, M., and Park, B. K. (1996) Characterisation of the toxic metabolite(s) of naphthalene. *Toxicology* 114, 233–242.
- (5) Bolton, J. L., Trush, M. A., Penning, T. M., Dryhurst, G., and Monks, T. J. (2000) Role of quinones in toxicology. *Chem. Res. Toxicol.* 13, 135–160.
- (6) Iwamoto, N., Sumi, D., Ishii, T., Uchida, K., Cho, A. K., Froines, J. R., and Kumagai, Y. (2007) Chemical knockdown of protein-tyrosine phosphatase 1B by 1,2-naphthoquinone through covalent modification causes persistent transactivation of epidermal growth factor receptor. *J. Biol. Chem.* 282, 33396–33404.
- (7) Endo, A., Sumi, D., and Kumagai, Y. (2007) 1,2-Naphthoquinone disrupts the function of cAMP response element-binding protein through covalent modification. *Biochem. Biophys. Res. Commun.* 361, 243–248.
- (8) Saeed, M., Higginbotham, S., Rogan, E., and Cavalieri, E. (2007) Formation of depurinating N3adenine and N7guanine adducts after reaction of 1,2-naphthoquinone or enzyme-activated 1,2-dihydroxy-naphthalene with DNA. Implications for the mechanism of tumor initiation by naphthalene. *Chem.-Biol. Interact.* 165, 175–188.
- (9) Sun, Y., Taguchi, K., Sumi, D., Yamano, S., and Kumagai, Y. (2006) Inhibition of endothelial nitric oxide synthase activity and suppression of endothelium-dependent vasorelaxation by 1,2-naphthoquinone, a component of diesel exhaust particles. *Arch. Toxicol.* 80, 280–285.
- (10) Kikuno, S., Taguchi, K., Iwamoto, N., Yamano, S., Cho, A. K., Froines, J. R., and Kumagai, Y. (2006) 1,2-Naphthoquinone activates vanilloid receptor 1 through increased protein tyrosine phosphorylation, leading to contraction of guinea pig trachea. *Toxicol. Appl. Pharmacol.* 210, 47–54.
- (11) Kumagai, Y., Koide, S., Taguchi, K., Endo, A., Nakai, Y., Yoshikawa, T., and Shimojo, N. (2002) Oxidation of proximal protein sulfhydryls by phenanthraquinone, a component of diesel exhaust particles. *Chem. Res. Toxicol.* 15, 483–489.
- (12) Taguchi, K., Fujii, S., Yamano, S., Cho, A. K., Kamisuki, S., Nakai, Y., Sugawara, F., Froines, J. R., and Kumagai, Y. (2007) An approach to evaluate two-electron reduction of 9,10-phenanthraquinone and redox activity of the hydroquinone associated with oxidative stress. *Free Radical Biol. Med.* 43, 789–799.
- (13) Rees, J. R., and Pirie, A. (1967) Possible reactions of 1,2-naphthoquinone in the eye. *Biochem. J.* 102, 853–863.
- (14) Itoh, K., Ishii, T., Wakabayashi, N., and Yamamoto, M. (1999) Regulatory mechanisms of cellular response to oxidative stress. *Free Radical Res.* 31, 319–324.
- (15) Ishii, T., Itoh, K., Takahashi, S., Sato, H., Yanagawa, T., Katoh, Y., Bannai, S., and Yamamoto, M. (2000) Transcription factor Nrf2 coordinately regulates a group of oxidative stress-inducible genes in macrophages. *J. Biol. Chem.* 275, 16023–16029.
- (16) Itoh, K., Chiba, T., Takahashi, S., Ishii, T., Igarashi, K., Katoh, Y., Oyake, T., Hayashi, N., Satoh, K., Hatayama, I., Yamamoto, M., and Nabeshima, Y. (1997) An Nrf2/small Maf heterodimer mediates the induction of phase II detoxifying enzyme genes through antioxidant response elements. *Biochem. Biophys. Res. Commun.* 236, 313–322.
- (17) Thimmulappa, R. K., Mai, K. H., Srisuma, S., Kensler, T. W., Yamamoto, M., and Biswal, S. (2002) Identification of Nrf2-regulated genes induced by the chemopreventive agent sulforaphane by oligonucleotide microarray. *Cancer Res.* 62, 5196–5203.
- (18) Enomoto, A., Itoh, K., Nagayoshi, E., Haruta, J., Kimura, T., O'Connor, T., Harada, T., and Yamamoto, M. (2001) High sensitivity of Nrf2 knockout mice to acetaminophen hepatotoxicity associated with decreased expression of ARE-regulated drug metabolizing enzymes and antioxidant genes. *Toxicol. Sci.* 59, 169–177.
- (19) Hayashi, A., Suzuki, H., Itoh, K., Yamamoto, M., and Sugiyama, Y. (2003) Transcription factor Nrf2 is required for the constitutive and inducible expression of multidrug resistance-associated protein 1 in mouse embryo fibroblasts. *Biochem. Biophys. Res. Commun.* 310, 824–829.
- (20) O'Brien, P. J. (1991) Molecular mechanisms of quinone cytotoxicity. *Chem.-Biol. Interact.* 80, 1–41.
- (21) Monks, T. J., Hanzlik, R. P., Cohen, G. M., Ross, D., and Graham, D. G. (1992) Quinone chemistry and toxicity. *Toxicol. Appl. Pharmacol.* 112, 2–16.
- (22) Taguchi, K., Shimada, M., Fujii, S., Sumi, D., Pan, X., Yamano, S., Nishiyama, T., Hiratsuka, A., Yamamoto, M., Cho, A. K., Froines, J. R., and Kumagai, Y. (2008) Redox cycling of 9,10-phenanthraquinone to cause oxidative stress is terminated through its monoglucuronide conjugation in human pulmonary epithelial A549 cells. *Free Radical Biol. Med.* 44, 1645–1655.
- (23) Miura, T., and Kumagai, Y. (2010) Immunochemical method to detect proteins that undergo selective modification by 1,2-naphthoquinone derived from naphthalene through metabolic activation. *J. Toxicol. Sci.* 35, 843–852.
- (24) Toyama, T., Sumi, D., Shinkai, Y., Yasutake, A., Taguchi, K., Tong, K. L., Yamamoto, M., and Kumagai, Y. (2007) Cytoprotective role of Nrf2/Keap1 system in methylmercury toxicity. *Biochem. Biophys. Res. Commun.* 363, 645–650.
- (25) Iwamoto, N., Nishiyama, A., Eiguren-Fernandez, A., Hinds, W., Kumagai, Y., Froines, J., Cho, A., and Shinyashiki, M. (2010) Biochemical and cellular effects of electrophiles present in ambient air samples. *Atmos. Environ.* 44, 1483–1489.
- (26) Okawa, H., Motohashi, H., Kobayashi, A., Aburatani, H., Kensler, T. W., and Yamamoto, M. (2006) Hepatocyte-specific deletion of the keap1 gene activates Nrf2 and confers potent resistance against acute drug toxicity. *Biochem. Biophys. Res. Commun.* 339, 79–88.
- (27) Shinkai, Y., Sumi, D., Toyama, T., Kaji, T., and Kumagai, Y. (2009) Role of aquaporin 9 in cellular accumulation of arsenic and its cytotoxicity in primary mouse hepatocytes. *Toxicol. Appl. Pharmacol.* 237, 232–236.
- (28) Laemmli, U. K. (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227, 680–685.
- (29) Kyhse-Andersen, J. (1984) Electrophoretic transfer of multiple gels: a simple apparatus without buffer tank for rapid transfer of proteins from polyacrylamide to nitrocellulose. *J. Biochem. Biophys. Methods* 10, 203–209.
- (30) Denizot, F., and Lang, R. (1986) Rapid colorimetric assay for cell growth and survival. Modifications to the tetrazolium dye procedure giving improved sensitivity and reliability. *J. Immunol. Methods* 89, 271–277.
- (31) Dinkova-Kostova, A. T., Holtzclaw, W. D., Cole, R. N., Itoh, K., Wakabayashi, N., Katoh, Y., Yamamoto, M., and Talalay, P. (2002) Direct evidence that sulfhydryl groups of Keap1 are the sensors regulating induction of phase 2 enzymes that protect against carcinogens and oxidants. *Proc. Natl. Acad. Sci. U.S.A.* 99, 11908–11913.
- (32) Wakabayashi, N., Dinkova-Kostova, A. T., Holtzclaw, W. D., Kang, M. I., Kobayashi, A., Yamamoto, M., Kensler, T. W., and Talalay, P. (2004) Protection against electrophile and oxidant stress by induction of the phase 2 response: fate of cysteines of the Keap1 sensor modified by inducers. *Proc. Natl. Acad. Sci. U.S.A.* 101, 2040–2045.
- (33) Chu, X. Y., Huskey, S. E., Braun, M. P., Sarkadi, B., Evans, D. C., and Evers, R. (2004) Transport of ethinylestradiol glucuronide and ethinylestradiol sulfate by the multidrug resistance proteins MRP1, MRP2, and MRP3. *J. Pharmacol. Exp. Ther.* 309, 156–164.

(34) Donato, M. T., Montero, S., Castell, J. V., Gomez-Lechon, M. J., and Lahoz, A. (2010) Validated assay for studying activity profiles of human liver UGTs after drug exposure: inhibition and induction studies. *Anal. Bioanal. Chem.* 396, 2251–2263.

(35) Morrow, C. S., Peklak-Scott, C., Bishwokarma, B., Kute, T. E., Smitherman, P. K., and Townsend, A. J. (2006) Multidrug resistance protein 1 (MRP1, ABCC1) mediates resistance to mitoxantrone via glutathione-dependent drug efflux. *Mol. Pharmacol.* 69, 1499–1505.

(36) Luo, Y., Eggler, A. L., Liu, D., Liu, G., Mesecar, A. D., and van Breemen, R. B. (2007) Sites of alkylation of human Keap1 by natural chemoprevention agents. *J. Am. Soc. Mass Spectrom.* 18, 2226–2232.

(37) Fujii, S., Sawa, T., Ihara, H., Tong, K. I., Ida, T., Okamoto, T., Ahtesham, A. K., Ishima, Y., Motohashi, H., Yamamoto, M., and Akaike, T. (2010) The critical role of nitric oxide signaling, via protein S-guanylation and nitrated cyclic GMP, in the antioxidant adaptive response. *J. Biol. Chem.* 285, 23970–23984.

(38) Kobayashi, M., Li, L., Iwamoto, N., Nakajima-Takagi, Y., Kaneko, H., Nakayama, Y., Eguchi, M., Wada, Y., Kumagai, Y., and Yamamoto, M. (2009) The antioxidant defense system Keap1-Nrf2 comprises a multiple sensing mechanism for responding to a wide range of chemical compounds. *Mol. Cell. Biol.* 29, 493–502.

(39) Sekhar, K. R., Rachakonda, G., and Freeman, M. L. (2010) Cysteine-based regulation of the CUL3 adaptor protein Keap1. *Toxicol. Appl. Pharmacol.* 244, 21–26.

(40) Okada, K., Shoda, J., Taguchi, K., Maher, J. M., Ishizaki, K., Inoue, Y., Ohtsuki, M., Goto, N., Takeda, K., Utsunomiya, H., Oda, K., Warabi, E., Ishii, T., Osaka, K., Hyodo, I., and Yamamoto, M. (2008) Ursodeoxycholic acid stimulates Nrf2-mediated hepatocellular transport, detoxification, and antioxidative stress systems in mice. *Am. J. Physiol. Gastrointest. Liver Physiol.* 295, G735–747.

(41) Taguchi, K., Maher, J. M., Suzuki, T., Kawatani, Y., Motohashi, H., and Yamamoto, M. (2010) Genetic analysis of cytoprotective functions supported by graded expression of Keap1. *Mol. Cell. Biol.* 30, 3016–3026.

(42) Kumagai, Y., Arimoto, T., Shinyashiki, M., Shimojo, N., Nakai, Y., Yoshikawa, T., and Sagai, M. (1997) Generation of reactive oxygen species during interaction of diesel exhaust particle components with NADPH-cytochrome P450 reductase and involvement of the bioactivation in the DNA damage. *Free Radical Biol. Med.* 22, 479–487.

(43) Brown, P. C., Dulik, D. M., and Jones, T. W. (1991) The toxicity of menadione (2-methyl-1,4-naphthoquinone) and two thioether conjugates studied with isolated renal epithelial cells. *Arch. Biochem. Biophys.* 285, 187–196.

(44) Buffinton, G. D., Ollinger, K., Brunmark, A., and Cadenas, E. (1989) DT-diaphorase-catalysed reduction of 1,4-naphthoquinone derivatives and glutathionyl-quinone conjugates. Effect of substituents on autoxidation rates. *Biochem. J.* 257, 561–571.

(45) Nishiyama, T., Izawa, T., Usami, M., Ohnuma, T., Ogura, K., and Hiratsuka, A. (2010) Cooperation of NAD(P)H:quinone oxidoreductase 1 and UDP-glucuronosyltransferases reduces menadione cytotoxicity in HEK293 cells. *Biochem. Biophys. Res. Commun.* 394, 459–463.

