

Figure 2. Suppression by phosphodiesterase (PDE) 3 inhibition of agonist-induced Ca²⁺ influx mediated by diacylglycerol (DAG)-activated transient receptor potential canonical (TRPC) channels. A to D, Average time courses of ATP-stimulated changes in intracellular Ca²⁺ concentration ([Ca²⁺]_i) determined by Fura-2 ratio (Ratio) in vector-, TRPC3-, TRPC6-, and TRPC7-overexpressing HEK293 cells. Ca²⁺ release-mediated [Ca²⁺]_i increase was first induced by ATP (100 μmol/L) in Ca²⁺-free external solution, and then Ca²⁺ influx-mediated [Ca²⁺]_i increase was induced by the addition of Ca²⁺ (2 mmol/L). Cells were pretreated with the indicated concentration of cilostazol (CLZ) for 30 minutes before ATP stimulation. E and F, Peak changes in intracellular Ca²⁺ concentration induced by ATP in the absence (E) or presence (F) of extracellular Ca²⁺. n = 48 to 74 cells.

was also abolished by cotreatment with KT5720, a PKA inhibitor, but not by KT5823 (a PKG inhibitor) or Go6976 (a protein kinase C inhibitor) (Supplemental Figure VII). In addition, treatment with 8-Br-cAMP and forskolin significantly suppressed Ca²⁺ influx-mediated [Ca²⁺]_i increases induced by OAG stimulation in TRPC6-wt-expressing cells, whereas both PKA activators did not suppress OAG-induced [Ca²⁺]_i increases in TRPC6-Thr69Ala-expressing cells (Supplemental Figure VI). Furthermore, stimulation of muscarinic receptor with carbachol induced inward currents in TRPC6-wt-expressing cells; these currents were completely suppressed by the pretreatment with cilostazol (Figure 3A to 3C). The suppression of TRPC6-induced currents by cilostazol was abolished in cells expressing TRPC6-Thr69Ala (Figure 3D). These results suggest that PKA-dependent phosphorylation of TRPC6 at Thr69 underlies inhibition of TRPC6 channel activity by PDE3 inhibition.

PKA-Dependent Phosphorylation of TRPC6 by PDE3 Inhibition

We have recently generated an anti-phospho-TRPC6 (Thr69) antibody³⁷ and found that activation of PKA by the treatment with forskolin and cilostazol dramatically increased the phosphorylation of TRPC6 at Thr69 in TRPC6-wt-expressing cells, which was not observed in TRPC6-Thr69Ala-expressing cells (Figure 3E). This result suggests the specific recognition of Thr69 phosphorylation by this antibody. Treatment of HEK293 cells with cilostazol increased TRPC6 phosphorylation levels ≈1.5-fold, which was completely suppressed by cotreatment with KT5720. Treatment with cilostazol gradually increased TRPC6 phosphorylation levels (Figure 3F). The time course of this phosphorylation was consistent with the time of pretreatment required to obtain the inhibition of TRPC6-mediated Ca²⁺ influx with cilostazol (Supplemental Figure IV). Although cilostazol completely suppressed TRPC6-mediated Ca²⁺ influx, as in the case of forskolin (Supplemental Figure VI), the extent of TRPC6 phosphorylation by cilostazol was much weaker than that by forskolin. However, phosphorylation levels of membrane-localized TRPC6 proteins induced by cilostazol were similar to those induced by forskolin (Figure 3G). These results suggest that phosphorylation of plasma membrane-localized TRPC6 proteins is essential for suppression of TRPC6 channel activity by PDE3 inhibition.

Phosphorylation of TRPC6 at Thr69 Is Essential for Suppression of Ang II-Induced Ca²⁺ Influx by PDE3 Inhibition in RAoSMCs

We next examined which endogenous TRPC subtype(s) is involved in Ang II-induced Ca²⁺ influx in RAoSMCs. Although Ang II-induced Ca²⁺ release was not suppressed by the knockdown of TRPC3/6/7, the Ca²⁺-influx-mediated Ca²⁺ response induced by Ang II was suppressed by knockdown of TRPC6 but not of TRPC3 and TRPC7 (Figure 4A and 4B). Pretreatment with cilostazol significantly reduced Ca²⁺ influx-mediated Ca²⁺ responses induced by Ang II, which was abolished by TRPC6 knockdown. We found that TRPC3 and TRPC6 proteins were actually expressed in RAoSMCs, and expression levels of TRPC proteins were actually suppressed by siRNA treatment (Figure 4C). Human TRPC3 has 2 sites (Thr11 and Ser263) phosphorylated by PKG or PKA, and Ser263 (position 2), but not Thr11 (position 1), is conserved in rodent TRPC3/6/7 proteins (Supplemental Figure VI). As the expression of mouse TRPC3 mutant (Ser251Ala) abolished the suppression of OAG-induced Ca²⁺ response by cilostazol, position 2 may be involved in suppression of TRPC3 channel activity by PDE3 inhibition. However, pretreatment of RAoSMCs with cilostazol actually induced phosphorylation of TRPC6 at Thr69 (position 1) but did not induce phosphorylation of TRPC3/6/7 at Ser (position 2), although forskolin cotreatment induced phosphorylation of these sites (Figure 4D). These results suggest that PDE3 inhibition suppresses Ang II-induced Ca²⁺ influx via local PKA-mediated phosphorylation of TRPC6 at Thr69 in RAoSMCs.




Figure 3. Requirement of phosphorylation of transient receptor potential (TRPC) 6 at Thr69 in suppression of TRPC6 currents by phosphodiesterase (PDE) 3 inhibition. **A** and **B**, Typical traces of inward cation currents induced by carbachol (CCh; 100 μ mol/L) without (**A**) or with (**B**) cilostazol (CLZ) pretreatment. Cells were pretreated with CLZ (5 μ mol/L) for 30 minutes before addition of CCh. WT indicates wild-type. **C**, Average TRPC6-induced current density at the holding potential of -50 mV with or without CLZ pretreatment in TRPC6-wt- and TRPC6-Thr69Ala-expressing cells. Numbers of experiments are shown in the columns. n.s. indicates not significant. **D**, Current-voltage relationship of wild-type and Thr69Ala-mutated TRPC6 channels with or without pretreatment with CLZ. To construct the latter, rising voltage ramps (-100 to $+100$ mV, 2 s) were applied. **E**, Effects of forskolin (FSK) and KT5720 on CLZ-induced TRPC6 phosphorylation in TRPC6-wt- and TRPC6-Thr69Ala-overexpressing cells. Cells were treated with CLZ (5 μ mol/L), forskolin (10 μ mol/L), and KT5720 (0.1 μ mol/L) for 30 minutes. **F**, Time courses of TRPC6 phosphorylation induced by CLZ (5 μ mol/L) in TRPC6-expressing HEK293 cells. **G**, Effects of forskolin on CLZ-induced phosphorylation of TRPC6 proteins expressed in the plasma membrane. $n=3$ experiments. * $P<0.05$, ** $P<0.01$ vs control (-), ### $P<0.01$ vs CLZ.

Formation of the TRPC-PKA-PDE3 Signal Complex in the Plasma Membrane

We further examined whether DAG-activated TRPC channels functionally interact with PKA and PDE3. TRPC3-GFP, TRPC6-GFP, and farnesylation signal-encoding GFP proteins are predominantly expressed in crude membrane fraction of HEK293 cells, whereas PKA is predominantly expressed in cytosol. Immunoprecipitation of GFP-containing proteins revealed that endogenous PDE3A could associate with TRPC3-GFP and TRPC6-GFP but not with farnesylation

signal-encoding GFP proteins. In addition, PKA catalytic subunit also associated with TRPC3-GFP and TRPC6-GFP (Figure 5A). Treatment with cilostazol actually induced phosphorylation of TRPC3/6/7 at Ser (position 2) (Figure 5B). These results suggest that TRPC3/6/7 could form a signal complex with PDE3 and PKA beneath the plasma membrane in HEK293 cells. However, endogenous TRPC6 associated with PDE3A and PKA more potently than TRPC3 in RAoSMCs (Figure 5C), suggesting that endogenous TRPC6 preferentially forms a ternary complex with PDE3 and PKA in RAoSMCs.




Figure 4. Requirement of transient receptor potential canonical (TRPC) 6 phosphorylation in the suppression of angiotensin II (Ang II)-induced Ca²⁺ influx by phosphodiesterase (PDE) 3 inhibition in rat aortic smooth muscle cells (RAoSMCs). A and B, Average time courses of Ca²⁺ responses (A) and peak [Ca²⁺]_i increases induced by Ang II (100 nmol/L) (B) in the presence or absence of cilostazol (CLZ) (1 μmol/L) in RAoSMCs treated with small interfering (si) RNAs for control or TRPC3, TRPC6, or TRPC7. n=24 to 38 cells. C, Expression levels of native TRPC proteins in respective TRPC-knocked-down RAoSMCs. D, Phosphorylation of TRPC6 but not TRPC3 and TRPC7 by CLZ (10 μmol/L) with or without forskolin (FSK) (30 μmol/L) for 1 hour. n=3 experiments.

Requirement of TRPC6 Phosphorylation for Cilostazol-Induced Suppression of Vasoconstriction Induced by Ang II in VSMC

To examine the functional relevance of TRPC6 phosphorylation at Thr69 to the vasodilating effect of cilostazol, we examined the effects of TRPC6-Thr69Ala in RAoSMCs. Treatment of RAoSMCs with Ang II induced a transient

increase in [Ca²⁺]_i, which was significantly suppressed by pretreatment with cilostazol (Figure 6A and 6B). Overexpression of TRPC6-wt slightly enhanced the Ang II-induced Ca²⁺ response, which was also suppressed by cilostazol. In contrast, overexpression of TRPC6-Thr69Ala abolished the suppression of Ang II-induced Ca²⁺ response by cilostazol pretreatment. We next examined the contractile response




Figure 5. Association of transient receptor potential canonical (TRPC) channels with protein kinase A (PKA) and phosphodiesterase (PDE) 3 proteins. **A**, Interaction of TRPC proteins with PKA catalytic subunit and PDE3A in vector-, TRPC3-, and TRPC6-overexpressing HEK293 cells. **B**, Phosphorylation of TRPC proteins by the treatment with cilostazol (CLZ) (10 μ mol/L) for 1 hour in HEK293 cells. **C**, Association of endogenous TRPC6 with PKA and PDE3 in rat aortic smooth muscle cells (RAoSMCs). $n=3$ experiments. GFP indicates green fluorescent protein; IP, immunoprecipitation; WB, Western blotting.

of the aortic rings reconstituted from RAoSMCs. Treatment of reconstituted aortic ring with Ang II induced contraction, which was significantly attenuated by pretreatment with cilostazol in the vector (GFP)-expressing rings (Figure 6C and 6D). In contrast, the expression of TRPC6-Thr69Ala abolished the suppression of Ang II-induced contraction by cilostazol pretreatment. A dominant-negative mutant of TRPC6 (TRPC6-3A) also abolished the cilostazol-induced suppression of Ang II-induced contraction (Supplemental Figure VIII). These results suggest that the phosphorylation of TRPC6 at Thr69 is required for the suppression of Ang II-induced contraction by cilostazol in RAoSMCs.

Discussion

In excitable cells, one of the main functions of receptor-activated TRPC channels is to induce membrane depolarization through sustained cation influx by receptor stimulation.^{23,34} The membrane depolarization subsequently induces activation of VDCCs, leading to activation of Ca^{2+} -dependent signaling required for muscle contraction. We revealed that inhibition of ROCCs and VDCCs significantly reduced the Ang II-induced contraction of rat thoracic aorta (Figure 1), suggesting that the VDCC-mediated Ca^{2+} influx via TRPC-mediated membrane depolarization is involved in Ang II-induced contraction of rat thoracic aorta. In contrast, as the magnitude of the suppression by cilostazol of phenylephrine- and U46619-induced vasoconstriction was smaller than that of Ang II-induced vasoconstriction, and U46619-induced contraction of reconstituted rings was weakly (not significantly) suppressed by PDE3 inhibition (Supplemental Figure VIII). These results suggest that Ang II preferentially induces DAG-mediated Ca^{2+} influx in aortic VSMCs.

We have shown the critical involvement of TRPC6 channels in Ang II-induced vasoconstriction through analyzing the mechanism of vasodilation by PDE3 inhibition. The PDE3 inhibition specifically suppresses DAG-mediated (but not IP_3 -mediated and store-operated) Ca^{2+} influx through PKA-dependent phosphorylation of TRPC3/6/7. TRPC proteins are known to function not only as a Ca^{2+} -permeable channel-forming subunit but also as a scaffold protein to form the Ca^{2+} signaling complex.³⁸ For example, TRPC3 channels are reported to associate with several signaling molecules, such as IP_3 receptors, calmodulin, vesicle-associated membrane protein 2, phospholipase C- γ , and protein kinase C.^{9,31} We found that exogenous TRPC3 and TRPC6 proteins could associate with PKA and PDE3 in HEK293 cells (Figure 5). However, endogenous TRPC6 preferentially forms a ternary complex with PKA and PDE3 rather than TRPC3 in RAoSMCs (Figure 5). This makes it possible to induce local PKA-mediated phosphorylation of TRPC6 proteins by PDE3 inhibition. Although the position 1 phosphorylation site of human TRPC3 is not conserved in mouse TRPC3, PDE3 inhibition actually induced phosphorylation of TRPC3 at Ser251 and of TRPC7 at Ser267 (Figure 5). As the inhibition of TRPC3 phosphorylation canceled suppression of TRPC3-mediated Ca^{2+} influx by PDE3 inhibition (Supplemental Figure VI), our results indicate that the PKG phosphorylation sites of DAG-activated TRPC channels are also targets of PKA.

It has been thought that phosphorylation of MLCK is essential for the PKA-mediated vasodilating action.³⁹ However, the present results suggest that PKA induces vasodilation through 2 mechanisms: one is to suppress agonist-induced Ca^{2+} influx through inhibition of DAG-activated TRPC channels, and the other is to inhibit MLCK activity by phosphorylation of MLCK. As PDE3 inhibition did not suppress the KCl-induced contraction, PKA-mediated inhibition of MLCK activity may not participate in the vasodilating effects of PDE3 inhibition. In addition, Rho functions as an essential mediator in agonist-induced vasoconstriction.^{33,39} Rho decreases myosin light chain phosphatase activity through Rho kinase-mediated phosphorylation. It has been reported that TRPC6-mediated Ca^{2+} influx induces RhoA




Figure 6. Requirement of transient receptor potential canonical (TRPC) 6 phosphorylation in the suppression of angiotensin II (Ang II)-induced aortic contraction by phosphodiesterase (PDE) 3 inhibition. **A**, Effects of cilostazol (CLZ) on Ang II-induced Ca^{2+} responses in primary cultured rat aortic smooth muscle cells (RAoSMCs) expressing vector (pcDNA3), TRPC6-wild-type (wt), or TRPC6-Thr69Ala. Cells were treated with CLZ for 30 minutes before addition of Ang II (100 nmol/L). **B**, Peak increase in $[Ca^{2+}]_i$ induced by Ang II (100 nmol/L) in the presence or absence of CLZ (10 μ mol/L). $n=40$ to 80 cells. **C**, Representative time courses of changes in tensions of aortic ring reconstituted from RAoSMCs expressing vector, TRPC6-wt, TRPC6-Thr69Ala, or dominant-negative mutant of TRPC6 (TRPC6-3A). CLZ (10 μ mol/L) was treated for 30 minutes before stimulation with Ang II (100 nmol/L). **D**, Peak contraction induced by Ang II in the absence (-) or presence (+) of CLZ. * $P<0.05$, ** $P<0.01$ vs (-) CLZ, ## $P<0.01$ vs C6-wild-type (WT) (-) CLZ.

activation in thrombin-stimulated endothelial cells.⁴⁰ However, PDE3 inhibition did not attenuate agonist-induced Rho activation in RAoSMCs (Supplemental Figure VIII). Thus, PDE3 inhibition may not affect the Rho-mediated signaling pathway.

Cilostazol has many pharmacological actions, including vasodilation, inhibition of platelet activation and aggregation, inhibition of thrombosis, increased blood flow to the limbs, improvement in serum lipids with lowering of triglycerides and elevation of high-density lipoprotein cholesterol, and inhibition of growth of VSMCs.^{10–12} Nakamura et al have previously reported the PKA-independent vasodilating effect of PDE3 inhibition using pressurized rabbit cerebral penetrating arterioles.⁴¹ It is expected that the effectiveness of vasodilation by PDE3 inhibition is greatly dependent on the kinds of blood vessel sampled, the agonist stimulation, and the timing of drug treatment. Thus, our present study is only to show a novel mechanism underlying PKA-dependent vasodilation by PDE3 inhibition. Cilostazol is clinically applicable to the patients with intermittent claudication and arteriosclerosis obliterata.^{10–12} It is generally thought that endothelium-dependent vasodilation and capillary-like for-

mation underlies protection of PDE3 inhibition against peripheral arterial disease. Although the relationship between TRPC channels and peripheral arterial disease is still unknown, our findings will provide a novel mechanism underlying endothelium-independent improvement of peripheral blood flow by PDE3 inhibition.

In conclusion, we demonstrated that PDE3 inhibition specifically suppresses DAG-mediated Ca^{2+} influx through PKA-dependent phosphorylation of TRPC3/6/7 channels in HEK293 cells. Furthermore, DAG-activated TRPC6 channels were found to mediate the Ang II-induced contraction of aortic VSMCs, and PDE3 inhibition preferentially induced phosphorylation of rodent TRPC6 at Thr69. Formation of TRPC6/PDE3/PKA ternary complex may be involved in this mechanism. These findings will indicate the novel mechanism underlying the vasorelaxant effects of PDE3 inhibition against vasoactive hormones.

Sources of Funding

This study was supported by grants from the Ministry of Education, Culture, Sports, Science, and Technology of Japan (to M. Nishida,

M. Nakaya, and H.K.); from the Naito and Mochida Memorial Foundation (to M. Nishida); and from the Vehicle Racing Commemorative Foundation (to T.I. and R.I.) and by a Grant-in-Aid for Scientific Research on Innovative Areas (to M. Nishida) and a Grant-in-Aid for Scientific Research on Priority Areas (H.K.).

Disclosures

None.

References

- Johnson PC. The myogenic response and the microcirculation. *Microvasc Res.* 1977;13:1–18.
- Akata T. Cellular and molecular mechanisms regulating vascular tone: part 1: basic mechanisms controlling cytosolic Ca^{2+} concentration and the Ca^{2+} -dependent regulation of vascular tone. *J Anesth.* 2007;21:220–231.
- Berridge MJ. Inositol trisphosphate and calcium signalling. *Nature.* 1993;361:315–325.
- Putney JW Jr. Capacitative calcium entry revisited. *Cell Calcium* 1990; 11:611–624.
- Leung FP, Yung LM, Yao X, Laher I, Huang Y. Store-operated calcium entry in vascular smooth muscle. *Brit J Pharmacol.* 2008;153:846–857.
- Nishizuka Y. Protein kinase C and lipid signaling for sustained cellular responses. *FASEB J.* 1995;9:484–496.
- Inoue R, Okada T, Onoue H, Hara Y, Shimizu S, Naitoh S, Ito Y, Mori Y. The transient receptor potential protein homologue TRP6 is the essential component of vascular α 1-adrenoceptor-activated Ca^{2+} -permeable cation channel. *Circ Res.* 2001;88:325–332.
- Welsh DG, Morielli AD, Nelson MT, Brayden JE. Transient receptor potential channels regulate myogenic tone of resistance arteries. *Circ Res.* 2002;90:248–250.
- Dietrich A, Mederos Y, Schnitzler M, Gollasch M, Gross V, Storch U, Dubrovskaya G, Obst M, Yildirim E, Salanova B, Kalwa H, Essin K, Pinkenburg O, Luft FC, Gudermann T, Birnbaumer L. Increased vascular smooth muscle contractility in TRPC6^{-/-} mice. *Mol Cell Biol.* 2005;25:6980–6989.
- Kimura Y, Tani T, Kanbe T, Watanabe K. Effect of cilostazol on platelet aggregation and experimental thrombosis. *Arzneimittelforschung.* 1985; 35(7A):1144–1149.
- Kambayashi J, Liu Y, Sun B, Shakur Y, Yoshitake M, Czerwiec F. Cilostazol as a unique antithrombotic agent. *Curr Pharm Des.* 2003;9: 2289–2302.
- Dawson DL, Cutler BS, Meissner MH, Strandness DE, Jr. Cilostazol has beneficial effects in treatment of intermittent claudication: results from a multicenter, randomized, prospective, double-blind trial. *Circulation.* 1998;98:678–686.
- Tanaka T, Ishikawa T, Hagiwara M, Onoda K, Itoh H, Hidaka H. Effects of cilostazol, a selective cAMP phosphodiesterase inhibitor on the contraction of vascular smooth muscle. *Pharmacology.* 1988;36:313–320.
- Shiraishi Y, Kanmura Y, Itoh T. Effect of cilostazol, a phosphodiesterase type III inhibitor, on histamine-induced increase in $[Ca^{2+}]_i$ and force in middle cerebral artery of the rabbit. *Brit J Pharmacol.* 1998;123: 869–878.
- Adelstein RS, Conti MA, Hathaway DR, Klee CB. Phosphorylation of smooth muscle myosin light chain kinase by the catalytic subunit of adenosine 3': 5'-monophosphate-dependent protein kinase. *J Biol Chem.* 1978;253:8347–8350.
- Huang J, Zhou H, Mahavadi S, Sriwai W, Murthy KS. Inhibition of $G\alpha_q$ -dependent PLC- β 1 activity by PKG and PKA is mediated by phosphorylation of RGS4 and GRK2. *Am J Physiol Cell Physiol.* 2007;292: C200–C208.
- Heximer SP, Knutsen RH, Sun X, Kaltenbronn KM, Rhee MH, Peng N, Oliveira-dos-Santos A, Penninger JM, Muslin AJ, Steinberg TH, Wyss JM, Mecham RP, Blumer KJ. Hypertension and prolonged vasoconstrictor signaling in RGS2-deficient mice. *J Clin Invest.* 2003;111: 445–452.
- Harris DM, Cohn HI, Pesant S, Eckhart AD. GPCR signalling in hypertension: role of GRKs. *Clin Sci (Lond).* 2008;115:79–89.
- Montell C, Rubin GM. Molecular characterization of the *Drosophila* *trp* locus: a putative integral membrane protein required for phototransduction. *Neuron.* 1989;2:1313–1323.
- Nishida M, Hara Y, Yoshida T, Inoue R, Mori Y. TRP channels: molecular diversity and physiological function. *Microcirculation.* 2006; 13:535–550.
- Hofmann T, Schaefer M, Schultz G, Gudermann T. Subunit composition of mammalian transient receptor potential channels in living cells. *Proc Natl Acad Sci U S A.* 2002;99:7461–7466.
- Lucas P, Ukhanov K, Leinders-Zufall T, Zufall F. A diacylglycerol-gated cation channel in vomeronasal neuron dendrites is impaired in TRPC2 mutant mice: mechanism of pheromone transduction. *Neuron.* 2003;40: 551–561.
- Onohara N, Nishida M, Inoue R, Kobayashi H, Sumimoto H, Sato Y, Mori Y, Nagao T, Kurose H. TRPC3 and TRPC6 are essential for angiotensin II-induced cardiac hypertrophy. *EMBO J.* 2006;25: 5305–5316.
- Poburko D, Liao C-H, Lemos VS, Lin E, Maruyama Y, Cole WC, van Breemen C. Transient receptor potential channel 6-mediated, localized cytosolic $[Na^+]_i$ transients drive Na^+/Ca^{2+} exchanger-mediated Ca^{2+} entry in purinergically stimulated aorta smooth muscle cells. *Circ Res.* 2007;101:1030–1038.
- Iwamoto T, Kita S, Zhang J, Blaustein MP, Arai Y, Yoshida S, Wakimoto K, Komuro I, Katsuragi T. Salt-sensitive hypertension is triggered by Ca^{2+} entry via Na^+/Ca^{2+} exchanger type-1 in vascular smooth muscle. *Nat Med.* 2004;10:1193–1199.
- Thebault S, Flourakis M, Vanoverberghe K, Vandermoere F, Roudbaraki M, Lehen'kyi V, Slomianny C, Beck B, Mariot P, Bonnal JL, Mauroy B, Shuba Y, Capiod T, Skryma R, Prevarskaya N. Differential role of transient receptor potential channels in Ca^{2+} entry and proliferation of prostate cancer epithelial cells. *Cancer Res.* 2006;66:2038–2047.
- Trebak M, Hempel N, Wedel BJ, Smyth JT, Bird GS, Putney JW, Jr. Negative regulation of TRPC3 channels by protein kinase C-mediated phosphorylation of serine 712. *Mol Pharmacol.* 2005;67:558–563.
- Kwan HY, Huang Y, Yao X. Protein kinase C can inhibit TRPC3 channels indirectly via stimulating protein kinase G. *J Cell Physiol.* 2006;207:315–321.
- Kwan HY, Huang Y, Yao X. Regulation of canonical transient receptor potential isoform 3 (TRPC3) channel by protein kinase G. *Proc Natl Acad Sci U S A.* 2004;101:2625–2630.
- Takahashi S, Lin H, Geshi N, Mori Y, Kawarabayashi Y, Takami N, Mori MX, Honda A, Inoue R. Nitric oxide-cGMP-protein kinase G pathway negatively regulates vascular transient receptor potential channel TRPC6. *J Physiol.* 2008;586:4209–4223.
- Nishida M, Sugimoto K, Hara Y, Mori E, Morii T, Kurosaki T, Mori Y. Amplification of receptor signalling by Ca^{2+} entry-mediated translocation and activation of PLC γ 2 in B lymphocytes. *EMBO J.* 2003;22: 4677–4688.
- Nishida M, Sato Y, Uemura A, Narita Y, Tozaki-Saitoh H, Nakaya M, Ide T, Suzuki K, Inoue K, Nagao T, and Kurose H. P2Y₆ Receptor- $G\alpha_{12/13}$ signaling in cardiomyocytes triggers pressure overload-induced cardiac fibrosis. *EMBO J.* 2008;27:3104–3115.
- Bi D, Nishimura J, Niuro N, Hirano K, Kanaide H. Contractile properties of the cultured vascular smooth muscle cells: the crucial role played by RhoA in the regulation of contractility. *Circ Res.* 2005;96:890–897.
- Large RA. Receptor-operated Ca^{2+} -permeable nonselective cation channels in vascular smooth muscle: a physiologic perspective. *J Cardiovasc Electrophysiol.* 2002;13:493–501.
- Inoue R, Jensen LJ, Shi J, Morita H, Nishida M, Honda A, Ito Y. Transient receptor potential channels in cardiovascular function and disease. *Circ Res.* 2006;99:119–131.
- Schwede F, Maronde E, Genieser H, Jastorff B. Cyclic nucleotide analogs as biochemical tools and prospective drugs. *Pharmacol Ther.* 2000;87: 199–226.
- Nishida M, Watanabe K, Sato Y, Nakaya M, Kitajima K, Ide T, Inoue R, and Kurose H. Phosphorylation of TRPC6 channels at Thr⁶⁹ is required for anti-hypertrophic effects of phosphodiesterase 5 inhibition. *J Biol Chem.* 2010;285:13244–13253.
- Ambudkar IS. Ca^{2+} signaling microdomains: platforms for the assembly and regulation of TRPC channels. *Trends Pharmacol Sci.* 2006;27:25–32.
- Hirano K, Hirano M, Kanaide H. Regulation of myosin phosphorylation and myofibrillar Ca^{2+} sensitivity in vascular smooth muscle. *J Smooth Muscle Res.* 2004;40:219–236.
- Singh I, Knezevic N, Ahmed GU, Kini V, Malik AB, Mehta D. $G\alpha_q$ -TRPC6-mediated Ca^{2+} entry induces RhoA activation and resultant endothelial cell shape change in response to thrombin. *J Biol Chem.* 2007;282:7833–7843.
- Nakamura K, Ikomi F, Ohhashi T. Cilostazol, an inhibitor of type 3 phosphodiesterase, produces endothelium-independent vasodilation in pressurized rabbit cerebral penetrating arterioles. *J Vasc Res.* 2006;43: 86–94.

Supplementary Materials and Methods

Animals, Materials and Antibodies

All experiments on male Sprague-Dawley rats (240-330 g) were performed in accordance with the Guide for the Care and Use of Laboratory Animals prepared by Kyushu University. Poly-L-lysine, cilostazol, 1-oleoyl-2-acetyl-*sn*-glycerol (OAG), adenosine triphosphate (ATP), phenylephrine (α 1 adrenergic receptor agonist), 9,11-dideoxy-11 α ,9 α -epoxymethano-prostaglandin F_{2 α} (U46619; thromboxane A₂ analogue), cilostazol, milrinone, ionomycin, SK&F96365, 8-Bromo-cAMP, forskolin, KT5823, KT5720, and ascorbic acid were purchased from Sigma. Go6976 and KB-R7943 were from Carbiochem. Fugene 6 was from Roche. Pentobarbital was from Dainippon-Sumitomo Pharm. Fura2-AM was from Dojindo. Ang II was from Peptide Inst. CV11974 was provided from Takeda Chemical Industry. EGTA was from Nacalai. Cellmatrix type I-P was from Nitta Gelatin. Neutralized formaldehyde was from Wako. The cDNAs coding mouse TRPC6 and TRPC6-T69A mutant were constructed as described (1, 2). Anti-GFP antibody was from Abcam, anti-TRPC6 antibody was from Alomone, and protein A sepharose beads were from GE Healthcare. Phospho-Thr⁶⁹ TRPC6 antiserum and phospho-Ser TRPC3/6/7 antiserum were generated against phospho-TRPC6 peptide (CHRRQ(p)TILREK) (3) and phospho-TRPC3/6/7 peptide (CKNDYRKL(p)SMQ). These phospho-specific antibodies were purified by antigen column. Anti-PDE3 antibody was from FabGennix Inc., and anti-PKA catalytic subunit (C α) antibody was from BD Transduction Laboratory. Anti-RhoA, anti-mouse IgG and anti-rabbit IgG antibodies were from Santa Cruz.

Isometric Force Measurements of Rat Aortic Rings and Reconstituted Rings

Rats were anaesthetized with sodium pentobarbital (50 mg/kg, i.p.) and descending thoracic aorta was carefully isolated, cleaned of any adhering connective tissue, and cut into 3-mm-long rings. Each ring was mounted between a pair of triangular stainless steel hooks, one of which was stationary and the other was connected to a strain-gauge transducer (UL10GR, Minebea), and the isometric tension was recorded (MC6625, Graphtec, Tokyo, Japan). The rings, placed in 20 ml organ baths containing Krebs Henseleit (KH) solution maintained at 37 °C and aerated continuously with 95% O₂ and 5% CO₂, were stretched to a resting tension of 1.5 g and allowed to equilibrate for at least 30 min with occasional changes of KH solution. After equilibration, rings were set at 1.5 g and treated with or without cilostazol for 20 min. Then, the rings were re-set at 1.5 g and incubated for 10 min. We confirmed that the tension before agonist stimulation was 1.5 ± 0.1 g. The endothelium was removed from some preparations by gentle rubbing of the intimal surface with a paper swab moistened with KH solution. The presence or absence of endothelium was confirmed by examining the relaxations in response to acetylcholine (1 μ M) of aortic rings precontracted by KCl (60 mM).

Production of a Reconstituted Ring and Measurement of Tension

Production of a reconstituted ring and measurement of tension were performed as described previously (4). The cultured primary rat aortic smooth muscle cells were collected with trypsinization and transfected with pCIneo vector encoding wild type TRPC6 or TRPC6 (T69A) mutant and pEGFP-N1 using electroporation (1400V, 10 msec x 3, Digital-Bio). Then cell suspensions were mixed with the culture medium (5 x

10^6 cells/ml) containing 0.6 mg/ml type I collagen (Cellmatrix Type I-P). For the expression of TRPC6 (3A) mutant (5), trypsinized cells were infected with adenoviruses encoding GFP or TRPC6 (3A) at 100 MOI in serum-free culture medium for 30 min and then mixed with collagen-containing culture medium. The mixture (100 μ l, 0.5×10^6 cells) was placed around a silicone column ($\phi = 2$ mm), which was attached on the center of the silicone disc ($\phi = 8$ mm). After the collagen gel became solid, the culture medium was added and cultured for 24-48 h. The ring preparation so obtained was used for measurements of isometric tension development. We counted GFP-positive RAoSMCs using fluorescent microscopy and determined the transfection efficiency of electroporation was $70 \pm 5\%$ and that of adenoviral infection was more than 90% in monolayer-cultured RAoSMCs plated on the collagen-coated glass-base dish.

For the tension recordings, the reconstituted rings were mounted onto two tungsten wires by passing the wires through the lumen of the rings. The resting tension was adjusted to 22 mg in Tyrode solution (NaCl 118 mM, KCl 5.4 mM, D-Glucose 10 mM, Hepes 10 mM, NaH_2PO_4 0.373 mM, MgCl_2 2 mM, CaCl_2 2 mM, taurine 1.5 mM). The rings were allowed to equilibrate for 60 min before starting experiments. Changes in tension were recorded isometrically through force displacement transducer (UL-2GR, Minebea).

Immunoprecipitation and Western Blot Analysis

HEK293 cells plated on 10 cm plates were harvested with 1 ml of RIPA buffer. Cell lysates were homogenized and supernatants were fractionated by centrifugation for

3 min at 10,000 rpm. GFP-fused proteins were immunoprecipitated in supernatants containing anti-GFP antibody (1.5 μ l) in the presence of a 20 μ l bed volume of protein A Sepharose beads rocking for 2 h at 4 °C (6). The immune complex were washed three times with RIPA buffer and resuspended with 80 μ l of SDS sample buffer. For Western blotting, TRPC6-expressing HEK293 cells (3×10^5 cells) plated on 6 well dishes were directly harvested with 2 x SDS sample buffer (200 μ l). After centrifugation, supernatants (20-40 μ l) were fractionated by 8% SDS-PAGE gel and then transferred onto PVDF membrane. The expression and phosphorylation of endogenous TRPC6 proteins were detected by anti-TRPC6 (dilution rate: 1/1000) and anti-phospho-TRPC6 (1/1000) antibodies (3). The reactive bands were visualized using Supersignal[®] West Pico Luminol/Enhancer solution (Pierce). The optical density of the film was scanned and measured with Scion Image Software.

Measurement of Rho Activity


Activation of small G proteins was determined as described previously (7, 8). Activated Rho was pulled down with 10 μ g of GST-fused Rho-binding domain of Rhotekin and 20 μ l bed volume of glutathione Sepharose beads. Pull-downed Rho was detected with anti-RhoA antibody.

Reference


1. Inoue R, Okada T, Onoue H, Hara Y, Shimizu S, Naitoh S, Ito Y, Mori Y. The transient receptor potential protein homologue TRP6 is the essential component of vascular α 1-adrenoceptor-activated Ca^{2+} -permeable cation channel. *Circ Res.* 2001; 88:325-32.
2. Takahashi S, Lin H, Geshi N, Mori Y, Kawarabayashi Y, Takami N, Mori MX, Honda A, Inoue R. Nitric oxide-cGMP-protein kinase G pathway negatively regulates vascular transient receptor potential channel TRPC6. *J Physiol.* 2008; 586:4209-23.
3. Nishida M, Watanabe K, Sato Y, Nakaya M, Kitajima N, Ide T, Inoue R, Kurose H. Phosphorylation of TRPC6 channels at Thr⁶⁹ is required for anti-hypertrophic effects of phosphodiesterase 5 inhibition. *J Biol Chem.* 2010; 285:13244-13253.
4. Bi D, Nishimura J, Niuro N, Hirano K, Kanaide H. Contractile properties of the cultured vascular smooth muscle cells: the crucial role played by RhoA in the regulation of contractility. *Circ Res.* 2005; 96:890-897.
5. Onohara N, Nishida M, Inoue R, Kobayashi H, Sumimoto H, Sato Y, Mori Y, Nagao T, Kurose H. TRPC3 and TRPC6 are essential for angiotensin II-induced cardiac hypertrophy. *EMBO J.* 2006; 25:5305-16.
6. Nishida M, Sugimoto K, Hara Y, Mori E, Morii T, Kurosaki T, Mori Y. Amplification of receptor signalling by Ca^{2+} entry-mediated translocation and activation of PLC γ 2 in B lymphocytes. *EMBO J.* 2003;22:4677-4688.
7. Nishida M, Tanabe S, Maruyama Y, Mangmool S, Urayama K, Nagamatsu Y, Takagahara S, Turner JH, Kozasa T, Kobayashi H, Sato Y, Kawanishi T, Inoue R, Nagao T & Kurose H. G $\alpha_{12/13}$ -and reactive oxygen species-dependent activation of

c-Jun NH₂-terminal kinase and p38 MAPK by angiotensin receptor stimulation in rat neonatal cardiomyocytes. *J Biol Chem.* 2005;280:18434-18441.


8. Nishida M, Sato Y, Uemura A, Narita Y, Tozaki-Saitoh H, Nakaya M, Ide T, Suzuki K, Inoue K, Nagao T & Kurose H. P2Y₆ Receptor-Gα_{12/13} Signaling in Cardiomyocytes Triggers Pressure Overload-induced Cardiac Fibrosis. *EMBO J.* 2008;27:3104-3115.




Supplemental Figure I. Milrinone suppresses Ang II-induced vasoconstriction by inhibiting of Ang II-induced Ca^{2+} influx. Ang II-induced contraction in the aortas pretreated with SK&F96365 (SK&F; 7 μM) with or without milrinone (Mil; 5 μM) for 30 min before Ang II stimulation. **: $P < 0.01$ vs control (none), n.s.; not significant.




Supplemental Figure II. Suppression of DAG-mediated Ca²⁺ influx by PDE3 inhibition. (A-D) Average time courses of Ca²⁺ responses induced by DAG (1-oleoyl-2-acetyl-sn-glycerol, OAG) (30 μM) in the absence or presence of extracellular Ca²⁺ in vector (A), TRPC3 (B), TRPC6 (C), and TRPC7 (D)-overexpressing HEK293 cells. Cells were pretreated with indicated concentration of cilostazol (CLZ) for 30 min before OAG treatment. (E, F) Peak Ca²⁺ release-mediated increases in [Ca²⁺]_i (E) and Ca²⁺ influx-mediated increases in [Ca²⁺]_i (F) induced by OAG in the presence or absence of CLZ. N=42-76 cells.




Supplemental Figure III. Inhibition of PDE3 does not suppress DAG-independent Ca²⁺ influx. (A-C) Effects of cilostazol (CLZ) on TRPC5-mediated Ca²⁺ influx induced by ATP stimulation. (A) Average time courses of Ca²⁺ responses induced by ATP in TRPC5-GFP-overexpressing HEK293 cells. (B, C) Peak changes in [Ca²⁺]_i induced by ATP in the absence (B) or presence (C) of extracellular Ca²⁺ (2 mM). (D, E) Effects of cilostazol on Ca²⁺ influx induced by store depletion. Cells were pretreated with CLZ (10 μM) for 30 min before ionomycin treatment. (D) Average time courses of Ca²⁺ responses induced by ionomycin (1 μM) in the absence or presence of extracellular Ca²⁺. (E) Peak increases in [Ca²⁺]_i induced by ionomycin. N=108-136 cells.




Supplemental Figure IV. Time-dependent inhibition of TRPC6-mediated Ca²⁺ influx by PDE3 inhibition. (A, C) Average time courses of Ca²⁺ responses induced by OAG (A) and ATP (C) in the absence or presence of extracellular Ca²⁺ in TRPC6-expressing HEK293 cells. Cells were pretreated with cilostazol (CLZ) for the indicated time before treatment with OAG (30 μM) or ATP (100 μM). (B, D) Peak Ca²⁺ influx-mediated increases in [Ca²⁺]_i induced by OAG (B) or ATP (D). N=69-92 cells.



Supplemental Figure V. Inhibition of PDE3 does not affect membrane localization of TRPC3 and TRPC6 proteins. (A-F) Visualization of GFP (A, D), GFP-fused TRPC3 (B, E) and GFP-fused TRPC6 (C, F) proteins in the absence (A-C) or presence (D-F) of cilostazol (CLZ). Cells were treated with CLZ (1 μ M) for 30 min. N=3 experiments.



Supplemental Figure VI. Involvement of phosphorylation of TRPC6 at Thr69 in suppression of DAG-mediated Ca²⁺ influx by PDE3 inhibition. (A, B) Effects of CLZ on DAG (1-oleoyl-2-acetyl-sn-glycerol, OAG)-induced Ca²⁺ influx in wild type TRPC6 (TRPC6-wt)- or TRPC6-T69A-expressing cells. (A) Average time courses of changes in [Ca²⁺]_i induced by OAG (30 μM). Cells were pretreated with CLZ for 30 min before OAG stimulation. (B) Peak increases in [Ca²⁺]_i induced by OAG in the presence of extracellular Ca²⁺. (C, D) Average time courses (C) and Peak changes (D) of [Ca²⁺]_i induced by OAG in TRPC6-wt- or TRPC6-T69A-expressing cells. Cells were pretreated with 8-Br-cAMP (30 μM) or forskolin (10 μM) for 30 min before OAG stimulation. (E) Conserved PKG/PKA-phosphorylation sites in mouse TRPC3 (836 a.a.), TRPC6 (930 a.a.), and TRPC7 (862 a.a.). (F) Peak Ca²⁺ influx-mediated Ca²⁺ increases induced by OAG (30 μM) in vector-, wild type TRPC3-, and TRPC3-S251A-expressing HEK293 cells. **, P < 0.01 vs control (-), **#, P < 0.01 vs TRPC6-wt. N=38-74 cells.



Supplemental Figure VII. PKA-dependent suppression of TRPC6-mediated Ca^{2+} influx by PDE3 inhibition. (A-C) Average time courses of Ca^{2+} responses induced by ATP in TRPC6-expressing cells in the presence of PKA inhibitor (KT5720) (A), PKG inhibitor (KT5823) (B), and PKC inhibitor (Go6976) (C). Cells were pretreated with indicated concentration of cilostazol (CLZ) with KT5823 (0.3 μM), KT5720 (0.1 μM), or Go6976 (20 nM) for 30 min before stimulation with ATP (100 μM). (D, E) Peak Ca^{2+} release-mediated $[\text{Ca}^{2+}]_i$ increases (D) and Ca^{2+} influx-mediated $[\text{Ca}^{2+}]_i$ increases (E) induced by ATP in the absence or presence of CLZ. *, $P < 0.05$, **, $P < 0.01$ vs CLZ (-), #, $P < 0.05$, ###, $P < 0.01$ vs none.

ヒト幹細胞を用いた細胞・組織加工医薬品等の品質・安全性確保に関する指針整備と主なポイント

早川 堯夫（近畿大学薬学総合研究所所長）

青井 貴之（京都大学iPS細胞研究所教授）

梅澤 明弘（国立成育医療センター生殖医療研究部部長）

小澤 敬也（自治医科大学医学部内科学講座血液学部門教授）

佐藤 陽治（国立医薬品食品衛生研究所遺伝子細胞医薬部第2室長）

澤 芳樹（大阪大学大学院医学系研究科外科学講座教授）

松山 晃文（(財)先端医療振興財団再生医療研究開発部門部門長補佐）

大和 雅之（東京女子医科大学先端生命医学研究所教授）

山中 伸弥（京都大学iPS細胞研究所所長）

はじめに

細胞・組織加工医薬品等による再生医療は、ヒトの臓器や組織の確保が難しいわが国の医療状況下において強く期待されており、研究の進歩に伴う技術的な実現可能性の高まりとともに、医療としての実用化を望む声がますます強くなっている。

また、平成19年11月の総合科学技術会議において、その研究成果が注目されている人工多能性幹細胞（iPS細胞）について意見交換が行われ、今後、再生医療研究の加速のための支援のあり方等を検討することが必要との意見が出された。その後わが国を挙げて再生医療の実用化に向けた動きが急速に進められる中で、ヒト由来の多能性細胞、すなわち間葉系幹細胞などの体性幹細胞、胚性幹細胞（ES細胞）、さらには人工多能性幹細胞（iPS細胞）について、近い将来に予想される製品の評価を円滑に進めるための準備を早期に行う必要があると考えられた。

再生医療におけるヒト幹細胞の活用について、ヒト胚性幹細胞は倫理上使用が限定され実用化は困難とされていたが、最近、米国ではその使用が認められ、製品も次々に開発されつつある。さらに間葉系幹細胞やiPS細胞由来製品については国際的に実用化に向けた研究開発が活発に行われている。各種幹細胞の実用化の為に必要な要件を行政当局が開発早期から示すことは、研究開発者、企業、規制側にとり有用であり、幹細胞を用いた再生医療をより迅速に実用化するために規制環境整備という面から支えるという意味で必要不可欠である。

1. ヒト幹細胞を用いる再生医療 関連指針整備

現在のところ、再生医療には主に2つのアプローチがある。1つは薬事法下においての細胞・組織加工製品の製造販売承認を目指したものである。すなわち、企業による研究開発、治験に入る前にヒトに適用する（First-in-Man）に際して支障がないという確認を公的に規制当局から得るための評価に供する「確認申請（注：現行制度上のステップ。但し平成22年11月1日付け906号通知一部改正通知（薬食発1101第3号2）により、後述するヒト幹臨床研究指針に適合していれば確認申請はスキップできることになった。また、きわめて近い将来に薬事戦略相談制度をもって代替することとなっているが、本稿中では同様の趣旨を示す用語として確認申請を用いている）」、そして「治験」、「製造販売承認」、「臨床利用」という段階を踏むアプローチである。このようなアプローチをとる際の基本的考え方や、基本的技術要件については、平成12年12月26日医薬発第1314号「ヒト又は動物由来成分を原料として製造される医薬品等の品質及び安全性確保について」に示されていた。2つ目は、ひとまず医師法下で行われる「ヒト幹細胞臨床研究」で、これは平成18年7月3日付け厚生労働省告示第425号「ヒト幹細胞を用いる臨床研究に関する指針」及び平成18年7月3日付け厚生労働省健康局長通知健発第0703003号「ヒト幹細胞を用いる臨床研究に関する指針の施行等について」（注：改訂版が平成22年11月に出された〔厚生労働省告示第380号〕）に従って行うことになっているが、この科学的内容は、おおそ医薬発第1314号及び後述する平成20年の「自己」及び「同種」改訂版に準じたものとなっている。「ヒト幹細胞臨床研究」の実施の是非は、厚生労働省