

HRMS. Zinc(II) or copper(II) complex formation was accomplished by treatment of the above compounds with 10 equiv of $ZnCl_2$ or $CuCl_2$ in PBS. All zinc(II) or copper(II) complexes were characterized by chemical shifts of their methylene protons in 1H NMR analysis. The pyridyl zinc(II) complex was characterized previously,^[37] and zinc(II) or copper(II) complex formation with these macrocyclic compounds has been reported elsewhere.^[41,42,48,49] Detailed procedures and data are provided in the Supporting Information.

Biological assays

A CXCR4 binding assay for compounds, based on the inhibition of [^{125}I]CXCL12 binding to Jurkat cells, was performed as reported by Tanaka et al.^[38] CXCR4 antagonistic activity was evaluated as described by Ichiyama et al.^[27], measuring inhibitory activity against Ca^{2+} mobilization induced by CXCL12 stimulation in HOS cells expressing CXCR4. Anti-HIV activity was determined by inhibitory activity against X4-HIV-1(NL4-3)-induced cytopathogenicity in MT-4 cells as reported by Tanaka et al.^[38] An X4 HIV-1 infectious molecular clone (pNL4-3) was obtained from the AIDS Research and Reference Reagent Program. The virus NL4-3 was obtained from the culture supernatant of 293T cells transfected with pNL4-3.

Molecular modeling

Molecular modeling calculations were performed using Sybyl (version 7.0, Tripos). Energy minimization was performed using the Tripos force field and Gasteiger-Hückel charge parameters. The lowest energy conformation was obtained by random search methods.

Acknowledgements

T.T. and N.O. are supported by research fellowships for young scientists from the Japan Society for the Promotion of Science. This work was supported in part by a Grant-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science, and Technology of Japan, and Health and Labor Sciences Research Grants from the Japanese Ministry of Health, Labor, and Welfare. The authors thank Mr. Wu Honggui (Tokyo University of Science) for his assistance with the anti-HIV assay.

Keywords: azamacrocycles • Ca^{2+} mobilization • CXCR4 • HIV • structure-activity relationships

- [1] T. Nagasawa, H. Kikutani, T. Kishimoto, *Proc. Natl. Acad. Sci. USA* **1994**, *91*, 2305–2309.
- [2] C. C. Bleul, M. Farzan, H. Choe, C. Parolin, I. Clark-Lewis, J. Sodroski, T. A. Springer, *Nature* **1996**, *382*, 829–833.
- [3] E. Oberlin, A. Amara, F. Bachelier, C. Bessia, J. L. Virelizier, F. Arenzana-Seisdedos, O. Schwartz, J. M. Heard, I. Clark-Lewis, D. L. Legler, M. Loetscher, M. Baggiolini, B. Moser, *Nature* **1996**, *382*, 833–835.
- [4] K. Tashiro, H. Tada, R. Heilker, M. Shirozu, T. Nakano, T. Honjo, *Science* **1993**, *261*, 600–603.
- [5] C. C. Bleul, R. C. Fuhlbrigge, J. M. Casanovas, A. Aiuti, T. A. Springer, *J. Exp. Med.* **1996**, *184*, 1101–1109.
- [6] K. Tachibana, S. Hirota, H. Iizasa, H. Yoshida, K. Kawabata, Y. Kataoka, Y. Kitamura, K. Matsushima, N. Yoshida, S. Nishikawa, T. Kishimoto, T. Nagasawa, *Nature* **1998**, *393*, 591–594.
- [7] T. Nagasawa, S. Hirota, K. Tachibana, N. Takakura, S. Nishikawa, Y. Kitamura, N. Yoshida, H. Kikutani, T. Kishimoto, *Nature* **1996**, *382*, 635–638.
- [8] Y. Zhu, Y. Yu, X. C. Zhang, T. Nagasawa, J. Y. Wu, Y. Rao, *Nat. Neurosci.* **2002**, *5*, 719–720.
- [9] R. K. Stumm, C. Zhou, T. Ara, F. Lazarini, M. Dubois-Dalcq, T. Nagasawa, V. Holtz, S. Schulz, *J. Neurosci.* **2003**, *23*, 5123–5130.
- [10] H. K. Deng, R. Liu, W. Ellmeier, S. Choe, D. Unutmaz, M. Burkhart, P. D. Marzio, S. Marmon, R. E. Sutton, C. M. Hill, C. B. Davis, S. C. Peiper, T. J. Schall, D. R. Littman, N. R. Landau, *Nature* **1996**, *381*, 661–666.
- [11] Y. Feng, C. C. Broder, P. E. Kennedy, E. A. Berger, *Science* **1996**, *272*, 872–877.
- [12] T. Koshiba, R. Hosotani, Y. Miyamoto, J. Ida, S. Tsuji, S. Nakajima, M. Kawaguchi, H. Kobayashi, R. Doi, T. Hori, N. Fujii, M. Imamura, *Clin. Cancer Res.* **2000**, *6*, 3530–3535.
- [13] A. Müller, B. Homey, H. Soto, N. Ge, D. Catron, M. E. Buchanan, T. McClanahan, E. Murphy, W. Yuan, S. N. Wagner, J. L. Barrera, A. Mohar, E. Verstegui, A. Zlotnik, *Nature* **2001**, *410*, 50–56.
- [14] H. Tamamura, A. Hori, N. Kanzaki, K. Hiramatsu, M. Mizumoto, H. Nakashima, N. Yamamoto, A. Otaka, N. Fujii, *FEBS Lett.* **2003**, *550*, 79–83.
- [15] N. Tsukada, J. A. Burger, N. J. Zvaifler, T. J. Kipps, *Blood* **2002**, *99*, 1030–1037.
- [16] J. Juarez, K. F. Bradstock, D. J. Gottlieb, L. J. Bendall, *Leukemia* **2003**, *17*, 1294–1300.
- [17] T. Nanki, K. Hayashida, H. S. El-Gabalawy, S. Suson, K. Shi, H. J. Girschick, S. Yavuz, P. E. Lipsky, *J. Immunol.* **2000**, *165*, 6590–6598.
- [18] H. Tamamura, M. Fujisawa, K. Hiramatsu, M. Mizumoto, H. Nakashima, N. Yamamoto, A. Otaka, N. Fujii, *FEBS Lett.* **2004**, *569*, 99–104.
- [19] T. Murakami, T. Nakajima, Y. Koyanagi, K. Tachibana, N. Fujii, H. Tamamura, N. Toshida, M. Waki, A. Matsumoto, O. Yoshie, T. Kishimoto, N. Yamamoto, T. Nagasawa, *J. Exp. Med.* **1997**, *186*, 1389–1393.
- [20] D. Schols, S. Struyf, J. Van Damme, J. A. Este, G. Henson, E. DeClarcq, *J. Exp. Med.* **1997**, *186*, 1383–1388.
- [21] B. J. Doranz, K. Grovit-Ferbas, M. P. Sharron, S.-H. Mao, M. Bidwell Goetz, E. S. Daar, R. W. Doms, W. A. O'Brien, *J. Exp. Med.* **1997**, *186*, 1395–1400.
- [22] G. A. Donzella, D. Schols, S. W. Lin, J. A. Este, K. A. Nagashima, *Nat. Med.* **1998**, *4*, 72–76.
- [23] O. M. Z. Howard, J. J. Oppenheim, M. G. Hollingshead, J. M. Covey, J. Bigelow, J. J. McCormack, R. W. Buckheit, Jr., D. J. Clanton, J. A. Turpin, W. G. Rice, *J. Med. Chem.* **1998**, *41*, 2184–2193.
- [24] H. Tamamura, Y. Xu, T. Hattori, X. Zhang, R. Arakaki, K. Kanbara, A. Omagari, A. Otaka, T. Ibuka, N. Yamamoto, H. Nakashima, N. Fujii, *Biochem. Biophys. Res. Commun.* **1998**, *253*, 877–882.
- [25] H. Tamamura, A. Omagari, S. Oishi, T. Kanamoto, N. Yamamoto, S. C. Peiper, H. Nakashima, A. Otaka, N. Fujii, *Bioorg. Med. Chem. Lett.* **2000**, *10*, 2633–2637.
- [26] N. Fujii, S. Oishi, K. Hiramatsu, T. Araki, S. Ueda, H. Tamamura, A. Otaka, S. Kusano, S. Terakubo, H. Nakashima, J. A. Broach, J. O. Trent, Z. Wang, S. C. Peiper, *Angew. Chem.* **2003**, *115*, 3373–3375; *Angew. Chem. Int. Ed.* **2003**, *42*, 3251–3253.
- [27] K. Ichiyama, S. Yokoyama-Kumakura, Y. Tanaka, R. Tanaka, K. Hirose, K. Bannai, T. Edamatsu, M. Yanaka, Y. Niitani, N. Miyano-Kurosaki, H. Takaku, Y. Koyanagi, N. Yamamoto, *Proc. Natl. Acad. Sci. USA* **2003**, *100*, 4185–4190.
- [28] H. Tamamura, N. Fujii, *Curr. Drug Targets-Infectious Disorders* **2004**, *4*, 103–110.
- [29] H. Tamamura, K. Hiramatsu, S. Ueda, Z. Wang, S. Kusano, S. Terakubo, J. O. Trent, S. C. Peiper, N. Yamamoto, H. Nakashima, A. Otaka, N. Fujii, *J. Med. Chem.* **2005**, *48*, 380–391.
- [30] H. Tamamura, T. Araki, S. Ueda, Z. Wang, S. Oishi, A. Esaka, J. O. Trent, H. Nakashima, N. Yamamoto, S. C. Peiper, A. Otaka, N. Fujii, *J. Med. Chem.* **2005**, *48*, 3280–3289.
- [31] G. C. Valks, G. McRobbie, E. A. Lewis, T. J. Hubin, T. M. Hunter, P. J. Sadler, C. Pannecouque, E. De Clercq, S. J. Archibald, *J. Med. Chem.* **2006**, *49*, 6162–6165.
- [32] W. Zhan, Z. Liang, A. Zhu, S. Kurtkaya, H. Shim, J. P. Snyder, D. C. Liotta, *J. Med. Chem.* **2007**, *50*, 5655–5664.
- [33] A. Khan, G. Nicholson, J. Greenman, L. Madden, G. McRobbie, C. Pannecouque, E. De Clercq, R. Ullom, D. L. Maples, R. D. Maples, J. D. Silverides, T. J. Hubin, S. J. Archibald, *J. Am. Chem. Soc.* **2009**, *131*, 3416–3417.
- [34] G. J. Bridger, R. T. Skerlj, P. E. Hernandez-Abad, D. E. Bogucki, Z. Wang, Y. Zhou, S. Nan, E. M. Boehringer, T. Wilson, J. Crawford, M. Metz, S. Hatse, K. Princen, E. De Clercq, D. Schols, *J. Med. Chem.* **2010**, *53*, 1250–1260.

- [35] R. T. Skerlj, G. J. Bridger, A. Kaller, E. J. McEachern, J. B. Crawford, Y. Zhou, B. Atsma, J. Langille, S. Nan, D. Veale, T. Wilson, C. Harwig, S. Hatse, K. Princen, E. De Clercq, D. Schols, *J. Med. Chem.* **2010**, *53*, 3376–3388.
- [36] M. Takenaga, H. Tamamura, K. Hiramatsu, N. Nakamura, Y. Yamaguchi, A. Kitagawa, S. Kawai, H. Nakashima, N. Fujii, R. Igarashi, *Biochem. Biophys. Res. Commun.* **2004**, *320*, 226–232.
- [37] H. Tamamura, A. Ojida, T. Ogawa, H. Tsutsumi, H. Masuno, H. Nakashima, N. Yamamoto, I. Hamachi, N. Fujii, *J. Med. Chem.* **2006**, *49*, 3412–3415.
- [38] T. Tanaka, H. Tsutsumi, W. Nomura, Y. Tanabe, N. Ohashi, A. Esaka, C. Ochiai, J. Sato, K. Itotani, T. Murakami, K. Ohba, N. Yamamoto, N. Fujii, H. Tamamura, *Org. Biomol. Chem.* **2008**, *6*, 4374–4377.
- [39] G. J. Bridger, R. T. Skerlj, D. Thornton, S. Padmanabhan, S. A. Martellucci, G. W. Henson, M. J. Abrams, N. Yamamoto, K. De Vreese, R. Pauwels, E. De Clercq, *J. Med. Chem.* **1995**, *38*, 366–378.
- [40] G. J. Bridger, R. T. Skerlj, S. Padmanabhan, S. A. Martellucci, G. W. Henson, M. J. Abrams, H. C. Joao, M. Witvrouw, K. De Vreese, R. Pauwels, E. De Clercq, *J. Med. Chem.* **1996**, *39*, 109–119.
- [41] Y. Inouye, T. Kanamori, T. Yoshida, T. Koike, M. Shionoya, H. Fujioka, E. Kimura, *Biol. Pharm. Bull.* **1996**, *19*, 456–458.
- [42] L. O. Gerlach, J. S. Jakobsen, K. P. Jensen, M. R. Rosenkilde, R. T. Skerlj, U. Ryde, G. J. Bridger, T. W. Schwartz, *Biochemistry* **2003**, *42*, 710–717.
- [43] H. F. Egberink, E. De Clercq, A. L. Van Vliet, J. Balzarini, G. J. Bridger, G. Henson, M. C. Horzinek, D. Schols, *J. Virol.* **1999**, *73*, 6346–6352.
- [44] M. Le Baccon, F. Chuburu, L. Toupet, H. Handel, M. Soibinet, I. De-champs-Olivier, J.-P. Barbier, M. Aplincourt, *New J. Chem.* **2001**, *25*, 1168–1174.
- [45] B. Antonioli, D. J. Bray, J. K. Clegg, K. Gloe, K. Gloe, O. Kataeva, L. F. Lindoy, J. C. McMurtrie, P. J. Steel, C. J. Sumby, M. Wenzel, *Dalton Trans.* **2006**, 4783–4794.
- [46] S. P. Foxon, D. Utz, J. Astner, S. Schindler, F. Thaler, F. W. Heinemann, G. Liehr, J. Mukherjee, V. Balamurugan, D. Ghosh, R. Mukherjee, *Dalton Trans.* **2004**, 2321–2328.
- [47] S. Mandal, F. Lloret, R. Mukherjee, *Inorg. Chim. Acta* **2009**, *362*, 27–37.
- [48] M. Soibinet, I. De-champs-Olivier, E. Guillon, J.-P. Barbier, M. Aplincourt, F. Chuburu, M. Le Baccon, H. Handel, *Eur. J. Inorg. Chem.* **2003**, 1984–1994.
- [49] R. W. Hay, M. T. Tarafder, *Transition Met. Chem.* **1990**, *15*, 490–492.

Received: December 19, 2010

Published online on February 10, 2011

Fluorescence-Quenching Screening of Protein Kinase C Ligands with an Environmentally Sensitive Fluorophore


Wataru Nomura,[†] Nami Ohashi,[†] Yoshiaki Okuda,^{†,*} Tetsuo Narumi,[†] Teikichi Ikura,[†] Nobutoshi Ito,[†] and Hirokazu Tamamura^{*,†,‡}

[†]Department of Medicinal Chemistry, Institute of Biomaterials and Bioengineering, Tokyo Medical and Dental University, 2-3-10 Kandasurugadai, Chiyoda-ku, Tokyo 101-0062, Japan

[‡]Graduate School of Biomedical Science, Tokyo Medical and Dental University, 1-5-45 Yushima, Bunkyo-ku, Tokyo 113-8510, Japan

S Supporting Information

ABSTRACT: A novel fluorescence-quenching screening method for protein kinase C (PKC) ligands was developed utilizing solvatochromic fluorophores. Solvatochromic dyes, highly sensitive to the presence or the absence of competitive ligands in their binding to the C1b domain of PKC δ (δ C1b), were combined with a known pharmacophoric moiety of 1,2-diacylglycerol (DAG) lactones, PKC ligands. Addition of δ C1b to the fluorescent compounds caused a gradual increase in the fluorescent intensity in proportion to the increase of δ C1b. As a competitive ligand was added to the complex of δ C1b domain and fluorescent compounds, a gradual decrease in the fluorescent intensity was observed. The relative binding affinities of known ligands were successfully determined by this fluorescent method and corresponded well to the K_i values measured by a radioisotope method. These results indicate that washing, which is a laborious step in binding evaluations, is not required for this environmentally sensitive fluorophore based system. Screening with the system was performed for 2560 preselected library compounds with possible pharmacophores, and some lead compounds were found. This fluorescence-based method could be applied widely to known ligand–receptor combinations.



INTRODUCTION

Solvatochromic fluorophores are sensitive to immediate changes in their environment such as binding of ligands to proteins,¹ and significant biological issues have been examined by chemical probes with fluorescent dyes.^{2,3} The major advantage of the fluorescence-based methods in the observation of physiological phenomena in cells is that detection and evaluation can be performed in real-time directly by microscopy. In these techniques, the signal/background ratio is critical to avoid the observation of artifacts or to obtain clearer images at high resolution. Screening methods of bioactive compounds have been developed as another application of fluorescence.⁴ From the aspect of sensitivity, radioisotope (RI) based methods are more efficient but involve laborious experimental procedures. For the development of convenient screening procedures, a simple method for evaluation of binding is needed. By attachment of solvatochromic fluorophores to known ligands, the binding affinity of unknown ligands can be evaluated and related to RI methods.⁵ Fluorescent intensity is highly dependent on the binding state to a target protein, and the signal/background ratio would be sufficient without washing steps, which are normally necessary in fluorescent analysis.^{6,7}

In this study, we have developed reporter compounds containing solvatochromic dyes for the detection of ligand binding to a target protein and applied them to the screening of chemical libraries. Protein kinase C δ (PKC δ) was chosen as a target

protein because of its importance in physiological phenomena. The 11 isozymes that constitute PKC play pivotal roles in physiological responses to growth factors, oxidative stress, 1,2-diacylglycerol (DAG), and tumor promoters, such as phorbol esters. These responses regulate numerous cellular processes,^{8,9} including proliferation,¹⁰ differentiation,¹¹ migration,¹² and apoptosis.^{13,14} Membrane translocation of PKC is caused by binding of ligands to the C1b domain and has been recognized as an important phenomenon in signal transduction because the localization of PKC is key in determining isozyme-specific functions by defining binding partners for downstream signaling.¹⁵ Despite the complex regulatory mechanisms of PKC activation, considerable progress has been made in understanding isozyme-specific functions¹⁶ and several ligands with high specificity for PKC isozymes have been developed as potential drugs.^{17–23} Through the development of DAG-lactones, whose design is based on the endogenous ligand DAG, the importance of maintaining intact the pharmacophore triad of two carbonyl groups (*sn*-1 and *sn*-2) and the primary alcohol has been established as a requirement in high affinity ligands for PKC (Figure 1).¹⁶ Various PKC ligands have been synthesized and found to be inhibitors of tumor promotion or of other diseases.

Received: December 13, 2010

Revised: March 13, 2011

Published: March 25, 2011

However, there are limitations to the design of compound templates so far. Thus, in this study, we have attempted to develop the screening methods aimed at discovery of novel template structures for PKC δ ligands as drug leads.

EXPERIMENTAL PROCEDURES

General. ^1H NMR spectra were recorded using a Bruker AV500 spectrometer. Chemical shifts were reported in δ (ppm) relative to Me_4Si (in CDCl_3) as an internal standard. Low- and




Figure 1. Binding mode of DAG-lactone derivatives to the δC1b domain. The *sn*-1 and *sn*-2 carbonyl groups are indicated. Three important pharmacophores are indicated by light blue spheres. Amino acid residues interacting with these pharmacophores (threonine at 242, leucine at 251, and glycine at 253) are indicated by red spheres.


high-resolution mass spectra were recorded on a JMS-T1000LC AccuTOF and Bruker Daltonics microTOF-2focus in both positive and negative detection modes. Wakogel C-200 (Wako Pure Chemical Industries, Ltd.) and silica gel 60 N (Kanto Chemical Co., Inc.) were employed for flash chromatography. Fluorescent spectra were recorded on a JASCO FP-6600 spectrofluorometer and JASCO V-650 spectrophotometer using a quartz cell with 1.0 cm path length. Fluorescent intensities of samples in 96-well plates were recorded on Wallac ARVO MX.

Standard Monoacylation Procedure. Under argon, Et_3N (3 equiv) was added at 0°C to a solution containing 5,5-bis(hydroxymethyl)oxalan-2-one in THF, and the mixture was stirred at 0°C for 30 min. Then 0.2–1.1 equiv of a carboxylic acid derivative (an acyl chloride or carboxylic anhydride) was added. This mixture was stirred at 0°C for a further 4 h. After evaporation of the solvent, the obtained residue was purified by flash column chromatography.

Preparation of Compounds 7, 10–13, 16, 17, and 19–21. Compounds 7, 10–12, and 16 were prepared as described in ref 17. Compounds 13 and 17 were prepared as described in refs 24 and 25. The synthesis of 19 is described elsewhere.²⁶ Compounds 20 and 21 are commercially available.

Expression and Purification of the δC1b Domain. DNA coding C1b domain of mouse PKC δ (231–280)²⁷ was subcloned into *Bam*HI and *Eco*RI sites of pGEX-2tk (GE Healthcare) and expressed as glutathione-S-transferase (GST) fusion protein in *Escherichia coli* C41 which contains extension sequences Gly-Ser-Arg-Arg-Ala-Ser-Val-Gly-Ser and Glu-Phe-Ile-Val-Thr-Asp at the N- and C-termini, respectively. The δC1b

Scheme 1. Synthesis of Fluorescent Compounds 1–3^a



^a (a) LDA, -78°C ; (b) MsCl , DBU , 0°C to room temp; (c) TBAF , 0°C to room temp; (d) $\text{C}_9\text{H}_{19}\text{COCl}$, Et_3N , 0°C ; (e) EDCl , DMAP , 0°C ; (f) SOCl_2 , Et_3N , 0°C .




Figure 2. Fluorescent spectra of compounds 1 (A), 2 (B), and 3 (C) in different solvents. The spectra show fluorescence in the solvents color-coded as follows: red, MeOH; blue, THF; black, CHCl₃; green, EtOAc. Each ligand was prepared as 1 mM solution in DMSO. In UV measurement, 10 μ L of ligand was added to 990 μ L of solvents.




Figure 3. Schematic representation of a fluorescence-quenching screening system. In the presence of the δ C1b domain, fluorescence of a solvatochromic fluorophore is enhanced. The replacement by candidate compounds results in fluorescence quenching.

Table 1. Changes of Fluorescent Intensity upon δ C1b Binding and Inhibition Constants to δ C1b Determined by Competition Assay against [³H]PDBu

compd	ΔF^a	K_i (nM) ^b
1	3.8	35.9
2	2.3	368
3	0.50	93.3

^a Fluorescent change by binding of the δ C1b domain. The values were determined by titration of the δ C1b domain. The ΔF was determined by dividing F_1 by F_0 at the maximum of fluorescent emission in each spectrum. ^b Inhibitory constants determined by competition analysis against [³H]PDBu.^{28,36,37}

domain after cleavage contains 65 amino acid residues. Cells were grown at 37 °C in LB medium and induced with 0.3 mM IPTG at growth phase. Cells were cultured overnight at 20 °C after induction. Cells were collected and lysed in 50 mM Tris·HCl buffer (pH 8.0) containing 100 mM NaCl, 1 μ g/mL leupeptine, 1 mM PMSF, and 1 mM DTT. Expressed protein was extracted by sonication, then purified by affinity chromatography utilizing glutathione-Sepharose 4B beads resin. GST moiety was cleaved by 300 units of thrombin at 4 °C overnight. The cleaved protein was eluted and further purified on a 2.6 cm \times 60 cm Superdex S-75 gel filtration column. Purification was by FPLC system at a flow rate of 1 mL/min utilizing 0.1 M triethanolamine·HCl (pH 7.0) containing 0.5 M NaCl.




Figure 4. Changes of fluorescent spectra of compound **1** during titration of the δ C1b domain (A) and during titration of PDBu (compound **8**) (B) after δ C1b binding. The concentration of the δ C1b domain was increased to $1.28 \mu\text{M}$ (6.4 equiv to compound **1**) by titration. The concentration of PDBu was increased to $10.2 \mu\text{M}$ (51.2 equiv to compound **1**) by titration.




Figure 5. Competitive assay against compound **1** based on fluorescent intensity (spectrofluorometer). (A) Structures of test compounds. (B) Inhibition curves indicate results of compounds **8** (\blacktriangle), **9** (\circ), **10** (\bullet), **11** (\blacksquare), **12** (\square), **13** (\diamond), and **14** (\blacktriangledown).

Determination of K_i of Compounds for Full Length PKC δ . Enzyme–ligand interaction was assessed in vitro as the ability of the ligand to displace bound [^3H]phorbol 12,13-dibutyrate ([^3H]PDBu) from recombinant human PKC δ in the presence of phosphatidylserine (PS) in an experimental procedure that was described previously.²⁸ In constant concentration analysis, the concentrations of compounds were fixed at 100 nM for known compounds and at $10 \mu\text{M}$ for library compounds. The other procedures were the same as those of the standard evaluation.

Fluorescent Titration of the δ C1b Domain. An amount of 25 μL of a solution of PS in chloroform (10 mg/mL) was evaporated to dryness under nitrogen. Then 50 mM Tris·HCl buffer (pH 7.4) was added and the PS was sonicated with a Microtip for a total of 15 s. A stock solution of compound **1** was diluted with 50 mM Tris·HCl buffer (pH 7.4) containing 100 $\mu\text{g}/\text{mL}$ PS to prepare a $0.2 \mu\text{M}$ solution. The recombinant δ C1b domain was added to the $0.2 \mu\text{M}$ solution of compound **1**, and fluorescent spectra ($\lambda_{\text{ex}} = 340 \text{ nm}$) were measured at 25 $^\circ\text{C}$. The concentration of the recombinant δ C1b domain started at 0.01 μM and increased to $1.28 \mu\text{M}$ by titration. The final concentrations of compounds **2** and **3** were 0.2 and 0.05 μM , respectively. Change of fluorescent intensity (ΔF) was calculated at the following wavelengths, which showed λ_{max} in each evaluation; compound **1**, 406 (F_0) and 394 (F_1) nm; compound **2**, 520 (F_0) and 480 (F_1) nm; compound **3**, 463 (F_0) and 461 (F_1) nm. Wavelengths recorded before and after the addition of the δ C1b domain.

Fluorescent Titration of Candidate Ligands. Compound **1** was diluted with 50 mM Tris·HCl (pH 7.4) containing

100 $\mu\text{g}/\text{mL}$ PS to obtain a solution with a $0.2 \mu\text{M}$ final concentration. The δ C1b domain was added to the above solution of compound **1** to be $0.96 \mu\text{M}$. A candidate ligand was added to the solution, and fluorescent spectra were recorded with a spectrofluorometer at 25 $^\circ\text{C}$. Fluorescent titration curves ($\lambda_{\text{em}} = 407 \text{ nm}$) were analyzed by nonlinear regression, and IC_{50} values were estimated by a nonlinear least-squares curve-fitting method using GraphPad Prism 5 (GraphPad Software, Inc., La Jolla, CA, U.S.). In the constant concentration analysis, the concentrations of known compounds were fixed at 100 nM and of library compounds at $10 \mu\text{M}$. The other procedures were the same as those in the standard evaluation.

Fluorescence-Quenching Analysis for Library Screening Recorded by Microplate Reader. Fluorescence intensity was recorded on Wallac ARVO MX (PerkinElmer) using a 96-well black plate. Compound **1** was diluted with 50 mM Tris·HCl (pH 7.4) containing 100 $\mu\text{g}/\text{mL}$ PS to obtain a solution with a $0.2 \mu\text{M}$ final concentration. The δ C1b domain was added to the above solution of compound **1** to be $0.96 \mu\text{M}$. A candidate ligand ($10 \mu\text{M}$) was added to the compound **1**– δ C1b domain complex solution, and fluorescent intensities ($\lambda_{\text{ex}} = 355 \text{ nm}$, $\lambda_{\text{em}} = 405 \text{ nm}$) were measured at 25 $^\circ\text{C}$.

RESULTS AND DISCUSSION

Synthesis and Fluorescent Properties of Compounds 1–3. Three representative solvatochromic fluorophores, involving 6-methoxynaphthalene,²⁹ 5-(dimethylamino)naphthalene-1-sulfonyl (dansyl),^{6,30,31} and diethylaminocoumarin,^{32–35} have




Figure 6. Comparison of inhibition percentages between the fluorescent assay (microplate reader) and the RI assay at a constant concentration of test compounds. (A) Structures of test compounds. (B) Inhibition of binding of compound 1 (fluorescent assay, left gray bars) or $[^3\text{H}]\text{PDBu}$ (RI assay, right black bars) in the presence of test compounds ($10 \mu\text{M}$ in the fluorescent assay and 100 nM in the RI assay). The final DMSO concentration is 4%. (C) Plots of IC_{50} values (μM) by the fluorescent assay versus K_i (nM) by the RI assay for compounds 8, 9, 10, 11, and 17.

been utilized. The 6-methoxynaphthalene was incorporated at the α -carbon of the *sn*-1 carbonyl group of the DAG-lactone structure (Scheme 1). The dansyl and diethylaminocoumarin were incorporated into the *sn*-2 carbonyl group. In comparison with the fluorescent spectra of these compounds, obtained in various solvents, compound 1 showed the most variation with solvent polarity (Figure 2). It is known that solvatochromic fluorophores show higher fluorescent intensity in hydrophobic environments and that the fluorophores are sensitive to the environment of a binding pocket in target proteins. The fluorescent intensity of labeled DAG-lactones should reflect its binding to a target protein, PKC δ (Figure 3), and as expected, our reporter compounds 1 and 2 containing solvatochromic fluorophores exhibited a remarkable increase of fluorescent intensity in the proximity of the C1b domain of PKC δ (δC1b), which is a specific target of DAG-lactone derivatives (Table 1 and Figure 4A). This indicates that in the binding of compounds 1 and 2 to the δC1b domain, the 6-methoxynaphthalene and dansyl moieties are located in the hydrophobic environment. However, for compound 3, the fluorescent intensity was decreased as the concentration of δC1b increased. The spectra in various solvents showed dependence of the fluorescent intensity of the compound 3 on solvent polarity, and it was suggested that in the binding of compound 3 to the δC1b domain, the diethylaminocoumarin moiety could be located in the hydrophilic environment. The changes of fluorescent intensity of the synthetic compounds are summarized in Table 1. The binding constants of fluorescent compounds were evaluated by $[^3\text{H}]\text{PDBu}$ competitive assay. Compound 1 showed the highest binding affinity for the δC1b domain. In the fluorescence-based analysis of binding, two factors are key to the sensitivity of the assay

system: (i) the fluorescent change as a function of the ligand binding and (ii) the binding affinity of a reporter compound.

Evaluation of Binding to PKC δ by Fluorescent Change of Compound 1. The results obtained supported the selection of compound 1 as a reporter compound for further study. In competitive assays, when candidate ligands of PKC δ are present, they replace the fluorescent compounds, which are formerly bound to δC1b , and the fluorescent intensity should decrease. Practically, addition of PDBu to the complex of compound 1 and δC1b resulted in a remarkable decrease of fluorescent intensity (Figure 4B). By utilization of compound 1 as a reporter, binding analysis of known PKC δ ligands was performed to assess the reliability of the assay system. Compounds 8–14 were prepared as test compounds (Figure 5A), and their IC_{50} values, determined by the fluorescence-based competitive assay with a fluorospectrophotometer against compound 1, were 0.60, 0.49, 1.95, 2.42, 10.0, 54.2, and $1.22 \times 10^4 \mu\text{M}$, respectively (Figure 5B). The order of the IC_{50} values of compounds 8–11 is identical to that of the K_i values (0.91, 0.26, 5.0, and 6.2 nM, respectively) determined by the $[^3\text{H}]\text{PDBu}$ competitive assay.

The Binding Analysis by Competition with Compound 1 Showed Linearity with the Classical RI Assay. To further assess the collinearity of the fluorescent assay on microplates and the $[^3\text{H}]\text{PDBu}$ assay, the inhibition percentages by compounds 8–21 in both assays (Figure 6A) at constant concentration were compared. The compound concentrations were set at $10 \mu\text{M}$ for the fluorescent assay and at 100 nM for the $[^3\text{H}]\text{PDBu}$ assay. As shown in Figure 6B, the inhibition percentages of compounds 8–21 are similar in both assays. Comparison of the IC_{50} and K_i of compounds 8, 9, 10, 11, and 17, possessing more than 60% inhibition percentages, showed acceptable linearity (Figure 6C).




Figure 7. The RI competition assay of positive hit compounds found in fluorescent screening utilizing compound 1. (A) Structures of test compounds obtained from screening. (B) Results of the inhibition assay based on RI methods. The concentration of compounds was fixed at $10\ \mu\text{M}$. The X axis indicates compound numbers. Columns show average inhibition obtained from triplicated experiments. Bars indicate standard errors.

In view of the above results, the present fluorescence-based evaluation of binding to the δC1b domain on microplates was deemed to be a promising alternative to the classical RI assay.

Expansion of the Fluorescence Evaluation to Screening of Library Compounds. As a proof-of-concept study, compound 1 was utilized in screening of a chemical library, supplied by the Screening Center at the Tokyo Medical and Dental University Screening Center. The structures of the library compounds were analyzed in advance of screening. For binding to the δC1b domain, pharmacophores including the carbonyl and hydroxy groups are necessary and only compounds including these functional groups were chosen. In the analysis of the library on microplates, compounds with more than 20% decrease of fluorescent intensity determined by triplicated assays were selected as

candidates for further study. Compounds 22–38 were found from 2560 library compounds (Figure 7A) as candidate δC1b ligands. To further assess the binding affinity of these compounds, evaluation of binding by competition with the $[^3\text{H}]\text{PDBu}$ was performed at a fixed concentration ($10\ \mu\text{M}$) (Figure 7B). In this analysis, compounds 30 and 26, with inhibition percentages at $10\ \mu\text{M}$ of 10.2% and of 9.4%, respectively, showed the most and the second most potent binding affinities for the δC1b domain. The binding affinities of these compounds are still very low compared to that of DAG-lactones, but an SAR study of these compounds could lead to novel structure templates for synthetic PKC ligands. This analysis also revealed that compounds 24, 25, 31, 32, 33, and 34 are false positive hits. The results obtained indicate that our screening system with a fluorescent

DAG-lactone derivative could be utilized for discovery of novel chemical leads for PKC δ .

CONCLUSION

A fluorescence-based method for chemical library screening has been developed. This method, which requires no washing steps, provides a linear response in relative binding affinity between compounds and is compatible with the classical RI assay for PKC ligands. By screening of more than 2500 library compounds, the method was proven to be a reliable means of discovery of compounds that bind to the C1b domain of PKC δ . The method would provide potent lead compounds that bind to the isozyme utilized in the assay. By utilization of the difference of the binding affinity of reporter compounds to different isozymes, it is possible that this screening system can provide an efficient selection of lead compounds highly specific to a target isozyme. This fluorescence-quenching method should be applicable to other known receptor–ligand combinations.

ASSOCIATED CONTENT

S Supporting Information. Synthesis procedures and NMR, HRMS, and IR data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

AUTHOR INFORMATION

Corresponding Author

*E-mail: tamamura.mr@tmd.ac.jp. Phone: +81-3-5280-8036. Fax: +81-3-5280-8039.

ACKNOWLEDGMENT

The authors thank Professor Masatoshi Hagiwara and Dr. Yukiko Okuno (Chemical Biology Screening Center, Tokyo Medical and Dental University) for their support in chemical library screening. This work was supported in part by Naito Foundation (W.N.) and Research Award of Graduate School of Biomedical Science, Tokyo Medical and Dental University (W.N. and T.I.).

REFERENCES

- (1) Loving, G. S., Sainlos, M., and Imperiali, B. (2010) Monitoring protein interactions and dynamics with solvatochromic fluorophores. *Trends Biotechnol.* 28, 73–83.
- (2) Rothman, D. M., Shults, M. D., and Imperiali, B. (2005) Chemical approaches for investigating phosphorylation in signal transduction networks. *Trends Cell Biol.* 15, 502–510.
- (3) Terai, T., and Nagano, T. (2008) Fluorescent probes for bioimaging applications. *Curr. Opin. Chem. Biol.* 12, 515–521.
- (4) Reymond, J. -L. (2008) Substrate arrays for fluorescence-based enzyme fingerprinting and high-throughput screening. *Ann. N.Y. Acad. Sci.* 1130, 12–20.
- (5) Nomura, W., Tanabe, Y., Tsutsumi, H., Tanaka, T., Ohba, K., Yamamoto, N., and Tamamura, H. (2008) Fluorophore labeling enables imaging and evaluation of specific CXCR4–ligand interaction at the cell membrane for fluorescence-based screening. *Bioconjugate Chem.* 19, 1917–1920.
- (6) Hayashida, O., Ogawa, N., and Uchiyama, M. (2007) Surface recognition and fluorescence sensing of histone by dansyl-appended cyclophane-based resorcinarene trimer. *J. Am. Chem. Soc.* 129, 13698–13705.
- (7) Tsutsumi, H., Nomura, W., Abe, S., Mino, T., Masuda, A., Ohashi, N., Tanaka, T., Ohba, K., Yamamoto, N., Akiyoshi, K., and Tamamura, H. (2009) Fluorogenically active leucine zipper peptides as tag-probe pairs for protein imaging in living cells. *Angew. Chem., Int. Ed.* 48, 9164–9166.
- (8) Nishizuka, Y. (1992) Intracellular signaling by hydrolysis of phospholipids and activation of protein kinase C. *Science* 258, 607–614.
- (9) Newton, A. C. (1995) Protein kinase C: structure, function, and regulation. *J. Biol. Chem.* 270, 28495–28498.
- (10) Watanabe, T., Ono, Y., Taniyama, Y., Hazama, K., Igarashi, K., Ogita, K., Kikkawa, U., and Nishizuka, Y. (1992) Cell division arrest induced by phorbol ester in CHO cells overexpressing protein kinase C-delta subspecies. *Proc. Natl. Acad. Sci. U.S.A.* 89, 10159–10163.
- (11) Mischak, H., Pierce, J. H., Goodnight, J., Kazanietz, M. G., Blumberg, P. M., and Mushinski, J. F. (1993) Phorbol ester-induced myeloid differentiation is mediated by protein kinase C-alpha and -delta and not by protein kinase C-beta II, -epsilon, -zeta, and -eta. *J. Biol. Chem.* 268, 20110–20115.
- (12) Li, C., Wernig, F., Leitges, M., Hu, Y., and Xu, Q. (2003) Mechanical stress-activated PKCdelta regulates smooth muscle cell migration. *FASEB J.* 17, 2106–2108.
- (13) Ghayur, T., Hugunin, M., Talanian, R. V., Ratnofsky, S., Quinlan, C., Emoto, Y., Pandey, P., Datta, R., Huang, Y., Kharbanda, S., Allen, H., Kamen, R., Wong, W., and Kufe, D. (1996) Proteolytic activation of protein kinase C delta by an ICE/CED 3-like protease induces characteristics of apoptosis. *J. Exp. Med.* 184, 2399–2404.
- (14) Alkon, D. L., Sun, M.-K., and Nelson, T. J. (2007) PKC signaling deficits: a mechanistic hypothesis for the origins of Alzheimer's disease. *Trends Pharmacol. Sci.* 28, 51–60.
- (15) Wang, Q. J. (2006) PKD at the crossroads of DAG and PKC signaling. *Trends Pharmacol. Sci.* 27, 317–323.
- (16) Marquez, V. E., and Blumberg, P. M. (2003) Synthetic diacylglycerols (DAG) and DAG-lactones as activators of protein kinase C (PK-C). *Acc. Chem. Res.* 36, 434–443.
- (17) Nacro, K., Bienfait, B., Lee, J., Han, K.-C., Kang, J.-H., Benzaria, S., Lewin, N. E., Bhattacharyya, D. K., Blumberg, P. M., and Marquez, V. E. (2000) Conformationally constrained analogues of diacylglycerol (DAG). 16. How much structural complexity is necessary for recognition and high binding affinity to protein kinase C?. *J. Med. Chem.* 43, 921–944.
- (18) Tamamura, H., Bienfait, B., Nacro, K., Lewin, N. E., Blumberg, P. M., and Marquez, V. E. (2000) Conformationally constrained analogues of diacylglycerol (DAG). 17. Contrast between *sn*-1 and *sn*-2 DAG lactones in binding to protein kinase C. *J. Med. Chem.* 43, 3209–3217.
- (19) Tamamura, H., Sigano, D. M., Lewin, N. E., Blumberg, P. M., and Marquez, V. E. (2004) Conformationally constrained analogues of diacylglycerol. 20. The search for an elusive binding site on protein kinase C through relocation of the carbonyl pharmacophore along the *sn*-1 side chain of 1,2-diacylglycerol lactones. *J. Med. Chem.* 47, 644–655.
- (20) Baba, Y., Ogoshi, Y., Hirai, G., Yanagisawa, T., Nagamatsu, K., Mayumi, S., Hashimoto, Y., and Sodeoka, M. (2004) Design, synthesis, and structure–activity relationship of new isobenzofuranone ligands of protein kinase C. *Bioorg. Med. Chem. Lett.* 14, 2963–2967.
- (21) Baba, Y., Mayumi, S., Hirai, G., Kawasaki, H., Ogoshi, Y., Yanagisawa, T., Hashimoto, Y., and Sodeoka, M. (2004) Evaluation of series of isobenzofuranone dimers as PKCalpha ligands: implication for the distance between the two ligand binding sites. *Bioorg. Med. Chem. Lett.* 14, 2969–2972.
- (22) Yanagita, R. C., Nakagawa, Y., Yamanaka, N., Kashiwagi, K., Saito, N., and Irie, K. (2008) Synthesis, conformational analysis, and biological evaluation of 1-hexylindolactam-V10 as a selective activator for novel protein kinase C isozymes. *J. Med. Chem.* 51, 46–56.
- (23) Nakagawa, Y., Yanagita, R. C., Hamada, N., Murakami, A., Takahashi, H., Saito, N., Nagai, H., and Irie, K. (2009) A simple analogue of tumor-promoting aplysiatoxin is an antineoplastic agent rather than a tumor promoter: development of a synthetically accessible protein kinase C activator with bryostatin-like activity. *J. Am. Chem. Soc.* 131, 7573–7579.

- (24) Martin, S. F., Josey, J. A., Wong, Y.-L., and Dean, D. W. (1994) General method for the synthesis of phospholipid derivatives of 1,2-*O*-diacyl-*sn*-glycerols. *J. Org. Chem.* *59*, 4805–4820.
- (25) Malolanarasimhan, K., Kedei, N., Sigano, D. M., Kelley, J. A., Lai, C. C., Lewin, N. E., Surawski, R. J., Pavlyukovets, V. A., Garfield, S. H., Wincovitch, S., Blumberg, P. M., and Marquez, V. E. (2007) Conformationally constrained analogues of diacylglycerol (DAG). 27. Modulation of membrane translocation of protein kinase C (PKC) isozymes alpha and delta by diacylglycerol lactones (DAG-lactones) containing rigid-rod acyl groups. *J. Med. Chem.* *50*, 962–978.
- (26) Nomura, W., Narumi, T., Ohashi, N., Serizawa, Y., Lewin, N. E., Blumberg, P. M., Furuta, T., and Tamamura, H. (2011) Synthetic caged DAG-lactones for photochemically-controlled activation of protein kinase C. *ChemBioChem* *12*, 535–539.
- (27) Mischak, H., Bodenteich, A., Kolch, W., Goodnight, J., Hofer, F., and Mushinski, J. F. (1991) Mouse protein kinase C-delta, the major isoform expressed in mouse hemopoietic cells: sequence of the cDNA, expression patterns, and characterization of the protein. *Biochemistry* *30*, 7925–7931.
- (28) Kazanietz, M. G., Krausz, K. W., and Blumberg, P. M. (1992) Differential irreversible insertion of protein kinase C into phospholipid vesicles by phorbol esters and related activators. *J. Biol. Chem.* *267*, 20878–20886.
- (29) List, B., Barbas, C. F., III, and Lerner, R. A. (1998) Aldol sensors for the rapid generation of tunable fluorescence by antibody catalysis. *Proc. Natl. Acad. Sci. U.S.A.* *95*, 15351–15355.
- (30) Davis, G. A. (1972) Dansylglycine as a fluorescent probe for aqueous solutions of cationic detergents. *J. Am. Chem. Soc.* *94*, 5089–5090.
- (31) Morii, T., Sugimoto, K., Makino, K., Otsuka, M., Imoto, K., and Mori, Y. (2002) A new fluorescent biosensor for inositol triphosphate. *J. Am. Chem. Soc.* *124*, 1138–1139.
- (32) Matlock, D. L., and Heyduk, T. (1999) A real-time fluorescence method to monitor the melting of duplex DNA during transcription initiation by RNA polymerase. *Anal. Biochem.* *270*, 140–147.
- (33) Kakio, A., Nishimoto, S., Yanagisawa, K., Kozutsumi, Y., and Matsuzaki, K. (2001) Cholesterol-dependent formation of GM1 ganglioside-bound amyloid beta-protein, an endogenous seed for Alzheimer amyloid. *J. Biol. Chem.* *276*, 24985–24990.
- (34) Timofeevski, S. L., Prusakiewicz, J. J., Rouzer, C. A., and Marnett, L. J. (2002) Isoform-selective interaction of cyclooxygenase-2 with indomethacin amides studied by real-time fluorescence, inhibition kinetics, and site-directed mutagenesis. *Biochemistry* *41*, 9654–9662.
- (35) Sakamoto, S., and Kudo, K. (2008) Supramolecular control of split-GFP reassembly by conjugation of beta-cyclodextrin and coumarin units. *J. Am. Chem. Soc.* *130*, 9574–9582.
- (36) Sharkey, N. A., and Blumberg, P. M. (1985) Highly lipophilic phorbol esters as inhibitors of specific [³H]phorbol 12,13-dibutyrate binding. *Cancer Res.* *45*, 19–24.
- (37) Ohashi, N., Nomura, W., Kato, M., Narumi, T., Lewin, N. E., Blumberg, P. M., and Tamamura, H. (2009) Synthesis of protein kinase Cdelta C1b domain by native chemical ligation methodology and characterization of its folding and ligand binding. *J. Pept. Sci.* *15*, 642–646.

Expert Opinion

1. Introduction
2. HIV fusion inhibitors such as Enfuvirtide
3. HIV co-receptor inhibitors such as Maraviroc
4. HIV integrase inhibitors such as Raltegravir
5. CD4 mimics as HIV entry inhibitors
6. Conclusion
7. Expert opinion

The successes and failures of HIV drug discovery

Chie Hashimoto, Tomohiro Tanaka, Tetsuo Narumi, Wataru Nomura* & Hirokazu Tamamura†

Institute of Biomaterials and Bioengineering, Tokyo Medical and Dental University, Chiyoda-ku, Tokyo, Japan

Introduction: To date, several anti-human immunodeficiency virus (HIV) drugs, including reverse transcriptase inhibitors and protease inhibitors, have been developed and used clinically for the treatment of patients infected with HIV. Recently, novel drugs have been discovered which have different mechanisms of action from those of the above inhibitors, including entry inhibitors and integrase (IN) inhibitors; the clinical use of three of these inhibitors has been approved. Other inhibitors are still in development.

Areas covered: This review article summarizes the history of the development of anti-HIV drugs and also focuses on successes in the development of these entry and IN inhibitors, along with looking at exploratory approaches for the development of other inhibitors.

Expert opinion: Currently used highly active antiretroviral therapy can be subject to a loss of efficacy, due to the emergence of multi-drug resistant (MDR) strains; a change of regimens of the drug combination is required to combat this, along with careful monitoring of the virus and CD4 in the blood, by methods such as cellular tropism testing. In such a situation, entry inhibitors such as CCR5/CXCR4 antagonists, CD4 mimics, fusion inhibitors and IN inhibitors might be optional agents for an expansion of the drug repertoire available to patients at all stages of HIV infection.

Keywords: AIDS, CCR5, CD4 mimic, chemokine receptor, CXCR4, fusion, HIV, integrase

Expert Opin. Drug Discov. (2011) 6(10):1067-1090

1. Introduction

The human immunodeficiency virus (HIV) is the cause of acquired immunodeficiency syndrome (AIDS), which was discovered by Montagnier and colleagues in 1983 [1]. HIV infects human host cells and destroys immune systems, subsequently causing immunodeficiency. To date, the number of people worldwide infected with HIV is certainly in excess of 30 million.

Several anti-HIV drugs have been reported in the last 25 years (Figure 1A). HIV is a retrovirus, which is an RNA virus that is replicated in a host cell via the enzyme reverse transcriptase to produce DNA from its RNA genome. This DNA is then incorporated into the host genome by an integrase (IN) enzyme. These inhibitors block the action of reverse transcriptase, and include nucleoside reverse transcriptase inhibitors (NRTIs) such as azidothymidine (AZT) and non-nucleoside reverse-transcriptase inhibitors (NNRTIs). The class of anti-HIV drugs that were initially approved for clinical use is reverse transcriptase inhibitors such as AZT [2]. The class of drugs that were subsequently approved is protease inhibitors, which prevent the cleavage of HIV precursor proteins into active proteins, a process that normally occurs in viral replication. This family of drugs includes Saquinavir (Invirase/Fortovase, Roche, Basel, Switzerland) and Ritonavir (Norvir, Abbott Laboratories, IL, USA), which have been used clinically in HIV/AIDS treatment. These drugs are usually administered as part of a two- or three-drug cocktail, accompanied by one or more

informa
healthcare

The successes and failures of HIV drug discovery

Article highlights.

- The highly active anti-retroviral therapy (HAART) involving use of reverse transcriptase inhibitors and protease inhibitors has led to great success in clinical treatment of human immunodeficiency virus (HIV)-infected patients.
- Brand-new drugs with different action mechanisms have been discovered to date.
- Enfuvirtide, a fusion inhibitor, Maraviroc, a co-receptor CCR5 antagonist and Raltegravir, an integrase (IN) inhibitor have successively been approved for clinical use.
- The potential of new inhibitors from novel drug categories such as entry inhibitors including CCR5/ CXCR4 antagonists and CD4 mimics, fusion inhibitors and IN inhibitors might be critical.
- Optional agents are valuable for an expansion of the drug repertoire available to patients because in case of loss of efficacy of HAART, change of regimens of the drug combination in HAART is required.

This box summarizes key points contained in the article.

NRTIs. Such cocktail therapies are known as highly active anti-retroviral therapy (HAART), which has brought great success and hope in the clinical treatment of HIV-infected patients [2]. HAART is capable of lowering the HIV level in the blood until it cannot be detected with current methods. Side effects associated with protease inhibitors include a lipodystrophy syndrome in which abnormal distribution of fat occurs and the face, arms and legs become thin but the therapy involves more serious clinical problems such as the emergence of multi-drug resistant (MDR) HIV-1 strains, and considerable expense. These drawbacks encouraged us to develop brand-new drugs with novel mechanisms of action.

Recently, the molecular mechanism of HIV-1 replication has been elucidated in detail. A dynamic supramolecular mechanism is associated with HIV entry/fusion steps. First, an HIV envelope protein gp120 interacts with a host-cell surface protein CD4, which causes gp120 to undergo a conformational change subsequently binding to the second cellular receptors, a chemokine receptor such as CCR5 [3-7] and CXCR4 [8]. The binding triggers the exposure of another envelope protein gp41 whose N-terminus penetrates the cell membrane. This is followed by the formation of the gp41 trimer-of-hairpins structure, which leads to fusion of HIV/cell membranes, completing the infection process [9]. The description of this dynamic molecular machinery has encouraged researchers to develop inhibitors which block the HIV entry/fusion steps targeting the receptors, CD4, CCR5 and CXCR4, and the viral protein gp41. In 2003, the Food and Drug Administration (FDA) approved Enfuvirtide (Fuzeon/T-20, Roche, Basel, Switzerland/Trimeris, Durham, NC, USA) as the first 'fusion inhibitor' for use in combination with other anti-HIV drugs to treat advanced HIV-1 infection [10]. In 2007, the FDA approved a CCR5

co-receptor antagonist, Maraviroc (Pfizer, New York, NY, USA), for use in combination with other anti-HIV drugs for the treatment of patients infected with CCR5-tropic HIV-1 [11]. In the same year, the FDA approved Raltegravir (Merck Sharp & Dohme Corp., NJ, USA) for use in combination with other antiretroviral agents in treatment-experienced adult patients who present with evidence of viral replication and HIV-1 strains resistant to multiple HAART agents [12,13]. Subsequently in 2009, the FDA granted expanded approval of Raltegravir for use in all patients. Numerous reviews exist concerning reverse transcriptase and protease inhibitors, and this review will focus on the success of a fusion inhibitor, Enfuvirtide, a CCR5 antagonist, Maraviroc and an IN inhibitor, Raltegravir, as well as the development of other anti-HIV agents including CXCR4 antagonists and CD4 mimics.

2. HIV fusion inhibitors such as Enfuvirtide

The binding of gp120 to CCR5 or CXCR4 triggers the formation of the trimer-of-hairpins structure of gp41 and subsequent fusion of the HIV/cell membranes, as described above. The trimer-of-hairpins structure is a six-helical bundle consisting of a central parallel trimer of the N-terminal helical region (HR1 region) surrounded by the C-terminal helical region (HR2 region) oriented in an antiparallel, hairpin fashion (Figure 1B) [9]. A subdomain is composed of two peptides, a 51-mer from the HR1 region and a 43-mer from the HR2 region, designated as N51 and C43, respectively [14]. There have been numerous reports that several HR2 region peptides inhibit bundle formation of six alpha-helices by the binding to the inner three-stranded coiled coils of the HR1 region thereby inhibiting membrane fusion (Figure 2A) [15]. An HR2 region peptide, C34, with 34 residues from the native sequence of gp41, has potent inhibitory activity against HIV-1 fusion [16]. In addition, a 36-residue peptide, T-20, which has the native sequence of gp41 and 24 residues in common with C34, shows potent anti-HIV activity, and its clinical use as the first entry/fusion inhibitor [10] in HIV/AIDS treatment was approved by the FDA in 2003 designated as Enfuvirtide. Enfuvirtide, in combination with other antiretroviral agents was approved for the treatment of advanced HIV-1 infection in adults and children aged 6 years or older with evidence of HIV-1 replication notwithstanding current therapy, and of resistance to current anti-HIV drugs. In view of the clinical use of Enfuvirtide, the dynamic supramolecular mechanism involving membrane fusion is a valid and rational target for inhibitors of HIV-1 replication. The success of Enfuvirtide has encouraged the development of entry/fusion inhibitors as a new class of anti-HIV drugs distinct from the first- and second-generation drugs. While reverse transcriptase inhibitors and protease inhibitors work internally in T cells to inhibit functions of viral enzymes, entry and fusion inhibitors work extracellularly, preventing HIV from invading cells. C34 has an exact interface that is capable of interacting with the inner three-stranded coiled coils of the gp41 HR1 region, compared with Enfuvirtide, which has a poorly delineated interface. However, a disadvantage




Figure 1. A. HIV-1 replication cycle and anti-HIV drugs that are effective at its various steps. **B.** Mechanisms of HIV-1 entry and fusion.

of C34 is its poor aqueous solubility and highly soluble C34 analogs, known as subcutaneous glucagon-like peptides (SC) peptides, were developed by artificial remodeling of C34 by Otaka *et al.* (Figure 2) [17]. In the design of SC peptides, the amino acid residues at the *a*, *d* and *e* positions of the helical wheel diagram of C34, which are essential for interaction with the inner coiled coils of HR1, were maintained with no substitution, while non-conserved residues at the *b*, *c*, *f* and *g* positions, which are located in the exterior region, were replaced by Glu or Lys. Several Glu-Lys side-chain ion charge pairs formed between *i* and *i* + 4 positions enhance solubility and alpha-helicity of the C34 analogs. The aqueous solubility of the SC peptides,

SC34 and SC34EK, is more than three orders of magnitude greater than that of the original C34 peptide. Anti-HIV activities of these SC peptides were superior or comparable with that of the original C34 peptide, and an order of magnitude greater than that of Enfuvirtide [18]. Furthermore, SC peptides were even effective against an Enfuvirtide-resistant strain. The C34 and SC peptide sequences lack the C-terminal lipid binding domain of Enfuvirtide and it has been suggested that SC peptides have a mechanism of action distinct from that of Enfuvirtide [19]. Thus, these SC peptides are leads for further refinement and clinical development. C34, T-649 [20], Enfuvirtide and SC peptides are all 34- to 36-mer peptides derived from the HR2 region of




Figure 2. A. Schematic representation of gp41 and sequences of HR2 region peptides. B. Helical wheel representation of the C34 peptide. Amino acid residues are numbered according to gp41 of NL4-3 strain. C. The design concept of introduction of the Glu-Lys motif to the solvent-accessible site. D. Remodeling of dynamic structures of HR1 regions leads to synthetic antigen molecules inducing neutralizing antibodies.




Figure 2. (continued). A. Schematic representation of gp41 and sequences of HR2 region peptides. B. Helical wheel representation of the C34 peptide. Amino acid residues are numbered according to gp41 of NL4-3 strain. C. The design concept of introduction of the Glu-Lys motif to the solvent-accessible site. D. Remodeling of dynamic structures of HR1 regions leads to synthetic antigen molecules inducing neutralizing antibodies.

The successes and failures of HIV drug discovery

gp41. T-1249 (Roche, Basel, Switzerland/Trimeris, Durham, NC, USA), a successor to Enfuvirtide [21], was much anticipated as its hydrophobic C-terminal sequence inhibits HIV-1 fusion by interacting with lipid bilayers in a manner similar to the interaction enjoyed by Enfuvirtide, but its clinical trial was discontinued in January 2005 because of formulation problems. In addition, attempts have been made to develop small non-peptide inhibitors that block gp41 activation [22-24]. To date, development of highly potent and useful low molecular weight inhibitors has been difficult, although non-natural binding elements that contribute to the formation of a stable complex with the inner coiled coils have been identified using a biased combinatorial chemistry library [25]. The development of small organic compounds as useful fusion inhibitors is continuing.

The HR1 region is critical for the development of AIDS vaccine antigens because the six-helical bundle structure is formed by interactive approach to the central trimer of the HR1 regions by the HR2 regions of three strands coiled coils. In the design of artificial antigens that induce broadly neutralizing antibodies, a useful strategy is to synthesize molecules that mimic the natural trimer on the virion surface. Using an original template with C3-symmetric linkers, we designed and synthesized a novel three-helical bundle mimetic corresponding to the trimeric form of N36 [26] (Figure 2D). The antiserum produced by immunization of the N36 trimeric form antigen in mice showed structural preference for binding to the N36 trimer and more potent neutralizing activity against HIV-1 infection than the N36 monomer. The exposed timing of epitopes was limited during HIV-1 entry, and carbohydrates, which could disturb accession of antibodies to epitopes, were not included in the amino acid sequences of the native protein [27]. These two advantages in the design based on the HR1 region of gp41 could further enhance the potential of a vaccine design based on the HR1 region.

3. HIV co-receptor inhibitors such as Maraviroc

Interaction of CD4 with the HIV envelope protein gp120 causes a conformational change in the latter and its subsequent binding to the second cellular receptors, CCR5 [3-7] and CXCR4 [8] as described above, in Section 1. Macrophage-tropic (R5) HIV-1 strains, which constitute a majority in the early stage of HIV infection, use CCR5 as a co-receptor, whereas T cell line-tropic (X4) HIV-1 strains, which are the major species in the late stage of HIV infection and AIDS, use CXCR4 as a co-receptor. CCR5 and CXCR4 play physiological roles as the receptors for endogenous ligands, or chemokines. Several chemokine antagonists of CCR5 and CXCR4 have been developed as entry inhibitors. The validity of development of CCR5 antagonists is supported by the finding that individuals with the CCR5-Δ32 deletion mutation are healthy and strongly resistant to HIV-1 infection [28]. To date, several pharmaceutical companies have investigated novel CCR5 antagonists with suitable pharmaceutical properties. One CCR5-selective antagonist,

Maraviroc (1) is the first CCR5 antagonist to be approved by the FDA as described above (Figure 3A) [11] and is used for the treatment of patients infected with CCR5-tropic HIV-1. Several other CCR5 antagonists are currently in clinical trials. Maraviroc has relatively high oral bioavailability (23%) with food effects in humans [29,30]. During the process of further improvement of Maraviroc's pharmacological profile, compound 2 was found to have potent fusion inhibitory activity ($IC_{50} < 0.1$ nM) and a pharmacological profile identical to that of Maraviroc [31]. Compound 2 showed high anti-HIV activity even against a Maraviroc-resistant mutant. Thus, compound 2 might be a desirable second-generation CCR5 antagonist. Takeda Pharmaceutical Co. Ltd., Osaka, Japan developed TAK-779 (3), a CCR5 antagonist with minimal bioavailability for i.v. use [32,33] but clinical development was discontinued because of local reactions at s.c. injection sites. High throughput screening has been used to develop an orally bioavailable CCR5 antagonist, and has led to the discovery of a novel lead compound 4 with a scaffold structure different from that of TAK-779 [34]. Subsequent optimization resulted in the development of TAK-220, a new CCR5 antagonist with a piperidine-4-carboxamide structure (5) (Takeda Pharmaceutical Co. Ltd., Osaka, Japan/Tobira Therapeutics, Inc.), which has high CCR5 binding activity and resistance to metabolic modification [35]. Merck/Schering-Plough Corp., which became Merck & Co., NJ, USA in a merger, reported a piperidinopiperazine series such as Vicriviroc (SCH-D/SCH417690) (6) [36]. In its Phase III studies, the safety and efficacy of Vicriviroc in the addition to optimized or HAART regimens was assessed based on established criteria. Vicriviroc, however, failed to show that its addition to the current regimens for the further treatment of HIV-1 infectious patients with evidence of HIV-1 replication is more effective than background regimens. As a result, Merck & Co. suspended its support of the FDA approval for Vicriviroc in treatment-experienced subjects [37,38]. A CCR5 antagonist with a spirodiketopiperazine scaffold, ONO-4128/873140 (7) (GSK, Middlesex, UK/Ono Pharmaceutical Co. Ltd., Osaka, Japan), has been developed by combinatorial chemistry utilizing solid-phase techniques [39]. ONO-4128/873140 has CCR5 binding activity in the nanomolar range and potent Ca^{2+} mobilization inhibitory activity. The spirodiketopiperazine is an attractive scaffold because it is likely to lead to more diverse derivatives. ONO-4128/873140 was advanced into the Phase II studies, but not into Phase III studies on account of its hepatotoxicity.

The other co-receptor for HIV-1 entry is CXCR4 [8], which is also relevant to mediation of the metastasis of a variety of cancer cells [40-42], leukemia [43,44] and rheumatoid arthritis [45,46]. Thus, CXCR4 is an important target for drug discovery, because CXCR4 antagonists might overcome these diseases. To date, several CXCR4 antagonists, peptidic and non-peptidic have been developed.

A 14-mer peptide T140 (8) derived from polyphemusin II, has been reported to be a potent CXCR4 antagonist (Figure 3B) [47] by us (Kyoto University, Kyoto, Japan). Biostable T140 analogs [48,49] (Biokine Therapeutics Ltd.,





Figure 3. A. Structures of CCR5 antagonists. **B.** Structures of CXCR4 antagonists. **C.** Development of non-peptidic CXCR4 antagonists. **D.** Structures of bivalent CXCR4 ligands. A maximum increase in binding affinity for CXCR4 was observed in (**25**), $n = 20$ and (**26**), $m = 12$.

Rehovot, Israel) have significant inhibitory activity not only against HIV infection *in vitro* but also against tumor metastasis *in vivo* [41,42]. FC131 (**9**) was developed by downsizing of **8** [50].


AMD3100 (**10**) (Genzyme Corp., Cambridge, MA, USA), a non-peptidic bicyclam-containing small molecular CXCR4 antagonist, was the first CXCR4 antagonist to enter clinical trials for the treatment of HIV-1-infected patients (Figure 3B) [51,52]. It was found, however, to have adverse cardiovascular effects and its use as an anti-AIDS drug was discontinued. AMD3100 has two cyclam moieties tethered by *p*-xylene templates. Teixidó and colleagues have constructed

combinatorial libraries based on the structure of AMD3100 and containing: 1) at least two nitrogen atoms on each side of the *p*-xylene template; and 2) separation between these nitrogen atoms similar to that in cyclam [53]. As a result, the non-cyclam compound **11** with potent anti-HIV activity was found. Liotta and colleagues have screened various compounds in which two basic centers (e.g., guanidine, hydrazone or pyridine groups) were connected by a phenyl-containing bridge. A bisymmetric pyridine-containing compound **12** was found to have 10-nM potency [54] but further preclinical studies showed that compound **12** failed to exhibit any *in vivo*


The successes and failures of HIV drug discovery



TAK-220 5



Vicriviroc/SCH-D (SCH417690) 6



ONO-4128/873140 7

Figure 3. (continued). A. Structures of CCR5 antagonists. B. Structures of CXCR4 antagonists. C. Development of non-peptidic CXCR4 antagonists. D. Structures of bivalent CXCR4 ligands. A maximum increase in binding affinity for CXCR4 was observed in (25), $n = 20$ and (26), $m = 12$.

efficacy due to poor biostability. On the hypothesis that a poor pharmacokinetic profile of compound 12 might be a result of rapid oxidative metabolism, further structure-activity relationship studies of compound 12 were conducted and led to the development of compound 13 with a nanomolar level of CXCR4 antagonistic activity [55]. To develop non-

cyclam CXCR4 antagonists, De Clercq and colleagues have carried out structure-activity relationship studies on AMD3100 (10) [56,57]. First, according to the hypothesis that both rings of 10 are not essential structural requirements, they synthesized and evaluated single cyclam analogs containing an aromatic ring instead of another cyclam ring. The




Figure 3. (continued). A. Structures of CCR5 antagonists. B. Structures of CXCR4 antagonists. C. Development of non-peptidic CXCR4 antagonists. D. Structures of bivalent CXCR4 ligands. A maximum increase in binding affinity for CXCR4 was observed in (25), $n = 20$ and (26), $m = 12$.

The successes and failures of HIV drug discovery




Figure 3. (continued). A. Structures of CCR5 antagonists. B. Structures of CXCR4 antagonists. C. Development of non-peptidic CXCR4 antagonists. D. Structures of bivalent CXCR4 ligands. A maximum increase in binding affinity for CXCR4 was observed in (25), $n = 20$ and (26), $m = 12$.