

8.36 mmol) in CH_2Cl_2 (15 ml) were added trichloroacetonitrile (2.79 ml, 27.8 mmol) and Cs_2CO_3 (908 mg, 2.73 mmol) at 0 °C. After being stirred for 2 h, the reaction mixture was filtered through a Celite pad. The filtrate was concentrated in vacuo to afford glycosyl imidate **24**, which was used to the next reaction without further purification.

3.22. Methyl [2-O-azido-6-O-tert-butylidiphenylsilyl-2-O-deoxy-4-O-levulinyl-3-O-(4-methoxybenzyl)- α -D-glucopyranosyl]-(1 \rightarrow 4)-(tert-butylidimethylsilyl 2-O-benzoyl-3-O-benzyl- β -D-glucopyranosid)uronate (25)

The glycosyl donor **24** (8.38 mmol), glycosyl acceptor **8** (2.85 g, 5.51 mmol), and MS4A powder (7.0 g) were then suspended in anhydrous toluene (70 ml) under Ar. After being stirred for 2 h, the mixture was cooled to -20 °C. TBDMSOTf (506 μ l, 2.20 mmol) was added at the same temperature. The reaction mixture was stirred and gradually warmed to 0 °C. After 5 h, the reaction was quenched by the addition of water. The resulting mixture was filtered through a Celite pad. The filtrate was washed with water, satd NaHCO_3 aq, and brine, dried over Na_2SO_4 , filtered, and concentrated in vacuo. Flash chromatography (silica gel: 300 g, toluene/AcOEt=100:1 \rightarrow 20:1) afforded compound **25** as a brown oil. Yield 5.29 g (83%). ^1H NMR (600 MHz, CDCl_3) δ 8.04 (2H, d, $J=8.1$ Hz, Bz), 7.65–7.19 (20H, m, aromatic), 6.88 (2H, d, $J=8.8$ Hz, MPM), 5.51 (1H, d, $J_{1,2'}=3.4$ Hz, H-1'), 5.36–5.31 (2H, m, H-2, H-4'), 4.87 (1H, d, $J_{1,2}=7.4$ Hz, H-1), 4.79 and 4.72 (each 1H, d, $J_{\text{gem}}=10.8$ Hz, PhCH_2), 4.72 and 4.61 (each 1H, d, $J_{\text{gem}}=10.8$ Hz, PhCH_2), 4.28 (1H, dd, $J_{4,3}=8.8$ Hz, $J_{4,5}=9.5$ Hz, H-4), 4.07 (1H, d, $J_{5,4}=9.5$ Hz, H-5), 4.01 (1H, dd, $J_{3,2}=8.8$ Hz, $J_{3,4}=8.8$ Hz, H-3), 3.90 (1H, dd, $J_{3,2'}=10.2$ Hz, $J_{3,4'}=9.5$ Hz, H-3'), 3.79 (3H, s, OCH_3), 3.69 (1H, dd, $J_{6a',5'}=1.3$ Hz, $J_{6a',6b'}=11.8$ Hz, H-6a'), 3.64 (1H, dd, $J_{6b',5'}=1.3$ Hz, $J_{6b',6a'}=11.8$ Hz, H-6b'), 3.56–3.53 (4H, m, OCH_3 , H-5'), 3.34 (1H, dd, $J_{2,1'}=3.4$ Hz, $J_{2,3'}=10.2$ Hz, H-2'), 2.66–2.63 (2H, m, $\text{CH}_3\text{C}(=\text{O})\text{C}_2\text{H}_4\text{C}(=\text{O})-$), 2.42–2.36 (2H, m, $\text{CH}_3\text{C}(=\text{O})\text{C}_2\text{H}_4\text{C}(=\text{O})-$), 2.17 (3H, s, $\text{CH}_3\text{C}(=\text{O})\text{C}_2\text{H}_4\text{C}(=\text{O})-$), 1.02 (9H, s, $\text{C}(\text{CH}_3)_3$), 0.74 (9H, s, $\text{C}(\text{CH}_3)_3$), 0.06 (3H, s, CH_3), 0.00 (3H, s, CH_3); ^{13}C NMR (150 MHz, CDCl_3) δ 206.3, 168.4, 163.7, 135.7, 135.6, 135.6, 129.8, 129.6, 129.6, 129.6, 129.5, 128.4, 128.3, 127.8, 127.8, 127.7, 127.6, 127.5, 113.7, 97.3, 95.9, 82.2, 77.3, 74.8, 74.7, 74.3, 74.1, 73.7, 70.8, 69.6, 62.8, 60.7, 55.2, 52.4, 37.7, 29.8, 27.8, 26.6, 25.2, -4.4, -5.4; HRMS (positive mode) found: m/z 1182.4793 $[\text{M}+\text{Na}]^+$, calcd for $\text{C}_{62}\text{H}_{77}\text{N}_3\text{O}_{15}\text{Si}_2\text{Na}$: 1182.4791.

3.23. Methyl [2-O-azido-6-O-tert-butylidiphenylsilyl-2-O-deoxy-4-O-levulinyl-3-O-(4-methoxybenzyl)- α -D-glucopyranosyl]-(1 \rightarrow 4)-(tert-butylidimethylsilyl 2-O-benzoyl-3-O-benzyl- β -D-idopyranosid)uronate (26)

The glycosyl imidate **24** (0.375 mmol), glycosyl acceptor **15** (97.1 mg, 0.187 mmol), and MS4A powder (800 mg) were then suspended in anhydrous CH_2Cl_2 (8.0 ml) under Ar. After being stirred for 3 h, the mixture was cooled to -20 °C. TBDMSOTf (21.5 μ l, 93.7 mmol) was added at the same temperature. The reaction mixture was stirred and gradually warmed to room temperature. After 18 h, the reaction was quenched by the addition of water. The resulting mixture was filtered through a Celite pad. The filtrate was washed with water, and brine, dried over Na_2SO_4 , filtered, and concentrated in vacuo. Flash chromatography (silica gel: 20 g, toluene/AcOEt=100:1 \rightarrow 60:1) afforded compound **26** as a brown oil. Yield 169 mg (77%). ^1H NMR (600 MHz, CDCl_3) δ 8.21 (2H, d, $J=7.4$ Hz, Bz), 7.66 (18H, m, aromatic), 7.02 (2H, d, $J=8.8$ Hz, MPM), 6.80 (2H, d, $J=8.8$ Hz, MPM), 5.28 (1H, dd, $J_{4',3'}=9.5$ Hz, $J_{4',5'}=9.5$ Hz, H-4'), 5.19 (1H, s, H-1), 5.09 (1H, s, H-2), 4.84 and 4.76 (each 1H, d, $J_{\text{gem}}=11.5$ Hz, PhCH_2), 4.68 (1H, d,

$J_{1,2'}=3.4$ Hz, H-1'), 4.51 (1H, d, $J_{5,4}=2.0$ Hz, H-5), 4.30 (1H, d, $J_{3,4}=3.4$ Hz, H-3), 3.92–3.90 (3H, m, H-4, PhCH_2), 3.79–3.75 (4H, m, H-5', OCH_3), 3.67–3.66 (2H, m, H-6a', H-6b'), 3.55 (1H, dd, $J_{3',2'}=9.5$ Hz, $J_{3',4'}=9.5$ Hz, H-3'), 3.45 (3H, s, OCH_3), 3.31 (1H, dd, $J_{2,1'}=3.4$ Hz, $J_{2,3'}=9.5$ Hz, H-2'), 2.63–2.61 (2H, m, $\text{CH}_3\text{C}(=\text{O})\text{C}_2\text{H}_4\text{C}(=\text{O})-$), 2.39–2.37 (2H, m, $\text{CH}_3\text{C}(=\text{O})\text{C}_2\text{H}_4\text{C}(=\text{O})-$), 2.17 (3H, s, $\text{CH}_3\text{C}(=\text{O})\text{C}_2\text{H}_4\text{C}(=\text{O})-$), 1.00 (9H, s, $\text{C}(\text{CH}_3)_3$), 0.80 (9H, s, $\text{C}(\text{CH}_3)_3$), 0.14 (3H, s, CH_3), 0.06 (3H, s, CH_3); ^{13}C NMR (150 MHz, CDCl_3) δ 206.1, 170.7, 168.3, 166.5, 135.8, 135.7, 130.1, 129.6, 129.5, 129.5, 129.0, 128.5, 128.4, 128.2, 128.0, 127.5, 127.4, 113.6, 99.2, 93.4, 77.9, 75.1, 74.1, 73.7, 73.7, 73.0, 71.1, 69.7, 68.9, 63.4, 61.5, 55.2, 51.8, 37.7, 30.9, 29.8, 29.2, 27.8, 26.6, 25.5, -4.0, -5.3; HRMS (positive mode) found: m/z 1182.4759 $[\text{M}+\text{Na}]^+$, calcd for $\text{C}_{62}\text{H}_{77}\text{N}_3\text{O}_{15}\text{Si}_2\text{Na}$: 1182.4791.

3.24. Methyl [2-O-azido-6-O-tert-butylidiphenylsilyl-2-O-deoxy-4-O-levulinyl-3-O-(4-methoxybenzyl)- α -D-glucopyranosyl]-(1 \rightarrow 4)-(2-O-benzoyl-3-O-benzyl- β -D-glucopyranosyl)uronate (27)

To a solution of compound **25** (55.7 mg, 48.0 μ mol) in THF (1 ml) were added AcOH (5.5 μ l, 96.0 μ mol) and 1 M TBAF in THF (48.0 μ l, 48.0 μ mol) at room temperature. After being stirred for 12 h, the reaction mixture was poured into AcOEt. The resulting mixture was washed with water and brine, dried over Na_2SO_4 , filtered, and concentrated in vacuo. Flash chromatography (silica gel: 5 g, toluene/EtOAc=10:1 \rightarrow 8:1) afforded compound **27** as a colorless oil. Yield 39.5 mg (78%, $\alpha/\beta=5:1$). ^1H NMR (600 MHz, CDCl_3) δ for α -anomer 8.11 (2H, d, $J=7.4$ Hz, Bz), 7.66–7.19 (20H, m, aromatic), 6.86 (2H, d, $J=8.8$ Hz, MPM), 5.65 (1H, dd, $J_{1,2}=3.4$ Hz, $J_{1,\text{OH}}=5.4$ Hz, H-1), 5.37 (1H, d, $J_{1,2'}=3.4$ Hz, H-1'), 5.26 (1H, dd, $J_{4',5'}=10.2$ Hz, $J_{4',3'}=9.5$ Hz, H-4'), 5.16 (1H, dd, $J_{2,3}=8.1$ Hz, $J_{2,1}=3.4$ Hz, H-2), 4.85 (2H, s, PhCH_2), 4.70 (1H, d, $J_{5,4}=7.1$ Hz, H-5), 4.46 and 4.39 (each 1H, d, $J_{\text{gem}}=10.2$ Hz, PhCH_2), 4.37 (1H, dd, $J_{3,4}=7.4$ Hz, $J_{3,2}=8.1$ Hz, H-3), 4.19 (1H, dd, $J_{4,5}=7.1$ Hz, $J_{4,3}=7.4$ Hz, H-4), 3.81–3.77 (4H, m, H-3', OCH_3), 3.69–3.62 (3H, m, H-5', H-6a', H-6b'), 3.54 (3H, s, OCH_3), 3.32 (1H, dd, $J_{2,3'}=10.2$ Hz, $J_{2,1'}=3.4$ Hz, H-2'), 3.23 (1H, d, $J_{\text{OH},1}=5.4$ Hz, OH), 2.65–2.62 (2H, m, $\text{CH}_3\text{C}(=\text{O})\text{C}_2\text{H}_4\text{C}(=\text{O})-$), 2.42–2.39 (2H, m, $\text{CH}_3\text{C}(=\text{O})\text{C}_2\text{H}_4\text{C}(=\text{O})-$), 2.16 (3H, s, $\text{CH}_3\text{C}(=\text{O})\text{C}_2\text{H}_4\text{C}(=\text{O})-$), 1.01 (9H, s, $\text{C}(\text{CH}_3)_3$); ^{13}C NMR (150 MHz, CDCl_3) δ 206.1, 170.9, 169.2, 159.2, 135.7, 135.6, 130.1, 129.8, 129.6, 129.6, 129.5, 129.3, 128.6, 128.4, 128.4, 127.8, 127.7, 127.5, 127.5, 113.7, 98.3, 89.9, 77.3, 75.1, 74.3, 74.1, 73.3, 72.4, 71.5, 71.0, 70.0, 62.8, 61.7, 55.2, 52.4, 37.7, 29.8, 27.8, 26.6; HRMS (positive mode) found: m/z 1068.3932 $[\text{M}+\text{Na}]^+$, calcd for $\text{C}_{56}\text{H}_{63}\text{N}_3\text{O}_{15}\text{SiNa}$: 1068.3926.

3.25. Benzyl (2-O-azido-6-O-tert-butylidiphenylsilyl-2-O-deoxy-4-O-levulinyl-3-O-(4-methoxybenzyl)- α -D-glucopyranosyl]-(1 \rightarrow 4)-(methyl 2-O-benzoyl-3-O-benzyl- β -D-glucopyranosyluronate)-(1 \rightarrow 6)-2,3,4-tetra-O-benzyl- β -D-glucopyranoside (30)

To a solution of **27** (54.1 mg, 51.7 μ mol) in CH_2Cl_2 (2 ml) were added trichloroacetonitrile (51.8 μ l, 517 mmol) and Cs_2CO_3 (16.8 mg, 51.7 μ mol) at room temperature. After being stirred for 1 h, the reaction mixture was filtered through Celite pad. The filtrate was concentrated in vacuo to give glycosyl imidate **28**. The glycosyl imidate **28** (51.7 μ mol), glucose moiety **29** (55.9 mg, 0.103 mmol), and MS4A powder (400 mg) were then suspended in anhydrous CH_2Cl_2 (4 ml) under Ar. After being stirred for 2.5 h, the mixture was cooled to -78 °C. TMSOTf (5.74 μ l, 25.8 μ mol) was then added at the same temperature. After 23 h, the reaction was quenched by the addition of water. The resulting mixture was filtered through a Celite pad. The filtrate

was washed with water, and brine, dried over Na_2SO_4 , filtered, and concentrated in vacuo. Flash chromatography (silica gel: 5 g, toluene/AcOEt=50:1 \rightarrow 10:1) afforded compound **30** as a colorless oil. Yield 54.7 mg (68%, two steps). ^1H NMR (600 MHz, CDCl_3) δ 7.98 (2H, d, $J=6.7$ Hz, Bz), 7.65–7.13 (40H, m, aromatic), 6.88 (2H, d, $J=8.8$ Hz, MPM), 5.46 (1H, d, $J_{1',2'}=3.4$ Hz, H-1'), 5.43 (1H, dd, $J_{2',3'}=8.8$ Hz, $J_{2',1'}=7.4$ Hz, H-2'), 5.35 (1H, dd, $J_{4',5'}=9.5$ Hz, $J_{4',3'}=9.5$ Hz, H-4'), 4.87 and 4.69 (each 1H, d, $J_{\text{gem}}=10.8$ Hz, PhCH_2), 4.86 and 4.68 (each 1H, d, $J_{\text{gem}}=12.2$ Hz, PhCH_2), 4.79 and 4.70 (each 1H, d, $J_{\text{gem}}=10.2$ Hz, PhCH_2), 4.73 (1H, d, $J_{1',2'}=7.4$ Hz, H-1'), 4.69 and 4.64 (each 1H, d, $J_{\text{gem}}=10.8$ Hz, PhCH_2), 4.65 and 4.41 (each 1H, d, $J_{\text{gem}}=12.2$ Hz, PhCH_2), 4.60 and 4.44 (each 1H, d, $J_{\text{gem}}=10.8$ Hz, PhCH_2), 4.36 (1H, d, $J_{1,2}=7.4$ Hz, H-1), 4.26 (1H, dd, $J_{4',5'}=9.5$ Hz, $J_{4',3'}=8.8$ Hz, H-4'), 4.10 (1H, dd, $J_{6a,6b}=10.8$ Hz, $J_{6a,5}=1.3$ Hz, H-6a), 4.05 (1H, d, $J_{5',4'}=9.5$ Hz, H-5'), 3.97 (1H, dd, $J_{3',4'}=8.8$ Hz, $J_{3',2'}=8.8$ Hz, H-3'), 3.90 (1H, dd, $J_{3'',4''}=9.5$ Hz, $J_{3'',2''}=9.5$ Hz, H-3''), 3.78 (3H, s, OCH_3), 3.69 (1H, dd, $J_{6a'',6b''}=11.5$ Hz, $J_{6a'',5''}=1.3$ Hz, H-6a''), 3.64 (1H, dd, $J_{6b'',6a''}=11.5$ Hz, $J_{6b'',5''}=2.7$ Hz, H-6b''), 3.59–3.52 (6H, m, H-3, H-6b, H-5'', OCH_3), 3.44–3.41 (1H, m, H-5), 3.40 (1H, dd, $J_{2,3}=8.1$ Hz, $J_{2,1}=7.4$ Hz, H-2), 3.34–3.30 (2H, m, H-4, H-2'), 2.66–2.63 (2H, m, $\text{CH}_3\text{C}(=\text{O})\text{C}_2\text{H}_4\text{C}(=\text{O})-$), 2.42–2.39 (2H, m, $\text{CH}_3\text{C}(=\text{O})\text{C}_2\text{H}_4\text{C}(=\text{O})-$), 2.16 (3H, s, $\text{CH}_3\text{C}(=\text{O})\text{C}_2\text{H}_4\text{C}(=\text{O})-$), 1.01 (9H, s, $\text{C}(\text{CH}_3)_3$); ^{13}C NMR (150 MHz, CDCl_3) δ 206.1, 168.3, 164.8, 159.3, 138.4, 135.7, 135.7, 129.8, 129.6, 129.6, 129.0, 128.4, 128.3, 128.3, 128.2, 128.2, 128.1, 127.9, 127.8, 127.8, 127.7, 127.7, 127.6, 127.5, 125.2, 113.7, 102.1, 101.0, 97.4, 84.4, 82.1, 82.0, 77.7, 77.2, 75.6, 74.8, 74.7, 74.7, 74.5, 74.4, 74.1, 73.9, 72.9, 70.9, 70.7, 69.8, 68.3, 62.8, 61.4, 55.2, 52.4, 37.7, 29.8, 27.8, 26.7; HRMS (positive mode) found: m/z 1590.6341 $[\text{M}+\text{Na}]^+$, calcd for $\text{C}_{90}\text{H}_{97}\text{N}_3\text{O}_{20}\text{SiNa}$: 1590.6332.

3.26. Methyl [2-O-azido-6-O-tert-butylidiphenylsilyl-2-O-deoxy-4-O-levulinyl-3-O-(4-methoxybenzyl)- α -D-glucopyranosyl]-(1 \rightarrow 4)-(2-O-benzoyl-3-O-benzyl- β -D-glucopyranosyl)uronate (**31**)

To a solution of compound **26** (19.9 mg, 17.1 μmol) in pyridine (1 ml) was added HF \cdot pyridine (44.4 μl , 1.71 mmol) at room temperature. After being stirred for 2 h, the reaction mixture was poured into AcOEt. The resulting mixture was washed with water and brine, dried over Na_2SO_4 , filtered, and concentrated in vacuo. Flash chromatography (silica gel: 5 g, toluene/EtOAc=10:1 \rightarrow 3:1) afforded compound **31** as a colorless oil. Yield 13.1 mg (73%, $\alpha/\beta=5:3$). ^1H NMR (600 MHz, CDCl_3) δ for α -anomer 8.18 (2H, d, $J=7.4$ Hz, Bz), 7.65–7.17 (18H, m, aromatic), 7.04 (2H, d, $J=8.8$ Hz, MPM), 6.82 (2H, d, $J=8.8$ Hz, MPM), 5.49 (1H, d, $J_{1,1-\text{OH}}=8.8$ Hz, H-1), 5.05 (1H, d, $J_{2,3}=2.7$ Hz, H-2), 4.89 (1H, d, $J_{5,4}=2.7$ Hz, H-5), 4.87 (1H, d, $J_{\text{gem}}=11.5$ Hz, PhCH_2), 4.77 (1H, d, $J_{\text{gem}}=11.5$ Hz, PhCH_2), 4.68 (1H, d, $J_{1',2'}=3.4$ Hz, H-1'), 4.34 (1H, dd, $J_{3,2}=2.7$ Hz, $J_{3,4}=3.4$ Hz, H-3), 4.09 (1H, d, $J_{1-\text{OH},1}=8.8$ Hz, 1-OH), 4.01 (1H, dd, $J_{4,3}=3.4$ Hz, $J_{4,5}=2.7$ Hz, H-4), 3.96 (1H, d, $J_{\text{gem}}=10.8$ Hz, PhCH_2), 3.83 (1H, d, $J_{\text{gem}}=10.8$ Hz, PhCH_2), 3.77–3.75 (4H, m, H-6a', OCH_3), 3.72–3.68 (2H, m, H-5', H-6b'), 3.50 (3H, s, OCH_3), 3.29 (1H, dd, $J_{2',1'}=3.4$ Hz, $J_{2',3'}=7.4$ Hz, H-2'), 2.66–2.59 (2H, m, $\text{CH}_3\text{C}(=\text{O})\text{C}_2\text{H}_4\text{C}(=\text{O})-$), 2.42–2.31 (2H, m, $\text{CH}_3\text{C}(=\text{O})\text{C}_2\text{H}_4\text{C}(=\text{O})-$), 2.16 (3H, s, $\text{CH}_3\text{C}(=\text{O})\text{C}_2\text{H}_4\text{C}(=\text{O})-$), 1.00 (9H, s, $\text{C}(\text{CH}_3)_3$); ^{13}C NMR (150 MHz, CDCl_3) δ for β -anomer 206.0, 170.7, 165.8, 159.2, 135.7, 135.7, 133.2, 130.0, 129.7, 129.6, 129.5, 128.6, 128.5, 128.5, 128.3, 127.5, 127.4, 127.4, 113.6, 99.2, 93.6, 77.9, 74.7, 74.0, 73.6, 72.5, 71.2, 69.7, 68.2, 67.4, 63.2, 61.4, 55.2, 52.1, 37.7, 29.8, 27.8, 26.6; HRMS (positive mode) found: m/z 1068.3912 $[\text{M}+\text{Na}]^+$, calcd for $\text{C}_{56}\text{H}_{63}\text{N}_3\text{O}_{15}\text{SiNa}$: 1068.3926.

3.27. Benzyl (2-O-azido-6-O-tert-butylidiphenylsilyl-2-O-deoxy-4-O-levulinyl-3-O-(4-methoxybenzyl)- α -D-glucopyranosyl)-(1 \rightarrow 4)-(methyl 2-O-benzoyl-3-O-benzyl- β -D-glucopyranosyluronate)-(1 \rightarrow 6)-2,3,4-tetra-O-benzyl- β -D-glucopyranoside (**33**)

To a solution of **31** (112 mg, 107 μmol) in CH_2Cl_2 (8 ml) were added trichloroacetonitrile (107 μl , 1.07 mmol) and Cs_2CO_3 (34.8 mg, 107 μmol) at 0 $^\circ\text{C}$. The reaction mixture was stirred and gradually warmed to room temperature. After being stirred for 6 h, the reaction mixture was filtered through Celite pad. The filtrate was concentrated in vacuo to give glycosyl imidate **32**. The glycosyl imidate **32** (107 μmol), glucose moiety **29** (116 mg, 214 mmol), and MS4A powder (1.0 g) were then suspended in anhydrous CH_2Cl_2 (10 ml) under Ar. After being stirred for 2.5 h, the mixture was cooled to -20 $^\circ\text{C}$. TMSOTf (3.8 μl , 21 μmol) was then added at the same temperature. After 2.5 h, the reaction was quenched by the addition of water. The resulting mixture was filtered through a Celite pad. The filtrate was washed with water, satd NaHCO_3 aq, and brine, dried over Na_2SO_4 , filtered, and concentrated in vacuo. Flash chromatography (silica gel: 15 g, toluene/AcOEt=50:1 \rightarrow 30:1) afforded compound **33** as a colorless oil. Yield 103 mg (62%, two steps). ^1H NMR (600 MHz, CDCl_3) δ 8.17 (2H, d, $J=8.1$ Hz, Bz), 7.64–7.17 (40H, m aromatic), 7.04 (2H, d, $J=8.8$ Hz, MPM), 6.81 (2H, d, $J=8.8$ Hz, MPM), 5.30 (1H, dd, $J_{4'',3''}=9.5$ Hz, $J_{4'',5''}=10.2$ Hz, H-4''), 5.20 (1H, s, H-1'), 5.10 (1H, s, H-2'), 4.98 (1H, s, H-5'), 4.91 (1H, d, $J_{\text{gem}}=10.8$ Hz, PhCH_2), 4.91 (1H, d, $J_{\text{gem}}=10.8$ Hz, PhCH_2), 4.79 (1H, d, $J_{\text{gem}}=11.5$ Hz, PhCH_2), 4.77 (1H, d, $J_{\text{gem}}=10.2$ Hz, PhCH_2), 4.74 (1H, d, $J_{\text{gem}}=12.2$ Hz, PhCH_2), 4.73 (1H, d, $J_{\text{gem}}=10.8$ Hz, PhCH_2), 4.72 (1H, d, $J_{1',2'}=4.0$ Hz, H-1''), 4.68 (1H, d, $J_{\text{gem}}=10.8$ Hz, PhCH_2), 4.56 (1H, d, $J_{\text{gem}}=11.5$ Hz, PhCH_2), 4.50 (1H, d, $J_{\text{gem}}=12.2$ Hz, PhCH_2), 4.44 (1H, d, $J_{1,2}=7.4$ Hz, H-1), 4.16 (1H, s, H-3'), 4.03 (1H, s, H-4'), 4.03 (1H, dd, $J_{6a,6b}=11.5$ Hz, $J_{6a,5}=1.3$ Hz, H-6a), 4.01 (1H, d, $J_{\text{gem}}=10.2$ Hz, PhCH_2), 3.92 (1H, d, $J_{\text{gem}}=10.2$ Hz, PhCH_2), 3.86 (1H, dd, $J_{6b,6a}=11.5$ Hz, $J_{6b,5}=4.7$ Hz, H-6b), 3.79–3.77 (4H, m, H-5'', OCH_3), 3.71 (1H, dd, $J_{6a'',6b''}=10.2$ Hz, $J_{6a'',5''}=4.0$ Hz, H-6a''), 3.67 (1H, dd, $J_{6b'',6a''}=10.2$ Hz, $J_{6b'',5''}=2.7$ Hz, H-6b''), 3.61–3.56 (4H, m, H-3, H-4, H-5, H-3''), 3.42 (1H, dd, $J_{2,1}=7.4$ Hz, $J_{2,3}=8.1$ Hz, H-2), 3.37 (3H, s, OCH_3), 3.30 (1H, dd, $J_{2'',1''}=4.0$ Hz, $J_{2'',3''}=9.5$ Hz, H-2''), 2.64–2.61 (2H, m, $\text{CH}_3\text{C}(=\text{O})\text{C}_2\text{H}_4\text{C}(=\text{O})-$), 2.41–2.35 (2H, m, $\text{CH}_3\text{C}(=\text{O})\text{C}_2\text{H}_4\text{C}(=\text{O})-$), 2.16 (3H, s, $\text{CH}_3\text{C}(=\text{O})\text{C}_2\text{H}_4\text{C}(=\text{O})-$), 0.99 (9H, s, $\text{C}(\text{CH}_3)_3$); ^{13}C NMR (150 MHz, CDCl_3) δ 206.0, 170.7, 159.2, 138.0, 137.3, 135.7, 135.7, 129.9, 129.6, 129.5, 128.5, 128.5, 128.4, 128.3, 128.3, 128.3, 128.2, 128.1, 127.9, 127.8, 127.8, 127.7, 127.6, 127.6, 127.5, 127.4, 113.6, 102.3, 99.2, 98.8, 84.6, 82.2, 78.0, 77.5, 75.7, 75.2, 74.9, 74.9, 74.1, 72.4, 72.3, 72.3, 71.0, 70.7, 69.7, 67.8, 67.3, 67.1, 63.3, 61.5, 55.2, 51.8, 37.7, 29.8, 27.8, 26.6; HRMS (positive mode) found: m/z 1590.6357 $[\text{M}+\text{Na}]^+$, calcd for $\text{C}_{90}\text{H}_{97}\text{N}_3\text{O}_{20}\text{SiNa}$: 1590.6332.

3.28. Benzyl (2-O-azido-6-O-tert-butylidiphenylsilyl-2-O-deoxy-3-O-(4-methoxybenzyl)- α -D-glucopyranosyl)-(1 \rightarrow 4)-(methyl 2-O-benzoyl-3-O-benzyl- β -D-glucopyranosyluronate)-(1 \rightarrow 6)-2,3,4-tetra-O-benzyl- β -D-glucopyranoside

Compound **30** (649 mg, 0.413 mmol) was dissolved in pyridine/AcOH (3:2, 10 ml) and hydrazine monohydrate (30.2 μl , 0.620 mmol) was added at room temperature. After 3 h, the reaction was quenched by addition of acetone and the mixture was diluted with AcOEt. The organic layer was washed with 1 M HCl aq, satd NaHCO_3 aq, and brine, dried over Na_2SO_4 , filtered, and concentrated in vacuo. Flash chromatography (silica gel: 30 g, toluene/EtOAc=20:1 \rightarrow 10:1) afforded benzyl (2-O-azido-6-O-tert-butylidiphenylsilyl-2-O-deoxy-3-O-(4-methoxybenzyl)- α -D-glucopyranosyl)-(1 \rightarrow 4)-(methyl 2-O-

benzoyl-3-*O*-benzyl- β -*D*-glucopyranosyluronate)-(1 \rightarrow 6)-2,3,4-tetra-*O*-benzyl- β -*D*-glucopyranoside as a white solid. Yield 540 mg (89%). ^1H NMR (600 MHz, CDCl_3) δ 7.96 (2H, d, $J=8.1$ Hz, Bz), 7.69–7.13 (40H, m, aromatic), 6.91 (2H, d, $J=8.1$ Hz, MPM), 5.40–5.38 (2H, m, H-1'', H-2''), 4.88 and 4.79 (each 1H, d, $J_{\text{gem}}=10.8$ Hz, PhCH_2), 4.84 and 4.80 (each 1H, d, $J_{\text{gem}}=10.8$ Hz, PhCH_2), 4.79 and 4.64 (each 1H, d, $J_{\text{gem}}=10.8$ Hz, PhCH_2), 4.72 and 4.71 (each 1H, d, $J_{\text{gem}}=12.2$ Hz, PhCH_2), 4.68 and 4.45 (each 1H, d, $J_{\text{gem}}=11.5$ Hz, PhCH_2), 4.67 (1H, d, $J_{1,2'}=8.1$ Hz, H-1'), 4.66 and 4.41 (each 1H, d, $J_{\text{gem}}=12.2$ Hz, PhCH_2), 4.36 (1H, d, $J_{1,2}=7.4$ Hz, H-1), 4.25 (1H, dd, $J_{4',5'}=8.8$ Hz, $J_{4',3'}=8.8$ Hz, H-4'), 4.11 (1H, d, $J_{6a'',6b''}=10.8$ Hz, H-6a''), 4.01 (1H, d, $J_{5',4'}=8.8$ Hz, H-5'), 3.94–3.90 (2H, m, H-6a, H-3'), 3.79–3.72 (6H, m, H-6b, H-3'', H-4'', OCH_3), 3.60–3.53 (5H, m, H-3, H-6b'', OCH_3), 3.47–3.40 (3H, m, H-2, H-5, H-5''), 3.35 (1H, dd, $J_{4,5}=9.5$ Hz, $J_{4,3}=9.5$ Hz, H-4), 3.22 (1H, dd, $J_{2'',3''}=9.5$ Hz, $J_{2'',1''}=3.3$ Hz, H-2''), 2.78 (1H, d, $J_{\text{OH},4''}=1.3$ Hz, OH), 1.05 (9H, s, $\text{C}(\text{CH}_3)_3$); ^{13}C NMR (150 MHz, CDCl_3) δ 168.3, 164.8, 159.3, 137.8, 137.2, 135.6, 129.9, 129.9, 129.8, 129.6, 128.4, 128.3, 128.3, 128.3, 128.2, 128.1, 127.9, 127.9, 127.8, 127.7, 127.7, 127.6, 113.9, 102.1, 101.1, 97.5, 84.4, 82.2, 82.0, 78.8, 77.7, 75.6, 74.8, 74.8, 74.7, 74.5, 74.4, 74.3, 73.1, 73.0, 70.9, 70.7, 68.2, 64.0, 62.5, 55.2, 52.5, 26.8; HRMS (positive mode) found: m/z 1492.5978 [$\text{M}+\text{Na}$] $^+$, calcd for $\text{C}_{85}\text{H}_{91}\text{N}_3\text{O}_{18}\text{SiNa}$: 1492.5965.

3.29. Benzyl [2-*O*-azido-6-*O*-*tert*-butyldiphenylsilyl-2-*O*-deoxy-3-*O*-(4-methoxybenzyl)-4-*O*-methyl- α -*D*-glucopyranosyl]-(1 \rightarrow 4)-(methyl 2-*O*-benzoyl-3-*O*-benzyl- β -*D*-glucopyranosyluronate)-(1 \rightarrow 6)-2,3,4-tetra-*O*-benzyl- β -*D*-glucopyranoside (34)

To a solution of benzyl (2-*O*-azido-6-*O*-*tert*-butyldiphenylsilyl-2-*O*-deoxy-3-*O*-(4-methoxybenzyl)- α -*D*-glucopyranosyl)-(1 \rightarrow 4)-(methyl 2-*O*-benzoyl-3-*O*-benzyl- β -*D*-glucopyranosyluronate)-(1 \rightarrow 6)-2,3,4-tetra-*O*-benzyl- β -*D*-glucopyranoside (11.0 mg, 7.48 μmol) in anhydrous DMF (2 ml) were added MeI (0.90 μl , 15 μmol) and MS4AP (200 mg) at room temperature under Ar and the mixture was stirred for 2 h. After cooling to -20°C , lithium hexamethyldisilazide (LHMDS) in THF (7.0 μl , 1.6 M) was added at the same temperature. The reaction mixture was further stirred and gradually warmed to 0°C . After 4 h, the reaction was quenched by the addition of methanol and the reaction mixture was filtered through a Celite pad. The filtrate was washed with 10% $\text{Na}_2\text{S}_2\text{O}_3$ aq and brine, dried over Na_2SO_4 , filtered, and concentrated in vacuo. Flash chromatography (silica gel: 5 g, toluene/EtOAc=30:1 \rightarrow 10:1) afforded compound **34** as a white solid. Yield 9.44 mg (85%). ^1H NMR (600 MHz, CDCl_3) δ 7.97 (2H, d, $J=6.8$ Hz, Bz), 7.70–7.12 (40H, m, aromatic), 6.91 (2H, d, $J=8.8$ Hz, MPM), 5.42 (1H, dd, $J_{2',1'}=8.1$ Hz, $J_{2',3'}=8.1$ Hz, H-2'), 5.37 (1H, d, $J_{1'',2''}=4.0$ Hz, H-1''), 4.87 and 4.81 (each 1H, d, $J_{\text{gem}}=10.8$ Hz, PhCH_2), 4.85 and 4.69 (each 1H, d, $J_{\text{gem}}=10.2$ Hz, PhCH_2), 4.82 and 4.78 (each 1H, d, $J_{\text{gem}}=13.6$ Hz, PhCH_2), 4.79 and 4.39 (each 1H, d, $J_{\text{gem}}=12.2$ Hz, PhCH_2), 4.70 (1H, d, $J_{1,2'}=8.1$ Hz, H-1'), 4.68 and 4.63 (each 1H, d, $J_{\text{gem}}=10.2$ Hz, PhCH_2), 4.67 and 4.42 (each 1H, d, $J_{\text{gem}}=10.8$ Hz, PhCH_2), 4.34 (1H, d, $J_{1,2}=7.4$ Hz, H-1), 4.20 (1H, dd, $J_{4',3'}=8.8$ Hz, $J_{4',5'}=8.8$ Hz, H-4'), 4.09 (1H, d, $J_{6a,6b}=10.8$ Hz, H-6a), 4.01 (1H, d, $J_{5',4'}=8.8$ Hz, H-5'), 3.94–3.89 (2H, m, H-3', H-6a''), 3.81–3.78 (5H, m, H-3'', H-6b'', OCH_3), 3.62–3.50 (9H, m, H-3, H-6b, H-4'', $\text{OCH}_3 \times 2$), 3.41–3.36 (2H, m, H-2, H-5), 3.32–3.28 (2H, m, H-4, H-5''), 3.25 (1H, dd, $J_{2'',1''}=4.0$ Hz, $J_{2'',3''}=10.8$ Hz, H-2''), 1.02 (9H, s, $\text{C}(\text{CH}_3)_3$); ^{13}C NMR (150 MHz, CDCl_3) δ 168.3, 164.8, 159.4, 138.2, 137.2, 137.2, 135.8, 135.5, 130.0, 129.9, 129.6, 129.6, 128.4, 128.3, 128.3, 128.2, 128.1, 127.9, 127.8, 127.8, 127.8, 127.7, 127.7, 127.6, 127.6, 127.5, 113.9, 102.0, 101.0, 97.6, 84.4, 82.2, 82.0, 79.6, 79.2, 77.7, 75.6, 75.2, 74.7, 74.6, 74.5, 74.4, 74.2, 74.2, 73.0, 72.3, 70.7, 68.3, 63.1, 61.6, 60.7, 55.2, 26.8; HRMS (positive mode) found: m/z 1506.6133 [$\text{M}+\text{Na}$] $^+$, calcd for $\text{C}_{86}\text{H}_{93}\text{N}_3\text{O}_{18}\text{SiNa}$: 1506.6121.

3.30. Benzyl (2-*O*-azido-2-*O*-deoxy-3-*O*-(4-methoxybenzyl)-4-*O*-methyl- α -*D*-glucopyranosyl)-(1 \rightarrow 4)-(methyl 2-*O*-benzoyl-3-*O*-benzyl- β -*D*-glucopyranosyluronate)-(1 \rightarrow 6)-2,3,4-tetra-*O*-benzyl- β -*D*-glucopyranoside (35)

To a solution of compound **34** (19.9 mg, 13.4 μmol) in pyridine (4 ml) was added HF-pyridine (174 μl , 6.70 μmol) at room temperature. The mixture was stirred for 2 days. The volatiles were removed in vacuo. Flash chromatography (silica gel: 5 g, toluene/EtOAc=10:1 \rightarrow 4:1) afforded compound **35** as a colorless oil. Yield 15.9 mg (95%). ^1H NMR (600 MHz, CDCl_3) δ 7.97 (2H, d, $J=6.8$ Hz, Bz), 7.47–7.13 (30H, m, aromatic), 6.90 (2H, d, $J=8.8$ Hz, MPM), 5.41 (1H, d, $J_{1'',2''}=3.3$ Hz, H-1''), 5.41 (1H, dd, $J_{2',1'}=7.4$ Hz, $J_{2',3'}=8.8$ Hz, H-2'), 4.88 and 4.86 (each 1H, d, $J_{\text{gem}}=10.8$ Hz, PhCH_2), 4.82 and 4.69 (each 1H, d, $J_{\text{gem}}=10.2$ Hz, PhCH_2), 4.77 and 4.75 (each 1H, d, $J_{\text{gem}}=9.5$ Hz, PhCH_2), 4.73 (1H, d, $J_{1,2'}=7.4$ Hz, H-1'), 4.70 and 4.65 (each 1H, d, $J_{\text{gem}}=10.8$ Hz, PhCH_2), 4.70 and 4.45 (each 1H, d, $J_{\text{gem}}=10.8$ Hz, PhCH_2), 4.69 and 4.42 (each 1H, d, $J_{\text{gem}}=10.2$ Hz, PhCH_2), 4.37 (1H, d, $J_{1,2}=8.1$ Hz, H-1), 4.31 (1H, dd, $J_{4',3'}=8.1$ Hz, $J_{4',5'}=8.8$ Hz, H-4'), 4.12 (1H, dd, $J_{6a,5}=1.3$ Hz, $J_{6a,6b}=10.8$ Hz, H-6a), 4.06 (1H, d, $J_{5',4'}=8.8$ Hz, H-5'), 3.94 (1H, dd, $J_{3',2'}=8.8$ Hz, $J_{3',4'}=8.1$ Hz, H-3'), 3.87–3.75 (8H, m, H-3'', H-6a'', $\text{OCH}_3 \times 2$), 3.69 (1H, dd, $J_{6b'',5''}=3.4$ Hz, $J_{6b'',6a''}=11.5$ Hz, H-6b''), 3.60–3.54 (5H, m, H-3, H-6b, OCH_3), 3.45–3.38 (3H, m, H-2, H-5, H-5''), 3.34 (1H, dd, $J_{4,3}=8.8$ Hz, $J_{4,5}=10.2$ Hz, H-4), 3.24–3.21 (2H, m, H-2'', H-4''); ^{13}C NMR (150 MHz, CDCl_3) δ 168.6, 164.8, 138.4, 138.2, 133.2, 129.8, 129.6, 128.4, 128.3, 128.3, 128.3, 128.1, 127.9, 127.8, 127.8, 127.7, 127.7, 127.6, 113.8, 102.1, 101.0, 97.5, 84.4, 82.2, 82.0, 80.0, 79.3, 77.7, 75.6, 75.0, 75.0, 74.7, 74.6, 74.4, 74.4, 73.3, 72.1, 70.7, 68.3, 63.0, 61.2, 60.8, 55.2, 52.8; HRMS (positive mode) found: m/z 1172.4349 [$\text{M}+\text{Na}$] $^+$, calcd for $\text{C}_{70}\text{H}_{75}\text{N}_3\text{O}_{18}\text{Na}$: 1172.4344.

3.31. 2-Deoxy-4-*O*-methyl-2-sulfoamino- α -*D*-glucopyranosyl-(1 \rightarrow 4)-(β -*D*-glucopyranosyluronic acid)-(1 \rightarrow 6)-*D*-glucopyranose disodium salt (38)

To a solution of compound **35** (70.6 mg, 60.9 μmol) in methanol/THF (1:1, 1.6 ml) was added 1 M NaOH aq (400 μl) at room temperature. After 24 h, the reaction was quenched by addition of Dowex 50W (H^+). After removal of insoluble materials by filtration, the filtrate was concentrated in vacuo. To a solution of the residue in methanol/THF (1:1, 3.2 ml) was added 0.1 M NaOH aq (974 μl) and 1 M PMe_3 in THF (487 μl , 0.487 μmol) at room temperature. After 2 days, the reaction was quenched by addition of 0.1 M HCl aq (974 μl). After the volatiles were removed in vacuo, the residue was dissolved in H_2O (10 ml). SO_3 -pyridine (970 mg, 6.09 μmol) was added. The pH of the reaction mixture was maintained at 9.5 by the addition of 5 M NaOH aq during the reaction. After 5 h, the volatiles were removed in vacuo. The residue was then suspended in MeOH. Insoluble materials were removed by filtration and the filtrate was concentrated in vacuo. Reversed phase chromatography (ODS: 6 g, $\text{H}_2\text{O}/\text{MeOH}=8:1 \rightarrow 0:1$) afforded compound **37**. Obtained **37** was dissolved in $\text{H}_2\text{O}/\text{MeOH}/\text{AcOH}$ (5:5:1, 16 ml). Pd/C (10%, 51.2 mg) was added to the mixture. The reaction mixture was stirred for 6 days at room temperature under H_2 (7 kg/cm 2). After filtration by membrane filter, the filtrate was concentrated in vacuo. Gel filtration (Sephadex G-25 fine: ϕ 1.5 \times 1000 mm, H_2O) afforded compound **38** as a white solid. Yield 4.52 mg (11%, $\alpha/\beta=1:2$, four steps). ^1H NMR (600 MHz, D_2O) δ for β -anomer 5.40 (1H, d, $J_{1'',2''}=3.4$ Hz, H-1''), 4.45 (1H, d, $J_{1,2}=8.1$ Hz, H-1), 4.39 (1H, d, $J_{1,2'}=8.1$ Hz, H-1'), 3.97 (1H, d, $J_{6a,6b}=11.5$ Hz, H-6a), 3.92 (1H, d, $J_{6a'',6b''}=11.5$ Hz, H-6a''), 3.82 (1H, d, $J_{5',4'}=8.8$ Hz, H-5'), 3.67–3.60 (6H, m, H-5, H-6b, H-3', H-4', H-5'', H-6b''), 3.48 (1H, dd, $J_{3'',2''}=10.2$ Hz, $J_{3'',4''}=9.5$ Hz, H-3''), 3.41 (1H, dd, $J_{4,3}=8.8$ Hz, $J_{4,5}=8.8$ Hz, H-4), 3.36 (3H, s, OCH_3), 3.28 (1H, dd, $J_{3,2}=8.8$ Hz, $J_{3,4}=8.8$ Hz, H-3), 3.22 (1H, dd,

$J_{2',1'}=8.1$ Hz, $J_{2',3'}=8.1$ Hz, H-2'), 3.13 (1H, dd, $J_{4'',3''}=9.5$ Hz, $J_{4'',5''}=10.2$ Hz, H-4''), 3.07 (1H, dd, $J_{2'',1''}=3.4$ Hz, $J_{2'',3''}=10.2$ Hz, H-2''), 3.60 (1H, dd, $J_{2,1}=8.1$ Hz, $J_{2,3}=8.8$ Hz, H-2); ^{13}C NMR (150 MHz, D_2O) δ 172.9, 102.6, 97.5, 95.8, 78.7, 76.4, 75.7, 75.5, 74.7, 73.9, 72.5, 70.8, 70.6, 70.2, 69.3, 68.9, 68.7, 59.6, 57.9, HRMS (negative mode) found: m/z 632.1133 $[\text{M}-\text{Na}]^-$, calcd for $\text{C}_{19}\text{H}_{31}\text{NO}_{19}\text{SNa}$: 632.1114.

Acknowledgements

The present work was financially supported in part by grants from Japan Science and Technology Agency (Evolutionally-venture program V07-05 to Y.S., CREST to Y.S.), and the Japan Ministry of Health, Labour and Welfare (MHLW) (Nanomedicine program to Y.S.), and the National Institutes of Health, USA (RO1 HL079182 to M.S.).

References and notes


1. *Chemistry and Biology of Heparin and Heparan Sulfate*; Garg, H. G., Linhardt, R. J., Hales, C. A., Eds.; Elsevier: Amsterdam, The Netherlands, 2005.
2. (a) Conrad, H. E. *Heparin-Binding Proteins*; Academic: San Diego, CA, USA, 1998; (b) Imberty, A.; Lortat-Jacob, H.; Pérez, S. *Carbohydr. Res.* **2007**, *342*, 430–439; (c) Raman, R.; Sasisekharan, V.; Sasisekharan, R. *Chem. Biol.* **2005**, *12*, 267–277; (d) Coombe, D. R.; Kett, W. C. *Cell. Mol. Life Sci.* **2005**, *62*, 410–424; (e) Petitou, M.; van Boeckel, C. A. A. *Angew. Chem., Int. Ed.* **2004**, *43*, 3118–3133; (f) Rabenstein, D. A. *Nat. Prod. Rep.* **2002**, *19*, 312–331; (g) Capila, I.; Linhardt, R. J. *Angew. Chem., Int. Ed.* **2002**, *41*, 390–412; (h) Turnbull, J.; Powell, A.; Guimond, S. *Trends Cell Biol.* **2001**, *11*, 75–82; (i) Casu, B.; Lindahl, U. *Adv. Carbohydr. Chem. Biochem.* **2001**, *57*, 159–206; (j) Esko, J. D.; Lindahl, U. *J. Clin. Invest.* **2001**, *108*, 169–173; (k) Bernfield, M.; Götte, M.; Park, P. W.; Reizes, O.; Fitzgerald, M. L.; Lincecum, J.; Zako, M. *Annu. Rev. Biochem.* **1999**, *68*, 729–777.
3. For comprehensive reviews on the synthesis of GAGs, see: (a) Noti, C.; Seeberger, P. H. *Chem. Biol.* **2005**, *12*, 731–756; (b) Karst, N. A.; Linhardt, R. J. *Curr. Med. Chem.* **2003**, *10*, 1993–2031; (c) Poletti, L.; Lay, L. *Eur. J. Org. Chem.* **2003**, 2999–3024; (d) Yeung, B. K. S.; Chong, P. Y. C.; Petillo, P. A. *J. Carbohydr. Chem.* **2002**, *21*, 799–865; (e) Tamura, J. *Trends Glycosci. Glycotechnol.* **2001**, *13*, 65–88.
4. For recent articles on the synthesis of GAGs see: (a) Polat, T.; Wong, C. H. *J. Am. Chem. Soc.* **2007**, *129*, 12795–12800; (b) Zhou, Y.; Lin, F.; Chen, J.; Yu, B. *Carbohydr. Res.* **2006**, *341*, 1619–1629; (c) Noti, C.; de Paz, J. L.; Polito, L.; Seeberger, P. H. *Chem.—Eur. J.* **2006**, *12*, 8664–8686; (d) de Paz, J. L.; Noti, C.; Seeberger, P. H. *J. Am. Chem. Soc.* **2006**, *128*, 2766–2767; (e) Gama, C. I.; Tully, S. E.; Sotogaku, N.; Clark, P. M.; Rawat, M.; Vaidehi, N.; Goddard, W. A.; Nishi, A.; Hsieh-Wilson, L. C. *Nat. Chem. Biol.* **2006**, *2*, 467–473; (f) Codée, J. D. C.; Stubba, B.; Schiattarella, M.; Overkleeft, H. S.; van Boeckel, C. A. A.; van Boom, J. H.; van der Marel, G. A. *J. Am. Chem. Soc.* **2005**, *127*, 3767–3773; (g) Lee, J. C.; Lu, X. A.; Kulkarni, S. S.; Wen, Y. S.; Hung, S. C. *J. Am. Chem. Soc.* **2004**, *126*, 476–477; (h) Orgueira, H. A.; Bartolozzi, A.; Schell, P.; Litjens, R. E.; Palmacci, E. R.; Seeberger, P. H. *Chem.—Eur. J.* **2003**, *9*, 140–169; (i) de Paz, J. L.; Ojeda, R.; Reichardt, Martin-Lomas, M. *Eur. J. Org. Chem.* **2003**, 3308–3324; (j) Kovensky, J.; Mallet, J. M.; Esnault, J.; Driguez, P. A.; Sizun, P.; Héroult, J. P.; Herbert, J. M.; Petitou, M.; Sinaý, P. *Eur. J. Org. Chem.* **2002**, 3595–3603; (k) de Paz, J. L.; Angulo, J.; Lassaletta, J. M.; Nieto, P. M.; Redondo-Horcajo, M.; Lozano, R. M.; Gimenez-Gallego, G.; Martin-Lomas, M. *ChemBioChem* **2001**, *2*, 673–685 and references cited therein.
5. For recent articles on HP/HS oligosaccharide library, see: (a) Hecht, M. L.; Rosental, B.; Horlacher, T.; Hershkovitz, O.; de Paz, J. L.; Noti, C.; Schauer, S.; Porgador, A.; Seeberger, P. J. *Proteome Res.* **2009**, *8*, 712–720; (b) Herczeg, M.; Lázár, L.; Borbás, A.; Lipták, A.; Antus, S. *Org. Lett.* **2009**, *11*, 2619–2622; (c) Dilhas, A.; Lucas, R.; Loureiro-Morais, L.; Hersant, Y.; Bonnaffé, D. *J. Comb. Chem.* **2008**, *10*, 166–169; (d) de Paz, J. L.; Moseman, E. A.; Noti, C.; Polito, L.; von Andrian, U. H.; Seeberger, P. J. *ACS Chem. Biol.* **2007**, *2*, 735–744; (e) Lu, L.-D.; Shie, C.-R.; Kulkarni, S. S.; Pan, G.-R.; Lu, X.-A.; Hung, S.-C. *Org. Lett.* **2006**, *8*, 5995–5998; (f) Fan, R.-H.; Achkar, J.; Hernández-Torres, J. M.; Wei, A. *Org. Lett.* **2005**, *7*, 5095–5098.
6. For our previous studies, see: (a) Wakao, M.; Saito, A.; Ohishi, K.; Kishimoto, Y.; Nishimura, T.; Sobel, M.; Suda, Y. *Bioorg. Med. Chem. Lett.* **2008**, *18*, 2499–2504; (b) Suda, Y.; Arano, A.; Fukui, Y.; Koshida, S.; Wakao, M.; Nishimura, T.; Kusumoto, S.; Sobel, M. *Bioconjugate Chem.* **2006**, *17*, 1125–1135; (c) Koshida, S.; Suda, Y.; Sobel, M.; Kusumoto, S. *Tetrahedron Lett.* **2001**, *42*, 1289–1292; (d) Koshida, S.; Suda, Y.; Fukui, Y.; Ormsby, J.; Sobel, M.; Kusumoto, S. *Tetrahedron Lett.* **1999**, *40*, 5725–5728; (e) Suda, Y.; Bird, K.; Shiyama, T.; Koshida, S.; Marques, D.; Fukase, K.; Sobel, M.; Kusumoto, S. *Tetrahedron Lett.* **1996**, *37*, 1053–1056.
7. For articles on the synthesis of uronic acids moieties, see: (a) Hansen, S. U.; Baráth, M.; Salameh, B. A. B.; Pritchard, R. G.; Stimpson, W. T.; Gardiner, J. M.; Jayson, G. C. *Org. Lett.* **2009**, *11*, 4528–4531; (b) Tatai, J.; Osztrovszky, G.; Kajtár-Perey, M.; Fügedi, P. *Carbohydr. Res.* **2008**, *343*, 596–606; (c) Hassan, H. H. A. M. *Mini-Rev. Org. Chem.* **2007**, *4*, 61–74; (d) Lohman, G. J. S.; Hunt, D. K.; Högermeier, J. A.; Seeberger, P. H. *J. Org. Chem.* **2003**, *68*, 7559–7561; (e) Ke, W.; Whitfield, D. M.; Gill, M.; Larocque, S.; Yu, S.-H. *Tetrahedron Lett.* **2003**, *44*, 7767–7770; (f) Barroca, N.; Jacquinet, J.-C. *Carbohydr. Res.* **2000**, *329*, 667–679; (g) Lubineau, A.; Alais, G. J.; Bonnaffé, D. *Tetrahedron Lett.* **2000**, *41*, 307–311; (h) Ojeda, R.; de Paz, J. L.; Martin-Lomas, M.; Lassaletta, J. M. *Synlett* **1999**, 1316–1318; (i) Hinou, H.; Kurosawa, H.; Matsuo, K.; Terunuma, D.; Kuzuhara, H. *Tetrahedron Lett.* **1999**, *40*, 1501–1504; (j) Rochepeau-Jobron, L.; Jacquinet, J.-C. *Carbohydr. Res.* **1997**, *303*, 395–406; (k) Medakovic, D. *Carbohydr. Res.* **1994**, *253*, 299–300; (l) Jacquinet, J.-C.; Petitou, M.; Duchaussoy, P.; Lederman, I.; Choay, J.; Torri, G.; Sinaý, P. *Carbohydr. Res.* **1984**, *130*, 221–241; (m) Czuk, R.; Hoening, H.; Nimpf, J.; Weidmann, H. *Tetrahedron Lett.* **1980**, *21*, 2135–2136.
8. Greene, T. W.; Wuts, P. G. M. *Protective Groups in Organic Synthesis*, 3rd ed.; John Wiley & Sons: New York, NY, 1999.
9. A trace amount of furanose form was observed by NMR analysis.
10. Alper, P. B.; Hung, S.-C.; Wong, C.-H. *Tetrahedron Lett.* **1996**, *37*, 6029–6032.
11. α -Selective glycosylation with 2-azido imidate in the presence of TBDMSOTf was previously reported, see: Kovensky, J.; Duchaussoy, P.; Petitou, M.; Sinaý, P. *Tetrahedron: Asymmetry* **1996**, *7*, 3119–3128.

Ganglioside GQ1b: Efficient Total Synthesis and the Expansion to Synthetic Derivatives To Elucidate Its Biological Roles

Akihiro Imamura, Hiromune Ando, Hideharu Ishida, and Makoto Kiso

J. Org. Chem., 2009, 74 (8), 3009-3023 • DOI: 10.1021/jo8027888 • Publication Date (Web): 18 March 2009

Downloaded from <http://pubs.acs.org> on May 10, 2009



More About This Article

Additional resources and features associated with this article are available within the HTML version:

- Supporting Information
- Links to the 1 articles that cite this article, as of the time of this article download
- Access to high resolution figures
- Links to articles and content related to this article
- Copyright permission to reproduce figures and/or text from this article

[View the Full Text HTML](#)



ACS Publications
High quality. High impact.

The Journal of Organic Chemistry is published by the American Chemical Society, 1155 Sixteenth Street N.W., Washington, DC 20036


Ganglioside GQ1b: Efficient Total Synthesis and the Expansion to Synthetic Derivatives To Elucidate Its Biological Roles[†]

Akihiro Imamura,^{*,‡,§,||} Hiromune Ando,^{‡,§} Hideharu Ishida,[‡] and Makoto Kiso^{*,‡,§}

Department of Applied Bioorganic Chemistry, Faculty of Applied Biological Sciences, Gifu University, 1-1 Yanagido, Gifu-shi, Gifu 501-1193, Japan, and Institute for Integrated Cell-Material Sciences (iCeMS), Kyoto University, 69 Konocho, Yoshida, Sakyo-ku, Kyoto 606-8501, Japan

aimamura@ualberta.ca; kiso@gifu-u.ac.jp

Received December 25, 2008



The convergent total synthesis of ganglioside GQ1b based on the “cassette approach” between the nonreducing end GQ1b-core heptasaccharide and glucosylceramide building blocks was accomplished in high overall yield. The use of a sialyl α (2→8)sialyl α (2→3)galactose sequence as the key building block enhanced the efficiency of the glycan assembly and led to preparative-scale synthesis readily applicable for large-scale preparation. In addition, a judicious choice of *p*-methoxybenzyl protecting groups on glucosylceramide provided a solution to the previous synthetic problems, including a decrease in the yield of the deprotection steps, and led to elevation of the total yield. Furthermore, unnatural-type GQ1b derivatives were synthesized systematically in good yields by capitalizing on a similar approach in order to elucidate their biological roles.

Introduction

Gangliosides, sialylated glycosphingolipids, are the components of the membranes in the cells of all living organisms and are particularly abundant in the nervous system on the vertebrate cell plasma membrane.² For the past two decades, gangliosides have attracted wide attention in various scientific fields, because

they play various important roles in vital processes. Ganglioside GQ1b, which is classified as one of the b-series gangliosides, exists abundantly in the mammalian central nervous system and participates in many physiological events, such as neurite extension,³ toxin binding,⁴ modulation of protein phosphorylation,⁵ cell adhesion and growth,^{6,7} and apoptosis.⁸ Despite its biological importance, advancement in the biological study of b-series gangliosides, including GQ1b, has been restricted due

[†] Synthetic studies on sialoglycoconjugates, part 147. For part 146, see ref 1.

* To whom correspondence should be addressed. A.I.: phone, 81-58-293-2918; fax, 81-58-293-2918. M.K.: phone, 81-58-293-2916; fax, 81-58-293-2918.

[‡] Gifu University.

[§] Kyoto University.

^{||} Current address: Department of Chemistry, E5-55 Gunning-Lemieux Chemistry Centre, The University of Alberta, Edmonton, AB, T6G 2G2, Canada.

(1) Yamaguchi, M.; Ishida, H.; Kiso, M. *Carbohydr. Res.* **2008**, *343*, 1849–1857.

(2) Wiegandt, H. In *Glycolipids*; Wiegandt, H., Ed.; Elsevier: New York, 1985; pp 199–260. (b) Lloyd, K. O.; Furukawa, K. *Glycoconjugate J.* **1998**, *15*, 627–636. (c) Sonnino, S.; Mauri, L.; Chigorno, V.; Prinetti, A. *Glycobiology* **2006**, *17*, 1R–13R.

(3) Tsuji, S.; Arita, M.; Nagai, Y. *J. Biochem.* **1983**, *94*, 303–306.

(4) Ångström, J.; Teneberg, S.; Karlsson, K.-L. *Proc. Natl. Acad. Sci. U.S.A.* **1994**, *91*, 11859–11863.




FIGURE 1. Structure of target compounds.

to a deficiency in the supply of pure gangliosides. In 1994, our group first achieved the total synthesis of ganglioside GQ1b, which led to the elucidation of the biological functions of GQ1b at the molecular level.⁹ However, there were problems yet to be solved with regard to the overall yields and stereoselectivity of the coupling reactions, which prevented large quantity synthesis. Recently, it has become essential to develop large-scale synthesis of gangliosides, because of multifaceted research needs, e.g., microarray, molecular imaging, and X-ray crystallographic studies. Therefore, we envisioned renewal of the total synthetic method for natural ganglioside GQ1b (**1**) (Figure 1). Furthermore, we decided to develop access to versatile unnatural GQ1b derivatives to elucidate their biological roles. The following were designed as unnatural GQ1b derivatives (Figure 1): (i) *SE* [2-(trimethylsilyl)ethyl]-type derivative (**2**) demonstrated that SE-type GT1b derivatives serve as good tools for X-ray crystallographic analysis with lectins such as Siglec-7,¹⁰ botulinum neurotoxin,¹¹ and tetanus toxin,¹² because replacement of ceramide with an SE group may render it more amenable to crystallization (being more water-soluble, less flexible, and less likely to form micelles) and (ii) *Glc-ended* derivative (**3**) allowed accomplishment of the synthesis of GM1,

GM2, and GM3 bearing D-glucose at the reducing terminal and succeeded in the assembly of the sugar microarray.¹³ The D-glucose residue in this derivative provided a reactive aldehyde functionality at the reducing end and served as a spacer between the targeted sugar chain and the scaffold for immobilization because of its appropriate hydrophilicity and flexibility.

Herein, we report the renewed efficient synthesis of natural ganglioside GQ1b and its unnatural derivatives.

Results and Discussion

To achieve the efficient and systematic synthesis of the targeted compounds bearing different reducing terminals, sialyl α (2 \rightarrow 8)sialyl α (2 \rightarrow 3)galactose sequences were focused on as the common trisaccharide building blocks. Using this common sequence, a synthetic scheme for the targets was devised, as shown in Figure 2. This included four important glycosylation steps: (1) sialylation to produce the common trisaccharide; (2) galactosaminidation; (3) coupling of the trisaccharide donor and the tetrasaccharide acceptor, derived from the common trisaccharide; and (4) final glycosylation of the reducing terminal acceptors.

According to the above-mentioned scheme, disialylgalactose unit **6** was selected as the key compound and was prepared by glycosylation of the known galactose acceptor **5**¹⁴ with sialyl α (2 \rightarrow 8)sialyl phenylthioglycoside donor **4**.⁹ We were eager to find out how stereo- and regioselectivity in sialylation increases, and therefore a series of sialylations under various conditions were conducted. The results are summarized in Table 1. First, the sialylation of **5** in the presence of dimethyl(methylthio)sulfonium triflate (DMTST)¹⁵ as a promoter in acetonitrile at -35 °C hardly gave **6** (entry 1). In entry 2, by using

(5) Tsuji, S.; Nakajima, T.; Sasaki, Y.; Nagai, Y. *J. Biochem.* **1985**, *97*, 969–972.

(6) (a) Kleinman, H. K.; Martin, G. R.; Fishman, P. H. *Proc. Natl. Acad. Sci. U.S.A.* **1979**, *76*, 3367–3371. (b) Cheresch, D. A.; Pierschbacher, M. D.; Herzig, M. A.; Mujoo, K. *J. Cell Biol.* **1986**, *102*, 688–696.

(7) (a) Nakano, J.; Raj, B. K.; Asagami, C.; Lloyd, K. O. *J. Invest. Dermatol.* **1996**, *107*, 543–548. (b) Birkle, S.; Ren, S.; Slominski, A.; Zeng, G.; Gao, L.; Yu, R. K. *J. Neurochem.* **1999**, *72*, 954–961.

(8) De Maria, R.; Lenti, L.; Malisan, F.; d'Agostino, F.; Tomassini, B.; Zeuner, A.; Rippon, M. R.; Testi, R. *Science* **1997**, *277*, 1652–1655.

(9) (a) Ishida, H.-K.; Ishida, H.; Kiso, M.; Hasegawa, A. *Carbohydr. Res.* **1994**, *260*, C1–C6. (b) Ishida, H.-K.; Ishida, H.; Kiso, M.; Hasegawa, A.; *Tetrahedron: Asymmetry* **1994**, *5*, 2493–2512. (c) Nunomura, S.; Matsuzaki, Y.; Ito, Y.; Ogawa, T. *Pure Appl. Chem.* **1994**, *66*, 2123–2126.

(10) (a) Attrill, H.; Imamura, A.; Sharma, R. S.; Kiso, M.; Crocker, P. R.; van Aalten, D. M. F. *J. Biol. Chem.* **2006**, *281*, 32774–32783. (b) Attrill, H.; Takazawa, H.; Witt, S.; Kelm, S.; Isecke, R.; Brossmer, R.; Ando, T.; Ishida, H.; Kiso, M.; Crocker, P. R.; van Aalten, D. M. F. *Biochem. J.* **2006**, *397*, 271–278.

(11) Stenmark, P.; Dupuy, J.; Imamura, A.; Kiso, M.; Stevens, R. C. *PLoS Pathog.* **2008**, *4*, e1000129.

(12) Fotinou, C.; Emmsley, P.; Black, I.; Ando, H.; Ishida, H.; Kiso, M.; Sinha, K. A.; Fairweather, N. F.; Isaacs, N. W. *J. Biol. Chem.* **2001**, *276*, 32274–32281.

(13) Imamura, A.; Yoshikawa, T.; Komori, T.; Ando, M.; Ando, H.; Wakao, M.; Suda, Y.; Ishida, H.; Kiso, M. *Glycoconjugate J.* **2008**, *25*, 269–278.

(14) Komba, S.; Ishida, H.; Kiso, M.; Hasegawa, A. *Glycoconjugate J.* **1996**, *13*, 241–254.

(15) (a) Ravenscroft, M.; Roberts, R. M. G.; Tillett, J. G. *J. Chem. Soc., Perkin Trans. 2* **1982**, 1569–1572. (b) Fugedi, P.; Garegg, P. *J. Carbohydr. Res.* **1986**, *149*, C9–C12.

(16) Veeneman, G. H.; van Leeuwen, S. H.; van Boom, J. H. *Tetrahedron Lett.* **1990**, *31*, 1331–1334.




FIGURE 2. Outline of renewed synthetic approach for ganglioside GQ1b and its analogs.

TABLE 1. Preparation of the Key Building Block 6, Sialyl α (2 \rightarrow 8)sialyl α (2 \rightarrow 3)galactose Unit

4 [1.0 eq] + 5 [2.0 eq] $\xrightarrow{\text{See Table 1}}$ 6

entry	promoter	solvent	T (°C)	(2 \rightarrow 3) glycoside		(2 \rightarrow 4) glycoside	2,3-ene of donor
				α	β		
1	DMTST ^b	MeCN	-35	trace	trace	trace	trace
2	NIS-TfOH	MeCN	-35	33	17	10	22
3	NIS-BF ₃ ·OEt ₂	MeCN	-35	21	19	11	23
4	NIS-TMSOTf	MeCN	-35	31	15	9	20
5	NIS-TfOH	EtCN	-50	44	14	8	16

^a Isolated yield. ^b DMTST: dimethyl(methylthio)sulfonium triflate.

NIS-TfOH¹⁶ as a promoter system with the assistance of the solvent effect of acetonitrile,¹⁷ the desired α -sialoside **6** was obtained in 33% yield with a scarcely better α/β -ratio (2/1). However, adverse side reactions also occurred and resulted in a 2,3-ene derivative of the donor as a byproduct. In entry 3, the use of BF₃·OEt₂ instead of TfOH led to a reduction in both the yield of **6** and its stereoselectivity. On the other hand, the use of NIS-TMSOTf (entry 4) as a promoter system gave results similar to NIS-TfOH. In entry 5, when the reaction was conducted in the presence of NIS-TfOH in propionitrile at -50 °C, the desired **6** was obtained in a rewarding 44% yield, and

stereoselectivity was improved with a slight reduction of the byproduct. The α -configuration of the glycoside of **6** was assigned according to empirical rules¹⁸ and was further confirmed by observation of a strong cross peak between H³-3ax and C¹-1 in the HMBC spectrum¹⁹ of **6**, because the α -sialosides has a larger heteronuclear coupling constant (³J_{C-1,H-3ax}) than β -sialoside (see the Supporting Information).

The difficulty of introducing a β -galactosaminyl glycoside at the C-4 position of galactose has been revealed in the literature²⁰ in relation to the assembly of a ganglio-series

(17) (a) Schmidt, R. R.; Rücker, E. *Tetrahedron Lett.* **1980**, *21*, 1421-1424. (b) Hashimoto, S.; Hayashi, M.; Noyori, R. *Tetrahedron Lett.* **1984**, *25*, 1379-1382. (c) Murase, T.; Kameyama, A.; Kartha, K. P. R.; Ishida, H.; Kiso, M.; Hasegawa, A. *J. Carbohydr. Chem.* **1989**, *8*, 265-283.

(18) (a) van der Vleugel, D. J. M.; van Heeswijk, W. A. R.; Vliegthart, J. F. G. *Carbohydr. Res.* **1982**, *102*, 121-130. (b) Paulsen, H.; Tietz, H. *Carbohydr. Res.* **1984**, *125*, 47-64.

(19) Xia, J.; Alderfer, J. L.; Piskorz, C. F.; Matta, K. L. *Chem.-Eur. J.* **2001**, *7*, 356-367.

SCHEME 1. Preparation of Novel GalNAc Glycosyl Donors





TABLE 2. Galactosaminidations of 6 and Subsequent Conversion into Tetrasaccharide Acceptor 11



entry	donor	promoter	solvent	T (°C)	% yield ^a
1	8	NIS-TfOH	CH ₂ Cl ₂	0	58
2	8	NIS-TfOH	CH ₂ Cl ₂	-35	50
3	9	SnCl ₂ -AgClO ₄	CH ₂ Cl ₂	rt	52
4	9	HNTf ₂	BTF-BuCN ^b	rt	trace
5	9	Cp ₂ ZrCl ₂ -AgOTf	CH ₂ Cl ₂	-20	85

^a Isolated yield. ^b BTF: benzotrifluoride.

ganglioside skeleton, and the choice of the neighboring group of the anomeric center and the mode of protection for GalNAc has been examined in several ways.²¹ In this study, two kinds of GalNAc donors **8** and **9**, which commonly carry a 2,2,2-trichloroethoxycarbonyl (Troc) group at C-2, were designed, as shown in Scheme 1. It is known that the Troc group should ensure β -selective glycosylation as well as increased reactivity as a glycosyl donor compared with the *N*-phthaloyl group.²² Additionally, the advantage of the Troc group lies in the high degree of efficiency in installment and chemoselective deprotection.

Acetalization of the previously described phenylthio-galactosaminide **7**²³ with PhCH(OMe)₂ and (\pm)-10-camphorsulfonic acid (CSA) and subsequent protection of the remaining C-3-hydroxyl group by the Troc group afforded novel GalNAc thioglycoside donor **8** in 90% yield over two steps. Thiogly-

coside **8** was then efficiently converted into fluoride **9** as another GalNAc donor.²⁴ In the design of GalNAc donors, the Troc group was also exploited for the protection of the C-3 hydroxyl group to facilitate access to the tetrasaccharide acceptor for the next coupling reaction. The obtained GalNAc donors **8** and **9** were subjected to galactosaminidation with **6**. The results are summarized in Table 2.

In entries 1 and 2, the glycosidations of thioglycoside **8** with **6** were carried out in the presence of NIS-TfOH in CH₂Cl₂, affording the tetrasaccharide **10** in moderate yield along with several byproducts. Although the byproducts were inseparable, the mass spectrum of the mixture of byproducts indicated the presence of a molecule with the same molecular weight of **10**, probably α -glycoside. This result is in accordance with the report²⁵ that the *galacto*-type donor bearing the benzylidene group at the C-4,6 positions gave α -glycoside, which is unusual even in the presence of the neighboring participating acyl group at C-2. Next, in entries 3–5, the glycosidation of fluoride donor **9** in the presence of various promoters was attempted. When SnCl₂-AgClO₄²⁶ was used (entry 3), the yield of **10** was not improved, and the stereoselectivity did not change appreciably. The use of HNTf₂²⁷ in the mixture solvent (BTF/BuCN) hardly gave **10** (entry 4). On the other hand, when the reaction was conducted in the presence of Cp₂ZrCl₂-AgOTf²⁸ at -20 °C (entry 5), both yield and stereoselectivity were dramatically improved to afford **10** in 85% yield. Furthermore, it was found

(20) (a) Wessel, H. P.; Iversen, T.; Bundle, D. R. *Carbohydr. Res.* **1984**, *130*, 5–21. (b) Sugimoto, M.; Horisaki, T.; Ogawa, T. *Glycoconjugate J.* **1985**, *2*, 11–15. (c) Paulsen, H.; Paal, M. *Carbohydr. Res.* **1985**, *137*, 39–62. (d) Stauch, T.; Greilich, U.; Schmidt, R. R. *Liebigs Ann.* **1995**, 2101–2111.

(21) (a) Banoub, J.; Boullanger, P.; Lafont, D. *Chem. Rev.* **1992**, *92*, 1167–1195. (b) Matsuzaki, Y.; Nunomura, S.; Ito, Y.; Sugimoto, M.; Nakahara, Y.; Ogawa, T. *Carbohydr. Res.* **1993**, *242*, C1–C6. (c) Greilich, U.; Brescello, R.; Jung, K.-H.; Schmidt, R. R. *Liebigs Ann.* **1996**, 663–672. (d) Castro-Palmino, J. C.; Ritter, G.; Fortunato, S. R.; Reinhardt, S.; Old, L. J.; Schmidt, R. R. *Angew. Chem., Int. Ed.* **1997**, *36*, 1998–2001. (e) Castro-Palmino, J. C.; Schmidt, R. R. *Tetrahedron Lett.* **1995**, *36*, 6871–6874. (f) Debenham, J. S.; Madsen, R.; Robert, C.; Fraser-Reid, B. *J. Am. Chem. Soc.* **1995**, *117*, 3302–3303. (g) Jiao, H.; Hindsgaul, O. *Angew. Chem., Int. Ed.* **1999**, *38*, 346–348. (h) Bhattacharya, S. K.; Danishefsky, S. J. *J. Org. Chem.* **2000**, *65*, 144–151. (i) Liao, W.; Piskorz, C. F.; Locke, R. D.; Matta, K. L. *Bioorg. Med. Chem. Lett.* **2000**, *10*, 793–795.

(22) (a) Dullenkopf, W.; Castro-Palmino, J. C.; Manzoni, L.; Schmidt, R. R. *Carbohydr. Res.* **1996**, *296*, 135–147. (b) Ellervik, U.; Magnusson, G. *Carbohydr. Res.* **1996**, *280*, 251–260.

(23) Imamura, A.; Ando, H.; Ishida, H.; Kiso, M. *Org. Lett.* **2005**, *7*, 4415–4418.


(24) Nicolau, K. C.; Dolle, R. E.; Papahatjis, D. P.; Randall, J. L. *J. Am. Chem. Soc.* **1984**, *106*, 4189–4192.

(25) (a) Yule, J. E.; Wong, T. C.; Ganghi, S. S.; Qiu, D.; Riopel, M. A.; Koganty, R. R. *Tetrahedron Lett.* **1995**, *36*, 6839–6842. (b) Bélot, F.; Jacquinet, J.-C. *Carbohydr. Res.* **2000**, *325*, 93–106. (c) Chen, L.; Kong, F. *Tetrahedron Lett.* **2003**, *44*, 3691–3695.


(26) Mukaiyama, T.; Murai, Y.; Shoda, S. *Chem. Lett.* **1981**, 431–432.

(27) (a) Mukaiyama, T.; Maeshima, H.; Jona, H. *Chem. Lett.* **2001**, 426–427. (b) Jona, H.; Mandai, H.; Chavasiri, W.; Takeuchi, K.; Mukaiyama, T. *Bull. Chem. Soc. Jpn.* **2002**, *75*, 291–309.

SCHEME 2. Conversion of 6 into Donor Form 15



SCHEME 3. Glycosylation of 11 with 15 and Conversion into GQ1b Epitope Donor 20



that these conditions were applicable in a large-scale synthesis. As expected, the obtained tetrasaccharide **10** could be efficiently converted into tetrasaccharide acceptor **11** through the removal of the Troc groups by treatment with preactivated zinc, followed by the selective acetylation of the liberated amine of the GalNAc residue at C-2.

Scheme 2 shows the facile conversion of the key common unit **6** into donor form **15**. Thus, hydrogenolysis of the benzyl groups afforded the triol **12** in excellent yield, and subsequent benzylation of the exposed three hydroxyl groups successfully proceeded to give **13** in 85% yield. In order to introduce a leaving group at the reducing end of **13**, the SE group was removed by exposure to trifluoroacetic acid.²⁹ The obtained hemiacetal **14** was then converted into α -trichloroacetimidate **15** by treatment with CCl_3CN and DBU.³⁰

Next, as shown in Scheme 3, the coupling reaction of **15** and **11** was conducted. The reaction was performed in the presence of TMSOTf in CH_2Cl_2 at 0°C . This block coupling

surprisingly afforded the desired GQ1b-core heptasaccharide **16** in excellent yield (93%). The reaction was also successful in large scale (~10 g). Undoubtedly, the branched tetrasaccharide acceptor **11** performed excellent couplings with other glycosyl donors. As shown in Scheme 4, the glycosylation of **11** with the known sialyl $\alpha(2\rightarrow3)$ galactose donor **21**³¹ and galactose donor **22**³² resulted in the delivery of other glycan structures of b-series gangliosides, **23** (GT1b-core hexasaccharide) and **24** (GD1b-core saccharide) in excellent 95% and 90% yields, respectively. These results demonstrated that **11** was the expedient common unit for the synthesis of the entire b-series gangliosides.

Referring back to Scheme 3, the cleavage of benzyl groups and the benzyldene group of **16** was executed by hydrogenolysis on Pd(OH)_2 , and the following benzylation of the resulting hydroxyl groups gave **18**. In this benzylation, the combination of Bz_2O and DMAP in pyridine could not completely protect

(28) (a) Matsumoto, T.; Maeta, H.; Suzuki, K.; Tsuchihashi, G. *Tetrahedron Lett.* **1988**, *29*, 3567–3570. (b) Matsumoto, T.; Maeta, H.; Suzuki, K.; Tsuchihashi, G. *Tetrahedron Lett.* **1988**, *29*, 3571–3574. (c) Nicolaou, K. C.; Caulfield, T. J.; Kataoka, H.; Stylianides, N. A. *J. Am. Chem. Soc.* **1990**, *112*, 3693–3695.


(29) Jansson, K.; Ahlfors, S.; Frejd, T.; Kihlberg, J.; Magnusson, G.; Dahmén, J.; Noori, G.; Stenvall, K. *J. Org. Chem.* **1988**, *53*, 5629–5647.

(30) (a) Grundler, G.; Schmidt, R. R. *Liebigs Ann. Chem.* **1984**, 1826–1847. (b) Schmidt, R. R. *Angew. Chem., Int. Ed.* **1986**, *25*, 212–235. (c) Schmidt, R. R.; Michel, J. *Angew. Chem., Int. Ed.* **1980**, *19*, 731–732.

(31) Hasegawa, A.; Suzuki, N.; Ishida, H.; Kiso, M. *J. Carbohydr. Chem.* **1996**, *15*, 623–637. Physical data of compound **21** was described in the Supporting Information.

(32) Nikolaev, A. V.; Watt, G. M.; Ferguson, M. A. J.; Brimacombe, J. S. *J. Chem. Soc., Perkin Trans. I* **1997**, 969–979.

SCHEME 4. Potential as an Acceptor of 11 for the Systematical Synthesis of B-Series Gangliosides



the hydroxyl groups. Therefore, to complete the reaction, BzCl and AgOTf were added, which generated more reactive BzOTf in situ, whereas the addition of BzCl and DMAP could not complete the reaction. The use of BzOTf from the beginning of the reaction gave a crude mixture of *O*- and *N*-benzoylated products. The reason underlying this phenomenon is yet unclear. Liberation of the anomeric hydroxyl group of 18 was achieved by treatment with TFA, and the resulting hemiacetal 19 was then converted into the α -trichloroacetimidate 20, which was ready for the final glycosylation.

Hashimoto et al. reported the useful convergent synthesis of ganglioside GM3 using a glucosylceramide (GlcCer) building block equipped with benzyl groups at the C-3 and C-6 positions of the Glc residue.³³ However, the use of benzyl groups impaired the efficiency of the deprotection steps, since conventional hydrogenolysis of benzyl groups cannot be utilized due to the presence of the olefin functionality at the ceramide moiety. As a result, it has caused great loss of the samples at the final stage. Therefore, in this study, we envisioned the use of the *p*-methoxybenzyl (PMB) group as a surrogate of the benzyl group, since it can be removed by nonreductive conditions without affecting olefin functionality.

Starting from the known glucose derivative 25,³⁴ the GlcCer acceptor with PMB groups at the C-3 and C-6 positions (36) was prepared via eight steps (Scheme 5). Derivative 25 was subjected to selective *p*-methoxybenzylzation by way of stannylation³⁵ with DBTO in toluene, followed by treatment of PMBCl with TBAB to afford the 3-PMB and 2-PMB derivatives in 61% and 35% yields, respectively. Benzoylation of 26 was achieved under standard conditions to give 28 in almost a quantitative yield. Reductive opening of the anisylidene ring of 28 by treatment with NaBH₃CN and TFA in DMF³⁶

afforded the 6-*O*-PMB derivative 29 (89%), along with a small amount (8%) of the 4-*O*-PMB derivative 30. Monochloroacetylation of 29 proceeded smoothly to provide 31 in 95% yield. Selective exposure of the anomeric hydroxyl group of 31 was executed by treatment of polymethylhydrosiloxane (PMHS), Pd(PPh₃)₄, and ZnCl₂ in THF,³⁷ which afforded the hemiacetal 32 in 85% yield. The obtained hemiacetal 32 was converted into the corresponding trichloroacetimidate 33 in 93% ($\alpha/\beta = 40/53$) yield upon reaction with CCl₃CN and DBU. The α -imidate 33 α was then subjected to glycosylation with the known ceramide acceptor 34³⁸ in the presence of TMSOTf in CH₂Cl₂ to afford 35 in 48% yield. The selective deprotection of the monochloroacetyl group was accomplished by treatment of DABCO in the mixed solvent of EtOH/1,2-dichloroethane³⁹ to furnish the novel GlcCer acceptor 36 in 96% yield.

The final couplings of GQ1b-core heptasaccharide donor 20 (1.0 equiv) with glucosyl acceptors 36, 37, 38, and 39 (1.5 equiv each) were conducted in the presence of TMSOTf (0.1 equiv) in CH₂Cl₂ at 0°C, and the results are summarized in Table 3. In entry 1, the glycosylation of the novel GlcCer acceptor 36 with 20 afforded the desired fully protected ganglioside GQ1b 40 in an excellent 91% yield. On the other hand, a Hashimoto-type GlcCer acceptor 37 bearing the corresponding benzyl groups at the C-3 and C-6 positions afforded 41 in 73% yield (entry 2). These results indicated that the PMB groups at the C-3 and C-6 positions of the Glc residue enhanced the reactivity of the 4-hydroxyl group owing to its electron-donating property and elevated the coupling yield. Next, the previously described glucose derivative 38⁴⁰ was used as a coupling partner of 20 for the preparation of the SE-type GQ1b derivative (entry 3). As a result, the desired compound 42 was obtained in 63% yield.

(37) Chandrasekhar, S.; Reddy, Ch. R.; Rao, R. J. *Tetrahedron* 2001, 57, 3435–3438.

(38) (a) Kiso, M.; Nakamura, A.; Tomita, Y.; Hasegawa, A. *Carbohydr. Res.* 1986, 158, 101–111. (b) Kiso, M.; Nakamura, A.; Nakamura, J.; Tomita, Y.; Hasegawa, A. *J. Carbohydr. Chem.* 1986, 5, 335–340.

(39) Lefebvre, D. J.; Kamerling, J. P.; Vliegthart, J. F. G. *Org. Lett.* 2000, 2, 701–703.

(40) Fuse, T.; Ando, H.; Imamura, A.; Sawada, N.; Ishida, H.; Kiso, M.; Ando, T.; Li, S.-C.; Li, Y.-T. *Glycoconjugate J.* 2006, 23, 329–343.


(33) Sakamoto, H.; Nakamura, S.; Tsuda, T.; Hashimoto, S. *Tetrahedron Lett.* 2000, 41, 7691–7695.

(34) Aspinall, G. O.; Gammon, D. W.; Sood, R. K.; Chatterjee, D.; Rivoire, B.; Brennan, P. J. *Carbohydr. Res.* 1992, 237, 57–77.

(35) Qin, H.; Grindley, B. J. *Carbohydr. Chem.* 1996, 15, 95–108.

(36) Johansson, R.; Samuelsson, B. J. *Chem. Soc., Perkin Trans. 1* 1984, 2371–2374.

SCHEME 5. Preparation of a Novel Glucosyl Ceramide Acceptor 36



In entry 4, glycosylation of the previously described gentiobiose acceptor **39**¹³ with **20** provided the nonasaccharide **43** in 80% yield. These results strongly suggested that our synthetic strategy is applicable to the assembly of both natural gangliosides and unnatural ganglioside derivatives.

Scheme 6 incorporated global deprotection steps toward target compounds. Selective removal of the PMB groups of **40** was performed in the presence of TFA in CH₂Cl₂ at 0 °C, and the reaction was completed within 1 h. After a facile workup and purification by silica gel column chromatography, the target compound **44** was obtained in almost a quantitative yield (97%). The outcome demonstrated that the use of the PMB group was effective for the synthesis of natural gangliosides. Importantly, this strategy can be applied to the syntheses of compounds with hydrogenolysis-sensitive functionalities, such as allyls, alkynes, azides, and so on. Furthermore, the sustainability of azide and alkyne groups enables their application to 1,3-dipolar cycloadditions. Finally, the conventional deacylation⁴¹ and subsequent saponification gave natural ganglioside GQ1b **1** in 97% yield. It is noteworthy that the synthesis of the complex ganglioside GQ1b was first accomplished in the scale of ca. 80 mg per batch. Global deprotections of **42** and **43** were also executed by conventional methods to provide the targeted unnatural GQ1b derivatives **2** and **3** in good yields, respectively.

Conclusion

In conclusion, we have accomplished the highly efficient and practical synthesis of complex ganglioside GQ1b and its unnatural derivatives by convergent synthetic routes, based on the key coupling reactions of a GQ1b-core heptasaccharide and derivatized-Glc building blocks. In particular, a judicious choice of a PMB group as a protecting group at the C-3 and C-6





positions of the GlcCer led to not only the elevation of coupling yields but also to facilitate global deprotection with almost quantitative yields. Furthermore, our approach can efficiently reach various unnatural ganglioside derivatives with azide or alkyne groups for 1,3-dipolar cycloaddition. We are currently undertaking the syntheses of various gangliosides using this approach, and the extension of synthesized ganglioside derivatives to the chemical biology field is in progress.

Experimental Section

Ganglioside GQ1b (1). To a mixture of **44** (122 mg, 0.0316 mmol) in MeOH (6.2 mL) was added catalytic amounts of sodium methoxide (28% solution in MeOH). After stirring for 24 h at room temperature as the reaction was monitored by TLC (1-BuOH/MeOH/5% CaCl₂ (aq) = 2:1:1), the mixture was heated at 50 °C, and stirring was continued for 2 days at 50 °C. Then, H₂O (0.3 mL) was added to the mixture. After stirring for 5 h at room temperature as the reaction was monitored by TLC (1-BuOH/MeOH/5% CaCl₂ (aq) = 2:1:1), the reaction was neutralized with IR-120 (H⁺) resin and filtered through cotton, and the resin was washed with mixed solvent (CHCl₃/MeOH/H₂O = 5:5:1). The combined filtrate and washings were concentrated. The resulting residue was purified by gel filtration column chromatography on Sephadex LH-20 using CHCl₃/MeOH/H₂O (5:5:1) as the eluent to give **1** (77 mg, 97%): [α]_D = -6.3° (c 0.3, CHCl₃/MeOH/H₂O = 5:5:1); ¹H NMR (600 MHz, CDCl₃/CD₃OD/D₂O = 5:5:1) δ 5.70 (m, 1 H, J_{4,5} = 15.1 Hz, J_{5,6} = 8.2 Hz, J_{5,6'} = 6.8 Hz, H-5^{Cer}), 5.43 (dd, 1 H, J_{4,5} = 15.1 Hz, J_{3,4} = 7.5 Hz, H-4^{Cer}), 4.88 (d, 1 H, J_{1,2} = 8.2 Hz, anomer H), 4.61 (d, 1 H, anomer H), 4.46 (d, 1 H, J_{1,2} = 7.5 Hz, anomer H), 4.34 (d, 1 H, J_{1,2} = 7.5 Hz, anomer H), 3.02, 2.80, 2.74, and 2.63 (4 br dd, 4 H, 4 H-3eq), 2.17 (t, 1 H, NHCOCH₂), 2.06, 2.03, 2.03, and 2.02 (4 s, 17 H, 5 NAc, H-6^{Cer}, and 6^{Cer}), 1.84 (br t, 1 H, H-3ax), 1.69 (m, 3 H, 3 H-3ax), 1.57 (m, 1 H, NHCOCH₂), 1.36 (m, 1 H, H-7^{Cer}), 1.27 (m, 51 H, H-7^{Cer}, 25 CH₂), 0.89 (t, 6 H, 2 Me); ¹³C NMR (150 MHz, CDCl₃/CD₃OD/D₂O = 5:5:1) δ 174.8, 174.0, 173.8, 173.7, 173.4, 173.3, 172.8, 172.6, 172.4, 172.3, 133.6, 128.8, 103.7, 102.7, 102.2, 101.9, 100.7, 100.1, 100.1, 99.0, 80.5, 78.5,

(41) Zemplén, G. *Ber. Dtsch. Chem. Ges.* 1927, 60, 1555–1564.

TABLE 3. Final Glycosylations

Entry	Acceptor	Product	% Yield ^[a]
$\begin{array}{c} \text{20} \\ [1.0 \text{ eq}] \end{array} + \begin{array}{c} \text{36, 37, 38, 39} \\ [1.5 \text{ eq}] \end{array} \xrightarrow[\text{0}^\circ\text{C}]{\begin{array}{c} \text{TMSOTf} \\ \text{AW-300} \\ \text{CH}_2\text{Cl}_2 \end{array}}$ Fully protected GQ1b and unnatural GQ1b derivatives			
1	36		91
2	37		73
3	38		63
4	39		80

^a Isolated yield.


74.3, 73.9, 73.8, 73.8, 73.5, 73.4, 73.3, 73.2, 73.0, 72.8, 72.5, 72.4, 70.9, 70.8, 69.1, 69.0, 68.9, 68.9, 68.3, 68.1, 68.0, 68.0, 67.9, 67.3, 67.3, 67.0, 66.6, 66.2, 66.1, 65.5, 65.5, 64.9, 62.6, 62.5, 61.2, 61.1, 61.0, 60.9, 60.7, 60.6, 60.5, 59.7, 59.6, 59.3, 59.3, 52.7, 52.4, 52.1, 51.9, 51.7, 50.9, 40.3, 37.9, 35.5, 31.5, 31.0, 31.0, 28.8, 28.8, 28.7, 28.7, 28.6, 28.6, 28.4, 28.4, 25.2, 22.2, 21.7, 21.4, 21.3, 21.2, 21.0, 12.7; HRMS (ESI) m/z found $[M - 2H]^+$ 1208.5783, $C_{106}H_{182}N_6O_{55}$ calcd for $[M - 2H]^+$ 1208.5736.

2-(Trimethylsilyl)ethyl (5-Acetamido-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylonic acid)-(2 \rightarrow 8)-(5-acetamido-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylonic acid)-(2 \rightarrow 3)- β -D-galactopyranosyl-(1 \rightarrow 3)-2-acetamido-2-deoxy- β -D-galactopyranosyl-(1 \rightarrow 4)-[(5-acetamido-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylonic acid)-(2 \rightarrow 8)-(5-acetamido-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylonic acid)-(2 \rightarrow 3)]- β -D-galactopyranosyl-(1 \rightarrow 4)- β -D-glucopyranoside (2). To a mixture of **45** (40 mg, 0.0121 mmol) in MeOH (1.0 mL) was added catalytic amounts of sodium methoxide (28% solution in MeOH). After stirring for 17 h at room temperature as the reaction was monitored by TLC (1-BuOH/MeOH/5% $CaCl_2$ (aq) = 2:1:1), the mixture was heated at 50 $^\circ C$, and the stirring was continued for 17 h at 50 $^\circ C$. Then, 0.2 M KOH (0.5 mL) was added to the mixture. After stirring for 30 h at 75 $^\circ C$ as the reaction was monitored by TLC (1-BuOH/MeOH/5% $CaCl_2$ (aq) = 2:1:1), the reaction was neutralized with IR-120 (H^+) resin and filtered through cotton, and the resin was washed with MeOH and H_2O . The

combined filtrate and washings were concentrated. The resulting residue was purified by gel filtration column chromatography on Sephadex LH-20 using H_2O as the eluent to give **2** (20 mg, 87%): $[\alpha]_D^{25} = -5.0^\circ$ (c 0.7, H_2O); 1H NMR (500 MHz, $D_2O/(CD_3)_2CO = 10:1$) δ 4.73 (d, 1 H, H-1h), 4.59 (d, 1 H, $J_{1,2} = 7.4$ Hz, H-1e), 4.49 (d, 1 H, $J_{1,2} = 8.6$ Hz, H-1c), 4.47 (d, 1 H, $J_{1,2} = 8.0$ Hz, H-1d), 3.41 (t, 1 H, $J_{1,2} = J_{2,3} = 8.6$ Hz, H-2c), 3.25 (near t, 1 H, $J_{1,2} = 8.0$ Hz, $J_{2,3} = 9.1$ Hz, H-2d), 2.77–2.63 (m, 4 H, H-3eq-a, 3eq-b, 3eq-f, and 3eq-g), 2.05, 2.04, and 2.01 (3 s, 15 H, 5 NAc), 1.76–1.69 (m, 4 H, H-3ax-a, 3ax-b, 3ax-f, and 3ax-g), 1.07–0.91 (m, 2 H, $OCH_2CH_2SiMe_3$), 0.00 (s, 9 H, $OCH_2CH_2SiMe_3$); ^{13}C NMR (125 MHz, $D_2O/(CD_3)_2CO = 10:1$) δ 175.2, 175.1, 175.0, 173.9, 173.6, 173.6, 173.5, 104.5, 103.1, 102.8, 101.7, 100.8, 100.6, 100.5, 100.5, 80.2, 78.8, 78.6, 75.6, 75.5, 75.1, 74.9, 74.7, 74.6, 74.5, 74.3, 74.0, 73.1, 72.9, 72.0, 69.8, 69.8, 69.7, 69.5, 68.8, 68.7, 68.5, 68.3, 68.3, 67.9, 67.6, 62.9, 61.9, 61.8, 61.3, 61.2, 60.9, 60.3, 52.6, 52.5, 52.1, 51.4, 40.9, 40.8, 39.9, 39.8, 22.9, 22.6, 22.6, 22.3, 17.8, -2.1; HRMS (ESI) m/z found $[M - 4H]^+$ 491.9194, $C_{75}H_{125}N_5O_{33}Si$ calcd for $[M - 4H]^+$ 491.9192.

(5-Acetamido-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylonic acid)-(2 \rightarrow 8)-(5-acetamido-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylonic acid)-(2 \rightarrow 3)- β -D-galactopyranosyl-(1 \rightarrow 3)-2-acetamido-2-deoxy- β -D-galactopyranosyl-(1 \rightarrow 4)-[(5-acetamido-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylonic acid)-(2 \rightarrow 8)-(5-acetamido-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylonic acid)-(2 \rightarrow 3)]- β -D-galacto-

SCHEME 6. Global Deprotections



pyranosyl-(1→4)-β-D-glucopyranosyl-(1→6)-D-glucopyranose (3). To a mixture of **43** (46 mg, 0.0118 mmol) in MeOH (0.8 mL) was added catalytic amounts of sodium methoxide (28% solution in MeOH). After stirring for 12 h at room temperature as the reaction was monitored by TLC (1-BuOH/MeOH/5% CaCl₂ (aq) = 2:1:1), the mixture was heated at 50 °C, and the stirring was continued for 24 h at 50 °C. Then 0.2 M KOH (0.2 mL) was added to the mixture. After stirring for 3 h at room temperature as the reaction was monitored by TLC (1-BuOH/MeOH/5% CaCl₂ (aq) = 2:1:1), the reaction was neutralized with IR-120 (H⁺) resin and filtered through cotton, and the resin was washed with H₂O. The combined filtrate and washings were concentrated. The resulting residue was purified by gel filtration column chromatography on Sephadex LH-20 using H₂O as the eluent. Then, to a solution of the obtained white solid **46** in EtOH/H₂O (1.0 mL/1.0 mL) was added palladium hydroxide [20 wt % Pd (dry basis) on carbon, wet] (90 mg) at room temperature. After stirring for 14 h at room temperature under a H₂ atmosphere as the reaction was monitored by TLC (1-BuOH/MeOH/5% CaCl₂ (aq) = 1:1:1), the mixture was filtered through a membrane filter and washed with H₂O. The combined filtrate and washings were concentrated. The resulting residue was purified by gel filtration column chromatography on Sephadex LH-20 using H₂O as the eluent to give **3** (20 mg, 83%): [α]_D = -3.7° (c 0.4, H₂O); ¹H NMR (600 MHz, D₂O) δ 5.20 (d, 1 H, J_{1,2} = 3.4 Hz, H-1 of Glc unit), 4.71 (m, anomer H), 4.63 (d, 1 H, J_{1,2} = 8.2 Hz, H-1 of Glc unit), 4.58 (m, anomer H), 4.52 (d, 1 H, J_{1,2} = 8.2 Hz, anomer H), 4.50 (d, 1 H, J_{1,2} = 7.5 Hz, anomer H), 4.47 (d, 1 H, J_{1,2} = 8.2 Hz, anomer H), 3.51 (dd, 1 H, J_{1,2} = 3.4 Hz, J_{2,3} = 10.3 Hz, H-2 of Glc unit), 3.44 (t, 1 H, J_{2,3} = J_{3,4} = 8.9 Hz, H-3 of Glc unit), 3.40–3.30 (m, 3 H, 3 H-2), 3.22 (near t, 1 H, J_{1,2} = 8.2 Hz, J_{2,3} = 8.9 Hz, H-2), 2.76–2.62 (m, 4 H, H-3eq-a, 3eq-b, 3eq-f,

and 3eq-g), 2.04, 2.03, 2.02, and 2.00 (4 s, 15 H, 5 NAc), 1.78–1.69 (m, 4 H, H-3ax-a, 3ax-b, 3ax-f, and 3ax-g); ¹³C NMR (150 MHz, D₂O/(CD₃)₂CO = 10:1) δ 165.9, 164.8–163.9, 95.3, 93.8, 93.6, 93.5, 92.0–90.9, 87.0, 83.1, 71.0, 69.2, 69.1, 66.7, 66.4, 65.9, 65.8, 65.8, 65.6, 65.3, 65.2, 65.1, 64.8, 63.7, 62.7, 62.4, 61.4, 60.6, 60.5, 60.4, 60.2, 59.8, 59.8, 59.6, 59.4, 59.3, 59.2, 59.0, 58.6, 58.6, 53.6, 52.5, 52.4, 52.0, 52.0, 51.6, 51.0, 43.3, 43.2, 42.8, 42.1, 31.6, 31.5, 30.7, 30.2, 13.6, 13.4, 13.3, 13.1; HRMS (ESI) *m/z* found [M - 4H]⁴⁺ 507.4145, C₇₆H₁₂₃N₅O₅₈ calcd for [M - 4H]⁴⁺ 507.4143.

2-(Trimethylsilyl)ethyl [Methyl 5-acetamido-8-O-(5-acetamido-4,7,8,9-tetra-O-acetyl-3,5-dideoxy-D-glycero-α-D-galacto-2-nonulopyranosylono-1',9-lactone)-4,7-di-O-acetyl-3,5-dideoxy-D-glycero-α-D-galacto-2-nonulopyranosylonate]-(2→3)-2,6-di-O-benzyl-β-D-galactopyranoside (6). To a mixture of **4** (200 mg, 210 μmol) and **5** (196 mg, 420 μmol) in propionitrile (4.0 mL) were added 3 Å molecular sieves (400 mg) at room temperature. After stirring for 1 h, NIS (96 mg, 420 μmol) was added to the mixture. After cooling to -50 °C, TfOH (11 μL, 130 μmol) was added to the mixture. After stirring for 96 h at -50 °C as the reaction was monitored by TLC (EtOAc/MeOH = 10:1), reagents (NIS, 2 × 96 mg, 192 mg in total; TfOH, 2 × 11 μL, 22 μL in total) were added to the mixture after 120 and 144 h. The mixture was stirred for a total of 168 h at the same temperature. Then, the reaction was quenched by triethylamine and filtered through a Celite pad, and the pad was washed with CHCl₃. The combined filtrate and washings were evaporated. The residue was extracted with CHCl₃ and washed with saturated Na₂CO₃ (aq), saturated Na₂S₂O₃ (aq), and brine, dried over Na₂SO₄, and concentrated. The residue was purified by flash column chromatography twice (first with CHCl₃/MeOH = 50:1 and second with EtOAc) to give **6** (120

mg, 44%) and its β -isomer **6b** (39 mg, 14%). **6a**: $[\alpha]_D = -28.1^\circ$ (*c* 3.7, CHCl₃); ¹H NMR (500 MHz, CDCl₃) δ 7.35–7.24 (m, 10 H, 2 Ph), 5.69 (d, 1 H, NH-a), 5.67 (d, 1 H, NH-b), 5.38 (dt, 1 H, H-4b), 5.35 (dd, 1 H, H-7b), 5.15 (dt, 1 H, H-8b), 5.09 (dd, 1 H, H-7a), 5.07 (dt, 1 H, H-4a), 4.90 and 4.62 (2 d, 2 H, $J_{gem} = 11.2$ Hz, PhCH₂), 4.56 (2 d, 2 H, $J_{gem} = 11.6$ Hz, PhCH₂), 4.40 (d, 1 H, $J_{1,2} = 8.0$ Hz, H-1c), 4.37 (dd, 1 H, H-9'a), 4.29–4.19 (m, 3 H, H-9a, 9'b, and 8a), 4.18 (q, 1 H, H-5b), 4.07–3.95 (m, 6 H, H-9b, 5a, 3c, 6a, 6b, and OCH₂CH₂SiMe₃), 3.87 (br s, 1 H, H-4c), 3.82 (s, 3 H, CO₂Me), 3.81 (dd, 1 H, H-6'c), 3.73 (dd, 1 H, H-6c), 3.61 (m, 2 H, H-5c and OCH₂CH₂SiMe₃), 3.54 (t, 1 H, $J_{1,2} = 8.0$ Hz, $J_{2,3} = 9.1$ Hz, H-2c), 2.80 (s, 1 H, OH), 2.46 (dt, 2 H, H-3eq-a and H-3eq-b), 2.14–1.88 (8 s and 2 t, 26 H, H-3ax-a, H-3ax-b, and 8 Ac), 1.03 (t, 2 H, OCH₂CH₂SiMe₃), 0.02 (s, 9 H, OCH₂CH₂SiMe₃); ¹³C NMR (100 MHz, CDCl₃) δ 170.8, 170.7, 170.6, 170.5, 170.2, 169.8, 169.7, 168.1, 165.4, 138.6, 138.1, 128.4, 128.1, 128.0, 127.7, 127.6, 127.5, 103.3, 99.4, 96.1, 75.8, 74.9, 73.6, 73.0, 72.8, 72.1, 70.7, 69.4, 69.2, 69.1, 68.9, 67.8, 67.4, 67.1, 67.1, 66.6, 61.8, 53.1, 49.3, 49.0, 38.3, 35.8, 23.1, 20.9, 20.8, 20.7, 20.6, 18.5; HRMS (ESI) *m/z* found [M + Na]⁺ 1313.4779, C₆₀H₈₂N₂O₂₇Si calcd for [M + Na]⁺ 1313.4771. **6b**: $[\alpha]_D = -31.9^\circ$ (*c* 1.0, CHCl₃); ¹H NMR (500 MHz, CDCl₃) δ 7.50–7.27 (m, 10 H, 2 Ph), 5.46 (dt, 1 H, $J_{3eq,4} = 5.4$ Hz, $J_{4,5} = 10.2$ Hz, H-4b), 5.38 (d, 1 H, NH-b), 5.35 (dd, 1 H, H-7b), 5.17–5.09 (m, 3 H, H-4a, 7a, and 8b), 5.06 and 4.62 (2 d, 2 H, $J_{gem} = 10.9$ Hz, PhCH₂), 4.58–4.52 (m, 4 H, H-9'a, PhCH₂, and H-9a), 4.45 (d, 1 H, $J_{1,2} = 7.5$ Hz, H-1c), 4.34–4.28 (m, 3 H, H-6a, 8a, and 9b), 4.17 (q, 1 H, $J_{4,5} = 10.2$ Hz, H-5b), 4.05–3.99 (m, 3 H, H-5a, 9b, and OCH₂CH₂SiMe₃), 3.83 (br s, 1 H, H-4c), 3.81–3.75 (m, 3 H, H-6b, 3c, and 6'c), 3.73 (s, 3 H, CO₂Me), 3.71 (dd, 1 H, $J_{5,6} = 9.6$ Hz, H-6c), 3.61 (m, 2 H, $J_{5,6} = 9.6$ Hz, H-5c, and OCH₂CH₂SiMe₃), 3.64 (t, 1 H, $J_{1,2} = 7.5$ Hz, H-2c), 3.62–3.58 (m, 2 H, H-5c and OCH₂CH₂SiMe₃), 3.30 (s, 1 H, OH), 2.56 (dd, 1 H, $J_{gem} = 13.7$ Hz, $J_{3eq,4} = 4.3$ Hz, H-3eq-a), 2.46 (dd, 1 H, $J_{gem} = 13.5$ Hz, $J_{3eq,4} = 5.4$ Hz, H-3eq-b), 2.12, 2.08, 2.04, 2.03, 2.02, and 2.00 (6 s, 18 H, 6 Ac), 2.02 (m, 1 H, $J_{gem} = 13.7$ Hz, H-3ax-a), 1.91 (t, 1 H, $J_{gem} = 13.5$ Hz, H-3ax-b), 1.89 and 1.73 (2 s, 6 H, 2 NAc), 1.02 (t, 2 H, OCH₂CH₂SiMe₃), 0.01 (s, 9 H, OCH₂CH₂SiMe₃); ¹³C NMR (100 MHz, CDCl₃) δ 171.0, 170.9, 170.6, 170.5, 170.3, 169.9, 169.8, 169.6, 168.2, 164.7, 138.9, 137.8, 128.5, 128.4, 128.3, 128.1, 128.1, 127.8, 127.7, 103.3, 99.3, 96.6, 77.9, 77.7, 74.9, 73.6, 72.7, 72.3, 72.2, 70.8, 69.8, 69.0, 68.7, 68.6, 68.2, 67.6, 67.5, 66.6, 61.6, 53.2, 49.0, 48.4, 37.9, 34.7, 30.3, 23.2, 23.1, 21.0, 20.9, 20.8, 20.7, 20.6, 18.5; HRMS (ESI) *m/z* found [M + Na]⁺ 1313.4773, C₆₀H₈₂N₂O₂₇Si calcd for [M + Na]⁺ 1313.4771.

2-(Trimethylsilyl)ethyl 2-Acetamido-4,6-O-benzylidene-2-deoxy- β -D-galactopyranosyl-(1 \rightarrow 4)-[methyl 5-acetamido-8-O-(5-acetamido-4,7,8,9-tetra-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylono-1',9-lactone)-4,7-di-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylono-1',9-lactone]-2(2 \rightarrow 3)-2,6-di-O-benzyl- β -D-galactopyranoside (11). To a solution of **10** (6.30 g, 3.33 mmol) in (CH₂Cl)₂ (30 mL) were added activated Zn powder (25.2 g) and acetic acid (60 mL) at room temperature. After stirring for 1 h at 40 °C as the reaction was monitored by TLC (CHCl₃/MeOH = 10:1), the mixture was filtered through a Celite pad and washed with CHCl₃. The combined filtrate and washings were diluted with CHCl₃ and washed with saturated NaHCO₃ (aq), H₂O, and brine. After drying over Na₂SO₄ and being concentrated, the resulting residue was dissolved in CH₂Cl₂ (50 mL). Acetic anhydride (471 μ L, 5.00 mmol) was then added to the mixture at room temperature. After stirring for 30 min as the reaction was monitored by TLC (CHCl₃/MeOH = 10:1), the reaction mixture was concentrated. The residue was purified by flash column chromatography using CHCl₃/MeOH (28:1 to 20:1) as the eluent to give **11** (5.04 g, 96%): $[\alpha]_D = -23.6^\circ$ (*c* 0.5, CHCl₃); ¹H NMR (500 MHz, CDCl₃) δ 7.53–7.23 (m, 15 H, 3 Ph), 6.86 (d, 1 H, NH-d), 5.59 (s, 1 H, PhCH<), 5.56 (d, 1 H, NH-b), 5.46 (near d, 1 H, NH-a), 5.40 (dt, 1 H, $J_{4,5} = 10.4$ Hz, H-4b), 5.33 (dd, 1 H, $J_{6,7} = 2.1$ Hz, H-7b), 5.12 (m, 2 H, H-8b and 4a), 4.98 and 4.51 (2

d, 2 H, $J_{gem} = 11.2$ Hz, PhCH₂), 4.96 (d, 1 H, H-4d), 4.92 (dd, 1 H, H-7a), 4.77 (d, 1 H, $J_{1,2} = 8.3$ Hz, H-1d), 4.60 and 4.52 (2 d, 2 H, $J_{gem} = 11.7$ Hz, PhCH₂), 4.44 (d, 1 H, $J_{1,2} = 7.5$ Hz, H-1c), 4.43 (dd, 1 H, $J_{gem} = 12.4$ Hz, H-9a), 4.26 (dd, 1 H, H-9'b), 4.19 (q, 1 H, $J_{4,5} = J_{5,6} = J_{5,NH} = 10.4$ Hz, H-5b), 4.17 (m, 2 H, H-2d and 4c), 4.15–3.98 (m, 14 H, H-6a, 5a, 6d, 6'd, 5d, 3c, 9b, 6'c, 8a, and OCH₂CH₂SiMe₃), 3.89 (s, 3 H, COOMe), 3.82 (dd, 1 H, $J_{gem} = 12.4$ Hz, H-9'a), 3.74 (dd, 1 H, $J_{5,6} = 10.4$ Hz, $J_{6,7} = 2.1$ Hz, H-6b), 3.70 (s, 3 H, COOMe), 3.66 (dd, 1 H, H-6c), 3.62–3.54 (m, 3 H, OCH₂CH₂SiMe₃, H-3d, and 5c), 3.51 (near t, 1 H, $J_{1,2} = 7.5$ Hz, H-2c), 3.02 (s, 1 H, OH), 2.32 (m, 3 H, H-3eq-b, 3eq-a, and 3ax-a), 2.12–1.89 (8 s, 24 H, 8 Ac), 1.81 (near t, 1 H, H-3ax-b), 1.04 (m, 2 H, OCH₂CH₂SiMe₃), 0.02 (s, 9 H, OCH₂CH₂SiMe₃); HRMS (ESI) *m/z* found [M + Na]⁺ 1604.5873, C₇₅H₉₉N₃O₃₂Si calcd for [M + Na]⁺ 1604.5878.

[Methyl 5-Acetamido-8-O-(5-acetamido-4,7,8,9-tetra-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylono-1',9-lactone)-4,7-di-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylate]-2(2 \rightarrow 3)-2,4,6-tri-O-benzoyl- α -D-galactopyranosyl Trichloroacetimidate (15). To a solution of **13** (2.50 g, 1.75 mmol) in CH₂Cl₂ (12 mL) was added trifluoroacetic acid (4.0 mL) at 0 °C. After stirring for 5 h at room temperature as the reaction was monitored by TLC (CHCl₃/MeOH = 10:1), the reaction mixture was coevaporated with toluene. The residue was purified by flash column chromatography using CHCl₃/MeOH (30:1) as the eluent to give the hemiacetal **14** (2.29 g, 98%): HRMS (ESI) *m/z* found [M + Na]⁺ 1345.3912, C₆₂H₇₀N₂O₃₀ calcd for [M + Na]⁺ 1345.3911.

To a solution of **14** (827 mg, 624 μ mol) in CH₂Cl₂ (6.2 mL) were added trichloroacetimidate (1.25 mL, 12.5 mmol) and DBU (112 μ L, 750 μ mol) at 0 °C. After stirring for 2 h at room temperature as the reaction was monitored by TLC (CHCl₃/MeOH = 10:1), the reaction mixture was concentrated. The resulting residue was purified by flash column chromatography using CHCl₃/MeOH (30:1) as the eluent to give **15** (862 mg, 94%): $[\alpha]_D = +21.4^\circ$ (*c* 1.0, CHCl₃); ¹H NMR (500 MHz, CDCl₃) δ 8.59 (s, 1 H, C=NH), 8.10–7.40 (m, 15 H, 3 Ph), 6.83 (d, 1 H, $J_{1,2} = 3.6$ Hz, H-1c), 5.94 (d, 1 H, $J_{3,4} = 3.6$ Hz, $J_{4,5} = 1.2$ Hz, H-4c), 5.68 (dd, 1 H, $J_{1,2} = 3.6$ Hz, $J_{2,3} = 10.4$ Hz, H-2c), 5.56 (dt, 1 H, $J_{3eq,4} = 5.3$ Hz, $J_{3ax,4} = 11.4$ Hz, $J_{4,5} = 10.4$ Hz, H-4b), 5.44 (d, 1 H, NH-a), 5.35 (dd, 1 H, $J_{6,7} = 2.1$ Hz, H-7b), 5.27 (d, 1 H, $J_{5,NH} = 10.4$ Hz, NH-b), 5.19 (m, 2 H, H-4a and 8b), 4.98 (dd, 1 H, $J_{6,7} = 1.7$ Hz, $J_{7,8} = 9.2$ Hz, H-7a), 4.93 (dd, 1 H, $J_{2,3} = 10.4$ Hz, $J_{3,4} = 3.6$ Hz, H-3c), 4.74 (near dt, 1 H, $J_{4,5} = 1.2$ Hz, H-5c), 4.65 (dt, 1 H, $J_{7,8} = 9.2$ Hz, $J_{8,9} = 2.6$ Hz, H-8a), 4.48–4.40 (m, 3 H, H-6'c, 9a, and 6c), 4.28 (dd, 1 H, $J_{gem} = 12.6$ Hz, H-9'b), 4.17 (q, 1 H, $J_{4,5} = J_{5,NH} = 10.4$ Hz, H-5b), 4.15 (q, 1 H, H-5a), 4.04 (dd, 1 H, $J_{8,9} = 2.6$ Hz, H-9'a), 4.02 (dd, 1 H, $J_{gem} = 12.6$ Hz, H-9b), 3.96 (dd, 1 H, $J_{6,7} = 1.7$ Hz, H-6a), 3.91 (dd, 1 H, $J_{6,7} = 2.1$ Hz, H-6b), 3.36 (s, 3 H, COOMe), 2.56 (dd, 1 H, $J_{gem} = 13.1$ Hz, $J_{3eq,4} = 5.3$ Hz, H-3eq-b), 2.24 (dd, 1 H, H-3eq-a), 2.12–1.89 (8 s and t, 25 H, 8 Ac and H-3ax-a), 1.70 (t, 1 H, $J_{gem} = 13.1$ Hz, $J_{3ax,4} = 11.4$ Hz, H-3ax-b); ¹³C NMR (100 MHz, CDCl₃) δ 170.6, 170.5, 170.4, 170.4, 170.0, 169.8, 169.8, 167.4, 166.0, 165.6, 165.2, 164.4, 160.2, 133.5, 133.3, 133.2, 129.9, 129.7, 129.5, 129.3, 129.3, 128.9, 128.5, 128.4, 98.9, 97.0, 93.9, 90.8, 77.1, 72.5, 71.9, 70.6, 69.8, 69.5, 69.5, 69.2, 68.8, 68.4, 68.3, 67.9, 67.7, 66.7, 62.9, 62.0, 52.5, 49.1, 38.2, 36.1, 23.1, 23.0, 20.7, 20.7, 20.7, 20.6, 20.4; HRMS (ESI) *m/z* found [M + Na]⁺ 1488.3004, C₆₄H₇₀Cl₃N₃O₃₀ calcd for [M + Na]⁺ 1488.3007.

2-(Trimethylsilyl)ethyl [Methyl 5-acetamido-8-O-(5-acetamido-4,7,8,9-tetra-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylono-1',9-lactone)-4,7-di-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylate]-2(2 \rightarrow 3)-2,4,6-tri-O-benzoyl- β -D-galactopyranosyl-(1 \rightarrow 3)-2-acetamido-4,6-O-benzylidene-2-deoxy- β -D-galactopyranosyl-(1 \rightarrow 4)-[methyl 5-acetamido-8-O-(5-acetamido-4,7,8,9-tetra-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylono-1',9-lactone)-4,7-di-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylate]-2(2 \rightarrow 3)-2,4,6-tri-O-benzoyl- β -D-galactopyranosyl-(1 \rightarrow 3)-2-acetamido-4,6-O-benzylidene-2-deoxy- β -D-galactopyranosyl-(1 \rightarrow 4)-[methyl 5-acetamido-8-O-(5-acetamido-4,7,8,9-tetra-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylono-1',9-lactone)-4,7-di-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylate]-2(2 \rightarrow 3)-2,4,6-tri-O-benzoyl- β -D-galactopyranosyl-(1 \rightarrow 3)-2-acetamido-4,6-O-benzylidene-2-deoxy- β -D-galactopyranosyl-(1 \rightarrow 4)-[methyl 5-acetamido-8-O-(5-acetamido-4,7,8,9-tetra-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylono-1',9-lactone)-4,7-di-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylate]-2(2 \rightarrow 3)-2,4,6-tri-O-benzoyl- β -D-galactopyranosyl-(1 \rightarrow 3)-2-acetamido-4,6-O-benzylidene-2-deoxy- β -D-galactopyranosyl-(1 \rightarrow 4)-[methyl 5-acetamido-8-O-(5-acetamido-4,7,8,9-tetra-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylono-1',9-lactone)-4,7-di-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylate]-2(2 \rightarrow 3)-2,4,6-tri-O-benzoyl- β -D-galactopyranosyl-(1 \rightarrow 3)-2-acetamido-4,6-O-benzylidene-2-deoxy- β -D-galactopyranosyl-(1 \rightarrow 4)-[methyl 5-acetamido-8-O-(5-acetamido-4,7,8,9-tetra-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylono-1',9-lactone)-4,7-di-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylate]-2(2 \rightarrow 3)-2,4,6-tri-O-benzoyl- β -D-galactopyranosyl-(1 \rightarrow 3)-2-acetamido-4,6-O-benzylidene-2-deoxy- β -D-galactopyranosyl-(1 \rightarrow 4)-[methyl 5-acetamido-8-O-(5-acetamido-4,7,8,9-tetra-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylono-1',9-lactone)-4,7-di-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylate]-2(2 \rightarrow 3)-2,4,6-tri-O-benzoyl- β -D-galactopyranosyl-(1 \rightarrow 3)-2-acetamido-4,6-O-benzylidene-2-deoxy- β -D-galactopyranosyl-(1 \rightarrow 4)-[methyl 5-acetamido-8-O-(5-acetamido-4,7,8,9-tetra-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylono-1',9-lactone)-4,7-di-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylate]-2(2 \rightarrow 3)-2,4,6-tri-O-benzoyl- β -D-galactopyranosyl-(1 \rightarrow 3)-2-acetamido-4,6-O-benzylidene-2-deoxy- β -D-galactopyranosyl-(1 \rightarrow 4)-[methyl 5-acetamido-8-O-(5-acetamido-4,7,8,9-tetra-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylono-1',9-lactone)-4,7-di-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylate]-2(2 \rightarrow 3)-2,4,6-tri-O-benzoyl- β -D-galactopyranosyl-(1 \rightarrow 3)-2-acetamido-4,6-O-benzylidene-2-deoxy- β -D-galactopyranosyl-(1 \rightarrow 4)-[methyl 5-acetamido-8-O-(5-acetamido-4,7,8,9-tetra-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylono-1',9-lactone)-4,7-di-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylate]-2(2 \rightarrow 3)-2,4,6-tri-O-benzoyl- β -D-galactopyranosyl-(1 \rightarrow 3)-2-acetamido-4,6-O-benzylidene-2-deoxy- β -D-galactopyranosyl-(1 \rightarrow 4)-[methyl 5-acetamido-8-O-(5-acetamido-4,7,8,9-tetra-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylono-1',9-lactone)-4,7-di-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylate]-2(2 \rightarrow 3)-2,4,6-tri-O-benzoyl- β -D-galactopyranosyl-(1 \rightarrow 3)-2-acetamido-4,6-O-benzylidene-2-deoxy- β -D-galactopyranosyl-(1 \rightarrow 4)-[methyl 5-acetamido-8-O-(5-acetamido-4,7,8,9-tetra-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylono-1',9-lactone)-4,7-di-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylate]-2(2 \rightarrow 3)-2,4,6-tri-O-benzoyl- β -D-galactopyranosyl-(1 \rightarrow 3)-2-acetamido-4,6-O-benzylidene-2-deoxy- β -D-galactopyranosyl-(1 \rightarrow 4)-[methyl 5-acetamido-8-O-(5-acetamido-4,7,8,9-tetra-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylono-1',9-lactone)-4,7-di-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylate]-2(2 \rightarrow 3)-2,4,6-tri-O-benzoyl- β -D-galactopyranosyl-(1 \rightarrow 3)-2-acetamido-4,6-O-benzylidene-2-deoxy- β -D-galactopyranosyl-(1 \rightarrow 4)-[methyl 5-acetamido-8-O-(5-acetamido-4,7,8,9-tetra-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylono-1',9-lactone)-4,7-di-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylate]-2(2 \rightarrow 3)-2,4,6-tri-O-benzoyl- β -D-galactopyranosyl-(1 \rightarrow 3)-2-acetamido-4,6-O-benzylidene-2-deoxy- β -D-galactopyranosyl-(1 \rightarrow 4)-[methyl 5-acetamido-8-O-(5-acetamido-4,7,8,9-tetra-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylono-1',9-lactone)-4,7-di-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylate]-2(2 \rightarrow 3)-2,4,6-tri-O-benzoyl- β -D-galactopyranosyl-(1 \rightarrow 3)-2-acetamido-4,6-O-benzylidene-2-deoxy- β -D-galactopyranosyl-(1 \rightarrow 4)-[methyl 5-acetamido-8-O-(5-acetamido-4,7,8,9-tetra-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylono-1',9-lactone)-4,7-di-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylate]-2(2 \rightarrow 3)-2,4,6-tri-O-benzoyl- β -D-galactopyranosyl-(1 \rightarrow 3)-2-acetamido-4,6-O-benzylidene-2-deoxy- β -D-galactopyranosyl-(1 \rightarrow 4)-[methyl 5-acetamido-8-O-(5-acetamido-4,7,8,9-tetra-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylono-1',9-lactone)-4,7-di-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylate]-2(2 \rightarrow 3)-2,4,6-tri-O-benzoyl- β -D-galactopyranosyl-(1 \rightarrow 3)-2-acetamido-4,6-O-benzylidene-2-deoxy- β -D-galactopyranosyl-(1 \rightarrow 4)-[methyl 5-acetamido-8-O-(5-acetamido-4,7,8,9-tetra-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylono-1',9-lactone)-4,7-di-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylate]-2(2 \rightarrow 3)-2,4,6-tri-O-benzoyl- β -D-galactopyranosyl-(1 \rightarrow 3)-2-acetamido-4,6-O-benzylidene-2-deoxy- β -D-galactopyranosyl-(1 \rightarrow 4)-[methyl 5-acetamido-8-O-(5-acetamido-4,7,8,9-tetra-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylono-1',9-lactone)-4,7-di-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylate]-2(2 \rightarrow 3)-2,4,6-tri-O-benzoyl- β -D-galactopyranosyl-(1 \rightarrow 3)-2-acetamido-4,6-O-benzylidene-2-deoxy- β -D-galactopyranosyl-(1 \rightarrow 4)-[methyl 5-acetamido-8-O-(5-acetamido-4,7,8,9-tetra-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylono-1',9-lactone)-4,7-di-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylate]-2(2 \rightarrow 3)-2,4,6-tri-O-benzoyl- β -D-galactopyranosyl-(1 \rightarrow 3)-2-acetamido-4,6-O-benzylidene-2-deoxy- β -D-galactopyranosyl-(1 \rightarrow 4)-[methyl 5-acetamido-8-O-(5-acetamido-4,7,8,9-tetra-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylono-1',9-lactone)-4,7-di-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylate]-2(2 \rightarrow 3)-2,4,6-tri-O-benzoyl- β -D-galactopyranosyl-(1 \rightarrow 3)-2-acetamido-4,6-O-benzylidene-2-deoxy- β -D-galactopyranosyl-(1 \rightarrow 4)-[methyl 5-acetamido-8-O-(5-acetamido-4,7,8,9-tetra-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylono-1',9-lactone)-4,7-di-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylate]-2(2 \rightarrow 3)-2,4,6-tri-O-benzoyl- β -D-galactopyranosyl-(1 \rightarrow 3)-2-acetamido-4,6-O-benzylidene-2-deoxy- β -D-galactopyranosyl-(1 \rightarrow 4)-[methyl 5-acetamido-8-O-(5-acetamido-4,7,8,9-tetra-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylono-1',9-lactone)-4,7-di-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylate]-2(2 \rightarrow 3)-2,4,6-tri-O-benzoyl- β -D-galactopyranosyl-(1 \rightarrow 3)-2-acetamido-4,6-O-benzylidene-2-deoxy- β -D-galactopyranosyl-(1 \rightarrow 4)-[methyl 5-acetamido-8-O-(5-acetamido-4,7,8,9-tetra-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylono-1',9-lactone)-4,7-di-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylate]-2(2 \rightarrow 3)-2,4,6-tri-O-benzoyl- β -D-galactopyranosyl-(1 \rightarrow 3)-2-acetamido-4,6-O-benzylidene-2-deoxy- β -D-galactopyranosyl-(1 \rightarrow 4)-[methyl 5-acetamido-8-O-(5-acetamido-4,7,8,9-tetra-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylono-1',9-lactone)-4,7-di-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylate]-2(2 \rightarrow 3)-2,4,6-tri-O-benzoyl- β -D-galactopyranosyl-(1 \rightarrow 3)-2-acetamido-4,6-O-benzylidene-2-deoxy- β -D-galactopyranosyl-(1 \rightarrow 4)-[methyl 5-acetamido-8-O-(5-acetamido-4,7,8,9-tetra-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylono-1',9-lactone)-4,7-di-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylate]-2(2 \rightarrow 3)-2,4,6-tri-O-benzoyl- β -D-galactopyranosyl-(1 $\$

ulopyranosylate]-(2 \rightarrow 3)-2,6-di-*O*-benzyl- β -D-galactopyranoside (16). To a mixture of **15** (2.71 g, 1.84 mmol) and **11** (1.93 g, 1.22 mmol) in CH₂Cl₂ (30 mL) was added 4 Å molecular sieves (AW-300) (4.6 g) at room temperature. After stirring for 1 h and then cooling to 0 °C, TMSOTf (6.7 μ L, 36.8 μ mol) was added to the mixture. After stirring for 22 h at 0 °C as the reaction was monitored by TLC (CHCl₃/MeOH/EtOAc = 15:1:1), the reaction was quenched by saturated NaHCO₃ (aq) and filtered through a Celite pad, and the pad was washed with CHCl₃. The combined filtrate and washings were washed with brine, dried over Na₂SO₄, and concentrated. The resulting residue was purified by flash column chromatography using CHCl₃/MeOH/EtOAc (20:1:1) to CHCl₃/MeOH (9:1) as the eluent to give **16** (3.29 g, 93%): [α]_D = -11.5° (c 1.0, CHCl₃); ¹H NMR (500 MHz, CDCl₃) δ 8.09–7.18 (m, 30 H, 6 Ph), 5.74 (d, 1 H, H-4e), 5.54 (dt, 1 H, $J_{3eq,4}$ = 5.2 Hz, $J_{4,5}$ = 10.5 Hz, H-4g), 5.48 (t, 1 H $J_{1,2}$ = 7.7 Hz, H-2e), 5.43 (m, 3 H, H-4b, NH-f, and PhCH), 5.34 (m, 6 H, H-7b, 7g, and 4 NH), 5.17 (m, 3 H, H-4f, 8b, and 8g), 5.14 (d, 1 H, $J_{1,2}$ = 8.0 Hz, H-1d), 5.03 (dd, 1 H, H-7a), 4.97 (br d, 1 H, $J_{1,2}$ = 7.7 Hz, H-1e), 4.95 (dd, 1 H, $J_{7,8}$ = 9.6 Hz, H-7f), 4.88 (m, 2 H, H-4a and 3d), 4.80 and 4.52 (2 d, 2 H, J_{gem} = 11.4 Hz, PhCH₂), 4.65 (dt, 1 H, $J_{7,8}$ = 9.6 Hz, H-8f), 4.57 (dd, 1 H, H-6e), 4.52 and 4.46 (2 d, 2 H, J_{gem} = 11.6 Hz, PhCH₂), 4.42 (m, 3 H, H-9'a, 6'e, and 3e), 4.38–4.19 (m, 8 H, H-8a, 9'f, 1c, 4d, 9'g, 9'b, 5b, and 5e), 4.11 (2 q, 2 H, H-5g, and 5f), 4.08–3.80 (m, 18 H, OCH₂CH₂SiMe₃, H-5a, 9a, 9b, 9g, 6a, 6'd, 4c, 9f, 3c, 6g, 6f, 6b, 6'c, 5d, and COOMe-a), 3.71 (d, 1 H, H-6d), 3.55 (m, 2 H, H-6c and OCH₂CH₂SiMe₃), 3.44 (t, 1 H, H-5c), 3.34 (near q, 1 H, $J_{1,2}$ = 8.0 Hz, H-2d), 3.33 (near t, 1 H, H-2c), 3.17 (s, 3 H, COOMe-f), 2.54 (dt, 2 H, H-3eq-a and 3eq-b), 2.44 (dd, 1 H, $J_{3eq,4}$ = 5.2 Hz, J_{gem} = 13.5 Hz, H-3eq-g), 2.13–1.88 (m, 55 H, 17 Ac, H-3ax-a, 3ax-b, 3ax-f, and 3eq-f), 1.59 (t, 1 H, J_{gem} = 13.5 Hz, H-3ax-g), 1.01 (near t, 2 H, OCH₂CH₂SiMe₃), 0.00 (s, 9 H, OCH₂CH₂SiMe₃); ¹³C NMR (100 MHz, CDCl₃) δ 170.8, 170.7, 170.6, 170.5, 170.5, 170.5, 170.4, 170.4, 170.3, 170.3, 169.9, 169.8, 169.8, 169.8, 169.7, 169.6, 167.7, 167.3, 166.0, 164.7, 164.4, 164.2, 138.7, 138.5, 138.4, 133.4, 133.2, 133.0, 130.2, 129.9, 129.6, 129.5, 128.7, 128.6, 128.4, 128.4, 128.1, 128.0, 128.0, 127.9, 127.6, 127.5, 127.3, 126.4, 126.2, 103.4, 102.3, 100.7, 99.4, 99.1, 98.1, 97.1, 96.7, 77.8, 77.2, 75.9, 75.5, 74.6, 73.3, 72.9, 72.6, 72.5, 72.4, 72.1, 71.8, 71.5, 71.0, 70.6, 70.1, 69.6, 69.5, 69.3, 68.9, 68.7, 68.6, 68.5, 67.8, 67.7, 67.4, 66.7, 66.6, 66.0, 62.7, 62.0, 61.8, 54.3, 52.3, 49.1, 49.0, 48.9, 38.2, 35.9, 35.7, 30.0, 29.6, 23.1, 22.8, 20.9, 20.7, 20.7, 20.6, 20.6, 20.6, 20.5, 20.5, 18.5, 14.1, -1.5; HRMS (ESI) m/z found [M + 2Na]²⁺ 1465.9833, C₁₃₇H₁₆₇N₅O₆₁Si calcd for [M + 2Na]²⁺ 1465.9838.

[Methyl 5-Acetamido-8-*O*-(5-acetamido-4,7,8,9-tetra-*O*-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylono-1',9-lactone)-4,7-di-*O*-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylate]-(2 \rightarrow 3)-2,4,6-tri-*O*-benzoyl- β -D-galactopyranosyl-(1 \rightarrow 3)-2-acetamido-4,6-di-*O*-benzoyl-2-deoxy- β -D-galactopyranosyl-(1 \rightarrow 4)-{[methyl 5-acetamido-8-*O*-(5-acetamido-4,7,8,9-tetra-*O*-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylono-1',9-lactone)-4,7-di-*O*-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylate]-(2 \rightarrow 3)-2,6-di-*O*-benzoyl- α -D-galactopyranosyl Trichloroacetimidate (20). To a solution of **18** (795 mg, 261 μ mol) in CH₂Cl₂ (3.9 mL) was added trifluoroacetic acid (1.3 mL) at 0 °C. After stirring for 3 h at room temperature as the reaction was monitored by TLC (PhMe/MeOH = 5:1), the reaction mixture was coevaporated with toluene. The resulting residue was purified by flash column chromatography using CHCl₃/MeOH (17.5:1 \rightarrow 15:1 \rightarrow 10:1) as the eluent to give **19** (765 mg, 99%): HRMS (ESI) m/z found [M + 2Na]⁺ 1489.9383, C₁₃₉H₁₅₅N₅O₆₅ calcd for [M + 2Na]⁺ 1489.9380.

To a solution of **19** (200 mg, 68.1 μ mol) in CH₂Cl₂ (0.68 mL) were added trichloroacetonitrile (136 μ L, 1.36 mmol) and DBU (12 μ L, 81.7 μ mol) at 0 °C. After stirring for 2 h at room temperature as the reaction was monitored by TLC (PhMe/MeOH = 5:1), the reaction mixture was evaporated. The resulting residue was purified by flash column chromatography using CHCl₃/MeOH

(20:1 to 13:1) as the eluent to give **20** (199 mg, 95%): [α]_D = +4.8° (c 0.9, CHCl₃); ¹H NMR (500 MHz, CDCl₃) δ 8.46 (s, 1 H, C=NH), 8.12–7.25 (m, 35 H, 7 Ph), 6.55 (d, 1 H, $J_{1,2}$ = 3.4 Hz, H-1c), 5.95 (br d, 1 H, H-4d), 5.89 (br d, 2 H, 2 NH), 5.70 (d, 1 H, H-4e), 5.63 (dd, 1 H, $J_{1,2}$ = 3.4 Hz, $J_{2,3}$ = 10.4 Hz, H-2c), 5.58 (d, 1 H, NH-b), 5.55 (dt, 1 H, H-4g), 5.46 (2 d, 2 H, 2 NH), 5.36–5.31 (m, 4 H, H-4b, 2e, 7g, and 7b), 5.29 (d, 1 H, $J_{1,2}$ = 8.5 Hz, H-1d), 5.24 (m, 1 H, H-8b), 5.16 (m, 1 H, H-8g), 5.10 (near q, 1 H, H-4f), 5.04 (br d, 1 H, H-1e), 5.20 (m, 2 H, H-4a and 7a), 4.93 (br dd, 1 H, H-3d), 4.86 (dd, 1 H, H-7f), 4.80 (dd, 1 H, $J_{2,3}$ = 10.4 Hz, H-3c), 4.63–4.58 (m, 3 H, H-8f, 6'e, and 6'd), 4.45–4.33 (m, 5 H, H-5d, 6'c, 3e, 6c, and 6d), 4.30–4.16 (m, 10 H, H-8a, 9g, 9b, 9f, 4c, 6b, 5b, 9'a, 9a, and 5e), 4.11 (q, 1 H, H-5g), 4.06–3.98 (m, 6 H, H-5f, 9b, 9g, 6b, 5c, and 5a), 3.91 (2 dd, 2 H, H-6a and 6g), 3.86 (dd, 1 H, H-6f), 3.82 (s, 3 H, COOMe-a), 3.75 (m, 2 H, H-2d and 9'f), 3.24 (s, 3 H, COOMe-f), 2.57–2.49 (m, 3 H, H-3eq-a, 3eq-b, and 3eq-g), 2.10–1.81 (m, 53 H, 17 Ac, H-3eq-f, and 3ax-f), 1.63 (t, 1 H, H-3ax-g); ¹³C NMR (150 MHz, CDCl₃) δ 174.5, 171.2, 170.6, 170.6, 170.4, 170.2, 170.2, 169.8, 169.7, 169.6, 167.7, 167.1, 166.5, 166.0, 165.9, 165.5, 165.3, 164.8, 164.2, 160.0, 133.4, 133.2, 133.1, 133.0, 132.8, 132.8, 130.0, 129.9, 129.8, 129.7, 129.6, 129.5, 129.4, 129.3, 129.0, 128.4, 128.3, 128.2, 101.3, 99.8, 99.0, 98.9, 96.8, 95.4, 95.1, 94.1, 90.8, 76.2, 75.0, 73.3, 72.5, 71.9, 71.5, 71.3, 71.1, 71.0, 70.6, 70.1, 69.8, 69.6, 69.3, 69.2, 69.1, 68.8, 68.6, 68.1, 67.5, 67.1, 66.8, 66.7, 66.4, 64.2, 62.3, 62.1, 62.1, 61.9, 54.3, 53.0, 52.3, 49.0, 48.9, 48.7, 38.8, 38.2, 36.9, 36.1, 35.4, 29.5, 22.9, 22.8, 22.6, 22.5, 20.7, 20.6, 20.6, 20.5, 20.4, 20.3, 20.2, 13.9; HRMS (ESI) m/z found [M + 2Na]²⁺ 1561.3934, C₁₄₁H₁₅₅Cl₃N₆O₆₅ calcd for [M + 2Na]²⁺ 1561.3934.

2-*O*-Benzoyl-4-*O*-chloroacetyl-3,6-di-*O*-*p*-methoxybenzyl-D-glucopyranosyl Trichloroacetimidate (33). To a solution of **31** (500 mg, 0.779 mmol) in degassed THF (7.8 mL) were added polymethylhydrosiloxane (PMHS) (140 μ L, 2.33 mmol), tetrakis(triphenylphosphine)palladium (179 mg, 0.155 mmol), and ZnCl₂ (106 mg, 0.779 mmol) at room temperature. After stirring for 32 h at room temperature, the completion of the reaction was confirmed by TLC (EtOAc/PhMe = 1:4). Then, H₂O was added to the mixture and filtered through a Celite pad, and the pad was washed with EtOAc. The combined filtrate and washings were washed with saturated NaHCO₃ (aq), and brine. After drying over Na₂SO₄ and being concentrated, the resulting residue was purified by flash column chromatography using EtOAc/PhMe (1:5) as the eluent to give the hemiacetal **32** (400 mg, 85%, α/β = 7/1): ¹H NMR (600 MHz, CDCl₃) δ 8.05–7.42 (m, 5 H, Ph), 7.21–6.75 (4 d, 8 H, Ar), 5.55 (br t, 1 H, $J_{1,2}$ = $J_{1,OH}$ = 3.4 Hz, H-1 α), 5.19 (t, 1 H, $J_{3,4}$ = 8.9 Hz, H-4 β), 5.15 (t, 1 H, $J_{1,2}$ = $J_{2,3}$ = 8.9 Hz, H-2 β), 5.09 (t, 1 H, $J_{3,4}$ = $J_{4,5}$ = 9.6 Hz, H-4 α), 5.08 (dd, 1 H, $J_{1,2}$ = 3.4 Hz, $J_{2,3}$ = 9.6 Hz, H-2 α), 4.73 (t, 1 H, $J_{1,2}$ = $J_{1,OH}$ = 8.9 Hz, H-1 β), 4.67 and 4.52 (2 d, 2 H, J_{gem} = 11.6 Hz, ArCH₂O), 4.44 and 4.39 (2 d, 2 H, J_{gem} = 11.7 Hz, ArCH₂O), 4.18 (m, 2 H, $J_{2,3}$ = $J_{3,4}$ = 9.6 Hz, H-3 α and 5 α), 3.92 (d, 1 H, $J_{1,OH}$ = 8.9 Hz, OH- β), 3.87 (t, 1 H, $J_{2,3}$ = $J_{3,4}$ = 8.9 Hz, H-3 β), 3.78 and 3.74 (2 s, 6 H, 2 MeO), 3.69 and 3.61 (2 d, 2 H, J_{gem} = 14.4 Hz, ClCH₂), 3.58 (d, 1 H, $J_{1,OH}$ = 3.4 Hz, OH- α), 3.47 (dd, 1 H, J_{gem} = 10.3 Hz, $J_{5,6}$ = 5.5 Hz, H-6 α), 3.43 (dd, 1 H, J_{gem} = 10.3 Hz, $J_{5,6}$ = 3.4 Hz, H-6 α); HRMS (ESI) m/z found [M + Na]⁺ 623.1658, C₃₁H₃₃ClO₁₀ calcd for [M + Na]⁺ 623.1659.

To a solution of **32** (352 mg, 0.585 mmol) in CH₂Cl₂ (5.8 mL) were added trichloroacetonitrile (1.2 mL, 11.7 mmol) and K₂CO₃ (243 mg, 1.76 mmol) at room temperature. After stirring for 3 h at room temperature as the reaction was monitored by TLC (EtOAc/hexane = 1:2), the reaction mixture was filtered through cotton, washed with CH₂Cl₂, and then evaporated. The resulting residue was purified by flash column chromatography using EtOAc/hexane (1:5 to 1:4) as the eluent to give **33**: [α -isomer, 176 mg (40%) and β -isomer, 230 mg (53%)]. **33 α** : [α]_D = +77.1° (c 0.7, CHCl₃); ¹H NMR (500 MHz, CDCl₃) δ 8.54 (s, 1 H, C=NH), 7.98–7.41 (m, 5 H, Ph), 7.23–6.75 (4 d, 8 H, Ar), 6.63 (d, 1 H, $J_{1,2}$ = 3.5 Hz, H-1), 5.38 (dd, 1 H, $J_{1,2}$ = 3.5 Hz, $J_{2,3}$ = 9.5 Hz, H-2), 5.35 (t, 1

H, $J_{3,4} = J_{4,5} = 10.0$ Hz, H-4), 4.65 and 4.54 (2 d, 2 H, $J_{\text{gem}} = 11.5$ Hz, ArCH₂O), 4.46 and 4.37 (2 d, 2 H, $J_{\text{gem}} = 11.5$ Hz, ArCH₂O), 4.23 (near t, 1 H, $J_{2,3} = 9.5$ Hz, $J_{3,4} = 10.0$ Hz, H-3), 4.11 (m, 1 H, $J_{4,5} = 10.0$ Hz, $J_{5,6} = J_{5,6'} = 4.0$ Hz, H-5), 3.80 and 3.75 (2 s, 6 H, 2 MeO), 3.71 and 3.66 (2 d, 2 H, $J_{\text{gem}} = 15.0$ Hz, ClCH₂), 3.57 (dd, 1 H, $J_{\text{gem}} = 11.0$ Hz, $J_{5,6} = 4.0$ Hz, H-6), 3.52 (dd, 1 H, $J_{\text{gem}} = 11.0$ Hz, $J_{5,6'} = 4.0$ Hz, H-6'); ¹³C NMR (125 MHz, CDCl₃) δ 165.8, 165.1, 160.3, 159.3, 133.4, 129.7, 129.7, 129.7, 129.4, 129.0, 128.4, 113.7, 93.5, 75.8, 74.3, 73.2, 72.3, 71.7, 71.1, 68.2, 55.2, 55.2, 40.5; HRMS (ESI) m/z found [M + Na]⁺ 766.0750, C₃₃H₃₃Cl₄NO₁₀ calcd for [M + Na]⁺ 766.0756. **33 β** : [α]_D = +43.5° (c 1.0, CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 8.63 (s, 1 H, C=NH), 7.98–7.40 (m, 5 H, Ph), 7.24–6.69 (4 d, 8 H, Ar), 5.96 (d, 1 H, $J_{1,2} = 7.8$ Hz, H-1), 5.58 (near t, 1 H, $J_{1,2} = 7.8$ Hz, $J_{2,3} = 8.6$ Hz, H-2), 5.33 (t, 1 H, $J_{3,4} = J_{4,5} = 9.1$ Hz, H-4), 4.58 and 4.50 (2 d, 2 H, $J_{\text{gem}} = 11.4$ Hz, ArCH₂O), 4.45 and 4.40 (2 d, 2 H, $J_{\text{gem}} = 11.2$ Hz, ArCH₂O), 3.93 (near t, 1 H, $J_{2,3} = 8.6$ Hz, $J_{3,4} = 9.1$ Hz, H-3), 3.85 (m, 1 H, $J_{4,5} = 9.1$ Hz, H-5), 3.79 and 3.71 (2 s, 6 H, 2 MeO), 3.74–3.59 (m, 4 H, ClCH₂, H-6, and 6'); ¹³C NMR (100 MHz, CDCl₃) δ 165.9, 164.5, 161.2, 159.3, 159.2, 133.3, 129.7, 129.6, 129.5, 129.4, 129.3, 128.4, 113.7, 113.6, 95.8, 90.2, 78.5, 73.7, 73.4, 73.2, 72.2, 71.8, 68.8, 55.2, 55.1, 40.5; HRMS (ESI) m/z found [M + Na]⁺ 766.0750, C₃₃H₃₃Cl₄NO₁₀ calcd for [M + Na]⁺ 766.0756.

2-O-Benzoyl-4-O-chloroacetyl-3,6-di-O-p-methoxybenzyl- β -D-glucopyranosyl-(1 \rightarrow 1)-(2S,3R,4E)-3-O-benzoyl-2-octadecanamido-4-octadecene-1,3-diol (35). To a mixture of **33 α** (176 mg, 0.236 mmol) and **34** (105 mg, 0.157 mmol) in CH₂Cl₂ (3.9 mL) was added 4 Å molecular sieves (AW-300) (300 mg) at room temperature. After stirring for 2 h, TMSOTf (2.1 μ L, 11.8 μ mol) was added to the mixture. After stirring for 3 h at room temperature as the reaction was monitored by TLC (EtOAc/PhMe = 1:5), the reaction was quenched by saturated NaHCO₃ (aq) and filtered through a Celite pad, and the pad was washed with CHCl₃. The combined filtrate and washings were washed with brine, dried over Na₂SO₄, and concentrated. The resulting residue was purified by flash column chromatography using EtOAc/PhMe (1:10) as the eluent to give **35** (95 mg, 48%): [α]_D = +10.7° (c 1.6, CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 8.03–7.39 (m, 10 H, 2 Ph), 7.16–6.68 (4 d, 8 H, Ar), 5.83 (m, 1 H, $J_{4,5} = 15.1$ Hz, $J_{5,6} = 7.3$ Hz, $J_{5,6'} = 6.8$ Hz, H-5i), 5.71 (d, 1 H, $J_{2,\text{NH}} = 9.1$ Hz, NH), 5.51 (t, 1 H, $J_{2,3} = J_{3,4} = 6.8$ Hz, H-3i), 5.44 (dd, 1 H, $J_{3,4} = 6.8$ Hz, $J_{4,5} = 15.1$ Hz, H-4i), 5.25 (near t, 1 H, $J_{1,2} = 8.2$ Hz, $J_{2,3} = 9.6$ Hz, H-2h), 5.15 (t, 1 H, $J_{3,4} = J_{4,5} = 9.6$ Hz, H-4h), 4.53 and 4.46 (2 d, 2 H, $J_{\text{gem}} = 11.4$ Hz, ArCH₂O), 4.50 (d, 1 H, $J_{1,2} = 8.2$ Hz, H-1h), 4.40 (m, 1 H, $J_{1,2} = 2.7$ Hz, $J_{2,3} = 6.8$ Hz, H-2i), 4.28 and 4.22 (2 d, 2 H, $J_{\text{gem}} = 11.0$ Hz, ArCH₂O), 4.10 (dd, 1 H, $J_{\text{gem}} = 9.6$ Hz, $J_{1,2} = 2.7$ Hz, H-1i), 3.82 (t, 1 H, $J_{2,3} = J_{3,4} = 9.6$ Hz, H-3h), 3.78 and 3.70 (2 s, 6 H, 2 MeO), 3.65 (m, 2 H, ClCH₂), 3.58 (m, 2 H, H-5h and H-1'i), 3.45 (dd, 1 H, $J_{\text{gem}} = 10.0$ Hz, $J_{5,6} = 5.0$ Hz, H-6h), 3.38 (dd, 1 H, $J_{\text{gem}} = 10.0$ Hz, $J_{5,6'} = 4.6$ Hz, H-6'h), 1.98 (br dd, 2 H, $J_{\text{gem}} = 14.1$ Hz, $J_{5,6} = 7.3$ Hz, $J_{5,6'} = 6.8$ Hz, H-6i and 6'i), 1.74 and 1.37 (2 m, 2 H, NHCOCH₂), 1.26 (m, 52 H, 26 CH₂), 0.87 (t, 6 H, 2 Me); ¹³C NMR (100 MHz, CDCl₃) δ 172.5, 165.9, 165.0, 164.9, 159.2, 159.1, 137.2, 133.3, 132.8, 130.2, 129.6, 129.5, 129.5, 129.3, 128.4, 128.2, 124.7, 113.7, 113.6, 100.6, 78.8, 74.3, 73.6, 73.5, 73.2, 72.9, 72.6, 69.3, 67.1, 55.1, 55.0, 50.3, 40.5, 36.3, 32.2, 31.8, 29.6, 29.4, 29.3, 29.2, 29.1, 28.8, 25.4, 22.6, 14.0; HRMS (ESI) m/z found [M + Na]⁺ 1274.7255, C₇₄H₁₀₆ClNO₁₃ calcd for [M + Na]⁺ 1274.7250.

2-O-Benzoyl-3,6-di-O-p-methoxybenzyl- β -D-glucopyranosyl-(1 \rightarrow 1)-(2S,3R,4E)-3-O-benzoyl-2-octadecanamido-4-octadecene-1,3-diol (36). To a solution of **35** (95 mg, 0.0758 mmol) in EtOH/CH₂Cl₂ (3.6 mL/1.2 mL) was added DABCO (127 mg, 1.13 mmol) at room temperature. After stirring for 5 h at 55 °C as the reaction was monitored by TLC (EtOAc/PhMe = 1:3), the mixture was diluted with EtOAc. The organic layer was washed with 2 M HCl, saturated NaHCO₃ (aq), and brine. After drying over Na₂SO₄ and being concentrated, the resulting residue was purified by flash

column chromatography using EtOAc/PhMe (1:5) as the eluent to give **36** (85 mg, 96%): [α]_D = +5.4° (c 1.7, CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 8.01–7.37 (m, 10 H, 2 Ph), 7.18–6.66 (4 d, 8 H, Ar), 5.81 (m, 1 H, $J_{4,5} = 15.1$ Hz, $J_{5,6} = 7.8$ Hz, $J_{5,6'} = 6.8$ Hz, H-5i), 5.70 (d, 1 H, $J_{2,\text{NH}} = 9.1$ Hz, NH), 5.48 (t, 1 H, $J_{2,3} = J_{3,4} = 7.3$ Hz, H-3i), 5.42 (dd, 1 H, $J_{3,4} = 7.3$ Hz, $J_{4,5} = 15.1$ Hz, H-4i), 5.16 (near t, 1 H, $J_{1,2} = 7.8$ Hz, $J_{2,3} = 9.2$ Hz, H-2h), 4.64 (t, 2 H, $J_{\text{gem}} = 11.4$ Hz, 2 ArCH₂O), 4.45 (d, 1 H, $J_{1,2} = 7.8$ Hz, H-1h), 4.40 and 4.33 (2 d, 2 H, $J_{\text{gem}} = 11.4$ Hz, 2 ArCH₂O), 4.38 (m, 1 H, H-2i), 4.08 (dd, 1 H, $J_{\text{gem}} = 9.6$ Hz, $J_{1,2} = 2.3$ Hz, H-1i), 3.78 and 3.74 (2 s, 6 H, 2 MeO), 3.74 (dt, 1 H, $J_{3,4} = J_{4,5} = 9.2$ Hz, $J_{4,\text{OH}} = 1.8$ Hz, H-4h), 3.63 (t, 1 H, $J_{2,3} = J_{3,4} = 9.2$ Hz, H-3h), 3.57 (m, 3 H, H-6h, 6'h, and H-1'i), 3.45 (m, 1 H, $J_{4,5} = 9.2$ Hz, H-5h), 2.93 (d, 1 H, $J_{4,\text{OH}} = 1.8$ Hz, OH), 1.97 (near dd, 2 H, $J_{\text{gem}} = 13.2$ Hz, $J_{5,6} = 7.8$ Hz, $J_{5,6'} = 6.8$ Hz, H-6i and 6'i), 1.72 (t, 1 H, NHCOCH₂), 1.37 (m, 1 H, NHCOCH₂), 1.26 (m, 52 H, 26 CH₂), 0.87 (t, 6 H, 2 Me); ¹³C NMR (100 MHz, CDCl₃) δ 172.5, 165.2, 165.0, 159.2, 159.1, 137.1, 133.2, 132.7, 130.3, 130.0, 129.6, 129.5, 129.3, 128.4, 128.2, 124.7, 113.7, 113.6, 100.8, 81.1, 74.3, 73.9, 73.8, 73.5, 73.3, 72.6, 70.1, 67.0, 55.1, 55.0, 50.3, 36.3, 32.2, 31.8, 29.6, 29.6, 29.4, 29.3, 29.2, 29.1, 28.8, 25.4, 22.6, 14.0; HRMS (ESI) m/z found [M + Na]⁺ 1198.7533, C₇₂H₁₀₅NO₁₂ calcd for [M + Na]⁺ 1198.7534.

[Methyl 5-acetamido-8-O-(5-acetamido-4,7,8,9-tetra-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylono-1',9-lactone)-4,7-di-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylono-1',9-lactone]-2-(2-3)-2,4,6-tri-O-benzoyl- β -D-galactopyranosyl-(1 \rightarrow 3)-2-acetamido-4,6-di-O-benzoyl-2-deoxy- β -D-galactopyranosyl-(1 \rightarrow 4)-{[methyl 5-acetamido-8-O-(5-acetamido-4,7,8,9-tetra-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylono-1',9-lactone)-4,7-di-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylono-1',9-lactone]-2-(2-3)}-2,6-di-O-benzoyl- β -D-galactopyranosyl-(1 \rightarrow 4)-2-O-benzoyl-3,6-di-O-p-methoxybenzyl- β -D-glucopyranosyl-(1 \rightarrow 1)-(2S,3R,4E)-3-O-benzoyl-2-octadecanamido-4-octadecene-1,3-diol (40). To a mixture of **20** (146 mg, 0.0474 mmol) and **36** (85 mg, 0.0722 mmol) in CH₂Cl₂ (1.2 mL) was added 4 Å molecular sieves (AW-300) (200 mg) at room temperature. After stirring for 2 h and then cooling to 0 °C, TMSOTf (1.0 μ L, 5.52 μ mol) was added to the mixture. After stirring for 1 h at 0 °C as the reaction was monitored by TLC (PhMe/MeOH = 4:1), the reaction was quenched by saturated NaHCO₃ (aq) and filtered through a Celite pad, and the pad was washed with CHCl₃. The combined filtrate and washings were washed with brine, dried over Na₂SO₄, and concentrated. The resulting residue was purified by flash column chromatography using PhMe/MeOH (8:1 to 7:1) as the eluent to give **40** (177 mg, 91%): [α]_D = +3.2° (c 0.5, CHCl₃); ¹H NMR (600 MHz, CDCl₃) δ 8.08–7.28 (m, 45 H, 9 Ph), 7.07–6.56 (4 d, 8 H, Ar), 5.97 (near s, 1 H, H-4d), 5.76 (m, 1 H, $J_{4,5} = 14.4$ Hz, $J_{5,6} = 7.5$ Hz, $J_{5,6'} = 6.9$ Hz, H-5i), 5.69 (d, 1 H, $J_{3,4} = 4.1$ Hz, H-4e), 5.66 (d, 1 H, $J_{2,\text{NH}} = 8.9$ Hz, NH-i), 5.55 (dt, 2 H, $J_{3\text{eq},4} = 4.0$ Hz, $J_{4,5} = 10.9$ Hz, H-4g and NH-d), 5.49 (br d, 1 H, NH-b), 5.45 (t, 1 H, H-3i), 5.39 (dd, 2 H, H-4i and NH-f), 5.35–5.26 (m, 5 H, H-7g, 7b, 4b, 2c, and 2e), 5.22 (d, 1 H, $J_{1,2} = 8.9$ Hz, H-1d), 5.15 (m, 4 H, H-8b, 8g, 2h, and NH-a), 5.08 (m, 1 H, H-4f), 4.98 (br dd, 1 H, H-7a), 4.95 (m, 1 H, H-4a), 4.84 (br dd, 1 H, H-7f), 4.82 (d, 1 H, $J_{1,2} = 9.6$ Hz, H-1c), 4.80 (m, 2 H, H-3d and ArCH₂O), 4.63 (m, 3 H, H-6c, 8f, and ArCH₂O), 4.55 (d, 1 H, $J_{\text{gem}} = 10.9$ Hz, ArCH₂O), 4.40 (br d, 2 H, $J_{1,2} = 7.5$ Hz, H-1e and 3e), 4.34 (d, 1 H, $J_{1,2} = 8.2$ Hz, H-1h), 4.32 (m, 1 H, H-2i), 4.28–3.96 (m, 24 H, H-9'b, 9'g, 9'f, 9'a, 4c, 5b, 3c, 6'c, 5g, 6b, 8a, 9a, 9b, 9g, 5a, 4h, 5e, 6e, 6'e, 5d, 6'd, 6d, 1i, and ArCH₂O), 3.89 (dd, 1 H, H-6g), 3.84 (dd, 2 H, H-6a and 6f), 3.80 (t, 1 H, $J_{2,3} = J_{3,4} = 8.9$ Hz, H-3h), 3.77 (s, 3 H, COOMe-a), 3.71 (s, 3 H, MeO), 3.70 (m, 2 H, H-2d and 9f), 3.62 (near t, 1 H, H-5c), 3.43 (m, 6 H, H-6'h, 6h, 1'i, and MeO), 3.33 (m, 1 H, H-5h), 3.23 (s, 3 H, COOMe-f), 2.57 (dd, 1 H, $J_{\text{gem}} = 13.0$ Hz, $J_{3\text{eq},4} = 4.1$ Hz, H-3eq-b), 2.50 (dd, 1 H, $J_{\text{gem}} = 12.2$ Hz, $J_{3\text{eq},4} = 4.0$ Hz, H-3eq-g), 2.33 (br dd, 1 H, H-3eq-a), 2.10–1.76 (m, 57 H, 17 Ac, H-3eq-f, 3ax-f, 3ax-a, 3ax-b, 6i, and

6'i), 1.67 (t, 1 H, NHCOCH_2), 1.63 (t, 1 H, $J_{\text{gem}} = J_{3\text{ax},4} = 12.2$ Hz, H-3ax-g), 1.34 (m, 1 H, NHCOCH_2), 1.26 (m, 52 H, 26 CH_2), 0.87 (2 t, 6 H, 2 Me); ^{13}C NMR (100 MHz, CDCl_3) δ 172.4, 170.8, 170.6, 170.5, 170.4, 170.4, 170.3, 170.3, 170.1, 170.1, 169.7, 169.6, 169.5, 167.3, 167.1, 166.3, 165.9, 165.8, 165.6, 165.4, 165.2, 165.0, 164.9, 164.7, 164.1, 164.0, 158.9, 158.7, 136.9, 133.1, 133.0, 132.9, 132.6, 130.1, 130.1, 130.0, 129.8, 129.6, 129.5, 129.4, 129.4, 129.3, 128.9, 128.8, 128.4, 128.3, 128.1, 124.6, 113.4, 113.3, 101.3, 100.8, 100.3, 99.6, 99.4, 98.9, 96.8, 95.7, 80.1, 77.2, 76.4, 75.1, 74.3, 73.9, 73.8, 73.4, 73.1, 72.8, 72.5, 72.1, 71.8, 71.5, 71.3, 71.2, 71.1, 70.8, 70.7, 70.3, 69.8, 69.4, 69.3, 69.2, 68.7, 68.5, 68.2, 67.8, 67.1, 66.8, 66.6, 63.3, 62.0, 61.7, 55.0, 54.6, 53.0, 52.2, 50.2, 48.9, 48.7, 38.8, 38.2, 36.9, 36.2, 35.4, 32.5, 32.1, 31.7, 29.8, 29.5, 29.4, 29.3, 29.2, 29.1, 29.0, 28.8, 25.3, 22.9, 22.8, 22.6, 22.5, 20.6, 20.6, 20.5, 20.4, 20.2, 20.1, 14.0; HRMS (ESI) m/z found $[\text{M} + 2\text{Na}]^{2+}$ 2068.8101, $\text{C}_{211}\text{H}_{258}\text{N}_6\text{O}_{76}$ calcd for $[\text{M} + 2\text{Na}]^{2+}$ 2068.8146.

[Methyl 5-acetamido-8-O-(5-acetamido-4,7,8,9-tetra-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylono-1',9-lactone)-4,7-di-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylate]-(2-3)-2,4,6-tri-O-benzoyl- β -D-galactopyranosyl-(1-3)-2-acetamido-4,6-di-O-benzoyl-2-deoxy- β -D-galactopyranosyl-(1-4)-{[methyl 5-acetamido-8-O-(5-acetamido-4,7,8,9-tetra-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylono-1',9-lactone)-4,7-di-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylate]-(2-3)-2,6-di-O-benzoyl- β -D-galactopyranosyl-(1-4)-2-O-benzoyl-3,6-di-O-benzyl- β -D-glucopyranosyl-(1-1)-(2S,3R,4E)-3-O-benzyl-2-octadecanamide-4-octadecene-1,3-diol (41)}. To a mixture of 20 (148 mg, 0.0480 mmol) and 37 (79 mg, 0.0720 mmol) in CH_2Cl_2 (1.0 mL) was added 4 Å molecular sieves (AW-300) (180 mg) at room temperature. After stirring for 2 h and then cooling to 0 °C, TMSOTf (1.0 μL , 5.52 μmol) was added to the mixture. After stirring for 2 h at 0 °C as the reaction was monitored by TLC (PhMe/MeOH = 4:1), additional TMSOTf (1.0 μL , 5.52 μmol) was added to the mixture. After the stirring was continued for 24 h, the reaction was quenched by saturated NaHCO_3 (aq) and filtered through a Celite pad, and the pad was washed with CHCl_3 . The combined filtrate and washings were washed with brine, dried over Na_2SO_4 and concentrated. The resulting residue was purified by flash column chromatography (PhMe/MeOH = 9:1 to 8:1) to give 41 (140 mg, 73%): $[\alpha]_{\text{D}} = -2.5^\circ$ (c 1.0, CHCl_3); ^1H NMR (600 MHz, CDCl_3) δ 8.01–6.89 (m, 55 H, 11 Ph), 5.97 (near s, 1 H, H-4d), 5.69 (d, 1 H, $J_{3,4} = 2.7$ Hz, H-4e), 5.54 (m, 4 H, H-4g, 5i, and 2 NH), 5.36 (m, 5 H, H-4b, 7b, and 3 NH), 5.32 (dd, 1 H, $J_{6,7} = 2.0$ Hz, $J_{7,8} = 8.2$ Hz, H-7g), 5.19 (m, 7 H, H-8b, 2e, 2c, 1d, 2h, 8g, and 4i), 5.08 (m, 1 H, H-4f), 5.01 (near d, 1 H, $J_{7,8} = 8.9$ Hz, H-7a), 4.98 (m, 1 H, H-4a), 4.89 (d, 1 H, $J_{\text{gem}} = 10.9$ Hz, PhCH_2), 4.83 (near dd, 1 H, $J_{7,8} = 9.6$ Hz, H-7f), 4.62 (m, 5 H, H-3d, 8f, 6c, 6'c, and PhCH_2), 4.60 (d, 1 H, $J_{1,2} = 7.5$ Hz, H-1c), 4.49 and 4.28 (2 d, 2 H, $J_{\text{gem}} = 11.6$ Hz, PhCH_2), 4.49 and 4.17 (2 d, 2 H, $J_{\text{gem}} = 11.6$ Hz, PhCH_2), 4.44 (m, 2 H, H-1e and 3e), 4.35 (d, 1 H, $J_{1,2} = 7.5$ Hz, H-1h), 4.31–4.20 (m, 8 H, H-9'g, 9'b, 9'f, 9f, 5b, 9'a, 1'i, and 8a), 4.13–3.97 (m, 10 H, H-6b, 5a, 5g, 4h, 4c, 3c, 5f, 2i, 9b, and 9g), 3.89 (dd, 1 H, $J_{5,6} = 10.3$ Hz, $J_{6,7} = 2.0$ Hz, H-6g), 3.84 (m, 2 H, H-6a and 6f), 3.78 (m, 5 H, H-3h, 2d, and COOMe-a), 3.71 (m, 2 H, H-3i and 9a), 3.61 (br t, 1 H, H-5c), 3.53 (br dd, 1 H, $J_{\text{gem}} = 10.9$ Hz, H-6'h), 3.46 (br d, 1 H, $J_{\text{gem}} = 10.9$ Hz, H-6h), 3.35 (br dd, 1 H, H-1i), 3.26 (m, 1 H, H-5h), 3.23 (s, 3 H, COOMe-f), 2.57 (br dd, 1 H, H-3eq-b), 2.48 (dd, 1 H, H-3eq-g), 2.22 (m, 1 H, H-3eq-a), 2.11–1.82 (13 s and m, 57 H, 17 Ac, H-3eq-f, 3ax-f, 3ax-a, 3ax-b, 6i, and 6'i), 1.61 (m, 3 H, H-3ax-g, NHCOCH_2), 1.26 (m, 52 H, 26 CH_2), 0.87 (t, 6 H, 2 Me); ^{13}C NMR (125 MHz, CDCl_3) δ 172.3, 170.8, 170.7, 170.6, 170.6, 170.5, 170.4, 170.4, 170.2, 169.8, 169.8, 169.7, 167.5, 167.3, 166.2, 166.0, 165.8, 165.4, 165.2, 164.6, 164.2, 164.1, 138.5, 138.2, 138.0, 137.0, 133.4, 133.3, 133.1, 133.1, 133.0, 132.8, 130.3, 130.0, 129.9, 129.8, 129.8, 129.7, 129.6, 129.5, 129.5, 129.3, 128.6, 128.5, 128.5, 128.4, 128.3, 128.1, 128.0, 128.0, 127.9, 127.6, 127.5, 127.4, 127.3, 127.2, 101.3, 100.2, 100.2, 99.9, 99.6, 99.0, 97.1, 96.1, 80.3, 79.0,

75.6, 74.8, 74.2, 74.0, 73.6, 73.3, 73.1, 72.5, 72.1, 72.0, 71.6, 71.4, 71.0, 70.9, 70.4, 70.1, 69.9, 69.5, 69.4, 69.4, 69.0, 68.7, 68.6, 68.5, 67.9, 67.7, 67.2, 67.0, 66.7, 63.7, 62.1, 62.0, 61.8, 54.0, 53.2, 52.4, 51.2, 49.1, 48.9, 48.8, 38.8, 38.3, 36.4, 35.4, 32.2, 31.9, 29.7, 29.6, 29.5, 29.3, 29.2, 29.2, 25.4, 23.1, 23.1, 23.0, 22.6, 20.8, 20.7, 20.6, 20.3, 14.1; HRMS (ESI) m/z found $[\text{M} + 2\text{Na}]^{2+}$ 2031.8146, $\text{C}_{209}\text{H}_{256}\text{N}_6\text{O}_{73}$ calcd for $[\text{M} + 2\text{Na}]^{2+}$ 2031.8149.

2-(Trimethylsilyl)ethyl [Methyl 5-acetamido-8-O-(5-acetamido-4,7,8,9-tetra-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylono-1',9-lactone)-4,7-di-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylate]-(2-3)-2,4,6-tri-O-benzoyl- β -D-galactopyranosyl-(1-3)-2-acetamido-4,6-di-O-benzoyl-2-deoxy- β -D-galactopyranosyl-(1-4)-{[methyl 5-acetamido-8-O-(5-acetamido-4,7,8,9-tetra-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylono-1',9-lactone)-4,7-di-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosylate]-(2-3)-2,6-di-O-benzoyl- β -D-galactopyranosyl-(1-4)-2-O-benzoyl-3,6-di-O-benzyl- β -D-glucopyranoside (42)}. To a mixture of 20 (189 mg, 0.0613 mmol) and 38 (52 mg, 0.0919 mmol) in CH_2Cl_2 (1.5 mL) was added 4 Å molecular sieves (AW-300) (250 mg) at room temperature. After stirring for 2 h and then cooling to 0 °C, TMSOTf (1.1 μL , 6.13 μmol) was added to the mixture. After stirring for 2 h at 0 °C as the reaction was monitored by TLC (PhMe/MeOH = 4:1), the reaction was quenched by saturated NaHCO_3 (aq) and filtered through a Celite pad, and the pad was washed with CHCl_3 . The combined filtrate and washings were washed with brine, dried over Na_2SO_4 , and concentrated. The residue was purified by flash column chromatography using PhMe/MeOH (6:1) as the eluent to give 42 (135 mg, 63%): $[\alpha]_{\text{D}} = -0.6^\circ$ (c 1.5, CHCl_3); ^1H NMR (500 MHz, CDCl_3) δ 8.13–6.89 (m, 50 H, 10 Ph), 5.95 (br s, 1 H, H-4d), 5.85 (br d, 1 H, NH-d), 5.70 (d, 1 H, $J_{3,4} = 3.4$ Hz, H-4e), 5.63 (d, 1 H, $J_{5,\text{NH}} = 9.0$ Hz, NH-b), 5.54 (dt, 1 H, $J_{3\text{eq},4} = 5.3$ Hz, $J_{3\text{ax},4} = 12.2$ Hz, $J_{4,5} = 10.2$ Hz, H-4g), 5.45 (d, 1 H, $J_{5,\text{NH}} = 8.7$ Hz, NH-g), 5.43 (d, 1 H, $J_{5,\text{NH}} = 8.5$ Hz, NH-a), 5.34 (m, 1 H, H-4b), 5.32 (dd, 3 H, $J_{6,7} = 2.1$ Hz, $J_{7,8} = 9.0$ Hz, $J_{1,2} = 7.8$ Hz, H-7b, 7g, and 2e), 5.27 (d, 1 H, NH-f), 5.24 (t, 1 H, $J_{1,2} = 7.8$ Hz, $J_{2,3} = 9.5$ Hz, H-2c), 5.21 (d, 1 H, $J_{1,2} = 8.5$ Hz, H-1d), 5.17 (m, 3 H, $J_{1,2} = 7.8$ Hz, $J_{7,8} = 9.0$ Hz, H-2h, 8b, and 8g), 5.08 (m, 1 H, H-4f), 5.01 (dd, 1 H, $J_{6,7} = 1.7$ Hz, $J_{7,8} = 11.2$ Hz, H-7a), 4.96 (br dt, 1 H, H-4a), 4.86 and 4.62 (2 d, 2 H, $J_{\text{gem}} = 10.9$ Hz, PhCH_2), 4.84 (dd, 1 H, $J_{6,7} = 2.9$ Hz, $J_{7,8} = 9.2$ Hz, H-7f), 4.75 (br dd, 1 H, H-3d), 4.67 (d, 1 H, $J_{1,2} = 7.8$ Hz, H-1c), 4.63 (m, 1 H, $J_{7,8} = 9.2$ Hz, H-8f), 4.55 (br d, 1 H, H-6c), 4.50 and 4.28 (2 d, 2 H, $J_{\text{gem}} = 12.2$ Hz, PhCH_2), 4.42 (d, 3 H, $J_{1,2} = 7.8$ Hz, H-1e, 3e, and 1h), 4.33–4.19 (m, 10 H, H-9'b, 9'g, 9'a, 9'f, 8a, 6e, 6'e, 6'd, 6d, and 5b), 4.11 (q, 1 H, $J_{4,5} = 10.2$ Hz, $J_{5,6} = 10.4$ Hz, $J_{5,\text{NH}} = 8.7$ Hz, H-5g), 4.06–3.97 (m, 12 H, H-5f, 6b, 5a, 3c, 9a, 9b, 4c, 4h, 6'c, 9b, 5e, and 5d), 3.92 (dd, 1 H, $J_{5,6} = 10.4$ Hz, $J_{6,7} = 2.1$ Hz, H-6g), 3.87 (m, 1 H, $\text{OCH}_2\text{CH}_2\text{SiMe}_3$), 3.85–3.72 (m, 8 H, H-6a, 6f, 3h, 9f, 2d, and COOMe-a), 3.56 (m, 2 H, $J_{\text{gem}} = 10.7$ Hz, H-6'h and 5c), 3.50 (dd, 1 H, $J_{\text{gem}} = 10.7$ Hz, H-6h), 3.41 (m, 1 H, $\text{OCH}_2\text{CH}_2\text{SiMe}_3$), 3.35 (m, 1 H, H-5h), 3.22 (s, 3 H, COOMe-f), 2.58 (br dd, 1 H, H-3eq-b), 2.50 (dd, 1 H, $J_{3\text{eq},4} = 5.3$ Hz, $J_{\text{gem}} = 12.2$ Hz, H-3eq-g), 2.27 (br dd, 1 H, H-3eq-a), 2.11–1.80 (m, 55 H, 17 Ac, H-3ax-b, 3ax-a, 3eq-f, and 3ax-f), 1.63 (t, 1 H, $J_{3\text{ax},4} = J_{\text{gem}} = 12.2$ Hz, H-3ax-g), 0.81 (m, 2 H, $\text{OCH}_2\text{CH}_2\text{SiMe}_3$), -0.12 (s, 9 H, $\text{OCH}_2\text{CH}_2\text{SiMe}_3$); ^{13}C NMR (100 MHz, CDCl_3) δ 170.9, 170.6, 170.5, 170.5, 170.5, 170.4, 170.3, 170.2, 170.2, 169.8, 169.7, 169.7, 169.6, 167.4, 167.2, 166.1, 165.9, 165.9, 165.7, 165.4, 164.9, 164.6, 164.1, 138.2, 138.2, 133.2, 133.1, 133.1, 132.9, 132.8, 132.7, 130.2, 130.1, 129.9, 129.9, 129.7, 129.6, 129.6, 129.5, 129.5, 129.4, 128.9, 128.5, 128.4, 128.4, 128.3, 128.2, 128.2, 127.9, 127.8, 127.7, 127.4, 127.1, 101.2, 100.4, 100.1, 99.8, 99.5, 99.0, 97.1, 96.0, 80.8, 77.2, 76.4, 76.2, 74.7, 74.0, 73.8, 73.2, 73.1, 72.5, 72.1, 72.0, 71.6, 71.3, 70.9, 70.4, 69.8, 69.5, 69.3, 69.2, 69.0, 68.7, 68.5, 68.4, 68.0, 67.2, 67.1, 67.0, 66.9, 66.7, 63.5, 62.1, 62.0, 61.8, 54.3, 53.1, 52.3, 49.1, 48.9, 38.7, 38.2, 35.5, 35.4, 29.6, 23.0, 23.0, 22.9, 22.7, 20.7, 20.7, 20.6, 20.6, 20.6, 20.5, 20.3, 20.2, 17.8, -1.5 ; HRMS (ESI) m/z found $[\text{M} + 2\text{Na}]^{2+}$ 1763.0599, $\text{C}_{171}\text{H}_{193}\text{N}_5\text{O}_{71}\text{Si}$ calcd for $[\text{M} + 2\text{Na}]^{2+}$ 1763.0599.

Benzyl [Methyl 5-acetamido-8-*O*-(5-acetamido-4,7,8,9-tetra-*O*-acetyl-3,5-dideoxy-*D*-glycero- α -*D*-galacto-2-nonulopyranosylono-1',9-lactone)-4,7-di-*O*-acetyl-3,5-dideoxy-*D*-glycero- α -*D*-galacto-2-nonulopyranosylono-(2 \rightarrow 3)-2,4,6-tri-*O*-benzoyl- β -*D*-galactopyranosyl-(1 \rightarrow 3)-2-acetamido-4,6-di-*O*-benzoyl-2-deoxy- β -*D*-galactopyranosyl-(1 \rightarrow 4)-[methyl 5-acetamido-8-*O*-(5-acetamido-4,7,8,9-tetra-*O*-acetyl-3,5-dideoxy-*D*-glycero- α -*D*-galacto-2-nonulopyranosylono-1',9-lactone)-4,7-di-*O*-acetyl-3,5-dideoxy-*D*-glycero- α -*D*-galacto-2-nonulopyranosylono-(2 \rightarrow 3)]-2,6-di-*O*-benzoyl- β -*D*-galactopyranosyl-(1 \rightarrow 4)-2,3,6-tri-*O*-benzyl- β -*D*-glucopyranosyl-(1 \rightarrow 6)-2,3,4-tri-*O*-benzyl- β -*D*-glucopyranoside (43). To a mixture of **20** (140 mg, 0.0454 mmol) and **39** (66 mg, 0.0681 mmol) in CH₂Cl₂ (1.1 mL) was added 4 Å molecular sieves (AW-300) (210 mg) at room temperature. After stirring for 2 h and then cooling to 0 °C, TMSOTf (1.0 μ L, 5.53 μ mol) was added to the mixture. After stirring for 5 h at 0 °C as the reaction was monitored by TLC (PhMe/MeOH = 4:1), the reaction was quenched by saturated NaHCO₃ (aq) and filtered through a Celite pad, and the pad was washed with CHCl₃. The combined filtrate and washings were washed with brine, dried over Na₂SO₄, and concentrated. The resulting residue was purified by flash column chromatography using PhMe/MeOH (8:1) as the eluent to give **43** (140 mg, 80%): [α]_D = -5.0 (c 2.0, CHCl₃); ¹H NMR (600 MHz, CDCl₃) δ 8.13–6.94 (m, 70 H, 14 Ph), 5.97 (br s, 1 H, H-4d), 5.78 (br d, 1 H, NH-d), 5.70 (d, 1 H, J_{3,4} = 3.4 Hz, H-4e), 5.55 (dt, 1 H, J_{3eq,4} = 4.8 Hz, J_{3ax,4} = 13.0 Hz, J_{4,5} = 10.3 Hz, H-4g), 5.52 (d, 1 H, J_{5,NH} = 9.6 Hz, NH-b), 5.39–5.31 (m, 6 H, H-4b, 7b, 7g, and 3 NH), 5.28 (m, 1 H, H-2e), 5.24 (d, 1 H, J_{1,2} = 8.9 Hz, H-1d), 5.20 (near t, 1 H, J_{1,2} = 7.5 Hz, J_{2,3} = 8.2 Hz, H-2c), 5.16 (m, 2 H, H-8b and 8g), 5.08 (m, 1 H, H-4f), 5.01 (dd, 1 H, J_{6,7} = 1.7 Hz, J_{7,8} = 10.3 Hz, H-7a), 4.95 (dt, 1 H, J_{3eq,4} = 4.1 Hz, H-4a), 4.92–4.20 (14 d, 14 H, PhCH₂), 4.84 (dd, 1 H, H-7f), 4.67 (d, 1 H, J_{1,2} = 7.5 Hz, H-1c), 4.63 (dt, 1 H, H-8f), 4.41 (d, 3 H, J_{1,2} = 7.5 Hz, H-1e, 3e, and H-1h), 4.36 (d, 1 H, J_{1,2} = 7.5 Hz, H-1h), 4.28–4.18 (m, 7 H, H-9'b, 9'g, 9'f, 9a, 8a, 6b, and 5b), 4.13–3.98 (m, 8 H, H-5g, 5a, 3c, 4c, 5f, 9'a, 9b, and 9g), 3.91–3.35 (m, 11 H, Glc unit), 3.89 (br dd, 1 H, H-6g), 3.85 (br dd, 1 H, H-6f), 3.83 (br dd, 1 H, J_{6,7} = 1.7 Hz, H-6a), 3.78 (s, 3 H, COOMe-a), 3.70 (br d, 1 H, H-9'f), 3.60 (m, 1 H, H-2d), 3.24 (s, 3 H, COOMe-f), 3.15 (m, 1 H, H-5h), 2.57 (dd, 1 H, J_{gem} = 13.0 Hz, J_{3eq,4} = 4.8 Hz, H-3eq-b), 2.50 (dd, 1 H, J_{gem} = 13.0 Hz, J_{3eq,4} = 4.8 Hz, H-3eq-g), 2.29 (br dd, 1 H, H-3eq-a), 2.11–1.78 (m, 55 H, 17 Ac, H-3ax-b, 3ax-a, 3ax-f, and 3eq-f), 1.63 (t, 1 H, J_{gem} = J_{3ax,4} = 13.0 Hz, H-3ax-g); ¹³C NMR (125 MHz, CDCl₃) δ 171.0, 170.7, 170.6, 170.6, 170.5, 170.5, 170.4, 170.3, 170.2, 169.8, 169.8, 169.7, 167.4, 167.3, 166.1, 166.0, 165.7, 165.7, 165.4, 164.7, 164.2, 138.8, 138.5, 138.5, 138.3, 138.2, 138.0, 137.5, 133.3, 133.2, 133.1, 133.0, 132.8, 132.8, 130.3, 129.9, 129.9, 129.7, 129.7, 129.6, 129.6, 129.5, 129.4, 129.3, 128.5, 128.3, 128.3, 128.2, 128.2, 128.0, 128.0, 127.8, 127.7, 127.6, 127.4, 127.3, 103.7, 102.6, 101.4, 100.0, 99.5, 99.0, 97.1, 96.0, 84.6, 82.8, 82.2, 81.5, 78.2, 76.2, 76.0, 75.6, 75.1, 75.0, 74.8, 74.8, 74.7, 74.3, 74.0, 73.3, 73.1, 72.5, 72.2, 72.0, 71.6, 71.4, 71.1, 70.8, 70.4, 69.9, 69.8, 69.5, 69.4, 69.4, 69.2, 69.0, 68.8, 68.6, 68.4, 68.4, 68.1, 67.3, 67.0, 66.7, 66.7, 63.4, 62.1, 62.0, 61.9, 54.6, 53.2, 52.4, 49.1, 48.9, 48.9, 38.7, 38.3, 35.7, 35.4, 29.6, 23.1, 23.0, 22.8, 20.8, 20.7, 20.7, 20.7, 20.6, 20.6, 20.3, 20.3; HRMS (ESI) *m/z* found [M + 2Na]²⁺ 1967.1553, C₂₀₀H₂₁₇N₅O₇₅ calcd for [M + 2Na]²⁺ 1967.1557.

[Methyl 5-acetamido-8-*O*-(5-acetamido-4,7,8,9-tetra-*O*-acetyl-3,5-dideoxy-*D*-glycero- α -*D*-galacto-2-nonulopyranosylono-1',9-lactone)-4,7-di-*O*-acetyl-3,5-dideoxy-*D*-glycero- α -*D*-galacto-2-nonulopyranosylono-(2 \rightarrow 3)-2,4,6-tri-*O*-benzoyl- β -*D*-galactopyranosyl-(1 \rightarrow 3)-2-acetamido-4,6-di-*O*-benzoyl-2-deoxy- β -*D*-galactopyranosyl-(1 \rightarrow 4)-[methyl 5-acetamido-8-*O*-(5-acetamido-4,7,8,9-tetra-*O*-acetyl-3,5-dideoxy-*D*-glycero- α -*D*-galacto-2-nonulopyranosylono-1',9-lactone)-4,7-di-*O*-acetyl-3,5-dideoxy-*D*-glycero- α -*D*-galacto-2-nonulopyranosylono-(2 \rightarrow 3)]-2,6-di-*O*-benzoyl- β -*D*-galactopyranosyl-(1 \rightarrow 4)-2-*O*-benzoyl- β -*D*-glucopyranosyl-(1 \rightarrow 1)-(2*S*,3*R*,4*E*)-3-*O*-benzoyl-2-octadecanamide-4-moctadecene-1,3-diol (44). To a solution of **40** (144 mg, 0.0351 mmol) in CH₂Cl₂ (4.0 mL) was added trifluoroacetic acid (2.0 mL)

at 0 °C. After stirring for 1 h at 0 °C as the reaction was monitored by TLC (PhMe/MeOH = 4:1), the reaction mixture was diluted with CHCl₃. The organic layer was washed with ice-cooled saturated Na₂CO₃ (aq) (twice) and brine. After drying over Na₂SO₄ and being concentrated, the resulting residue was purified by flash column chromatography using CHCl₃/MeOH (25:1 to 10:1) to give **44** (131 mg, 97%): [α]_D = +4.7 (c 0.5, CHCl₃); ¹H NMR (600 MHz, CDCl₃) δ 8.08–7.28 (m, 45 H, 9 Ph), 5.94 (br s, 1 H, H-4d), 5.86 (m, 1 H, J_{4,5} = 15.1 Hz, J_{5,6} = 7.5 Hz, J_{5,6'} = 6.9 Hz, H-5i), 5.74 (d, 1 H, J_{2,NH} = 9.6 Hz, NH-i), 5.70 (d, 1 H, J_{3,4} = 3.4 Hz, H-4e), 5.57 (m, 1 H, H-4g), 5.54 (t, 1 H, J_{2,3} = J_{3,4} = 8.2 Hz, H-3i), 5.42 (dd, 1 H, J_{3,4} = 8.2 Hz, J_{4,5} = 15.1 Hz, H-4i), 5.38 (d, 1 H, J_{5,NH} = 10.9 Hz, NH-f), 5.34 (m, 3 H, H-7b, 7g, and 2c), 5.28 (m, 1 H, H-2e), 5.20 (m, 2 H, H-8b and 8g), 5.16 (d, 1 H, J_{1,2} = 8.2 Hz, H-1d), 5.10 (m, 2 H, H-4f and 2h), 5.00 (br dd, 1 H, H-7a), 4.92 (m, 1 H, H-4a), 4.86 (d, 2 H, H-7f and 3d), 4.77 (d, 1 H, J_{1,2} = 7.5 Hz, H-1c), 4.72 (m, 1 H, H-6'c), 4.65 (br dt, 1 H, H-8f), 4.61 (m, 1 H, H-6'e), 4.45–4.39 (m, 7 H, H-1e, 3e, 1h, 4c, 5d, 5e, and 2i), 4.28–4.21 (m, 9 H, H-3c, 8a, 9'b, 9'g, 9'a, 9f, 5c, 6c, and 6e), 4.12 (q, 2 H, J_{4,5} = 10.3 Hz, H-5g and 5b), 4.07–3.94 (m, 6 H, H-5f, 9b, 9g, 9a, 6d, and 6'd), 3.92–3.83 (m, 7 H, H-3h, 6g, 6b, 6f, 6a, 5a, and 1i), 3.77 (t, 1 H, J_{4,5} = 9.6 Hz, H-4h), 3.76 (m, 4 H, COOMe-a and H-9'f), 3.64 (m, 1 H, H-2d), 3.50 (dd, 1 H, J_{gem} = 9.6 Hz, J_{1,2} = 3.4 Hz, H-1'i), 3.24 (s, 3 H, COOMe-f), 3.16 (t, 1 H, J_{gem} = 10.9 Hz, H-6h), 3.05 (d, 1 H, J_{4,5} = 9.6 Hz, H-5h), 2.91 (br dd, 1 H, J_{gem} = 10.9 Hz, H-6'h), 2.66 (dd, 1 H, 6h-OH), 2.59 (br dd, 1 H, H-3eq-b), 2.50 (dd, 1 H, J_{3eq,4} = 5.4 Hz, J_{gem} = 12.3 Hz, H-3eq-g), 2.41 (br dd, 1 H, H-3eq-a), 2.11–1.81 (m, 58 H, 17 Ac, H-3eq-f, 3ax-f, 3ax-a, 3ax-b, 6i, 6'i, and NHCOCH₂), 1.63 (t, 1 H, J_{gem} = J_{3ax,4} = 12.3 Hz, H-3ax-g), 1.43 (m, 1 H, NHCOCH₂), 1.26 (m, 52 H, 26 CH₂), 0.88 (t, 6 H, 2 Me); ¹³C NMR (125 MHz, CDCl₃) δ 172.5, 171.1, 170.6, 170.5, 170.3, 170.1, 169.7, 169.6, 167.4, 167.2, 166.1, 166.0, 165.9, 165.7, 165.5, 165.4, 165.3, 164.7, 164.3, 164.1, 138.3, 133.2, 133.1, 133.0, 132.9, 132.7, 130.0, 129.9, 129.8, 129.7, 129.6, 129.5, 129.4, 129.4, 129.2, 129.1, 128.4, 128.3, 128.2, 124.6, 101.5, 101.3, 99.6, 99.4, 99.2, 98.9, 96.8, 95.8, 80.4, 75.8, 73.8, 73.6, 73.0, 72.7, 72.5, 72.3, 71.9, 71.5, 71.3, 71.7, 71.0, 70.5, 69.8, 69.7, 69.3, 69.3, 68.9, 68.7, 68.5, 68.2, 67.1, 66.9, 66.6, 66.6, 66.1, 64.0, 63.9, 62.3, 62.3, 62.2, 62.1, 62.0, 61.9, 61.9, 59.4, 54.7, 54.6, 54.6, 54.5, 54.5, 53.1, 52.2, 50.2, 49.1, 49.1, 48.9, 48.8, 48.7, 38.6, 38.2, 36.5, 35.4, 32.1, 31.7, 29.5, 29.5, 29.4, 29.4, 29.3, 29.2, 29.1, 28.7, 25.4, 22.9, 22.9, 22.7, 22.5, 20.7, 20.6, 20.6, 20.5, 20.5, 20.3, 20.2, 14.0; HRMS (ESI) *m/z* found [M + 2Na]²⁺ 1948.7547, C₁₉₅H₂₄₂N₅O₇₄ calcd for [M + 2Na]²⁺ 1948.7571.

2-(Trimethylsilyl)ethyl [Methyl 5-acetamido-8-*O*-(5-acetamido-4,7,8,9-tetra-*O*-acetyl-3,5-dideoxy-*D*-glycero- α -*D*-galacto-2-nonulopyranosylono-1',9-lactone)-4,7-di-*O*-acetyl-3,5-dideoxy-*D*-glycero- α -*D*-galacto-2-nonulopyranosylono-(2 \rightarrow 3)-2,4,6-tri-*O*-benzoyl- β -*D*-galactopyranosyl-(1 \rightarrow 3)-2-acetamido-4,6-di-*O*-benzoyl-2-deoxy- β -*D*-galactopyranosyl-(1 \rightarrow 4)-[methyl 5-acetamido-8-*O*-(5-acetamido-4,7,8,9-tetra-*O*-acetyl-3,5-dideoxy-*D*-glycero- α -*D*-galacto-2-nonulopyranosylono-1',9-lactone)-4,7-di-*O*-acetyl-3,5-dideoxy-*D*-glycero- α -*D*-galacto-2-nonulopyranosylono-(2 \rightarrow 3)]-2,6-di-*O*-benzoyl- β -*D*-galactopyranosyl-(1 \rightarrow 4)-2-*O*-benzoyl- β -*D*-glucopyranoside (45). To a solution of **42** (75 mg, 0.0215 mmol) in EtOH (2.1 mL) was added palladium hydroxide [20 wt % Pd (dry basis) on carbon, wet] (75 mg) at room temperature. After stirring for 33 h at 40 °C under a H₂ atmosphere as the reaction was monitored by TLC (PhMe/MeOH = 4:1), the mixture was filtered through a Celite pad and washed with CHCl₃. The combined filtrate and washings were concentrated. The resulting residue was purified by flash column chromatography using CHCl₃/MeOH (20:1 to 15:1) as the eluent to give **45** (63 mg, 90%): [α]_D = -2.3° (c 1.2, CHCl₃); ¹H NMR (500 MHz, CDCl₃) δ 8.10–7.26 (m, 40 H, 8 Ph), 5.93 (br s, 1 H, H-4d), 5.74 (br d, 1 H, NH-b), 5.70 (d, 1 H, J_{3,4} = 3.6 Hz, H-4e), 5.55 (dt, 1 H, J_{4,5} = 10.4 Hz, J_{3eq,4} = 5.1 Hz, J_{3ax,4} = 12.2 Hz, H-4g), 5.45 (d, 1 H, J_{5,NH} = 10.0 Hz, NH-b), 5.34 (m, 7 H, H-4b, 2c, 2e, 7g, 7b, NH-g, and NH-f), 5.18 (d, 1 H, J_{1,2} = 8.0 Hz, H-1d), 5.17 (m, 2 H, H-8b and 8g), 5.11 (t, 1 H, J_{1,2}

= 8.0 Hz, H-2h), 5.10 (m, 2 H, H-4f and NH-a), 5.01 (dd, 1 H, $J_{6,7}$ = 1.7 Hz, $J_{7,8}$ = 8.5 Hz, H-7a), 4.92 (m, 2 H, H-3d and 4a), 4.87 (dd, 1 H, $J_{6,7}$ = 1.7 Hz, $J_{7,8}$ = 9.5 Hz, H-7f), 4.77 (d, 1 H, $J_{1,2}$ = 8.0 Hz, H-1c), 4.71 (br dd, 1 H, H-6c), 4.63 (dt, 1 H, $J_{7,8}$ = 9.5 Hz, H-8f), 4.59 (m, 1 H, H-6e), 4.50 (d, 1 H, $J_{1,2}$ = 8.0 Hz, H-1h), 4.42 (m, 4 H, H-1e, 3e, 4c, and 6d), 4.29–4.19 (m, 11 H, H-5b, 8a, 3c, 6'e, 6b, 5e, 6'c, 9b, 9g, 9'f, and 9a), 4.12 (q, 1 H, $J_{4,5}$ = 10.4 Hz, $J_{5,6}$ = 10.2 Hz, H-5g), 4.06–3.98 (m, 5 H, H-5f, 9'a, 9'b, 9'g, and 6'd), 3.94–3.85 (m, 7 H, H-3h, 6g, 6f, 5a, 5c, 5d, and OCH₂CH₂SiMe₃), 3.81 (dd, 1 H, $J_{6,7}$ = 1.7 Hz, H-6a), 3.75 (m, 4 H, H-9f and COOMe-a), 3.69 (t, 1 H, $J_{3,4}$ = $J_{4,5}$ = 8.7 Hz, H-4h), 3.63 (br q, 1 H, $J_{1,2}$ = 8.0 Hz, H-2d), 3.46 (m, 2 H, H-6h and OCH₂CH₂SiMe₃), 3.43 (m, 1 H, H-6'h), 3.30 (m, 1 H, H-5h), 3.24 (s, 3 H, COOMe-f), 2.57 (br dd, 1 H, H-3eq-b), 2.51 (dd, 1 H, J_{gem} = 12.2 Hz, $J_{3\text{eq},4}$ = 5.1 Hz, H-3eq-g), 2.45 (m, 1 H, H-3eq-a), 2.11–1.70 (m, 55 H, 17 Ac, H-3ax-b, 3ax-a, 3eq-f, and 3ax-f), 1.64 (t, 1 H, J_{gem} = $J_{3\text{ax},4}$ = 12.2 Hz, H-3ax-g), 0.82 (m, 2 H, OCH₂CH₂SiMe₃), 0.00 and -0.12 (2 s, 9 H, OCH₂CH₂SiMe₃); ¹³C NMR (100 MHz, CDCl₃) δ 171.3, 170.7, 170.7, 170.6, 170.5, 170.4, 170.4, 170.4, 170.3, 169.9, 169.8, 169.8, 169.8, 169.7, 167.6, 167.3, 166.2, 166.1, 166.0, 165.6, 165.4, 165.1, 164.6, 164.5, 164.2, 133.5, 133.3, 133.3, 133.1, 133.0, 132.8, 130.3, 130.2, 129.9, 128.8, 129.7,

129.6, 129.6, 129.2, 128.6, 128.5, 128.4, 128.4, 128.2, 101.6, 100.5, 99.4, 99.2, 99.0, 97.1, 96.2, 81.2, 77.2, 75.8, 74.0, 73.8, 73.4, 73.2, 73.0, 72.8, 72.5, 72.1, 71.7, 71.4, 71.2, 70.6, 70.1, 69.9, 69.6, 69.4, 68.9, 68.8, 68.6, 68.5, 67.4, 67.3, 66.7, 64.0, 62.1, 62.0, 61.8, 60.6, 54.8, 53.2, 52.4, 49.3, 49.1, 49.1, 48.9, 38.6, 38.5, 38.3, 36.1, 35.6, 29.6, 23.1, 23.1, 23.0, 22.9, 20.8, 20.7, 20.7, 20.7, 20.6, 20.4, 20.3, 17.9, -1.1; HRMS (ESI) *m/z* found [M + 2Na]²⁺ 1673.0130, C₁₅₇H₁₈₁N₅O₇₁Si calcd for [M + 2Na]²⁺ 1673.0130.

Acknowledgment. This work was financially supported by the Ministry of Education, Culture, Sports, Science, and Technology (MEXT) of Japan (Grant-in-Aid for Scientific Research to M.K., No. 17101007).

Supporting Information Available: Experimental data for compounds **8–10**, **12**, **13**, **17–19**, **23**, **24**, and **26–31**, and copies of ¹H and ¹³C NMR spectra for all new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

JO8027888