

Review Article

Asbestos Induces Reduction of Tumor Immunity

Naoko Kumagai-Takei,¹ Megumi Maeda,^{1,2} Ying Chen,^{1,3} Hidenori Matsuzaki,¹ Suni Lee,¹
Yasumitsu Nishimura,¹ Junichi Hiratsuka,⁴ and Takemi Otsuki¹

¹ Department of Hygiene, Kawasaki Medical School, 577 Matsushima, Kurashiki 701-0192, Japan

² Division of Bioscience, Department of Biofunctional Chemistry, Okayama University Graduate School of Natural Science and Technology, 3-1-1 Tsushima-naka, Okayama 700-8530, Japan

³ Division of Pneumoconiosis, School of Public Health, China Medical University, 92 North 2nd, Heping District, Shenyang 110001, China

⁴ Department of Radiation Oncology, Kawasaki Medical School, 577 Matsushima, Kurashiki 701-0192, Japan

Correspondence should be addressed to Takemi Otsuki, takemi@med.kawasaki-m.ac.jp

Received 15 June 2011; Accepted 18 July 2011

Academic Editor: E. Shevach

Copyright © 2011 Naoko Kumagai-Takei et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Asbestos-related cancers such as malignant mesothelioma and lung cancer are an important issue in the world. There are many conflicts concerning economical considerations and medical evidence for these cancers and much confusion regarding details of the pathological mechanisms of asbestos-induced cancers. For example, there is uncertainty concerning the degree of danger of the iron-absent chrysotile compared with iron-containing crocidolite and amosite. However, regarding bad prognosis of mesothelioma, medical approaches to ensure the recognition of the biological effects of asbestos and the pathological mechanisms of asbestos-induced carcinogenesis, as well as clinical trials to detect the early stage of mesothelioma, should result in better preventions and the cure of these malignancies. We have been investigating the immunological effects of asbestos in relation to the reduction of tumor immunity. In this paper, cellular and molecular approaches to clarify the immunological effects of asbestos are described, and all the findings indicate that the reduction of tumor immunity is caused by asbestos exposure and involvement in asbestos-induced cancers. These investigations may not only allow the clear recognition of the biological effects of asbestos, but also present a novel procedure for early detection of previous asbestos exposure and the presence of mesothelioma as well as the chemoprevention of asbestos-related cancers.

1. Introduction

The fact that inhaled asbestos causes malignant mesothelioma and lung cancer is an enormous medical and social problem. Canada's decision to expand asbestos mining and export to developing countries in which asbestos has not been banned is unsettling [1]. People are sometimes influenced by economical forces even though they know many people have suffered from malignant diseases caused by these minerals, and their decisions appear to be made for financial reasons.

In Japan, the asbestos issue erupted in the summer of 2005 [2–4]. Residents were suddenly informed that asbestos, which was used in large amounts from the early 1950s up to the early 1990s in Japan with a maximum usage

of approximately 352,000 tons in 1974, caused malignant mesothelioma (MM). Residents that lived near the asbestos handling manufacturer Kubota Corporation, in Amasaka City, Hyogo Prefecture, developed MM. They had never worked in the asbestos-handling manufacture industry. In addition, medical information regarding MM induced anxiety in the Japanese people, since the prognosis is very poor, and there is no certain way to detect the cancer in the very early stage of the disease. Furthermore, people could not remember being exposed to asbestos 30 to 40 years ago. To reduce the anxieties of the Japanese people, epidemiological analyses regarding the Amagasaki area proceeded, and clinical and basic research was conducted on the biological effects of asbestos and early detection of mesothelioma. It is in this context that the authors

became involved in the project “Comprehensive Approach on Asbestos-Related Diseases”, supported by the “Special Coordination Funds for Promoting Science and Technology” (Dr. Takemi Otsuki, Department of Hygiene, Kawasaki Medical School, Kurashiki, Japan) from 2006 to 2010. In this project, a case and clinical specimen registration system was established. A feasibility clinical trial was established and involved a combined trimodality therapy using anticancer chemotherapy with cisplatin and pemetrexed, following by extrapleural pneumonectomy and postoperative radiation therapy for early-stage mesothelioma patients [5, 6]. Furthermore, early detection procedures were developed using serum or pleural effusions to measure soluble mesothelin-related peptide (SMRP) and other markers such as osteopontin, vascular endothelial growth factor (VEGF) and angiotropin-1 [7–9], as well as procedures for detection of circulating mesothelioma cells and circulating epithelial cells using peripheral blood [10, 11].

For the basic research, the project “Comprehensive Approach on Asbestos-Related Diseases” included three subgroups: (1) analyses of cellular and molecular characteristics using mesothelioma cell lines, (2) investigation of asbestos-induced carcinogenesis using an animal model, and (3) study of the immunological effects of silica/asbestos.

The first subgroup explored novel tumor suppressor gene(s) in mesothelioma cells and found that the serine/threonine-protein kinase (LATS2) gene is inactivated in approximately one-third of mesothelioma cell lines and is a candidate for a novel tumor suppressor in MM [12]. In addition, they found the possibility that the Yes-associated protein (YAP) involved the NF2/Merlin-hippo signaling pathway and that LATS2 may constitutively dephosphorylate and act as an oncogene to bind with the TEAD transcription factor to enhance the cell cycle and resistance to apoptosis [13]. In addition, mesothelioma-specific epigenetic profiles were identified for differential diagnosis with lung adenomatous cancers [14].

The second subgroup confirmed the importance of iron in asbestos-induced carcinogenesis. Findings showed that not only iron-containing crocidolite and amosite, but also chrysotile asbestos caused mesothelioma when these materials were injected into the peritoneal region of a rat. Even individual rats having mesothelioma caused by the injection of iron-absent chrysotile showed numerous depositions of iron in the spleen, liver, and kidney. In addition, adding nitrilotriacetate (NTA) to chrysotile-injected rats induced the acceleration of mesothelioma formation, suggesting the critical participation of iron for asbestos-induced carcinogenesis even for chrysotile. Although the detailed mechanisms of this phenomenon are now being explored, the binding ability of chrysotile to hemoglobin and other proteins and the induced hemolysis is a concern [15–18]. Moreover, the importance of a homozygous deletion of CDKN2A/2B was found in rat mesothelioma with the suggestion that this deletion seems to be fundamental for the development of mesothelioma, since these genes are also known to be homozygously deleted in human mesothelioma [19].

We have performed the third subtheme concerning the “immunological effects of silica/asbestos”. In this paper, we introduce our findings and considerations regarding involvement of reduced tumor immunity caused by asbestos exposure to immunocompetent cells as the basic condition in asbestos-exposed people who may develop MM.

2. Immunological Effects of Asbestos

Asbestos comprises a set of six naturally occurring silicate minerals (chrysotile as Serpentine and crocidolite, amosite, actinolite, anthophyllite, and tremolite as Amphibole) exploited commercially for their desirable physical properties. They all have in common their asbestiform structure, possessing long (having more than 1:3 aspect ratio, usually approximately 1:20) and thin fibrous crystals [20, 21]. Silica (SiO₂) certainly affects the human immune system, because people exposed to silica not only suffer from respiratory disorders known as silicosis, but also experience complications with autoimmune disorders such as rheumatoid arthritis (known as Caplan’s syndrome), systemic sclerosis, systemic lupus erythematosus, and antineutrophil cytoplasmic antibody- (ANCA-) related vasculitis/nephritis [22–27]. We have, therefore, been exploring the mechanisms involved in silica-induced dysregulation of autoimmunity using case peripheral blood specimens. We had found that there is dysregulated expression of the CD95/Fas molecule, which is very important for the survival of self-recognizing T cell clones. Additionally, analyses of Fas and Fas-related molecules in silicosis patients suggested that there are two populations of T cells: one is the long-term surviving populations probably including self-recognizing clones, and the other is a population repeating apoptosis caused by silica and recruiting from the bone marrow [28, 29]. In addition, our recent studies regarding CD4+25+ and forkhead box P3 (FoxP3)+ regulatory T cells (Treg) suggested that (1) silica activates both responder T cells (Tresp) and Treg, (2) Tresp chronically-activated by silica becomes CD4+25+ (and programmed cell death-1 (PD-1) + as an activated cell marker) expressers, (3) Treg activated by silica express higher CD95/Fas and are sensitive to Fas-mediated apoptosis, and (4) after the ongoing progression of these events, the composition of the peripheral CD4+25+ fraction in silicosis patients changes to reflect a loss of Treg and a gain of activated Tresp, and this reduction of Treg function results in activation of autoimmunity in silicosis patients [30–32].

Since silica influences the human immune system, its mineral silicate, an asbestos, may also have an effect. As we considered silica’s immunological effects from the complications of silicosis and autoimmune diseases, the most important complication of asbestos-exposed people is the occurrence of malignant disease such as MM and lung cancer. In addition, some epidemiological studies suggested a relationship between asbestos exposure and other cancers of the gastrointestinal tract, larynx, kidney, liver, pancreas, ovary, and hematopoietic systems [33–35]. Thus, if asbestos affects the immune system, a reduction of tumor immunity may result and then make people exposed to asbestos

sensitive to the development of malignancies. Of course, asbestos itself possesses carcinogenic activities. As shown in Figure 1, asbestos fibers having iron (or even chrysotile as mentioned above) produce reactive oxygen/nitrogen species (ROS/RNS) causing DNA damage to nearby cells, and fibers are sometimes directly inserted into the cells and injure chromosomes, while retained fibers may adsorb other carcinogens on their surface (known as an asbestos body) [15, 16, 18, 36, 37]. As a result, specific DNA alterations may result, such as inactivation (mostly homozygous deletion) of *p16^{INK4a}/p14^{ARF}*, *NF2/Merlin*, and *LATS2*, and the activation of *YAP* as mentioned above [12, 13]. However, it is difficult to explain why the development of mesothelioma requires 30 to 40 years, and how asbestos-exposed people possess sensitive features for other cancers.

We have been considering that asbestos may affect immunocompetent cells such as CD4+ Tresp, Treg, Th17 T cells, CD8+ cytotoxic T cells (CTL), monocyte-macrophage, natural killer (NK) cells, natural killer T (NKT) cells, and dendritic cells (DC). Firstly, to observe the effects of low-dose and continuous exposure to asbestos (we initially chose chrysotile because this is the most frequently used asbestos in Japan, and our investigations suggested it was not carcinogenic), we employed a human adult leukemia/lymphoma virus-1 (HTLV-1) immortalized polyclonal T cell line, MT-2 [38, 39]. In the next part of this paper, analysis of asbestos exposure to the MT-2 cell line is documented.

3. Transient and Continuous Exposure to Asbestos on a Human T Cell Line

Initially, the cellular alteration of MT-2 cells exposed to transient and high-dose chrysotile was observed to compare various published investigations showing the ability of asbestos exposure to induce ROS production and mitochondrial-pathway-dependent apoptosis in normal alveolar epithelial cells and mesothelial cells, which are the target cells of asbestos-induced carcinogenesis. As shown on the left side of Figure 2, transient and relatively high-dose exposure (25–50 $\mu\text{g}/\text{mL}$, not likely to comprise adhesive cells such as alveolar epithelial or mesothelial cells, since we are using suspended cells, and thus, $\mu\text{g}/\text{mL}$ was used instead of $\mu\text{g}/\text{cm}^2$) caused production of ROS as measured by production of O_2^- using flow cytometry, phosphorylation of proapoptotic molecules in the mitogen-activated protein kinase (MAPK) pathway such as p38 and c-Jun N-terminal kinase (JNK), release of cytochrome-c from mitochondria to the cytosol, BAX overexpression, cleavage of caspase-9 and -3, and thereafter the appearance of apoptosis [40]. These findings resembled the effects of asbestos on alveolar epithelia and mesothelial cells [41–44].

We then conducted a trial to establish a low-dose and continuous exposure cell line model by adding 5 or 10 $\mu\text{g}/\text{mL}$ of chrysotile (doses which cause apoptosis in less than half of cells exposed transiently) to the MT-2 cell culture. After more than eight months exposure with monthly monitoring for the occurrence of apoptosis, and when these cells were re-exposed to fibers one week after

being released from continuously exposed chrysotile, an MT-2 subline which showed resistance to chrysotile-induced apoptosis had been established. As shown on the right side of Figure 2, the continuously exposed subline of MT-2 showed activation of Src-family kinase, increased expression and production of interleukin (IL)-10, phosphorylation of signal transducer and activator of transcription 3 (STAT3) with overexpression of BCL-2 (located downstream of STAT3) [45, 46]. In addition, transforming growth factor (TGF)- β was upregulated [47, 48]. Actually, we had established three independent continuously exposed sublines to chrysotile B and three other sublines exposed to chrysotile A. The altered gene expression of these six continuously exposed sublines in comparison with the original MT-2 cell line was very similar, and the cellular and molecular characteristics of these cell lines in regard to tumor immunity with the *ex vivo* chrysotile exposure model using freshly isolated peripheral blood CD4+ T cells derived from healthy donors was investigated and confirmed using peripheral blood specimens derived from asbestos-exposed patients such as patients with pleural plaque (PP) or MM.

Chemokine receptor, CXC chemokine receptor (CXCR)3, expression and relation with interferon (IFN)- γ .

Using the above-mentioned MT-2 original cell line and the continuously exposed chrysotile sublines, molecules related to tumor immunity were investigated. For example, CXCR3 expression was a focus of investigations, since CXCR3 downregulation in sublines was detected in comparison with the original line using cDNA microarray analysis. It is known that CXCR3 expression and IFN- γ production are induced by T-cell activation and lead to the enhancement of antitumor immune function [49, 50].

From findings using the MT-2 cell line model, as shown in Figure 3(a), all six continuously exposed sublines showed reduced CXCR3 expression on their surface and mRNA expression levels with reduced production and expression of IFN- γ . Production of the Th1-type CXCR3 ligand CXCL10/IP10 was also significantly reduced in all six continuously exposed sublines when compared with the original line. In addition, another Th1-type chemokine, CCL4/MIP-1 β mRNA, was also expressed at low levels in all six sublines compared with the MT-2 original line as previously reported. However, CCR5, the Th1-type receptor for CCL4/MIP-1 β , was not reduced significantly through mRNA expression in MT-2Rsts cells. These results indicated that continuous exposure of MT-2 original cells to asbestos altered the expression of Th1-related chemokines (CXCL10/IP10 and CCL4/MIP-1 β) and chemokine receptors (CXCR3) [51].

Thereafter, we tried to determine whether freshly isolated human peripheral CD4+ T cells show a similar alteration *ex vivo* when proliferation is maintained by IL-2-containing medium in the presence of chrysotile as shown in Figure 3(b). After 40 days of coculture supplemented with IL-2 in the presence or absence of chrysotile, cell surface CXCR3 expression decreased in a dose-dependent manner. Thus, we examined cell surface expression of CXCR3 and CCR5 in CD4+ T cells derived from six healthy donors, since both receptors are preferentially expressed in Th1/effector T cells. The expression of CXCR3 was significantly reduced




FIGURE 1: Schematic model showing mechanisms of asbestos-induced carcinogenesis and genomic/epigenetic changes found in mesothelioma cells and the relationship of the immunological effects of asbestos in regard to reduced tumor immunity.




FIGURE 2: Summarized findings of cellular and molecular events caused by high-dose and transient exposure (left side) and low-dose and continuous exposure (CB1: one of the sublines established) (right side) to chrysotile asbestos using an HTLV-1 immortalized human polyclonal T cell line, MT-2.

following exposure to 10 µg/mL of chrysotile for 28 days although this difference seemed to depend on one case in which the expression decreased remarkably. Even if the culture conditions for the CD4+ T cells was limited to a period of around four weeks, four of the six healthy

donors showed a decrease of CXCR3 expression to various degrees, and it might be concluded that asbestos exposure potentiates reduction of CXCR3 expression in CD4+ T cells. On the other hand, the expression of CCR5 varied among all healthy donors, and there were no significant changes after




FIGURE 3: Schematic representation of asbestos-induced reduction of expression of a chemokine receptor, CXCR3, and expression and production of IFN- γ using the MT-2 cell line model (Org; MT-2 original cell line, and Rst: sublines exposed continuously to a low-dose of chrysotile), an *ex vivo* exposure model using freshly isolated CD4+ T cells from healthy donors (HD), as well as analyses of freshly isolated CD4+ T cells from healthy donors and patients with pleural plaque (PP) and malignant mesothelioma (MM).

seven and 28 days of coculture with chrysotile, as shown previously by the cell line model. These results indicated that CXCR3 expression might be specifically reduced by asbestos exposure. In addition, these experiments revealed decreased IFN- γ expression and production when CD4+ T cells from healthy donors were cultured with chrysotile for 28 days [52].

Finally, analyses of changes in surface CXCR3 expression on freshly isolated CD4+ T cells from asbestos-exposed patients such as PP or MM were compared with those from healthy donors. In addition, IFN- γ expression of CD4+ T cells from these patients and healthy donors was measured with stimulation using anti-CD3/CD28 antibodies with IL-2. As summarized in Figure 3(c), CXCR3 expression was reduced in CD4+ T cells from asbestos patients. A comparison of PP and MM patients showed that the expression level of CXCR3 on CD4+ T cells from MM was decreased although the difference was not statistically significant. Moreover, IFN- γ expression was only reduced in stimulated CD4+ T cells from MM patients, not in those from PP patients [52].

With the findings that CD4+CXCR3+ T cells in lymphocytes from MMs showed a tendency for an inverse correlation with CXCR3's ligand, CXCL10/IP10 in plasma, our results indicate a reduction of tumor immune function in asbestos-exposed patients and suggest that CXCR3, IFN- γ , and

CXCL10/IP10 may be candidates to detect and monitor disease status.

4. Alteration of NK Cells and Others

As shown in Figure 4, the effects of asbestos on other immunocompetent cells such as Treg, CD8+ CTL, and NK cells were investigated. As mentioned above with the MT-2 cell line model, sublines continuously exposed to chrysotile showed overproduction of IL-10 and TGF- β . It is well known that these cytokines are a typical soluble factor produced from Treg to function with a suppressive effect on activated responder T cells. On the other hand, it is also reported that MT-2 cells possess a Treg function, since cells express CD4 and CD25 with nuclear expression of FoxP3. Taken together, continuous exposure to chrysotile produces a stronger Treg function, at least with the capacity to produce soluble functional factors (i.e., IL-10 and TGF- β) [47, 48]. At present, we have been studying alteration of Treg function using the MT-2 cell line model, and preliminary findings indicate asbestos may enhance Treg function.

In regard to tumor immunity, CD8+ CTL and NK cells are very important players, since they directly kill tumor cells even when individually restricted with major histocompatibility complexes. Investigations have just started with




FIGURE 4: Schematic representation of findings showing asbestos-induced reduction of tumor immunity on CD4+ T cells, CD4+25+FoxP3+ regulatory T cells (Treg), T helper (Th)17, CD8+ cytotoxic T cells (CTL), natural killer (NK) cells, monocyte-macrophage, dendritic cells (DC), and natural killer T cells (NKT).

CD8+ CTL, but in *ex vivo* experimental conditions designed to produce CD8+ CTL proliferation and differentiation using freshly isolated peripheral blood mononuclear cells from healthy donors, the addition of asbestos seems to result in reduced proliferation and differentiation of CTL. Although detailed analyses concerning the roles of cytokines surrounding CTL differentiation are being performed, our ongoing studies suggest that asbestos reduces CTL activities.

Regarding NK cells, cellular and molecular analyses have been conducted using a human NK cell line, YT-A1, exposed continuously to asbestos in an *ex vivo* exposure model using freshly isolated NK cells from health donors, as well as asbestos-exposed patients such as PP and MM.

Focusing on the NK cell-activating receptors, including NKG2D (also known as KLRK1 (killer cell lectin-like receptor subfamily K, member 1), klr and CD314, binding to a diverse family of ligands that include MHC class I chain-related A and B proteins and UL-16 binding proteins, where ligand-receptor interactions can result in the activation of NK and T cells), 2B4 (also known as NAIL; Nmrk; NKR2B4; SLAMF4 and CD244, mediate nonmajor histocompatibility complex (MHC) restricted killing), and NKp46 (also known as NCR1 (natural cytotoxicity triggering receptor 1), LY94 and CD335, constituting a natural cytotoxic receptor family with NKp44 and NKp30, and being important in killing tumor cells and dendritic cells), the YT-A1 human NK cell line exposed continuously to chrysotile asbestos revealed reduced expression of NKG2D and 2B4 [53]. The reduced phosphorylation of extracellular signal-regulated kinase

(ERK) and subsequent reduction of degranulation of perforin and granzyme B resulting from reduced cytotoxicity were observed in this cell line model [54]. Similar to the cell line model, the *ex vivo* exposure model using freshly isolated NK cells from healthy donors revealed a reduction of NKp46 expression. Furthermore, freshly isolated NK cells from MM patients showed reduced killing function compared with those from healthy donors and revealed a lower expression of NKp46 [53]. Moreover, the expression level of NKp46, but not NKG2D or 2B4, and the cytotoxic activity of individual freshly isolated NK cells from health donors and MM patients clearly showed a reverse correlation, indicating that the target molecule of asbestos-induced dysfunction of NK cells is NKp46 [54]. Although further analyses are required regarding the interaction between asbestos-exposed NK cells and other immunocompetent cells such as dendritic cells, monocytes, and macrophages, molecular mechanisms to reduce NKp46 expression and other aspects need to be explored, and surface NKp46 expression levels may be the candidate to monitor the level of tumor immunity in asbestos-exposed patients [55].

Further investigations are needed to examine the effects of asbestos exposure on other types of immunocompetent cells such as Th17 dendritic cells, NKT, and the monocyte-macrophage lineage, and to investigate why asbestos seems to reduce tumor immunity in the total network of the immunological surveillance system.

In addition, although we have mainly analyzed the effects of chrysotile asbestos on the human immune system,

differences and similarities between the different types of fibers should also be investigated.

5. Conclusion

We have been investigating the effects of asbestos exposure on the human immune system in regard to tumor immunity and found that people exposed to asbestos possess reduced tumor immunity, making them sensitive to cancer development. Although these studies may contribute to the clear recognition of the biological effects of asbestos, the variety of alterations in immunocompetent cells may be the factor that allows detection of previous asbestos exposure and the occurrence of cancer in people that live or have lived near asbestos-handling manufacturers. Furthermore, to recover tumor immunity using physiologically active substances in foods or derived from plants may be an effective method for the chemical prevention of asbestos-induced cancers.

Acknowledgments

The authors specially thank Dr. Masayasu Kusaka (Kusaka Hospital, 1122 Nishikatagami, Bizen, 705-0121, Japan), Dr. Kozo Urakami (Hinase Urakami Iin, 243-4 Hinase, Hinasecho, Bizen, 701-3204, Japan), Professors Takashi Nakano and Kazuya Fukuoka (Department of Respiratory Medicine, Hyogo College of Medicine, 1-1, Mukogawa-cho, Nishinomiya, 663-8501, Japan), and Drs. Takumi Kishimoto, Rika Tabata, Kenichi Genma, and Ms. Yoko Kojima (Okayama Rosai Hospital, 1-10-25, Chikkou Midori-machi, Minamiku, Okayama 702-8055, Japan) for their particular contribution to the organization of patients. They also thank Ms. Tamayo Hatayama, Yoshiko Yamashita, Minako Kato, Tomoko Sueishi, Keiko Kimura, Misao Kuroki, Naomi Miyahara, and Shoko Yamamoto for their technical help. The experimental results performed by the authors and presented partly in this paper were supported by Special Funds for Promoting Science and Technology (H18-1-3-3-1), JSPS KAKENHI (22790550, 22700933, 20390178, 20890270, 19689153, 19790431, 19790411, 18390186, 16390175, and 09670500), the Takeda Science Foundation (Tokutei Kenkyu Josei I, 2008), and Kawasaki Medical School Project Grants (22-B1, 22-A58, 22-A29, 21-401, 21-201, 21-107, 20-411I, 20-210O, 20-109N, 20-402O, and 20-410I).

References

- [1] "Asbestos scandal," *Nature*, vol. 468, no. 7326, 868 pages, 2010.
- [2] S. Kumagai and N. Kurumatani, "Mapping the risk of mesothelioma due to neighborhood asbestos exposure," *American Journal of Respiratory and Critical Care Medicine*, vol. 178, no. 6, pp. 624–629, 2008.
- [3] S. Kumagai and N. Kurumatani, "Asbestos fiber concentration in the area surrounding a former asbestos cement plant and excess mesothelioma deaths in residents," *American Journal of Industrial Medicine*, vol. 52, no. 10, pp. 790–798, 2009.
- [4] S. Kumagai, N. Kurumatani, T. Tsuda, T. Yorifuji, and E. Suzuki, "Increased risk of lung cancer mortality among residents near an asbestos product manufacturing plant," *International Journal of Occupational and Environmental Health*, vol. 16, no. 3, pp. 268–278, 2010.
- [5] T. Yamanaka, F. Tanaka, S. Hasegawa et al., "A feasibility study of induction pemetrexed plus cisplatin followed by extrapleural pneumonectomy and postoperative hemithoracic radiation for malignant pleural mesothelioma," *Japanese Journal of Clinical Oncology*, vol. 39, no. 3, pp. 186–188, 2009.
- [6] S. Hasegawa and F. Tanaka, "Malignant mesothelioma: current status and perspective in Japan and the world," *General Thoracic and Cardiovascular Surgery*, vol. 56, no. 7, pp. 317–323, 2008.
- [7] R. Ohashi, K. Tajima, F. Takahashi et al., "Osteopontin modulates malignant pleural mesothelioma cell functions in vitro," *Anticancer Research*, vol. 29, no. 6, pp. 2205–2214, 2009.
- [8] C. Tabata, N. Hirayama, R. Tabata et al., "A novel clinical role for angiotensin-1 in malignant pleural mesothelioma," *European Respiratory Journal*, vol. 36, no. 5, pp. 1099–1105, 2010.
- [9] A. Yasumitsu, C. Tabata, R. Tabata et al., "Clinical significance of serum vascular endothelial growth factor in malignant pleural mesothelioma," *Journal of Thoracic Oncology*, vol. 5, no. 4, pp. 479–483, 2010.
- [10] Y. Okumura, F. Tanaka, K. Yoneda et al., "Circulating tumor cells in pulmonary venous blood of primary lung cancer patients," *Annals of Thoracic Surgery*, vol. 87, no. 6, pp. 1669–1675, 2009.
- [11] F. Tanaka, K. Yoneda, N. Kondo et al., "Circulating tumor cell as a diagnostic marker in primary lung cancer," *Clinical Cancer Research*, vol. 15, no. 22, pp. 6980–6986, 2009.
- [12] H. Murakami, T. Mizuno, T. Taniguchi et al., "LATS2 is a tumor suppressor gene of malignant mesothelioma," *Cancer Research*, vol. 71, no. 3, pp. 873–883, 2011.
- [13] Y. Sekido, "Inactivation of Merlin in malignant mesothelioma cells and the Hippo signaling cascade dysregulation," *Pathology International*, vol. 61, no. 6, pp. 331–344, 2011.
- [14] Y. Goto, K. Shinjo, Y. Kondo et al., "Epigenetic profiles distinguish malignant pleural mesothelioma from lung adenocarcinoma," *Cancer Research*, vol. 69, no. 23, pp. 9073–9082, 2009.
- [15] S. Toyokuni, "Mechanisms of asbestos-induced carcinogenesis," *Nagoya Journal of Medical Science*, vol. 71, no. 1-2, pp. 1–10, 2009.
- [16] S. Toyokuni, "Role of iron in carcinogenesis: cancer as a ferrotoxic disease," *Cancer Science*, vol. 100, no. 1, pp. 9–16, 2009.
- [17] Q. Hu, S. Akatsuka, Y. Yamashita et al., "Homozygous deletion of CDKN2A/2B is a hallmark of iron-induced high-grade rat mesothelioma," *Laboratory Investigation*, vol. 90, no. 3, pp. 360–373, 2010.
- [18] S. Toyokuni, "Iron as a target of chemoprevention for longevity in humans," *Free Radical Research*. In press.
- [19] H. Nagai and S. Toyokuni, "Biopersistent fiber-induced inflammation and carcinogenesis: lessons learned from asbestos toward safety of fibrous nanomaterials," *Archives of Biochemistry and Biophysics*, vol. 502, no. 1, pp. 1–7, 2010.
- [20] J. E. Craighead, A. I. Gibbs, and F. Pooley, "Mineralogy of asbestos," in *Asbestos and Its Diseases*, J. E. Craighead and A. R. Gibbs, Eds., pp. 23–38, Oxford University Press, New York, NY, USA, 2008.
- [21] V. L. Roggli and P. Coin, "Mineralogy of asbestos," in *Pathology of Asbestos-Associated Diseases*, V. L. Roggli, T. D. Oury, and T. A. Sporn, Eds., pp. 1–16, Springer Science and Business Media, New York, NY, USA, 2nd edition, 2004.

- [22] C. L. Uber and R. A. McReynolds, "Immunotoxicology of silica," *Critical Reviews in Toxicology*, vol. 10, no. 4, pp. 303–319, 1982.
- [23] K. Steenland and D. F. Goldsmith, "Silica exposure and autoimmune diseases," *American Journal of Industrial Medicine*, vol. 28, no. 5, pp. 603–608, 1995.
- [24] H. Reyes, E. A. Ojo-Amaize, and J. B. Peter, "Silicates, silicones and autoimmunity," *Israel Journal of Medical Sciences*, vol. 33, no. 4, pp. 239–242, 1997.
- [25] D. D'Cruz, "Autoimmune diseases associated with drugs, chemicals and environmental factors," *Toxicology Letters*, vol. 112–113, pp. 421–432, 2000.
- [26] M. D. Mayes, "Epidemiologic studies of environmental agents and systemic autoimmune diseases," *Environmental Health Perspectives*, vol. 107, no. 5, pp. 743–748, 1999.
- [27] J. Schreiber, D. Koschel, J. Kekow, N. Waldburg, A. Goette, and R. Merget, "Rheumatoid pneumoconiosis (Caplan's syndrome)," *European Journal of Internal Medicine*, vol. 21, no. 3, pp. 168–172, 2010.
- [28] T. Otsuki, Y. Miura, Y. Nishimura et al., "Alterations of Fas and Fas-related molecules in patients with silicosis," *Experimental Biology and Medicine*, vol. 231, no. 5, pp. 522–533, 2006.
- [29] T. Otsuki, M. Maeda, S. Murakami et al., "Immunological effects of silica and asbestos," *Cellular and Molecular Immunology*, vol. 4, no. 4, pp. 261–268, 2007.
- [30] P. Wu, Y. Miura, F. Hyodoh et al., "Reduced function of CD4+25+ regulatory T cell fraction in silicosis patients," *International Journal of Immunopathology and Pharmacology*, vol. 19, no. 2, pp. 357–368, 2006.
- [31] T. Otsuki, H. Hayashi, Y. Nishimura et al., "Dysregulation of autoimmunity caused by silica exposure and alteration of Fas-mediated apoptosis in T lymphocytes derived from silicosis patients," *International Journal of Immunopathology and Pharmacology*, vol. 24, supplement 1, pp. 11S–16S, 2011.
- [32] H. Hayashi, Y. Miura, M. Maeda et al., "Reductive alteration of the regulatory function of the CD4+CD25+ T cell fraction in silicosis patients," *International Journal of Immunopathology and Pharmacology*, vol. 23, no. 4, pp. 1099–1109, 2010.
- [33] R. Rolston and D. Oury, "Other neoplasia," in *Pathology of Asbestos-Associated Diseases*, V. L. Roggli, T. D. Oury, and T. A. Sporn, Eds., pp. 217–230, Springer Science and Business Media, New York, NY, USA, 2nd edition, 2004.
- [34] R. A. Lemen, "Epidemiology of asbestos-related diseases and the knowledge that led to what is known today," in *Asbestos, Risk Assessment, Epidemiology, and Health Effects*, R. F. Dodson and S. P. Hammared, Eds., pp. 201–308, CRC Press, Taylor and Francis Group, Boca Raton, Fla, USA, 2006.
- [35] J. E. Craughea, "Nonthoracic cancers possibly resulting from asbestos exposure," in *Asbestos and Its Diseases*, J. E. Craighead and A. R. Gibbs, Eds., pp. 230–252, Oxford University Press, New York, NY, USA, 2008.
- [36] J. C. Barrett, P. W. Lamb, and R. W. Wiseman, "Multiple mechanisms for the carcinogenic effects of asbestos and other mineral fibers," *Environmental Health Perspectives*, vol. 81, pp. 81–89, 1989.
- [37] J. M. Davis, "Mineral fibre carcinogenesis: experimental data relating to the importance of fibre type, size, deposition, dissolution and migration," *IARC Scientific Publications*, no. 90, pp. 33–45, 1989.
- [38] I. Miyoshi, I. Kubonishi, S. Yoshimoto, and Y. Shiraishi, "A T-cell line derived from normal human cord-leukocytes by co-culturing with human leukemic T-cells," *Gann*, vol. 72, no. 6, pp. 978–981, 1981.
- [39] I. Miyoshi, I. Kubonishi, S. Yoshimoto et al., "Type C virus particles in a cord T-cell line derived by co-cultivating normal human cord leukocytes and human leukaemic T cells," *Nature*, vol. 294, no. 5843, pp. 770–771, 1981.
- [40] F. Hyodoh, A. Takata-Tomokuni, Y. Miura et al., "Inhibitory effects of anti-oxidants on apoptosis of a human polyclonal T-cell line, MT-2, induced by an asbestos, chrysotile-A," *Scandinavian Journal of Immunology*, vol. 61, no. 5, pp. 442–448, 2005.
- [41] G. Liu, R. Beri, A. Mueller, and D. W. Kamp, "Molecular mechanisms of asbestos-induced lung epithelial cell apoptosis," *Chemico-Biological Interactions*, vol. 188, no. 2, pp. 309–318, 2010.
- [42] B. Fubini and A. Hubbard, "Reactive oxygen species (ROS) and reactive nitrogen species (RNS) generation by silica in inflammation and fibrosis," *Free Radical Biology and Medicine*, vol. 34, no. 12, pp. 1507–1516, 2003.
- [43] D. Upadhyay and D. W. Kamp, "Asbestos-induced pulmonary toxicity: role of DNA damage and apoptosis," *Experimental Biology and Medicine*, vol. 228, no. 6, pp. 650–659, 2003.
- [44] A. Shukla, M. Gulumian, T. K. Hei, D. Kamp, Q. Rahman, and B. T. Mossman, "Multiple roles of oxidants in the pathogenesis of asbestos-induced diseases," *Free Radical Biology and Medicine*, vol. 34, no. 9, pp. 1117–1129, 2003.
- [45] Y. Miura, Y. Nishimura, H. Katsuyama et al., "Involvement of IL-10 and Bcl-2 in resistance against an asbestos-induced apoptosis of T cells," *Apoptosis*, vol. 11, no. 10, pp. 1825–1835, 2006.
- [46] Y. Miura, Y. Nishimura, M. Maeda et al., "Immunological alterations found in mesothelioma patients and supporting experimental evidence," *Environmental Health and Preventive Medicine*, vol. 13, no. 2, pp. 55–59, 2008.
- [47] S. Murakami, Y. Nishimura, M. Maeda et al., "Cytokine alteration and speculated immunological pathophysiology in silicosis and asbestos-related diseases," *Environmental Health and Preventive Medicine*, vol. 14, no. 4, pp. 216–222, 2009.
- [48] M. Maeda, Y. Nishimura, N. Kumagai et al., "Dysregulation of the immune system caused by silica and asbestos," *Journal of Immunotoxicology*, vol. 7, no. 4, pp. 268–278, 2010.
- [49] J. Wenzel and T. Tüting, "An IFN-associated cytotoxic cellular immune response against viral, self-, or tumor antigens is a common pathogenetic feature in "interface dermatitis"," *Journal of Investigative Dermatology*, vol. 128, no. 10, pp. 2392–2402, 2008.
- [50] R. M. Strieter, M. D. Burdick, J. Mestas, B. Gomperts, M. P. Keane, and J. A. Belperio, "Cancer CXC chemokine networks and tumour angiogenesis," *European Journal of Cancer*, vol. 42, no. 6, pp. 768–778, 2006.
- [51] M. Maeda, Y. Nishimura, H. Hayashi et al., "Reduction of CXCR3 in an in vitro model of continuous asbestos exposure on a human T-cell line, MT-2," *American Journal of Respiratory Cell and Molecular Biology*. In press.
- [52] M. Maeda, Y. Nishimura, H. Hayashi et al., "Decreased CXCR3 expression in CD4+ T cells exposed to asbestos or derived from asbestos-exposed patients," *American Journal of Respiratory Cell and Molecular Biology*. In press.
- [53] Y. Nishimura, Y. Miura, M. Maeda et al., "Impairment in cytotoxicity and expression of NK cell-activating receptors on human NK cells following exposure to asbestos fibers," *International Journal of Immunopathology and Pharmacology*, vol. 22, no. 3, pp. 579–590, 2009.
- [54] Y. Nishimura, M. Maeda, N. Kumagai, H. Hayashi, Y. Miura, and T. Otsuki, "Decrease in phosphorylation of ERK following decreased expression of NK cell-activating receptors in human

NK cell line exposed to asbestos," *International Journal of Immunopathology and Pharmacology*, vol. 22, no. 4, pp. 879–888, 2009.

- [55] Y. Nishimura, N. Kumagai, M. Maeda et al., "Suppressive effect of asbestos on cytotoxicity of human NK cells," *International Journal of Immunopathology and Pharmacology*, vol. 24, supplement 1, pp. 5S–10S, 2011.

Decreased CXCR3 Expression in CD4⁺ T Cells Exposed to Asbestos or Derived from Asbestos-Exposed Patients

Megumi Maeda^{1,2}, Yasumitsu Nishimura¹, Hiroaki Hayashi¹, Naoko Kumagai¹, Ying Chen¹, Shuko Murakami¹, Yoshie Miura³, Jun-ichi Hiratsuka⁴, Takumi Kishimoto⁵, and Takemi Otsuki¹

¹Department of Hygiene, Kawasaki Medical School, Okayama, Japan; ²Division of Bioscience, Department of Biofunctional Chemistry, Graduate School of Natural Science and Technology, Okayama University, Okayama, Japan; ³Division of Molecular and Clinical Genetics, Department of Molecular Genetics, Medical Institute of Bioregulation, Kyushu University, Fukuoka, Japan; ⁴Department of Radiation Oncology, Kawasaki Medical School, Okayama, Japan; and ⁵Okayama Rosai Hospital, Okayama, Japan

Asbestos causes malignant tumors such as lung cancer and malignant mesothelioma (MM). To determine whether asbestos exposure causes reduction of antitumor immunity, we established an *in vitro* T-cell line model of low-dose and continuous exposure to asbestos using a human adult T-cell leukemia virus-1 immortalized human polyclonal T-cell line, MT-2, and revealed that MT-2 cells exposed continuously to asbestos showed resistance to asbestos-induced apoptosis. In addition, the cells presented reduction of surface CXCR3 chemokine receptor expression and IFN- γ production. In this study, to confirm that these findings are suitable for clinical translation, surface CXCR3 and IFN- γ expression were analyzed using freshly isolated human CD4⁺ T cells derived from healthy donors and patients with pleural plaque (PP) or MM. The results revealed that CXCR3 and IFN- γ expression in the *ex vivo* model were reduced in some cases. Additionally, CXCR3 expression in CD4⁺ T cells from PPs and MMs was significantly reduced compared with that from healthy donors, and CD4⁺ T cells from patients with MMs exhibited a marked reduction in IFN- γ mRNA levels after stimulation *in vitro*. Moreover, CD4⁺CXCR3⁺ T cells in lymphocytes from MMs showed a tendency for an inverse correlation with its ligand CXCL10/IP10 in plasma. These findings show reduction of antitumor immune function in asbestos-exposed patients and indicate that CXCR3, IFN- γ , and CXCL10/IP10 may be candidates to detect and monitor disease status.

Keywords: asbestos; pleural plaque; malignant mesothelioma; CXCR3; IFN- γ

The cellular and molecular biologic effects of asbestos fibers on alveolar epithelial cells and pleural mesothelial cells have been investigated using various animal models and culture cells (1–5). Studies of fibrogenesis have shown that the initial recognition of asbestos by alveolar macrophages (AMs) is important, and AM-releasing cytokines, such as TNF- α , TGF- β , and IL-1 β and IL-8, play a critical role in activating mesenchymal cells to induce the proliferation of collagen fibers (5, 6). Moreover, recent advances resulting from cellular and molecular investigations regarding activation of the nucleotide

CLINICAL RELEVANCE

In this study, we show that chrysotile asbestos has the potential to reduce chemokine receptor CXCR3 expression in human peripheral CD4⁺ T cells using an *ex vivo* model, and CXCR3 expression is decreased in peripheral CD4⁺ T cells from patients with asbestos-related diseases, such as pleural plaque or malignant mesothelioma. Our findings suggest that antitumor immunity might not function normally in patients with asbestos-related disease because the low expression of CXCR3 induces depressed chemotaxis and indicate that CXCR3 may be effective tool for the prediction of impaired antitumor immune status in these patients.

oligomerization domain-like receptor family, the pryin domain-containing 3 (NLRP3)-inflammasome caused by asbestos, are leading to a better understanding of this recognition (7–9). Reactive oxygen species and reactive nitrogen species play important roles in carcinogenesis (1–6), in which substances that damage DNA are generated through the activity of iron contained in the asbestos. Alveolar and mesothelial cells exposed to asbestos then develop apoptosis with activation of a mitochondrial pathway and a B-cell lymphoma 2 (Bcl-2)-associated x protein (Bax) dominant balance of the Bax/Bcl-2 complex, cytosomal release of cytochrome-c from mitochondria, and activation of caspase 9 and 3 (1–6).

The above findings are based on studies using animal models and cell culture experiments in which exposure conditions were temporary and involved high doses. In human asbestos-related diseases, the latency is quite long and may range from 10 to 30 years for fibrogenic changes known as asbestosis and around 40 years for malignant mesothelioma (MM) (10–12).

To investigate the immunological effects of asbestos on human immunocompetent cells, we constructed an *in vitro* experimental T-cell line model of continuous asbestos exposure using a human adult T-cell leukemia virus type 1-immortalized human polyclonal T-cell line (MT-2) because this cell line is not tumor-cell derived and is reported to have a normal karyotype (13–16). In our experiments, chrysotile was used as the asbestos because it may be important to compare this substance with silica (SiO₂), which causes silicosis and alteration of autoimmunity (17, 18). In addition, the dose of chrysotile used in industries around the world is much higher than that of other fibers, and carcinogenicity is considered much lower for chrysotile (10–12).

As we reported recently, our experimental models, which use the MT-2 original cell line and six sublines exposed to chrysotile independently and continuously, revealed that all of the sublines exhibited reduction of surface expression of CXCR3 chemokine receptor 3 (CXCR3) and production of IFN- γ (19). It is well known that IFN- γ is important in antitumor immunity

(Received in original form October 21, 2010 and in final form February 8, 2011)

This study was supported in part by Special Coordination Funds for Promoting Science and Technology (H18-1-3-3-1, Comprehensive approach on asbestos-related diseases); grants from the Ministry of Education, Culture, Sports, Science and Technology of Japan (18390186, 19659153, 19790411, 20390178 and 22700933); and Kawasaki Medical School Project Grants (18-209T, 19-205Y, and 20-210O).

Correspondence and requests for reprints should be addressed to Megumi Maeda, Ph.D., Department of Biofunctional Chemistry, Graduate School of Natural Science and Technology, Okayama University, 1-1-1 Tsushima-naka Kita-ku, Okayama 700-8530, Japan. E-mail: mmaeda@cc.okayama-u.ac.jp

This article has an online supplement, which is accessible from this issue's table of contents at www.atsjournals.org.

Am J Respir Cell Mol Biol Vol 45, pp 795–803, 2011

Originally Published in Press as DOI: 10.1165/rcmb.2010-0435OC on February 25, 2011
Internet address: www.atsjournals.org

and antiviral immunity (20–24). In addition, CXCR3 is thought to be involved in antitumor immune function. Some reports have investigated the state of chemokines surrounding mesothelioma tumor cells (25–28). It has been found that the ligand of CXCR3, C-X-C motif chemokine 10 (CXCL10)/IFN- γ -induced protein 10 kD (IP10), is present in pleural fluid collected from patients with MM and that human mesothelioma tumors are abundantly infiltrated with CD4⁺ T cells (29). Furthermore, CXCL10/IP10 mRNA expression is significantly higher in MM compared with normal mesothelial cell lines and pleural mesothelium (30). CXCL10/IP10 inhibits tumor growth and metastasis through a decrease in tumor-associated angiogenesis and recruits Th1/effector T cells by binding the chemokine receptor CXCR3 (20, 31, 32). Therefore, it is important to regulate lymphocyte transport mediated by chemokine receptors to enhance antitumor immune responses because CXCR3-positive Th1/effector CD4⁺ T cells are recruited to CXCL10/IP10-enriched tumor sites, where IFN- γ is secreted and subsequent suppression of tumor growth is induced (33).

We examined CXCR3 and IFN- γ expression in CD4⁺ T cells co-cultured with chrysotile and showed the potential of chrysotile to inhibit CXCR3 and IFN- γ expression. In addition, we investigated CXCR3 and IFN- γ expression in CD4⁺ T cells from healthy donors (HDs) and asbestos-exposed patients with pleural plaque (PP) or MM and found low-level expression of CXCR3 in CD4⁺ T cells from patients with PPs and MMs and decreased mRNA levels of IFN- γ in cultured CD4⁺ T cells from patients with MMs. These findings highlight parameters that are effective in predicting the risk of impaired antitumor immune status in patients with PP and MM.

MATERIALS AND METHODS

Ex Vivo Activation and Exposure to Asbestos on Human CD4⁺ T Cells

After receiving informed consent, blood samples were obtained from six HDs (mean age, 34.8 \pm 9.7 yr; range, 25–50 yr; three men and three women), and peripheral blood mononuclear cells (PBMCs) were isolated using the Ficoll-Paque method. CD4⁺ T cells were isolated by positive selection using anti-CD4-coated beads (Miltenyi Biotec GmbH, Bergisch Gladbach, Germany) in accordance with the manufacturer's instructions. Freshly isolated CD4⁺ T cells (2×10^5 cells/well) in 96-well, U-bottomed plates were stimulated with anti-CD3 (clone UCHT1) and anti-CD28 (clone CD28.2) antibodies (Beckman Coulter, Inc., Fullerton, CA) at 2 μ g/ml and cultured in RPMI 1640 medium containing 10% FBS and recombinant IL-2 (rIL-2) (10 ng/ml) (PeproTech Inc., Rocky Hill, NJ). After 3 days, activated CD4⁺ T cells were transferred to 24-well plates at 1×10^6 cells/well and expanded in medium supplemented with 10 ng/ml rIL-2 for 1 week and further cultured in the absence or presence of 2 to 10 μ g/cm² of chrysotile-B. Chrysotile-B was kindly provided by the Department of Occupational Health, National Institute for Occupational Health, South Africa (34). Before analysis, cultured CD4⁺ T cells were released from chrysotile-B by the Ficoll-Paque method and cultured for 3 days. For intracellular staining, cultured CD4⁺ T cells (1×10^6 cells/ml) were stimulated with 5 ng/ml phorbol 12-myristate 13-acetate and 250 ng/ml ionomycin for 6 hours in the presence of monesin.

Flow Cytometry

Surface antigens were stained as reported previously (19) with the following antibodies: anti-CD4 (clone RPA-T4), CXCR3 (clone 1C6), and CC chemokine receptor 5 (CCR5) (clone 2D7) (BD Biosciences Pharmingen, San Diego, CA). Intracellular staining was performed using a Cytofix/Cytoperm kit (BD Biosciences) in accordance with the manufacturer's instructions. Cells were stained with anti-IFN- γ (clone 4S.B3) antibody (BD Biosciences Pharmingen) for 30 minutes at 4°C and analyzed on a flow cytometer (FACSCalibur; BD Biosciences, Franklin Lakes, NJ).

Patients and Isolation of PBMCs

Blood from a different series of 19 HDs (mean age, 42.3 \pm 12.8 yr; range, 29–70 yr; 9 men and 10 women), 13 patients with PP (mean age, 71.5 \pm 5.3 yr; range, 58–79 yr; 11 men and 2 women), and eight patients with MM (mean age, 67.4 \pm 8.5 yr; range, 59–79 yr; seven men and one woman) was collected in citrate phosphate dextrose. Plasma was collected from supernatants after centrifugation at 3,000 rpm for 5 minutes. PBMCs were isolated as described above and subjected to FACS analysis. For analysis of mRNA expression and cytokine production, CD4⁺ T cells were isolated as described above. Freshly isolated CD4⁺ T cells were cultured in RPMI 1640 medium containing 10% FBS at 2×10^4 cells/100 μ l in 96-well, U-bottomed plates and stimulated with anti-CD3 and anti-CD28 antibodies at 2 μ g/ml. After 5 days, the supernatants and cells were collected. Informed consent was obtained from all donors, and the project and procedures used were approved by the Institutional Ethics Committees of Kawasaki Medical School and Okayama Rosai Hospital.

RESULTS

Changes of CXCR3 Cell Surface Expression in CD4⁺ T Cells Cultured with Chrysotile-B

As reported recently, our *in vitro* T-cell line model of low-level and continuous exposure to chrysotile asbestos showed reduction of Th1-type cell surface expression of CXCR3 (19). Therefore, we tried to determine whether freshly isolated human peripheral CD4⁺ T cells show a similar alteration *ex vivo* when proliferation is maintained by IL-2-containing medium in the presence of chrysotile-B. After 40 days of co-culture, cell surface CXCR3 expression decreased in a dose-dependent manner (2–10 μ g/cm²) (Figure 1A). Thus, we examined cell surface expression of CXCR3 and CCR5 in CD4⁺ T cells derived from six HDs because both receptors are preferentially expressed in Th1/effector T cells. The expression of CXCR3 was significantly reduced after exposure to 10 μ g/cm² of chrysotile for 28 days, although this difference seemed to depend on one case in which the expression decreased remarkably (Figure 2, upper panel). Even if the culture conditions for the CD4⁺ T cells was limited to a period of around 4 weeks, four of six HDs showed a decrease of CXCR3 expression to various degrees, and it might be concluded that asbestos exposure potentiates reduction of CXCR3 expression in CD4⁺ T cells. On the other hand, the expression of CCR5 varied among all HDs, and there were no significant changes after 7 and 28 days of co-culture with chrysotile-B (Figure 1B, lower panel), as shown previously by the cell line model (19). These results indicated that CXCR3 expression might be specifically reduced by asbestos exposure.

IFN- γ Expression under *Ex Vivo* Exposure Conditions

We also showed previously that IFN- γ production in the T-cell line model was decreased by chronic exposure to chrysotile (19). Thus, we examined intracellular expression of IFN- γ in CD4⁺ T cells exposed to chrysotile-B (10 μ g/cm²) from three healthy subjects under the same *ex vivo* exposure conditions used for the analysis of surface CXCR3 expression. After 28 days of co-culture, mRNA expression of IFN- γ was reduced in all three cases, although there were no significant differences ($P = 0.3226$) (Figure 2A). In addition, intracellular staining showed that IFN- γ -positive cells tended to be reduced in CD4⁺ T cells from all three subjects when cultured with chrysotile-B ($P = 0.0511$) (Figure 2B).

These findings from the *ex vivo* study indicated that chronic exposure to asbestos reduced CXCR3 and IFN- γ expression in CD4⁺ T cells, although the reductions were not statistically significant.




Figure 1. Chronic exposure to chrysotile reduces expression of CXCR3 in IL-2-cultured human CD4⁺ T cells. Peripheral CD4⁺ T cells from healthy donors were stimulated with anti-CD3/CD28 antibodies and maintained with IL-2-containing medium. Ten days later, exposure to chrysotile-B (CB) was initiated. (A) Histogram shows representative example of CXCR3 expression on CD4⁺ T cells after 40 days of exposure to various doses of CB. (B) Graphs show expression of CXCR3 (upper panel) and CCR5 (lower panel) after 7 and 28 days of culture in the absence or presence of 10 µg/cm² CB. Data are presented for six healthy subjects. The *P* value was obtained using the Wilcoxon signed-rank test. **P* < 0.05.

Expression of Cell Surface CXCR3 in Human CD4⁺ T Cells from Patients with Asbestos-Related Disease

To investigate whether peripheral CD4⁺ T cells in people exposed to asbestos also show reduction of CXCR3, we analyzed cell surface CXCR3 and CCR5 expression in peripheral CD4⁺ T cells from patients with asbestos-related PP and MM by flow cytometry (Figure 3A). Given that the population of lymphocytes in MMs was significantly lower than that in the other groups (Figure 3B, left panel), gated lymphocytes were analyzed for CXCR3 and CCR5 expression in CD4⁺ T cells. The percentage of CD4⁺ T cells in lymphocytes revealed no differences among the three groups (Figure 3B, right panel).

The percentage of CD4⁺CXCR3⁺ T cells in lymphocytes from PPs and MMs were significantly lower than that of HDs, and MMs showed the lowest percentage compared with the other groups (Figure 3C, left panel). In contrast, the percentage of CD4⁺CCR5⁺ T cells in lymphocytes showed no differences (Figure 3C, right panel). Furthermore, the expression of CXCR3 in CD4⁺ T cells from PPs and MMs was significantly lower than

that of HDs (Figure 3D, left panel), although CCR5 expression in CD4⁺ T cells showed no differences (Figure 3D, right panel).

To confirm whether there is an aging effect concerning decreased expression of CXCR3, 19 HDs were divided into two groups: younger (mean age, 34.9 ± 5.8 yr; range, 25–45 yr; *n* = 13) and older (mean age, 58.3 ± 7.6 yr; range, 51–70 yr; *n* = 6). The difference between these groups regarding age was significant (Figure 4A). However, our findings indicated that there were no significant differences in the percentage of CD4⁺CXCR3⁺ T cells in lymphocytes (Figure 4B, upper panel) and CXCR3 expression in CD4⁺ T cells (Figure 4B, lower panel) between the two groups. Thus, these results confirmed that aging is not associated with a reduction of CXCR3 expression and that decreased expression of CXCR3 in peripheral CD4⁺ T cells occurs in people exposed to asbestos.

IFN-γ mRNA Levels in Stimulated Peripheral CD4⁺ T Cells from Patients with Asbestos-Related Diseases

Real-time RT-PCR was used to detect the mRNA levels of IFN-γ in stimulated peripheral CD4⁺ T cells from 13 HDs, 13




Figure 2. IFN-γ production in IL-2-cultured human CD4⁺ T cells from patients showing reduced CXCR3 expression by exposure to chrysotile. CD4⁺ T cells from three subjects were activated with anti-CD3/CD28 antibodies and cultured with IL-2-containing medium in the absence or presence of 10 µg/cm² CB for 30 days. (A) IFN-γ mRNA in subjects who showed decreased expression of surface CXCR3 in CD4⁺ T cells by CB exposure was measured by real-time RT-PCR. (B) Histogram shows representative staining for intracellular IFN-γ. Graph indicates the percentage of IFN-γ-positive cells. Data are shown for three subjects. The *P* value was obtained using the paired *t* test. **P* < 0.05.




Figure 3. Peripheral CD4⁺ T cells in patients with pleural plaque (PP) or malignant mesothelioma (MM) show decreased CXCR3 expression. (A) A forward/side scatter dot plot was used to gate lymphocytes, and CXCR3 and CCR5 expressions in peripheral CD4⁺ T cells from HDs ($n = 19$), PPs ($n = 13$), and MMs ($n = 8$) were analyzed by FACS. Representative FACS profiles are indicated. Numbers in each dot plot indicate percentage of lymphocytes (top), CD4⁺CXCR3⁺ T cells (middle), and CD4⁺CCR5⁺ T cells (bottom). (B) The percentage of gated lymphocytes in total cells (left) and CD4⁺ T cells in gated lymphocytes (right). (C) The percentage of CD4⁺CXCR3⁺ T cells (left) and CD4⁺CCR5⁺ T cells (right) in gated lymphocytes. (D) The percentage of CXCR3-positive (left) and CCR5-positive (right) cells in gated CD4⁺ T cells. The box plot shows 10% (lower whisker), 25% (lower hinge of box), the median (line in the box), 75% (upper hinge of box), 90% (upper whisker), and the extreme points (black dots). The P value was obtained using the Student-Newman-Keuls test. * $P < 0.01$; ** $P < 0.05$.

PPs, and 7 MMs. mRNA expression levels of IFN- γ relative to GAPDH were significantly decreased only in MMs compared with HDs and PPs (Figure 5A). Additionally, the Spearman's rank correlation test showed that there was no significant correlation between levels of CXCR3 expression in peripheral CD4⁺ T cells and IFN- γ mRNA levels of stimulated CD4⁺ T cells in all three groups (Figures 5B–5D), although HDs showed a tendency for a positive correlation ($P = 0.1295$) and PPs showed a tendency for a negative correlation ($P = 0.0502$). Unlike suppressed CXCR3 expression, these results suggested that decreased mRNA expression of IFN- γ was dependent on the occurrence of mesothelioma and was not necessarily associated with asbestos exposure.

IFN- γ and Production of other Cytokines in Human CD4⁺ T Cells from Patients with Asbestos-Related Disease

The cytometric bead array assay was performed to measure the concentration of Th1/Th2/Th17 cytokine (IL-2, IL-4, IL-6, IL-10, TNF- α , IFN- γ , and IL-17A) in the supernatants from cultured peripheral CD4⁺ T cells. There were no significant differences in IFN- γ production from cultured peripheral CD4⁺ T cells (Figure 6), although there were statistically significant differences in IFN- γ mRNA expression (Figure 5A). Thus, the expected result for IFN- γ -producing activity was not obtained under these conditions. The results for other cytokines showed that the IL-6 concen-

tration in PPs and MMs was significantly higher than that of HDs (Figure 6). Although the IL-4 concentration in MMs was significantly higher than that of HDs, the level might be too low for the cytokine to exert its physiological function (Figure 6).

Correlation between the Percentage of CD4⁺CXCR3⁺ T Cells in Lymphocytes and Levels of Plasma CXCL10/IP10

Because the ligand for CXCR3, CXCL10/IP10, is secreted not only from monocytes, endothelial cells, and fibroblasts (20, 31, 32) but also from mesothelial cells and transformed (mesothelioma) cells (30), the plasma levels of CXCL10/IP10 in HDs, PPs, and MMs were measured by CBA assay. The levels of plasma CXCL10/IP10 had no correlation with CXCR3 expression in CD4⁺ T cells (data not shown). Therefore, the correlation between the percentage of CD4⁺CXCR3⁺ T cells in lymphocytes and plasma chemokine levels was determined using Spearman's rank correlation test. The plasma concentration of CXCL10/IP10 in PPs or MMs tended to be higher when compared with that of HDs, although there were no significant differences (Figure 7A). Therefore, there was no significant correlation between the percentage of CD4⁺CXCR3⁺ T cells and CXCL10/IP10 concentration in all three groups (Figures 7B–7D), although MMs showed a tendency for an inverse correlation ($P = 0.081$) in comparison with HDs ($P = 0.927$) and PPs ($P = 0.578$).




Figure 4. Age of subjects does not affect CXCR3 expression in $CD4^+$ T cells. (A) Nineteen HDs were divided into younger (< 50 yr old; $n = 13$) and older (≥ 50 yr old; $n = 6$) groups. (B) The percentage of $CD4^+CXCR3^+$ T cells in gated lymphocytes (*top*) and CXCR3-positive cells in gated $CD4^+$ T cells (*bottom*). The box plot is presented as described in Figure 3. The *P* value was obtained using the Student-Newman-Keuls test. * $P < 0.01$.

DISCUSSION

There are few publications regarding the effects of asbestos on the general immune system, particularly antitumor immunity. However, the mechanisms by which local immune cells, such as alveolar macrophages, recognize asbestos fibers have been investigated, particularly the discovery of the NLRP3-inflammasome and its ability to recognize and process asbestos fibers and activate IL-1 β and caspase 1 (7–9). Asbestos is a mineral silicate that contains magnesium, iron, and calcium. Patients exposed to silica particles may develop a disease

known as silicosis, which is often complicated with autoimmune diseases such as rheumatoid arthritis (known as Caplan’s syndrome), systemic sclerosis, and anti-neutrophil cytoplasmic autoantibody-related vasculitis/nephritis (35–37). However, the most important complication resulting in asbestos-exposed patients is the occurrence of cancer, such as lung cancer and MM (10–12). In particular, MM is known to be caused by low-level and long-term exposure to asbestos (10–12).

Therefore, asbestos fibers may affect the human immune system, particularly antitumor immunity. In fact, we have been reporting impaired NK cell function with reduced expression of




Figure 5. Peripheral $CD4^+$ T cells in MMs show reduced IFN- γ mRNA expression. Freshly isolated peripheral $CD4^+$ T cells from HDs ($n = 13$), PPs ($n = 13$), and MMs ($n = 7$) were stimulated with anti-CD3/CD28 antibodies for 5 days. (A) Total RNA was isolated from $CD4^+$ T cells stimulated with anti-CD3/CD28 antibodies for 5 days, and relative mRNA expression of IFN- γ was estimated by real-time RT-PCR. The box plot is presented as described in Figure 3. The *P* value was obtained using the Mann-Whitney U test. * $P < 0.01$; ** $P < 0.05$. (B) Correlations between mRNA expression of IFN- γ and cell surface expression of CXCR3 in $CD4^+$ T cells in HDs (B), PPs (C), and MMs (D) were analyzed using Spearman’s rank correlation test.




Figure 6. Peripheral CD4⁺ T cells in patients with PP or MM show increased IL-6 production. Freshly isolated peripheral CD4⁺ T cells from HDs ($n = 13$), PPs ($n = 13$), and MMs ($n = 7$) were stimulated with anti-CD3/CD28 antibodies for 5 days, and the concentration of various cytokines indicated in the graphs were measured using a CBA Human Th1/Th2/Th17 Cytokine Kit. The box plot is presented as described in Figure 3. The P value was obtained using the Mann-Whitney U test. $^{***}P < 0.05$.

the NK-activating receptor NKp46 (38, 39). In addition, we attempted to establish an *in vitro* T-cell model of continuous and low-level exposure to asbestos and reported that the MT-2 subline exhibited resistance to asbestos-induced apoptosis with activating Src-family kinases, up-regulation of IL-10, phosphorylation of STAT3, and subsequent up-regulation of Bcl-2 (15, 16). Furthermore, six independent sublines of MT-2 original cells continuously exposed to chrysotile showed reduced Th1-related IFN- γ production, CXCL10/IP10 production, and cell surface expression of the CXCL10/IP10 receptor CXCR3 (19).

If these findings obtained using the *in vitro* cell line model are compatible with those pertaining to asbestos-exposed patients, such as those possessing PPs and MMs, these molecules may represent clinical markers for asbestos exposure, disease progression, and even therapeutic targets. Thus, we tried to confirm alteration of surface CXCR3 expression and IFN- γ production using freshly isolated human CD4⁺ T cells derived from HDs and patients with PP and MM. To use the *ex vivo* model of low-dose and long-term exposure to chrysotile asbestos, these CD4⁺ T cells should be cultured for several months. However, because IL-2-dependent proliferation of CD4⁺ T cells is not permanent, our experiment was limited to culture for several weeks in the presence of IL-2 after stimulation of anti-CD3/CD28 antibodies. The results showed that four of six cases exhibited reduced CXCR3 expression within the limitation of the *ex vivo* model. Moreover, intracellular IFN- γ expression was clearly

reduced in this *ex vivo* experimental situation, although the number of applied cases was small. These findings encouraged us to examine CXCR3 expression and IFN- γ production using the samples derived from asbestos-exposed patients, such as those possessing PPs and MMs.

Analysis of the case samples revealed that the expression of CXCR3 in CD4⁺ T cells from PPs and MMs was decreased when compared with that of HDs, even though another Th1-type chemokine receptor, CCR5, did not differ between the three groups. Because CXCR3 is implicated in the migration of Th1/effector T cells, decreased CXCR3 expression in CD4⁺ T cells by asbestos exposure may reduce normal Th1/effector T-cell recruitment to tumor sites, resulting in impairment of anti-tumor immune functions. Meanwhile, asbestos is accumulated in several lymph nodes (40–42). Thus, it seems that CXCR3 expression in CD4⁺ T cells is inhibited by contact with asbestos in the lymph nodes of patients with PP and MM. In addition, CD4⁺CXCR3⁺ T cells in lymphocytes were significantly reduced in MMs rather than PPs because the population of lymphocytes was significantly decreased in MMs compared with PPs. This suggests that CD4⁺CXCR3⁺ T cells are gradually decreased and that they depend not only on asbestos exposure but also on disease status, such as the occurrence of mesothelioma.

IFN- γ mRNA levels in CD4⁺ T cells derived from MMs stimulated *in vitro* were lower than those of HDs or PPs, whereas there were no significant differences in IFN- γ pro-




Figure 7. Modest inverse correlation between the population size of CD4⁺CXCR3⁺ T cells and plasma CXCL10 levels in MMs. (A) Plasma concentration of CXCL10/IP10 from HDs ($n = 10$), PPs ($n = 8$), and MMs ($n = 8$). The box plot is presented as described in Figure 3. Correlations between the percentage of CD4⁺CXCR3⁺ T cells in gated lymphocytes and plasma concentration of CXCL10/IP10 from HDs (B), PPs (C), and MMs (D) were analyzed using Spearman's rank correlation test.

duction. These results suggest that CD4⁺ T cells from MMs already possess a reduced capacity to produce IFN- γ , which is important for activation of antitumor immune function due to long-term asbestos exposure and the presence of tumor. A correlation test showed that mRNA levels of IFN- γ in PPs tended to increase in stimulated CD4⁺ T cells with decreasing surface CXCR3 expression levels, although mRNA levels of IFN- γ in HDs were positively correlated with CXCR3 expression. Given that asbestos exposure is partially associated with autoimmune responses (43), asbestos exposure in PPs might change the character of CD4⁺ T cells to enhance antitumor immune function, and IFN- γ mRNA expression might not be suppressed in patients with PP who do not have MM.

Alternatively, the producible capacities of IL-6 in stimulated CD4⁺ T cells tended to be higher in PPs and MMs compared with HDs, suggesting that asbestos exposure enhances inflammatory cytokine producibility in CD4⁺ T cells. We showed previously that the plasma concentration of IL-6 in MM was higher than that of HDs and PPs (44). Given that the levels of plasma IL-6 were not high in PPs, it seems that plasma IL-6 in MMs is mainly derived from mesothelioma cells (45–47). Furthermore, IL-6 inhibits Th1-cell differentiation and promotes Th2-cell differentiation (48). Thus, plasma IL-6 might induce decreased IFN- γ mRNA expression and increased IL-4 production in stimulated CD4⁺ T cells from MMs. On the other hand, although there were no significant differences in IL-10 and IL-17A production under these experimental conditions, there are several reports regarding the relationship between IL-10/IL-17 and cancer status, which indicates that IL-10 may suppress antitumor immune function and that IL-17 may promote inflammation in cancer (49, 50). Future studies should therefore investigate the pathophysiological roles of IL-10 and IL-17 in asbestos-exposed patients such as those with PPs or MMs.

Finally, we found that the plasma concentration of CXCL10/IP10 tended to be higher for PPs and MMs than HDs. Furthermore, there tended to be an inverse correlation between the population of CD4⁺CXCR3⁺ T cells in lymphocytes and the plasma concentration of CXCL10/IP10 in MMs but not in HDs or PPs. These results indicated that antitumor immune function in MMs may be impaired after reduction of CD4⁺CXCR3⁺ T cells in lymphocytes because CXCL10/IP10 from

monocytes, endothelial cells, fibroblasts, mesothelial cells, and mesothelioma in the region of existing tethered asbestos and a cancerous lesion attracts antiinflammatory or antitumor T cells.

In conclusion, this study revealed that reduction of CXCR3 expression in human peripheral CD4⁺ T cells occurred after chronic exposure to chrysotile. This finding suggests that antitumor immunity might not function normally in patients with asbestos-related disease because the low expression of CXCR3 induces depressed chemotaxis. Additionally, it seems that IFN- γ expression is inhibited by the occurrence of mesothelioma rather than asbestos exposure, and the weak production of IFN- γ leads to suppression of antitumor activity. Further research of chemokine-mediated migration dependent on CXCR3 expression is required to determine the mechanism of immune dysfunction via chemokine receptors after exposure to asbestos and to investigate the use of the chemokine receptor CXCR3 as a key molecule in immunotherapy strategies for asbestos-related disease. In addition, these molecules may be effective tools for the detection of asbestos exposure, considering our previous findings of enhanced Bcl-2 expression in CD4⁺ T cells from MMs (16) and higher plasma concentration of IL-6, IL-10, and TGF- β (44). For example, higher plasma IL-10 and TGF- β with lower INF- γ expression may be interpreted as an enhanced regulatory T-cell activity overcoming antitumor T-cell activity in MMs.

Further studies regarding the size and function of regulatory T cells in MMs and the *in vitro* model using the cell line should be performed to better understand immunological status among asbestos-exposed patients with or without mesothelioma. Moreover, a device to evaluate total antitumor activities in these patients using the degree of reduced expression in CD4⁺ T cells of an NK cell-activating surface receptor such as NKp46, including CXCR3 and IFN- γ expression, may assist in the diagnosis of MMs and reveal good prognostic factors for these patients.

Author Disclosure: None of the authors has a financial relationship with a commercial entity that has an interest in the subject of this manuscript.

Acknowledgments: The authors thank Ms. Tamayo Hatayama, Minako Kato, Naomi Miyahara, Shoko Yamamoto, Misao Kuroki, Keiko Kimura, Yoshiko Yamashita, and Tomoko Sueishi for technical help.

References

- Upadhyay D, Kamp DW. Asbestos-induced pulmonary toxicity: role of DNA damage and apoptosis. *Exp Biol Med (Maywood)* 2003;228:650–659.
- Panduri V, Surapureddi S, Soberanes S, Weitzman SA, Chandel N, Kamp DW. P53 mediates amosite asbestos-induced alveolar epithelial cell mitochondria-regulated apoptosis. *Am J Respir Cell Mol Biol* 2006;34:443–452.
- Upadhyay D, Panduri V, Kamp DW. Fibroblast growth factor-10 prevents asbestos-induced alveolar epithelial cell apoptosis by a mitogen-activated protein kinase-dependent mechanism. *Am J Respir Cell Mol Biol* 2005;32:232–238.
- Mossman BT. Introduction to serial reviews on the role of reactive oxygen and nitrogen species (ROS/RNS) in lung injury and diseases. *Free Radic Biol Med* 2003;34:1115–1116.
- Kamp DW, Panduri V, Weitzman SA, Chandel N. Asbestos-induced alveolar epithelial cell apoptosis: role of mitochondrial dysfunction caused by iron-derived free radicals. *Mol Cell Biochem* 2002;234–235:153–160.
- Jiang L, Nagai H, Ohara H, Hara S, Tachibana M, Hirano S, Shinohara Y, Kohyama N, Akatsuka S, Toyokuni S. Characteristics and modifying factors of asbestos-induced oxidative DNA damage. *Cancer Sci* 2008;99:2142–2151.
- Pettrilli V, Dostert C, Muruve DA, Tschopp J. The inflammasome: a danger sensing complex triggering innate immunity. *Curr Opin Immunol* 2007;19:615–622.
- Sutterwala FS, Ogura Y, Flavell RA. The inflammasome in pathogen recognition and inflammation. *J Leukoc Biol* 2007;82:259–264.
- Cook GP, Savic S, Wittmann M, McDermott MF. The NLRP3 inflammasome, a target for therapy in diverse disease states. *Eur J Immunol* 2010;40:631–634.
- Kannerstein M, Chung J, McCaughey E, Selikoff IJ. Pathogenic effects of asbestos. *Arch Pathol Lab Med* 1977;101:623–628.
- Kannerstein M, Chung J, McCaughey WT. Asbestos and mesothelioma: a review. *Pathol Annu* 1978;13:81–129.
- Craighead JE, Mossman BT. The pathogenesis of asbestos-associated diseases. *N Engl J Med* 1982;306:1446–1455.
- Miyoshi I, Kubonishi I, Yoshimoto S, Shiraishi Y. A T-cell line derived from normal human cord leukocytes by co-culturing with human leukemic T-cells. *Gann* 1981;72:978–981.
- Miyoshi I, Kubonishi I, Yoshimoto S, Akagi T, Ohtsuki Y, Shiraishi Y, Nagata K, Hinuma Y. Type c virus particles in a cord T-cell line derived by co-cultivating normal human cord leukocytes and human leukaemic T cells. *Nature* 1981;294:770–771.
- Hyodoh F, Takata-Tomokuni A, Miura Y, Sakaguchi H, Hatayama T, Hatada S, Katsuyama H, Matsuo Y, Otsuki T. Inhibitory effects of anti-oxidants on apoptosis of a human polyclonal T-cell line, MT-2, induced by an asbestos, chrysotile-A. *Scand J Immunol* 2005;61:442–448.
- Miura Y, Nishimura Y, Katsuyama H, Maeda M, Hayashi H, Dong M, Hyodoh F, Tomita M, Matsuo Y, Uesaka A, et al. Involvement of IL-10 and Bcl-2 in resistance against an asbestos-induced apoptosis of T cells. *Apoptosis* 2006;11:1825–1835.
- Otsuki T, Miura Y, Nishimura Y, Hyodoh F, Takata A, Kusaka M, Katsuyama H, Tomita M, Ueki A, Kishimoto T. Alterations of Fas and Fas-related molecules in patients with silicosis. *Exp Biol Med (Maywood)* 2006;231:522–533.
- Otsuki T, Maeda M, Murakami S, Hayashi H, Miura Y, Kusaka M, Nakano T, Fukuoka K, Kishimoto T, Hyodoh F, et al. Immunological effects of silica and asbestos. *Cell Mol Immunol* 2007;4:261–268.
- Maeda M, Nishimura Y, Hayashi H, Kumagai N, Chen Y, Murakami S, Miura Y, Hiratsuka JI, Kishimoto T, Otsuki T. Reduction of CXCR3 in an *in vitro* model of continuous asbestos exposure on a human T-cell line, MT-2. *Am J Respir Cell Mol Biol* (In press)
- Homey B, Muller A, Zlotnik A. Chemokines: agents for the immunotherapy of cancer? *Nat Rev Immunol* 2002;2:175–184.
- Rudge G, Barrett SP, Scott B, van Driel IR. Infiltration of a mesothelioma by IFN-gamma-producing cells and tumor rejection after depletion of regulatory T cells. *J Immunol* 2007;178:4089–4096.
- Miller CH, Maher SG, Young HA. Clinical use of interferon-gamma. *Ann N Y Acad Sci* 2009;1182:69–79.
- Rockley PF, Tying SK. Interferons alpha, beta and gamma therapy of anogenital human papillomavirus infections. *Pharmacol Ther* 1995;65:265–287.
- Welsh RM, Lin MY, Lohman BL, Varga SM, Zarozinski CC, Selin LK. Alpha beta and gamma delta T-cell networks and their roles in natural resistance to viral infections. *Immunol Rev* 1997;159:79–93.
- Pan J, Burdick MD, Belperio JA, Xue YY, Gerard C, Sharma S, Dubinett SM, Strieter RM. CXCR3/CXCR3 ligand biological axis impairs RENCA tumor growth by a mechanism of immunoangiostasis. *J Immunol* 2006;176:1456–1464.
- Liu Y, Poon RT, Feng X, Yu WC, Luk JM, Fan ST. Reduced expression of chemokine receptors on peripheral blood lymphocytes in patients with hepatocellular carcinoma. *Am J Gastroenterol* 2004;99:1111–1121.
- Liu Y, Poon RT, Hughes J, Feng X, Yu WC, Fan ST. Chemokine receptors support infiltration of lymphocyte subpopulations in human hepatocellular carcinoma. *Clin Immunol* 2005;114:174–182.
- Winter D, Moser J, Kriehuber E, Wiesner C, Knobler R, Trautinger F, Bombosi P, Stingl G, Petzelbauer P, Rot A, et al. Down-modulation of CXCR3 surface expression and function in CD8+ T cells from cutaneous T cell lymphoma patients. *J Immunol* 2007;179:4272–4282.
- Hegmans JP, Hemmes A, Hammad H, Boon L, Hoogsteden HC, Lambrecht BN. Mesothelioma environment comprises cytokines and T-regulatory cells that suppress immune responses. *Eur Respir J* 2006;27:1086–1095.
- Kettunen E, Nicholson AG, Nagy B, Wikman H, Seppanen JK, Stjernvall T, Ollikainen T, Kinnula V, Nordling S, Hollmen J, et al. L1CAM, INP10, P-cadherin, tPA and ITGB4 over-expression in malignant pleural mesotheliomas revealed by combined use of cDNA and tissue microarray. *Carcinogenesis* 2005;26:17–25.
- Dufour JH, Dziejman M, Liu MT, Leung JH, Lane TE, Luster AD. IFN-gamma-inducible protein 10 (IP-10; CXCL10)-deficient mice reveal a role for IP-10 in effector T cell generation and trafficking. *J Immunol* 2002;168:3195–3204.
- Kieseier BC, Tani M, Mahad D, Oka N, Ho T, Woodroffe N, Griffin JW, Toyka KV, Ransohoff RM, Hartung HP. Chemokines and chemokine receptors in inflammatory demyelinating neuropathies: a central role for IP-10. *Brain* 2002;125:823–834.
- Liu L, Huang D, Matsui M, He TT, Hu T, Demartino J, Lu B, Gerard C, Ransohoff RM. Severe disease, unaltered leukocyte migration, and reduced IFN-gamma production in CXCR3^{-/-} mice with experimental autoimmune encephalomyelitis. *J Immunol* 2006;176:4399–4409.
- Kohyama N, Shinohara Y, Suzuki Y. Mineral phases and some reexamined characteristics of the International Union Against Cancer standard asbestos samples. *Am J Ind Med* 1996;30:515–528.
- Scheule RK, Holian A. Immunologic aspects of pneumoconiosis. *Exp Lung Res* 1991;17:661–685.
- Uber CL, McReynolds RA. Immunotoxicology of silica. *Crit Rev Toxicol* 1982;10:303–319.
- Steenland K, Goldsmith DF. Silica exposure and autoimmune diseases. *Am J Ind Med* 1995;28:603–608.
- Nishimura Y, Miura Y, Maeda M, Kumagai N, Murakami S, Hayashi H, Fukuoka K, Nakano T, Otsuki T. Impairment in cytotoxicity and expression of NK cell-activating receptors on human NK cells following exposure to asbestos fibers. *Int J Immunopathol Pharmacol* 2009;22:579–590.
- Nishimura Y, Maeda M, Kumagai N, Hayashi H, Miura Y, Otsuki T. Decrease in phosphorylation of ERK following decreased expression of NK cell-activating receptors in human NK cell line exposed to asbestos. *Int J Immunopathol Pharmacol* 2009;22:879–888.
- Tossavainen A, Karjalainen A, Karhunen PJ. Retention of asbestos fibers in the human body. *Environ Health Perspect* 1994;102:253–255.
- Dodson RF, O'Sullivan MF, Huang J, Holiday DB, Hammar SP. Asbestos in extrapulmonary sites: omentum and mesentery. *Chest* 2000;117:486–493.
- Uibu T, Vanhala E, Sajantila A, Lunetta P, Makela-Bengs P, Goebeler S, Jantti M, Tossavainen A. Asbestos fibers in para-aortic and mesenteric lymph nodes. *Am J Ind Med* 2009;52:464–470.
- Noonan CW, Pfau JC, Larson TC, Spence MR. Nested case-control study of autoimmune disease in an asbestos-exposed population. *Environ Health Perspect* 2006;114:1243–1247.
- Murakami S, Nishimura Y, Maeda M, Kumagai N, Hayashi H, Chen Y, Kusaka M, Kishimoto T, Otsuki T. Cytokine alteration and speculated immunological pathophysiology in silicosis and asbestos-related diseases. *Environ Health Perspect* 2009;117:216–222.
- Nishimoto N. Interleukin-6 as a therapeutic target in candidate inflammatory diseases. *Clin Pharmacol Ther* 2010;87:483–487.

46. Nowak AK, Lake RA, Kindler HL, Robinson BW. New approaches for mesothelioma: biologics, vaccines, gene therapy, and other novel agents. *Semin Oncol* 2002;29:82-96.
47. Fitzpatrick DR, Peroni DJ, Bielefeldt-Ohmann H. The role of growth factors and cytokines in the tumorigenesis and immunobiology of malignant mesothelioma. *Am J Respir Cell Mol Biol* 1995;12:455-460.
48. Diehl S, Anguita J, Hoffmeyer A, Zapton T, Ihle JN, Fikrig E, Rincon M. Inhibition of Th1 differentiation by IL-6 is mediated by SOCS1. *Immunity* 2000;13:805-815.
49. Mantovani A, Allavena P, Sica A, Balkwill F. Cancer-related inflammation. *Nature* 2008;454:436-444.
50. Maniati E, Soper R, Hagemann T. Up for mischief? IL-17/Th17 in the tumour microenvironment. *Oncogene* 2010;29:5653-5662.

Reduction of CXC Chemokine Receptor 3 in an *In Vitro* Model of Continuous Exposure to Asbestos in a Human T-Cell Line, MT-2

Megumi Maeda¹, Yasumitsu Nishimura¹, Hiroaki Hayashi¹, Naoko Kumagai¹, Ying Chen¹, Shuko Murakami¹, Yoshie Miura³, Jun-ichi Hiratsuka², Takumi Kishimoto⁴, and Takemi Otsuki¹

¹Department of Hygiene, and ²Department of Radiation Oncology, Kawasaki Medical School, Okayama, Japan; ³Division of Molecular and Clinical Genetics, Department of Molecular Genetics, Medical Institute of Bioregulation, Kyushu University, Fukuoka, Japan; and ⁴Department at Okayama Rosai Hospital, Okayama, Japan

Because patients with silicosis who are chronically exposed to silica particles develop not only pulmonary fibrosis, but also complications involving autoimmune diseases such as rheumatoid arthritis and systemic sclerosis, exposure to asbestos may affect the human immune system. This immunologic effect may impair antitumor immune function because cancer complications such as lung cancer and malignant mesothelioma are found in patients exposed to asbestos. To elucidate the antitumor immune status caused by CD4⁺ T cells exposed to asbestos, an *in vitro* T-cell model of long-term and low-level exposure to chrysotile asbestos was established from a human adult T-cell leukemia virus-1-immortalized human polyclonal T cell line, MT-2, and the resulting six sublines showed resistance to asbestos-induced apoptosis after more than 8 months of continuous exposure. The results of DNA microarray analysis showed that the expression of 139 genes was altered by long-term and low-level exposure to asbestos, and the profile was almost similar among the six sublines when compared with the original MT-2 cells that had never been exposed to asbestos. Pathway and network analysis indicated a down-regulation of IFN- γ signaling and expression of CXC chemokine receptor 3 (CXCR3) in the sublines, whereas ELISA and flow cytometry analysis demonstrated a reduction in Th1-related IFN- γ production and cell-surface CXCR3 expression. These findings suggest that chronic exposure to asbestos may reduce antitumor immune status in CD4⁺ T cells, and that an *in vitro* T-cell model may be useful in identifying molecules related to the impairment of antitumor immune function.

Keywords: asbestos; malignant mesothelioma; CXCR3; IFN- γ

Exposure to asbestos (i.e., chrysotile, crocidolite, or amosite) leads to the development of asbestos-related diseases such as asbestos-related pleural plaque (PP) and malignant mesothelioma (MM) (1–3). Both diseases arise from exposure to asbestos, but MM has a poor prognosis, whereas PP is benign. Given that asbestos-related MM possesses a latency period ranging from 20–50 years, the peak of annual deaths from these diseases is predicted to occur around 2030 in Japan (4). Many investigations sought to elucidate the mechanisms underlying these diseases, and reports show that asbestos induces DNA damage

CLINICAL RELEVANCE

We show that long-term and low-level exposures to asbestos decrease the expression of Th1-related molecules (CXCR3, IFN- γ , and CXCL10/IP10) in the CD4⁺ T-cell line, MT-2, suggesting that exposure to asbestos may induce an impairment of antitumor immune responses. Therefore, our findings may be of use in detecting patients exposed to asbestos, identifying prognostic factors, and designing therapeutic devices to prevent the reduction of antitumor immune function found in immunocompetent cells exposed to asbestos.

and apoptosis in alveolar epithelial and mesothelial cells through a process mediated by reactive oxygen and nitrogen species of the mitochondrial dysfunction pathway (5–10).

MM is caused by exposure to asbestos, including conditions of long-term and low-level exposure. We reported that exposure to asbestos decreases the cytotoxicity of human natural killer (NK) cells, and that the cytotoxicity of NK cells is impaired in patients with MM (11, 12), suggesting that long-term and low-level exposures to asbestos may lead to a reduction of antitumor immune function. On the other hand, we showed that high-level exposures to asbestos induced the apoptosis of CD4⁺ T cells in peripheral blood mononuclear cells *in vitro* because of activation-induced cell death (13, 14). Therefore, in an effort to determine whether long-term and low-level exposures of human immune cells to asbestos can induce a reduction in antitumor immune function, we developed an *in vitro* experimental model of chronic exposure to asbestos (chrysotile), using a human T-cell leukemia virus type-1 (HTLV-1)-immortalized human polyclonal T-cell line, MT-2 (15, 16), and we successfully established an asbestos-induced, apoptosis-resistant subline (MT-2Rst) (17). Because the original MT-2 cells (MT-2Org) constitute an HTLV-1-immortalized cell line, this line can continue to divide for many generations. In previous studies, we showed that long-term and low-level exposures to chrysotile induced an up-regulation of Src-family kinase-mediated IL-10 production, with a subsequent activation of the signal transducer and activator of transcription 3 (STAT3), and an overexpression of the antiapoptotic protein Bcl-2, located downstream from STAT3 (17). In addition, short-term and high-level exposures to chrysotile promoted the production of reactive oxygen species (ROS) and triggered apoptosis via a caspase-dependent mitochondrial pathway in the original MT-2 cells (MT-2Org) (18). These mechanisms are summarized in Figure 1.

We found that patients with MM manifest a high expression of Bcl-2 in peripheral CD4⁺ T cells (17), a high level of IL-10 and transforming growth factor (TGF)- β 1 in plasma, and the multiple overrepresentation of the T-cell receptor V β in peripheral CD3⁺ T cells (19, 20). Therefore, an analysis of the immunologic effects

(Received in original form May 26, 2010 and in final form November 16, 2010)

This study was supported by Special Coordination Funds for Promoting Science and Technology grant H18-1-3-3-1 (Comprehensive Approach on Asbestos-Related Diseases), grants from the Ministry of Education, Culture, Sports, Science and Technology of Japan (18390186, 19659153, 19790411, 20390178, and 22700933), and Kawasaki Medical School Project Grants (18-209T, 19-205Y, and 20-210O).

Correspondence and requests for reprints should be addressed to Megumi Maeda, Ph.D., Department of Hygiene, Kawasaki Medical School, 577 Matsushima, Kurashiki, Okayama 7010192, Japan. E-mail: mmaeda@med.kawasaki-m.ac.jp

Am J Respir Cell Mol Biol Vol 45, pp 470–479, 2011

Originally Published in Press as DOI: 10.1165/rcmb.2010-0213QC on December 10, 2010
Internet address: www.atsjournals.org




Figure 1. Schematic representation of molecular mechanism leading to resistance of MT-2Rst cells to asbestos-induced apoptosis. MT-2Rst cells continuously exposed to low-level chrysotile-B (CB) for more than 8 months showed resistance against asbestos-induced apoptosis, accompanied by the up-regulation of Src-family kinases, IL-10, signal transducer and activator of transcription 3 (STAT3), and Bcl-2, as previously reported (17, 18). HTLV-1, human T-cell leukemia virus type-1; JNK, c-Jun N-terminal kinase; MAPK, mitogen-activated protein kinase.

of asbestos on experimental T-cell models may help explain the reduced antitumor immune function in patients with MM.

Here, we investigated differences in gene expression between MT-2Org and six independent asbestos-induced, apoptosis-resistant sublines (MT-2Rsts) in an effort to identify genes altered by long-term and low-dose exposures to asbestos in T cells. Because three of the sublines were exposed to chrysotile-A (CA) and the other three were exposed to chrysotile-B (CB), they are designated as MT-2CA1-3 and MT-2CB1-3 (the initial MT-2Rst was CB1), respectively. Using these *in vitro* models of low-level and continuous exposure to asbestos, we found a down-regulation of Th1-type molecules such as CXC chemokine receptor 3 (CXCR3), chemokine (C-X-C motif) ligand 10 (CXCL10)/IFN- γ -induced protein 10 kD (IP10), and IFN- γ in MT-2Rsts cells.

These findings may be useful in detecting patients exposed to asbestos, in identifying prognostic factors, and in designing therapeutic devices to prevent the reduction of antitumor immune function found in immunocompetent cells exposed to asbestos.

MATERIALS AND METHODS

Cell Lines and Asbestos

MT-2Org and MT-2Rsts cells were passaged many times in RPMI-1640 medium supplemented with 10% FBS, streptomycin, and penicillin at 37°C, and maintained in a humidified atmosphere of 5% CO₂. The International Union against Cancer standard CA and CB were kindly provided by the Department of Occupational Health at the National Institute for Occupational Health of South Africa (21). Chrysotile asbestos is composed of Mg₃Si₂O₅(OH)₄. CA from Zimbabwe contains 2% fibrous anthophyllite, although CB from Canada does not contain any fibrous impurities.

Real-Time RT-PCR

Real-time RT-PCR was performed using the SYBER Green method (TaKaRa, Shiga, Japan) with the Mx3000P QPCR System (Agilent Technologies, Inc., Santa Clara, CA), as previously described (17), to amplify CXCR3, chemokine (C-C motif) ligand 4 (CCL4)/macrophage inflammatory protein-1 β (Mip-1 β), and CC chemokine receptor 5 (CCR5). We used the primers CXCR3 forward (ACACCTCCTGCTCCACCTA), CXCR3 reverse (GTTTCAGGTAGCGGTCAAAGC), CCL4/MIP-1 β forward (GAAACCTCTTTGCCACCAA), CCL4/MI

P-1 β reverse (TCACTGGGATCAGCACAGAC), CCR5 forward (TAGTCATCTTGGGGCTGGTC), and CCR5 reverse (TGTAGGGA GCCAGAAGAGA).

Flow Cytometry

Cells were stained with fluorescent conjugated antibodies for 30 minutes at 4°C. After washing with PBS, cells were analyzed on a flow cytometer (FACSCalibur; BD Biosciences, Franklin Lakes, NJ). The antibody used in this study was CXCR3-PE (clone 1C6; BD Biosciences Pharmingen, San Diego, CA).

ELISA

MT-2Org and MT-2Rsts cells (1×10^5 /ml) were cultured in 24-well plates for 72 hours. The culture supernatants were then collected and assessed for the production of IFN- γ and CXCL10/IP10 by immunoassay, using Quantikine ELISA kits (R&D Systems, Minneapolis, MN).

DNA Microarray Analysis

Total RNA was isolated using the RNeasy Mini Kit (Qiagen GmbH, Hilden, Germany), and the quality of the RNA was assessed by examining the integrity of ribosomal RNA peaks, using an Agilent 2100 Bioanalyzer (Agilent Technologies, Inc.). Purified RNA (0.5 μ g) was reverse transcribed using moloney murine leukemia virus reverse transcriptase (Agilent Technologies, Inc.) and a T7-oligo(-dT) promoter primer. After synthesis of the cDNA second strand, this product was employed to generate labeled complementary RNA (cRNA), using T7 RNA polymerase with cyanine 3-cytidine triphosphate (Low RNA Input Fluorescent Linear Amplification Kit; Agilent Technologies, Inc.). Labeled cRNA (1.5 μ g) was then fragmented and hybridized to a 60-mer oligonucleotide microarray containing approximately 41,000 human genes (Human Whole Genome Oligo Microarray; Agilent Technologies, Inc.) for 17 hours at 65°C. After washing, the array was scanned using an Agilent DNA microarray scanner.

Data analysis was performed using Genespring (Agilent Technologies, Inc.). For statistical evaluation, expression profiles were normalized for the MT-2Org cell expression ratio to unity. After the removal of saturated and low-signal genes, genes that were twofold up-regulated or down-regulated in MT-2Rsts cells compared with MT-2Org cells were listed. The resultant signal information was analyzed using the Student *t* test ($P < 0.05$), and was clustered based on correlation coefficients. The resulting sets of differentially expressed genes were examined by pathway and network analysis, using the MetaCore Analytical Suite (<http://www.genego.com>; GeneGo, St. Joseph, MI).