

# The role of p53 in an apoptotic process caused by an oral malodorous compound in periodontal tissues: a review

Izumi Aoyama, Ken Yaegaki<sup>1</sup>, Bogdan Calenic, Hisataka Ii, Nikolay Ishkitiev and Toshio Imai

Department of Oral Health, Nippon Dental University, 1-9-20 Fujimi, Chiyodaku, Tokyo, Japan

E-mail: yaegaki-k@tky.ndu.ac.jp

Received 1 November 2011

Accepted for publication 3 January 2012

Published 27 February 2012

Online at stacks.iop.org/JBR/6/017104

## Abstract

Oral malodor is caused by volatile sulfur compounds (VSCs) composed mainly of hydrogen sulfide (H<sub>2</sub>S) and methyl mercaptan. In particular, H<sub>2</sub>S is an important compound, since it is a major component of physiologic halitosis. The toxicity of VSCs is similar to that of hydrogen cyanide, and is well investigated. The role of VSCs in reducing collagen in human gingival fibroblasts is one of the main sources of their toxicity to human oral tissues. It has been reported recently that H<sub>2</sub>S may cause apoptosis in several periodontal tissues. In human gingival fibroblasts, H<sub>2</sub>S inhibits not only cytochrome c oxidase activity but also superoxide dismutase activity. The levels of reactive oxygen species are markedly increased, which causes the release of cytochrome c into the cytoplasm, resulting in caspase-9 activation; finally, the executor caspase, caspase-3, is activated. This pathway is commonly observed in cells from all periodontal tissues. Moreover, p53, an apoptotic factor, and phosphorylated p53, which is the activated form, are increased by H<sub>2</sub>S in keratinocyte stem cells and osteoblasts. H<sub>2</sub>S also increases the expression of Bax, a primary response gene playing an important role in p53-mediated apoptosis, but maintains a lower expression of Bcl-2, an anti-apoptotic factor, in osteoblasts. It is concluded that the Bax apoptotic pathway and the mitochondrial pathway are activated by H<sub>2</sub>S.

## Introduction

Oral malodor is caused by volatile sulfur compounds (VSCs), mainly hydrogen sulfide (H<sub>2</sub>S) and methyl mercaptan [1, 2]. Indeed, H<sub>2</sub>S is always present in both physiological and oral pathological halitosis, and is a major component of physiological halitosis [3–5]. The toxicity of H<sub>2</sub>S is similar to that of hydrogen cyanide; H<sub>2</sub>S is recognized as a potent inhibitor of cytochrome-c oxidase (COX), a key enzyme of the respiratory chain in the mitochondria that produces adenosine triphosphate (ATP) [6]. COX inhibition by H<sub>2</sub>S has been observed in some tissues and in purified COX [6, 7]. The key mechanism by which H<sub>2</sub>S inhibits COX is in its binding

to the heme iron of the enzyme, thus completely inhibiting the aerobic metabolism producing ATP [7].

Therefore, some studies have focused not only on the esthetic problems of VSCs but also on their toxicity to oral tissues, especially periodontal tissues. Studies have shown that increased levels of H<sub>2</sub>S in the oral environment demonstrate highly toxic effects on periodontal tissues and play a role in the etiology and development of periodontitis [8–11]. Moreover, the activation of osteoclasts *in vitro* and in rat alveolar bone [10, 11] has been reported.

The apoptotic process is also involved in periodontal pathology; apoptosis plays an important role in the onset and progress of periodontal conditions [12–17]. A disruption of this between cell loss and cell production might induce periodontal conditions. The two main mechanisms in the apoptotic process are an intrinsic pathway involving the

<sup>1</sup> Author to whom any correspondence should be addressed.

mitochondria and an extrinsic pathway involving a receptor–ligand-mediated mechanism [16, 17].


Recently, it has been shown that H<sub>2</sub>S induces apoptosis in cells from several human periodontal tissues. Apoptosis in human periodontal tissues may be able to develop or control inflammation as well as cell destruction in periodontitis [18, 19]. If apoptosis of the cells is delayed, inflammatory cells may stay longer in periodontal tissues. Furthermore, excessive cytokine secretion following the process described above may cause periodontal destruction [20–22].

p53, a Bcl-2 family member, is one of the triggers of apoptosis in periodontal tissues [23]. p53, a typical tumor suppressor gene, functions as a transcriptional activator. Apoptosis, DNA damage and its repair are controlled by p53 [24]. p53 can also initiate cell-cycle control and avert further degradation of genomic DNA; p53 prevents replication of damaged DNA by apoptosis [25]. Since H<sub>2</sub>S strongly inhibits superoxide dismutase, and as huge amounts of superoxide species are produced, it is clear that the mitochondrial membrane becomes depolarized and thus the intrinsic pathway is activated [23]. In the extrinsic pathway, caspase-8 is activated by a death-inducing signal complex with specific membrane receptors, and also by a pathway requiring p53 [14]. It has been shown that specific phosphorylation of p53 regulates biological activities such as growth arrest or apoptosis [26]. Phosphorylation of serine-46 is involved in the p53-dependent apoptotic process after genomic DNA damage. In periodontal tissues, it has been shown that the main pathway causing apoptosis is the intrinsic pathway which may involve p53 [23]. Thus, p53 might play an important role in the process. In this paper, we refer to the papers reporting apoptosis in periodontal tissue cells caused by H<sub>2</sub>S in headspace air, and the current discussion is also focused on the role of p53 in the apoptosis of periodontal tissues.


### Apoptosis of gingival fibroblasts

After fibroblasts are incubated with H<sub>2</sub>S in headspace air, severe DNA fragmentation is found and the percentage of apoptotic cells increases in a time-dependent manner (figures 1 and 2) [27, 28]. H<sub>2</sub>S inhibits COX, resulting in increased levels of reactive oxygen species (ROS) in the mitochondria, and collapses the mitochondrial membrane potential; the increased production of ROS in mitochondria causes disruption of the electrochemical gradient across the inner mitochondrial membrane [9]. H<sub>2</sub>S at the lowest concentration in periodontal pockets or relatively healthy gingival crevices strongly inhibits both CuZn-SOD (superoxide dismutase) and Mn-SOD. Moreover, the percentage inhibition of hepatocyte growth factor (HGF)-SOD by H<sub>2</sub>S was 79.6 ± 5.1 and a significantly increased amount of ROS was measured [27], indicating that an increase in the amount of ROS may be induced by SOD inhibition as well as COX inhibition caused by H<sub>2</sub>S. Furthermore, it has been suggested that the increased ROS caused by SOD inhibition with H<sub>2</sub>S might be carcinogenic (figures 3 and 4).


Increased ROS causes a significant loss of the mitochondrial inner transmembrane potential as described




**Figure 1.** DNA fragmentation and hydrogen sulfide incubation time in human gingival fibroblasts. DNA fragmentation was assayed using the cell death detection ELISA kit™. (A) DNA fragmentation in the cells. (B) DNA fragmentation in the medium (\**p* < 0.05, ANOVA). Reused under license from [27].



**Figure 2.** Comparison of apoptotic to necrotic and late-apoptotic cells induced by H<sub>2</sub>S in human gingival fibroblasts. Apoptosis levels were detected by flow cytometry. (A) Percentage of early apoptotic cells. After 72 h, a significant difference was found between samples and the corresponding control. (B) Late apoptotic and necrotic cells. After 72 h, the levels of necrosis and late apoptosis were greatly increased. Each bar represents the mean ± SD of five independent experiments (\**p* < 0.05, ANOVA). Reused under license from [27].



**Figure 3.** SOD inhibition in human gingival fibroblasts. %SOD inhibition was determined by the SOD Assay Kit-WST. Reused under license from [27].



**Figure 4.** ROS produced by H<sub>2</sub>S, and MitoSOX positive cells producing ROS. After H<sub>2</sub>S incubation, a significant difference was found among the test groups. Each bar represents the mean ± SD of five independent experiments (\**p* < 0.01, Mann–Whitney test). Reused under license from [27].

**Table 1.** Caspase activity in human gingival fibroblasts. Reused under license from [28].

	24 h		48 h	
	Caspase-9	Caspase-3	Caspase-9	Caspase-3
Control	0.2 ± 0.1	0.22 ± 0.1	0.3 ± 0.1	0.3 ± 0.1
Sample	2.9 ± 1.3 <sup>a</sup>	2.9 ± 1.3 <sup>a</sup>	4.1 ± 3.2 <sup>a</sup>	4.1 ± 3.2 <sup>a</sup>


Each bar represents the mean ± SD of five independent experiments (<sup>a</sup>*p* < 0.05, ANOVA).

above. This is followed by the release of cytochrome c from the mitochondrial inner membrane into the cytosol. Cytochrome c activates caspase-9, an upstream caspase, which in turn activates caspase-3. In HGF, caspase-9 was activated, followed by caspase-3 (table 1) [28]. Caspase-3 activation ultimately leads to DNA damage and apoptosis. The inactivity of caspase-8 suggests that the extrinsic pathway is not involved in this process.

Gingival epithelial tissues, especially the gingival crevicular epithelial tissue, play an important function in periodontal pathogenesis by producing a barrier against infiltration by periodontal pathogens and the destructive products of micro-organisms [29]. It is known that VSCs increase the permeability of the gingival crevicular epithelial model [29]. Other toxic effects of VSCs to HGF are reported to include the enhancement of the secretion of prostaglandin, cAMP and procollagenase [30]. VSCs increase collagen degradation and reduce collagen synthesis in HGF [31, 32], and increased levels of VSCs also delay the wound healing process *in vivo*, thus playing an important role in periodontal pathogenesis [33].

### Apoptosis of gingival epithelial cells

VSCs also inhibit the proliferation of human gingival epithelial cells (HGECs) [34]. Among gingival tissues, epithelium plays a key function as a barrier against pathogens or toxic



**Figure 5.** Hydrogen sulfide triggers apoptosis in epithelial cells derived from human gingiva. Apoptosis levels were detected by flow cytometry. (A) Percentage of early apoptotic cells. After 24 and 48 h, a significant difference was found between samples and the corresponding control. (B) Late apoptotic and necrotic cells. After 72 h, the levels of necrosis and late apoptosis were greatly increased. Each bar represents the mean ± SD of five independent experiments (\**p* < 0.05, ANOVA). Reused under license from [35].

compounds as described above. The conservation of the epithelial barrier is therefore essential for maintaining the normal barrier function. During the progression of periodontal disease, this barrier can be affected.

In HGECs, both the levels of ROS, depolarization of the mitochondrial membrane and early apoptosis, were increased by H<sub>2</sub>S exposure (table 2, figure 5) [35]. Caspase-9 levels were also markedly increased. Finally, H<sub>2</sub>S exposure increased the caspase-3 activity, an executioner caspase, leading to the final stage of the apoptotic process (table 3).

However, caspase-8 activity stayed low. Increased caspase-8 activity initiates receptor–ligand-mediated apoptosis. The low activity of the enzyme indicates that the extrinsic pathway, receptor–ligand-mediated apoptosis, is not involved in H<sub>2</sub>S-induced apoptosis in gingival epithelial cells (table 3) [35]. These results suggest that H<sub>2</sub>S induces apoptosis by activating the intrinsic pathway, mitochondria-mediated apoptosis, in HGEC (figure 5).

Genomic DNA damage was detected using single-cell gel electrophoresis (Comet Assay; Trevigen, Gaithersburg, MD, USA). DNA involves a highly organized structure which deteriorates when damaged. The procedure is very practical and involves a simple theory. Briefly, the cells lysed using a lysis solution are placed in an electrophoretic field. After electrophoresis, DNA in the cells is stained with SYBR green, and then image analysis of the damaged DNA is carried out using both the fluorescence microscope and imaging software. Image analysis is based on the principle that intact DNA migrates extremely slowly during electrophoresis compared to the damaged DNA, which moves much faster than intact DNA because of the small DNA fragments. In image analysis, the cell is shaped like a comet; the head of the comet corresponds to the intact nucleus DNA region and its tail to the fragments of damaged DNA. The number of DNA strand breaks is correlated with the DNA comet tail shape; the size or shape of the tail represents the level of DNA damage, i.e. the number of DNA strand breaks is correlated with comet-tail parameters, such as the tail length and percentage of DNA in the tail, and their product, the tail moment. Genomic DNA damage as well as apoptosis was found in HGECs after exposure to H<sub>2</sub>S

**Table 2.** Mitochondrial changes in epithelial cells derived from human gingiva. Reused under license from [35].


	ROS(%)		Depolarization (%)		Cytochrome c (ng ml <sup>-1</sup> )	
	24 h	48 h	24 h	48 h	24 h	48 h
Control	7.6 ± 3.6	9.1 ± 3.7	14.5 ± 5.0	19.6 ± 5.5	0.02 ± 0.01	0.02 ± 0.01
Sample	56.6 ± 4.1 <sup>a</sup>	56.6 ± 4.1 <sup>a</sup>	45.8 ± 11 <sup>a</sup>	38.1 ± 10.5 <sup>a</sup>	0.12 ± 0.02 <sup>b</sup>	0.21 ± 0.02 <sup>b</sup>

Each data point represents SD of five independent experiments. (<sup>a</sup>*p* < 0.01, <sup>b</sup>*p* < 0.05, ANOVA).


**Table 3.** Caspase activities in epithelial cells derived from human gingiva. Reused under license from [35].

	24 h			48 h		
	Caspase-9	Caspase-3	Caspase-8	Caspase-9	Caspase-3	Caspase-8
Control	5.3 ± 2.5	3.3 ± 1.1	4.5 ± 2.3	4.6 ± 1.1	3.0 ± 1.2	2.8 ± 2.1
Sample	18.7 ± 3.8 <sup>a</sup>	25.6 ± 8.4 <sup>a</sup>	5.5 ± 2.1	51.1 ± 2.8 <sup>a</sup>	46.1 ± 8.0 <sup>a</sup>	7.0 ± 1.5

Each bar represents the mean ± SD of five independent experiments (<sup>a</sup>*p* < 0.05, ANOVA).



**Figure 6.** Genomic DNA damage in human gingival epithelial cells. Genomic DNA damage was detected using three different parameters. (A) Tail length, expressing distance of damaged DNA migration from the nucleoid, was significantly increased after 24 and 48 h compared with their controls. (B) Percentage of DNA in tail was increased after both 24 and 48 h. (C) Tail moment, representing the product of the first two parameters, was significantly increased at each time point compared with their controls. Each bar represents the mean ± SD of five independent experiments; 75 nuclei analyzed per experiment (<sup>\*</sup>*p* < 0.05, ANOVA). Reused under license from [35].



**Figure 7.** H<sub>2</sub>S triggers apoptosis in keratinocyte stem cells. H<sub>2</sub>S triggers apoptosis in keratinocyte stem cells. (A) Percentage of early apoptotic cells after both 24 and 48 h, a significant difference was found between samples and their corresponding controls. (B) Late apoptotic and necrotic cells were <10% at each time point. Each bar represents the mean ± SD of five independent experiments (<sup>\*</sup>*p* < 0.05, ANOVA). Reproduced with permission from the American Academy of Periodontology [23].

(figures 6(A)–(C)) [35]. It is also suggested that the p53 pathway, which is initiated by DNA damage, might be involved in the apoptotic process and that H<sub>2</sub>S may have pathological effects on human gingival epithelia at the genomic level [23].

### Apoptosis of keratinocyte stem cells: the role of p53

Keratinocyte stem cells play a key role in maintaining gingival crevicular epithelium which composes the periodontal tissues against invasion of periodontal pathogens and their products. Thus, crevicular epithelium is the first barrier against bacterial infiltration. It has been found that H<sub>2</sub>S induces apoptosis in these cells (figure 7). As can be seen in HGFs or HGECs, caspase-8 activity levels remained very low after H<sub>2</sub>S exposure

**Table 4.** Caspase-3, 8, 9 activities in keratinocyte stem cells. Reused with permission from the American Academy of Periodontology [23].


	24 h			48 h		
	Caspase-9	Caspase-8	Caspase-3	Caspase-9	Caspase-8	Caspase-3
Control	0.05 ± 0.01	0.24 ± 0.05	0.04 ± 0.01	0.06 ± 0.02	0.26 ± 0.05	0.05 ± 0.02
Sample	3.08 ± 0.41 <sup>a</sup>	0.17 ± 0.06	0.49 ± 0.13 <sup>a</sup>	5.67 ± 0.42 <sup>a</sup>	0.19 ± 0.08	0.76 ± 0.18 <sup>a</sup>

Each bar represents the mean ± SD of five independent experiments (<sup>a</sup>*p* < 0.05, ANOVA).

**Table 5.** Genomic DNA damage in keratinocyte stem cells. Reused with permission from the American Academy of Periodontology [23].

	24 h			48 h		
	Tail Length (px)	%DNA (Percentage)	Tail Moment	Tail Length (px)	%DNA (Percentage)	Tail Moment
Control	2.17 ± 1.01	2.26 ± 0.05	0.09 ± 0.03	1.02 ± 0.09	2.72 ± 0.4	0.11 ± 0.05
Sample	7.82 ± 2.94 <sup>a</sup>	12.92 ± 4.12 <sup>a</sup>	1.51 ± 0.55 <sup>a</sup>	16.71 ± 6.86 <sup>a</sup>	16.52 ± 4.14 <sup>a</sup>	3.95 ± 0.71 <sup>a</sup>

Each data point represents means ± standard deviations of five independent experiments; 75 nuclei analyzed per experiment (<sup>a</sup>*p* < 0.05).



**Figure 8.** Mitochondrial changes in keratinocyte stem cells. (A) ROS was significantly increased compared to their controls. (B) Mitochondrial membrane electrical gradient was disrupted after 24 and 48 h. (C) Cytochrome c into cytosol was significantly increased compared to the corresponding control groups. Each bar represents the mean ± SD of five independent experiments (<sup>a</sup>*p* < 0.05, ANOVA). Reused with permission from the American Academy of Periodontology [23].

of keratinocyte stem cells, while caspase-9 and caspase-3 were increased in a time-dependent manner (table 4) [23]. DNA fragmentation parameters, such as the tail length, percentage


of DNA in the tail and the tail moment, were also increased. As a result, DNA strand breaks were dramatically increased after H<sub>2</sub>S exposure (table 5) [23].

After H<sub>2</sub>S exposure ROS was significantly increased (figure 8(A)), and the number of mitochondrial membrane-depolarized cells was also increased (figure 8(B)). Consequently, a significant amount of cytochrome c was released into the cytosol (figure 8(C)).


Therefore, the extrinsic pathway is not involved in H<sub>2</sub>S-induced apoptosis. Bax, a member of the Bcl-2 protein family, is a p53 primary response gene causing p53-mediated apoptosis. The Bcl-2 family is composed of two groups involving exactly opposite functions: one is an initiator of apoptosis like Bax and another suppresses apoptosis as Bcl-2 does. A tumor-suppressor gene, p53, can act as the main defense mechanism against cancer by activating apoptosis and exterminating the cancer cell. Both total and phosphorylated p53 are increased by H<sub>2</sub>S exposure (figure 9), and Bax also increases in a time-dependent manner (figure 10) [23]. Bax, in general an apoptosis-promotion gene, usually exists in the cytoplasm; it shifts to mitochondria as a response to the apoptosis-inducing signal. An increase in ROS levels collapses the mitochondrial transmembrane potential, which is a key process for the release of cytochrome c, and then activates caspase-9 and caspase-3. H<sub>2</sub>S triggers apoptosis in keratinocyte stem cells by activating a p53-mediated apoptotic pathway, followed by the activation of the mitochondrial pathway as described above.

### Apoptosis of osteoblasts: the role of p53

Alveolar bone loss is one of the most important changes in the development of periodontitis as well as the destruction of connective tissues. H<sub>2</sub>S also induces apoptosis in osteoblasts. The total p53 level in H<sub>2</sub>S-exposed cells was significantly increased compared to their respective controls (48.45 ± 4.76 versus 15.04 ± 1.44 pg ml<sup>-1</sup> for 24 h; 70.99 ± 5.70 versus 12.57 ± 2.62 pg ml<sup>-1</sup> for 36 h; and 72.50 ± 6.65 versus 15.51 ± 3.82 pg ml<sup>-1</sup> for



**Figure 9.** Effect of H<sub>2</sub>S on p53 levels in keratinocyte stem cells. Total p53 (A) and p53 phosphorylated at serine 46 (B) were analyzed using enzyme-linked immunosorbent assay and were increased after 24 and 8 h of incubation. Each bar represents the mean  $\pm$  SD of five independent experiments (\**p* < 0.05, ANOVA). Reused with permission from the American Academy of Periodontology [23].



**Figure 10.** Effects of H<sub>2</sub>S on Bax gene expression. Gene expression was measured using quantitative RT-PCR to compare cells treated with H<sub>2</sub>S to their controls. Each bar represents the mean  $\pm$  SD of five independent experiments (\**p* < 0.05, ANOVA). Reused with permission from the American Academy of Periodontology [23].

48 h, for *p* < 0.01 and *n* = 5) [36]. p53 initiates the cell-death ligand/receptor pathway including caspase-8, one of the major pathways involved in apoptosis.

In the intrinsic pathway, p53-primary-response molecule is Bax [37]. Consequently, Bax causes mitochondrial membrane depolarization, which induces the release of cytochrome c from the mitochondria [23, 37].

After H<sub>2</sub>S exposure, the Bax gene was upregulated in a time-dependent manner, whereas Bcl-2, an anti-apoptotic gene, stayed low. Following H<sub>2</sub>S incubation, depolarization of the mitochondrial membrane was significantly increased

at 24, 36 and 48 h, respectively [36]. Following membrane depolarization, the cytochrome-c level in the cytoplasm of H<sub>2</sub>S-treated cells was significantly higher than in the controls at 24, 36 and 48 h, respectively. Cytochrome c in cytosol starts the apoptotic cascade, and activates the initiator caspase-9. Downstream, the cascade activates caspase-3, an executor enzyme. Also, a significant increase in the activity of both caspase-9 and caspase-3 was found after H<sub>2</sub>S exposure [36]. The mitochondrial pathway was activated by H<sub>2</sub>S exposure during the apoptotic process in osteoblasts as well as in other periodontal cells.

Both caspase-9 and caspase-8 are initiator caspases activating caspase-3, the executor of apoptosis. The receptor-ligand-mediated apoptosis mechanism was also examined; in this study, caspase-8 is an initiator caspase in the extrinsic pathway mentioned above. The activation of the caspase is associated with Fas, TNF-R1 and TRAILs [38–41]. Then, caspase-8 is activated with both inducible dimerization and inducible cleavage. It was inferred that caspase-8 was activated in a time-dependent manner following H<sub>2</sub>S exposure; however, TRAILs never activate caspase-8 in osteoblasts [42, 43].

Therefore, caspase-8 activation in this study could not have been caused by TRAILs. Also, both Fas and Fas-ligand activity levels remained low, suggesting that they are not involved in H<sub>2</sub>S-induced apoptosis. Many cytokines including TNF- $\alpha$  activate caspase-8 and start the apoptotic process [44–51]. Therefore, the activities of TNF- $\alpha$  and other several inflammatory cytokines were determined after H<sub>2</sub>S exposure [52, 53]. There was no difference in TNF- $\alpha$ , IL- $\beta$ , IL-2, IL-4, IL-10, IFN- $\gamma$ , G-CSF and GM-CSF levels between the tests and their respective controls.

Consequently, the cell-death ligand and receptors in the extrinsic pathway may not be involved in osteoblasts apoptosis caused by H<sub>2</sub>S, whereas caspase-8 is activated during the apoptotic process. Previous studies clearly demonstrated that p53 directly initiates caspase-8 independent of the extrinsic pathways [38–40]. Both p53 and caspase-8 were significantly activated after H<sub>2</sub>S exposure, while no cell-death ligand and receptors were activated [36]. This result suggests that caspase-8 can be activated in the absence of a death-receptor signaling and caused by p53 as reported previously [38–40]. Caspase-8 activation by H<sub>2</sub>S might be directly associated with p53.

## Conclusion

In conclusion, H<sub>2</sub>S at concentrations normally found in human gingival crevicular fluid induces apoptosis in gingival fibroblasts, gingival epithelial cells, keratinocyte stem cells and osteoblasts. H<sub>2</sub>S triggers apoptosis by activating a p53-mediated apoptotic pathway in keratinocyte stem cells and osteoblasts, and it is suggested that H<sub>2</sub>S may also activate p53 in other cells. Only in osteoblasts among these periodontal tissues do the molecular mechanisms underlie those apoptotic processes that include caspase-8 activation. In such a process, p53 activates both caspase-8 and caspase-9.

## References

- [1] Tonzetich J 1971 Direct gas chromatographic analysis of sulphur compounds in mouth air in man *Arch. Oral Biol.* **16** 587–97
- [2] Van den Velde S, van Steenberghe D, Van Hee P and Quirynen M 2009 Detection of odorous compounds in breath *J. Dent. Res.* **88** 285–9
- [3] Yaegaki K and Sanada K 1992 Biochemical and clinical factors in fluencing oral malodour in periodontal patients *J. Periodontol.* **63** 783–9
- [4] Yaegaki K and Sanada K 1992 Volatile sulfur compounds in mouth air from clinically healthy subjects and patients with periodontal disease *J. Periodontol. Res.* **27** 233–8
- [5] Murata T, Yamaga T, Iida T, Miyazaki H and Yaegaki K 2002 Classification and examination of halitosis *Int. Dent. J.* **52** (Suppl. 3) 181–6
- [6] Dorman D C, Moulin F J, McManus B E, Mahle K C, James R A and Struve M F 2002 Cytochrome oxidase inhibition induced by acute hydrogen sulfide inhalation: correlation with tissue sulfide concentrations in the rat brain, liver, lung, and nasal epithelium *Toxicol. Sci.* **65** 18–25
- [7] Beauchamp R O Jr, Bus J S, Popp J A, Boreiko C J and Andjelkovich D A 1984 A critical review of the literature on hydrogen sulfide toxicity *Crit. Rev. Toxicol.* **13** 25–97
- [8] Yaegaki K and Sanada K 1992 Volatile sulfur compounds in mouth air from clinically healthy subjects and patients with periodontal disease *J. Periodontol. Res.* **27** 233–8
- [9] Yaegaki K 2008 Oral malodorous compounds are periodontally pathogenic and carcinogenic *Japan Dent. Sci. Rev.* **44** 100–8
- [10] Irie K, Ekuni D, Yamamoto T, Morita M, Yaegaki K, Ii H and Imai T 2009 A single application of hydrogen sulfide application induces a transient osteoclast differentiation with RANKL expression in the rat model *Arch. Oral Biol.* **54** 723–9
- [11] Ii H, Imai T, Yaegaki K, Irie K, Ekuni D and Morita M 2010 Oral malodorous compound induces osteoclast differentiation without receptor activator of nuclear factor  $\kappa$ B Ligand *J. Periodontol.* **81** 1691–7
- [12] Kato S, Nakashima K, Inoue M, Tomioka J, Nonaka K, Nishihara T and Kowashi Y 2000 Human epithelial cell death caused by *Actinobacillus actinomycetemcomitans* infection *J. Med. Microbiol.* **49** 739–45
- [13] Hasebe A et al 2004 Biological activities of bacteroides forsythus lipoproteins and their possible pathological roles in periodontal disease *Infect. Immun.* **72** 1318–25
- [14] Ekuni D, Tomofuji T, Yamanaka R, Tachibana K, Yamamoto T and Watanabe T 2005 Initial apical migration of junctional epithelium in rats following application of lipopolysaccharide and proteases *J. Periodontol.* **76** 43–8
- [15] Bantel H, Beikler T, Flemmig T F and Schulze-Osthoff K 2005 Caspase activation is involved in chronic periodontitis *FEBS Lett.* **579** 5559–64
- [16] Vitkov L, Krautgartner W D and Hannig M 2005 Bacterial internalization in periodontitis *Oral Microbiol. Immunol.* **20** 317–21
- [17] Urnowey S, Ansai T and Bitko V 2006 Temporal activation of anti- and pro-apoptotic factors in human gingival fibroblasts infected with the periodontal pathogen, porphyromonas gingivalis: potential role of bacterial proteases in host signaling *BMC Microbiol.* **6** 26
- [18] Jewett A, Hume W R, Le H, Huynh T N, Han Y W, Cheng G and Shi W 2000 Induction of apoptotic cell death in peripheral blood mononuclear and polymorphonuclear cells by an oral bacterium, *Fusobacterium nucleatum* *Infect. Immun.* **68** 1893–8
- [19] Gamonal J, Bascones A, Acevedo A, Blanco E and Silva A 2001 Apoptosis in chronic adult periodontitis analyzed by *in situ* DNA breaks, electron microscopy, and immunohistochemistry *J. Periodontol.* **72** 517–25
- [20] Johnson P, Yaegaki K and Tonzetich J 1996 Effect of methyl mercaptan on synthesis and degradation of collagen *J. Periodontol. Res.* **31** 323–9
- [21] Johnson P W, Yaegaki K and Tonzetich J 1992 Effect of volatile thiol compounds on protein metabolism by human gingival fibroblasts *J. Periodontol. Res.* **27** 553–61
- [22] Ng W and Tonzetich J 1984 Effect of hydrogen sulfide and methyl mercaptan on the permeability of oral mucosa *J. Dent. Res.* **63** 994–7
- [23] Calenic B, Yaegaki K, Kozhuharova A and Imai T 2010 Oral malodorous compound causes oxidative stress and p53-mediated programmed cell death in keratinocyte stem cells *J. Periodontol.* **81** 1317–23
- [24] Gotz C and Montenarh M 1995 P53 and its implication in apoptosis *Int. J. Oncol.* **6** 1129–35
- [25] Zhang X P, Liu F and Wang W 2011 Two-phase dynamics of p53 in the DNA damage response *Proc. Natl. Acad. Sci. USA* **108** 8990–5
- [26] Sluss H K, Gannon H, Coles A H, Shen Q, Eischen C M and Jones S N 2010 Phosphorylation of p53 serine 18 upregulates apoptosis to suppress Myc-induced tumorigenesis *Mol. Cancer Res.* **8** 216–22
- [27] Yaegaki K, Qian W, Murata T, Imai T, Sato T, Tanaka T and Kamoda T 2008 Oral malodorous compound causes apoptosis and genomic DNA damage in human gingival fibroblasts *J. Periodontol. Res.* **43** 391–9
- [28] Fujimura M, Calenic B, Yaegaki K, Murata T, Ii H, Imai T, Sato T and Izumi Y 2010 Oral malodorous compound activates mitochondrial pathway inducing apoptosis in human gingival fibroblasts *Clin. Oral Invest.* **14** 367–73
- [29] Ng W and Tonzetich J 1984 Effect of hydrogen sulfide and methyl mercaptan on the permeability of oral mucosa *J. Dent. Res.* **63** 994–7
- [30] Ratkay L G, Waterfield J D and Tonzetich J 1995 Stimulation of enzyme and cytokine production by methyl mercaptan in human gingival fibroblast and monocyte cell cultures *Arch. Oral Biol.* **40** 337–44
- [31] Johnson P, Yaegaki K and Tonzetich J 1996 Effect of methyl mercaptan on synthesis and degradation of collagen *J. Periodontol. Res.* **31** 323–9
- [32] Johnson P W, Yaegaki K and Tonzetich J 1992 Effect of volatile thiol compounds on protein metabolism by human gingival fibroblasts *J. Periodontol. Res.* **27** 553–61
- [33] Shimizu H, Nakagami H, Morita S, Tsukamoto I, Osako M K, Nakagami F, Shimosato T, Minobe N and Morishita R 2009 New treatment of periodontal diseases by using NF- $\kappa$ B decoy oligodeoxynucleotides via prevention of bone resorption and promotion of wound healing *Antioxid. Redox. Signal* **11** 2065–75
- [34] Setoguchi T, Machigashira M, Yamamoto M, Yotsumoto Y, Yoshimori M, Izumi Y and Yaegaki K 2002 The effects of methyl mercaptan on epithelial cell growth and proliferation *Int. Dent. J.* **52** 241–6
- [35] Calenic B, Yaegaki K, Murata T, Imai T, Aoyama I, Sato T and Ii H 2010 Oral malodorous compound triggers mitochondrial-dependent apoptosis and causes genomic DNA damage in human gingival epithelial cells *J. Periodontol. Res.* **45** 31–7
- [36] Aoyama I, Calenic B, Imai T, Ii H and Yaegaki K 2011 Oral malodorous compound causes both caspase-8 and -9 mediated programmed cell death in osteoblasts *J. Periodontol. Res.* at press
- [37] Wesselborg S, Engels I H, Rossmann E, Los M and Schulze-Osthoff K 1999 Anticancer drugs induce

- caspase-8/FLICE activation and apoptosis in the absence of CD95 receptor/ligand interaction *Blood* **93** 3053–63
- [38] Liedtke C, Groger N, Manns M P and Trautwein C 2003 The human caspase-8 promoter sustains basal activity through SP1 and ETS-like transcription factors and can be up-regulated by a p53-dependent mechanism *J. Biol. Chem.* **278** 27593–604
- [39] Chipuk J E, Kuwana T, Bouchier-Hayes L, Droin N M, Newmeyer D D, Schuler M and Green D 2004 Direct activation of Bax by p53 mediates mitochondrial membrane permeabilization and apoptosis *Science* **303** 1010–4
- [40] Ding H F, McGill G, Rowan S, Schmaltz C, Shimamura A and Fisher D E 1998 Oncogene-dependent regulation of caspase activation by p53 protein in a cell-free system *J. Biol. Chem.* **273** 28378–83
- [41] Ding H F, Lin Y L, McGill G, Joo P, Zhu H, Blenis J, Yuan J and Fisher D E 2000 Essential role for caspase-8 in transcription-independent apoptosis triggered by p53 *J. Biol. Chem.* **275** 38905–11
- [42] Haupt S, Berger M, Goldberg Z and Haupt Y 2003 Apoptosis—the p53 network *J. Cell Sci.* **116** 4077–85
- [43] Oberst A, Pop C, Tremblay A G, Blais V, Denault J B, Salvesen G S and Green D R 2010 Inducible dimerization and inducible cleavage reveal a requirement for both processes in caspase-8 activation *J. Biol. Chem.* **285** 16632–42
- [44] Atkins G J, Bouralexis S, Evdokiou A, Hay S, Labrinidis A, Zannettino A C, Haynes D R and Findlay D M 2002 Human osteoblasts are resistant to Apo2L/TRAIL-mediated apoptosis *Bone* **31** 448–56
- [45] Bu R, Borysenko C W, Li Y, Cao L, Sabokbar A and Blair H C 2003 Expression and function of TNF-family proteins and receptors in human osteoblasts *Bone* **33** 760–70
- [46] Son Y O et al 2006 Quercetin, a bioflavonoid, accelerates TNF-alpha-induced growth inhibition and apoptosis in MC3T3-E1 osteoblastic cells *Eur. J. Pharmacol.* **529** 24–32
- [47] Chua C C, Chua B H, Chen Z, Landy C and Hamdy R C 2002 TGF-beta1 inhibits multiple caspases induced by TNF-alpha in murine osteoblastic MC3T3-E1 cells *Biochem. Biophys. Acta* **1593** 1–8
- [48] Suh K S, Koh G, Park C Y, Woo J T, Kim S W, Kim J W, Park I K and Kim Y S 2003 Soybean isoflavones inhibit tumor necrosis factor-alpha-induced apoptosis and the production of interleukin-6 and prostaglandin E2 in osteoblastic cells *Phytochemistry* **63** 209–15
- [49] Kim K B, Choi Y H, Kim I K, Chung C W, Kim B J, Park Y M and Jung Y K 2002 Potentiation of Fas- and TRAIL-mediated apoptosis by IFN-gamma in A549 lung epithelial cells: enhancement of caspase-8 expression through IFN-response element *Cytokine* **20** 283–38
- [50] Hopkins-Donaldson S, Ziegler A, Kurtz S, Bigosch C, Kandioler D, Ludwig C, Zangemeister-Wittke U and Stahel R 2003 Silencing of death receptor and caspase-8 expression in small cell lung carcinoma cell lines and tumors by DNA methylation *Cell Death Differ.* **10** 356–64
- [51] Shankaranarayanan P and Nigam S 2003 IL-4 induces apoptosis in A549 lung adenocarcinoma cells: evidence for the pivotal role of 15-hydroxyeicosatetraenoic acid binding to activated peroxisome proliferator-activated receptor gamma transcription factor *J. Immunol.* **70** 887–94
- [52] Adachi Y, Taketani S, Ikehara S, Ikebukuro K, Tokunaga R and Ikehara S 1999 Apoptosis of colorectal adenocarcinoma induced by 5-FU and/or IFN-gamma through caspase 3 and caspase 8 *Int. J. Oncol.* **15** 1191–6
- [53] Refaeli Y, Van Parijs L, London C A, Tschopp J and Abbas A K 1998 Biochemical mechanisms of IL-2-regulated Fas-mediated T cell apoptosis *Immunity* **8** 615–23



## Advances in breath odor research: re-evaluation and newly-arising sciences

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2012 J. Breath Res. 6 010201

(<http://iopscience.iop.org/1752-7163/6/1/010201>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 202.246.68.10

The article was downloaded on 08/05/2012 at 07:10

Please note that [terms and conditions apply](#).

## EDITORIAL

# Advances in breath odor research: re-evaluation and newly-arising sciences

**Ken Yaegaki**

*President of the International Society for Breath Odor Research, Nippon Dental University, Tokyo, Japan*

The articles in this special section describe the most recent advances in halitosis research presented at the Ninth International Conference on Breath Odor Research, a joint conference with the XXIV CONBRAPE (Brazilian Congress of Periodontology) held at Bahia Othon Palace Hotel in Salvador, Bahia, Brazil on 25–28 May 2011. It has been almost half a century since Joseph Tonzetich of the Faculty of Dentistry at the University of British Columbia, Vancouver, Canada (the first honorable member of the International Society for Breath Odor Research (ISBOR) who died in 2000, and is known as ‘the godfather of halitosis research’) published his first halitosis paper entitled ‘Evaluation of volatile odoriferous components of saliva’ in *Archives of Oral Biology* in 1964 [1]. This was the starting point for breath-odor research, long before ISBOR was established, although research in this area had declined by the time we convened global collaboration in halitosis research.

One must ask the question; what progress have we made during the past half-century? Although a few bad-breath detectors have appeared on the market, organoleptic scoring is still likely to remain as the ‘gold standard’. In the 21st century, people in modern dentistry or medicine are still sniffing their patients’ breath in order to diagnose halitosis, a very subjective method! Halitosis is widely believed to be simple to diagnose and treat, but this is certainly not the case, considering we have not yet even completed an objective detection measurement protocol for oral malodor.

In the treatment of halitosis, for many years before Tonzetich came into the field, tongue coating had been suspected to be the main cause, and this was scientifically proven a quarter of a century ago. However, people still visually and subjectively measure the amount of tongue coating to diagnose halitosis and an objective and more accurate method is required and needs to be developed. Those of us that work in this field employ mainly subjective protocols for diagnosis, but scientists are well aware that the data obtained by subjective procedures are not universal and therefore it is not possible to compare data obtained by a subjective measure with data from a different paper. It has been suggested that our procedures in halitosis clinics are not scientific and are almost the same as they were hundreds of years ago. Why have we not made progress in these fundamental measures for halitosis clinics? Everybody involved in halitosis clinics or the related sciences should ask themselves this question.

Removing tongue coating (TC), which consists mostly of bacteria and exfoliated keratinized epithelial cells, is a radical remedy and is considered as the most effective way to treat halitosis. For many years, people employed mechanical tongue-cleaning methods, such as using a tongue scraper. In the early 1970s it was reported that mechanical stimulation promotes tongue cancer [2]. This hypothesis may not be absolutely correct, since most tongue cancers are found on the side (lateral margin) of middle third of the tongue, and so the relationship between tongue scraping and cancer has not yet been confirmed. However, there is still a possibility that mechanical stimulation is one of the causes of tongue cancer. To effectively and safely remove tongue coating there is no doubt that we must develop a novel technology without using mechanical

stimulation, but no such research efforts have yet been reported since the early 1970s.

The protocols for halitosis clinical research can be divided into two sub-classifications: short-term and long-term. For both groups, subjects must abstain from ingestion of food, drinking and oral hygiene for four hours prior to evaluation of oral malodor, as described in [3], since this is the best time to measure volatile sulfur compounds (VSCs), as defined by the ADA guidelines. A recent paper has suggested that this protocol might not be correct, since eating and oral hygiene may affect VSCs in mouth air for longer than four hours after these activities [4]. In long-term studies, three weeks' intervention is recommended by the ADA, during which time they measure malodor strength or VSC concentration at the baseline each day. The premise of this protocol is that the baseline of VSC concentration or malodor strength is constant over certain days, but there are those who doubt that the baseline is consistent.

The organoleptic procedure is perceived as the gold standard for diagnosing oral malodor; however, for clinical work on halitosis, not only detection but also quantification is required. Organoleptic measurements do not require purpose-built apparatus, which explains why this has been a popular method among clinicians. But there are many drawbacks to this type of diagnosis. Quantification of odor sensations is very difficult, and the most difficult feature of the organoleptic procedures is stimulus presentation, while objective measurements directly determine the concentration of stimulus. Therefore, halitosis detectors should be the method of choice for clinicians.

A number of halitosis detectors have been used in the past 30 years. The portable sulfide monitor was very popular, but it also reacts with other compounds which cannot be accurately detected. Recently, portable gas chromatographs (GCs) were introduced, but this exciting technology could only report a few evaluations. Even using portable or regular GCs, the ADA recommends that their protocols be employed in clinical research.

The papers selected for this special section provide some answers to the above questions or demonstrate a novel aspect of halitosis pathology. Moreover, they illustrate the diversity of approaches to the pathological and physiological activities of VSCs. It is a challenging field.

The main challenges, as noted by K Yeagaki *et al* [5], are that problems have been identified in current, widely-used protocols. (1) The baselines of VSC concentrations in mouth air varied considerably over the course of a week. (2) When subjects refrained from eating, drinking and oral hygiene, including mouth rinsing, the VSC concentrations remained constant until the subject began eating again. (3) Over a six-hour period after a meal and oral hygiene, VSC concentrations decreased significantly. The above data point to optimal times and conditions for sampling subjects. Yeagaki *et al* also compared measurements obtained using several portable devices with measurements obtained using GCs, showing that portable devices demonstrate capabilities similar to those of GCs. Thus, a recommended protocol has been established. The proposed protocol includes the following recommendations: (a) a short-term rather than long-term study is strongly recommended, since VSC concentrations are constant in the short term; (b) a crossover study would best avoid the effects of individual specificities on each clinical intervention; and (c) measurements of VSCs should preferably be carried out using either a GC or a portable GC.

To control VSC production, removing the TC is essential for the maintenance of oral hygiene and tongue-brushing or a scraper is utilized for this purpose. Mechanical stimulation needs to be eliminated as much as possible for the reasons mentioned above or to avoid unpleasant side effects. Nohno *et al* [6] have developed candy tablets containing a protease, actinidine, and effect of long-term use of these on both TC accumulation and the concentration of VSCs in mouth air has been determined. This is a novel procedure that removes the TC safely.

Although most researchers and clinicians evaluate TC accumulation subjectively and visually, this is neither scientific nor objective. Nohno *et al* have also demonstrated an objective method of evaluating TC using a digital camera and the software ImageJ (NIH, USA). Moreover, they suggest that a long-term research design might not be appropriate. To develop new halitosis sciences, ISBOR would never to nip new knowledge in the bud—we should nurture such a promising young bud.

Tangerman *et al* [7] found no association between halitosis and *H. pylori* infection of the stomach, although *H. pylori* is believed to be a cause of extra-oral pathologic halitosis. Suzuki *et al* [8] found that the use of probiotics, a now popular way for controlling oral malodor, is effective in reducing TC accumulation, but they also described its limitations. Determining the limitations of a remedy is essential for both clinicians and patients. We must encourage the adoption of such a presentation style as demonstrated by Suzuki *et al* among breath-odor clinical scientists or clinicians.

Possible chemosensory dysfunction can elicit halitosis complaints. Falcão *et al* [9] found that chemosensory dysfunction may cause pseudo-halitosis that proves very difficult to treat.

Aoyama *et al* [10] reviewed the role of p53 in the apoptosis of periodontal tissues caused by oral malodorous compounds, and emphasized its toxicity. This information could be very useful in enabling people to prevent halitosis. Ishkitiev *et al* [11] established the protocol for differentiation of human dental pulp stem cells to hepatic cells, and found that hydrogen sulfide increases hepatic differentiation. This is a positive effect of oral malodorous compounds. However, they still need to establish universally accepted standards for evaluating the toxicity or positive effects of VSCs as shown in their report, since those topics are not yet well understood among halitosis scientists.

I would like to emphasise that this editorial represents only my own views on some of the most interesting findings in halitosis research. However, I must stress again that it is time to re-evaluate the halitosis clinical sciences, and to invest in the newly-arising sciences of halitosis. We look forward to our next ISBOR conference in two years' time.

## References

- [1] Tonzetich J and Richter V J 1964 *Arch. Oral. Biol.* 9 39–45
- [2] Yaegaki K, Coil J M, Kamemizu T, and Miyazaki H 2002 *Int. Dent. J.* 52 192–6
- [3] American Dental Association 2003 *Acceptance Program Guidelines: Products Used in the Management of Oral Malodor* (Chicago, IL: American Dental Association) pp 1–14
- [4] Fukui Y, Yaegaki K, Murata T, Sato T, Tanaka T, Imai, T Kamoda T and Herai M 2008 *Int. Dent. J.* 58 159–66
- [5] Yaegaki K *et al* 2012 *J. Breath. Res.* 6 017101
- [6] Nohno K, Yamaga T, Kaneko N and Miyazaki H 2012 *J. Breath. Res.* 6 017107
- [7] Tangerman A, Winkel E G, de Laat L, van Oijen A H and de Boer W A 2012 *J. Breath. Res.* 6 017102
- [8] Suzuki N, Tanabe K, Takeshita T, Yoneda M, Iwamoto T, Oshiro S, Yamashita Y and Hirofuji T 2012 *J. Breath. Res.* 6 017106
- [9] Falcão D P, Vieira C N and Batista de Amorim R F 2012 *J. Breath. Res.* 6 017105
- [10] Aoyama I, Yaegaki K, Calenic B, Ii H, Ishkitiev N and Imai T 2012 *J. Breath. Res.* 6 017104
- [11] Ishkitiev N, Calenic B, Aoyama I, Ii H, Yaegaki K and Imai T 2012 *J. Breath. Res.* 6 017103

# Preventing Aspiration Pneumonia by Oral Health Care

JMAJ 54(1): 39–43, 2011

Koichiro UEDA\*<sup>1</sup>

## Abstract

In recent years, the need for oral care in preventing aspiration pneumonia has been recognized across the academic disciplines. Pursuing both oral cleaning and improvements in food ingestion is necessary to achieve the objective. At a minimum, cleaning should include mechanical cleaning of the tongue and palatal mucus and removal of pathogenic bacteria causing aspiration pneumonia (typically oral anaerobic bacteria). As for the improvement of eating function, the patient should be first trained in breathing and swallowing saliva so that he/she gradually regains swallowing function. Even if the patient has a gastric fistula, successful oral ingestion of several mouthfuls not only prevents aspiration pneumonia, but can also raise the quality of the social life by enhancing the patient and the family's motivation for everyday living.

Regardless of the patient's clinical condition, the oral cavity becomes moist and healthy in color as soon as it is cared for—and its care is reflected in the entire body. I believe that the oral cavity deserves greater recognition, and hope less people will have to suffer from ingestion problems and aspiration pneumonia through proper oral care in the future.

**Key words** Aspiration pneumonia, Eating function training, Oral care, Oral intake, Oral moisture retention in the oral cavity

## Introduction

Pneumonia is the fourth most common cause of death among Japanese, and is the leading cause among the elderly in need of long-term care. Moreover, 30% of those who die of pneumonia are diagnosed with aspiration pneumonia.<sup>1</sup> Thanks to intervention studies and research, it is now recognized across the academic disciplines that regular oral care by a professional can prevent aspiration pneumonia and reduce the onset of fever.<sup>2</sup> However, even oral microorganisms that are not ordinary pathogenic tend to develop into infectious diseases as a result of opportunistic infections. Culture tests of bacteria obtained from patients who contracted aspiration pneumonia using the percutaneous pneumocentesis revealed that oral resident bacteria were most common.<sup>3</sup>

Moreover, oral care not only prevents pneumonia, but has also been reported to have a positive impact on activities in daily life and cognitive functions.<sup>4</sup> Currently, the concept of oral care is broadly interpreted as *oral health care*, including not only oral cleaning but also eating function training. This paper will examine the characteristics of the oral cavities of elderly people in need of long-term care who developed aspiration pneumonia and the concept and techniques for oral care.

## Characteristics of Oral Cavity Which Causes Onset of Aspiration Pneumonia


The oral cavities of patients who contract aspiration pneumonia have characteristic mucous membranes, soft tissue, teeth, and oral func-

\*1 Professor, Dysphagia Rehabilitation, School of Dentistry, Nihon University, Tokyo, Japan (ueda-k@dent.nihon-u.ac.jp).

This article is a revised English version of a paper originally published in the Journal of the Japan Medical Association (Vol.138, No.9, 2009, pages 1785–1788).



**Fig. 1** Mucous membranes residue, remaining like an oblate



**Fig. 2** Coated tongue

tion. The key points to observe to prevent aspiration pneumonia or improve the prognosis are described below.

#### Residue in oral mucous epithelium and oral dryness

When oral functions are almost completely suppressed due to depressed consciousness or tube feeding such as intravenous drips and gastric fistula, the secretion of mucosal resting saliva begins to dominate, mixing with the residue in the oral epithelium to form a sticky paste that adheres to the oral cavity membranes and teeth. Since the self-cleaning function of the oral cavity is not working, the oral mucous membrane is not regenerated and replaced. Instead, the mucous remains on the palate like an oblate as shown in **Fig. 1**, and the tongue becomes coated (**Fig. 2**). The bacterial flora of such palate or tongue contains different bacteria that do not normally appear in the healthy mouth, raising the risk of upper respiratory tract infections and thus aspiration pneumonia.

The main causative microorganism of aspiration pneumonia is thought to be gram-negative anaerobic bacteria, which live in secreta (phlegm), epithelial residue, crusts, and saliva. A silent aspiration of these microorganisms (called microaspiration) in the process of normal respiration causes pneumonia.

#### Morphological damage to oral hard tissue (teeth)

Inadequate oral cleaning can result in multiple and simultaneous cases of dental caries at the



**Fig. 3** Remaining roots after losing the crown portion

tooth cervix. If oral hygiene continues to be neglected, the crown can rot away with only residual roots left as shown in **Fig. 3**. Food residue tends to adhere to the residual root of the decayed teeth, creating bacterial plaque, and forming bacterial flora that cannot be removed by mere gargling. If the condition reaches the point that bad breath pervades the entire room, it is evident that decaying matter is remaining in the mouth, meaning there is a grave risk of aspiration pneumonia.

#### Impaired oral function

When using a cerebral stroke as an example, more than 50% of patients in the acute phase have aspiration, but only 7% still experience aspiration during the recovery phase. But because their oral cavity organs such as the lips, tongue, cheek, and soft palate remain paralyzed even during the recovery phase, 30% of chronic patients experi-



Fig. 4 Moisturizing agents




Fig. 5 Cleaning the tongue and palate

ence impaired mastication.

As the patient recover from acute phase and move on to chronic phase, impairment of eating functions shifts from the pharyngeal phase to oral phase.<sup>5</sup> Even if the patient succeeds in ingesting food orally, poor mastication prevents the formation of alimentary bolus, and food adheres to the surface of the teeth in its original shape. This kind of functional impairment of the oral phase is one cause of poor oral hygiene and consequently induces silent aspiration or miswallowing.

### Approaches and Techniques for Oral Care

Aspiration pneumonia can be broadly divided into 2 groups: one caused by *micro-aspiration* (the aspiration of microorganisms into the trachea) and *macro-aspiration* (the aspiration of food). Micro-aspiration can be prevented by cleaning of the oral cavity. Macro-aspiration can be addressed by improving eating and swallowing functions. When pursuing both oral hygiene management and eating function restoration, the original objective of preventing aspiration pneumonia can be achieved.

#### Oral hygiene management

##### Moisture retention

The first step is to retain moisture in the oral cavity. There are 3 types of moisturizing agents: gels, creams, and liquids (Fig. 4). One may use a piece of gauze or cotton wrapped around the

finger as a cleaning tool, but it is better to use an ordinary sponge brush or special brush specifically designed for the mucous membranes in order to more scrupulously and reliably enhance the effects of moisture retention and cleaning. After spreading a moisturizing agent to a brush, apply it to the upper and lower lips, left and right buccal mucous membranes, palate, and tongue, to care for the mucous membranes.


Moisturizing agents in gel form include a polymer (poly glyceryl methacrylate), which achieves a swelling effect by absorbing moisture and softening the mucous membranes.

Moisturizing creams include substances that promote a natural healing for oral inflammation with a combination of milk protein extracts containing antimicrobial ingredients and immunoglobulin. In particular, using a toothbrush to remove epithelial residue (shown in Fig. 1) or crusts can have an adverse result of bleeding. However, applying a moisturizing agent to the dry lining in advance softens the epithelium within a few minutes, making removal easier.

Some liquid moisturizing agents contain hyaluronic acid. Once the oral cleaning is finished, spraying the liquid moisturizer and then applying a moisturizing gel can extend the moisturizing effect for a longer period.

##### Areas that must be cleaned

In the oral cavity of a tube-fed patient, microorganisms are most often stuck to 2 areas: the tongue and palate. This means that even just mechanically cleaning the tongue and palate can help to prevent aspiration pneumonia (Fig. 5).



**Fig. 6** Frequency of pneumonia in a 3-year period in the group that received both professional oral cleaning and swallowing training (11 patients)

#### Regarding eating function training

##### Key points for eating function

Mechanical stimulation from oral cleaning can stimulate the secretion of saliva and activate the swallowing and coughing reflexes. Thus, oral cleaning can be a very important anti-infection measure. When cleaning the mouth, attention should be paid to the way in which the patient disposes of saliva. In other words, if the visual observation confirms the closing of the lips and elevating of the larynx when the patient is swallowing saliva, no residual sound is noted in the pharyngeal region with a stethoscope, and the patient can expectorate when choking, then, the patient can proceed to aggressive swallowing training. Tests that utilize mechanical devices, such as videofluorography (VF) or video endoscopy (VE), should also be used for diagnosis as necessary.

As long as the patient is unable to control his/her breathing, mastication and deglutition movements will be difficult. So, first the swallowing training should start with the practice in abdominal breathing on the bed. Next, the patient can start training in swallowing saliva by licking candies to stimulate saliva secretion and induce saliva swallowing.

##### “A spoon of life”

In this section, I will discuss a comparative case study I conducted with my colleagues.<sup>6</sup> At an intensive care home for the elderly, we provided oral care to patients with dementia once a week since 1999. Of these patients, a dental professional provided weekly oral cleaning to 10 dysphagia patients with feeding tube who needed assistance

for all of their daily life activities. The outbreak frequency of pneumonia among these 10 patients was tracked over a period of 3 years. After 3 years, there was an overall decline in the frequency of pneumonia, but since there were several patients who experienced pneumonia repeatedly, the difference was not statistically significant in these 3-year period data.

However, at another facility, we provided functional training of dysphagia in addition to oral cleaning to 11 patients with similar physical conditions. The functional training was the ingestion of less than 40 ml of gelatin jelly once a week. Endoscopic diagnosis, visual examination, auscultation, and palpation were used to check for aspiration, and the patient only orally ingested the jelly when the dentist visited. Two of the 11 patients died of heart failure during the 3-year period. But as illustrated in **Fig. 6**, the remaining 9 patients showed a decrease in the frequency of pneumonia or no change, which was statistically significant.

In addition to daily oral cleaning by care givers, by providing weekly professional oral cleaning and the ingestion of a spoonful of jelly once a week, this study succeeded in reducing the frequency of pneumonia.<sup>6</sup> Even though it was only a mouthful, this ingestion of jelly was “a spoon of life.” In the latter facility, more than 15 patients a year died of pneumonia prior to 1999, but the pneumonia mortality has decreased to less than 6 since 2001. These figures confirmed the importance of using eating function in patient care, even to ingest just a spoonful of gelatin jelly, in addition to proper cleaning.

#### Conclusion

The oral problems among elderly patients in need of long-term care become progressively worse as the period of systemic illness prolongs, to the point of disintegration of the crown, stagnation of food residue, adhesion of epithelial residue, and loss of artificial teeth. Before reaching such dismal conditions, at some point in the transition—from the acute to recovery phase or from the recovery to maintenance phase, there must have been a turning point somewhere.

In recent years, the need for oral care in preventing aspiration pneumonia has been recognized across the academic disciplines. In order to successfully prevent aspiration pneumonia, a



patient must: 1) improve his/her ability to ingest food items, and 2) receive sufficient oral care including proper cleaning. To this end, oral cleaning should at least include mechanical cleaning of the tongue and palate to remove pathogenic bacteria (e.g., anaerobic bacteria) that can lead to aspiration pneumonia. As for improving eating function, the patient should first be trained in breathing and swallowing saliva in order to gradually regain swallowing function. Ingestion of several mouthfuls can successfully prevent aspiration pneumonia, even if the patient is being tube fed. The act of eating, even as little as a spoonful, can also raise motivation for life for both the patient and the family and can enhance

the quality of their social life.

The oral cavity is an organ that responds promptly to proper care, regardless of the clinical condition. It becomes moist and healthy in color as soon as it is cared for, and the improved oral cavity can also influence the entire body. "Eating" is such a simple, everyday activity for healthy people, and so is oral care. But for those who have problem with eating function, "it is only oral care, but it is part of the care." I would like to see greater recognition of oral care, so that less patients and people in recuperation will have to suffer from ingestion problems and aspiration pneumonia.

#### References

1. Kida K. Respiratory disease of the elderly: on clinical condition of aspiration pneumonia. *Geriatric Dentistry*. 1995;10:3–10. (in Japanese)
2. Yoneyama T, Yoshida M, Matsui T, et al. Oral care and pneumonia. *Lancet*. 1999;354:515.
3. Bartlett JG, Gorbach SL, Finegold SM. The bacteriology of aspiration pneumonia. *Am J Med*. 1974;56:202–207.
4. Suzuki M, Saito E, Koguchi K, et al. Dental treatment for the disabled elderly and its effect on disability. *The Journal of the Japan Dental Association*. 1999;52:608–617. (in Japanese)
5. Ueda K. Oral maladies of people with cerebrovascular disorders and prosthetic treatment. *Japanese Journal of Gerodontology*. 2002;16:320–326. (in Japanese)
6. Ueda K, Yamada Y, Toyosato A, et al. Effects of functional training of dysphagia to prevent pneumonia for patients on tube feeding. *Gerodontology*. 2004;21:108–111. (in Japanese)

## 簡易な開口力測定器の開発

## —第1報：健常者の開口力，握力および年齢との比較—

Development of a Jaw-Opening Sthenometer to Assess Swallowing Functions  
First Report : Jaw Opening Muscle Strength of Healthy Volunteers

戸原 玄<sup>1)</sup>，和田 聡子<sup>1)</sup>，三瓶 龍一<sup>1)</sup>，井上 統温<sup>1)</sup>  
佐藤 光保<sup>1)</sup>，飯田 貴俊<sup>1)</sup>，鰐原 賀子<sup>1)</sup>，岡田 猛司<sup>1)</sup>  
島野 嵩也<sup>1)</sup>，石山 寿子<sup>1)</sup>，中川 量晴<sup>2)</sup>，植田耕一郎<sup>1)</sup>

Haruka Tohara<sup>1)</sup>, Satoko Wada<sup>1)</sup>, Ryuichi Sanpei<sup>1)</sup>, Motoharu Inoue<sup>1)</sup>  
Mitsuyasu Sato<sup>1)</sup>, Takatoshi Iida<sup>1)</sup>, Katsuko Ebihara<sup>1)</sup>, Takeshi Okada<sup>1)</sup>  
Takaya Shimano<sup>1)</sup>, Hisako Ishiyama<sup>1)</sup>, Kazuharu Nakagawa<sup>2)</sup> and Koichiro Ueda<sup>1)</sup>

**抄録：**超高齢社会である日本では摂食・嚥下障害への対応は急務であるため，簡便な評価方法が過去に複数考案されてきた。ただし，いずれの評価方法も嚥下もしくは咳反射の有無や可否などを観察するものであり，嚥下の強さの判定を目的とする簡便なテストはなかった。嚥下時に収縮する舌骨上筋のうち複数の筋肉が開口筋であることに着目して，嚥下機能の評価を目的として開口力測定器を作成し，基礎的なデータを採取するために平均年齢44.7±12.6歳，64名の健常者に対して開口力の測定を行った。対象者全員の開口力の平均は約8kg，男性は約10kg，女性は約6kgであり，男性の筋力が有意に高いが年齢との開口力の相関は認められず，過去に報告された結果とも同様の数値が得られた。また，身体他部位の筋力である握力と開口力は有意に相関したことから，開発した筋力計を用いて有効な評価ができていると考えられた。また，健常な60代の舌骨上筋には加齢に伴う筋力低下がみられなかった。

**キーワード：**摂食・嚥下機能，摂食・嚥下障害，開口力測定器，舌骨上筋，開口筋

## 緒 言

超高齢社会を迎え，高齢者の主たる死因である誤嚥性肺炎を直接引き起こし得る摂食・嚥下障害への対応は急務である。特に，摂食・嚥下障害の精査には嚥下造影検査や嚥下内視鏡検査があげられるが，機器のみならずそれらの実施に必要な技術および知識が必要である。よって簡便な評価方法の必要性の高さから，これまで多くの摂食・嚥下障害のスクリーニングテストが考案されてきた<sup>1~9)</sup>。それらス

クリーニングテストの評価対象は，自発的な唾液の反復嚥下回数測定による誤嚥有無判定<sup>1,2)</sup>，水分の嚥下<sup>3~7)</sup>もしくは半固形物の嚥下<sup>7)</sup>などを用いた誤嚥有無判定，また刺激物吸入後の咳反射の出現有無観察<sup>8,9)</sup>による不顕性誤嚥有無判定であり，嚥下もしくは咳反射の有無や可否などを観察するものである。それぞれがすでに臨床応用されているが，嚥下に関連する筋力の判定を目的とする簡便なテストはなかった。

また，正常な嚥下を完了するためには舌骨挙上が必要となる。舌骨上筋が弱くなると，舌骨や喉頭の挙上量が減少するために食道入口部の開大が障害されて，咽頭残留が増加するだけではなく，誤嚥のリスクも高まる<sup>10)</sup>。よって同部への訓練的なアプローチとして，臥位で頭を持ち上げる負荷を利用して舌

<sup>1)</sup> 日本大学歯学部摂食機能療法学講座

<sup>2)</sup> 昭和大学歯学部口腔衛生学教室

<sup>1)</sup> Department of Dysphagia Rehabilitation, Nihon University School of Dentistry

<sup>2)</sup> Department of Hygiene and Oral Health, Showa University School of Dentistry

骨上筋を鍛える頭部挙上訓練も考案されている<sup>11)</sup>。この訓練を行うことで、舌骨および喉頭挙上量が増

加するために、嚥下時の食道入口部の開大が改善する<sup>11)</sup>ことから、同部の筋力は嚥下の強さを押し量るために重要な指標となる。これら舌骨挙上に関わる筋肉は顎舌骨筋、顎二腹筋、オトガイ舌骨筋、莖突舌骨筋であるが、このうち顎舌骨筋、顎二腹筋前腹、オトガイ舌骨筋は開口筋でもあることに着目し、強さをパラメーターとした嚥下に関する簡便な評価方法の確立を目的として、簡易な開口力の測定器の開発を試みた。まずは、基礎的なデータ採取の目的で健康な成人を対象に開口力を測定し、多少の知見を得たのでここに報告する。

**対象および方法**

布製ベルト、マジックテープ、熱可塑性スプリント剤 (LMB ブレンド, 株式会社 Hogy medical), および小型の等尺性筋力計 (ミュータス F1, アニマ株式会社) を用いて開口力測定器を開発した (図1)。バンドは頭部周囲に1本 (図1-①)、頭部周囲との頭頂部固定用に2本 (図1-②)、頭部周囲との下顎固定用に2本用いた (図1-③)。熱可塑性スプリント材はそれぞれのベルト固定部 (図1-④) や




図1 開口力測定器

頭部および下顎の固定用バンド, チンキャップ, および筋力計 (ミュータス F1, アニマ株式会社製) からなる。頭部と下顎のバンドはマジックテープで長さを変えられるようにし、チンキャップの下部に筋力計を備えつけてある。

- ①頭部周囲用バンド, ②頭部周囲と頭頂部の固定用バンド (2本), ③頭部周囲と下顎の固定用バンド (2本), ④ベルト固定部 (4つ), ⑤チンキャップ, ⑥筋力計 (ミュータス F1, アニマ株式会社製), ⑦モニター (ミュータス F1, アニマ株式会社製)




図2 開口力測定器装着時

頭部および下顎にバンドおよびチンキャップを合わせて、開口時に口があかないようにバンドを締めて固定する。下顎のバンドを頭部のバンドに固定する位置は被検者のこめかみおよび耳の後ろとした。

- ①頭頂部のバンドはクロスする。
- ②下顎のバンドはチンキャップ固定部直前でクロスする。

表1 健常者の握力および開口力

	年齢 (歳)	握力 (kg)	開口力 (kg)
全員 (62名)	44.7±12.6	37.5±8.7	8.2±3.0
男性 (32名)	44.6±12.8	44.2±5.3	10.3±2.6
女性 (30名)	44.8±12.6	30.2±4.8	5.9±1.5

男女の握力および開口力には有意に差がみられた。  
( $p < 0.01$ , Mann-Whitney の U 検定)




図3 健常者の開口力と握力の関係  
健常者の開口力と握力には有意な相関がみられた。

チンキャップ部 (図1-⑤) に使用し, 開口時の筋力を測定するために, チンキャップの直下に筋力計 (図1-⑥) を固定し筋力計とモニターを接続した (図1-⑦)。そのチンキャップおよび筋力計は下顎のバンドにて頭部周囲のバンドに固定した。

開口力測定器を装着した様子を示す (図2)。頭部周囲径および, 頭部から下顎までの距離は被検者によって異なるため, それぞれの長さをマジックテープで調節可能とし, また計測時にはバンドが緩んで開口しないように, 可及的にバンドを締めて固定した。なお, 頭頂部のバンドは頭頂でクロスするように位置させた (図2-①)。チンキャップ部のバンドは, 頭部周囲のバンド固定部より耳を前後からはさんでチンキャップ直前でクロスするように位置させて (図2-②), チンキャップを超えた後は同様に対側のバンド固定部に固定した。

対象は摂食・嚥下障害に関連する症状, 既往, 訴えない健常のボランティア 62 名 (男性 32 名, 女




図4 健常者の開口力と年齢の関係  
健常者の開口力と年齢には相関はみられなかった。

性 30 名, 平均年齢  $44.7 \pm 12.6$  歳, 24~64 歳) を対象とした。上記の機器を用いて開口筋群筋力測定で得られた結果を開口力とし, さらに身体他部位との筋力との整合性をみるために利き手の握力 (デジタル握力計 TL110, トーエイライト株式会社) を測定した。それぞれ 3 回測定し, その最大値を計測値とした。

本研究は日本大学歯学部倫理委員会の承認を受けて行った。得られたデータの解析には SPSS statistics 17.0 (SPSS Inc.) を用いた。

## 結 果

対象者全員 (平均年齢  $44.7 \pm 12.6$  歳, 24~64 歳) の握力は平均  $37.5 \pm 8.7$  kg, 開口力は平均  $8.2 \pm 3.0$  kg, 男性 (平均年齢  $44.6 \pm 12.8$  歳, 26~64 歳) の握力は平均  $44.2 \pm 5.3$  kg, 開口力は  $10.3 \pm 2.6$  kg, 女性 (平均年齢  $44.8 \pm 12.6$  歳, 24~61 歳) の握力は平均  $30.2 \pm 4.8$  kg, 開口力は  $5.9 \pm 1.5$  kg であり, 男女の握力および開口力には有意に差がみられた (表1)。

開口力と握力との間には有意な相関がみられた (図3)。また, 開口力と年齢との間には有意な相関は認められなかった (図4)。