

Research Article

Bactericidal Effects of Diode Laser Irradiation on *Enterococcus faecalis* Using Periapical Lesion Defect Model

Masato Nagayoshi,¹ Tatsuji Nishihara,² Keisuke Nakashima,³ Shigetsugu Iwaki,⁴ Ker-Kong Chen,⁵ Masamichi Terashita,⁶ and Chiaki Kitamura¹

¹ Division of Pulp Biology, Operative Dentistry, and Endodontics, Department of Cariology and Periodontology, Kyushu Dental College, 2-6-1 Manazuru, Kokurakita, Kitakyushu 803-8580, Japan

² Division of Infections and Molecular Biology, Department of Health Promotion, Kyushu Dental College, 2-6-1 Manazuru, Kokurakita, Kitakyushu 803-8580, Japan

³ Division of Periodontology, Department of Cariology and Periodontology, Kyushu Dental College, 2-6-1 Manazuru, Kokurakita, Kitakyushu 803-8580, Japan

⁴ Marketing Research Section, R&D Department, NISSIN DENTAL PRODUCTS INC., 22-1 miyabayashi, asahi, Kameoka, Kyoto 621-0001, Japan

⁵ Department of Conservative Dentistry, Kaohsiung Medical University Hospital and College of Dental Medicine, Kaohsiung Medical University, Shih-Chuan 1st Road, Kaohsiung 80708, Taiwan

⁶ Division of Comprehensive Dentistry, Department of Clinical Communication and Practice, Kyushu Dental College, 2-6-1 Manazuru, Kokurakita, Kitakyushu 803-8580, Japan

Correspondence should be addressed to Chiaki Kitamura, chi-aki-k@kyu-dent.ac.jp

Received 13 May 2011; Accepted 19 June 2011

Academic Editor: A. Carrassi

Copyright © 2011 Masato Nagayoshi et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Objective. Photodynamic therapy has been expanded for use in endodontic treatment. The aim of this study was to investigate the antimicrobial effects of diode laser irradiation on endodontic pathogens in periapical lesions using an *in vitro* apical lesion model. **Study Design.** *Enterococcus faecalis* in 0.5% semisolid agar with a photosensitizer was injected into apical lesion area of *in vitro* apical lesion model. The direct effects of irradiation with a diode laser as well as heat produced by irradiation on the viability of microorganisms in the lesions were analyzed. **Results.** The viability of *E. faecalis* was significantly reduced by the combination of a photosensitizer and laser irradiation. The temperature caused by irradiation rose, however, there were no cytotoxic effects of heat on the viability of *E. faecalis*. **Conclusion.** Our results suggest that utilization of a diode laser in combination with a photosensitizer may be useful for clinical treatment of periapical lesions.

1. Introduction

It is generally accepted that disinfecting processes are essential for successful root canal treatment, and antimicrobial irrigants to remove microorganisms are important for chemomechanical preparation of a root canal [1, 2]. Endodontic irrigants are required to have a broad spectrum of antimicrobial activities, as well as a relative lack of toxicity against sound periapical tissue. Sodium hypochlorite (NaOCl) is a major endodontic irrigant, however, it has cytotoxic and neurotoxic effects when extruded into periapical tissues [3, 4]. To develop a safe endodontic irrigants, ozonated water was previously examined as an endodontic

irrigant and demonstrated to be a useful irrigant for removal of microorganisms from root canals without damage to other tissues [5]. However, it is also known that complete elimination of microorganisms from root canals and periapical lesions by antimicrobial irrigants only is difficult, because of the anatomical complexities of root canals, deep invasion of microorganisms into dentinal tubules, and formation of biofilms on the surface of root apex, resulting in persistent apical periodontitis [6–8]. In addition, several studies have reported that the lack of response of refractory periapical lesion is due to the inaccessibility of the extraradicular microorganisms or to the presence of microorganisms [9–11].




FIGURE 1: Schematic representation of present *in vitro* model of apical periodontitis. Resin block A, root part with a single root canal with an apical foramen (diameter 600 μm). Resin block B, periapical tissue portion with a space (Space C, diameter 5 mm) fashioned as a periapical lesion defect.

Recently, photodynamic antimicrobial chemotherapy has received focus as an alternate antibacterial, antifungal, and antiviral treatment for drug-resistant microorganisms [12, 13]. Along that line, disinfection of root canals by laser irradiation has been demonstrated [14–16]. When using laser irradiation for disinfection of root canals and periapical lesions, damage to periapical tissues by heat produced by the irradiation procedure should be avoided [17–19]. Several studies have reported the antimicrobial effectiveness and safety of laser irradiation of root canals. However, there are no known reports regarding the antimicrobial effects of laser irradiation on microorganisms in periapical lesions without corresponding heat damage to sound periapical tissues, because of difficulties in mimicking the related environment.

A few studies have reported that use of a photosensitizer in combination with laser irradiation was effective for selective elimination of microorganisms from root canals and periapical lesions [20–22]. The aim of this study was to investigate the antimicrobial effects of diode laser irradiation in combination with a photosensitizer against *Enterococcus faecalis*, one of the major organisms related to persistent apical periodontitis [10, 11, 23, 24] in periapical lesions using an *in vitro* apical lesion model. In addition, irradiation-induced heat changes and the effects of heat on *E. faecalis* were examined using our periapical lesion model.

2. Material and Methods

2.1. Light Source and Photosensitizer. The irradiation source was a diode laser (P-Laser; Panasonic Dental Co., Ltd., Osaka, Japan). Its wavelength, output power, and duty were 805 nm, 5 watts (W), and 20%, respectively, while the diameter of quartz optical fiber was 400 μm . Indocyanine green (Ophthogreen®; Santen Pharmaceutical Co., Ltd., Osaka, Japan) (12.5 mg/mL), a commonly used fluorescent fundus contrast medium, was used as the photosensitizer.

2.2. Antimicrobial Activity of Laser Irradiation. *E. faecalis* ATCC 29212 was cultured in brain-heart infusion (BHI)

broth (Difco, Detroit, MI) at 37°C for 18 hours in an atmosphere of 5% CO₂. The organisms were harvested by centrifugation at 10,000 $\times g$ for 5 minutes then suspended in saline and adjusted to 3 $\times 10^6$ cells/mL using a spectrophotometer. Bacterial cell suspensions were mixed with sterilized saline or indocyanine green solution in sterilized test tubes for a final suspension of 1.5 $\times 10^6$ cells/mL, then individually irradiated by the diode laser at a distance of 1 mm for 30, 60, or 120 seconds. Following irradiation, each suspension (20 μL including 3 $\times 10^2$ cells) was cultured on a BHI agar plate. The inoculated plates were incubated at 37°C for 18 hours in an atmosphere of 5% CO₂ and colony-forming units (CFU) counted using the spread plate method.

2.3. Laser Irradiation Using In Vitro Model of Periapical Lesion Defect. An *in vitro* model of a periapical lesion defect was fashioned from resin blocks (Nissin Dental Products, Inc., Kyoto, Japan) (Figure 1). It was made up of a single root canal with an apical foramen (diameter; 600 μm) in resin block A, which formed the root part of the model, while resin block B, which was used as the periapical tissue portion, had a space (Space C, diameter; 5 mm) fashioned as a periapical lesion defect. The model was sterilized before each experiment. *E. faecalis* was diluted to 2 $\times 10^6$ cells/mL with prewarmed (45°C) BHI broth containing 1% agar, then 20 μL of the bacterial cell suspension was mixed with 20 μL of distilled saline or indocyanine green solution in Space C (final concentration of *E. faecalis*, 1 $\times 10^6$ cells/mL). Resin block A was then inserted into resin block B and kept at room temperature.

The optical fiber of the diode laser was inserted into the root canal to reach its apex then microorganisms in Space C were irradiated for 30, 60, or 120 seconds. In addition, the model was also irrigated with sterilized saline (4 mL) or 2.5% NaOCl (4 mL) for 120 seconds as negative and positive controls, respectively. After each treatment, *E. faecalis* in Space C were diluted 1/100 with sterilized saline and subjected to vortexing for 5 minutes, then 20 μL of each sample (2 $\times 10^2$ cells) was cultured on a BHI agar plate. CFU were determined after incubation of the inoculated plates at 37°C for 18 hours in an atmosphere of 5% CO₂.

2.4. Exposure of Microorganisms to Heat. *E. faecalis* suspensions were mixed with a sterilized indocyanine green solution (final suspension 1.5 $\times 10^6$ cells/mL) and incubated in a heat block (TERMO ALUMI BATH ALB-220: IWAKI GLASS Co., Ltd. Japan) at 65°C for 60 seconds, after which heat-treated *E. faecalis* (20 μL ; 3 $\times 10^2$ cells) were cultured on BHI agar plates. The inoculated plates were then incubated at 37°C for 18 hours in an atmosphere of 5% CO₂ in air and CFU determined.

2.5. Temperature Monitoring during Laser Irradiation. Temperatures in the periapical lesion (Space C) and surrounding areas of the *in vitro* model were monitored during laser irradiation using infrared thermography (TH9100 WV: NEC Avio Infrared Technologies Co., Ltd. Japan). Just after performing laser irradiation, the inner temperature of the periapical lesion area (Space C) was also measured




FIGURE 2: Antimicrobial effects of diode laser on *E. faecalis* in test tubes. The bacteria were exposed to laser irradiation with or without an added photosensitizer (closed diamond, without photosensitizer; closed square, with photosensitizer) for 30, 60, and 120 seconds. (A) Elimination of *E. faecalis* in model of apical periodontitis. *E. faecalis* in the presence and absence of a photosensitizer [PS(-), without photosensitizer; PS(+), with photosensitizer] in the lesion defect area of the model were exposed to diode laser irradiation for 30, 60, and 120 seconds. In addition, one sample was irrigated with sterilized saline for 120 seconds as a negative control (saline-120 s) and another with 2.5% NaOCl for 120 seconds as a positive control (2.5% NaOCl-120 s). (B) The number of viable cells after each treatment was counted. Data are expressed as the mean \pm standard deviation of triplicate determinations. The experiment was performed three times, with similar results obtained in each.

by immediate insertion of a thermocouple thermometer (Digital Thermometer CT-800: As One Corp., Japan).

2.6. *Statistical Analysis.* Differences among variables in the experiments were compared using Student's *t*-test.

3. Results

3.1. *Antimicrobial Effects of Diode Laser with Photosensitizer and Disinfection of Periapical Lesion Defect in In Vitro Model.* There were no antimicrobial effects seen without the photosensitizer, whereas in its presence the cell viability of *E. faecalis* decreased to 72% after irradiation for 30 seconds and was not detected after irradiation of 60 seconds or more in test tubes (Figure 2(A)).

Figure 2(B) shows the antimicrobial effects of diode laser irradiation on *E. faecalis* in the *in vitro* model. Irrigation with saline did not have an antimicrobial effect, while that with NaOCl completely eliminated *E. faecalis*. When the diode laser was used without the photosensitizer for 120 seconds, the decrease in viability was only 25%. In contrast, a significant decrease in the viability of *E. faecalis* was observed following laser irradiation with the photosensitizer for 60 and 120 seconds.

3.2. *Effects of Heat Produced by Laser Irradiation on Bacterial Cell Viability.* The inner temperature of the apical lesion area of the *in vitro* model was determined with a thermocouple thermometer just after laser irradiation (Figure 3(A)). Irradiation for 60 seconds increased the temperature of the periapical lesion defect to 65°C. To analyze the cytotoxic effects

of that elevated temperature (65°C) on *E. faecalis*, bacterial cells were incubated in a heat block at 65°C for 60 seconds (Figure 3(B)). There was no effect on bacterial cell viability by exposure to heating at 65°C, whereas the viability was significantly reduced when the bacterial cells were irradiated for 60 seconds in the presence of the photosensitizer. The rise in temperature in the periapical lesion and surrounding area of the *in vitro* model during laser irradiation was monitored using a thermotracer (Figure 3(C)). There was no change in temperature in either the lesion or surrounding area caused by irradiation when the photosensitizer was not added to the bacterial cell suspension (Figures 3(D)–3(F)). However, the thermotracer indicated an increase in temperature in the periapical lesion area with addition of the photosensitizer (Figures 3(G)–3(I)), which was 2°C after irradiation for 60 seconds and 6°C after that for 120 seconds (Figure 3(J)). In contrast, there was no rise in temperature observed in the area surrounding the lesion area even with addition of the photosensitizer (Figures 3(G)–3(I)).

4. Discussion

It is known that various photosensitizers are taken up by tumor cells and microorganisms, and those activated by laser irradiation interact with oxygen to produce radical species that have a toxic effect on those cells and microorganisms [25]. In the present study, indocyanine green, which was a photosensitizer commonly used as fluorescent fundus contrast medium that does not have toxicity toward surrounding tissues, increased the antimicrobial effects of laser irradiation




FIGURE 3: Effects of heat produced by laser irradiation. (A) Change of inner temperature in apical lesion area of *in vitro* model following laser irradiation. (B) Antimicrobial effects of heat on *E. faecalis*. The microorganisms were exposed to diode laser irradiation in the presence of a photosensitizer for 60 seconds [5 W-60 s/PS(+)] or exposed to heat at 65°C for 60 seconds (65°C-60 s). (C). Photograph of temperature change monitoring using a thermotracer, which was used to monitor 4 points (a, b, c, d) of the *in vitro* model. (D–F). Temperature change without photosensitizer. (G–I). Temperature change with photosensitizer. (J) Temperature change in periapical lesion area of the model during irradiation (closed diamond, with photosensitizer; closed square, without photosensitizer). The rise in temperature was calculated as the difference in temperature between points a and b. Data are expressed as the mean \pm standard deviation of triplicate determinations. The experiment was performed three times, with similar results obtained in each.

in test tubes. However, that environment is different from that of a periapical lesion, which is surrounded by sound periapical tissues, and it takes many efforts to make a standardized apical lesion model *in vivo*.

To simulate biological conditions, we prepared an *in vitro* apical lesion model and examined the antimicrobial effects of diode laser irradiation with or without the addition of a photosensitizer. When a bacterial cell suspension with indocyanine green was irradiated with the diode laser using periapical lesion model, *E. faecalis* CFU were significantly decreased to the same amount following irrigation with NaOCl. In a study of bactericidal activity against *E. faecalis* biofilm in extracted human teeth, it was reported that the total energy output of a diode laser was 36J [26]. In the present study, an irradiation time of 60 seconds was employed to achieve a total energy level of 76J and that with an added photosensitizer was adequate for a high level of sterilization. On the other hand, application of the present diode laser for 60 seconds increased the temperature of the lesion area to 65°C. However, exposure of *E. faecalis* to 65°C for 60 seconds in a heat block did not have any effects on bacterial viability. These results indicate that the reduction in viability of *E. faecalis* was from the laser irradiation itself, not from the heat produced by the irradiation, and that addition of a photosensitizer is essential for the observed antimicrobial effects.

Adverse effects of intense heat produced by laser irradiation on sound periapical tissues are a concern in clinical applications. It has been reported that use of a diode laser resulted in the least amount of temperature increase among several types of lasers tested [19]. In the present study, it was found that irradiation at 5 W for 60 seconds in the presence of a photosensitizer raised it by 2°C in the lesion defect area, whereas no change in temperature was observed in the surrounding area. Furthermore, it has been demonstrated that the same diode laser irradiation with various levels of laser power (0.5 W to 5 W) for 2 minutes did not affect proliferation of mammalian cells *in vitro* and enhanced BMP-induced osteoblast differentiation by stimulating BMP/Smad signaling pathway [27]. Taken together, the present laser technique, which was adequate for elimination of *E. faecalis*, may have no cytotoxic effects on sound periapical tissues.

In conclusion, diode laser irradiation in combination with a photosensitizer had nearly the same antimicrobial effect as 2.5% NaOCl. Many clinicians prefer a diluted concentration to reduce the irritation potential of NaOCl, with 2.5% commonly recommended [28]. However, fibroblasts were found damaged by 2.5% NaOCl [3, 5, 28–30]. The present results indicated no cytotoxic effects from heat produced by laser irradiation, thus use of a diode laser with a photosensitizer is useful for treatment of periapical lesions without adverse effects on surrounding tissues. This standardized *in vitro* apical lesion model needs more improvement, but it may be useful for an endodontic study.

Acknowledgments

This investigation was supported in part by a Grant-in-Aid for Scientific Research (21592428) from The Ministry

of Education, Culture, Sports, Science and Technology of Japan.

References

- [1] T. W. Chow, "Mechanical effectiveness of root canal irrigation," *Journal of Endodontics*, vol. 9, no. 11, pp. 475–479, 1983.
- [2] J. F. Siqueira Jr., I. N. Rôças, S. R. Santos, K. C. Lima, F. A. Magalhães, and M. de Uzeda, "Efficacy of instrumentation techniques and irrigation regimens in reducing the bacterial population within root canals," *Journal of Endodontics*, vol. 28, no. 3, pp. 181–184, 2002.
- [3] A. Gatot, J. Arbelles, A. Leiberman, and I. Yanai-Inbar, "Effects of sodium hypochlorite on soft tissues after its inadvertent injection beyond the root apex," *Journal of Endodontics*, vol. 17, no. 11, pp. 573–574, 1991.
- [4] E. J. Neaverth and R. Swindle, "A serious complication following the inadvertent injection of sodium hypochlorite outside the root canal system," *Compendium*, vol. 11, no. 8, pp. 474–481, 1990.
- [5] M. Nagayoshi, C. Kitamura, T. Fukuizumi, T. Nishihara, and M. Terashita, "Antimicrobial effect of ozonated water on bacteria invading dentinal tubules," *Journal of Endodontics*, vol. 30, no. 11, pp. 778–781, 2004.
- [6] N. Noguchi, Y. Noiri, M. Narimatsu, and S. Ebisu, "Identification and localization of extraradicular biofilm-forming bacteria associated with refractory endodontic pathogens," *Applied and Environmental Microbiology*, vol. 71, no. 12, pp. 8738–8743, 2005.
- [7] Y. Noiri, A. Ehara, T. Kawahara, N. Takemura, and S. Ebisu, "Participation of bacterial biofilms in refractory and chronic periapical periodontitis," *Journal of Endodontics*, vol. 28, no. 10, pp. 679–683, 2002.
- [8] F. J. Vertucci, "Root canal anatomy of the human permanent teeth," *Oral Surgery Oral Medicine and Oral Pathology*, vol. 58, no. 5, pp. 589–599, 1984.
- [9] R. Fujii, Y. Saito, Y. Tokura, K. I. Nakagawa, K. Okuda, and K. Ishihara, "Characterization of bacterial flora in persistent apical periodontitis lesions," *Oral Microbiology and Immunology*, vol. 24, no. 6, pp. 502–505, 2009.
- [10] K. Subramanian and A. K. Mickel, "Molecular analysis of persistent periradicular lesions and root ends reveals a diverse microbial profile," *Journal of Endodontics*, vol. 35, no. 7, pp. 950–957, 2009.
- [11] P. T. Sunde, I. Olsen, G. J. Debelian, and L. Tronstad, "Microbiota of periapical lesions refractory to endodontic therapy," *Journal of Endodontics*, vol. 28, no. 4, pp. 304–310, 2002.
- [12] M. Wainwright, "Photodynamic antimicrobial chemotherapy (PACT)," *Journal of Antimicrobial Chemotherapy*, vol. 42, no. 1, pp. 13–28, 1998.
- [13] M. Wainwright and K. B. Crossley, "Photosensitising agents—circumventing resistance and breaking down biofilms: a review," *International Biodeterioration and Biodegradation*, vol. 53, no. 2, pp. 119–126, 2004.
- [14] E. B. de Souza, S. Cai, M. R.L. Simionato, and J. L. Lage-Marques, "High-power diode laser in the disinfection in depth of the root canal dentin," *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontology*, vol. 106, no. 1, pp. e68–e72, 2008.
- [15] J. A. Williams, G. J. Pearson, and M. John Colles, "Antibacterial action of photoactivated disinfection {PAD} used on endodontic bacteria in planktonic suspension and in artificial

- and human root canals,” *Journal of Dentistry*, vol. 34, no. 6, pp. 363–371, 2006.
- [16] L. Zhu, M. Tolba, D. Arola, M. Salloum, and F. Meza, “Evaluation of effectiveness of Er,Cr:YSGG laser for root canal disinfection: theoretical simulation of temperature elevations in root dentin,” *Journal of Biomechanical Engineering*, vol. 131, no. 7, Article ID 710041, 2009.
- [17] M. Kreisler, W. Kohnen, M. Beck et al., “Efficacy of NaOCl/H₂O₂ irrigation and GaAlAs laser in decontamination of root canals *in vitro*,” *Lasers in Surgery and Medicine*, vol. 32, no. 3, pp. 189–196, 2003.
- [18] U. Schoop, W. Kluger, S. Dervisbegovic et al., “Innovative wavelengths in endodontic treatment,” *Lasers in Surgery and Medicine*, vol. 38, no. 6, pp. 624–630, 2006.
- [19] U. Schoop, W. Kluger, A. Moritz, N. Nedjelic, A. Georgopoulos, and W. Sperr, “Bactericidal effect of different laser systems in the deep layers of dentin,” *Lasers in Surgery and Medicine*, vol. 35, no. 2, pp. 111–116, 2004.
- [20] J. L. Fimple, C. R. Fontana, F. Foschi et al., “Photodynamic treatment of endodontic polymicrobial infection *in vitro*,” *Journal of Endodontics*, vol. 34, no. 6, pp. 728–734, 2008.
- [21] A. Silva Garcez, S. C. Núñez, J. L. Lage-Marques, A. O. C. Jorge, and M. S. Ribeiro, “Efficiency of NaOCl and laser-assisted photosensitization on the reduction of *Enterococcus faecalis in vitro*,” *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontology*, vol. 102, no. 4, pp. e93–e98, 2006.
- [22] N. S. Soukos, P. S. Y. Chen, J. T. Morris et al., “Photodynamic therapy for endodontic disinfection,” *Journal of Endodontics*, vol. 32, no. 10, pp. 979–984, 2006.
- [23] R. M. Love, “*Enterococcus faecalis*—a mechanism for its role in endodontic failure,” *International Endodontic Journal*, vol. 34, no. 5, pp. 399–405, 2001.
- [24] I. N. Rôças, J. F. Siqueira Jr., and K. R. N. Santos, “Association of *Enterococcus faecalis* with different forms of periradicular diseases,” *Journal of Endodontics*, vol. 30, no. 5, pp. 315–320, 2004.
- [25] K. Konopka and T. Goslinski, “Photodynamic therapy in dentistry,” *Journal of Dental Research*, vol. 86, no. 8, pp. 694–707, 2007.
- [26] Z. Lim, J. L. Cheng, T. W. Lim et al., “Light activated disinfection: an alternative endodontic disinfection strategy,” *Australian Dental Journal*, vol. 54, no. 2, pp. 108–114, 2009.
- [27] S. Hirata, C. Kitamura, H. Fukushima et al., “Low-level laser irradiation enhances BMP-induced osteoblast differentiation by stimulating the BMP/Smad signaling pathway,” *Journal of Cellular Biochemistry*, vol. 111, no. 6, pp. 1445–1452, 2010.
- [28] G. L. Becker, S. Cohen, and R. Borer, “The sequelae of accidentally injecting sodium hypochlorite beyond the root apex: report of a case,” *Oral Surgery Oral Medicine and Oral Pathology*, vol. 38, no. 4, pp. 633–638, 1974.
- [29] R. E. Beltz, M. Torabinejad, and M. Pouresmail, “Quantitative analysis of the solubilizing action of MTAD, sodium hypochlorite, and EDTA on bovine pulp and dentin,” *Journal of Endodontics*, vol. 29, no. 5, pp. 334–337, 2003.
- [30] I. Heling, I. Rotstein, T. Dinur, Y. Schwec-Levine, and D. Steinberg, “Bactericidal and cytotoxic effects of sodium hypochlorite and sodium dichloroisocyanurate solutions *in vitro*,” *Journal of Endodontics*, vol. 27, no. 4, pp. 278–280, 2001.

Apoptosis and survivability of human dental pulp cells under exposure to Bis-GMA

Junya YANO¹, Chiaki KITAMURA¹, Tatsuji NISHIHARA², Masayuki TOKUDA³, Ayako WASHIO¹, Ker-Kong CHEN⁴, Masamichi TERASHITA⁵

1- DDS, PhD, Division of Pulp Biology, Operative Dentistry, and Endodontics, Department of Cariology and Periodontology, Kyushu Dental College, Japan.

2- DDS, PhD, Division of Infections and Molecular Biology, Department of Health Promotion, Kyushu Dental College, Japan.

3- DDS, PhD, Department of Restorative Dentistry and Endodontology, Kagoshima University Graduate School of Medical and Dental Sciences, Japan.

4- DDS, PhD, Department of Conservative Dentistry, Kaohsiung Medical University Hospital and College of Dental Medicine, Kaohsiung Medical University, Kaohsiung, Taiwan.

5- DDS, PhD, Division of Comprehensive Dentistry, Department of Clinical Communication Practice, Kyushu Dental College, Japan.

Corresponding address: Chiaki Kitamura, DDS, PhD - Division of Pulp Biology, Operative Dentistry and Endodontics, Department of Cariology and Periodontology, Kyushu Dental College, 2-6-1 - Manazuru - Kokurakita - Kitakyushu - 803-8580 - Japan - Phone: 81-93-582-1131 - Fax: 81-93-581-5399 - e-mail: chi-aki-k@kyu-dent.ac.jp

Received: June 19, 2009 - Modification: March 19, 2010 - Accepted: March 27, 2010

ABSTRACT

Objective: In the present study, we examined whether 2, 2-bis [4-(2-hydroxy-3-methacryloxypropoxy) phenyl] propane (Bis-GMA) has effects on LSC2 cells, human dental pulp cell line. Material and Methods: The viability, cell cycle, and morphology of LSC2 cells were analyzed after exposure to several different concentrations of Bis-GMA. The recovery of viability of Bis-GMA exposed cells was also analyzed in the condition without Bis-GMA. Further, penetration of Bis-GMA to dentin disc was examined using isocratic high-performance liquid chromatography. Results: There was a concentration-dependent decrease in cell proliferation and an increase in cell number in the sub-G1 population after exposure to Bis-GMA. Furthermore, the cells showed typical characteristics of apoptotic cells after the exposure to high concentration of Bis-GMA. In contrast, cells exposed to lower concentrations of Bis-GMA recovered their viability after being cultured without Bis-GMA. We also found that Bis-GMA is capable of penetrating 1-mm-thick dentin discs, though the penetrated concentration was lower than that showing cytotoxicity. Conclusion: These results suggest that Bis-GMA has cytotoxic effects, though dental pulp exposed to lower concentrations is able to recover their viability when Bis-GMA is removed.

Key words: Apoptosis. Bis-GMA. Cell viability. Dental pulp.

INTRODUCTION

During restorative procedures for a carious tooth, dental pulp is exposed to a variety of stimulations such as heat stress produced by cavity preparation², and hypoxia produced by local anesthesia¹.

When composite resin is used as a restorative material, dentists sometimes note inflammatory responses of dental pulp, which are considered to be caused by microleakage at the interface between cavity walls and filled composite resin^{8,10}. Other studies have suggested that unpolymerized resin monomers remaining on the cavity floor may irritate dental pulp cells directly or indirectly through dentin, resulting in inflammatory responses^{9,12}.

It has previously been reported that apoptosis of

pulp cells was induced during pulp wound healing after cavity preparation⁵, and that apoptosis of pulp cell line was induced by heat stress⁶ and hypoxia¹³. It was also found that capping agents modified pulp apoptosis induction⁷. Apoptotic cells induced in dental pulp disappear during pulp wound healing processes, suggesting cytotoxic effects of external stimulations via residual dentin and the recovery of dental pulp viability. However, it is largely unknown whether dental pulp cells adversely affected by resin monomers can recover their viability. In the present study, we examined effects of resin monomer, 2, 2-bis [4-(2-hydroxy-3-methacryloxypropoxy) phenyl] propane (Bis-GMA), on the viability of human dental pulp cell line.

MATERIAL AND METHODS

Cell culture

LSC2 cells, a dental pulp cell line derived from human dental pulp¹¹, were maintained in Dulbecco's modified Eagle's medium (DMEM; Invitrogen Corp., Carlsbad, CA, USA) containing 10% heat-inactivated fetal calf serum (FCS), 100 µg/mL of streptomycin, and 100 U/mL of penicillin, and then incubated in a humidified atmosphere of 5% CO₂ at 37°C.

Cell proliferation assay

LSC2 cells (2x10⁴ cells/well) were cultured in 96 well plate with a medium containing 5% FCS for 24 h, then the medium was exchanged to that containing 1% FCS. Several different concentrations (0.01, 0.02, 0.03, and 0.04 mmol/L) of Bis-GMA (Polyscience Inc., Warrington, PA, USA) diluted by dimethylsulfoxide (DMSO; Wako Pure Chemical Inc, Osaka, Japan) were added to the culture medium. At 48 h after exposure to Bis-GMA, cell proliferation was examined using the 5-bromo-2'-deoxyuridine (BrdU) assay (The Biotrak cell proliferation ELISA system version 2; GE Healthcare UK Ltd, UK). Cell viability was analyzed by measuring optical density (OD) using a test wavelength of 540 nm and a reference wavelength of 620 nm with a multiscan biochromatic microplate reader (Multiscan JX; Thermo Fisher Scientific K.K., Yokohama, Japan).

Cell cycle analysis

LSC2 cells were exposed to Bis-GMA for 48 h, then suspended in a hypotonic solution (0.1% Triton X-100, 1 mmol/L Tris/HCl, pH 8.0, 3.4 mmol/L sodium citrate, 0.1 mmol/L EDTA) and stained with 5 µg/mL of propidium iodide, after which cell-cycle distribution was analyzed with a FACScalibur flow cytometer EPICS XL (Beckman Coulter Inc., Brea, CA, USA).

Detection of cell death by hoechst staining

LSC2 cells were exposed to Bis-GMA in a chamber slides for 48 h, then fixed with 1% glutaraldehyde for 30 min, washed with 1x phosphate buffered saline (PBS), and stained with 5 g/mL of Hoechst dye 33342 (Life Technologies Co., CA, USA). Cells detached from the coverslips into the medium were also recovered by centrifugation and stained. Nuclei were observed under a fluorescence microscope (OLYMPUS BX50/BX-FLA/DP70; Olympus Co., Tokyo, Japan).

Cell recovery after removal of Bis-GMA from culture medium

After exposure of LSC2 cells to Bis-GMA for 48 h, the wells were rinsed by PBS, and added a medium containing 10% FCS without Bis-GMA. After the culture of surviving cells in a humidified atmosphere

of 5% CO₂ at 37°C for 96 h, cell proliferation and cell cycle were analyzed.

Dentin penetration assay

Dentin discs (1 mm thick, minimum diameter: 6 mm) were cut from freshly extracted sound human third molars using a low speed diamond-coated saw (Isomet; Buehler, Lake Bluff, IL, USA) under water coolant, then etched for 60 s with 35% phosphoric acid to completely open dentinal tubules, rinsed with ultrasonication for 3 min in distilled water, and placed in the PFA tube (Sanplatec, Osaka, Japan). The lower chamber was filled with distilled water and the upper chamber with 0.2 mmol/L of Bis-GMA. After standing in a humid chamber at room temperature for 2 weeks, the distilled water was collected and mixed with chloroform to isolate Bis-GMA. Bis-GMA was detected using isocratic high-performance liquid chromatography (HPLC) (Waters, Milford, MA, USA) with a reverse-phase column (µ-Bondashere C18, 5 µm, 300 A column, JLC010032Y) (Waters) at 280 nm. The peak area




Figure 1- Cytotoxic effects of 2, 2-bis [4-(2-hydroxy-3-methacryloxypropoxy) phenyl] propane (Bis-GMA) on LSC2 cells. (A). Viability of LSC2 cells after exposure to Bis-GMA. (B). Cell cycle of LSC2 cells after exposure to Bis-GMA




Figure 2- Morphological changes of LSC2 cells promoted by 2, 2-bis [4-(2-hydroxy-3-methacryloxypropoxy) phenyl] propane (Bis-GMA). (A). Phase-contrast microphotographs of LSC2 cells after exposure to Bis-GMA. Original magnification, x100. (B). Fluorescent microphotographs of LSC2 cells after exposure to Bis-GMA. Original magnification, x400

of Bis-GMA shown by HPLC tracing was analyzed using the Waters 2487 Dual λ absorbance detector (Waters), and the amount of Bis-GMA was calculated from the calibration curve of known concentrations from control experiments.

Statistical analysis

Statistical analysis of the data was performed using one-way ANOVA followed by a multiple-comparison Sheffe's test with Statview 5.0 software (SAS software Inc., Cary, NC, USA). Statistical significance was determined at $p < 0.05$.

RESULTS

Cell death of human dental pulp cells after exposure to Bis-GMA

The viability of LSC2 cells decreased after exposure to Bis-GMA in a dose-dependent manner (Figure 1A). There was no significant difference ($p > 0.05$) in viability between 0 and 0.01 mmol/L of Bis-GMA, whereas significant decreases of cell viability were observed after being cultured with 0.02, 0.03, and 0.04 mmol/L of Bis-GMA ($p < 0.01$). The effects of exposure to Bis-GMA on cell cycle progression of LSC2 cells are shown in Figure 1B.

Exposure to Bis-GMA reduced the number of the cells in G1, S, and G2/M phases in a dose-dependent manner, with a concomitant increase of those in the sub-G1 population, implying cell death such as apoptosis.

Morphological changes of LSC2 cells following exposure to Bis-GMA are shown in Figure 2. The structures of those cells exposed to 0.01 mmol/L of Bis-GMA were nearly the same as those of the control, whereas the rounding and the detachment were observed in cells exposed to 0.04 mmol/L of Bis-GMA (Figure 2A). Fluorescent microscopic analysis revealed that the nuclei of adhesive cells had normal structures, whereas those of cells that had fallen into the medium after exposure to 0.04 mmol/L of Bis-GMA showed fragmentation, a typical characteristic of apoptotic cells (Figure 2B).

Recovery of cell viability after the removal of Bis-GMA

Following exposure to Bis-GMA, surviving cells attached to the bottom of the well were cultured in normal condition without Bis-GMA (Figure 3A). The proliferation of surviving cells following exposure to 0.01 and 0.02 mmol/L of Bis-GMA was not different after 24 h of cell culture in normal condition, and




Figure 3- Survival of LSC2 cells against 2, 2-bis [4-(2-hydroxy-3-methacryloxypropoxy) phenyl] propane (Bis-GMA). (A). Recovery of viability of LSC2 cells that survived after exposure to Bis-GMA. Concentrations of Bis-GMA (mmol/L); 0 (black circle), 0.01 (white circle), 0.02 (white rectangle), 0.03 (black triangle), and 0.04 (white triangle). (B). Cell cycle of LSC2 cells that survived after exposure to Bis-GMA

then increased at 48 and 96 h, while surviving cells exposed to 0.03 mmol/L of Bis-GMA showed proliferation at 96 h. In contrast, exposure to 0.04 mmol/L of Bis-GMA showed great cytotoxicity on the viability of the cells, and proliferation was not recovered during the culture in the normal condition. The recovery of cell-cycle progression in surviving cells exposed to 0.01, 0.02, and 0.03 mmol/L of Bis-GMA was also observed (Figure 3B). When Bis-GMA-exposed cells were cultured in normal condition without Bis-GMA for 96 h, the cell cycle distribution pattern of Bis-GMA-insulted cells returned to that prior to exposure to Bis-GMA.

Dentin penetration of Bis-GMA through dentinal tubules

In the penetration model, Bis-GMA penetrated through 1 mm thick dentin disc. From the calibration




Figure 4- Calibration curve of known concentrations of 2, 2-bis [4-(2-hydroxy-3-methacryloxypropoxy) phenyl] propane (Bis-GMA) and concentrations of Bis-GMA that penetrated 1 mm dentin discs. White rectangles indicate known concentrations of control Bis-GMA for the calibration curve. Black rectangles indicate concentrations calculated from HPLC results of penetrated Bis-GMA

curve of known concentrations of Bis-GMA, the average concentration of penetrated Bis-GMA was 0.0043 mmol/L (Figure 4).

DISCUSSION

In the present study, it was found that exposure to high concentrations of Bis-GMA induced an increase in cell number in the sub-G1 population in cell-cycle, and the cell death with typical structures of apoptosis, suggesting that direct exposure to a high concentration of Bis-GMA induced apoptosis in human dental pulp cells.

It was also examined whether LSC2 cells exposed to Bis-GMA can recover their viability after the removal of Bis-GMA. We cultured surviving cells after exposure to Bis-GMA in the normal medium without Bis-GMA, and found that the cells exposed to 0.01, 0.02, and 0.03 mmol/L of Bis-GMA showed the recovery of the cell proliferation, whereas those exposed to 0.04 mmol/L of Bis-GMA did not recover their proliferation. In addition, cell-cycle of surviving cells after exposure to 0.01, 0.02, and 0.03 mmol/L of Bis-GMA returned to the same level as the control. These results suggest that even after the exposure to Bis-GMA, LSC2 cells have an ability to recover their viability when the concentration of Bis-GMA is low, whereas the cells exposed to high concentration of Bis-GMA lose such recovery ability.

Another finding of the present study was that the average concentration of Bis-GMA that penetrated 1 mm thick dentin was 0.0043 mmol/L, which was lower than the concentration used in the present study. These results are consistent with previous studies^{3,4}, and suggest that an adequate volume of

residual dentin thickness would be able to suppress the penetration of a harmful concentration of Bis-GMA.

Taken together, the findings of the present study suggest that dental pulp cells have the tolerance against lower concentrations of resin monomers, and that polymerized composite resin on the sufficient thickness of residual dentin may not be harmful to dental pulp.

CONCLUSION

These results produced evidence that Bis-GMA has cytotoxic effects, though dental pulp exposed to lower concentrations is able to recover their viability when Bis-GMA is removed.

ACKNOWLEDGEMENTS

This research was supported by Grants in Aid for Scientific Research 18209057 and 20592231 from The Ministry of Education, Science, and Culture of Japan, Tokyo, Japan.

REFERENCES

- 1- Agata H, Kagami H, Watanabe N, Ueda M. Effect of ischemic culture conditions on the survival and differentiation of porcine dental pulp-derived cells. *Differentiation*. 2008;76:981-93.
- 2- Amano T, Muramatsu T, Amemiya K, Kubo K, Shimono M. Responses of rat pulp cells to heat stress *in vitro*. *J Dent Res*. 2006;85:432-5.
- 3- Bouillaguet S, Wataha JC, Hanks CT, Ciucchi B, Holz J. *In vitro* cytotoxicity and dentin permeability of HEMA. *J Endod*. 1996;22:244-8.
- 4- Hanks CT, Wataha JC, Parsell RR, Strawn SE, Fat JC. Permeability of biological and synthetic molecules through dentine. *J Oral Rehabil*. 1994;21:475-87.
- 5- Kitamura C, Kimura K, Nakayama T, Toyoshima K, Terashita M. Primary and secondary induction of apoptosis in odontoblasts after cavity preparation of rat molars. *J Dent Res*. 2001;80:1530-4.
- 6- Kitamura C, Nishihara T, Ueno Y, Chen K-K, Morotomi T, Yano J, et al. Effects of sequential exposure to lipopolysaccharide and heat stress on dental pulp cells. *J Cell Biochem*. 2006;99:797-806.
- 7- Kitamura C, Ogawa Y, Morotomi T, Terashita M. Differential induction of apoptosis by capping agents during pulp wound healing. *J Endod*. 2003;29:41-3.
- 8- Lu Y, Liu T, Li H, Pi G. Histological evaluation of direct pulp capping with a self-etching adhesive and calcium hydroxide on human pulp tissue. *Int Endod J*. 2008;41:643-50.
- 9- Mantellini MG, Botero TM, Yaman P, Dennison JB, Hanks CT, Nör JE. Adhesive resin induces apoptosis and cell-cycle arrest of pulp cells. *J Dent Res*. 2003;82(8):592-6.
- 10- Murray PE, Smyth TW, About I, Remusat R, Franquin JC, Smith AJ. The effect of etching on bacterial microleakage of an adhesive composite restoration. *J Dent*. 2002;30:29-36.
- 11- Rikitake Y. immortalization of human dental pulp cells with transfecting of the plasmid, pMT1-neo. *Kokubyo Gakkai Zasshi*. 1989;56:540-61.
- 12- Schweikl H, Spagnuolo G, Schmalz G. Genetic and cellular toxicology of dental resin monomers. *J Dent Res*. 2006;85:870-7.
- 13- Ueno Y, Kitamura C, Terashita M, Nishihara T. Re-oxygenation improves hypoxia-induced pulp cell arrest. *J Dent Res*. 2006;85:824-8.

Promoting effects of thymosin β 4 on granulation tissue and new bone formation after tooth extraction in rats

Kou Matsuo, DDS, PhD,^a Yosuke Akasaki, DDS,^b Kazutaka Adachi, DDS,^b
Min Zhang, MD, PhD,^c Ayataka Ishikawa, DDS, PhD,^c Eijiro Jimi, DDS, PhD,^d
Tatsuji Nishihara, DDS, PhD,^e and Ryuji Hosokawa, DDS, PhD,^f Kitakyushu, Japan
KYUSHU DENTAL COLLEGE

Objectives. The aim of this study was to evaluate the effects of thymosin β 4 (TB4) on wound healing after tooth extraction in rats.

Study design. After extraction of the rats' mandibular first molar teeth, a synthetic partial peptide of TB4 was injected intraperitoneally at the time of extraction and every day thereafter for 6 days. Control subjects for the treatment received identical amounts of phosphate-buffered saline solution in the same manner. Histologic analysis, apoptosis assay, and reverse-transcription polymerase chain reaction were performed.

Results. The overall data showed that TB4 treatment suppressed apoptosis and inflammation; it accelerated the process of wound healing, including new bone formation.

Conclusions. The findings demonstrated not only the usefulness of the TB4 partial peptide in wound healing of tooth extraction sockets, but also its potential application for bone regeneration and osteogenesis in bone and bone-related tissues. (*Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2011;xx:xxx)

Thymosin was originally isolated from calf thymus and was considered to be a thymic hormone consisting of a single polypeptide,¹⁻³ but it is now known to consist of a mixture of small polypeptides.⁴⁻⁶ Thymosins are now divided into 3 main groups, i.e., α -, β -, and γ -thymosins, according to their isoelectric points. The β -thymosins (TBs), whose isoelectric points range from 5.0 to 7.0, are a family of highly conserved polar 5-kDa polypeptides consisting of 40-44 amino acids (AA).^{5,6} To date, 15 TB have been isolated from various vertebrates and invertebrates. Thymosin β 4 (TB4) is the most abundant type of TB in most mammalian tissues, representing ~70%-80% of the total TB content. The complete TB4 amino acid sequence of 43 residues is invariant among most known mammalian sources, such as human, rat, mouse, cat, calf, horse, pig, sheep, and chicken, with the exception of rabbit.⁵ Thymosins β 10

(TB10) and β 15 (TB15) are also present in human, rat, and mouse tissues.

TBs are often referred to as "G-actin-sequestering peptides," meaning that TBs bind monomeric globular actin (G-actin) in a 1:1 complex and prevent polymerization into actin filaments.⁵⁻⁸ Besides being simple actin monomer-buffering proteins, recent studies, especially regarding TB4, have revealed that TBs are multifunctional polypeptides involved in angiogenesis,⁹⁻¹¹ wound healing,¹²⁻¹⁸ antiapoptotic activity,¹⁸⁻²⁰ and antiinflammatory reactions.^{13,21} Nonetheless, there have been very few reports of TBs relating directly to the fields of dental research or clinical dentistry.²²⁻²⁴

Based on the background described above, we initiated a basic experiment to investigate the functional analysis of TB4 in wound healing after tooth extraction, with the ultimate aim of clinical dental application, including regenerative therapy. This report shows a new functional aspect of TB4, including its possible role in bone regeneration and osteogenesis.

MATERIAL AND METHODS

All protocols for the present experiment were reviewed and approved by the Council on Animal Care at Kyushu Dental College (approval number 07-051).

Synthetic peptide

A 20-AA partial peptide was synthesized (Sigma Genosys, Hokkaido, Japan), corresponding to AAs 17-36 in the full-length (AAs 1-43) TB4 polypeptide^{5,6} (Fig. 1).

Supported by a grant-in-aid for Scientific Research (C) 20592179 from the Japan Society for the Promotion of Science (to K.M.), a grant-in-aid from Kyushu Dental College Internal Grants (to K.M.), and a grant-in-aid from Kyushu Dental College Alumni Association Grants (to K.M.).

^aAssociate Professor, Department of Biosciences.

^bResident, Department of Oral Functional Reconstruction.

^cAssistant Professor, Department of Biosciences.

^dProfessor, Department of Biosciences.

^eProfessor, Department of Health Promotion.

^fProfessor, Department of Oral Functional Reconstruction.

Received for publication Apr. 27 2011; accepted for publication May 28, 2011.

1079-2104/\$ - see front matter

© 2011 Mosby, Inc. All rights reserved.

doi:10.1016/j.tripleo.2011.05.025




Fig. 1. Amino acid sequences and activities of thymosin β 4 and its partial peptides.

Table 1. PCR primer sets used in this study

Gene (accession number)*	Forward primer sequences (5'-3')	Reverse primer sequences (5'-3')	Size (bp)
<i>G3PDH</i> (M17701)	TGAAGGTCGGGTGCAACGGATTGGC	CATGTAGGCCATGAGGTCCACCAC	983
<i>VEGF-A</i> (NM_031836)	GCCCATGAAGTGGTGAAGTT	ACTCCAGGGCTTCATCATTG	172
<i>VEGF-B</i> (NM_053549)	AGGGAGGTGGTGGTACCTCTGAGCA	TGGTCTGCATTACATTGGCTGTGT	231
<i>VEGF-C</i> (NM_053653)	GATGTGGGGAAAGGAGTTTGGAGCAG	ATTGTGACTGGTTTGGGCCTTGAG	188
<i>VEGF-D</i> (AF014827)	TTGGGGCTTCAAACCTTGTCTCTGG	ACTGAAGCCCTGCACCAGGTACACA	173
<i>Angpt1</i> (NM_053546)	AAGGCGAGTGTGGCAGTACAATGA	CCCTTTGCAGAGCGTTGGTGTGTGA	229
<i>Angpt2</i> (NM_134454)	TGCAGATAAATCCCGGGACCACATT	TGTCAAGCACAAGACGGAACAACGA	168
<i>VEGFR-1</i> (NM_019306)	CATCGGCCATCATCTGAATGTGGTT	GCTGACACTGTCTAGCGGGGTTTC	232
<i>VEGFR-2</i> (U93306)	ATTGGGCAGATGCCTTTGGAATCG	ACCCAGCAGGTTCCACCATTGAGA	165
<i>VEGFR-3</i> (NM_053652)	GTCATTGACACCGGGGACAGCCTAT	GCCCTCAATCCGGGCCTTGATATAC	249
<i>Tie2</i> (NM_001105737)	CAGGGCAAAAATGAAGACCAGCACA	CCTCCAAGGTCTTTAGGGGCTGGAA	191
<i>MMP2</i> (NM_031054)	GGAGAAGGCTGTGTTCTTCG	CGGGTCCATTTTCTTCTCA	214
<i>ILK</i> (NM_133409)	AATGTCCCCGGCTCAGGATTTTCTC	TTGGCTCTGGTCCACAACGAAATTG	167
<i>HGF</i> (NM_017017)	GCATGATGTGGGGACCAAACCTTCT	GCCCTTGTTTCTGATGCACCTGTTG	242
<i>TGF-β1</i> (NM_021578)	AAAGAAGTCACCCGCGTGCTAATGG	CCAAGTAACGCCAGGAATTGTTGC	233
<i>IL-1α</i> (NM_017019)	ATCCGCAGCTTCCCAGAGCTGTTA	ATGCCATGCGAGTACTTAGGACGA	200
<i>IL-1β</i> (NM_031512)	GGGATCCTCTCCAGTCAGGCTTCTC	TGCTCTCATCTGGACAGCCCAAGTC	186
<i>IL-6</i> (NM_012589)	GCCCACCAGGAACGAAAGTCAACTC	TGGTCTGTTGTGGGTGGTATCCTCTG	179
<i>TNF-α</i> (NM_012675)	TCCTTCCTTCCCTCCCTCAGA	GTTGGCCTGAAGACAGCTTCCCAAC	209
<i>BMP-2</i> (NM_017178)	TTTGGCCACGACGGTAAAGGACAT	AGTTCAGGTGATCAGCCAGGGGAAA	218
<i>BMP-4</i> (NM_012827)	GGGGAGGAGGAGGAAGAAGAGCAGA	CACCTCATTCTCTGGGATGCTGCTG	186
<i>BMP-7</i> (XM_001053727)	CCAGAACCCTCAAGACTCCAAG	AGGGAAGGCACACTCTCCCTCACAG	199
<i>Osterix</i> (AY177399)	CCCAGCATGTCTTCCACCCAAAGATGT	CCAGCCGCTCTAGCTCCTGACAGTT	186
<i>ALP</i> (NM_013059)	GACAAGAAGCCCTTACAGCCATCC	GCCATGACGTGGGGGATGTAGTTCT	239
<i>Osteocalcin</i> (NM_013414)	AGCAGGAGGGCAGTAAGGTGGTGAA	ATGCCCTAAACGGTGGTCCATAGA	196

*Accession number of GenBank. The primer sequences were designed based on rat mRNA sequences obtained from GenBank. *G3PDH*, glyceraldehyde-3-phosphate dehydrogenase; *VEGF*, vascular endothelial growth factor; *Angpt*, angiopoietin; *VEGFR*, vascular endothelial growth factor receptor; *Tie*, tyrosine kinase with Ig-like loops and epidermal growth factor homology domains; *MMP*, matrix metalloproteinase; *ILK*, integrin-linked kinase; *HGF*, hepatocyte growth factor; *TGF*, transforming growth factor; *IL*, interleukin; *TNF*, tumor necrosis factor; *BMP*, bone morphogenetic protein; *ALP*, alkaline phosphatase; *bp*, base pair.

Animals

Ninety-six 6-week-old male Wistar rats (Charles River Laboratories, Yokohama, Japan) were used. After tooth extraction, they were given a powdered diet and water ad libitum.

Tooth extraction and administration of thymosin β 4

Under pentobarbital anesthesia, mandibular first molars were extracted using custom-made dental forceps (Seiwa Riken, Tokyo, Japan). The rats received an




Fig. 2. Representative hematoxylin-eosin (A and B) and terminal deoxynucleotide transferase–mediated dUTP nick-end labeling (TUNEL) (C and D) stained sections of tooth sockets on day 1 from the control (A and C) and TB4-treated (B and D) groups. On day 1, the control group socket is filled mostly with a blood clot (A). A slightly greater amount of granulation tissue is noted in the bottom of the socket in the TB4-treated group (B). A lower number of TUNEL-positive apoptotic cells are observed in the TB4-treated group (D) than in the control group (C). The TUNEL specimens were counterstained with methyl green.




Fig. 3. Number of apoptotic cells (A) and percentage of granulation tissue area (GA) relative to the tooth extraction socket area (TA) (B). Data are expressed as mean and SD. Significant differences between control and TB4-treated groups are indicated by asterisks (* $P < .05$; ** $P < .01$; *** $P < .001$).

intraperitoneal injection of TB4 synthetic peptide solution (50- μ g peptide in 300 μ L phosphate-buffered saline solution [PBS], pH 7.4) immediately after tooth

extraction and every day thereafter for 6 days. Control subjects received an equivalent amount of PBS intraperitoneally under the same schedule.




Fig. 4. Representative phosphotungstic acid hematoxylin (PTAH) staining (A and B) and immunostaining for osterix (C and D) on day 3 from control (A and C) and TB4-treated (B and D) groups. GA, granulation tissue area; BA, blood clot area; PA, periodontal ligament area. The dashed line (*) denotes the upper limit of the tooth extraction socket area (TA). Arrows indicate the border between GA and BA. Arrowheads indicate the border between PA and GA or PA and BA. Accelerated formation of granulation tissue on day 3 in the TB4-treated group is seen (B) compared with the control group (A). Osterix-positive cells are mainly observed on day 3 at the bottom of the socket (C). More osterix-positive cells have spread into the center of the GA from the bottom of the socket in the TB4-treated group (D). The immunostained sections were counterstained with methyl green.

Histologic and immunohistochemical analyses

Treated rats were killed under pentobarbital anesthesia on days 1, 2, 3, 4, and 6 after tooth extraction. The mandibles were removed, fixed in 10% neutral-buffered formalin overnight, and then decalcified in 10% EDTA containing 7% sucrose for up to 21 days. The completely decalcified mandibles were sectioned

vertically in the buccolingual direction to obtain the maximum exposed surface of the socket corresponding to the distal root of the extracted tooth. The samples were then paraffin embedded and sectioned at a thickness of 5 μm . These serial sections were stained with hematoxylin-eosin (H-E) and phosphotungstic acid hematoxylin (PTAH). Immunohisto-




Fig. 5. Number of osterix-positive cells (A) and percentage of newly formed bone trabeculae area (TBA) relative to the tooth extraction socket area (TA) (B). Data are expressed as mean and SD. Significant differences between control and TB4-treated groups are indicated by asterisks (* $P < .05$; ** $P < .01$).

chemical staining was performed using anti-Sp7/osterix rabbit polyclonal antibody (ab22552, 1:200; Abcam, Cambridge, U.K.) and antiperiostin rabbit polyclonal antibody (ab14041, 1:150; Abcam) as the first antibodies. Histofine Simple Stain Rat Max-PO (Multi) (Nichirei Biosciences, Tokyo, Japan) was used as the second antibody. Detection was performed with the Impact DAB peroxidase substrate kit (Vector Laboratories, Burlingame, CA). Nuclear counterstaining was carried out with the use of either methyl green or hematoxylin.

The digital images of the histologic sections were analyzed by Adobe Photoshop CS4 Extended (Adobe Systems, San Jose, CA) and ImageJ, version 1.44a (National Institutes of Health, Bethesda, MD) software. To simplify the calculation of the area, the tooth extraction socket area (TA) was defined as the area bounded by a curve along the medial border of the alveolar bone and a straight line (upper limit of TA) connecting the 2 ends of the curve. The granulation tissue area (GA) was distinguished from the blood clot area (BA) and periodontal ligament area (PA). BA was defined as the area stained deep blue by PTAH. PA was confirmed by immunohistochemical staining for periostin on the serial sections. The percentage of GA relative to TA was calculated for each PTAH-stained section. When we assessed granulation tissue formation, we did not exclude areas of newly formed bone trabeculae or osteoid tissue from GA. For the assessment of new bone and osteoid formation in the tooth extraction socket, the percentage of newly formed bone trabecular area (BTA) relative to TA was calculated from the digital images of the H-E sections. The BTA, including osteoid, was confirmed by immunostaining for osterix on serial

sections. For further assessment of new bone and osteoid formation, all cells positively immunostaining for osterix were counted on the digital images.

Apoptosis assay based on terminal deoxynucleotide transferase-mediated dUTP nick-end labeling method

Serial unstained sections from the paraffin-embedded tissues were used for apoptosis assay with the ApoMark Apoptosis Detection Kit (Exalpa Biologicals, Maynard, MA) according to the manufacturer's protocol. Apoptotic cells were counted on the digital images.

Preparation of total RNA and reverse-transcription polymerase chain reaction

Immediately after the rats were killed, at either 9 hours or 3 days after tooth extraction, granulation tissue was isolated from the socket with a 27-gauge needle, and then total RNAs were prepared from this with the use of the SV total RNA isolation system (Promega, Madison, WI).

Reverse-transcription polymerase chain reaction (RT-PCR) was performed using the Titanium One-Step RT-PCR kit (Clontech, Mountain View, CA) according to the manufacturer's instructions. The primers used in our study are summarized in Table I. The RT-PCR products after 25-35 PCR cycles were electrophoresed on 1%-2% agarose gel, visualized with ethidium bromide staining under ultraviolet light exposure, photographed using a digital camera (PowerShot A640; Canon, Tokyo, Japan), and stored as digital images. The integrated density (IntDen) of each band was calculated with the use of ImageJ. The relative mRNA levels of target genes were expressed as a percentage of




Fig. 6. Representative hematoxylin-eosin sections from days 4 (A and B) and 6 (C and D) in the control (A and C) and TB4-treated (B and D) groups. A greater amount of newly formed bone trabeculae is noted in the TB4-treated group on day 4 (B) than in the control group (A). On day 6, the socket in the TB4-treated group (D) still shows a greater amount of bone trabeculae than in the control group (C).

the ratio of target gene *IntDen* relative to glyceraldehyde-3-phosphate dehydrogenase *IntDen*. Analysis was performed 4 times using different batches of total RNA samples. The results are shown as mean values with standard deviations ($n = 4$).

Data analysis

The statistical significance of differences in the mean values between the 2 groups was assessed by Student *t* test if variances were equal as determined by an *F* test, otherwise by Welch *t* test. All tests were 2 tailed, with differences reported as significant when $P < .05$.

RESULTS

Comparative histologic findings in the healing process after tooth extraction

The extraction sockets on day 1 in the control group were filled mostly with blood clots, although remaining periodontal tissue and only a very small amount of granulation tissue were observed at the bottom. In contrast, there was a slightly greater amount of granulation tissue in the bottom of sockets in the TB4-treated group. In addition, the TB4-treated group had fewer apoptotic cells than the control group (Figs. 2 and 3). On day 2, the sockets in the control group still contained only a small amount of granulation tissue at the




Fig. 7. Representative data from reverse-transcription polymerase chain reaction using total RNAs from granulation tissue in the tooth sockets after 9 hours and 3 days. ND, Not detected by amplification of up to 35 cycles. For abbreviations of target genes, see Table I.

bottom, while a greater amount of granulation tissue was present in the TB4-treated group, with or without a small amount of bone matrix-like tissue. Although the granulation tissue grew rapidly in both groups from days 3 to 4, the TB4-treated group always showed a greater amount of granulation tissue than the control group (Figs. 3 and 4). Furthermore, a greater amount of osteoid tissue and new bone trabeculae appeared in the TB4-treated group on days 3 and 4 than in the control group (Figs. 4-6). Immunohistochemical staining with osterix revealed more osterix-positive granulation tissue in the TB4-treated animals than in control animals from days 2 to 4 (Figs. 4 and 5). The TB4-treated group had a greater amount of bone trabeculae on day 6 than the control group (Figs. 5 and 6).

Differential expression of mRNA between thymosin β 4-treated and control groups

Compared with the control group, the TB4-treated group showed increased expression of the following genes associated with angiogenesis, cell proliferation, and migration on day 3: vascular endothelial growth factor B (*Vegf-B*), *Vegf-C*, *Vegf-D*, *angiopoietin-1* (*Angpt1*), *Angpt2*, *VEGF receptor 1* (*Vegfr-1*), *Vegfr-2*, *Vegfr-3*, *tyrosine kinase with Ig-like loops and epidermal growth factor homology domains 2* (*Tie2*), *matrix metalloproteinase 2* (*Mmp2*), *integrin-linked kinase* (*Ilk*), *hepatocyte growth factor* (*Hgf*), and *transforming growth factor β 1* (*Tgf- β 1*) (Fig. 7; Table II). However, the expression level of *Vegf-A* in the TB4-treated group was lower than in the control group. The expression levels of representative genes related to inflammation, such as *interleukin-1 α* (*Il-1 α*), *Il-1 β* , *Il-6*, and *tumor necrosis factor α* (*Tnf- α*), were significantly decreased in the TB4-treated group than in the control group. With regard to genes related to osteogenesis, significantly higher expression levels of *osterix* and *bone morphogenetic protein 4* (*Bmp-4*) were observed on day 3 in the TB4-treated group than in the control group.

There were significant differences between the 2 groups regarding gene expression levels at 9 hours after tooth extraction (Fig. 7; Table II). In terms of angiogenesis, only 3 genes, *Vegf-B*, *Vegfr-1*, and *Vegfr-2*, were significantly expressed more in the TB4-treated group than in the control group. Regarding inflammatory cytokines, *Il-6* and *Tnf- α* were significantly decreased in the TB4-treated group than in the control group. Most osteogenesis-related genes were not detected by amplification of up to 35 cycles. Interestingly, *Bmp-7* was significantly expressed more in the TB4-treated group 9 hours after extraction than in the control group, although no significant difference for samples on day 3 was noted between the 2 groups.

Table II. Summary of reverse-transcription polymerase chain reaction (RT-PCR) results

Target gene	Day 3 RNAs, ratio to G3PDH (SD)		9-h RNAs, ratio to G3PDH (SD)	
	Control (n = 4)		TB4 (n = 4)	
<i>VEGF-A</i>	92.4 (20.2)	>**	43.7 (8.8)	89.0 (20.2) NS
<i>VEGF-B</i>	52.4 (8.0)	<***	150.1 (12.7)	16.5 (4.1) <**
<i>VEGF-C</i>	9.6 (2.2)	<*	40.2 (15.9)	ND —
<i>VEGF-D</i>	13.4 (3.6)	<*	41.6 (16.6)	25.7 (4.3) NS
<i>Angpt1</i>	17.8 (6.5)	<*	41.3 (13.4)	ND —
<i>Angpt2</i>	13.9 (3.2)	<*	48.9 (19.4)	ND —
<i>VEGFR-1</i>	14.3 (3.3)	<*	58.3 (23.1)	18.1 (1.0) <*
<i>VEGFR-2</i>	20.6 (3.6)	<***	93.1 (24.4)	24.2 (7.7) <*
<i>VEGFR-3</i>	17.9 (3.5)	<***	66.5 (10.6)	ND —
<i>Tie2</i>	13.5 (5.8)	<*	75.3 (24.9)	ND —
<i>MMP2</i>	88.7 (46.1)	<***	287.2 (68.9)	ND —
<i>ILK</i>	73.5 (20.5)	<***	187.3 (36.6)	66.1 (14.3) NS
<i>HGF</i>	29.7 (4.3)	<***	189.2 (4.7)	ND —
<i>TGF-β1</i>	106.0 (49.4)	<***	331.4 (80.3)	23.9 (5.4) NS
<i>IL-1α</i>	294.6 (22.7)	>***	115.7 (6.9)	282.5 (9.2) NS
<i>IL-1β</i>	302.7 (16.3)	>***	152.6 (16.7)	207.2 (12.4) NS
<i>IL-6</i>	87.5 (24.5)	>*	28.6 (5.6)	182.6 (22.0) >**
<i>TNF-α</i>	74.5 (22.1)	>*	29.7 (5.3)	118.8 (18.9) >**
<i>BMP-2</i>	25.4 (3.8)	NS	35.5 (7.2)	ND —
<i>BMP-4</i>	18.7 (4.2)	<*	56.5 (14.3)	ND —
<i>BMP-7</i>	26.4 (8.9)	NS	25.3 (13.2)	18.0 (4.2) <*
<i>Osterix</i>	30.9 (3.9)	<***	53.6 (8.7)	ND —
<i>ALP</i>	58.1 (12.3)	NS	67.3 (13.4)	41.1 (5.2) NS
<i>Osteocalcin</i>	41.4 (8.4)	NS	45.8 (11.1)	ND —

Ratios of target gene to G3PDH expressed as mean percentage values.

NS, No significant difference; ND, not detected by RT-PCR (up to 35 cycles). For abbreviations of target genes, see Table I.

Significant differences between control and TB4-treated groups are indicated by asterisks: * $P < .05$; ** $P < .01$; *** $P < .001$.

DISCUSSION

The full-length TB4 is composed of 43 AA residues. Its functional sequence motif and certain biologic activities of TB4-derived partial peptides have been reported¹⁹; the main functions are shown in Fig. 1. In this study, we used a 20-AA synthetic peptide that included the 9-AA G-actin-binding motif (LKKTETQEK)⁵⁻⁸ and 7-AA essential sites for angiogenic activity (LKKTETQ),²⁵ because it is not easy to synthesize a full-length TB4 polypeptide owing to the distribution of proline throughout the TB4 sequence. Our 20-AA synthetic peptide included 2 glutamines corresponding to the sites where TB4 is cross-linked to fibrin by factor XIIIa, a transglutaminase.^{26,27} We anticipated that these 2 glutamine residues might increase the local concentration of the TB4 partial peptide near the site of the blood clot in the tooth extraction socket after intraperitoneal injection of the peptide.

Histologic analysis showed that TB4 treatment promoted granulation tissue formation in the tooth socket after extraction (Figs. 2-4). This finding supports the previously reported promoting effects of TB4 on angiogenesis, cell proliferation, and migration,^{9-16,18} because the characteristic histologic feature of granulation tissue is the presence of new small blood vessels (an-

giogenesis); cell proliferation and migration are also essential for granulation tissue formation. Meanwhile, gene expression analysis revealed increased expression of the following genes associated with angiogenesis, cell proliferation, and migration on day 3 in the TB4-treated group: *Vegf-B*, *-C*, and *-D*, *Angpt1* and 2, *Vegfr1*, *-2*, and *-3*, *Tie2*, *Mmp2*, *Ilk*, *Hgf*, and *Tgf-β1* (Fig. 7; Table II). This finding from gene expression analysis corroborates the promoting effect of TB4 on granulation tissue formation shown by histologic analysis (Figs. 3 and 4). Among those genes listed above, TB4 has been reported to up-regulate the expression levels of mRNA and/or proteins of *Mmp2*, *Angpt 2*, *Hgf*, *Ilk*, and *Tgf-β1* in various in vitro and/or in vivo experiments.^{10,28-33} Regarding all other genes under study (*Vegf-B*, *-C*, and *-D*, *Angpt1*, *Vegfr1*, *-2*, and *-3*, and *Tie2*), which are angiogenic factors and their receptors, for the first time the up-regulation of these genes by TB4 is suggested from the findings of our in vivo experiment, although it is unclear whether TB4 directly up-regulated the expression levels of the genes in the granulation tissue on day 3.

On the other hand, only 3 genes from those described above (*Vegf-B* and *Vegfr-1* and *-2*) were up-regulated by TB4 treatment in the extraction socket at 9 hours

after tooth extraction (Fig. 7; Table II). Because even the extraction socket at 9 hours in the TB4-treated group should contain only remaining periodontal tissue and, if at all, a very small amount of granulation tissue, many genes may not be up-regulated by TB4 treatment. Nevertheless, up-regulation of these 3 genes may be convincing evidence for the augmentation of angiogenesis by TB4 administration during the early stage of granulation tissue formation. Recent studies have revealed that VEGF-B acts as a survival factor for many types of cells, such as vascular endothelial cells, pericytes, smooth muscle cells, neurons, and cardiac myocytes, by inhibition of apoptosis via both VEGFR-1 and neuropilin-1.³⁴⁻³⁷ TB4-induced expression of VEGF-B not only may have contributed to angiogenesis but also may have decreased the number of apoptotic cells in the TB4-treated group on day 1 (Figs. 2 and 3).

The full-length TB4 polypeptide has been shown to be effective in reducing inflammation.^{13,15} It is also reported that only the 4-AA, amino-terminal peptide of TB4, known as Ac-SDKP, can block inflammation.³⁸ Other than that, however, there has been no information reported on the active sites of TB4 regarding antiinflammatory activity.¹⁹ We have showed in the present study that the 20-AA TB4 partial peptide suppressed the expression levels of the representative inflammatory cytokines IL-1 α , IL-1 β , IL-6, and TNF- α mRNA (Fig. 7; Table II). This finding suggests that AA residues 17-36 may include other functional sites for antiinflammatory activity. In addition, the down-regulation of *Tnf- α* mRNA 9 hours after extraction when treated by the peptide may have suppressed TNF- α -induced apoptosis and resulted in a decreased number of apoptotic cells in the experimental group (Figs. 2 and 3).

TB4 promotes wound healing in various tissues, such as skin, cornea, and heart.^{12-16,18} There has been, however, no report to date on the effects of TB4 regarding bone regeneration or osteogenesis. Our results show that a greater quantity of osterix-positive cells, osteoid, and bone trabeculae appeared in the TB4-treated group than in the control group (Figs. 4-6). mRNA expression of the osteogenesis-promoting factors BMP-4 and TGF- β 1 was significantly higher in the TB4-treated group on day 3 than in the control group, although there were no significant differences between the 2 groups regarding the expression levels of BMP-2 and BMP-7 mRNA on day 3 (Fig. 7; Table II). Regarding the osteogenic differentiation markers alkaline phosphatase (ALP) and osteocalcin mRNA, there were no significant differences between the 2 groups. Although regarding the comparison of expression levels of osteogenic differentiation markers, it would be reasonable to compare those between days 4 and 6, it was hard to obtain sufficient RNA, especially from the TB4-treated

group, where newly formed bone trabeculae occupied the socket from day 4 onward.

Expression of osteogenesis-related genes, apart from *Bmp7*, at 9 hours showed no significant differences between the 2 groups and was undetectable after 35 PCR cycles (Fig. 7; Table II). Although the up-regulation of *Bmp7* mRNA after 9 hours in the TB4-treated group is an interesting finding, it is hard to speculate what role BMP-7 plays in the early healing process after tooth extraction. From our *in vivo* experiments, it remains unclear whether TB4 affects osteogenic progenitor cells directly or indirectly during osteogenesis. In any case, our findings raise the possibility that TB4 accelerates bone regeneration and osteogenesis during the healing of bone and certain bone-associated tissues.

Taken together, we have shown that the 20-AA partial peptide (AA residues 17-36) has effects on angiogenesis, cell migration, granulation tissue formation, inflammation, and apoptosis similar to the full-length 43-AA TB4 polypeptide. Because the shorter peptide is easier and less expensive to synthesize, the 20-AA partial peptide of TB4 has an advantage regarding cost. Furthermore, our findings implicate a potential therapeutic value for TB4 in treating problems of bone and bone-associated tissues.

REFERENCES

1. Goldstein AL, Guha A, Zatz MM, Hardy MA, White A. Purification and biological activity of thymosin, a hormone of the thymus gland. *Proc Natl Acad Sci U S A* 1972;69:1800-3.
2. Goldstein AL, Slater FD, White A. Preparation, assay, and partial purification of a thymic lymphocytopoietic factor (thymosin). *Proc Natl Acad Sci U S A* 1966;56:1010-7.
3. Klein JJ, Goldstein AL, White A. Enhancement of *in vivo* incorporation of labeled precursors into DNA and total protein of mouse lymph nodes after administration of thymic extracts. *Proc Natl Acad Sci U S A* 1965;53:812-7.
4. Low TL, Goldstein AL. Thymosin fraction 5 and 5A. *Methods Enzymol* 1985;116:219-33.
5. Hannappel E, Huff T. The thymosins. Prothymosin alpha, parathymosin, and beta-thymosins: structure and function. *Vitam Horm* 2003;66:257-96.
6. Huff T, Muller CS, Otto AM, Netzker R, Hannappel E. Beta-Thymosins, small acidic peptides with multiple functions. *Int J Biochem Cell Biol* 2001;33:205-20.
7. Van Troys M, Dewitte D, Goethals M, Carlier MF, Vandekerckhove J, Ampe C. The actin binding site of thymosin beta 4 mapped by mutational analysis. *EMBO J* 1996;15:201-10.
8. Safer D, Elzinga M, Nachmias VT. Thymosin beta 4 and Fx, an actin-sequestering peptide, are indistinguishable. *J Biol Chem* 1991;266:4029-32.
9. Grant DS, Rose W, Yaen C, Goldstein A, Martinez J, Kleinman H. Thymosin beta4 enhances endothelial cell differentiation and angiogenesis. *Angiogenesis* 1999;3:125-35.
10. Malinda KM, Goldstein AL, Kleinman HK. Thymosin beta 4 stimulates directional migration of human umbilical vein endothelial cells. *FASEB J* 1997;11:474-81.
11. Grant DS, Kinsella JL, Kibbey MC, LaFlamme S, Burbelo PD, Goldstein AL, et al. Matrigel induces thymosin beta 4 gene in differentiating endothelial cells. *J Cell Sci* 1995;108:3685-94.