



Supplementary Figure II.




Supplementary Figure III.


A.




B.




C.



D.



Supplementary Figure IV.






Figure 1. (A) Ulcerations in the left colon (arrows), seen on the first colonoscopy. (B) Colonoscopy after antiviral treatment, ulcerations are in phase of remission (arrows).

DISCUSSION

CMV colitis is a rare cause of diarrhea in older adults; it is more commonly seen in people who are immunosuppressed (with the human immunodeficiency virus or after bone marrow transplant), in whom it is often due to a virus reactivation, or in patients with preexisting inflammatory bowel disease.⁴ Nevertheless, although it may be considered an invasive diagnostic test in a frail elderly patient, a sigmoidoscopy with biopsy should be considered as a necessary investigation if culture-negative diarrhea persists. Although some cases of CMV colitis are described in immunocompetent patients, when a diagnosis of CMV colitis is made, screening to exclude the presence of concomitant immunomodulating conditions or inflammatory bowel disease is necessary. In the clinical history of this patient, different coexisting immune-modulating conditions (diabetes mellitus, previous HCV infection, probable essential thrombocythemia) can be identified.

Most cases of CMV infection described in the literature are limited to the left colon, but the infection could theoretically involve all of the digestive tract. In this case, an ulceration was also found in the bulbar duodenum; because the ulcer was bloody, histology was not done, so it was not possible to confirm whether it was a location of CMV infection. Anyway, in a meta-analysis, in which the authors identified 44 cases of CMV bowel infection in immunocompetent patients, the extent of disease was not an independent predictor of survival.³

No conclusive statement regarding the need for specific antiviral treatment can be made from the available data in the literature. Although patients with no associated comorbidities seem to have a good rate of spontaneous remission, a trend for higher mortality has been reported in patients aged 55 and older and in patients with diseases affecting immune responses,³ with a mortality rate of 31.8% in patients aged 55 and older. The patient in the current case belongs to this latter group at high risk of mortality, so it was thought that the antiviral treatment was mandatory. Nevertheless, randomized controlled trials are needed for a more-conclusive answer about antiviral treatment in immunocompetent patients suffering from severe CMV infection.

Margherita Azzini, MD
 Claudia Bozzini, MD, PhD
 Claudio Bellamoli, MD
 Ospedale Civile Maggiore
 Prima Geriatria
 Verona, Italy

ACKNOWLEDGMENTS

We are grateful to Dr. Carlo Manfrini for giving us the images of the colonoscopy.

Conflict of Interest: The editor in chief has reviewed the conflict of interest checklist provided by the authors and has determined that the authors have no financial or any other kind of personal conflicts with this paper.

Author Contributions: Margherita Azzini contributed to the collection of clinical data and the preparation of manuscript. Claudia Bozzini wrote the manuscript. Claudio Bellamoli contributed to the collection of clinical data and reviewed the manuscript.

Sponsor's Role: None.

REFERENCES

- Rafailidis PI, Mourtouzoukou EG, Varbobitis IC et al. Severe cytomegalovirus infection in apparently immunocompetent patients: A systematic review. *Virology* 2008;5:47.
- Lockwood MR, Liddle J, Kitsanta P. Cytomegalovirus colitis—an unusual cause for diarrhoea in an elderly woman. *Age Ageing* 2006;35:198–200.
- Galiatsatos P, Shrier I, Lamoureux E et al. Meta-analysis of outcome of Cytomegalovirus colitis in immunocompetent hosts. *Digestive Dis Sci* 2005;50:609–616.
- Julka K, Surawicz CM. Cytomegalovirus in inflammatory bowel disease: Time for another look? *Gastroenterology* 2009;137:1163–1166.

EFFECTS OF TESTOSTERONE IN OLDER MEN WITH MILD-TO-MODERATE COGNITIVE IMPAIRMENT

To the Editor: Some population-based studies have found that endogenous testosterone levels are associated with general cognitive function,^{1,2} and it has also been reported that testosterone levels are associated with physical and psychological functions, including cognition, in disabled older men,³ but there have been few studies evaluating the effects of testosterone supplementation in men with cognitive impairment, and the results were inconsistent.^{4–7} Thus, additional information is needed for frail or disabled older men with cognitive impairment as the targets of testosterone supplementation. Here, a pilot study to investigate the effect of oral testosterone supplementation for 6 months on

Table 1. Changes in Functional Parameters According to Treatment Group

Functional Parameters	Mean \pm Standard Deviation										P-Value
	Testosterone					Control					
	Baseline	3 Months	6 Months	Difference: 0 to 6 Months	Baseline	3 Months	6 Months	Difference: 0 to 6 Months			
Mini-Mental State Examination	20.2 \pm 4.5	21.8 \pm 4.7	22.6 \pm 6.5*	2.4 \pm 3.1	21.9 \pm 5.3	22.0 \pm 4.6	22.0 \pm 4.1	0.1 \pm 2.7	.03		
Hasegawa Dementia Scale, Revised	17.6 \pm 5.9	18.2 \pm 7.1	20.6 \pm 7.3*	3.0 \pm 4.3	19.6 \pm 5.6	20.1 \pm 7.0	18.8 \pm 7.7	-0.8 \pm 2.3	.02		
Barthel Index	91 \pm 12	89 \pm 17	91 \pm 15	0.5 \pm 7.1	92 \pm 10	91 \pm 10	92 \pm 7	0.4 \pm 7.6	.70		
Vitality Index	9.0 \pm 0.9	9.3 \pm 0.9	7.9 \pm 1.3	-1.1 \pm 1.0	9.0 \pm 1.0	9.4 \pm 1.0	9.4 \pm 0.9	0.4 \pm 1.0	.35		

P-values are based on repeated-measures analysis of variance comparing the 6-month change between the groups.

*P < .05 compared with baseline.

cognitive function in Japanese older men with mild to moderate cognitive decline is reported.

Eleven men with cognitive impairment, mean age 81 ± 6 , receiving long-term care, were assigned to take oral testosterone undecanoate 40 mg daily for 6 months after a breakfast containing 15 to 20 g of fat. The control group of 13 men matched for age and cognitive function were followed without testosterone treatment. Cognitive function was evaluated using the Mini-Mental State Examination (MMSE) and Hasegawa Dementia Scale, Revised (HDS-R) at baseline and at 3 and 6 months. Plasma hormone levels were also measured. The institutional review board approved the study protocol, and all participants or their families gave written informed consent.

At baseline, mean total and free testosterone levels, calculated using the Vermeulen equation,⁸ were 14 ± 4 nmol/L and 246 ± 47 pmol/L, respectively. There were no significant differences between the groups in age, length of education, nutritional parameters, functional parameters, or plasma hormone levels. Fasting plasma testosterone levels in the morning did not change significantly during the study, whereas the post-dose levels increased up to 30 ± 8 nmol/L 6 hours after testosterone administration, as reported previously.⁹ The changes in functional parameters in each group from baseline to 6 months are shown in Table 1. At 3 months, subjects who received testosterone treatment showed a nonsignificant increase in MMSE and HDS-R scores, whereas at 6 months, cognitive scores were significantly greater than at baseline. In the control group, both cognitive scores remained unchanged. The difference between the groups was significant at 6 months. Prostate-specific antigen and liver function were unchanged, and no adverse effects were observed.

No significant changes were observed in basic activities of daily living (ADL) and ADL-related vitality in either group (Table 1), possibly because these scores were preserved in most subjects at baseline; the Barthel Index and Vitality Index¹⁰ were 91 ± 10 (full score = 100) and 9.0 ± 1.0 (full score = 10), respectively.

This preliminary study needs to be confirmed in a randomized controlled trial with a large sample size. Nevertheless, these results indicate the effects of testosterone treatment on cognitive function in frail elderly men.

Shiho Fukai, MD
Masahiro Akishita, MD, PhD
Department of Geriatric Medicine
Graduate School of Medicine
University of Tokyo
Tokyo, Japan

Shizuru Yamada, MD, PhD
Kenji Toba, MD, PhD
Department of Geriatric Medicine
School of Medicine
Kyorin University, Tokyo, Japan

Yasuyoshi Ouchi, MD, PhD
Department of Geriatric Medicine
Graduate School of Medicine
University of Tokyo
Tokyo, Japan

ACKNOWLEDGMENTS

Conflict of Interest: The editor in chief has reviewed the conflict of interest checklist provided by the authors and has determined that the authors have no financial or any other kind of personal conflicts with this paper.

Author Contributions: Shiho Fukai: acquisition of subjects and data, data analysis and interpretation of data, preparation of manuscript. Masahiro Akishita: coordinator of study concept and design, acquisition of subjects and data, data analysis and interpretation of data, preparation of manuscript. Shizuru Yamada: acquisition of subjects and data. Kenji Toba and Yasuyoshi Ouchi: coordinator of study concept and design.

Sponsor's Role: None.

REFERENCES

1. Yaffe K, Lui LY, Zmuda J et al. Sex hormones and cognitive function in older men. *J Am Geriatr Soc* 2002;50:707-712.
2. Moffat SD, Zonderman AB, Metter EJ et al. Longitudinal assessment of serum free testosterone concentration predicts memory performance and cognitive status in elderly men. *J Clin Endocrinol Metab* 2002;87:5001-5007.
3. Fukai S, Akishita M, Yamada S et al. Association of plasma sex hormone levels with functional decline in elderly men and women. *Geriatr Gerontol Int* 2009;9:282-289.
4. Cherrier MM, Matsumoto AM, Amory JK et al. Testosterone improves spatial memory in men with Alzheimer disease and mild cognitive impairment. *Neurology* 2005;64:2063-2068.
5. Tan RS, Pu SJ. A pilot study on the effects of testosterone in hypogonadal aging male patients with Alzheimer's disease. *Aging Male* 2003;6:13-17.
6. Kenny AM, Fabregas G, Song C et al. Effects of testosterone on behavior, depression, and cognitive function in older men with mild cognitive loss. *J Gerontol A Biol Sci Med Sci* 2004;59A:75-78.
7. Lu PH, Masterman DA, Mulnard R et al. Effects of testosterone on cognition and mood in male patients with mild Alzheimer disease and healthy elderly men. *Arch Neurol* 2006;63:177-185.
8. Vermeulen A, Verdonck L, Kaufman JM. A critical evaluation of simple methods for the estimation of free testosterone in serum. *J Clin Endocrinol Metab* 1999;84:3666-3672.
9. Houwing NS, Maris F, Schnabel PG et al. Pharmacokinetic study in women of three different doses of a new formulation of oral testosterone undecanoate, Andriol Testocaps. *Pharmacotherapy* 2003;23:1257-1265.
10. Toba K, Nakai R, Akishita M et al. Vitality Index as a useful tool to assess elderly with dementia. *Geriatr Gerontol Int* 2002;2:23-29.

A CASE OF SELECTIVE SEROTONIN REUPTAKE INHIBITOR-INDUCED RAPID EYE MOVEMENT BEHAVIOR DISORDER

To the Editor: Rapid eye movement (REM) sleep behavior disorder (RBD) is often seen in older patients and is characterized by a loss of normal skeletal muscle atonia during REM sleep.^{1,2} As a result, the disease manifests as nocturnal motor activity consistent with the enactment of dream content, for example grabbing the bed partner in response to a dream about falling from a cliff. RBD often results in injury to the patient, bed partner, or both.^{1,2}

In perhaps up to two-thirds of cases, RBD is associated with neurodegenerative disorders, most notably the alpha-synucleinopathies (Parkinson's disease, Lewy body disease, multiple systems atrophy), often antedating other manifestations of these disorders by many years.¹⁻⁴ Other cases seem to be idiopathic, although it has been suggested that various medications, notably selective serotonin reuptake inhibitors (SSRI) and other antidepressants, may commonly

induce RBD.^{1,4,5} In spite of this assertion, there have been few supporting case reports.^{5,6} The authors recently cared for a man who clearly developed RBD as a result of SSRI treatment; the use of the SSRI for posttraumatic stress disorder (PTSD) complicated the clinical picture.

CASE REPORT

An 87-year-old male World War II veteran had been treated for PTSD with associated nightmares but no nocturnal motor activity with bupropion and lorazepam. Past medical history was significant only for essential hypertension. In 1998, after many years of treatment, sertraline was added because of increasing symptoms. Within 6 months of adding sertraline, the patient developed frequent nocturnal motor behavior consistent with the content of his dreams and nightmares, for example punching and choking his wife in the context of a dream about being in a fight. As a result, he and his wife had suffered lacerations and contusions. Other behaviors included running out of his bedroom or running into a window. Upon awakening, he was able to recall portions of the dreams but was unaware of the motor behaviors.

Trials of temazepam, zolpidem, and trazodone were ineffective in improving these behaviors. Ultimately, a diagnosis of RBD was made based on the clinical presentation. Clonazepam 1 mg at bedtime was added, which resulted in a moderate decrease in the frequency of the nocturnal motor activity, from nightly to two or three times per week. After 3 months, sertraline was slowly tapered and discontinued, which resulted in a complete cessation of all nocturnal motor behavior. He remained free of nocturnal motor activity for 5 months, until sertraline was inadvertently restarted after the loss of his wife. Within 1 month of restarting sertraline, the nocturnal motor behavior returned. There has thus far been no evidence of dementia or of parkinsonism.

This patient's clinical presentation was typical of RBD; unfortunately, his and his wife's injuries were also typical. It seems clear that his RBD was SSRI induced; it developed after sertraline was started, did not definitively improve until it was stopped, and recurred after it was inadvertently restarted, and there was no evidence of parkinsonism or dementia over the previous 12 years. Although there are few published cases of SSRI-induced overt RBD, increased electromyography activity during REM sleep has been demonstrated in patients taking SSRIs. (None of the patients were being treated for PTSD.)⁷

The relationship between RBD and PTSD is complex and not fully investigated. There is clinical and polysomnographic evidence of greater motor activity during REM sleep in patients with PTSD,⁸ and greater prevalence of RBD was noted in a cohort of patients with PTSD.⁹ SSRIs are effective for PTSD-related nightmares¹⁰ but may cause RBD, clonazepam is effective for RBD^{1,2,4} but not for PTSD-related nightmares,¹⁰ and RBD is not associated with the typical diurnal symptoms of PTSD. In spite of his long history of PTSD and related nightmares, this patient had never exhibited any significant motor activity during sleep until the SSRI was started.

RBD is relatively common in geriatric practice and should be explored in any patient with nocturnal injuries or motor activity. RBD responds well to treatment, generally with clonazepam. Discontinuation of SSRIs or changing to

ORIGINAL ARTICLE: EPIDEMIOLOGY,
CLINICAL PRACTICE AND HEALTH

Effects of dehydroepiandrosterone supplementation on cognitive function and activities of daily living in older women with mild to moderate cognitive impairment

Shizuru Yamada,¹ Masahiro Akishita,² Shiho Fukai,² Sumito Ogawa,²
Kiyoshi Yamaguchi,² Jun Matsuyama,³ Koichi Kozaki,¹ Kenji Toba¹ and
Yasuyoshi Ouchi²

¹Department of Geriatric Medicine, Kyorin University School of Medicine, ²Department of Geriatric
Medicine, Graduate School of Medicine, The University of Tokyo, Tokyo, and ³Medical Corp.
Shojyu-kai, Matsuyama Hospital, Gunma, Japan

Aim: There is little evidence that dehydroepiandrosterone (DHEA) has beneficial effects on physical and psychological functions in older women. We investigated the effect of DHEA supplementation on cognitive function and ADL in older women with cognitive impairment.

Methods: A total of 27 women aged 65–90 years (mean \pm standard deviation, 83 ± 6) with mild to moderate cognitive impairment (Mini-Mental State Examination, MMSE; 10–28/30 points), receiving long-term care at a facility in Japan were enrolled. Twelve women were assigned to receive DHEA 25 mg/day p.o. for 6 months. The control group ($n = 15$) matched for age and cognitive function was followed without hormone replacement. Cognitive function was assessed by MMSE and Hasegawa Dementia Scale-Revised (HDS-R), and basic activities of daily living (ADL) by Barthel Index at baseline, 3 and 6 months. Plasma hormone levels including testosterone, DHEA, DHEA-sulfate and estradiol were also followed up.

Results: After 6 months, DHEA treatment significantly increased plasma testosterone, DHEA and DHEA-sulfate levels by 2–3-fold but not estradiol level compared to baseline. DHEA administration increased cognitive scores and maintained basic ADL score, while cognition and basic ADL deteriorated in the control group (6-month change in DHEA group vs control group; MMSE, $+0.6 \pm 3.2$ vs -2.1 ± 2.2 , $P < 0.05$; HDS-R, $+2.8 \pm 2.8$ vs -0.3 ± 4.1 , $P < 0.05$; Barthel Index, $+3.7 \pm 7.1$ vs -2.7 ± 4.6 , $P = 0.05$). Among the cognitive domains, DHEA treatment improved verbal fluency ($P < 0.05$).

Correspondence: Dr Masahiro Akishita MD PhD, Department
of Geriatric Medicine, Graduate School of Medicine, The
University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo
113-8655, Japan. Email: akishita-tky@umin.ac.jp

Accepted for publication 10 March 2010.

Conclusion: DHEA supplementation in older women with cognitive impairment may have beneficial effects on cognitive function and ADL. *Geriatr Gerontol Int* 2010; 10: 280–287.

Keywords: activities of daily living, cognitive function, dehydroepiandrosterone.

Introduction

Dehydroepiandrosterone (DHEA) and its sulfate (DHEA-S) are the most abundant circulating steroids mainly produced by the adrenal zona reticularis in both sexes.¹ Their circulating levels decline with advancing age,^{1–4} and there has been growing public interest in DHEA supplementation to prevent age-associated physical and cognitive impairment. DHEA is considered a crucial precursor of human sex steroid biosynthesis, and to exert indirect androgenic and estrogenic effects following conversion into smaller amounts of testosterone and estradiol.^{5,6} While this conversion contributes to a part of testosterone production in men, its role may be much more significant in postmenopausal women whose ovarian production of androgen and estrogen has waned. Importantly, postmenopausal women with intact ovaries continue to produce androgens; DHEA(-S), testosterone and androstenedione, while their production of estradiol is minimal.⁷ However, the role of androgens in older women's health is not fully understood.

Clinical trials of the effects of estrogen replacement therapy on cognitive function have shown a lack of efficacy in postmenopausal women initiating hormone replacement therapy after the age of 65 years.^{8,9} On the other hand, previous reports have suggested that DHEA may have neuroprotective effects, and the age-associated DHEA(-S) decline is associated with cognitive impairment in older women.^{2,10–12} One longitudinal study observed lower DHEA-S levels in patients who subsequently developed Alzheimer's disease.¹³ However, controlled trials with DHEA supplementation have failed to show beneficial effects on cognition in healthy middle-aged to older women.^{14–16} In these studies, the participants were limited to those who did not have cognitive impairment; therefore, it is reasonable to hypothesize that DHEA supplementation may be effective in much older women with cognitive decline as well as lower DHEA levels.

Dehydroepiandrosterone deficiency is also considered to be involved in the development of physical frailty.¹⁷ Clinical experience with DHEA supplementation in older women is limited, and the few clinical trials examining its effect on physical function and activity of daily living (ADL) have yielded inconsistent results.^{18–20} Evidence is lacking for much older women in whom physical impairment becomes more apparent and is

accompanied by an age-associated DHEA decline. In our previous study, plasma DHEA and DHEA-S levels, but not estradiol level, were independently related to higher basic ADL in older women aged 70–93 years with functional decline receiving long-term care.²¹ We hypothesized that in older women, DHEA replacement could be effective for the age-related decline of physical as well as psychological function.

This study therefore examined the effect of relatively low-dose (25 mg daily) p.o. DHEA supplementation for 6 months on cognitive function and ADL in older women with cognitive impairment.

Methods

Subjects and study design

In this open, non-randomized controlled study, 27 women aged 65 years or older who attended a health service facility for the elderly (a facility that provides nursing care and rehabilitation services to elderly people with disability, Mahoroba-no-Sato, located in Nagano Prefecture, Japan) were enrolled. The participants were in a chronic stable condition and receiving Long-term Care Insurance service either for admission to the facility or day-care services. The principal inclusion criteria were mild to moderate cognitive decline; both Mini-Mental State Examination (MMSE)²² and Hasegawa Dementia Scale-Revised (HDS-R)²³ scores were between 10 and 28. The subjects were diagnosed as having a mild cognitive impairment²⁴ or Alzheimer's disease according to the Diagnostic and Statistical Manual of Mental Disorders IV.²⁵ The participants had never been treated with hormone replacement therapy, and plasma DHEA-S concentration was less than 3.0 $\mu\text{mol/L}$. The exclusion criteria were history of stroke, extremely low ADL status (Barthel Index²⁶ <50), malnutrition (serum albumin <3.5 mg/dL), malignancy, acute inflammation (fever, white blood cell count >10 000/ μL , or other signs of infection within 4 weeks before enrollment) and overt endocrine diseases, because these diseases may affect both plasma sex hormone levels and functions. None of the subjects were taking a cholinesterase inhibitor (donepezil hydrochloride) or glucocorticoid, opiate or hormone supplement.

Twelve women were assigned to receive DHEA capsule (25 mg/day, Athena Clinics International,

Honolulu, HI, USA) and 15 women were followed up without any additive medication. Medications that could influence cognitive function and plasma hormone levels were not changed during the study period. Outcome measures were cognitive function, ADL, plasma hormone levels, blood cell counts, blood chemical parameters and subjective adverse events. They were assessed at baseline, and after 3 and 6 months. The institutional review board of Mahoroba-no-Sato approved the study protocol, and all participants or their families gave written informed consent.

Hormone measurements

Blood samples were obtained from the participants in the morning after an overnight fast, and plasma hormone levels in addition to blood cell counts and blood chemical parameters were determined by a commercial laboratory (Health Sciences Research Institute, Yokohama, Japan). DHEA and DHEA-S were assayed using sensitive radioimmunoassays with minimum detection limits of 0.04 ng/mL (0.14 nmol/L) and 2.0 µg/dL (0.05 µmol/L), respectively. Total testosterone and estradiol were assayed using chemiluminescent immunoassays with minimum detection limits of 7 ng/dL (0.2 nmol/L) and 4 pg/mL (14.7 pmol/L), respectively. The intra-assay coefficients of variation for these measurements were less than 5%.

Cognitive function

Trained examiners administered two standardized cognitive function tests, MMSE²² and HDS-R,²³ to assess multiple, diverse aspects of cognitive function at baseline and at the 3- and 6-month visits. Both scores range 0–30, with higher scores indicating better performance. HDS-R includes questions about the subject's age, orientation, immediate recall, serial subtraction of 7 s, reciting digits backward, recalling three words, recalling five objects and word fluency (generating names of vegetables). MMSE evaluates five aspects of cognition: (i) orientation; (ii) registration; (iii) attention and calculation; (iv) recall; and (v) comprehension of spoken language (naming objects, spoken language ability, following commands). MMSE, but not HDS-R, includes four performance tests: (i) three-stage command; (ii) reading and following a command; (iii) writing; and (iv) construction drawing). Based on the results of HDS-R and MMSE, we evaluated seven cognitive domains (points) as follows: (i) orientation (10); (ii) verbal memory (9); (iii) attention and calculation (5); (iv) visual memory (5); (v) spoken-language comprehension (9); (vi) verbal fluency (5); and (vii) performance (7).

Other functional parameters and anthropometric measures

Trained nurses and physical therapists visited the participants at the facility and performed the assessments. Basic ADL was assessed by Barthel Index,²⁶ mood by Geriatric Depression Scale (GDS, 15 items),²⁷ and ADL-related vitality by Vitality Index (10-point scale).²⁸ Higher GDS scores indicate a more marked self-reported depressive status, while higher Vitality Index scores indicate greater willingness.

Adverse events

Information regarding adverse events was obtained by questioning or examining the subjects. At each visit during the treatment period, all new complaints and symptoms were recorded. The safety of DHEA supplementation was assessed from the symptoms and by measuring blood chemical parameters including liver and kidney function, electrolyte levels and hematological parameters. Preexisting complaints or symptoms that increased in intensity or frequency during the treatment period also were examined.

Statistical analysis

Data were analyzed using SPSS statistical software ver. 17.0. Changes in outcome measures at 3 and 6 months were calculated by comparing the values at baseline with those at each measurement. Within each group, the significance of the change from baseline to 6 months was tested using paired Student's *t*-test. Repeated-measures ANOVA was used to test the statistical significance of the effects of DHEA versus control. Significance tests were two-sided, with an α -level of 0.05.

Results

Hormone changes and adverse effects

Characteristics and hormone levels at baseline according to treatment groups are shown in Table 1. There were no significant differences between the DHEA group and the control group in age, length of education, nutritional parameters, functional parameters and plasma hormone levels. DHEA supplementation was well tolerated, with high adherence, and there were no detectable adverse events and none of the subjects dropped out during the study. Measures of liver function, kidney function, electrolyte levels and hemoglobin level were not significantly altered by treatment with DHEA (data not shown). Body mass index remained unchanged in both groups.

Subjects in the DHEA group showed a significant increase from baseline to 3 and 6 months in levels of

Table 1 Participant characteristics at baseline

	DHEA	Control
No. of subjects	12	15
Age, years	82 ± 6 (69–90)	83 ± 6 (65–89)
Education, years	8 ± 2	8 ± 2
Nutritional parameters		
Body mass index, kg/m ²	22.0 ± 2.4 (18.8–26.4)	22.4 ± 3.2 (17.6–27.1)
Albumin, g/dL	4.4 ± 0.3 (3.7–4.9)	4.3 ± 3.2 (3.8–4.7)
Total cholesterol, mg/dL	227 ± 39 (166–294)	203 ± 22 (173–250)
Functional parameters		
MMSE	24.0 ± 4.2 (18–28)	23.4 ± 4.4 (14–28)
HDS-R	19.9 ± 5.8 (10–28)	21.7 ± 5.6 (10–28)
Barthel Index	89.6 ± 9.4 (55–100)	89.7 ± 6.4 (75–100)
Vitality Index	9.8 ± 0.6 (8–10)	9.9 ± 0.3 (9–10)
GDS	7.0 ± 4.4 (1–15)	7.0 ± 4.0 (1–13)
Hormones		
DHEA-S, µmol/L	1.8 ± 0.6 (0.7–2.4)	1.6 ± 0.8 (0.3–2.9)
DHEA, nmol/L	7.6 ± 4.7 (2.4–19.1)	6.6 ± 3.1 (2.1–11.5)
Testosterone, nmol/L	1.4 ± 0.4 (0.9–2.3)	1.3 ± 0.9 (0.2–3.8)
Estradiol, pmol/L	88 ± 52 (15–187)	70 ± 26 (45–115)

Values are shown as mean ± standard deviation (range). HDS-R, Hasegawa Dementia Scale-Revised; MMSE, Mini-Mental State Examination; GDS, Geriatric Depression Scale; DHEA-S, dehydroepiandrosterone sulfate; DHEA, dehydroepiandrosterone. There was no significant difference in each parameter between the groups.

circulating DHEA, DHEA-S and testosterone, with levels reaching approximately 2–3-fold higher than those at baseline, whereas the increase in estradiol level was not significant (Table 2). Subjects in the control group showed no significant change in hormone levels.

Changes in cognitive function and ADL

The changes in functional parameters in each group from baseline to 6 months are shown in Table 2. After 6 months, mean HDS-R score significantly improved in the DHEA group while it remained unchanged in the control group. Mean MMSE score significantly declined in the control group while it remained unchanged in the DHEA group. As a result, significant differences were found in these scores between the groups. DHEA treatment maintained Barthel Index score, whereas the score deteriorated significantly during 6 months in the control group, although the between-group difference at 6 months was not statistically significant. Regarding the components of Barthel Index, in the control group, the sum score of mobility deteriorated significantly after 6 months compared to baseline, while no significant change was observed in the sum score of self care (Table 3). Neither Vitality Index nor GDS changed significantly in both groups.

Table 4 shows the cognitive domain scores at baseline and at 3- and 6-month follow up. Among the seven cognitive domains, DHEA treatment improved verbal fluency ($P < 0.05$), resulting in a significant difference at 6 months between the groups. Verbal memory showed a non-significant trend towards improvement in the DHEA group. Performance test scores significantly declined over time in both groups. There were no differences between the groups in the scores of orientation, attention and calculation, visual memory and spoken-language comprehension.

Discussion

Daily administration of DHEA 25 mg for 6 months in elderly women with mild to moderate cognitive impairment improved cognitive function and maintained basic ADL, compared to the control group. Among the cognitive domains, DHEA significantly improved verbal fluency. At baseline, DHEA and DHEA-S levels were lower than those reported in healthy postmenopausal women in both groups,^{2,4} and DHEA treatment increased DHEA, DHEA-S and testosterone levels by 2–3-fold to the mid-normal range for premenopausal

Table 2 Changes in hormone levels and functional parameters by treatment group

	DHEA					Control		P	
	Baseline	3 months	6 months	0-6-month difference	Baseline	3 months	6 months		0-6-month difference
Hormones									
DHEA-S, $\mu\text{mol/L}$	1.8 \pm 0.6	4.5 \pm 1.3*	5.6 \pm 2.9*	3.8 \pm 2.8	1.6 \pm 0.8	1.8 \pm 1.0	1.7 \pm 0.8	-0.02 \pm 0.4	<0.01
DHEA, nmol/L	7.6 \pm 4.7	12.2 \pm 4.8*	13.7 \pm 7.7*	6.1 \pm 8.2	6.6 \pm 3.1	7.3 \pm 3.7	7.4 \pm 4.5	0.9 \pm 2.8	0.04
Testosterone, nmol/L	1.4 \pm 0.4	2.3 \pm 0.7*	2.3 \pm 0.8*	0.9 \pm 0.8	1.4 \pm 0.7	1.4 \pm 0.7	1.6 \pm 0.8	0.2 \pm 0.5	<0.01
Estradiol, pmol/L	88 \pm 52	92 \pm 48	101 \pm 37	13 \pm 51	70 \pm 26	68 \pm 20	67 \pm 42	-4.0 \pm 38	0.17
Functional parameters									
MMSE	24.0 \pm 4.2	24.1 \pm 4.6	24.6 \pm 4.3	0.6 \pm 3.2	23.4 \pm 4.4	23.1 \pm 5.4	21.3 \pm 5.0**	-2.1 \pm 2.2	0.04
HDS-R	19.9 \pm 5.8	20.5 \pm 7.3	22.7 \pm 6.3**	2.8 \pm 2.8	21.7 \pm 5.6	22.1 \pm 5.6	21.3 \pm 6.4	-0.3 \pm 4.1	0.04
Barthel Index	89.6 \pm 9.4	92.7 \pm 6.5	93.3 \pm 6.8	3.7 \pm 7.1	89.7 \pm 6.4	86.9 \pm 7.2	87.0 \pm 6.7*	-2.7 \pm 4.6	0.04
Vitality Index	9.8 \pm 0.6	9.7 \pm 0.5	9.7 \pm 0.7	-0.1 \pm 1.0	9.9 \pm 0.3	9.8 \pm 0.5	9.7 \pm 1.0	-0.3 \pm 1.0	0.80
GDS	7.0 \pm 4.4	6.2 \pm 3.4	6.6 \pm 3.7	-0.4 \pm 1.7	7.0 \pm 4.0	8.3 \pm 3.9	7.5 \pm 3.5	0.5 \pm 3.3	0.60

Values are shown as mean \pm standard deviation (range). P-values are for repeated-measure ANOVA over all three time points: DHEA, dehydroepiandrosterone; HDS-R, Hasegawa Dementia Scale-Revised; MMSE, Mini-Mental State Examination; GDS, Geriatric Depression Scale; DHEA-S, dehydroepiandrosterone sulfate; DHEA, dehydroepiandrosterone. ** $P < 0.01$ compared to baseline, * $P < 0.05$ compared to baseline.

women.² No detectable adverse effects were observed throughout the study.

According to the previous trials, DHEA supplementation of 50 mg or more daily does not provide beneficial effects on cognition in healthy middle-aged to elderly women without cognitive impairment.¹⁴⁻¹⁶ However, in a small-scale randomized double-blind placebo-controlled study, DHEA transiently improved cognition (after 3 months) in subjects with Alzheimer's disease while the improvement was not significant at 6 months.²⁹ Preliminary analysis of the small number of subjects in the present study suggested that DHEA treatment was no less effective in subjects with low baseline cognitive function than those with higher cognitive function (data not shown). Whether the effects of DHEA might be influenced by baseline cognitive function should be further investigated.

It is noteworthy that the 6-month effect of donepezil hydrochloride (5 or 10 mg), the only cholinesterase inhibitor used in Japan, in patients with Alzheimer's disease ranged from no change to less than 1 point improvement in MMSE score,²⁹⁻³³ which is not so different from the effect of DHEA observed in the present study.

In the present study, not only the participants' cognitive function was impaired, but baseline plasma DHEA(-S) level was also low compared to that in postmenopausal or perimenopausal women.^{2,4,10} Regarding DHEA-S levels, according to a report in which healthy pre- and postmenopausal women were studied, DHEA-S levels in women aged 35-44 years and 45-55 years were as follows: 4.31 \pm 2.11, 3.90 (mean \pm standard deviation) and 3.42 \pm 2.01 $\mu\text{mol/L}$.² In this study, DHEA-S was measured using chemiluminescent enzyme immunoassay; although the measurements by this method and those by radioimmunoassay have been reported to be comparable. In our study, DHEA treatment increased DHEA-S levels to the mid-normal range for premenopausal women.² Also, the subjects with lower baseline DHEA-S levels showed non-significant trend towards more improvement in cognitive scores (data not shown). Thus, future studies are needed to explore whether the effects of DHEA might be influenced by baseline DHEA levels.

Because the DHEA receptor has not been identified, DHEA may act after conversion to testosterone and subsequently estradiol through estrogen receptors and androgen receptors, both of which are found in the hippocampus and frontal lobes and subserve verbal memory and working memory in women.^{34,35} Further, hippocampal volume and perfusion have been shown to correlate with serum DHEA-S level in demented patients.^{36,37} It has also been suggested that estrogenic and androgenic derivatives of DHEA might have different effects on cognitive functions.³⁸ However, the mechanism by which DHEA improves cognitive

Table 3 Changes in mobility and self-care scores in Barthel Index during the study

Domains (points)	Mean \pm SD				<i>P</i>
	Baseline	3 months	6 months	Change (0–6 months)	
Mobility (55)					
DHEA	46.9 \pm 9.2	48.2 \pm 6.0	49.2 \pm 5.2	2.3 \pm 5.4	0.01
Control	47.5 \pm 5.4	46.2 \pm 5.5	45.0 \pm 4.3*	-3.7 \pm 3.9	
Self care (45)					
DHEA	42.7 \pm 6.1	44.5 \pm 1.5	43.1 \pm 2.5	0.4 \pm 6.9	0.96
Control	41.8 \pm 4.2	42.5 \pm 3.4	41.2 \pm 4.3	0.7 \pm 3.2	

Mobility is the sum score of five domains: (i) transfer (moving from a bed to a wheelchair and back); (ii) walking on a level surface; (iii) propelling a wheel chair; (iv) ascending and descending stairs; and (v) bathing and toilet use. Self care includes feeding, grooming, dressing, bowels and bladder. *P*-values are for repeated-measure ANOVA over all three time points. **P* < 0.05 compared to baseline. SD, standard deviation.

Table 4 Changes in cognitive domain scores during study

Domains (points)	Mean \pm SD				<i>P</i>
	Baseline	3 months	6 months	Change (0–6 months)	
Orientation (10)					
DHEA	8.3 \pm 1.9	8.0 \pm 2.7	7.5 \pm 3.0	-0.1 \pm 1.2	0.28
Control	8.3 \pm 1.9	8.0 \pm 2.8	7.5 \pm 2.9	-0.7 \pm 1.7	
Verbal memory (9)					
DHEA	5.7 \pm 2.1	6.5 \pm 2.3	6.7 \pm 2.5†	1.0 \pm 1.9	0.79
Control	6.5 \pm 1.7	7.5 \pm 1.8	7.0 \pm 1.9	0.5 \pm 1.7	
Attention and calculation (5)					
DHEA	2.3 \pm 1.9	2.8 \pm 2.0	2.7 \pm 1.8	0 \pm 2.3	0.79
Control	2.0 \pm 1.7	1.9 \pm 1.2	1.8 \pm 1.5	-0.5 \pm 1.4	
Visual memory (5)					
DHEA	3.6 \pm 0.9	3.6 \pm 1.3	3.8 \pm 1.2	0.3 \pm 1.1	0.91
Control	3.6 \pm 1.3	3.9 \pm 0.9	3.9 \pm 1.0	0.5 \pm 1.1	
Language comprehension (9)					
DHEA	8.5 \pm 0.8	7.8 \pm 2.5	8.7 \pm 0.7	0.1 \pm 0.3	0.12
Control	8.5 \pm 0.8	8.5 \pm 0.8	8.4 \pm 1.1	-0.1 \pm 0.9	
Verbal fluency (5)					
DHEA	2.8 \pm 3.3	2.5 \pm 2.0	4.3 \pm 1.1*	1.5 \pm 1.7	0.01
Control	3.2 \pm 1.9	3.8 \pm 1.6	3.3 \pm 1.9	0.1 \pm 2.1	
Performance (7)					
DHEA	5.7 \pm 0.7	5.5 \pm 0.7	4.8 \pm 0.4**	-0.8 \pm 0.6	0.36
Control	5.6 \pm 0.6	5.1 \pm 0.6	4.5 \pm 0.9**	-1.1 \pm 0.8	

Change refers to score change during 0–6 months for each parameter in each treatment group. *P*-values are for repeated-measure ANOVA over all three time points. DHEA, dehydroepiandrosterone. **P* < 0.05, ***P* < 0.01, †*P* < 0.1 vs baseline. SD, standard deviation.

function is unknown. In the present study, plasma estradiol level was not significantly increased after DHEA treatment, implying that its beneficial effects on cognition might be androgen-dependent. Unfortunately, free testosterone levels were not measured, because they were considered to be undetectable in many cases in older women. In addition, sex hormone-binding globulin (SHBG) measurement was not available; however, it has

been reported that DHEA 50 mg treatment for 3 months in postmenopausal women did not significantly change SHBG levels,³⁹ suggesting that the change in SHBG-bound hormone levels after DHEA treatment might be minimal. Given the local aromatization of androgen to estradiol in the brain, the effect of DHEA on cognition might be indirect, complex and heterogeneous. The molecular mechanism underlying the association

between DHEA and cognitive function needs to be clarified, and active forms of testosterone and estradiol should also be examined to investigate whether they would change after DHEA administration.

In our previous study, plasma DHEA and DHEA-S levels were independently related to higher basic ADL in older women aged 70–93 years with functional decline,²¹ and other reports have shown a correlation between DHEA level and muscle mass, strength and physical performance.^{40,41} In the present study, DHEA treatment maintained the Barthel Index score, while the score deteriorated significantly in the control group. Regarding body composition and strength, DHEA administration in postmenopausal older women aged up to 80 years did not alter body composition, physical performance or strength.^{18–20} However, in one small-scale open-label trial, DHEA treatment for 4 weeks improved ADL in three out of seven patients (both men and women) with multi-infarct dementia.⁴² All these studies are preliminary, and large-scale and long-term studies are required to ascertain whether DHEA could have a beneficial effect on ADL in older women.

In the present study, no effect of DHEA on depressive mood or vitality was observed, consistent with most clinical trials in older women.^{15,43,44} This might be attributable to the participants' relatively low depressive status and high vitality status, namely, ceiling effects.

The limitations of our study should be acknowledged. First, this study was neither blinded nor randomized. Second, the number of participants was too small to confirm the results. Thus, results need to be confirmed by large-scale randomized trials to exclude possible selection bias. Third, considering the sensitivity and accuracy, a standard test like the Alzheimer's Disease Assessment Scale should be used in clinical trials to ascertain the effect of DHEA. Finally, our study duration was 6 months so it does not provide any information on the effects of longer-term DHEA supplementation.

In summary, this small study showed that supplementation of DHEA 25 mg for 6 months to older women with mild to moderate cognitive impairment improved cognitive scores and maintained basic ADL. The results should be confirmed in large-scale randomized trials.

Acknowledgments

This study was supported by a Health and Labor Sciences Research Grant (H17-Choju-046) from the Ministry of Health, Labor and Welfare of Japan; Grants-in-Aid for Scientific Research from the Ministry of Education, Science, Culture and Sports of Japan (20249041, 21390220); and grants from the NOVARTIS Foundation for Gerontological Research, Yamaguchi Endocrine Research Association and Mitsui Sumitomo Insurance Welfare Foundation.

References

- Orentreich N, Brind L, Rizer R, Vogelman JH. Age changes and sex differences in serum dehydroepiandrosterone sulfate concentrations throughout adulthood. *J Clin Endocrinol Metab* 1984; **59**: 551–555.
- Davison S, Bell R, Donath S, Montalto J, Davis S. Androgen levels in adult females: changes with age, menopause and oophorectomy. *J Clin Endocrinol Metab* 2005; **90**: 3847–3853.
- Kaufman JM, Vermeulen A. The decline of androgen levels in elderly men and its clinical and therapeutic implications. *Endocr Rev* 2005; **26**: 833–876.
- Tannenbaum C, Barrett-Connor E, Laughlin GA, Platt RW. A longitudinal study of dehydroepiandrosterone sulphate (DHEAS) change in older men and women: the Rancho Bernardo Study. *Eur J Endocrinol* 2004; **151**: 717–725.
- Webb SJ, Geoghegan TE, Prough RA, Michael Miller KK. The biological actions of dehydroepiandrosterone involves multiple receptors. *Drug Metab Rev* 2006; **38**: 89–116.
- Labrie F, Luu-The V, Labrie C, Simard J. DHEA and its transformation into androgens and estrogens in peripheral target tissues: intracrinology. *Front Neuroendocrinol* 2001; **22**: 185–212.
- Arlt W. Androgen therapy in women. *Eur J Endocrinol* 2006; **154**: 1–11.
- Shumaker SA, Legault C, Rapp SR *et al.*, WHIMS Investigators. Estrogen plus progestin and the incidence of dementia and mild cognitive impairment in postmenopausal women: the Women's Health Initiative Memory Study: a randomized controlled trial. *JAMA* 2003; **289**: 2651–2662.
- Shumaker SA, Legault C, Kuller L *et al.* Women's Health Initiative Memory Study. Conjugated equine estrogens and incidence of probable dementia and mild cognitive impairment in postmenopausal women: Women's Health Initiative Memory Study. *JAMA* 2004; **291**: 2947–2958.
- Davis SR, Shah SM, McKenzie DP, Kulkarni J, Davison SL, Bell RJ. Dehydroepiandrosterone sulfate levels are associated with more favorable cognitive function in women. *J Clin Endocrinol Metab* 2008; **93**: 801–808.
- Goldman N, Gleib DA. Sex differences in the relationship between DHEAS and health. *Exp Gerontol* 2007; **42**: 979–987.
- Valle'e M, Mayo W, Le Moal M. Role of pregnenolone, dehydroepiandrosterone and their sulfate esters on learning and memory in cognitive aging. *Brain Res Rev* 2001; **37**: 301–312.
- Hillen T, Lun A, Reischies FM, Borchelt M, Steinhagen-Thiessen E, Schaub RT. DHEA-S plasma levels and incidence of Alzheimer's disease. *Biol Psychiatry* 2000; **47**: 161–163.
- Kritz-Silverstein D, von Mühlen D, Laughlin GA, Bettencourt R. Effects of dehydroepiandrosterone supplementation on cognitive function and quality of life: the DHEA and Well-Ness (DAWN) Trial. *J Am Geriatr Soc* 2008; **56**: 1292–1298.
- Barnhart KT, Freeman E, Grisso JA *et al.* The effect of dehydroepiandrosterone supplementation to symptomatic perimenopausal women on serum endocrine profiles, lipid parameters, and health-related quality of life. *J Clin Endocrinol Metab* 1999; **84**: 3896–3902.
- Wolf OT, Kudielka BM, Hellhammer DH, Hellhammer J, Kirschbaum C. Opposing effects of DHEA replacement in

- elderly subjects on declarative memory and attention after exposure to a laboratory stressor. *Psychoneuroendocrinology* 1998; **23**: 617–629.
- 17 Fried LP, Tangen CM, Walston J *et al*. Cardiovascular Health Study Collaborative Research Group. Frailty in older adults: evidence for a phenotype. *J Gerontol A Biol Sci Med Sci* 2001; **56**: M146–M156.
 - 18 Morales AJ, Haubrich RH, Hwang JY, Asakura H, Yen SS. The effect of six months treatment with a 100 mg daily dose of dehydroepiandrosterone (DHEA) on circulating sex steroids, body composition and muscle strength in age-advanced men and women. *Clin Endocrinol (Oxf)* 1998; **49**: 421–432.
 - 19 Percheron G, Hogrel JY, Denot-Ledunois S *et al*. Double-blind placebo-controlled trial. Effect of 1-year oral administration of dehydroepiandrosterone to 60- to 80-year-old individuals on muscle function and cross-sectional area: a double-blind placebo-controlled trial. *Arch Intern Med* 2003; **163**: 720–727.
 - 20 Nair KS, Rizza RA, O'Brien P *et al*. DHEA in elderly women and DHEA or testosterone in elderly men. *N Engl J Med* 2006; **355**: 1647–1659.
 - 21 Fukai S, Akishita M, Yamada S *et al*. Association of plasma sex hormone levels with functional decline in elderly men and women. *Geriatr Gerontol Int* 2009; **9**: 282–289.
 - 22 Folstein MF, Folstein SE, McHugh PR. "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *J Psychiat Res* 1975; **12**: 189–198.
 - 23 Kato S, Shimogaki M, Onodera H. Revised Hasegawa Dementia Scale (HDS-R). *Jpn J Geriatr Psychiatr* 1991; **2**: 1339–1347.
 - 24 Winblad B, Palmer K, Kivipelto M *et al*. Mild cognitive impairment – beyond controversies, towards a consensus: report of the International Working Group on Mild Cognitive Impairment. *J Intern Med* 2004; **256**: 240–246.
 - 25 American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*, 4th edn. Washington, DC: American Psychiatric Association, 1994.
 - 26 Mahoney FI, Barthel DW. Functional evaluation: Barthel Index. *Md State Med J* 1965; **14**: 61–65.
 - 27 Yesavage JA. Geriatric Depression Scale. *Psychopharmacol Bull* 1988; **24**: 709–711.
 - 28 Toba K, Nakai R, Akishita M *et al*. Vitality Index as a useful tool to assess elderly with dementia. *Geriatr Gerontol Int* 2002; **2**: 23–29.
 - 29 Wolkowitz OM, Kramer JH, Reus VI *et al*. DHEA treatment of Alzheimer's disease: a randomized, double-blind, placebo-controlled study. *Neurology* 2003; **60**: 1071–1076.
 - 30 Winblad B, Engedal K, Soininen H *et al*. Donepezil Nordic Study Group. A 1-year, randomized, placebo-controlled study of donepezil in patients with mild to moderate AD. *Neurology* 2001; **57**: 489–495.
 - 31 Rogers SL, Farlow MR, Doody RS, Mohs R, Friedhoff LT. A 24-week, double-blind, placebo-controlled trial of donepezil in patients with Alzheimer's disease. Donepezil Study Group [see comments]. *Neurology* 1998; **50**: 136–145.
 - 32 Feldman H, Gauthier S, Hecker J, Vellas B, Subbiah P, Whalen E. A 24-week, randomized, double-blind study of donepezil in moderate to severe Alzheimer's disease. *Neurology* 2001; **57**: 613–620.
 - 33 Nozawa M, Ichimiya Y, Nozawa E *et al*. Clinical effects of high oral dose of donepezil for patients with Alzheimer's disease in Japan. *Psychogeriatrics* 2009; **9**: 50–55.
 - 34 Janowsky JS. Thinking with your gonads: testosterone and cognition. *Trends Cogn Sci* 2006; **10**: 77–82.
 - 35 Genazzani AR, Pluchino N, Luisi S, Luisi M. Estrogen, cognition and female ageing. *Hum Reprod Update* 2007; **13**: 175–187.
 - 36 Magri F, Terenzi F, Ricciardi T *et al*. Association between changes in adrenal secretion and cerebral morphometric correlates in normal aging and senile dementia. *Dement Geriatr Cogn Disord* 2000; **11**: 90–99.
 - 37 Magri F, Terenzi F, Ricciardi T *et al*. Hippocampal perfusion and pituitary-adrenal axis in Alzheimer's disease. *Neuropsychobiology* 2000; **42**: 51–57.
 - 38 Hirschman E, Merritt P, Wang CC *et al*. Evidence that androgenic and estrogenic metabolites contribute to the effects of dehydroepiandrosterone on cognition in postmenopausal women. *Horm Behav* 2004; **45**: 144–155.
 - 39 Stomati M, Rubino S, Spinetti A *et al*. Endocrine, neuroendocrine and behavioral effects of oral dehydroepiandrosterone sulfate supplementation in postmenopausal women. *Gynecol Endocrinol* 1999; **13**: 15–25.
 - 40 Valenti G, Denti L, Maggio M *et al*. Effect of DHEAS on skeletal muscle over the life span: the InCHIANTI study. *J Gerontol A Biol Sci Med Sci* 2004; **59**: 466–472.
 - 41 O'Donnell AB, Travison TG, Harris SS, Tenover JL, McKinlay JB. Testosterone, dehydroepiandrosterone, and physical performance in older men: results from the Massachusetts Male Aging Study. *J Clin Endocrinol Metab* 2006; **91**: 425–431.
 - 42 Azuma T, Nagai Y, Saito T, Funauchi M, Matsubara T, Sakoda S. The effect of dehydroepiandrosterone sulfate administration to patients with multi-infarct dementia. *J Neurol Sci* 1999; **162**: 69–73.
 - 43 Wolf OT, Neumann O, Hellhammer DH *et al*. Effects of a two week physiological dehydroepiandrosterone substitution on cognitive performance and well-being in healthy elderly women and men. *J Clin Endocrinol Metab* 1997; **82**: 2363–2367.
 - 44 Arlt W, Callies F, Koehler I *et al*. Dehydroepiandrosterone supplementation in healthy men with an age-related decline of dehydroepiandrosterone secretion. *J Clin Endocrinol Metab* 2001; **86**: 4686–4692.

ORIGINAL ARTICLE

Association of low testosterone with metabolic syndrome and its components in middle-aged Japanese men

Masahiro Akishita¹, Shiho Fukai¹, Masayoshi Hashimoto², Yumi Kameyama¹, Kazushi Nomura¹, Tetsuro Nakamura³, Sumito Ogawa¹, Katsuya Iijima¹, Masato Eto¹ and Yasuyoshi Ouchi¹

Epidemiological studies have shown that low testosterone is associated with metabolic syndrome (MetS) in Caucasian men. We investigated whether testosterone level is related to the prevalence of MetS in middle-aged Japanese men. A cross-sectional survey was conducted in 194 men aged 30–64 years (49 ± 9). Blood sampling was performed in the morning after a 12-h fast, and the relationship between plasma hormone and MetS was analyzed. Low total testosterone was associated with MetS according to the Japanese criteria (HRs of 2.02 by quartile of testosterone; 95% CI=1.43–2.87) and the International Diabetes Federation criteria (HRs of 1.68 by quartile of testosterone; 95% CI=1.25–2.25). Age-adjusted regression analyses revealed that testosterone was significantly related to the MetS parameters of obesity ($\beta=-0.365$ and -0.343 for waist circumference and body mass index, respectively), hypertension ($\beta=-0.278$ and -0.157 for systolic and diastolic blood pressure, respectively), dyslipidemia ($\beta=-0.242$ and 0.228 for triglycerides and high-density lipoprotein cholesterol, respectively), insulin resistance ($\beta=-0.253$ and -0.333 for fasting plasma glucose and homeostasis model assessment of insulin resistance, respectively) and adiponectin ($\beta=0.216$). Inclusion of waist circumference into the model largely weakened the association of testosterone with other metabolic risk factors. In contrast, high estradiol was associated with MetS and its parameters, mostly attributing to the positive correlation between estradiol and obesity. Dehydroepiandrosterone sulfate was not associated with MetS or its parameters. These results suggest that low testosterone is associated with MetS and its parameters in middle-aged Japanese men. The association between estradiol and MetS needs further investigation.

Hypertension Research (2010) 33, 587–591; doi:10.1038/hr.2010.43; published online 26 March 2010

Keywords: androgen; estrogen; insulin resistance; obesity; sex hormone

INTRODUCTION

There is growing awareness that metabolic syndrome (MetS) is one of the most important threats to public health because of its association with type 2 diabetes mellitus, cardiovascular disease and mortality.^{1–3} In men, it is well established that endogenous androgens decline with advancing age,⁴ and low testosterone levels have been associated with insulin resistance,⁵ type 2 diabetes,^{6,7} hypertension⁸ and increased cardiovascular and all-cause mortality.^{9,10} Moreover, men with low testosterone are likely to have more components of MetS in cross-sectional studies,^{11–13} and longitudinal studies show that lower total testosterone predicts higher frequency of MetS.^{14,15} These data were mostly from studies with Caucasian men in western countries. Regarding Japanese men, one study showed that testosterone was positively correlated with plasma adiponectin.¹⁶ However, there are no reports showing a relationship between testosterone and MetS or its components in Japanese men.

Recently, we reported that low testosterone is an independent determinant of endothelial dysfunction in middle-aged men¹⁷ and is

a predictor of cardiovascular events in men with coronary risk factors,¹⁸ suggesting a link between testosterone and cardiovascular pathology. Given these findings, this study investigated the relationship of endogenous testosterone with MetS in middle-aged Japanese men.

METHODS

Subjects

Enrollment screening included consecutive, apparently healthy male subjects aged 30–64 years who underwent medical examinations at either our department or at two clinics located in Tokyo. After exclusion of subjects who met the exclusion criteria, 194 subjects (104 from our department and 90 from the clinics) were enrolled. Exclusion criteria included history of cardiovascular disease (stroke, coronary heart disease, congestive heart failure and peripheral arterial disease), malignancy or overt endocrine disease or use of steroid hormones, because these conditions may influence plasma sex hormones and/or the components of MetS. Other exclusion criteria were diabetic subjects

¹Department of Geriatric Medicine, Graduate School of Medicine, University of Tokyo, Tokyo, Japan; ²Department of General Internal Medicine, Kobe University School of Medicine, Kobe, Japan and ³Research Institute of Aging Science, Tokyo, Japan
Correspondence: Dr M Akishita, Department of Geriatric Medicine, Graduate School of Medicine, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-8655, Japan.
E-mail: akishita-ky@umin.ac.jp

Received 17 November 2009; revised 6 January 2010; accepted 3 February 2010; published online 26 March 2010

on insulin injection or hypoglycemic agent drugs or with hemoglobin A1c >8%, and subjects on β -blockers¹⁹ or fibrates. History, physical examination and laboratory tests were performed for all subjects. Of the included subjects, 23% ($n=44$) were taking anti-hypertensive drugs (angiotensin-converting enzyme inhibitors, angiotensin receptor blockers, calcium channel blockers and diuretics), and 22% were taking statins. Each subject gave written, informed consent before study enrollment. The study protocol was approved by the ethics committee of the Graduate School of Medicine at the University of Tokyo.

Assays of metabolic risk factors and plasma hormones

Clinical information was collected at baseline when each patient attended the initial medical examination. Blood sampling and measurement of height, weight, waist circumference and blood pressure were performed in the morning after a 12-h overnight fast. Blood pressure was measured at least twice using an automated, digital electrophygmomanometer (Omron Healthcare, Kyoto, Japan) on the non-dominant arm in a sitting position, and the average was used for statistical analysis.

Serum total cholesterol and triglyceride were measured enzymatically, and serum high-density lipoprotein (HDL) cholesterol was measured by the heparin- $\text{Ca}^{2+}\text{Ni}^{2+}$ precipitation method. Low-density lipoprotein cholesterol was determined using the Friedewald formula or the direct, liquid, selective detergent method when triglycerides were >400 mg per 100 ml. Plasma glucose was assayed by the glucose oxidase method, and hemoglobin A1c was measured by high-performance liquid chromatography. Plasma total testosterone, dehydroepiandrosterone sulfate and estradiol were determined using sensitive radioimmunoassays. Homeostasis model assessment of insulin resistance (HOMA-IR) was calculated as fasting insulin ($\mu\text{IU ml}^{-1}$) \times fasting plasma glucose (mg per 100 ml)/405. Patients with a fasting plasma glucose >140 mg per 100 ml were excluded from the HOMA-IR calculation because of a lack of data reliability. Serum adiponectin was measured using an enzyme-linked immunosorbent assay (Human Adiponectin ELISA kit, Otsuka Pharmaceutical, Tokyo, Japan). These assays were performed by a commercial laboratory (SRL, Tokyo, Japan). The intra-assay coefficients of variation for the measurements were <5%.

Definition of MetS

We applied both the Japanese criteria²⁰ and the International Diabetes Federation (IDF) criteria for Japanese ethnicity²¹ for the diagnosis of MetS. In the Japanese criteria, MetS was diagnosed when waist circumference ≥ 85 cm and two or more of the following three components were present: (1) HDL cholesterol <40 mg per 100 ml and/or triglyceride ≥ 150 mg per 100 ml; (2) systolic blood pressure ≥ 130 mm Hg and/or diastolic blood pressure ≥ 85 mm Hg and (3) fasting plasma glucose ≥ 110 mg per 100 ml. Subjects taking anti-hypertensive medications were considered hypertensive for statistical purposes.

In the IDF criteria for Japanese ethnicity, MetS was diagnosed when waist circumference ≥ 85 cm and two or more of the following four components were present: (1) HDL cholesterol <40 mg per 100 ml; (2) triglyceride ≥ 150 mg per 100 ml; (3) systolic blood pressure ≥ 130 mm Hg and/or diastolic blood pressure ≥ 85 mm Hg and (4) fasting plasma glucose ≥ 100 mg per 100 ml. Subjects taking anti-hypertensive medications were considered hypertensive for statistical purposes.

Data analysis

Values are expressed as the mean \pm s.d. in the text unless otherwise stated. Pearson's simple correlation coefficients were calculated between plasma hormones and the number of MetS components. Differences between the quartile groups of sex hormones were analyzed using one-factor ANOVA followed by the Newman-Keuls' test. Logistic regression analysis was performed to determine the association of sex hormones with the diagnosis of MetS. Furthermore, multiple regression analysis was performed to determine the association between sex hormones and metabolic risk factors for MetS. A value of $P < 0.05$ was considered statistically significant. The data were analyzed using SPSS (Version 17.0, SPSS, Chicago, IL, USA).

RESULTS

Sex hormones and MetS criteria

Characteristics of the study subjects are shown in Table 1. Twenty-three and 32% of the subjects were diagnosed with MetS according to the Japanese criteria and the IDF criteria, respectively. The prevalence is comparable with that reported in middle-aged Japanese men.^{22,23}

As plasma total testosterone was negatively correlated with the number of MetS components (Figure 1a), the association of testosterone with MetS was analyzed by quartile of testosterone. As shown in Figure 2a, lower testosterone was associated with a step-wise increase in the number of MetS components. Age-adjusted logistic regression analysis revealed that the hazard ratios for MetS diagnosis by quartile decline of testosterone were 2.02 (95% CI=1.43–2.87) and 1.68 (95% CI=1.25–2.25) according to the Japanese criteria and the IDF criteria, respectively.

Interestingly, plasma estradiol was positively correlated with the number of MetS components ($R=0.285$, $P < 0.001$); therefore, the association with MetS was also analyzed by quartile of estradiol. As shown in Figure 2b, higher estradiol was associated with a step-wise increase in the number of MetS components. Age-adjusted logistic regression analysis revealed that the hazard ratios for MetS diagnosis by quartile increment of estradiol were 1.48 (95% CI=1.06–2.06) and 1.63 (95% CI=1.20–2.21) according to the Japanese criteria and the IDF criteria, respectively. Dehydroepiandrosterone sulfate was not associated with MetS components or diagnosis (data not shown).

Table 1 Characteristics of study subjects ($N=194$)

Age (years)	49 \pm 9	[30–64]
Body mass index (kg m^{-2})	25.2 \pm 4.0	[17.3–41.9]
Waist circumference (cm)	87 \pm 10	[69–125]
Hip circumference (cm)	96 \pm 7	[80–125]
Waist/hip ratio	0.94 \pm 0.06	[0.78–1.09]
Systolic blood pressure (mm Hg)	126 \pm 14	[95–183]
Diastolic blood pressure (mm Hg)	79 \pm 11	[50–128]
Triglycerides (mg per 100 ml)	162 \pm 135	[32–880]
HDL cholesterol (mg per 100 ml)	54 \pm 16	[26–110]
Free fatty acids (mEq l ⁻¹)	0.53 \pm 0.28	[0.08–2.08]
LDL cholesterol (mg per 100 ml)	128 \pm 29	[54–213]
Fasting plasma glucose (mg per 100 ml)	98 \pm 13	[76–158]
Hemoglobin A1c (%)	5.2 \pm 0.6	[4.0–8.0]
Insulin ($\mu\text{U ml}^{-1}$)	6.7 \pm 4.0	[1.0–21.2]
HOMA-IR	1.64 \pm 1.04	[0.21–5.50]
Total testosterone (nmol l ⁻¹)	19.1 \pm 6.2	[4.6–38.2]
DHEA-S ($\mu\text{mol l}^{-1}$)	5.89 \pm 2.37	[1.12–12.0]
Estradiol (pmol l ⁻¹)	92.5 \pm 43.7	[18.4–216.6]
<i>Metabolic syndrome (MetS) and its components</i>		
MetS (Japanese criteria), n (%)		44 (23)
MetS (IDF criteria), n (%)		62 (32)
Waist circumference ≥ 85 cm, n (%)		110 (56)
High blood pressure, n (%)		89 (46)
HDL cholesterol <40 mg per 100 ml, n (%)		34 (18)
Triglycerides ≥ 150 mg per 100 ml, n (%)		79 (41)
Fasting plasma glucose ≥ 110 mg per 100 ml, n (%)		23 (12)
Fasting plasma glucose ≥ 100 mg per 100 ml, n (%)		73 (38)

Abbreviations: DHEA-S, dehydroepiandrosterone sulfate; HDL, high-density lipoprotein; HOMA-IR, homeostasis model assessment of insulin resistance; IDF, International Diabetes Federation; LDL, low-density lipoprotein.

Values are expressed as the mean \pm s.d. (range). High blood pressure was defined if subjects showed systolic blood pressure ≥ 130 mm Hg and/or diastolic blood pressure ≥ 85 mm Hg, or were taking antihypertensive medications.




Figure 1 Scattergrams and regression lines (dotted lines) showing the correlation between testosterone and the number of metabolic syndrome (MetS) components (a) or waist circumference (b).




Figure 2 Number of metabolic syndrome (MetS) components according to quartiles of plasma testosterone (a) and estradiol (b). MetS components were defined according to the Japanese criteria (closed bars) and the IDF criteria for Japanese ethnicity (open bars). Values are expressed as the mean \pm s.e.m. * P <0.05, ** P <0.01 vs. Q1. Cut offs of the quartiles were 14.1, 18.7 and 23.4 nmol l⁻¹ (405, 540 and 674 ng per 100 ml) for testosterone, and 55, 101 and 125 pmol l⁻¹ (15.0, 27.5 and 34.0 pg ml⁻¹) for estradiol.

Sex hormones and metabolic risk factors

The associations of plasma sex hormones with each of the metabolic risk factors were analyzed. As shown in Table 2, the unadjusted model shows that testosterone was significantly related to parameters of MetS except for diastolic blood pressure. Testosterone was not related to low-density lipoprotein cholesterol, but this parameter is not included in the definitions of MetS used here. Adjustment for age did not considerably influence the results of the regression analysis, but the association between testosterone and diastolic blood pressure became significant after adjustment for age. In contrast, inclusion of waist circumference into the model weakened the association of testosterone with metabolic risk factors. As a result, systolic blood pressure, triglycerides, fasting plasma glucose and HOMA-IR were significantly related to testosterone. The significant association for diastolic blood pressure, HDL cholesterol, free fatty acids, hemoglobin A1c, insulin and adiponectin were attenuated after adjustment for age and waist circumference. Adjustment for body mass index or waist/hip ratio instead of waist circumference showed similar results (data not shown).

As shown in Table 3, estradiol showed weaker association than testosterone with parameters of MetS, but was significantly related to body mass index, waist circumference, systolic blood pressure, HDL

Table 2 Multiple regression analysis determining the impact of plasma testosterone on metabolic risk factors

	Unadjusted	Age adjusted	Age+waist adjusted
Body mass index	-0.376*	-0.343*	ND
Waist circumference	-0.378*	-0.365*	ND
Waist/hip ratio	-0.353*	-0.384*	ND
Systolic blood pressure	-0.230**	-0.278*	-0.169***
Diastolic blood pressure	-0.114	-0.157***	-0.098
Triglycerides	-0.247*	-0.242*	-0.182***
HDL cholesterol	0.252*	0.228**	0.065
Free fatty acids	-0.208**	-0.209**	-0.137
LDL cholesterol	-0.054	-0.056	-0.020
Fasting plasma glucose	-0.231**	-0.253**	-0.228**
Hemoglobin A1c	-0.166***	-0.220**	-0.137
Insulin	-0.331*	-0.307*	-0.129
HOMA-IR	-0.349*	-0.333*	-0.159***
Adiponectin	0.222**	0.216**	0.046

Abbreviations: HDL, high-density lipoprotein; HOMA-IR, homeostasis model assessment of insulin resistance; LDL, low-density lipoprotein; ND, not determined. Regression coefficients with plasma testosterone as an independent variable and each of risk factors as a dependent variable are shown. Age and/or waist circumference were included in multiple regression models as indicated. * P <0.001, ** P <0.01, *** P <0.05.

Table 3 Multiple regression analysis determining the impact of plasma estradiol on metabolic risk factors

	Unadjusted	Age adjusted	Age+waist adjusted
Body mass index	0.279*	0.260*	ND
Waist circumference	0.346*	0.338*	ND
Waist/hip ratio	0.102	0.082	ND
Systolic blood pressure	0.133	0.158**	0.042
Diastolic blood pressure	0.036	0.058	-0.002
Triglycerides	0.105	0.094	-0.012
HDL cholesterol	-0.207***	-0.193***	-0.040
Free fatty acids	0.087	0.091	0.049
LDL cholesterol	-0.056	-0.056	-0.094
Fasting plasma glucose	0.130	0.141	0.095
Hemoglobin A1c	0.040	0.067	-0.030
Insulin	0.240***	0.228***	0.038
HOMA-IR	0.250***	0.243***	0.060
Adiponectin	-0.267*	-0.262*	-0.114

Abbreviations: HDL, high-density lipoprotein; HOMA-IR, homeostasis model assessment of insulin resistance; LDL, low-density lipoprotein; ND, not determined. Regression coefficients with plasma estradiol as an independent variable and each of risk factors as a dependent variable are shown. Age and/or waist circumference were included in multiple regression models as indicated. * $P < 0.001$, ** $P < 0.05$, *** $P < 0.01$.

cholesterol, insulin, HOMA-IR and adiponectin after adjustment for age. Further adjustment for waist circumference, body mass index or waist/hip ratio (Table 3 and data not shown) eliminated the significant associations between estradiol and these metabolic parameters. Dehydroepiandrosterone sulfate was not significantly related to parameters of MetS in unadjusted or adjusted analyses (data not shown).

DISCUSSION

In this study, cross-sectional analysis of 194 middle-aged Japanese men showed that low testosterone is positively related to MetS, MetS components and additional metabolic risk factors. Adjustment for obesity parameters such as waist circumference, body mass index and waist/hip ratio greatly diminished the association, but low testosterone retained weak associations with some metabolic risk factors including systolic blood pressure, triglycerides, fasting plasma glucose and HOMA-IR. Taken together, results in this statistical model suggest that abdominal obesity is an important contributor to the association between low testosterone and MetS, but additional factors may also impact testosterone. To our knowledge, this is the first report showing the significant association between low testosterone and MetS in Japanese men.

Several mechanisms have been suggested for the causal relationship between low testosterone and abdominal obesity. Activation of the lipoprotein lipase and lipolysis²⁴ may explain the effect of testosterone on adipose tissue. Many studies including a medium-sized, randomized controlled trial²⁵ and a meta-analysis²⁶ showed the inverse effect of testosterone on adiposity. Conversely, it has been reported that men with MetS are prone to hypogonadism.²⁷ This finding might be due to elevated leptin levels that interfere with gonadotropin-stimulated androgen production²⁸ and to increased aromatase activity in adipose tissue that leads to higher circulating estradiol and suppression of testosterone production by negative feedback.²⁹ These findings suggest a bi-directional causal relationship between low testosterone and obesity.

After adjustment for waist circumference, testosterone was weakly but significantly related to some metabolic risk factors including systolic blood pressure, triglycerides, fasting plasma glucose and

HOMA-IR, which is consistent with earlier reports.^{5,6,8,12} Testosterone is likely to be involved in the pathogenesis of MetS, irrespective of obesity. For example, testosterone increases the hepatic production of apolipoprotein A-1 and consequently increases HDL cholesterol,³⁰ improves insulin sensitivity and increases muscle strength.³¹ There was no significant correlation between age and testosterone ($R=0.114$, $P=0.12$). This result may be because the cohort was limited to middle-aged men (30–64 years old). However, age was included in the multivariate analyses in this study, because it is well established that testosterone declines with age.⁴

The positive association found between testosterone and adiponectin is in agreement with earlier reports.^{16,32,33} However, the direct action of testosterone on adiponectin production/secretion might be different from these findings, because testosterone decreases adiponectin secretion in mice and in adipocytes.^{34,35} Accordingly, abdominal obesity may underlie the positive correlation between testosterone and adiponectin in men.

In this study, estradiol was associated positively with MetS and its components, consistent with an earlier report.¹² This relationship may be independent of testosterone because estradiol was not correlated with testosterone by simple regression analysis ($R=-0.019$, $P=0.80$), and the inclusion of both testosterone and estradiol into the multiple regression model as covariates did not influence the association of each other with MetS parameters (data not shown). The relationship between estradiol and MetS might be attributed to increased aromatase activity and subsequent elevation of circulating estradiol in obese subjects.²⁹ Increased estradiol may subsequently suppress pituitary function,²⁹ and lead to a further decrease in testosterone. Comprehensive assessment of sex hormone, gonadotropin and components of MetS reveal a causal relationship. Unfortunately, we could not measure gonadotropin because of limited plasma. Further investigation is needed to address the mechanistic and pathophysiological interactions between sex hormones and MetS.

There are some limitations to our study. First, the cross-sectional design does not clarify the causal relationship between sex hormones and MetS. As there may be bi-directional causalities as mentioned above, longitudinal follow-up studies and hormone replacement studies should be performed in Japanese populations. Second, active forms of testosterone such as bioavailable and calculated free testosterone were not measured. A direct assay of bioavailable testosterone or of sex hormone-binding globulin (required for free testosterone calculation) was not available for the study. Third, the potential influence of medications on the measured parameters cannot be denied, although the exclusion of subjects on statins ($n=40$) or anti-hypertensive drugs ($n=44$) did not seriously affect the association of testosterone with waist circumference (statins, $R=-0.304$, $P < 0.01$; anti-hypertensives, $R=-0.337$, $P < 0.01$) and the number of MetS components (statins, $R=-0.274$, $P < 0.01$; anti-hypertensives, $R=-0.278$, $P < 0.01$). Fourth, because the sample size ($n=194$) is relatively small, the finding needs to be confirmed in a larger cohort.

In summary, this study suggests that low testosterone is associated with MetS and its parameters in middle-aged Japanese men. We also found a positive but weaker association between estradiol and MetS. These associations were largely attenuated by adjustment for waist circumference. Our results reinforce the need to address the causal relationship and pathophysiological interactions between sex hormones and MetS.

ACKNOWLEDGEMENTS

We thank Ms Yuki Ito for her excellent technical assistance. This study was supported by a Health and Labor Sciences Research Grant (H17-Choju-046)

from the Ministry of Health, Labor and Welfare of Japan, Grants-in-Aid for Scientific Research from the Ministry of Education, Science, Culture and Sports of Japan (21390220, 20249041) and grants from the NOVARTIS Foundation for Gerontological Research and the Yamaguchi Endocrine Research Association.

- 1 Taskinen MR. Is metabolic syndrome the main threat to human health in the twenty-first century? *Arterioscler Thromb Vasc Biol* 2007; **27**: 2275.
- 2 Oda E. The metabolic syndrome as a concept of adipose tissue disease. *Hypertens Res* 2008; **31**: 1283–1291.
- 3 Noda H, Iso H, Saito I, Konishi M, Inoue M, Tsugane S, JPHC Study Group. The impact of the metabolic syndrome and its components on the incidence of ischemic heart disease and stroke: the Japan public health center-based study. *Hypertens Res* 2009; **32**: 289–298.
- 4 Muller M, den Tonkelaar I, Thijssen JH, Grobbee DE, van der Schouw YT. Endogenous sex hormones in men aged 40–80 years. *Eur J Endocrinol* 2003; **149**: 583–589.
- 5 Tsai EC, Matsumoto AM, Fujimoto WY, Boyko EJ. Association of bioavailable, free, and total testosterone with insulin resistance: influence of sex hormone-binding globulin and body fat. *Diabetes Care* 2004; **27**: 861–868.
- 6 Stellato RK, Feldman HA, Hamdy O, Horton ES, McKinlay JB. Testosterone, sex hormone-binding globulin, and the development of type 2 diabetes in middle-aged men: prospective results from the Massachusetts male aging study. *Diabetes Care* 2000; **23**: 490–494.
- 7 Simon D, Charles MA, Nahoul K, Orssaud G, Kremiski J, Hully V, Joubert E, Papoz L, Eschwège E. Association between plasma total testosterone and cardiovascular risk factors in healthy adult men: the Telecom Study. *J Clin Endocrinol Metab* 1997; **82**: 682–685.
- 8 Fogari R, Preti P, Zoppi A, Fogari E, Rinaldi A, Corradi L, Mugellini A. Serum testosterone levels and arterial blood pressure in the elderly. *Hypertens Res* 2005; **28**: 625–630.
- 9 Khaw K-T, Dowsett M, Folkard E, Bingham S, Wareham N, Luben R, Welch A, Day N. Endogenous testosterone and mortality due to all causes, cardiovascular disease, and cancer in men. *Circulation* 2007; **116**: 2694–2701.
- 10 Laughlin GA, Barrett-Connor E, Bergstrom J. Low serum testosterone and mortality in older men. *J Clin Endocrinol Metab* 2008; **93**: 68–75.
- 11 Laaksonen DE, Niskanen L, Punnonen K, Nyyssönen K, Tuomainen TP, Salonen R, Rauramaa R, Salonen JT. Sex hormones, inflammation and the metabolic syndrome: a population-based study. *Eur J Endocrinol* 2003; **149**: 601–608.
- 12 Muller M, Grobbee DE, den Tonkelaar I, Lamberts SW, van der Schouw YT. Endogenous sex hormones and metabolic syndrome in aging men. *J Clin Endocrinol Metab* 2005; **90**: 2618–2623.
- 13 Rodriguez A, Muller DC, Metter EJ, Maggio M, Harman SM, Blackman MR, Andres R. Aging, androgens, and the metabolic syndrome in a longitudinal study of aging. *J Clin Endocrinol Metab* 2007; **92**: 3568–3572.
- 14 Laaksonen DE, Niskanen L, Punnonen K, Nyyssönen K, Tuomainen TP, Valkonen VP, Salonen R, Salonen JT. Testosterone and sex hormone-binding globulin predict the metabolic syndrome and diabetes in middle-aged men. *Diabetes Care* 2004; **27**: 1036–1041.
- 15 Kupelian V, Page ST, Araujo AB, Travison TG, Bremner WJ, McKinlay JB. Low sex hormone-binding globulin, total testosterone, and symptomatic androgen deficiency are associated with development of the metabolic syndrome in nonobese men. *J Clin Endocrinol Metab* 2006; **91**: 843–850.
- 16 Isobe T, Saitoh S, Takagi S, Takeuchi H, Chiba Y, Katoh N, Shimamoto K. Influence of gender, age and renal function on plasma adiponectin level: the Tanno and Sobetsu study. *Eur J Endocrinol* 2005; **153**: 91–98.
- 17 Akishita M, Hashimoto M, Ohike Y, Ogawa S, Iijima K, Eto M, Ouchi Y. Low testosterone level is an independent determinant of endothelial dysfunction in men. *Hypertens Res* 2007; **30**: 1029–1034.
- 18 Akishita M, Hashimoto M, Ohike Y, Ogawa S, Iijima K, Eto M, Ouchi Y. Low testosterone level as a predictor of cardiovascular events in Japanese men with coronary risk factors. *Atherosclerosis* 2009 (e-pub ahead of print).
- 19 Fogari R, Preti P, Derosa G, Marasi G, Zoppi A, Rinaldi A, Mugellini A. Effect of antihypertensive treatment with valsartan or atenolol on sexual activity and plasma testosterone in hypertensive men. *Eur J Clin Pharmacol* 2002; **58**: 177–180.
- 20 Matsuzawa Y. Metabolic syndrome—definition and diagnostic criteria in Japan. *J Atheroscler Thromb* 2005; **12**: 301.
- 21 Alberti KG, Zimmet P, Shaw J, IDF Epidemiology Task Force Consensus Group. The metabolic syndrome—a new worldwide definition. *Lancet* 2005; **366**: 1059–1062.
- 22 Takeuchi H, Saitoh S, Takagi S, Ohnishi H, Ohhata J, Isobe T, Shimamoto K. Metabolic syndrome and cardiac disease in Japanese men: applicability of the concept of metabolic syndrome defined by the national cholesterol education program-adult treatment panel III to Japanese men—the Tanno and Sobetsu study. *Hypertens Res* 2005; **28**: 203–208.
- 23 Nishimura R, Nakagami T, Tominaga M, Yoshiike N, Tajima N. Prevalence of metabolic syndrome and optimal waist circumference cut-off values in Japan. *Diabetes Res Clin Pract* 2007; **78**: 77–84.
- 24 Rebuffé-Scrive M, Mårin P, Björntorp P. Effect of testosterone on abdominal adipose tissue. *Int J Obes* 1991; **15**: 791–795.
- 25 Emmelot-Vonk MH, Verhaar HJ, Nakhai Pour HR, Aleman A, Lock TM, Bosch JL, Grobbee DE, van der Schouw YT. Effect of testosterone supplementation on functional mobility, cognition, and other parameters in older men: a randomized controlled trial. *JAMA* 2008; **299**: 39–52.
- 26 Isidori AM, Giannetta E, Greco EA, Gianfrilli D, Bonifacio V, Isidori A, Lenzi A, Fabbri A. Effects of testosterone on body composition, bone metabolism and serum lipid profile in middle-aged men: a meta-analysis. *Clin Endocrinol (Oxf)* 2005; **63**: 280–293.
- 27 Laaksonen DE, Niskanen L, Punnonen K, Nyyssönen K, Tuomainen TP, Valkonen VP, Salonen JT. The metabolic syndrome and smoking in relation to hypogonadism in middle-aged men: a prospective cohort study. *J Clin Endocrinol Metab* 2005; **90**: 712–719.
- 28 Isidori AM, Caprio M, Strollo F, Moretti C, Frajese G, Isidori A, Fabbri A. Leptin and androgens in male obesity: evidence for leptin contribution to reduced androgen levels. *J Clin Endocrinol Metab* 1999; **84**: 3673–3680.
- 29 Kalyani RR, Dobs AS. Androgen deficiency, diabetes, and the metabolic syndrome in men. *Curr Opin Endocrinol Diabetes Obes* 2007; **14**: 226–234.
- 30 Zmuda JM, Caucey JA, Kriska A, Glynn NW, Gutai JP, Kuller LH. Longitudinal relation between endogenous testosterone and cardiovascular disease risk factors in middle-aged men. A 13 year follow-up of former multiple risk factor intervention trial participants. *Am J Epidemiol* 1997; **146**: 609–617.
- 31 Carniniti G, Volterrani M, Iellamo F, Marazzi G, Massaro R, Miceli M, Mammì C, Piepoli M, Fini M, Rosano GM. Effect of long-acting testosterone treatment on functional exercise capacity, skeletal muscle performance, insulin resistance, and baroreflex sensitivity in elderly patients with chronic heart failure: a double-blind, placebo-controlled, randomized study. *J Am Coll Cardiol* 2009; **54**: 919–927.
- 32 Laughlin GA, Barrett-Connor E, May S. Sex-specific determinants of serum adiponectin in older adults: the role of endogenous sex hormones. *Int J Obes (Lond)* 2007; **31**: 457–465.
- 33 Gannagé-Yared MH, Khalife S, Semaan M, Fares F, Jambart S, Halaby G. Serum adiponectin and leptin levels in relation to the metabolic syndrome, androgenic profile and somatotrophic axis in healthy non-diabetic elderly men. *Eur J Endocrinol* 2006; **155**: 167–176.
- 34 Nishizawa H, Shimomura I, Kishida K, Maeda N, Kuriyama H, Nagaretani H, Matsuda M, Kondo H, Furuyama N, Kihara S, Nakamura T, Tochino Y, Funahashi T, Matsuzawa Y. Androgens decrease plasma adiponectin, an insulin-sensitizing adipocytederived protein. *Diabetes* 2002; **51**: 2734–2741.
- 35 Xu A, Chan KW, Hoo RL, Wang Y, Tan KC, Zhang J, Chen B, Lam MC, Tse C, Cooper GJ, Lam KS. Testosterone selectively reduces the high molecular weight form of adiponectin by inhibiting its secretion from adipocytes. *J Biol Chem* 2005; **280**: 18073–18080.