

Conflict of interest The authors have no conflicts or disclosures to declare regarding the present manuscript.

References

1. Yoshimura N, Muraki S, Oka H, Mabuchi A, En-yo Y, Yoshida M, Saika A, Suzuki T, Yoshida H, Kawaguchi H, Nakamura K, Akune T (2009) Prevalence of knee osteoarthritis, lumbar spondylosis and osteoporosis in Japanese men and women: the Research on Osteoarthritis/Osteoporosis Against Disability (ROAD). *J Bone Miner Metab* 27:620–628
2. Ministry of Health, Labour and Welfare (2007) The outline of the results of National Livelihood Survey 2007. <http://www.mhlw.go.jp/toukei/list/20-19-1.html>. Accessed 7 Jan 2010
3. Sakata T, Yaegashi Y, Hosoi T, Orimo H (2009) The nationwide survey on the incidence of hip fracture in Japan and 20-year trends. *Osteoporos Int* 20(suppl 1):S63
4. Delmas PD, Eastell R, Garnero P, Seibel MJ, Stepan J, Committee of Scientific Advisors of the International Osteoporosis Foundation (2000) The use of biochemical markers of bone turnover in osteoporosis. *Osteoporos Int* 11(suppl 6):S2–S17
5. Szulc P, Delmas PD (2008) Biochemical markers of bone turnover: potential use in the investigation, management of postmenopausal osteoporosis. *Osteoporos Int* 19:1683–1704
6. Bergmann P, Body JJ, Boonen S, Boutsen Y, Devogelaer JP, Goemaere S, Kaufman JM, Reginster JY, Gangji V, Members of Advisory Board on Bone Markers (2009) Evidence-based guidelines for the use of biochemical markers of bone turnover in the selection and monitoring of bisphosphonate treatment in osteoporosis: a consensus document of the Belgian Bone Club. *Int J Clin Pract* 63:19–26
7. Garnero P, Sornay-Rendu E, Duboeuf F, Delmas PD (1999) Markers of bone turnover predict postmenopausal forearm bone loss over 4 years: the OFELY study. *J Bone Miner Res* 14:1614–1621
8. Chapurlat RD, Garnero P, Sprnay-Rendu E, Arlot ME, Claustrat B, Delmas PD (2000) Longitudinal study of bone loss in pre- and perimenopausal women: evidence for bone loss in perimenopausal women. *Osteoporos Int* 11:493–498
9. Stepan JJ (2000) Prediction of bone loss in postmenopausal women. *Osteoporos Int* 11(suppl 6):S45–S54
10. Iki M, Morita A, Ikeda Y et al (2006) Biochemical markers of bone turnover predict bone loss in perimenopausal women but not in postmenopausal women—the Japanese Population-based Osteoporosis (JPOS) cohort study. *Osteoporosis Int* 17:1086–1095
11. Yoshimura N, Hashimoto T, Sakata K, Morioka S, Kasamatsu T, Cooper C (1999) Biochemical markers of bone turnover and bone loss at the lumbar spine and femoral neck. The Taiji study. *Calcif Tissue Int* 65:198–202
12. Bauer DC, Garnero P, Harrison SL, Cauley JA, Eastell R, Ensrud KE, Orwoll E, Osteoporotic Fractures in Men (MrOS) Research Group (2009) Biochemical markers of bone turnover, hip bone loss, and fracture in older men: the MrOS study. *J Bone Miner Res* 24:2032–2038
13. Yoshimura N, Kinoshita H, Danjoh S, Takijiri T, Morioka S, Kasamatsu T, Sakata K, Hashimoto T (2002) Bone loss at the lumbar spine and the proximal femur in a rural Japanese community, 1990–2000: the Miyama study. *Osteoporos Int* 13:803–808
14. Kasamatsu T, Yoshimura N, Morioka S, Sugita K, Hashimoto T (1996) A population survey on bone mineral density in a fishing village in Wakayama Prefecture (Part 1). Distribution of bone mineral density by sex and age on a representative sample of the community. *Jpn J Hyg* 50:1084–1092 (in Japanese)
15. Yoshimura N, Kasamatsu T, Morioka S, Hashimoto T (1996) A population survey on bone mineral density in a fishing village in Wakayama Prefecture (Part 2). The analysis of the risk factors affecting the bone mineral density. *Jpn J Hyg* 51:677–684 (in Japanese)
16. Yoshimura N, Hashimoto T, Morioka S, Sakata K, Kasamatsu T, Cooper C (1998) Determinants of bone loss in a rural Japanese community. The Taiji Study. *Osteoporos Int* 8:604–610
17. Yoshimura N, Kakimoto T, Nishioka M, Kishi T, Iwasaki H, Niwa T, Morioka S, Sakata T, Hashimoto T (1997) Evaluation of reproducibility of bone mineral density measured by dual energy X-ray absorptiometry (DPX-L). *J Wakayama Med Soc* 48:461–466
18. Schmidt-Gayk H, Spanuth E, Kotting J, Bartl R, Felsenberg D, Pfeilshifter J, Raue F, Roth HJ (2004) Performance evaluation of automated assays for beta-Crosslaps, N-MID Osteocalcin and intact parathyroid hormone (BIROSE Multicenter Study). *Clin Chem Lab Med* 42:90–95
19. Clemens JD, Herrick MV, Singer FR, Eyre DR (1997) Evidence that serum NTx (collagen-type I N-telopeptides) can act as an immunochemical marker of bone resorption. *Clin Chem* 43:2058–2063
20. Gertz BJ, Clemens JD, Holland SD, Yuan W, Greenspan S (1998) Application of a new serum assay for type I collagen cross-linked N-telopeptides: assessment of diurnal changes in bone turnover with and without alendronate treatment. *Calcif Tissue Int* 63:102–106
21. Ebeling PR, Atley LM, Guthrie JR, Burger HG, Dennerstein L, Hopper JL, Wark JD (1996) Bone turnover markers and bone density across the menopausal transition. *J Clin Endocrinol Metab* 81:3366–3371
22. Sowers MR, Greendale GA, Bondarenko I, Finkelstein JS, Cauley JA, Neer RM, Ettinger B (2003) Endogenous hormones and bone turnover markers in pre- and perimenopausal women: SWAN. *Osteoporos Int* 14:191–197
23. Yoshimura N, Kasamatsu T, Sakata K, Hashimoto T, Cooper C (2002) The relationship between endogenous estrogen, sex hormone binding globulin and bone loss in female residents of a rural Japanese community: the Taiji study. *J Bone Miner Metab* 20:303–310
24. Yoshimura N, Takijiri T, Kinoshita H, Danjoh S, Kasamatsu T, Morioka S, Sakata K, Hashimoto T, Takeshita T (2004) Characteristics and course of bone mineral densities among fast bone losers in a rural Japanese community: the Miyama study. *Osteoporos Int* 15:139–144
25. Fujiwara S, Mizuno S, Ochi Y, Sasaki H, Kodama K, Russell WJ, Hosoda Y (1991) The incidence of thoracic vertebral fractures in a Japanese population, Hiroshima and Nagasaki, 1958–86. *J Clin Epidemiol* 44:1007–1014
26. Hannon R, Eastell R (2000) Preanalytical variability of biochemical markers of bone turnover. *Osteoporos Int* 11(suppl 6):S30–S44
27. Seibel MJ (2000) Molecular markers of bone turnover: biochemical, technical and analytical aspects. *Osteoporos Int* 11(suppl 6):S18–S29

Reference values for hand grip strength, muscle mass, walking time, and one-leg standing time as indices for locomotive syndrome and associated disability: the second survey of the ROAD study

Noriko Yoshimura · Hiroyuki Oka · Shigeyuki Muraki · Toru Akune · Naoki Hirabayashi · Shinji Matsuda · Takako Nojiri · Kazuhiro Hatanaka · Yuyu Ishimoto · Keiji Nagata · Munehito Yoshida · Fumiaki Tokimura · Hiroshi Kawaguchi · Kozo Nakamura

Received: 4 April 2011 / Accepted: 30 August 2011 / Published online: 6 October 2011
© The Japanese Orthopaedic Association 2011

Abstract

Background We established reference values for hand grip strength, muscle mass, walking time, and one-leg standing time as indices reflecting components of locomotive syndrome and associated disability using a large-scale population-based sample from the second survey of the Research on Osteoarthritis/Osteoporosis Against Disability (ROAD) cohort.

Methods We measured the above-mentioned indices in 2,468 individuals ≥ 40 years old (826 men, 1,642 women; mean age 71.8 years) during the second visit of the ROAD study. Disability was defined as certified disability according to the long-term care insurance system through public health centres of each municipality.

Results Mean values for hand grip strength (weaker side), muscle mass of the thighs, walking time for 6 m at the

usual pace, and the fastest pace for men were 32.7 kg, 7.0 kg, 5.6 s, and 3.7 s, respectively, and those for women were 20.8 kg, 5.2 kg, 5.9 s, and 4.1 s, respectively. The median values for one-leg standing time (weaker side) were 14 s for men and 12 s for women. The prevalence of disability in men aged 65–69, 70–74, 75–79, and ≥ 80 was 0.0, 1.0, 6.3, and 8.8%, respectively, and in women was 3.4, 3.5, 9.2, and 14.7%, respectively. There were significant associations between the presence of disability and walking time for 6 m at the usual pace and at the fastest pace, and between the presence of disability and walking speed.

Conclusions We established reference values for indices reflecting components of locomotive syndrome, and identified significant associations between walking ability and disability.

N. Yoshimura (✉) · H. Oka
Department of Joint Disease Research,
22nd Century Medical and Research Center,
The University of Tokyo, Hongo 7-3-1,
Bunkyo-ku, Tokyo 113-8655, Japan
e-mail: yoshimuran-ort@h.u-tokyo.ac.jp

S. Muraki · T. Akune
Department of Clinical Motor System Medicine,
22nd Century Medical and Research Center,
The University of Tokyo, Tokyo, Japan

N. Hirabayashi
Kawakami Clinic, Hidakagawa-cho, Wakayama, Japan

S. Matsuda
Iwade Public Health Center, Wakayama, Japan

T. Nojiri · K. Hatanaka
Gobo Public Health Center, Wakayama, Japan

Y. Ishimoto · K. Nagata · M. Yoshida
Department of Orthopaedic Surgery,
Wakayama Medical University School of Medicine,
Wakayama, Japan

F. Tokimura
Tokyo Metropolitan Institute of Gerontology, Tokyo, Japan

H. Kawaguchi
Department of Orthopaedic Surgery,
Sensory and Motor System Medicine,
Graduate School of Medicine,
The University of Tokyo, Tokyo, Japan

K. Nakamura
National Rehabilitation Center for Persons with Disabilities,
Saitama, Japan

Introduction

Musculoskeletal diseases, including osteoarthritis (OA) and osteoporosis (OP), are major public health problems among the elderly that affect activities of daily living (ADL) and quality of life (QOL), leading to increased morbidity and mortality. According to the recent National Livelihood Survey by the Ministry of Health, Labour, and Welfare in Japan, OA is ranked fourth, while falls and osteoporotic fractures are ranked fifth among diseases that cause disabilities and subsequently require support for ADL [1]. Previous studies have reported increased mortality following osteoporotic fractures at the hip and other sites [2], and have estimated that a total of 47,000,000 people (21,000,000 men and 26,000,000 women) aged ≥ 40 years will eventually be affected by either OA or OP. Considering that the population of Japan is aging very rapidly—more than 22% of the population is aged ≥ 65 years [3]—a comprehensive and evidence-based prevention strategy for musculoskeletal diseases is urgently needed.

The Japanese Orthopaedic Association has proposed the term ‘locomotive syndrome’ to designate a condition in high-risk groups with musculoskeletal diseases who are highly likely to require nursing care [4]. Locomotive syndrome is caused by weakening of musculoskeletal organs such as bone, joint, and muscle, which in turn interferes with physical performance, especially self-transportation. Loss of locomotor abilities such as walking causes disabilities requiring support. Therefore, to prevent decline into disability, it is important to maintain a healthy range of bone, joint, muscle, and physical performance.

These four components, bone, joint, muscle, and physical performance, each have objective measurements that can be used as indices to evaluate their present condition. For example, bone mineral density (BMD) is a representative index of the condition of the bone. Joint space width (JSW), joint space area (JSA), and osteophyte area (OPA) are indices reflecting the condition of the joint. Regarding muscle, although the best index remains controversial, hand grip strength can be used to reflect muscle strength [5], and muscle mass is one index of muscle volume [6]. In addition, as objective indices of physical performance, walking speed and/or one-leg standing times are candidates [7, 8]. However, at present, it is difficult to use such indices for evaluating, diagnosing, or predicting the future occurrence and progression of locomotive syndrome in Japan, because there is little information on reference values for such indices to distinguish patients at risk from normal individuals in a large population-based cohort.

In 2005–2007, we began a large-scale population-based cohort study entitled Research on Osteoarthritis/

Osteoporosis Against Disability (ROAD), consisting of 3,040 participants in three communities located in urban, mountainous, and coastal areas (baseline study). Following the baseline study, a second survey was performed in the same communities in 2008–2010, in which 2,674 inhabitants participated (second visit).

Through analysis of the baseline data of the ROAD, the age-gender distribution of BMD has been reported as an index for bone mass [3], and the medial and lateral JSW, medial and lateral JSA, OPA, and femorotibial angle of the knee have been reported as indices of the health of joints [9] in these populations. However, there is still scant information regarding the condition of the muscles and physical performance. Therefore, in the present study, we aimed to establish reference values for hand grip strength as an index of muscle power, muscle mass as an index of muscle volume, and walking time and one-leg standing time as indices of physical performance, classified by age and gender, using the data from the second visit of the ROAD study. This information is expected to be valuable for early diagnosis and prevention of locomotive syndrome. In addition, we evaluated the prevalence of disabilities in participants in the ROAD study second visit, and identified associations between hand grip strength, muscle mass, walking time, and one-leg standing time and the presence of disability.

Participants and methods

Participants

Reference values were obtained from the results of cross-sectional measurements of participants enrolled in the second visit of the ROAD study. The ROAD study, which began in 2005, is a nationwide prospective study comprising population-based cohorts established in three communities, such as urban, mountainous, and coastal regions in Japan. Recruitment methods for this study have been described in detail elsewhere [3]. To date, participants in the urban region, aged ≥ 60 years, were recruited from among those enrolled in a randomly selected cohort study from the previously established Itabashi Ward resident’s registration database. The response rate was 75.6%. Participants in the mountainous and coastal regions, aged ≥ 40 years, were recruited from listings of resident registration. Residents aged ≤ 60 years in the urban area and ≤ 40 years in the mountainous and coastal areas who were interested in participating in the study were invited. We have completed development of a baseline database including clinical and genetic information for 3,040 inhabitants aged 23–95 years (1,061 men, 1,979 women).

The second visit of the ROAD study began in 2008 and was completed in 2010. All the participants in the baseline study were invited to participate in the second visit. In addition to the former participants, inhabitants aged ≥ 60 years in the urban area and those aged ≥ 40 years in the mountainous and coastal areas who were willing to participate in the ROAD survey performed in 2008–2010 were also included in the second visit. In addition, residents aged ≤ 60 years in the urban area and ≤ 40 years in the mountainous and coastal areas who were interested in participating in the study were invited to be examined as well as the baseline.

The inclusion criteria of participants were as follows: (1) ability to walk to the clinic where the survey was performed, (2) ability to provide self-reported data, and (3) ability to understand and sign an informed consent form. No other exclusion criteria were used.

Thus, a total of 2,674 residents (892 men, 1,782 women) aged 21–97 years participated in the second visit. In the present study, we analysed the data for 2,468 individuals (826 men, 1,642 women; mean age 71.8 years); this population comprised 956 individuals (318 men, 638 women) in the urban region, 726 individuals (258 men, 468 women) in the mountainous region, and 786 individuals (250 men, 536 women) in the coastal region who participated in the second visit and were ≥ 40 years old.

All the participants provided written informed consent, and the study was conducted with the approval of the ethics committees of the participating institutions.

Hand grip strength, muscle mass, walking time, and one-leg standing time

Hand grip strength was measured bilaterally using a Toei Light handgrip dynamometer (Toei Light Co., Ltd., Saitama, Japan). Both hands were tested, and the better value was used to characterise the maximum muscle strength of the subject.

Among the 2,468 participants who participated in the second visit of the ROAD study, 778 residents (248 men, 530 women) in the coastal town of Taiji were examined to determine their segmental muscle mass using the bioelectrical impedance method (BIP; Physion MD; Physion Inc., Kyoto, Japan). We obtained values for the muscle masses of the right and left forearms, upper arms, upper limbs, quadriceps, thighs, lower legs, and lower limbs. This method had previously been validated as having a close correlation to muscle volume as measured by magnetic resonance imaging [10].

Among the 2,468 participants who participated in the second visit of the ROAD study, 1,637 residents (559 men, 1,078 women) of the mountainous town of Hidakagawa and the coastal town of Taiji were examined to determine

their walking time. Walking time was measured as the time required to complete a 6-m course. All participants walked the 6-m course twice; they first walked at their usual walking speed and then repeated the course at their fastest pace.

Among the 2,468 participants who participated in the second visit of the ROAD study, one-leg standing time with eyes open was measured on both sides for 2,433 individuals (816 men, 1,617 women). The time until the raised leg was set down on the floor was measured, with a maximum time of 60 s recorded for those who could stand on one leg for at least that length of time. The shorter value of the two measurements was used as the worse side and the longer measurement as the better side for the one-leg standing time of the subject.

Mean values and standard deviations (SDs) of hand grip strength, muscle mass, and walking time were classified by gender and age group (40, 50, 60, 70, and ≥ 80 s) to establish age-gender reference values for the general population. However, reference values classified by gender and age group for one-leg standing time were established using median (50th percentile) values and 25th–75th percentile ranges. These values were recorded using a maximum time of 60 s for anyone who could exceed that time; thus, the data do not fit a normal distribution, and use of means and SDs is unsuitable for one-leg standing time reference values.

Presence of disability

Disability in the present study was defined as ‘cases requiring long-term care’ as determined by the long-term care insurance system based on the Long-Term Care Insurance Act of 1997 in Japan. The procedure for identifying cases requiring long-term care is as follows: (1) each municipality establishes a long-term care approval board consisting of clinical experts, physicians, and specialists at the Division of Health and Welfare in each municipal office; (2) the long-term care approval board investigates the insured person using an interviewer-administered questionnaire consisting of 82 items regarding mental and physical condition and makes a screening judgement based on the opinion of a regular doctor; and (3) ‘cases requiring long-term care’ are determined according to standards for long-term care certification uniformly and objectively applied nationwide [11].

During the 3 years between the baseline and the second visit of the ROAD study, we annually obtained information on the participating residents regarding deaths, changes of residence, and presence or absence of certified disability according to the long-term care insurance system from the public health centres of the participating municipalities.

Statistical analysis

All statistical analyses were performed using Stata statistical software (Stata, College Station, TX, USA). Differences in the values of the indices were tested for significance using analysis of variance for comparisons among multiple groups. Scheffé’s least significant difference test was then used for pairs of age groups.

To ascertain associations between the presence of disability and hand grip strength, muscle mass, walking time, and one-leg standing time, logistic regression analyses were performed using the presence of disability (yes, 1; no, 0) as an objective factor, and values for hand grip strength, muscle mass, walking time, and one-leg standing time as the explanatory factor after adjusting for age, gender, and body mass index (BMI, kg/m²).

Results

Characteristics of participants

Summary characteristics including age, height, weight, and BMI of the participants in the present study are shown in Table 1. Two-thirds of the 2,468 subjects were women, and the mean age of the female participants was 1 year younger than that of the male participants. Height and weight were

significantly lower for women than for men, but no significant difference in BMI was noted between the genders. All anthropometric measurements other than BMI of females tended to decrease with age. BMI of women in their 80s and older was significantly lower than that in younger age groups, while there were no significant differences among age groups 40–70 years old.

Reference values for hand grip strength, muscle mass, walking time, and one-leg standing time

Table 1 also shows the age-gender distribution of hand grip strength for both the better and worse sides. Mean hand grip strength in men was significantly higher than that in women ($p < 0.001$) and decreased with age in both men and women ($p < 0.001$).

Mean muscle mass for both forearms, both upper arms, both upper limbs, both quadriceps, both thighs, both lower legs, and both lower limbs are shown in Table 2. Muscle masses for all parts of the body were significantly higher in men than in women ($p < 0.001$). Mean muscle mass in men decreased with age for all areas except the lower leg. Particularly in the quadriceps and thighs, muscle masses in men aged ≥ 70 were significantly lower than those in their 40s–50s ($p < 0.05$). By contrast, although muscle mass for women aged ≥ 80 and older tended to be lower than those of younger age groups (other than the lower legs), there

Table 1 Mean values (standard deviation) of anthropometric measurements and hand grip strength of the participants classified by sex and gender

Age strata (years)	Number of subjects	Weight (kg)	Height (cm)	Body mass index (g/cm ²)	Grip strength (better side) (kg)	Grip strength (worse side) (kg)
Men						
40–49	32	73.5 (10.2)	170.3 (7.3)	25.4 (3.6)	49.5 (8.2)	49.3 (8.4)
50–59	100	68.8 (10.6)	168.0 (5.2)	24.3 (3.3)	47.3 (7.0)	42.6 (6.9)
60–69	137	65.4 (11.1) ^a	165.2 (6.2) ^{a,b}	23.9 (3.5)	41.4 (6.6) ^{a,b}	36.9 (7.9) ^{a,b}
70–79	308	60.0 (8.1) ^{a,b,c}	161.1 (5.7) ^{a,b,c}	23.1 (2.7) ^{a,b}	35.4 (6.8) ^{a,b,c}	31.5 (7.1) ^{a,b,c}
80 and older	249	57.2 (8.9) ^{a,b,c,d}	159.7 (6.0) ^{a,b,c}	22.4 (2.9) ^{a,b,c}	29.7 (6.2) ^{a,b,c,d}	26.3 (6.3) ^{a,b,c,d}
Total	826	61.6 (10.3)	162.5 (6.7)	23.3 (3.1)	36.6 (9.1)	32.7 (9.1)
Women						
40–49	93	55.9 (9.5)	157.0 (4.4)	22.6 (3.5)	31.2 (4.3)	28.2 (4.4)
50–59	191	55.3 (8.9)	154.4 (5.8) ^a	23.2 (3.7)	28.7 (4.9) ^a	25.4 (4.9) ^a
60–69	316	54.2 (8.0)	152.0 (5.5) ^{a,b}	23.4 (3.2)	26.6 (4.3) ^{a,b}	23.77 (4.5) ^{a,b}
70–79	599	51.3 (8.5) ^{a,b,c}	148.4 (5.9) ^{a,b,c}	23.3 (3.5)	22.6 (4.6) ^{a,b,c}	19.7 (4.7) ^{a,b,c}
80 and older	443	47.4 (8.3) ^{a,b,c,d}	145.5 (5.9) ^{a,b,c,d}	22.4 (3.6) ^{c,d}	19.4 (4.4) ^{a,b,c,d}	16.6 (4.6) ^{a,b,c,d}
Total	1,642	51.6 (8.9)	149.5 (6.7)	23.0 (3.5)	23.7 (5.8)	20.8 (5.8)

^a Significantly different ($p < 0.05$) from values of the age group in their 40s
^b Significantly different ($p < 0.05$) from values of the age group in their 50s
^c Significantly different ($p < 0.05$) from values of the age group in their 60s
^d Significantly different ($p < 0.05$) from values of the age group in their 70s

Table 2 Mean values (standard deviation) of segmental muscle mass (kg) in total right and left sides classified by age and gender

Age strata (years)	Number of subjects	Forearm	Upper arm	Upper limb	Quadriceps	Thigh	Lower leg	Lower limb
Men								
40–49	25	1.20 (0.19)	1.59 (0.36)	2.79 (0.54)	3.91 (0.64)	7.76 (1.19)	3.41 (0.66)	11.16 (1.69)
50–59	60	1.18 (0.16)	1.53 (0.28)	2.71 (0.41)	3.73 (0.64)	7.45 (1.22)	3.41 (0.66)	10.86 (1.60)
60–69	67	1.15 (0.17)	1.50 (0.28)	2.65 (0.42)	3.50 (0.68)	7.02 (1.28)	3.52 (0.86)	10.54 (1.89)
70–79	66	1.17 (0.20)	1.43 (0.28)	2.60 (0.46)	3.37 (0.66) ^a	6.78 (1.26) ^a	3.51 (0.68)	10.29 (1.70)
80 and older	30	1.11 (0.17)	1.37 (0.26)	2.48 (0.38)	3.10 (0.62) ^{a,b}	6.27 (1.18) ^{a,b}	3.92 (1.11)	10.18 (2.05)
Total	248	1.16 (0.18)	1.48 (0.29)	2.65 (0.44)	3.52 (0.69)	7.04 (1.30)	3.53 (0.80)	10.57 (1.79)
Women								
40–49	67	0.77 (0.12)	0.86 (0.19)	1.63 (0.30)	2.65 (0.60)	5.37 (1.12)	2.65 (0.47)	8.02 (1.45)
50–59	124	0.76 (0.10)	0.82 (0.16)	1.58 (0.24)	2.56 (0.44)	5.20 (0.82)	2.58 (0.51)	7.78 (1.20)
60–69	161	0.78 (0.11)	0.84 (0.16)	1.62 (0.25)	2.55 (0.45)	5.18 (0.84)	2.57 (0.42)	7.74 (1.10)
70–79	130	0.80 (0.12) ^b	0.85 (0.16)	1.66 (0.27)	2.54 (0.46)	5.17 (0.85)	2.66 (0.53)	7.83 (1.24)
80 and older	48	0.79 (0.43)	0.82 (0.16)	1.61 (0.28)	2.39 (0.45)	4.90 (0.84)	2.91 (0.69) ^{b,c}	7.81 (1.38)
Total	530	0.78 (0.11)	0.84 (0.16)	1.62 (0.26)	2.55 (0.47)	5.18 (0.88)	2.63 (0.51)	7.81 (1.23)

^a Significantly different ($p < 0.05$) from values of the age group in their 40s

^b Significantly different ($p < 0.05$) from values of the age group in their 50s

^c Significantly different ($p < 0.05$) from values of the age group in their 60s

were no specific trends in muscle mass among age groups ≤ 79 years old. However, as for men, the muscle mass of the quadriceps in women tended to decline with age, although the difference was not statistically significant.

Mean 6-m walking time and the calculated walking speed (m/s) using the walking time, classified by age and gender, are shown in Table 3. Six-meter walking time was significantly lower in men than in women ($p < 0.05$), indicating that men tended to walk faster than women in this study population. Mean 6-m walking time for both men and women increased with age. In particular, 6-m walking times for men and women ≥ 70 years old were significantly higher than those in younger age groups ($p < 0.05$).

Table 4 shows median one-leg standing time classified by age and gender with 25th–75th percentile ranges. For both men and women in their 40s–50s, all median, 25th percentile, and 75th percentile values were 60 s, with no gender difference. One-leg standing times for men ≥ 60 years old tended to be higher than those for women, and median values declined with age in both men and women.

Prevalence of disability among subjects ≥ 65 years old

Among the 2,468 participants in the second visit of the ROAD study, we surveyed 1,845 subjects (625 men, 1,220 women) ≥ 65 years old and obtained information on the presence or absence of disability certified for long-term care insurance. We found a total of 149 individuals (8.1%;

36 men, 5.8%; 113 women, 9.3%) that were certified as requiring support. Figure 1 shows the prevalence of disability classified by gender and age. The prevalence of disability in men 65–69, 70–74, 75–79, and ≥ 80 years old was 0.0, 1.0, 6.3, and 8.8%, respectively, and that in women in the same age groups was 3.4, 3.5, 9.2, and 14.7%, respectively (Fig. 1). The prevalence of disability in women was significantly higher than that in men ($p < 0.05$) and increased with age in both genders ($p < 0.01$).

Associations between disability and hand grip strength, muscle mass, walking speed, and one-leg standing time

Logistic regression analysis was performed using the presence of disability (1, yes; 0, no) as an objective factor, and hand grip strength on the better side and the worse side; muscle mass of the forearms, upper arms, upper limbs, quadriceps, thighs, lower legs, and lower limbs; walking time for 6 m at the usual pace and at the fastest pace; and quartile of one-leg standing time [0: 0–25% (highest quartile), 1: 25–50% (higher quartile), 2: 50–75% (lower quartile), 3: 75–100% (the lowest quartile)] on the better and worse sides as explanatory factors, after adjusting for age, gender, and BMI. No significant associations were found between the presence of disability and hand grip strength, muscle mass, or one-leg standing time. However, there were significant associations between the presence of disability and 6-m walking time at the usual

Table 3 Mean values (standard deviation) of 6-m walking time (s) and walking speed (m/s) with usual pace and the fastest pace classified by age and gender

Age strata (years)	Number of subjects	Usual pace		Fastest pace	
		Time for 6 m (s)	Walking speed (m/s)	Time for 6 m (s)	Walking speed (m/s)
Men					
40–49	32	4.4 (0.6)	1.38 (0.19)	3.0 (0.5)	2.09 (0.43)
50–59	100	4.8 (0.9)	1.29 (0.20)	3.2 (0.6)	1.97 (0.36)
60–69	134	5.1 (0.9)	1.21 (0.20) ^a	3.4 (0.7)	1.82 (0.33) ^a
70–79	196	5.9 (1.8) ^{a,b,c}	1.09 (0.25) ^{a,b,c}	4.0 (1.4) ^{a,b,c}	1.62 (0.39) ^{a,b,c}
80 and older	97	6.8 (3.0) ^{a,b,c,d}	0.99 (0.33) ^{a,b,c,d}	4.5 (1.8) ^{a,b,c,d}	1.48 (0.44) ^{a,b,c}
Total	559	5.6 (1.9)	1.15 (0.27)	3.7 (1.3)	1.73 (0.42)
Women					
40–49	92	4.7 (1.0)	1.32 (0.24)	3.2 (0.6)	1.95 (0.31)
50–59	190	4.9 (0.9)	1.27 (0.23)	3.3 (0.7)	1.87 (0.33)
60–69	299	5.1 (1.1)	1.22 (0.23)	3.7 (0.8)	1.71 (0.32) ^{a,b}
70–79	345	6.3 (2.4) ^{a,b,c}	1.03 (0.25) ^{a,b,c}	4.4 (1.5) ^{a,b,c}	1.46 (0.36) ^{a,b,c}
80 and older	152	8.4 (3.9) ^{a,b,c,d}	0.82 (0.27) ^{a,b,c,d}	5.8 (2.7) ^{a,b,c,d}	1.17 (0.36) ^{a,b,c,d}
Total	1,078	5.9 (2.4)	1.12 (0.29)	4.1 (1.6)	1.60 (0.42)

^a Significantly different ($p < 0.05$) from values of the age group in their 40s

^b Significantly different ($p < 0.05$) from values of the age group in their 50s

^c Significantly different ($p < 0.05$) from values of the age group in their 60s

^d Significantly different ($p < 0.05$) from values of the age group in their 70s

Table 4 Values of median (25–75 percentile) of one-leg standing time (s, maximum = 60 s) in a better side and a worse side classified by age and gender


Age strata (years)	Number of subjects	One-leg standing time (better side) (s)	One-leg standing time (worse side) (s)
Men			
40–49	32	60 (60–60)	60 (60–60)
50–59	99	60 (60–60)	60 (60–60)
60–69	136	60 (34.5–60)	45 (14.25–60)
70–79	303	27 (9–60)	9 (4–35)
80 and older	246	8 (4–32)	4 (2–12)
Total	816	39.5 (8–30)	14 (4–60)
Women			
40–49	92	60 (60–60)	60 (60–60)
50–59	191	60 (60–60)	60 (43–60)
60–69	317	60 (41.5–60)	43 (13–60)
70–79	593	21 (8–57.5)	8 (3–25)
80 and older	424	7 (3–18.8)	3 (2–7)
Total	1,617	31 (8–60)	12 (4–60)

pace [+1 s, odds ratio (OR) 1.15, 95% confidential interval (CI) 1.07–1.24, $p < 0.001$] and at the fastest pace (+1 s, OR 1.22, 95% CI 1.08–1.38, $p < 0.01$). In addition, there were significant associations between the presence of disability and walking speed at the usual pace (+1 m/s, OR 0.07, 95% CI 0.02–0.27, $p < 0.001$) and at the fastest pace (+1 m/s, OR 0.16, 95% CI 0.06–0.41, $p < 0.001$).

Discussion

In this study, we established age-gender-classified mean values for hand grip strength as an index of muscle strength, muscle mass as an index of muscle volume, and walking time and median one-leg standing time as indices of physical performance, using data for a large-scale

Fig. 1 Prevalence of disability among subjects ≥ 65 years old classified by gender and age



population-based cohort. We found that mean hand grip strength, muscle mass, walking time, and median one-leg standing time were higher in men than in women, and decreased with age (with the exception of the muscle mass of the lower legs).

The Japanese Ministry of Education, Culture, Sports, Science, and Technology has reported ranges for physical strength and sporting ability in 69,745 Japanese men and women 6–79 years old. Mean hand grip strength in both men and women reaches peak values between the ages of 20–40, and decreases with age after 40 [12]. In the present study of a population aged ≥ 40 years, hand grip strength declined significantly with age, consistent with the previous report of the Japanese government.

Although computed tomography or MRI scans are the most reliable methods of measuring segmental muscle mass, these methods are not suitable for a large-scale population-based study. The BIA method is rapid, inexpensive, portable, and importantly, a noninvasive measuring method. Previous studies have shown that there is a strong correlation between BIA resistance and measurements of skeletal muscle mass in the arms [13], legs [13], and whole body [14]. For the BIA method used in the present study, Miyatani et al. [10] had previously compared values for muscle mass using a series of cross-sectional images of the forearm, upper arm, lower leg, and thigh on the right side of 22 male subjects as determined by the BIA and MRI methods. The BIA impedance index (L^2/Z) for every segment, calculated as the ratio of the segment length squared to the impedance, was significantly correlated with the muscle volume measured by MRI, with $r = 0.902$ – 0.976 ($p < 0.05$). These previous results demonstrate that the BIA method used in the present study is sufficiently reliable as an index of muscle volume.

In the present study, age-related differences were observed in the thighs and quadriceps. In addition, the age-related decreases in muscle mass were greater in the lower limbs than in the upper limbs, and in proximal sites than in distal sites. Yamada et al. [15] estimated the bioelectrical impedance (BI) index, calculated as the ratio of the square of segmental length to impedance in a Japanese population consisting of 1,006 individuals (374 men, 632 women) and reported that the BI index decreased most with age in the thighs, whereas there were no significant age-related changes in the forearms or lower limbs. Miyatani et al. [16] studied muscle thickness at nine sites, the forearm, anterior and posterior upper arm, abdomen, subscapular, anterior and posterior thigh, and anterior and posterior lower leg, using brightness-mode ultrasonography in 348 Japanese men aged 20–79 years. They found a greater decrease in muscle thickness in the trunk and anterior thigh than at other sites, consistent with our results. They speculated that site-related differences in muscle loss with aging may be attributed to age-related changes in the patterns of loading to and/or activation of individual muscles in daily life. However, because these results were obtained from a cross-sectional study, longitudinal data would be required to determine the mechanism of these differences. We have begun the third visit of the ROAD study, 6 years after the baseline and 3 years after the second visit, to measure losses of skeletal muscle mass at various sites. Losses of muscle mass in the quadriceps and/or thighs may result in a decrease in walking ability, including walking speed. Therefore, establishment of reference values for muscle mass is useful for prediction of future disability.

Walking ability is regarded as the most important activity for the elderly to maintain an independent life in the community, and walking speed is an important index of

walking ability. Reference values have been published for populations in western countries [17, 18]; however, there has been little information available for the Japanese population. Takahashi et al. [19] surveyed walking speed at 130 crosswalks and reported that at least 1.0 m/s was required to safely cross the street. In the present study, we determined that the mean 6-m walking time at the usual pace at an age of ≥ 40 years was 5.6 s for men and 5.9 s for women, and the mean 6-m walking time at the fastest pace was 3.7 s in men and 4.1 s in women. These walking speeds for both the usual pace and the fastest pace provide a baseline for clinical judgments of patient performance and could be used to determine which subjects would benefit from therapeutic intervention to improve locomotive function.

The Japanese Ministry of Education, Culture, Sports, Science and Technology published mean values for one-leg standing time with a maximum time of 120 s using 5,500 individuals (2,741 men, 2,759 women) with an age range of 65–79 years in each prefecture who participated in an examination of sporting ability, including walking ability [12]. They reported that mean one-leg standing times for men 65–69, 70–74, and 75–79 years old were 79.9, 66.5, and 50.5 s, respectively, and those for women were 80.8, 62.1, and 45.0 s, respectively. These values were measured up to 120 s, and ours were measured up to 60 s. Because the measuring method was different and their outcomes are means while our results are medians, the results cannot be compared directly. However, one-leg standing time was significantly lower with age in both studies. Again, establishment of reference values for physical performance, including walking and standing ability, would be useful for prediction of future disability.

We then evaluated associations between hand grip strength, muscle mass, walking time, and one-leg standing time, as indices reflecting components of locomotive syndrome and the presence of disability. We found that the 6-m walking time may be a useful index for detection of disability. To evaluate the independence of elderly persons in daily life, physical performance has been measured using various outcomes. Walking speed has been reported to be one important index that can predict future disability, hospitalisation, and mortality in the general geriatric population [20, 21]. In a Japanese population, Shinkai et al. [22] demonstrated that lower scores on baseline performance measures, particularly maximum walking speed, predicted an increased risk of onset of functional dependence, based on their 6-year follow-up of a cohort in a rural community consisting of 736 participants. In the present study, a 1-s slower normal walking time for 6 m was associated with a 15% increase in the presence of disability, and a 1-s slower fastest walking time for 6-m was associated with a 22% increase in the presence of disability. Our study evaluated

only walking ability and the presence of disability, not the occurrence of disability; however, we expect to follow these populations and clarify the predictive ability of walking speed for the occurrence of disability over the next few years.

On the other hand, no associations were found between indices such as hand grip strength, muscle mass, and one-leg standing time and the presence of disability. There is growing evidence that reduced hand grip strength is associated with adverse outcomes in older years, including morbidity, lower quality of life, higher fracture rates, increased length of hospital stay, and mortality [23–25]. Progressive decline in muscle mass has been defined as sarcopenia, which represents an impaired state of health associated with morbidity disorders, increased risk of falls and fractures, impaired ADL, loss of independence, and increased risk of death [6, 26–29]. Lang et al. [29] stated that loss of muscle mass and power increases the difficulties associated with procuring adequate nutrition and the effort required to undertake exercise; the combination of nutritional loss and reduced physical activity levels results in further loss of muscle mass and power. The resulting decrements in power, endurance, and physical performance lead to a loss of independence. In addition to muscle strength and mass, balance appears to be an important index of disability. Shinkai et al. [22] measured the one-leg standing time of 736 participants in a cohort established in a rural community, and the individuals in the lowest performance quartile had a significantly higher occurrence of disability.

Self-selection bias is suggested as a possible reason for the lack of associations between hand grip strength, muscle mass, and one-leg standing time and disability observed here, compared with previous reports. Self-selection bias is one type of sampling bias exhibited by subjects who voluntarily enrol in an epidemiological study. In this second visit of the ROAD study, volunteers who could walk to the clinic where the survey was performed, and could understand and sign an informed consent form, and who wanted to learn about their bone and joint conditions were welcomed. Therefore, the participants in the second survey may have been healthier than the general Japanese population. In fact, the estimated number of persons with disability in Japan using the age-gender prevalence of the second visit and the age-gender distribution of the Japanese population based on the national census in 2007 would be estimated at 1,510,000 (350,000 men, 1,160,000 women), considerably lower than the 4,940,000 reported by the government in 2010. Thus, self-selection bias likely affected the reference values; the reference values for hand grip strength, muscle mass, and one-leg standing time obtained from the present study may be higher, and walking speed faster, than the actual values. However, self-selection bias is somewhat unavoidable in such an

examination, because it is impossible to obtain measurements for individuals who cannot grasp a handgrip dynamometer or walk 6 m. This bias should be taken into consideration when reference values are used, including not only those obtained from the present study, but also from the national survey of physical strength and sporting ability published by the government.

In addition to self-selection bias, this study has several limitations. First, our results were obtained from a cross-sectional study of the second visit of the ROAD study; thus, we can not conclude causal relationships between such indices and disability, since some of the indices, such as muscle mass and one-leg standing time, were first introduced and performed during the second visit. We have begun the third visit of the ROAD study to clarify the relationships between physical performance and the occurrence of disability. Once the significance of indices reflecting components of locomotive syndrome can be determined as predictors for the occurrence of disability, appropriate thresholds can be developed as predictors of future disability. In addition, because of the lack of sufficient information, we could not determine the disorders that caused the disability. Thus, the disabled status of the participants in the present study might have been affected by various diseases such as cardiovascular diseases, dementia, or other diseases. However, regardless of the cause of disability, we found that walking ability was significantly associated with the presence of disability.

Conclusions

We have established reference values for hand grip strength, muscle mass, walking time, and one-leg standing time using data for a large-scale population-based cohort, and identified gender and age differences in the reference values. In addition, we determined that walking ability, including walking time and walking speed at the usual and maximum pace, was significantly associated with the presence of disability.

Acknowledgments This work was supported by Grants-in-Aid for Scientific Research B20390182 (Noriko Yoshimura), C20591737 (Toru Akune), and C20591774 (Shigeyuki Muraki); for Young Scientists A18689031 (Hiroyuki Oka); and Collaborating Research with NSF 08033011-00262 (Director, Noriko Yoshimura) from the Ministry of Education, Culture, Sports, Science, and Technology; it was also supported by grants H17-Men-eki-009 (Director, Kozo Nakamura), H18-Choujyu-037 (Director, Toshitaka Nakamura), and H20-Choujyu-009 (Director, Noriko Yoshimura), H23-Choujyu-002 (Director, Toru Akune) from the Ministry of Health, Labour, and Welfare of Japan. Further, this study was supported by grants from the Japan Osteoporosis Society (Noriko Yoshimura), Nakatomi Foundation (Noriko Yoshimura), and research aid from the Japanese Orthopaedic Association (JOA-Subsidized Science Project Research 2006-1, 2010-2, Director, Hiroshi Kawaguchi). The authors wish to

thank Mrs. Tomoko Takijiri, Mrs. Kumiko Shinou, Mrs. Rie Takiguchi, Mrs. Kyoko Maeda, Ms. Ikuyo Ueyama, Mrs. Michiko Mori, Mrs. Hisayo Sugimoto, and other members of the public office in Hidakagawa Town; and Mrs. Tamako Tsutsumi, Mrs. Kanami Maeda, Mr. Shoichi Shimoichi, Mrs. Megumi Takino, Mrs. Shuko Okada, Mrs. Kazuyo Setoh, Mrs. Chise Ryouno, Mrs. Miki Shimosaki, Mrs. Chika Yamaguchi, Mrs. Yuki Shimoji, and other members of the public office in Taiji Town for their assistance in locating and scheduling participants for examination. We also thank Ms. Kyoko Yoshimura, Mrs. Toki Sakurai, and Mrs. Saeko Sahara for their assistance in data reduction and administration.

Conflict of interest No conflict of interest has been declared by the authors.

References

1. Ministry of Health, Labour and Welfare. The outline of the results of National Livelihood Survey 2007. Available at <http://www.mhlw.go.jp/toukei/list/20-19-1.html>.
2. Muraki S, Yamamoto S, Ishibashi H, Nakamura K. Factors associated with mortality following hip fracture in Japan. *J Bone Miner Metab.* 2006;24:100–4.
3. Yoshimura N, Muraki S, Oka H, Mabuchi A, En-yo Y, Yoshida M, Saika A, Suzuki T, Yoshida H, Kawaguchi H, Nakamura K, Akune T. Prevalence of knee osteoarthritis, lumbar spondylosis and osteoporosis in Japanese men and women: the Research on Osteoarthritis/Osteoporosis Against Disability study. *J Bone Miner Metab.* 2009;27:620–8.
4. Nakamura K. Locomotive syndrome in an aging society. *J Jpn Orthop Assoc.* 2008;82:1–2. in Japanese.
5. Ministry of Internal Affairs and Communications. Vital statistics in Japan: The number of inhabitants and households based on resident registration in Japan 2010. Available at http://www.soumu.go.jp/menu_news/s-news/01gyosei02_01000001.html (in Japanese).
6. Cruz-Jentoft AJ, Baeyens JP, Bauer JM, Boirie Y, Cederholm T, Landi F, Martin FC, Michel JP, Rolland Y, Schneider SM, Topinková E, Vandewoude M, Zamboni M. European consensus on definition and diagnosis: report of the European Working Group on Sarcopenia in Older People. *Age Ageing.* 2010;39:412–23.
7. Ostir GV, Kuo YF, Berges IM, Markides KS, Ottenbacher KJ. Measures of lower body function and risk of mortality over 7 years of follow-up. *Am J Epidemiol.* 2007;166:599–605.
8. Michikawa T, Nishiwaki Y, Takebayashi T, Toyama Y. One-leg standing test for elderly populations. *J Orthop Sci.* 2009;14:675–85.
9. Oka H, Muraki S, Akune T, Nakamura K, Kawaguchi H, Yoshimura N. Normal and threshold values of radiographic parameters for knee osteoarthritis using a computer-assisted measuring system (KOACAD): The ROAD Study. *J Orthop Sci.* 2010;15:784–9.
10. Miyatani M, Kanehisa H, Masuo Y, Ito M, Fukunaga T. Validity of estimating limb muscle volume by bioelectrical impedance. *J Appl Physiol.* 2001;91:386–94.
11. Ministry of Health, Labour and Welfare. Long-term care insurance in Japan. Available at <http://www.mhlw.go.jp/english/topics/elderly/care/index.html>.
12. Ministry of Education, Culture, Sports, Science and Technology. The outline of the results of physical strength and sporting ability 2008. Available at http://www.mext.go.jp/b_menu/toukei/001/1285611.htm (in Japanese).

13. Petrobelli A, Morini P, Battistini N, Chiumello G, Nuñez C, Heymsfield SB. Appendicular skeletal muscle mass: prediction from multiple frequency segmental bioimpedance analysis. *Eur J Clin Nutr.* 1998;52:507–11.
14. Janssen I, Heymsfield SB, Baumgartner RN, Ross R. Estimation of skeletal muscle mass by bioelectrical impedance analysis. *J Appl Physiol.* 2000;89:465–71.
15. Yamada Y, Kimura M, Nakamura E, Masuo Y, Oda S. Limb muscle mass decrease with aging in Japanese men and women aged 15–97 yr. *Jpn J Phys Fitness Sports Med.* 2007;56:461–72. (in Japanese).
16. Miyatani M, Kanehisa H, Azuma K, Kuno S, Fukunaga T. Site-related differences in muscle loss with aging: a cross-sectional survey on the muscle thickness in Japanese men aged 20 to 79 years. *Int J Sport Health Sci.* 2003;1:34–40.
17. Hinmann JE, Cunningham DA, Rechnitzcr PA, Paterson DH. Age related changes in speed of walking. *Med Sci Sports Exerc.* 1988;20:161–6.
18. Bohannon RW. Comfortable and maximum walking speed of adults aged 20–79 years: reference values and determinants. *Age Ageing.* 1997;26:15–9.
19. Takahashi S, Toriida M, Tayama H. A study on the standard of gait evaluation—by an actual investigation of crosswalks. *Rigaku Ryoho Gaku.* 1989;16:261–6. in Japanese.
20. Cesari M, Kritchevsky SB, Penninx BW, Nicklas BJ, Simonsick EM, Newman AB, Tylavsky FA, Brach JS, Satterfield S, Bauer DC, Visser M, Rubin SM, Harris TB, Pahor M. Prognostic value of usual gait speed in well-functioning older people—results from the Health, Aging and Body Composition Study. *J Am Geriatr Soc.* 2005;53:1675–80.
21. Guralnik JM, Ferrucci L, Simonsick EM, Salive ME, Wallace RB. Lower-extremity function in persons over the age of 70 years as a predictor of subsequent disability. *N Engl J Med.* 1995;332:556–61.
22. Shinkai S, Watanabe S, Kumagai S, Fujiwara Y, Amano H, Yoshida H, Ishizaki T, Yukawa H, Suzuki T, Shibata H. Walking speed as a good predictor for the onset of functional dependence in a Japanese rural community population. *Age Ageing.* 2000;29:441–6.
23. Muraki S, Akune T, Oka H, En-yo Y, Yoshida M, Saika A, Suzuki T, Yoshida H, Ishibashi H, Tokimura F, Yamamoto S, Nakamura K, Kawaguchi H, Yoshimura N. Association of radiographic and symptomatic knee osteoarthritis with health-related quality of life in a population-based cohort study in Japan: The ROAD Study. *Osteoarthr Cartil.* 2010;18:1227–34.
24. Sirola J, Rikkonen T, Tuppurainen M, Jurvelin JS, Kroger H. Association of grip strength change with menopausal bone loss and related fractures: a population based follow-up study. *Calcif Tissue Int.* 2006;78:218–26.
25. Bohannon RW. Hand-grip dynamometry predicts future outcomes in aging adults. *J Geriatr Phys Ther.* 2008;31:10–313.
26. Lauretani F, Russo CR, Bandinelli S, Bartali B, Cavazzini C, Di Iorio A, Corsi AM, Rantanen T, Guralnik JM, Ferrucci L. Age-associated changes in skeletal muscles and their effect on mobility: an operational diagnosis of sarcopenia. *J Appl Physiol.* 2003;95:1851–60.
27. Rolland Y, Czerwinski S, Abellan Van Kan G, Morley JE, Cesari M, Onder G, Woo J, Baumgartner R, Pillard F, Boirie Y, Chumlea WM, Vellas B. Sarcopenia: its assessment, etiology, pathogenesis, consequences and future perspectives. *J Nutr Health Aging.* 2008;12:433–50.
28. Topinková E. Aging, disability and frailty. *Ann Nutr Metab.* 2008;52(Suppl 1):s6–11.
29. Lang T, Streeper T, Cawthon P, Baldwin K, Taaffe DR, Harris TB. Sarcopenia: etiology, clinical consequences, intervention, and assessment. *Osteoporos Int.* 2010;21:543–59.

Biochemical markers of bone turnover as predictors of osteoporosis and osteoporotic fractures in men and women: 10-year follow-up of the Taiji cohort

Noriko Yoshimura · Shigeyuki Muraki ·
Hiroyuki Oka · Hiroshi Kawaguchi ·
Kozo Nakamura · Toru Akune

Received: 24 December 2010 / Accepted: 17 March 2011 / Published online: 22 April 2011
© Japan College of Rheumatology 2011

Abstract We aimed to assess the capacity of biochemical markers of bone turnover (BTMs) to predict bone loss, osteoporosis (OP), and osteoporotic fractures. We randomly selected 400 individuals (age 40–79 years in 1993; 50 of each gender and age stratum) from a list of registered residents. In the years 1993, 1996, 2000, and 2003, bone mineral density (BMD) of the spine and hip were measured by dual-energy X-ray absorptiometry. The BTMs assessed at baseline were serum intact osteocalcin (OC), total OC, bone-specific alkaline phosphatase, C-terminal propeptide of type I procollagen, N-terminal propeptide of type I procollagen (PINP), C-terminal cross-linking telopeptide of type I collagen generated by matrix metalloproteinase, C-terminal cross-linking telopeptide of type I collagen (beta-CTX), N-terminal cross-linking telopeptide of type I collagen (NTX), urinary pyridinoline, and deoxypyridinoline (DPD). For 307 completers, multivariate analysis after adjusting for confounders revealed that serum PINP levels in men [hazard ratio (HR) 2.80, $P < 0.05$] and serum PINP (HR 1.65, $P < 0.05$), beta-CTX (HR 1.80, $P < 0.001$), NTX (HR 1.96, $P < 0.01$), and urinary DPD levels (HR

1.40, $P < 0.05$) in women were significantly related to the occurrence of spinal OP. In addition to adjustment for the baseline status of BMD, i.e., osteopenia or normal range, PINP, beta-CTX, and NTX in women could significantly predict the future occurrence of spinal OP. BTMs were not significant predictors of bone loss, femoral OP, or osteoporotic fractures. In conclusion, various BTMs in women can predict the occurrence of spinal OP.

Keywords Biochemical markers of bone turnover · Bone resorption · Bone formation · Bone mineral density · Osteoporotic fracture

Introduction

Osteoporosis (OP) impairs the activities of daily living (ADL) and quality of life (QOL), leading to increased morbidity and mortality in the elderly [1, 2]. With the rapid aging of the population, an urgent need has been identified for the development of methods to prevent OP. The estimated number of patients with OP in Japan is about 11 million [3], and osteoporotic fractures are ranked fifth among the diseases responsible for disabilities requiring support in Japan [4].

As the restoration of diminished bone volume seems quite difficult to achieve, the early diagnosis of OP is the most valuable strategy for preventing osteoporotic fractures. However, the prediction of rapid bone loss, incidence of OP, and osteoporotic fractures remains difficult.

Biochemical markers of bone turnover (BTMs) reflect the status of bone metabolism in various processes coupled with bone resorption and formation, and are widely used in clinical situations to evaluate the efficacy of treatments for OP [5–8]. Several population-based epidemiological

N. Yoshimura (✉) · H. Oka
Department of Joint Disease Research, 22nd Century Medical
and Research Center, Graduate School of Medicine,
The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku,
Tokyo 113-8655, Japan
e-mail: YOSHIMURAN-ORT@h.u-tokyo.ac.jp

S. Muraki · T. Akune
Department of Clinical Motor System Medicine, 22nd Century
Medical and Research Center, Graduate School of Medicine,
The University of Tokyo, Tokyo, Japan

H. Kawaguchi · K. Nakamura
Department of Orthopaedic Surgery, Faculty of Medicine,
The University of Tokyo, Tokyo, Japan

studies have shown that BTMs can predict bone loss and the incidence of osteoporotic fractures in women [9–15], but the effectiveness of BTMs for predicting such epidemiological indices over the long-term, such as 10 years, is unclear. In addition, few reports, besides our own previous reports [16, 17], have evaluated BTM values and bone loss or osteoporotic fractures in men.

We established a cohort comprising men and women in a rural area in Japan, and followed this cohort for 10 years. The present study was performed for the purpose of evaluating the capacity of baseline urinary and serum concentrations of various BTMs to predict future bone loss and the occurrence of OP and osteoporotic fractures in men and women randomly selected from a rural population in Japan.

Subjects, materials, and methods

Establishment of the baseline cohort

Details of the cohort survey at the Japanese town of Taiji have already been reported [18–20], and the Taiji cohort is therefore described here only in brief. The town of Taiji is located in the southern coastal area of Wakayama Prefecture in the south-western area of the main island of Japan. A list of all inhabitants born between 1913 and 1952, and therefore aged between 40 and 79 years in 1993, was compiled based on resident registrations as of the end of 1992. A cohort of 2,261 inhabitants (1,028 men, 1,233 women) was identified, and all members of the cohort completed a self-administered, 125-item questionnaire addressing lifestyle factors such as dietary habits, smoking habits, alcohol consumption, and physical exercise (whole cohort).

From this whole cohort, 50 men and 50 women in each of four age groups between 40 and 79 years by decade of birth year (1913–1922, 1923–1932, 1933–1942, and 1943–1952), for a total of 400 participants, were selected randomly, using a table of random numbers, and underwent measurement of bone mineral density (BMD) in 1993. At this time, blood and urine samples were taken from all participants. An interviewer administered a second questionnaire to these 400 participants, covering items of past medical history, including questions related to OP, osteoporotic fractures and falls, family history, calcium intake, dietary habits, physical exercise, occupational activities, and sun exposure, and, for women, additional questions about reproductive variables such as menstrual status (premenopause, perimenopause, or menopause), age at menopause, age at menarche, number of childbirths, lactation, use of estrogen for treatment, history of ovariectomy, and history of uterectomy (BMD cohort, baseline study).

All examinations were performed with the full consent of the participants. These studies were approved by the ethics committees of both Wakayama Medical University and the University of Tokyo.

BMD measurements at baseline and follow-up surveys

Baseline BMD was measured in 1993, using dual-energy X-ray absorptiometry (DXA) (QDR 1000; Hologic, Bedford, MA, USA), providing antero-posterior images of lumbar vertebrae L2–4 and the proximal femur (femoral neck, Ward's triangle, trochanter). These measurements were repeated on the same participants after 3 (2nd visit, 1996), 7 (3rd visit, 2000) and 10 years (4th visit, 2003).

At each follow-up survey, an interviewer-administered questionnaire survey was performed regarding changes in lifestyle factors during the observation period, and covering items of medical history, including questions related to OP, osteoporotic fractures and falls, anti-OP treatment, calcium intake, dietary habits, physical exercise, occupational activities, and sun exposure, and, for women, additional questions about reproductive variables such as menstrual status (premenopause, perimenopause, or menopause), age at menopause, use of estrogen for treatment, history of ovariectomy, and history of uterectomy.

To control for the precision of DXA, the equipment was checked at all examinations using the same phantom, and the BMD of the phantom was regulated to 1.030 ± 0.016 g/cm² (1.5%) during all examinations. The same physician (N.Y.) obtained all BMD measurements. Intra-observer variability for DXA scans done by this investigator was 0.35%, using the phantom as described [21].

Detection of the occurrence of OP

OP was defined based on World Health Organization criteria, according to which OP is diagnosed based on *T* scores of BMD ≤ 2.5 standard deviations (SDs) lower than peak bone mass [22]. The mean L2–4 BMD for young adult men and women measured using the Hologic QDR 1000 in Japan is reported as 1.011 g/cm², and the SD is 0.119 g/cm² [23]. The mean femoral neck BMD (SD) in Japan is reported as 0.863 (0.127) for young men and 0.787 (0.109) for young women [23]. This study therefore defined OP, using these indices, as lumbar spine BMD < 0.714 g/cm² for both men and women, and as femoral neck BMD < 0.546 g/cm² for men and < 0.515 g/cm² for women.

To define the incidence of OP among the 400 participants at the initial survey, individuals with spinal or femoral neck OP were excluded. Among the remaining participants without OP at the lumbar spine and/or femoral neck at baseline, the number of new cases of OP was

counted at the 3, 7, and 10-year follow-up surveys. Incidences of OP were estimated using the number of new cases divided by the person-years, consisting of years of individuals diagnosed with OP and years of drop-outs. The annual incidence of lumbar and femoral neck OP was then estimated.

Measurements of BTMs

All blood and urine samples were collected between 09:00 and 15:00. After centrifugation of the blood samples, sera were immediately placed in dry ice and transferred to a deep freezer within 24 h. Spot urine samples were frozen using the same procedure. These samples were kept at -80°C until needed for assays.

From the samples of participants in the baseline study, the following BTMs were measured to establish values. As markers of bone formation, serum intact osteocalcin (OC), serum total OC, serum bone-specific alkaline phosphatase (BAP), serum C-terminal propeptide of type I procollagen (PICP), and serum N-terminal propeptide of type I procollagen (PINP) were utilized. To monitor bone resorption, products of collagen breakdown, i.e., serum C-terminal cross-linking telopeptide of type I collagen generated by matrix metalloproteinase (ICTP), serum beta-isomerized C-terminal cross-linking telopeptide of type I collagen (beta-CTX), serum N-terminal cross-linking telopeptide of type I collagen (NTX), urinary pyridinoline cross-links of collagen (PYR), and urinary deoxypyridinoline cross-links of collagen (DPD) were used.

Reference values classified by age and gender for serum intact OC, PICP, and ICTP and urinary PYR and DPD have already been described [17]. Measurement methods for these compounds are therefore only described in brief. Serum intact OC was measured using an immunoradiometric assay (Osteocalcin IRMA kit; Mitsubishi Kagaku BCL, Tokyo, Japan) [24]. Serum PICP and ICTP were measured using a radioimmunoassay (RIA) (Orion Diagnostics, Espoo, Finland) [25]. Urinary PYR and DPD in hydrolyzed urine specimens were analyzed by high-performance liquid chromatography followed by fluorescent detection using essentially the same methods [26]. The values of these urinary markers were standardized to urinary creatinine concentrations.

Total OC was measured using an electrochemiluminescence immunoassay (ECLIA) (Elecsys N-MID Osteocalcin; Roche Diagnostics, Mannheim, Germany) [27] with an intraassay coefficient of variation (CV) of 0.5%, and sensitivity of 0.5 ng/mL. We measured BAP using an enzyme immunoassay (Metra BAP; Quidel, San Diego, CA, USA) [28] with an intraassay CV of 3.9–5.2% and sensitivity of 0.7 U/L. Serum PINP was measured using an

RIA (Orion Diagnostics) with an intraassay CV of 3.1–9.3% and sensitivity of 2 ng/mL [29, 30].

As markers of bone resorption, serum beta-CTX was measured using an ECLIA (Elecsys beta-CrossLaps; Roche Diagnostics) with an intraassay CV of 2.0% and sensitivity of 0.01 ng/mL [27]. Serum NTX was measured using an enzyme-linked immunosorbent assay (Osteomark NTX serum; Ostex International, Seattle, WA, USA) [31, 32] with an intraassay CV of 4.6% and sensitivity of 3.2 nM BCE/L.

Fracture assessment

All participants completed a detailed questionnaire at baseline, including a history of fragility fractures (that is, fractures resulting from low-impact trauma) that had occurred since the age of 40 years. Thereafter, at each subsequent examination, information about the occurrence of fractures since the previous visit was extracted from interviewer-registered questionnaires and registered. Information about fractures considered to be osteoporotic was analyzed. Osteoporotic fractures comprised those of the spine, pelvis, ribs, distal radius, forearm, humerus, and hip that occurred in the absence of high-impact trauma.

Statistical analysis

All statistical analyses were performed using STATA statistical software (STATA, College Station, TX, USA). Differences were tested for significance using analysis of variance for comparisons among multiple groups and Scheffe's least significant difference test for pairs of groups. Causal relationships between bone changes and serum and urinary concentrations of BTMs at baseline were clarified using multiple regression analysis with the rate of change of BMD (% per year) as an objective factor, and values of BTMs (/SD) at baseline after adjusting for age, weight, and menstrual status (0 pre- and perimenopause; 1 menopause) in women at baseline. Causal relationships between the incidence of OP and osteoporotic fractures, and serum and urinary concentrations of BTMs at baseline were clarified using Cox proportional hazards modeling using the occurrence of OP (yes 1; no 0) and occurrence of osteoporotic fractures (yes 1; no 0) as objective factors, and BTM level (/SD) at baseline as an explanatory factor after adjusting for age and weight, and menstrual status in women at baseline. Regarding anti-OP drugs, during the observation period from 1993 to 2003, bisphosphonates such as alendronate and risedronate and selective estrogen receptor modulator (SERM) agents had not been approved for use in Japan for the treatment of OP. In addition, we asked participants, in the questionnaires at each follow-up at the 2nd, 3rd, and 4th visits, whether they

had been prescribed calcitonin or alfacalcidol, and confirmed that all participants without OP in our examinations had not been given these treatments. For this reason, we did not include the presence or absence of anti-OP treatment into the analysis as an adjustment factor. Moreover, we did not exclude from our analysis subjects with experience of using calcium supplements, because we decided that such supplements should be regarded as a kind of food intake and that the pharmacological effects on BTMs and BMD were small.

Results

Eligible participants and baseline characteristics

Table 1 shows the background data, including physical characteristics, for all 400 participants at baseline. The mean body weight of men in their 40s was significantly higher than that in men in their 50s, 60s, and 70s ($P < 0.05$), whereas that of women was significantly lower in their 40s, 60s, and 70s than during their 50s ($P < 0.05$).

Among the 400 participants at baseline, one man in his 60s refused to undergo blood and urinary examinations for BTMs. As a result, BTMs at baseline were examined in 399 participants (199 men, 200 women). At the 2nd examination, to evaluate changes in BMDs over 3 years, 369 (92.3%; 181 men, 188 women) of the 400 initially recruited individuals participated. At the 3rd examination,

to evaluate changes in BMDs and BTMs over 7 years, 338 (84.5%; 170 men, 168 women) of the 400 initial recruits participated. At the 4th (10-year) follow-up in 2003, 322 (80.5%; 153 men, 169 women) of the original 400 recruits participated. Among the 322 participants at the 4th follow-up, 6 men and 9 women who had missed the 3rd visit were excluded from analysis. Detailed reasons for drop-outs at each visit are summarized in Fig. 1. We also specifically searched for participants who had been treated for primary hyperparathyroidism, or who had undergone estrogen or steroid therapy for more than 3 months between 1993 and 2003, but no such individuals were identified. In addition, we confirmed that the main reason for illness or death was not attributable to osteoporotic fracture in any cases. The remaining 307 participants (76.8%; 147 men, 160 women) completed all examinations over the 10-year study period.

Changes in BMD over 10 years

Table 2 shows the initial mean values and rates of change in L2–4 BMD classified by sex and age stratum over the 10-year study period. BMD values at L2–4 for men in their 50s and 60s had slightly increased by the 10-year follow-up, but had decreased slightly for those in their 40s and 70s. By contrast, BMD at L2–4 had decreased in all age strata for women over the 10 years, at a mean rate of $-7.5\%/10$ years. BMD at the femoral neck had increased for men in their 40s and 50s, and had considerably increased for those in their 70s, while BMD at the same site

Table 1 Summary of participants’ characteristics at baseline, classified by age and gender

Birth cohort	Age group (years)	<i>n</i>	Age (years)	Height (cm)	Weight (kg)	BMI (kg/m ²)
Men						
1943–1952	40–49	50	44.2 (2.6)	168.8 (5.2)	69.0 (10.4)	24.2 (3.2)
1933–1942	50–59	50	54.8 (2.7)	165.6 (5.0) ^a	63.5 (9.4) ^a	23.1 (2.9)
1923–1932	60–69	50	64.6 (2.5)	163.0 (4.8) ^a	62.9 (9.6) ^a	23.6 (3.2)
1913–1922	70–79	50	74.0 (2.7)	160.7 (5.4) ^{a,b}	57.5 (8.3) ^{a,b,c}	22.2 (2.8)
1913–1952	40–79	200	59.4 (11.4)	164.5 (5.9)	63.2 (10.2)	23.3 (3.1)
Women						
1943–1952	40–49	50	44.0 (2.8)	154.4 (5.0)	54.1 (8.3)	22.7 (3.1)
1933–1942	50–59	50	55.8 (2.8)	154.9 (5.3)	59.4 (10.0) ^a	24.8 (4.0) ^a
1923–1932	60–69	50	64.8 (2.6)	151.1 (4.6) ^{a,b}	52.1 (9.1) ^b	22.8 (3.5) ^b
1913–1922	70–79	50	74.4 (2.8)	147.7 (5.4) ^{a,b,c}	48.4 (8.2) ^{a,b}	22.2 (3.4) ^b
1913–1952	40–79	200	59.8 (11.6)	152.0 (5.8)	53.5 (9.7)	23.1 (3.6)

Values are given as means with standard deviations in parentheses


BMI body mass index, *n* number of participants

^a Significantly different ($P < 0.05$) from values of participants in their 40s

^b Significantly different ($P < 0.05$) from values of participants in their 50s

^c Significantly different ($P < 0.05$) from values of participants in their 60s

Fig. 1 Flow chart of participants in the cohort. *BTM* Biochemical marker of bone turnover



in women had decreased in all age strata. Rates of change did not differ significantly among age strata.

Incidence of OP and osteoporotic fractures over 10 years

Among the 400 participants at the initial survey, 47 (9 men, 38 women) with spinal OP were excluded from estimation of the incidence of OP. Among the remaining 353 participants (191 men, 162 women), 29 (4 men, 25 women) developed OP at the lumbar spine over the 10-year period. Incidences of OP at the lumbar spine over the 10-year period in men and women aged 40–79 years were 23.8 and 176.0 per 10,000 person-years, respectively. Similarly, 22 (3 men, 19 women) of 383 participants (200 men, 183 women) developed OP at the femoral neck over the 10-year period. Incidences of OP at the femoral neck in men and women aged 40–79 years were 17.1 and 114.5 per 10,000 person-years, respectively. The annual incidence of lumbar

and femoral neck OP was thus approximately sevenfold higher among women than among men.

Incidence of OP classified by age was then examined in detail. Incidences of lumbar OP in men aged in their 40s, 50s, 60s, and 70s were 0, 22.1, 0, and 96.2 per 10,000 person-years, respectively, with a peak in the oldest stratum. By contrast, these values for women were 63.8, 205.5, 380.2, and 120.5 per 10,000 person-years, respectively, with peaks in the 50s and 60s. Incidences of OP at the femoral neck in men in their 40s, 50s, 60s, and 70s were 0, 0, 42.9, and 28.4 per 10,000 person-years, respectively, with peaks in the 60s and 70s. These values in women were 0, 20.4, 224.4, and 301.0 per 10,000 person-years, respectively, with the highest peak in the oldest stratum, followed by the 60s.

During the 10-year observation period, we detected 32 osteoporotic fractures (10 in men, 22 in women) after the exclusion of traumatic fractures (results of traffic accidents and falls from more than the subject's height). These 32

Table 2 Mean values (standard deviation) of bone mineral density (g/cm²) and change rate (%) at lumbar spine L2–4 and femoral neck over 3, 7, and 10 years, classified by age and gender

Birth cohort	Age group (years)	L2–4								Femoral neck			
		Baseline		2nd visit (3-year follow-up)		3rd visit (7-year follow-up)		4th visit (10-year follow-up)		Baseline	2nd visit	3rd visit	4th visit
		<i>n</i>	BMD (g/cm ²)	<i>n</i>	Change rate (%/3 years)	<i>n</i>	Change rate (%/7 years)	<i>n</i>	Change rate (%/10 years)	BMD (g/cm ²)	Change rate (%/3 years)	Change rate (%/7 years)	Change rate (%/10 years)
Men													
1943–1952	40–49	50	1.05 (0.15)	48	0.6 (3.8)	46	−0.6 (5.1)	42	−0.2 (5.8)	0.86 (0.09)	0.3 (4.6)	−1.8 (4.8)	−1.3 (10.9)
1933–1942	50–59	50	0.98 (0.17)	47	1.0 (3.3)	46	−0.0 (6.3)	43	2.1 (8.0)	0.80 (0.13) ^a	−0.2 (4.9)	0.7 (10.0)	−2.6 (6.8)
1923–1932	60–69	50	1.04 (0.21)	49	1.3 (3.6)	47	1.4 (7.1)	41	2.3 (9.4)	0.77 (0.11) ^a	1.0 (7.0)	−0.1 (9.3)	0.3 (12.5)
1913–1922	70–79	50	0.97 (0.19)	37	0.1 (5.3)	31	−1.2 (7.9)	21	−1.1 (9.2)	0.71 (0.08) ^{a,b,c}	0.9 (6.3)	4.6 (10.2)	7.1 (16.7) ^b
1913–1952	40–79	200	1.01 (0.18)	181	0.8 (4.0)	170	0.0 (6.6)	147	1.1 (8.1)	0.79 (0.12)	0.5 (5.7)	0.5 (8.9)	−0.4 (11.7)
Women													
1943–1952	40–49	50	1.07 (0.14)	48	−1.1 (4.2)	47	−8.2 (9.3)	45	−11.6 (10.0)	0.79 (0.10)	−1.7 (5.0)	−3.0 (9.5)	−8.8 (9.3)
1933–1942	50–59	50	0.92 (0.16) ^a	50	−3.1 (5.7)	47	−8.5 (9.4)	47	−7.9 (11.8)	0.70 (0.11) ^a	0.1 (4.9)	−4.5 (8.1)	−6.4 (9.4)
1923–1932	60–69	50	0.78 (0.17) ^{ab}	47	−0.3 (3.9) ^b	42	−3.9 (5.3)	39	−3.5 (7.1) ^a	0.62 (0.09) ^{ab}	1.5 (5.8)	−3.3 (8.3)	−5.0 (7.9)
1913–1922	70–79	50	0.77 (0.12) ^{ab}	43	−0.6 (4.9)	32	−2.8 (7.1) ^{ab}	29	−6.0 (9.4)	0.59 (0.10) ^{ab}	−0.7 (6.7)	−3.8 (9.1)	−6.8 (10.7)
1913–1952	40–79	200	0.89 (0.19)	188	−1.3 (4.8)	168	−6.2 (8.4)	160	−7.5 (10.2)	0.68 (0.13)	−0.2 (5.7)	−3.6 (8.7)	−6.8 (9.3)

BMD bone mineral density, *n* number of subjects

^a Significantly different ($P < 0.05$) from values of subjects in their 40s

^b Significantly different ($P < 0.05$) from values of subjects in their 50s

^c Significantly different ($P < 0.05$) from values of subjects in their 60s

osteoporotic fractures included 6 hip fractures (1 man, 5 women), 5 clinical vertebral fractures (1 man, 4 women), 2 wrist fractures (2 men), and 9 costal fractures (4 men, 5 women). Incidences of osteoporotic fractures in men in their 40s, 50s, 60s, and 70s were 20.8, 90.1, 44.4, and 82.6 per 10,000 person-years, respectively, and incidences in women were 20.2, 157.0, 158.2, and 170.3 per 10,000 person-years, respectively. Osteoporotic fractures tended to increase with age in women, whereas an age-related but non-significant tendency was seen in men.

Capacity of BTMs at baseline to predict rates of change of BMD and the occurrence of OP and osteoporotic fractures

Age-gender distributions of mean BTM levels at the initial survey are shown in Table 3. No significant difference was seen among age groups for BTM levels in men, while significant differences were seen for each marker between the 40s and 50–70s in women ($P < 0.05$ each).

Table 4 shows mean BTM levels in women at the initial survey classified by menstrual status, categorized into the following three groups: premenopausal group with regular period; perimenopausal group with irregular period; and postmenopausal group with no period within at least 1 year. No significant differences in BTM levels were seen between the pre- and perimenopausal groups, with the exception of serum PINP. Although a tendency toward increased levels in the perimenopausal group was seen compared to the premenopausal group, this was slight and non-significant. Conversely, all BTM levels measured in the present study were significantly higher in the postmenopausal group than in the premenopausal group ($P < 0.001$). In addition, serum total OC, PICP, PINP, beta-CTX, NTX, and urinary DPD were significantly lower in the perimenopausal group than in the postmenopausal group ($P < 0.05$). These results suggest that BTM levels were significantly accelerated after menopause.

We clarified whether values of BTMs at baseline could predict rates of change of BMD over 10 years, using multiple regression analysis with rate of change of BMD (%/year) as an objective factor, and values of BTMs (/SD) at baseline after adjusting for age, weight, and menstrual status (0 pre- and perimenopause; 1 menopause) in women at baseline. No associations were identified between any BTMs and rate of change in BMDs at L2–4 or the femoral neck over 10 years.

Table 5 shows the hazard ratios (HRs) of BTMs (/SD) for the incidence of OP at L2–4 and the femoral neck. Cox proportional hazards modeling using the occurrence of OP (yes 1; no 0) as an objective factor and BTM level (/SD) at baseline as an explanatory factor after adjusting for age and weight, and menstrual status in women, at baseline

identified only the PINP level as being significantly related to the occurrence of OP at L2–4 in men ($P < 0.05$). By contrast, serum PINP, beta-CTX, and NTX and urinary DPD levels were significantly related to the occurrence of OP at L2–4 in women (PINP, $P < 0.05$; beta-CTX, $P < 0.001$; NTX, $P < 0.01$; DPD, $P < 0.05$). In addition to the above-mentioned analysis, we then added the baseline BMD status, i.e., osteopenia or normal range, as an adjusted factor. Using the baseline L2–4 BMD, 36 men and 71 women were categorized into the group of spinal osteopenia. After adjusting for the baseline status of L2–4 BMD (1 osteopenia; 0 normal) in addition to age, weight, and menstrual status in women, the association between PINP and the occurrence of spinal OP was diluted in men (HR 3.88, 95% confidence interval [CI] 0.92–16.4, $P = 0.066$), but the association of the BTMs, with the exception of DPD, remained significant in women (PINP, HR 1.57, 95% CI 1.02–2.43, $P < 0.05$; beta-CTX, HR 1.99, 95% CI 1.30–3.05, $P < 0.01$; NTX, HR 1.68, 95% CI 1.69–2.65, $P < 0.05$; DPD, HR 1.42, 95% CI 0.99–2.04, $P = 0.056$). However, no BTMs were identified as significant predictors of the incidence of OP at the femoral neck in either men or women.

We estimated the HRs of BTMs (/SD) for the incidence of osteoporotic fracture by Cox proportional hazards modeling, using the occurrence of osteoporotic fractures (yes 1; no 0) as an objective factor, and BTM levels (/SD) at baseline as explanatory factors after adjusting for age and weight, and menstrual status in women, at baseline. No BTMs were identified as significant predictors of the incidence of osteoporotic fractures in men or women.

Discussion

The present study first clarified rates of bone loss at the lumbar spine and femoral neck over 10 years in the general population. BMD values for men had changed slightly by the 10-year follow-up, with the exception of femoral neck BMD for men in their 70s. Although the reason of the considerable increase in femoral neck BMD among men in their 70s is uncertain, it might be partially attributable to bone proliferative degeneration, such as hip osteoarthritis. Conversely, BMD values had decreased in all age strata for women over the 10 years, at an approximate mean rate of $-7\%/10$ years, at both L2–4 and the femoral neck. BMD values decreased most rapidly among women in their 40s, suggesting a menopausal effect, whereas rates of change did not differ significantly among age strata.

We then clarified the incidence of OP and osteoporotic fractures over the 10 years among the general population. We have previously reported the incidence of OP in individuals aged 40–79 years living in a mountain village [33].

Table 3 Mean values of biochemical markers of bone turnover for participants at baseline, classified by age and gender

Age group (years)	n	Sera								Urine	
		Intact OC (ng/mL)	Total OC (ng/mL)	BAP (U/L)	PICP (µg/L)	PINP (ng/mL)	ICTP (µg/L)	β-CTX (ng/mL)	NTX (nmol BCE/L)	PYR (pmol/µmol Cr)	DPD (pmol/µmol Cr)
Men											
40–49	50	3.63 (1.70)	18.8 (7.5)	26.1 (9.3)	126.8 (36.1)	41.4 (17.3)	2.85 (1.71)	0.190 (0.107)	13.3 (2.8)	18.0 (6.5)	3.02 (1.28)
50–59	50	3.80 (4.28)	19.7 (18.3)	26.7 (16.6)	115.5 (33.0)	39.4 (28.1)	2.61 (0.83)	0.197 (0.162)	14.0 (6.0)	18.9 (7.7)	3.24 (2.41)
60–69	49	3.58 (1.55)	16.4 (6.1)	26.4 (7.1)	113.1 (28.5)	34.5 (13.3)	3.00 (1.07)	0.174 (0.107)	13.6 (4.1)	19.0 (7.5)	2.81 (0.97)
70–79	50	3.41 (1.77)	18.9 (8.1)	26.9 (10.7)	121.8 (34.3)	37.4 (16.0)	3.39 (1.06) ^b	0.187 (0.099)	13.5 (3.4)	21.5 (4.9)	3.17 (0.94)
40–79	199	3.60 (2.54)	18.5 (11.1)	26.5 (10.7)	119.3 (33.3)	38.2 (19.5)	2.96 (1.24)	0.187 (0.121)	13.6 (4.2)	19.3 (6.8)	3.06 (1.53)
Women											
40–49	50	3.58 (1.72)	14.9 (5.7)	19.7 (5.2)	96.1 (25.6)	30.6 (13.3)	2.51 (0.60)	0.103 (0.066)	11.6 (2.3)	20.7 (5.9)	3.20 (1.18)
50–59	50	5.68 (2.42) ^a	28.1 (8.8) ^a	30.2 (7.5) ^a	130.2 (41.1) ^a	57.8 (20.9) ^a	3.07 (0.62) ^a	0.255 (0.121) ^a	15.4 (3.3) ^a	27.5 (5.3) ^a	5.05 (1.34) ^a
60–69	50	6.61 (3.10) ^a	32.6 (12.4) ^a	32.3 (11.2) ^a	136.2 (35.7) ^a	59.2 (20.0) ^a	3.51 (1.25) ^a	0.301 (0.136) ^a	17.8 (4.4) ^{a,b}	32.0 (10.8) ^a	5.79 (2.14) ^a
70–79	50	6.09 (3.51) ^a	28.3 (10.3) ^a	32.2 (10.8) ^a	143.3 (39.2) ^a	52.8(20.0) ^a	3.56 (1.22) ^a	0.275 (0.153) ^a	16.0 (3.2) ^a	29.7 (8.9) ^a	4.95 (1.84) ^a
40–79	200	5.60 (3.01)	26.0 (11.6)	28.6 (10.4)	126.4 (40.0)	50.1 (21.9)	3.16 (1.06)	0.234 (0.145)	15.2 (4.0)	27.5 (9.0)	4.76 (1.91)

Values are given as means with standard deviations in parentheses

OC osteocalcin, BAP bone-specific alkaline phosphatase, PICP C-terminal propeptide of type I procollagen, PINP N-terminal propeptide of type I procollagen, ICTP C-terminal cross-linking telopeptide of type I collagen generated by matrix metalloproteinase, β-CTX β-isomerized C-terminal cross-linking telopeptide of type I collagen, NTX N-terminal cross-linking telopeptide of type I collagen, PYR pyridinoline cross-links of collagen, DPD deoxypyridinoline cross-links of collagen, Cr creatinine

^a Significantly different ($P < 0.05$) from values of participants in their 40s

^b Significantly different ($P < 0.05$) from values of participants in their 50s

Table 4 Mean values of biochemical markers of bone turnover at baseline classified by menstrual status in women

Menstrual status	n	Age (years)	Sera					Urine				
			Intact OC (ng/mL)	Total OC (ng/mL)	BAP (U/L)	PICP (μg/L)	PINP (ng/mL)	ICTP (μg/L)	β-CTX (ng/mL)	NTX (nmol BCE/L)	PYR (pmol/μmol Cr)	DPD (pmol/μmol Cr)
Premenopause	41	44.2 (3.0)	3.35 (1.28)	14.5 (4.4)	19.0 (5.1)	96.5 (26.6)	29.5 (11.1)	2.49 (0.56)	0.099 (0.060)	11.3 (2.0)	20.7 (4.9)	3.12 (0.89)
Perimenopause	14	47.4 (4.9)	4.83 (2.75)	20.4 (10.2)	25.7 (7.0)	94.4 (24.8)	45.2 (25.9) ^a	2.88 (0.82)	0.165 (0.120)	13.0 (2.8)	24.1 (7.8)	4.09 (1.84)
Menopause	145	65.3 (8.1) ^{a,b}	6.20 (3.07) ^a	29.8 (10.8) ^{a,b}	31.6 (10.1) ^a	138.0 (38.3) ^{a,b}	56.4 (20.2) ^{a,b}	3.38 (1.10) ^a	0.278 (0.138) ^{a,b}	16.5 (3.7) ^{a,b}	29.7 (9.0) ^a	5.28 (1.86) ^{a,b}
Total	200	59.7 (11.6)	5.60 (3.01)	26.0 (11.6)	28.6 (10.4)	126.4 (40.0)	50.1 (21.9)	3.16 (1.06)	0.234 (0.145)	15.2 (4.0)	27.5 (9.0)	4.76 (1.91)

Values are given as means with standard deviations in parentheses

n number of subjects, OC osteocalcin, BAP bone-specific alkaline phosphatase, PICP C-terminal propeptide of type I procollagen, PINP N-terminal propeptide of type I procollagen, ICTP C-terminal cross-linking telopeptide of type I collagen generated by matrix metalloproteinase, β-CTX β-isomerized C-terminal cross-linking telopeptide of type I collagen, NTX N-terminal cross-linking telopeptide of type I collagen, PYR pyridinoline cross-links of collagen, DPD deoxypyridinoline cross-links of collagen, Cr creatinine

^a Significantly different ($P < 0.05$) from values of participants in the premenopausal group

^b Significantly different ($P < 0.05$) from values of participants in the perimenopausal group

Estimated incidences of lumbar spine OP in men and women in that village over 10 years were 55.6 and 231.7 per 10,000 person-years, respectively. The incidences of spinal OP in the present study were approximately half of those found in the mountain village, indicating regional differences in the incidence of OP among Japanese populations. We have already reported on regional differences between cohorts in mountainous and seaside (the present cohort) areas [34–36]. Residents of the mountainous area tended to show lower BMDs [34], and proportions of fast bone losers were higher [35] than those in residents of the seaside area. In addition, we found that levels of BTMs, including serum intact OC and urinary PYR and DPD, in the residents of the mountainous area were significantly higher than those in residents of the seaside area [36]. These differences in the incidence of OP between regions support the concept of regional differences in bone metabolism in Japan, and might be due to environmental differences. From meteorological data in 1990, when the cohort from the mountainous area was started, the total annual duration of sunlight exposure was 1340.9 h in the mountainous area, lower than the 2224.4 h in the seaside area. In addition, based on the results of a nutrition survey of Wakayama Prefecture in 1993, total calcium intake for inhabitants of the mountainous area was 542 mg/day, compared to 563 mg/day for the seaside area [37]. However, this finding was only a result of an ecological study, not a direct survey of participants from our two cohorts. In addition, cohorts in both studies comprised only 400 individuals each, so these differences need to be confirmed in larger population-based cohorts. We have therefore established larger-scale cohorts based on the cohort in the present study, entitled the ROAD study [3, 38], in which BMD and X-ray examinations were performed in all 1,690 participants, and serum and urinary samples were collected. This enlarged population-based cohort study may confirm regional differences in bone metabolism and in OP and osteoporotic fractures.

The present study found no significant differences in baseline levels of the various BTMs among men in any age groups, with the exception of serum ICTP. By contrast, each marker showed significant differences between women in their 40s and those in their 50–70s ($P < 0.05$). In addition, the values of the BTMs in women started to increase in the perimenopausal period, with rapidly accelerating elevations after menopause, according to estrogen deficiency. We had already measured levels of endogenous estrogen and sex hormone-binding globulin in the present cohort, and reported serum estradiol (E2) and BMD levels among postmenopausal women at the 3-year follow-up [39], but cannot confirm any association between BTMs and endogenous sex steroids yet. Further studies to clarify levels of E2 and BTMs in women are warranted. In