









## Analysis results on infants born in 2010






I-12/16





No.	Resources of participating hospitals	All hospitals	n
<b>I</b>			
<b>Gastrointestinal problem</b>			
1101	Intravenous hyperalimentation (among infants with live birth and remained)	 <p>1:Yes 72% 2:No 28%</p>	3591
1102	NEC (among infants with live birth and remained)	 <p>1:Yes 2% 2:No 98%</p>	3591
1103	Idiopathic intestinal perforation (among infants with live birth and remained)	 <p>1:Yes 3% 2:No 97%</p>	3591
<b>J</b>			
<b>Hearing screening</b>			
1201	Hearing loss screening (among infants with live birth and remained)	 <p>1:Pass 75% 2:Refer 8% 3:not done 17%</p>	3591
<b>K</b>			
<b>Retinopathy of prematurity</b>			
1301	ROP(worst stage) (among infants with live birth and remained)	 <p>1:&lt;II 66% 2:III(early) 9% 3:III(intermediate) 9% 4:III(late) 2% 5:not done 15%</p>	3591

## Analysis results on infants born in 2010

I-13/16

No.	Resources of participating hospitals	All hospitals	n
1302	Treatment for ROP (among infants with live birth and remained)	 <p>1:Yes 14% 2:No 86%</p>	3591
<b>L</b>			
<b>Diagnosis</b>			
1411	Congenital anomaly	 <p>1:Yes 7% 2:No 93%</p>	3689
1412	Diagnosis of congenital anomaly (among infants with congenital anomaly)		264
1413	Operation for congenital anomaly (among infants with live birth, remained and congenital anomaly)	 <p>1:Yes 25% 2:No 75%</p>	252
<b>M</b>			
<b>Summary</b>			
1501	Age at enteral feeding exceed 100ml/kg (median) (among infants with live birth and remained)	11.0	3094
	lower quartile	8.0	
	upper quartile	16.0	

Analysis results on infants born in 2010		I-14/16	
No.	Resources of participating hospitals	All hospitals	n
1511	Blood transfusion (among infants with live birth and remained)	 <p>1: Yes 31% 2: No 69%</p>	3591
1512	Erythropoietin (among infants with live birth and remained)	 <p>1: Yes 71% 2: No 29%</p>	3591
<b>N</b>	<b>Condition at discharge</b>		
1601	Age at discharge (mean) (among infants with live birth and remained)	86.4	3573
	SD	55.1	
	95% confidence interval	84.6-88.2	
1602A	Dead at discharge (among infants with live birth and remained)	 <p>1: Yes 7% 2: No 93%</p>	3591
1602B	Dead at discharge (among infants with live birth)	 <p>1: Yes 7% 2: No 93%</p>	3610
1603	Autopsy (among infants with live birth, remained and dead at discharge)	 <p>1: Yes 15% 2: No 85%</p>	250

Analysis results on infants born in 2010		I-15/16	
No.	Resources of participating hospitals	All hospitals	n
1604	Cause of death (among infants with live birth, remained and dead at discharge)	<p>90 63Number 10 44Number 31 34Number 50 19Number 41 10Number</p>	212
1605	Discharge home (among infants with live birth, remained and alive at discharge)	 <p>1: Yes 87% 2: No 13%</p>	3341
1606	Disposition (among infants with live birth, remained, alive at discharge, and transferred)	 <p>1: Delivered hospital 64% 2: Other NICU 13% 3: Pediatric ward 17% 4: Other hospital 1% 5: Facility for disabled children 1% 6: Orphanage 2%</p>	429
1607	HOT (among infants with live birth, remained and alive at discharge)	 <p>1: Yes 6% 2: No 94%</p>	3341
1608	Tracheostomy (among infants with live birth and alive at discharge)	 <p>1: Yes 1% 2: No 99%</p>	3341
1609	Body weight at discharge (mean) (among infants with alive at discharge)	2614.0	3546
	SD	840.5	
	95% confidence interval	2586.4-2641.7	



Analysis results on infants born in 2010





I-16/16

No.	Resources of participating hospitals	All hospitals	n
1610	Body length at discharge (mean) (among infants with alive at discharge)	45.7	3322
	SD	5.0	
	95% confidence interval	45.6-45.9	
1611	Head circumference at discharge (mean) (among infants with alive at discharge)	33.8	3308
	SD	3.2	
	95% confidence interval	33.7-33.9	

Analysis results on infants born in 2010

II-1/56






資料21

Analysis results on infants born in 2010



II-4/56



資料22

Analysis results on infants born in 2010


II-5/56



資料23

Analysis results on infants born in 2010




II-6/56







資料24

Analysis results on infants born in 2010

II-7/56






資料27

Analysis results on infants born in 2010

II-10/56




among infants with positive histologic CAM


資料28

Analysis results on infants born in 2010

II-11/56




Grade of histologic CAM







among infants with positive histologic CAM

PROM








資料31

Analysis results on infants born in 2010




II-14/56







資料32

Analysis results on infants born in 2010

II-15/56






資料35

Analysis results on infants born in 2010


II-18/56





資料36

Analysis results on infants born in 2010





II-19/56



among infants with inborn




among infants with inborn



資料39

Analysis results on infants born in 2010


II-22/56




資料40

Analysis results on infants born in 2010

II-23/56



among infants with live birth and remained




among infants with live birth and remained

資料41

Analysis results on infants born in 2010

II-24/56




among infants with live birth and remained

資料42


Analysis results on infants born in 2010

II-25/56




among infants with live birth and remained


—211—




among infants with live birth and remained




among infants with live birth and remained




among infants with live birth, remained and mechanical ventilation



among infants with live birth, remained and alive at 28 days of age




among infants with live birth, remained and mechanical ventilation




among infants with live birth, remained and alive at 28 days of age







among infants with CLD




among infants with live birth, remained, alive at 36 wk(corrected age), and CLD




among infants with CLD




among infants with live birth, remained, alive at 36 wk(corrected age), and CLD




among infants with live birth and remained




among infants with live birth and remained




among infants with live birth and remained




among infants with live birth and remained




among infants with symptomatic PDA



among infants with live birth, remained and alive at 7 d



among infants with symptomatic PDA




among infants with live birth, remained and alive at 7 d

資料51

Analysis results on infants born in 2010

II-34/56




among infants with live birth and remained


資料52

Analysis results on infants born in 2010


II-35/56



among infants with live birth and remained



among infants with live birth and remained



among infants with live birth and remained