








among infants with inborn




among infants with inborn









among infants with live birth and remained




among infants with live birth and remained




among infants with live birth and remained




among infants with live birth and remained




among infants with live birth and remained



among infants with live birth and remained




among infants with live birth and remained




among infants with live birth and remained







among infants with live birth, remained and mechanical ventilation




among infants with live birth, remained and alive at 28 days of age




among infants with live birth, remained and mechanical ventilation




among infants with live birth, remained and alive at 28 days of age




among infants with CLD




among infants with live birth, remained, alive at 36 wk(corrected age), and CLD




among infants with CLD




among infants with live birth, remained, alive at 36 wk(corrected age), and CLD




among infants with live birth and remained




among infants with live birth and remained




among infants with live birth and remained




among infants with live birth and remained




among infants with symptomatic PDA



among infants with live birth, remained and alive at 7 d




among infants with symptomatic PDA



among infants with live birth, remained and alive at 7 d

Analysis results on infants born in 2010


IV-31/53




among infants with live birth and remained

Analysis results on infants born in 2010


IV-32/53



among infants with live birth and remained




among infants with live birth and remained



among infants with live birth and remained




Analysis results on infants born in 2010


IV-33/53




Analysis results on infants born in 2010

IV-34/53







among infants with live birth and remained



among infants with live birth and remained




among infants with live birth and remained



among infants with live birth and remained

Analysis results on infants born in 2010


IV-37/53




among infants with live birth and remained

Analysis results on infants born in 2010


IV-38/53



among infants with live birth and remained




among infants with live birth and remained




among infants with live birth and remained







among infants with live birth and remained




among infants with live birth and remained




among infants with live birth and remained




among infants with live birth and remained




among infants with live birth and remained




among infants with live birth and remained





among infants with live birth and remained




among infants with live birth and remained




among infants with live birth and remained




among infants with live birth and remained







among infants with live birth, remained and congenital anomaly



among infants with live birth and remained



among infants with live birth, remained and congenital anomaly



among infants with live birth and remained