

Figure 1. (a) Structure of a *c19orf10* gene and a *c19orf10* protein. The DNA sequence of *c19orf10* and amino acid alignment of the encoded *c19orf10* protein are shown. *C19orf10* is predicted to have a molecular weight of 17 kDa and contain a signal peptide cleavage site (indicated as a black arrow). (b) *C19orf10* gene expression profiles in various tissues by SAGE. Y-axis indicates the number of tags corresponding to *c19orf10* in each tissue. (c, d) RTD-PCR analysis of *c19orf10*. RNA was isolated from 34 tissue samples: 15 HCC, 15 corresponding noncancerous liver samples and four normal liver samples. Differential expression of each gene among normal liver tissues, noncancerous liver tissues and HCC tissues was examined using the Kruskal–Wallis test and unpaired *t*-test. The mean value of gene expression data in each group is indicated (c). *C19orf10* was overexpressed in 10 of 15 examined HCC tissues compared to the noncancerous liver tissues (d).

sequence presumably encoded a secretory protein with a signal peptide sequence (Fig. 1a).

When we examined the expression profiles of *c19orf10* using retrieved SAGE data from various cancers and their normal counterparts, we identified that *c19orf10* was abundantly expressed in human HCC (Fig. 1b). We further examined the publicly available EST profiles of *c19orf10* (<http://www.ncbi.nlm.nih.gov/unigene>) and confirmed its tendency to be overexpressed in HCC compared to the normal liver (data not shown). We validated the overexpression of *c19orf10* in 15 independent HCC tissues and adjacent noncancerous liver tissues by RTD-PCR. Gene expression of *c19orf10* was significantly higher in the HCC tissues than in

the normal liver tissues and adjacent noncancerous liver tissues ($p = 0.014$ and 0.048 , respectively; Fig. 1c). *C19orf10* expression was elevated in HCC tissues compared to the adjacent noncancerous liver tissues in 10 of 15 patients (66.7%; Fig. 1d).

Overexpression of *C19ORF10* in AFP-positive HCC

As HCC is a heterogeneous mixture of cancer epithelial cells and stromal cells, and a previous report indicated that *c19orf10* is expressed in fibroblast-like synoviocytes. We, therefore, evaluated the expression of *c19orf10* in tumor epithelial cells and stromal cells separately using LCM and RTD-PCR in 20 HCC tissues (Fig. 2a). Although tumor




Figure 2. (a) Representative photomicrographs of an HCC tissue used for LCM (toluidine blue staining). Inflammatory mononuclear cells and stromal cells were separately captured (left: Pre-LCM, right: Post-LCM). (b) RTD-PCR analysis of *c19orf10* expression in inflammatory mononuclear cells and tumor epithelial cells in 20 HCV-related HCC tissues. Tumor-inflammatory mononuclear cells and stromal cells were isolated using LCM. RNAs were isolated from these cells as well as parenchymal tissues from the same liver, followed by RTD-PCR for *c19orf10* gene expression. Expression of the *c19orf10* gene was higher than that observed in HCC-infiltrating inflammatory mononuclear cells. * $p < 0.05$. (c-e) Scatter plot analysis of *c19orf10*, *AFP*, *KRT19* and *GPC3* expression in HCC. RNA was isolated from 17 HBV-related HCC and 19 HCV-related HCC. (f) RTD-PCR analysis of *c19orf10* in AFP-negative (HLE and HLF) and -positive (HuH1, HuH7 and Hep3B) liver cancer cell lines.

stromal cells expressed *c19orf10* at some level, the expression levels were significantly higher in tumor epithelial cells than in stromal cells ($p = 0.006$) (Fig. 2b).

To explore the relationship of *c19orf10* with other established HCC markers, we investigated the gene expression of *c19orf10*, *AFP* (alpha-fetoprotein), *KRT19* (cytokeratin 19) and *GPC3* (glypican 3). Because only 1 of 15 HCC tissues analyzed above (Fig. 1d) was AFP positive (data not shown), we further investigated the expression of *c19orf10* in an additional 36 HCC tissues using RTD-PCR. Interestingly, *c19orf10* expression was significantly positively correlated with *AFP* ($r = 0.44$, $p = 0.008$), but not with *KRT19* ($r = 0.08$, $p = 0.66$) nor *GPC3* ($r = 0.11$, $p = 0.54$) (Figs. 2c-2e).

Furthermore, when we examined the expression of *c19orf10* in AFP-positive (HuH1, HuH7 and Hep3B) and -negative (HLE and HLF) HCC cell lines, we identified the overexpression of *c19orf10* in AFP-positive HCC cell lines (Fig. 2f). These data suggested that *c19orf10* is overexpressed and may play some role in AFP-positive HCCs.

***C19orf10* regulates MAPK/Akt pathways and activates cell proliferation**

To explore the functional role of *c19orf10* in HCC, we performed *c19orf10* overexpression and knockdown studies using *c19orf10*-low HLE cells and *c19orf10*-high Hep3B and HuH7 cells, respectively. When we transfected HLE cells with




Figure 3. (a) RTD-PCR analysis of *c19orf10* expression in HLE cells transfected with pcDNA3.1 or pcDNA3.1-*c19orf10* plasmids. (b) Cell proliferation assay of HLE cells transfected with pcDNA3.1 or pcDNA3.1-*c19orf10* plasmids. Cell proliferation was evaluated 72 hr after each plasmid transfection. (c) RTD-PCR analysis of *c19orf10* expression in Hep3B cells transfected with Si-Control or Si-*c19orf10*. Gene expression was measured in triplicates 48 hr after transfection. (d) Cell proliferation assay of Hep3B cells transfected with Si-Control or Si-*c19orf10*. Cell proliferation was evaluated 72 hr after siRNA transfection. (e) Cell cycle analysis of HuH7 cells transfected with Si-Control or Si-*c19orf10*. Cell cycle was evaluated 72 hr after siRNA transfection. A black arrow indicates the G2 phase peak. (f) Western blotting analysis of Huh7 cells transfected with Si-Control or Si-*c19orf10*. Cells were lysed by RIPA buffer 72 hr after siRNA transfection.

pcDNA3.1 or pcDNA3.1-*c19orf10* plasmids, we identified an approximately sixfold overexpression of *c19orf10* when compared to the control 48 hr after transfection ($p < 0.0001$) (Fig. 3a). Interestingly, cell proliferation was modestly, but

significantly, enhanced compared to the control 72 hr after transfection ($p = 0.0015$) (Fig. 3b).

We also transfected siRNAs targeting an irrelevant sequence (Si-Control) or *c19orf10* (Si-*c19orf10*) in Hep3B and




Figure 4. (a) Coomassie blue staining and Western blotting of culture supernatant of NIH3T3 cells transfected with pSI-*c19orf10*-FLAG. A black arrow indicates the 17-kDa *c19orf10* protein. (b) Cell proliferation assay of HLE cells supplemented with recombinant *c19orf10*-FLAG, FLAG peptides or BSA. Cell proliferation was measured in quadruplicates 72 hr after supplementation. (c) Western blotting of HLE cells supplemented with *c19orf10*-FLAG (40 ng/ml). Cells were lysed at indicated time after *c19orf10* supplementation. (d) Cell proliferation assay of HLE cells supplemented with control BSA (40 ng/ml) (white bar), *c19orf10*-FLAG (40 ng/ml) (light gray bar), *c19orf10*-FLAG (40 ng/ml) + anti-*c19orf10* antibodies (gray bar) and *c19orf10*-FLAG (40 ng/ml) + control mouse IgG (black bar).

HuH7 cells. We observed an ~50% decrease in *c19orf10* expression in Hep3B cells transfected with Si-*c19orf10* compared to the control 48 hr after transfection with statistical significance ($p < 0.0001$). In this condition, cell proliferation was suppressed to 50% compared to the control 72 hr after transfection ($p < 0.0001$) (Figs. 3c and 3d). When we performed cell cycle analysis of HuH7 cells transfected with Si-*Control* or Si-*c19orf10*, we identified an increase of G1-phase cells and a decrease of S- and G2-phase cells by *c19orf10* knockdown, suggesting that the G1 cycle arrest was caused by the knockdown of *c19orf10* (Fig. 3e).

We examined the representative MAPK/Akt pathway-associated proteins and cell cycle regulators using Western blotting 72 hr after siRNAs transfection (Fig. 3f). Interestingly, phosphorylation of c-Raf, MEK, MAPK, PI3K and pAkt was inhibited by knockdown of *c19orf10*, suggesting the involvement of *c19orf10* in the MAPK/Akt pathways. Furthermore, phosphorylation of Rb, CDK4 and CDK6 was also inhibited by knockdown of *c19orf10*, consistent with the

observation of G1 cell cycle arrest by *C19ORF10* knockdown. PTEN, p53 and phosphorylated CDC2 protein expression was not affected by knockdown of *c19orf10*.

***C19orf10* encodes the secretory protein and stimulates cell proliferation**

As the sequence of *c19orf10* suggested that it encodes a secretory protein, we transfected pSI-*c19orf10*-FLAG in NIH3T3 cells and examined the culture supernatant. Immunoprecipitation of the collected culture supernatant 48 hr after transfection using anti-FLAG antibodies indicated the existence of a 17-kDa protein (*c19orf10*), compatible with the molecular weight of the 142 amino acids protein encoded by *c19orf10* (Fig. 4a). We purified *c19orf10*-FLAG protein from the supernatant of HEK293 cells infected with Ad-*c19orf10*-FLAG using an anti-FLAG column. Supplementation of purified *c19orf10*-FLAG into the culture media for 72 hr enhanced the proliferation of HLE cells in a dose-dependent manner with statistical significance, whereas control FLAG peptides

and BSA had no effects on cell proliferation (Fig. 4b). Western blot analysis of HLE cells cultured with purified c19orf10-FLAG (40 ng/ml) or BSA control (40 ng/ml) indicated the immediate strong phosphorylation of Akt peaked 5 min after supplementation (Fig. 4c). The modest phosphorylation of GSK3 β (Ser9) and p44/42 MAPK also followed and peaked 60 min after c19orf10 supplementation. These data suggest that Akt pathway might be directly involved in the c19orf10-mediated cell proliferation signaling with the subsequent activation of MAPK pathway. Furthermore, addition of antibodies against c19orf10 to the culture media abolished the cell proliferation induced by c19orf10, whereas control IgG had no effects (Fig. 4d). Taken together, these data suggest that c19orf10 may be a growth factor overexpressed in AFP-positive HCCs and activates the Akt/MAPK pathways, potentially through the activation of an unidentified c19orf10 receptor.

Discussion

SAGE facilitates the measurement of transcripts from normal and malignant tissues in a nonbiased and highly accurate, quantitative manner. Indeed, SAGE produces a comprehensive gene expression profile without *a priori* gene sequence information, leading to the identification of novel transcripts potentially involved in the pathogenesis of human cancer.¹⁹ In our study, we identified seven SAGE tags potentially corresponding to novel genes activated in HCC. Among them, we identified the secretory protein c19orf10 activated in a subset of HCCs.

Several serum markers including AFP, DCP and Glypican 3 are currently used for the detection and/or the evaluation of the treatment for HCCs in the clinic.^{15–18,35} These markers are known as oncofetal proteins, that is, expressed in the fetus, transcriptionally suppressed in the adult organ and reactivated in the tumor. We identified that the expression of *c19orf10* positively correlated with *AFP* expression but did not correlate with the expression of *GPC3* or the biliary marker *KRT19*. As *c19orf10* was rarely detected in the normal liver, it is possible that c19orf10 is also an oncofetal protein activated in HCC. We are currently developing a system to detect serum c19orf10 in HCC patients, and the significance of the serum c19orf10 value as an HCC marker should be clarified.

Recent advancement in molecular biology has revealed the considerable diversity of transcription initiation and/or termination of genes altered in the process of carcinogenesis.

Indeed, using 5' SAGE approach, we recently discovered the novel intronic transcripts activated in HCC.³⁶ Interestingly, when we investigated the transcription initiation of *c19orf10* using the 5' SAGE database, we identified a potential 5' splice variant initiated from the second exon of *c19orf10* (data not shown). Although we have not yet validated the presence of 5' splice variants in *c19orf10* by PCR, examination of 5' EST database also suggested the presence of the similar splice variants (GenBank Accession Number CR980295, BQ680744, BQ648461, *etc.*). Alteration of transcription initiation/termination in *c19orf10* might affect the abundance or function of c19orf10 protein, and the details of 5' splice variants in *c19orf10* should be clarified in future studies.

Molecular targeting therapy has rapidly emerged for solid tumors as well as for leukemia.^{37–39} Sorafenib is a multikinase inhibitor targeting Raf kinase in the MAPK pathway as well as VEGFR and the platelet-derived growth factor receptor.^{40,41} In our study, we identified that c19orf10 activates the MAPK and Akt/PI3K pathways and contributes to the proliferation of HCC cell lines, although we still could not discover the potential receptor of c19orf10. Development of a neutralizing c19orf10 antibody may provide novel therapeutic options for HCC patients to inhibit these signaling pathways, and its efficacy should be evaluated in the future.

Recently, c19orf10 was found to be expressed in fibroblast-like synoviocytes in the synovium using a proteomics approach.²⁹ In addition, a recent article indicated that c19orf10 was expressed in preadipocyte cells and involved in adipogenesis using two-dimensional electrophoresis mass spectrometry analysis.²⁸ Thus, c19orf10 may have pleiotropic effects on various lineages of normal organs in various developmental stages, and the clarification of its distribution and biological properties in the whole body may provide more detailed information about the function of c19orf10.

In conclusion, we have identified the protein c19orf10 that regulates the Akt/MAPK pathways and cell cycle through an unidentified mechanism in HCC. Although further studies should be conducted to detect the potential c19orf10 receptor or signaling molecules binding to c19orf10, our study suggests that c19orf10 may be a novel growth factor, a potential tumor marker and also a potential target molecule for HCC treatment.

Acknowledgements

The authors thank Ms. Mikie Kakiuchi, Ms. Masayo Baba and Ms. Nami Nishiyama for their excellent technical assistance.

References

- Befeler AS, Di Bisceglie AM. Hepatocellular carcinoma: diagnosis and treatment. *Gastroenterology* 2002;122: 1609–19.
- Tsukuma H, Hiyama T, Tanaka S, Nakao M, Yabuuchi T, Kitamura T, Nakanishi K, Fujimoto I, Inoue A, Yamazaki H, Kawashima T. Risk factors for hepatocellular carcinoma among patients with chronic liver disease. *N Engl J Med* 1993;328:1797–801.
- Liang TJ, Jeffers LJ, Reddy KR, De Medina M, Parker IT, Cheinquer H, Idrovo V, Rabassa A, Schiff ER. Viral pathogenesis of hepatocellular carcinoma in the United States. *Hepatology* 1993;18:1326–33.
- Mayans MV, Calvet X, Bruix J, Bruguera M, Costa J, Esteve J, Bosch FX, Bru C, Rodes J. Risk factors for hepatocellular carcinoma in Catalonia, Spain. *Int J Cancer* 1990;46:378–81.

5. Mohamed AE, Kew MC, Groeneveld HT. Alcohol consumption as a risk factor for hepatocellular carcinoma in urban southern African blacks. *Int J Cancer* 1992; 51:537-41.
6. Smedile A, Bugianesi E. Steatosis and hepatocellular carcinoma risk. *Eur Rev Med Pharmacol Sci* 2005;9:291-3.
7. Floreani A, Baragiotta A, Baldo V, Menegon T, Farinati F, Naccarato R. Hepatic and extrahepatic malignancies in primary biliary cirrhosis. *Hepatology* 1999; 29:1425-8.
8. Tradati F, Colombo M, Mannucci PM, Rumi MG, De Fazio C, Gamba G, Ciavarella N, Rocino A, Morfini M, Scaraggi A, Taioli E. A prospective multicenter study of hepatocellular carcinoma in Italian hemophiliacs with chronic hepatitis C. The Study Group of the Association of Italian Hemophilia Centers. *Blood* 1998;91:1173-7.
9. Jones DE, Metcalf JV, Collier JD, Bassendine MF, James OF. Hepatocellular carcinoma in primary biliary cirrhosis and its impact on outcomes. *Hepatology* 1997; 26:1138-42.
10. Caballeria L, Pares A, Castells A, Gines A, Bru C, Rodes J. Hepatocellular carcinoma in primary biliary cirrhosis: similar incidence to that in hepatitis C virus-related cirrhosis. *Am J Gastroenterol* 2001; 96:1160-3.
11. Yoshida H, Shiratori Y, Moriyama M, Arakawa Y, Ide T, Sata M, Inoue O, Yano M, Tanaka M, Fujiyama S, Nishiguchi S, Kuroki T, et al. Interferon therapy reduces the risk for hepatocellular carcinoma: national surveillance program of cirrhotic and noncirrhotic patients with chronic hepatitis C in Japan. IHIT Study Group. Inhibition of Hepatocarcinogenesis by Interferon Therapy. *Ann Intern Med* 1999; 131:174-81.
12. Yuen MF, Cheng CC, Laufer IJ, Lam SK, Ooi CG, Lai CL. Early detection of hepatocellular carcinoma increases the chance of treatment: Hong Kong experience. *Hepatology* 2000;31:330-5.
13. Peterson MS, Baron RL. Radiologic diagnosis of hepatocellular carcinoma. *Clin Liver Dis* 2001;5:123-44.
14. Choi BI. The current status of imaging diagnosis of hepatocellular carcinoma. *Liver Transpl* 2004;10:S20-S25.
15. Fujiyama S, Tanaka M, Maeda S, Ashihara H, Hirata R, Tomita K. Tumor markers in early diagnosis, follow-up and management of patients with hepatocellular carcinoma. *Oncology* 2002;62 (Suppl 1):57-63.
16. Tsai SL, Huang GT, Yang PM, Sheu JC, Sung JL, Chen DS. Plasma des-gamma-carboxyprothrombin in the early stage of hepatocellular carcinoma. *Hepatology* 1990; 11:481-8.
17. Ikoma J, Kaito M, Ishihara T, Nakagawa N, Kamei A, Fujita N, Iwasa M, Tamaki S, Watanabe S, Adachi Y. Early diagnosis of hepatocellular carcinoma using a sensitive assay for serum des-gamma-carboxy prothrombin: a prospective study. *Hepatogastroenterology* 2002; 49:235-8.
18. Kasahara A, Hayashi N, Fusamoto H, Kawada Y, Imai Y, Yamamoto H, Hayashi E, Ogihara T, Kamada T. Clinical evaluation of plasma des-gamma-carboxy prothrombin as a marker protein of hepatocellular carcinoma in patients with tumors of various sizes. *Dig Dis Sci* 1993; 38:2170-6.
19. Yamashita T, Honda M, Kaneko S. Application of serial analysis of gene expression in cancer research. *Curr Pharm Biotechnol* 2008;9:375-82.
20. Cheng AL, Kang YK, Chen Z, Tsao CJ, Qin S, Kim JS, Luo R, Feng J, Ye S, Yang TS, Xu J, Sun Y, et al. Efficacy and safety of sorafenib in patients in the Asia-Pacific region with advanced hepatocellular carcinoma: a phase III randomised, double-blind, placebo-controlled trial. *Lancet Oncol* 2009;10:25-34.
21. Llovet JM, Ricci S, Mazzaferro V, Hilgard P, Gane E, Blanc JF, de Oliveira AC, Santoro A, Raoul JL, Forner A, Schwartz M, Porta C, et al. Sorafenib in advanced hepatocellular carcinoma. *N Engl J Med* 2008;359:378-90.
22. Yamashita T, Hashimoto S, Kaneko S, Nagai S, Toyoda N, Suzuki T, Kobayashi K, Matsushima K. Comprehensive gene expression profile of a normal human liver. *Biochem Biophys Res Commun* 2000;269: 110-16.
23. Yamashita T, Honda M, Takatori H, Nishino R, Hoshino N, Kaneko S. Genome-wide transcriptome mapping analysis identifies organ-specific gene expression patterns along human chromosomes. *Genomics* 2004;84:867-75.
24. Yamashita T, Honda M, Takatori H, Nishino R, Minato H, Takamura H, Ohta T, Kaneko S. Activation of lipogenic pathway correlates with cell proliferation and poor prognosis in hepatocellular carcinoma. *J Hepatol* 2009;50:100-10.
25. Yamashita T, Kaneko S, Hashimoto S, Sato T, Nagai S, Toyoda N, Suzuki T, Kobayashi K, Matsushima K. Serial analysis of gene expression in chronic hepatitis C and hepatocellular carcinoma. *Biochem Biophys Res Commun* 2001;282:647-54.
26. Tulin EE, Onoda N, Nakata Y, Maeda M, Hasegawa M, Nomura H, Kitamura T. SF20/IL-25, a novel bone marrow stroma-derived growth factor that binds to mouse thymic shared antigen-1 and supports lymphoid cell proliferation. *J Immunol* 2001;167:6338-47.
27. Tulin EE, Onoda N, Nakata Y, Maeda M, Hasegawa M, Nomura H, Kitamura T. SF20/IL-25, a novel bone marrow stroma-derived growth factor that binds to mouse thymic shared antigen-1 and supports lymphoid cell proliferation. *J Immunol* 2003;170:1593.
28. Wang P, Mariman E, Keijer J, Bouwman F, Noben JP, Robben J, Renes J. Profiling of the secreted proteins during 3T3-L1 adipocyte differentiation leads to the identification of novel adipokines. *Cell Mol Life Sci* 2004;61:2405-17.
29. Weiler T, Du Q, Krokhin O, Ens W, Standing K, El-Gabalawy H, Wilkins JA. The identification and characterization of a novel protein, c19orf10, in the synovium. *Arthritis Res Ther* 2007;9:R30.
30. Takatori H, Yamashita T, Honda M, Nishino R, Arai K, Takamura H, Ohta T, Zen Y, Kaneko S. dUTP pyrophosphatase expression correlates with a poor prognosis in hepatocellular carcinoma. *Liver Int* 2010; 30:438-46.
31. Sakai Y, Honda M, Fujinaga H, Tatsumi I, Mizukoshi E, Nakamoto Y, Kaneko S. Common transcriptional signature of tumor-infiltrating mononuclear inflammatory cells and peripheral blood mononuclear cells in hepatocellular carcinoma patients. *Cancer Res* 2008;68: 10267-79.
32. Honda M, Yamashita T, Ueda T, Takatori H, Nishino R, Kaneko S. Different signaling pathways in the livers of patients with chronic hepatitis B or chronic hepatitis C. *Hepatology* 2006;44:1122-38.
33. Sakai Y, Morrison BJ, Burke JD, Park JM, Terabe M, Janik JE, Forni G, Berzofsky JA, Morris JC. Vaccination by genetically modified dendritic cells expressing a truncated neu oncogene prevents development of breast cancer in transgenic mice. *Cancer Res* 2004;64:8022-8.
34. Sakai Y, Kaneko S, Nakamoto Y, Kagaya T, Mukaida N, Kobayashi K. Enhanced anti-tumor effects of herpes simplex virus thymidine kinase/ganciclovir system by codelivering monocyte chemoattractant protein-1 in hepatocellular carcinoma. *Cancer Gene Ther* 2001;8:695-704.
35. Capurro M, Wanless IR, Sherman M, Deboer G, Shi W, Miyoshi E, Filmus J. Glypican-3: a novel serum and histochemical marker for hepatocellular carcinoma. *Gastroenterology* 2003;125: 89-97.
36. Hodo Y, Hashimoto S, Honda M, Yamashita T, Suzuki Y, Sugano S, Kaneko S, Matsushima K. Comprehensive gene expression analysis of 5'-end of mRNA identified novel intronic transcripts associated with hepatocellular carcinoma. *Genomics* 2010;95:217-23.

37. Romond EH, Perez EA, Bryant J, Suman VJ, Geyer CE, Jr, Davidson NE, Tan-Chiu E, Martino S, Paik S, Kaufman PA, Swain SM, Pisansky TM, et al. Trastuzumab plus adjuvant chemotherapy for operable HER2-positive breast cancer. *N Engl J Med* 2005;353:1673-84.
38. Hurwitz H, Fehrenbacher L, Novotny W, Cartwright T, Hainsworth J, Heim W, Berlin J, Baron A, Griffing S, Holmgren E, Ferrara N, Fyfe G, et al. Bevacizumab plus irinotecan, fluorouracil, and leucovorin for metastatic colorectal cancer. *N Engl J Med* 2004;350:2335-42.
39. Shepherd FA, Rodrigues Pereira J, Ciuleanu T, Tan EH, Hirsh V, Thongprasert S, Campos D, Maoleekoonpiroj S, Smylie M, Martins R, van Kooten M, Dediu M, et al. Erlotinib in previously treated non-small-cell lung cancer. *N Engl J Med* 2005; 353:123-32.
40. Llovet JM, Bruix J. Molecular targeted therapies in hepatocellular carcinoma. *Hepatology* 2008;48:1312-27.
41. Liu L, Cao Y, Chen C, Zhang X, McNabola A, Wilkie D, Wilhelm S, Lynch M, Carter C. Sorafenib blocks the RAF/MEK/ERK pathway, inhibits tumor angiogenesis, and induces tumor cell apoptosis in hepatocellular carcinoma model PLC/PRF/5. *Cancer Res* 2006;66: 11851-8.

Prolonged recurrence-free survival following OK432-stimulated dendritic cell transfer into hepatocellular carcinoma during transarterial embolization

Y. Nakamoto,* E. Mizukoshi,*
M. Kitahara,* F. Arihara,* Y. Sakai,*
K. Kakinoki,* Y. Fujita,*
Y. Marukawa,* K. Arai,*
T. Yamashita,* N. Mukaida,[†]
K. Matsushima,* O. Matsui[§] and
S. Kaneko*

*Disease Control and Homeostasis, Graduate School of Medicine, [†]Division of Molecular Bioregulation, Cancer Research Institute, Kanazawa University, [§]Department of Radiology, Graduate School of Medicine, Kanazawa University, Kanazawa, and [‡]Department of Molecular Preventive Medicine, Graduate School of Medicine, University of Tokyo, Tokyo, Japan

Accepted for publication 19 July 2010

Correspondences: S. Kaneko, Disease Control and Homeostasis, Graduate School of Medical Science, Kanazawa University, 13-1 Takara-machi, Kanazawa 920-8641, Japan.
E-mail: skaneko@m-kanazawa.jp

Introduction

Many locoregional therapeutic approaches including surgical resection, radiofrequency ablation (RFA) and transcatheter hepatic arterial embolization (TAE) have been taken in the search for curative treatments of hepatocellular carcinoma (HCC). Despite these efforts, tumour recurrence rates remain high [1,2], probably because active hepatitis and cirrhosis in the surrounding non-tumour liver tissues causes *de novo* development of HCC [3,4]. One strategy to reduce tumour recurrence is to enhance anti-tumour immune responses that may induce sufficient inhibitory effects to prevent tumour cell growth and survival [5,6]. Dendritic

Summary

Despite curative locoregional treatments for hepatocellular carcinoma (HCC), tumour recurrence rates remain high. The current study was designed to assess the safety and bioactivity of infusion of dendritic cells (DCs) stimulated with OK432, a streptococcus-derived anti-cancer immunotherapeutic agent, into tumour tissues following transcatheter hepatic arterial embolization (TAE) treatment in patients with HCC. DCs were derived from peripheral blood monocytes of patients with hepatitis C virus-related cirrhosis and HCC in the presence of interleukin (IL)-4 and granulocyte-macrophage colony-stimulating factor and stimulated with 0.1 KE/ml OK432 for 2 days. Thirteen patients were administered with 5×10^6 of DCs through arterial catheter during the procedures of TAE treatment on day 7. The immunomodulatory effects and clinical responses were evaluated in comparison with a group of 22 historical controls treated with TAE but without DC transfer. OK432 stimulation of immature DCs promoted their maturation towards cells with activated phenotypes, high expression of a homing receptor, fairly well-preserved phagocytic capacity, greatly enhanced cytokine production and effective tumoricidal activity. Administration of OK432-stimulated DCs to patients was found to be feasible and safe. Kaplan–Meier analysis revealed prolonged recurrence-free survival of patients treated in this manner compared with the historical controls ($P = 0.046$, log-rank test). The bioactivity of the transferred DCs was reflected in higher serum concentrations of the cytokines IL-9, IL-15 and tumour necrosis factor- α and the chemokines CCL4 and CCL11. Collectively, this study suggests that a DC-based, active immunotherapeutic strategy in combination with locoregional treatments exerts beneficial anti-tumour effects against liver cancer.

Keywords: dendritic cells, hepatocellular carcinoma, immunotherapy, recurrence-free survival, transcatheter hepatic arterial embolization

cells (DCs) are the most potent type of antigen-presenting cells in the human body, and are involved in the regulation of both innate and adaptive immune responses [7]. DC-based immunotherapies are believed to contribute to the eradication of residual and recurrent tumour cells.

To enhance tumour antigen presentation to T lymphocytes, DCs have been transferred with major histocompatibility complex (MHC) class I and class II genes [8] and co-stimulatory molecules, e.g. CD40, CD80 and CD86 [9,10], and loaded with tumour-associated antigens, including tumour lysates, peptides and RNA transfection [11]. To induce natural killer (NK) and natural killer T (NK T) cell activation, DCs have been stimulated and modified to

Table 1. Patient characteristics.

Patient no.	Gender	Age (years)	HLA	TNM stages	No. of tumours	Largest tumour (mm)	Child–Pugh	KPS	Post-TAE Rx
1	M	60	A11 A33	III	5	35	B	100	RFA
2	M	57	A11 A24	III	1	21	B	100	RFA
3	M	57	A11 A31	III	2	39	B	100	RFA
4	M	77	A2 A24	III	2	35	A	100	RFA
5	F	83	A11 A24	III	3	29	B	100	RFA
6	F	74	A2 A24	II	1	35	A	100	RFA
7	F	72	A24 A33	III	3	41	B	100	RFA
8	F	65	A2 A11	II	4	12	B	100	RFA
9	M	71	A2 A11	II	4	16	A	100	RFA
10	M	79	A11 A24	III	2	40	A	100	RFA
11	M	71	A2 A24	II	1	28	A	100	RFA
12	M	56	A2 A26	III	2	25	B	100	RFA
13	M	64	A2 A33	III	2	37	B	100	RFA

M, male; F, female; TNM, tumour–node–metastasis; Child–Pugh, Child–Pugh classification; KPS, Karnofsky performance scores; TAE, transcatheter arterial embolization; Rx, treatment; HCC, hepatocellular carcinoma; HLA, human leucocyte antigen; RFA, percutaneous radiofrequency ablation.

produce larger amounts of cytokines, e.g. interleukin (IL)-12, IL-18 and type I interferons (IFNs) [10,12]. Furthermore, DC migration into secondary lymphoid organs could be induced by expression of chemokine genes, e.g. C-C chemokine receptor-7 (CCR7) [13], and by maturation using inflammatory cytokines [14], matrix metalloproteinases and Toll-like receptor (TLR) ligands [15].

DCs stimulated with OK432, a penicillin-inactivated and lyophilized preparation of *Streptococcus pyogenes*, were suggested recently to produce large amounts of T helper type 1 (Th1) cytokines, including IL-12 and IFN- γ and enhance cytotoxic T lymphocyte activity compared to a standard mixture of cytokines [tumour necrosis factor- α (TNF- α), IL-1 β , IL-6 and prostaglandin E₂ (PGE₂)] [16]. Furthermore, because OK432 modulates DC maturation through TLR-4 and the β_2 integrin system [16,17] and TLR-4-stimulated DCs can abrogate the activity of regulatory T cells [18], OK432-stimulated DCs may contribute to the induction of anti-tumour immune responses partly by reducing the activity of suppressor cells. Recently, in addition to the orchestration of immune responses, OK432-activated DCs have themselves been shown to mediate strong, specific cytotoxicity towards tumour cells via CD40/CD40 ligand interactions [19].

We have reported recently that combination therapy using TAE together with immature DC infusion is safe for patients with cirrhosis and HCC [20]. DCs were infused precisely into tumour tissues and contributed to the recruitment and activation of immune cells *in situ*. However, this approach by itself yielded limited anti-tumour effects due probably to insufficient stimulation of immature DCs (the preparation of which seems closely related to therapeutic outcome [21,22]). The current study was designed to assess the safety and bioactivity of OK432-stimulated DC infusion into tumour tissues following TAE treatment in patients with cirrhosis and HCC. In addition to documenting the safety of

this approach, we found that patients treated with OK432-stimulated DCs displayed unique cytokine and chemokine profiles and, most importantly, experienced prolonged recurrence-free survival.

Patients and methods

Patients

Inclusion criteria were a radiological diagnosis of primary HCC by computed tomography (CT) angiography, hepatitis C virus (HCV)-related HCC, a Karnofsky score of $\geq 70\%$, an age of ≥ 20 years, informed consent and the following normal baseline haematological parameters (within 1 week before DC administration): haemoglobin ≥ 8.5 g/dl; white cell count $\geq 2000/\mu\text{l}$; platelet count $\geq 50\,000/\mu\text{l}$; creatinine < 1.5 mg/dl and liver damage A or B [23].

Exclusion criteria included severe cardiac, renal, pulmonary, haematological or other systemic disease associated with a discontinuation risk; human immunodeficiency virus (HIV) infection; prior history of other malignancies; history of surgery, chemotherapy or radiation therapy within 4 weeks; immunological disorders including splenectomy and radiation to the spleen; corticosteroid or anti-histamine therapy; current lactation; pregnancy; history of organ transplantation; or difficulty in follow-up.

Thirteen patients (four women and nine men) presenting at Kanazawa University Hospital between March 2004 and June 2006 were enrolled into the study, with an age range from 56 to 83 years (Table 1). Patients with verified radiological diagnoses of HCC stage II or more were eligible and enrolled in this study. In addition, a group of 22 historical controls (nine women and 13 men) treated with TAE without DC administration between July 2000 and September 2007 was included in this study. All patients received RFA therapy to increase the locoregional effects 1 week later [24].

They underwent ultrasound, computed tomography (CT) scan or magnetic resonance imaging (MRI) of the abdomen about 1 month after treatment and at a minimum of once every 3 months thereafter, and tumour recurrences were followed for up to 360 days. The Institutional Review Board reviewed and approved the study protocol. This study complied with ethical standards outlined in the Declaration of Helsinki. Adverse events were monitored for 1 month after the DC infusion in terms of fever, vomiting, abdominal pain, encephalopathy, myalgia, ascites, gastrointestinal disorder, bleeding, hepatic abscess and autoimmune diseases.

Preparation and injection of autologous DCs

DCs were generated from blood monocyte precursors, as reported previously [25]. Briefly, peripheral blood mononuclear cells (PBMCs) were isolated by centrifugation in Lymphoprep™ Tubes (Nycomed, Roskilde, Denmark). For generating DCs, PBMCs were plated in six-well tissue culture dishes (Costar, Cambridge, MA, USA) at 1.4×10^7 cells in 2 ml per well and allowed to adhere to plastic for 2 h. Adherent cells were cultured in serum-free media (GMP CellGro® DC Medium; CellGro, Manassas, VA, USA) with 50 ng/ml recombinant human IL-4 (GMP grade; CellGro®) and 100 ng/ml recombinant human granulocyte-macrophage colony-stimulating factor (GM-CSF) (GMP grade; CellGro®) for 5 days to generate immature DC, and matured for a further 2 days in 0.1 KE/ml OK432 (Chugai Pharmaceuticals, Tokyo, Japan) to generate OK-DC. On day 7, the cells were harvested for injection, 5×10^6 cells were suspended in 5 ml normal saline containing 1% autologous plasma, mixed with absorbable gelatin sponge (Gelfoam; Pharmacia & Upjohn, Peapack, NJ, USA) and infused through an arterial catheter following Lipiodol (iodized oil) (Lipiodol Ultrafluide, Laboratoire Guerbet, Aulnay-Sous-Bois, France) injection during selective TAE therapy. Release criteria for DCs were viability > 80%, purity > 30%, negative Gram stain and endotoxin polymerase chain reaction (PCR) and negative in process cultures from samples sent 48 h before release. All products met all release criteria, and the DCs had a typical phenotype of CD14⁻ and human leucocyte antigen (HLA)-DR⁺.

Flow cytometry analysis

The DC preparation was assessed by staining with the following monoclonal antibodies for 30 min on ice: anti-lineage cocktail 1 (lin-1; CD3, CD14, CD16, CD19, CD20 and CD56)-fluorescein isothiocyanate (FITC), anti-HLA-DR-peridinin chlorophyll protein (PerCP) (L243), anti-CCR7-phycoerythrin (PE) (3D12) (BD Pharmingen, San Diego, CA, USA), anti-CD80-PE (MAB104), anti-CD83-PE (HB15a) and anti-CD86-PE (HA5.2B7) (Beckman Coulter, Fullerton, CA, USA). Cells were analysed on a fluorescence activated cell sorter (FACSocalibur™ flow cytometer. Data

analysis was performed with CELLQuest™ software (Becton Dickinson, San Jose, CA, USA).

DC phagocytosis

Immature DCs and OK432-stimulated DCs were incubated with 1 mg/ml FITC dextran (Sigma-Aldrich, St Louis, MO, USA) for 30 min at 37°C and the cells were washed three times in FACS buffer before cell acquisition using a FACS-Calibur™ cytometer. Control DCs (not incubated with FITC dextran) were acquired at the same time to allow background levels of fluorescence to be determined.

Enzyme-linked immunosorbent assay (ELISA)

DCs were seeded at 200 000 cells/ml, and supernatant collected after 48 h. IL-12p40 and IFN- γ were detected using matched paired antibodies (BD Pharmingen) following standard protocols.

Cytotoxicity assays

The ability of DCs to exert cytotoxicity was assessed in a standard ⁵¹Cr release assay [19]. We used the HCC cell lines Hep3B and PLC/PRF/5 [American Type Culture Collection (ATCC), Manassas, VA, USA] and a lymphoblastoid cell line T2 that expresses HLA-A*0201 (ATCC) as target cells. Target cells were labelled with ⁵¹Cr. In a 96-well plate, 2.5×10^3 target cells per well were incubated with DCs for 8 h at different effector/target (E/T) ratios in triplicate. Percentage of specific lysis was calculated as follows: (experimental release – spontaneous release)/(maximum release – spontaneous release) \times 100. Spontaneous release was always < 20% of the total.

NK cell activity

NK cell cytotoxicity against K562 erythroleukemia target cells was measured by using ⁵¹Cr-release assay, according to previously published methods [26], with PBMCs obtained from the patients. All experiments were performed in triplicate. Percentage of cytotoxicity was calculated as follows: {[experimental counts per minute (cpm) – spontaneous cpm]/[total cpm – spontaneous cpm]} \times 100.

Intracellular cytokine expression

Freshly isolated PBMCs were stimulated with 25 ng/ml phorbol 12-myristate 13-acetate (PMA; Sigma-Aldrich) and 1 μ g/ml ionomycin (Sigma-Aldrich) at 37°C in humidified 7% CO₂ for 4 h. To block cytokine secretion, brefeldin A (Sigma) [27] was added to a final concentration of 10 μ g/ml. After addition of stimuli, the surface staining was performed with anti-CD4-PC5 (13B8.2), anti-CD8-PerCP (SK1) and anti-CD56-PC5 (N901) (Beckman

Coulter). Subsequently, the cells were permeabilized, stained for intracellular IFN- γ and IL-4 using the FastImmune™ system (BD Pharmingen), resuspended in phosphate-buffered saline (PBS) containing 1% paraformaldehyde (PFA), and analysed on a flow cytometer ($\approx 10\,000$ gated events acquired per sample).

IFN- γ enzyme-linked immunospot (ELISPOT) assay

ELISPOT assays were performed as described previously with the following modifications [28–30]. HLA-A24 restricted peptide epitopes, squamous cell carcinoma antigen recognized by T cells 2 (SART2)₈₉₉ (SYTRLFLIL), SART3₁₀₉ (VYDYNCHVDL), multi-drug resistance protein 3 (MRP3)₇₆₅ (VYSDADIFL), MRP3₅₀₃ (LYAWEPSFL), MRP3₆₉₂ (AYVPQQAWI), alpha-fetoprotein (AFP)₄₀₃ (KYIQESQAL), AFP₄₃₄ (AYTKKAPQL), AFP₃₅₇ (EYSRRHPQL), human telomerase reverse transcriptase (hTERT)₁₆₇ (AYQVCGPPL) (unpublished), hTERT₄₆₁ (VYGFVRACL) and hTERT₃₂₄ (VYAETKHFL) were used in this study. Negative controls consisted of an HIV envelope-derived peptide (HIVenv₅₈₄). Positive controls consisted of 10 ng/ml PMA (Sigma) or a CMV pp65-derived peptide (CMVpp65₃₂₈). The coloured spots were counted with a KS ELISPOT Reader (Zeiss, Tokyo, Japan). The number of specific spots was determined by subtracting the number of spots in the absence of antigen from the number of spots in its presence. Responses were considered positive if more than 10 specific spots were detected and if the number of spots in the presence of antigen was at least twofold greater than the number of spots in the absence of antigen.

Cytokine and chemokine profiling

Serum cytokine and chemokine levels were measured using the Bioplex assay (Bio-Rad, Hercules, CA, USA). Briefly, frozen serum samples were thawed at room temperature, diluted 1:4 in sample diluents, and 50 μ l aliquots of diluted sample were added in duplicate to the wells of a 96-well microtitre plate containing the coated beads for a validated panel of 27 human cytokines and chemokines (cytokine 27-plex antibody bead kit) according to the manufacturer's instructions. These included IL-1 β , IL-1Ra, IL-2, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-12p70, IL-13, IL-15, IL-17, basic fibroblast growth factor (FGF), eotaxin, G-CSF, GM-CSF, IFN- γ , interferon gamma-induced protein (IP)-10, monocyte chemoattractant protein (MCP)-1, MIP-1 α , MIP-1 β , platelet-derived growth factor (PDGF)-BB, regulated upon activation normal T cell-expressed and secreted (RANTES), TNF- α and vascular endothelial growth factor (VEGF). Eight standards (ranging from 2 to 32 000 pg/ml) were used to generate calibration curves for each cytokine. Data acquisition and analysis were performed using Bio-Plex Manager software version 4.1.1.

Arginase activity

Serum samples were tested for arginase activity by conversion of L-arginine to L-ornithine [31] using a kit supplied by the manufacturer (BioAssay Systems, Hayward, CA, USA). Briefly, sera were treated with a membrane filter (Millipore, Billerica, MA, USA) to remove urea, combined with the sample buffer in wells of a 96-well plate, and incubated at 37°C for 2 h. Subsequently, the urea reagent was added to stop the arginase reaction. The colour produced was read at 520 nm using a microtitre plate reader.

Statistical analysis

Results are expressed as means \pm standard deviation (s.d.). Differences between groups were analysed for statistical significance by the Mann–Whitney *U*-test. Qualitative variables were compared by means of Fisher's exact test. The estimated probability of tumour recurrence-free survival was determined using the Kaplan–Meier method. The Mantel–Cox log-rank test was used to compare curves between groups. Any *P*-values less than 0.05 were considered statistically significant. All statistical tests were two-sided.


Results

Preparation of OK432-stimulated DCs

Adherent cells isolated from PBMCs of patients with cirrhosis and HCC (Table 1) were differentiated into DCs in the presence of IL-4 and GM-CSF. The cells were stimulated with 0.1 KE/ml OK432 for 3 days; 54.6 \pm 9.5% (mean \pm s.d.; *n* = 13) of OK432-stimulated cells showed high levels of MHC class II (HLA-DR) and the absence of lineage markers including CD3, CD14, CD16, CD19, CD20 and CD56, in which 30.9 \pm 14.2% were CD11c-positive (myeloid DC subset) and 14.8 \pm 11.2 were CD123-positive (plasmacytoid DC subset), consistent with our previous observations [20]. As reported [32,33], greater proportions of the cells developed high levels of expression of the co-stimulatory molecules B7-1 (CD80) and B7-2 (CD86) and an activation marker (CD83) compared to DCs prepared without OK432 stimulation (Fig. 1a). Furthermore, the chemokine receptor CCR7 which leads to homing to lymph nodes [13,34] was also induced following OK432 stimulation.

To evaluate the endocytic and phagocytic ability of the OK432-stimulated cells, uptake of FITC-dextran was quantitated by flow cytometry (Fig. 1b). The cells showed lower levels of uptake due to maturation compared to DCs prepared without OK432 stimulation, while the OK432-stimulated cells derived from HCC patients preserved a moderate uptake capacity. As expected, the OK432-stimulated cells produced large amounts of cytokines IL-12 and IFN- γ (Fig. 1c). In addition, they displayed high cyto-

Fig. 1. Effects of OK432 stimulation on the properties of dendritic cells (DCs) generated from blood monocyte precursors in patients with cirrhosis and hepatocellular carcinoma (HCC) ($n = 13$). (a) Lineage cocktail 1 (lin^{-}) human leucocyte antigen D-related (HLA-DR $^{-}$) subsets with [OK432(+)] and without [OK432(-)] stimulation were analysed for surface expression of CD80, CD83, CD86 and CCR7. Dot plots of a representative case are shown in the left-hand panel. Mean percentages [\pm standard deviation (s.d.)] of positive cells are indicated in the right-hand panel. OK432 stimulation resulted in the expression of high levels of CD80, CD83, CD86 and CCR7 in the lin^{-} human leucocyte antigen D-related (HLA-DR $^{-}$) DC subset. (b) DC subsets with and without OK432 stimulation were incubated with fluorescein isothiocyanate (FITC) dextran for 30 min and the uptake was determined by flow cytometry. A representative analysis is shown in the upper panel. Mean fluorescence intensities (MFIs) (\pm s.d.) of the positive cells are indicated in the lower panel. OK432-stimulated cells showed lower levels of uptake due to maturation. (c) DC supernatants were harvested and the concentrations of interleukin (IL)-12 and interferon (IFN)- γ measured by enzyme-linked immunosorbent assay (ELISA). OK432-stimulated cells produced large amounts of the cytokines. The data indicate means \pm s.d. of the groups with and without the stimulation. All comparisons in (a-c) [OK432(+)] versus [OK432(-)] were statistically significant by the Mann-Whitney U -test ($P < 0.005$). (d) Tumoricidal activity of DCs assessed by incubation with ^{51}Cr -labelled Hep3B, PLC/PRF/5 and T2 targets for 8 h at the indicated effector/target (E/T) cell ratios. OK432-stimulated cells displayed high cytotoxic activity against the target cells. The results are representative of the cases studied.



toxic activity against HCC cell lines (Hep3B and PLC/PRF/5) and a lymphoblastoid cell line (T2) although DCs without OK432 stimulation lysed none of the target cells to any great degree (Fig. 1d). Taken together, these results demonstrate that OK432 stimulation of IL-4 and GM-CSF-induced immature DCs derived from HCC patients promoted their maturation towards cells with activated phenotypes, high expression of a homing receptor, fairly well-preserved phagocytic capacity, greatly enhanced cytokine production and effective tumoricidal activity, consistent with previous observations [16,19].

Safety of OK432-stimulated DC administration

Prior to the administration of OK432-stimulated DCs to patients, the cells were confirmed to be safe in athymic nude mice to which 100-fold cell numbers/weight were injected subcutaneously (data not shown). Subsequently, OK432-stimulated DC administration was performed during TAE therapy in humans, in which DCs were mixed together with absorbable gelatin sponge (Gelfoam) and infused through an arterial catheter following iodized oil (Lipiodol) injection, as reported previously [20]. Adverse events were




Fig. 1. *Continued*

monitored clinically and biochemically after DC infusion (Table 2). A larger proportion (12 of 13) of the patients were complicated with high fever compared to those treated previously with immature DCs (five of 10) [20], due probably to the proinflammatory responses induced by OK432-stimulated DCs. However, there were no grades III or IV

National Cancer Institute Common Toxicity Criteria adverse events, including vomiting, abdominal pain, encephalopathy, myalgia, ascites, gastrointestinal disorders, bleeding, hepatic abscess or autoimmune diseases associated with DC infusion and TAE in this study. There was also no clinical or serological evidence of hepatic failure or autoimmune

Table 2. Adverse events.

Patient no.	Fever (days)	Vomiting	Abdominal pain	Encephalopathy	Others [†]
1	2	No	No	No	No
2	2	No	No	No	No
3	1	No	No	No	No
4	3	No	No	No	No
5	3	No	No	No	No
6	4	No	No	No	No
7	10	No	No	No	No
8	No	No	No	No	No
9	2	No	No	No	No
10	1	No	No	No	No
11	2	No	No	No	No
12	2	No	No	No	No
13	1	No	No	No	No

[†]Other adverse events include myalgia, ascites, gastrointestinal disorder, bleeding, hepatic abscess and autoimmune diseases.

response in any patients. Thus, concurrent treatment with OK432-stimulated DC infusions can be performed safely at the same time as TAE in patients with cirrhosis and HCC.

Recurrence-free survival following DC infusion

A further objective of this study was to determine clinical response following DC infusion. A group of historical controls treated with TAE without DC administration was reviewed for this study (Table 3). The clinical characteristics including tumour burden and hepatic reserve were comparable between patients treated with TAE and OK432-stimulated DC transfer ($n = 13$) and those historical controls with TAE but without DC administration ($n = 22$). We com-

pared the recurrence-free survival between these patient groups. Kaplan–Meier analysis indicated that patients treated with TAE and OK432-stimulated DC transfer had prolonged recurrence-free survival compared with the historical controls that had been treated with TAE alone (recurrence rates 360 days after the treatments; two of 13 and 12 of 22, respectively; $P = 0.046$, log-rank test) (Fig. 2). The results demonstrated that OK432-stimulated DC transfer during TAE therapy reduces tumour recurrence in HCC patients.

NK cell activity and intracellular cytokine responses in PBMCs

To assess systemic immunomodulatory effects of OK432-stimulated DC transfer, PBMCs were isolated 1 and 3 months after treatment and NK cell cytotoxicity against K562 erythroleukaemia target cells measured using the ⁵¹Cr-release assay (Fig. 3). The level of NK cell was unaltered following treatment. In addition, cytokine production capacity of lymphocyte subsets was quantitated by measuring intracellular IFN- γ and IL-4 using flow cytometry. There were also no significant changes in terms of cytokine production capacity in the CD4⁺, CD8⁺ and CD56⁺ subsets in the patients treated with OK432-stimulated DCs.

Immune responses to peptide epitopes derived from tumour antigens

To assess the effects on T cell responses to tumour antigens, PBMCs were obtained 4 weeks after DC infusion, pulsed with peptides derived from AFP, MRP3, SART2, SART3 and hTERT. IFN- γ production was then quantitated in an

Table 3. Clinical characteristics of patients treated with TAE + OK-DC and TAE alone.

	TAE + OK-DC	TAE	P
No. of patients	13	22	
Age (years)	68.2 \pm 9.1	70.0 \pm 7.6	n.s. [†]
Gender (M/F)	9/4	13/9	n.s. [‡]
White cell count ($\times 10^2/\mu\text{l}$)	34.4 \pm 11.6	41.4 \pm 18.9	n.s. [†]
Lymphocytes ($\times 10^2/\mu\text{l}$)	10.4 \pm 3.6	12.4 \pm 4.7	n.s. [†]
Platelets ($\times 10^4/\mu\text{l}$)	11.5 \pm 10.2	10.3 \pm 5.8	n.s. [†]
Hepaplastin test (%)	64.6 \pm 11.6	75.5 \pm 24.3	n.s. [†]
ALT (IU/l)	56.7 \pm 38.9	67.9 \pm 44.6	n.s. [†]
Total bilirubin (mg/dl)	1.3 \pm 0.7	1.1 \pm 0.6	n.s. [†]
Albumin (g/dl)	3.4 \pm 0.6	3.6 \pm 0.4	n.s. [†]
Non-cancerous liver parenchyma (no.)			
Chronic hepatitis	0	8	
Cirrhosis (Child–Pugh A/B/C)	13 (5/8/0)	14 (6/8/0)	n.s. [‡]
TNM stages (I/II/III/IV-A/IV-B)	0/4/9/0/0	3/8/11/0/0	n.s. [‡]
No. of tumours	2.5 \pm 1.3	1.9 \pm 1.3	n.s. [†]
Largest tumour (mm)	30.2 \pm 9.4	32.6 \pm 15.2	n.s. [†]
AFP	204.8 \pm 404.1	201.8 \pm 544.2	n.s. [†]

Results are expressed as means \pm standard deviation. [†]Mann–Whitney *U*-test. [‡]Fisher's exact test. TAE, transcatheter arterial embolization; OK-DC, OK432-stimulated dendritic cells; ALT, alanine transaminase; TNM, tumour–node–metastasis; AFP, alpha-fetoprotein; Child–Pugh, Child–Pugh classification; n.s., not significant.




Fig. 2. Recurrence-free survival of patients treated with transcatheter hepatic arterial embolization (TAE) with [TAE + OK-stimulated dendritic cells (DC); $n = 13$] and without (TAE: historical controls; $n = 22$) OK432-stimulated DC administration. Time zero is the date of TAE. All patients underwent ultrasound, computed tomography (CT) scan or magnetic resonance imaging (MRI) of the abdomen about 1 month after treatment and at a minimum of once every 3 months thereafter. Kaplan–Meier analysis indicated that TAE + OK-DC treatment prolonged recurrence-free survival compared with the TAE-alone group (recurrence rates 360 days after the treatments; two of 13 and 12 of 22, respectively; $P = 0.046$, log-rank test).

ELISPOT assay. Cells producing IFN- γ in response to stimulation with HLA-A24 [the most common HLA-A antigen (58.1%) in Japanese populations [35]]-restricted peptide epitopes derived from tumour antigens MRP3 and hTERT were induced in three of six HLA-A24-positive patients (numbers 2, 6 and 11) after treatment with TAE and OK432-stimulated DCs (Fig. 4). To understand the immunological and clinical significance of the T lymphocyte responses, PBMCs obtained from the historical control patients who had been treated with TAE without DC administration were also evaluated by ELISPOT. Similarly, positive reactions were observed in four (numbers t8, t19, t20 and t22) of six HLA-A24-positive patients. These data indicate that T lymphocyte responses to HLA-A24 restricted peptide epitopes of tumour antigens were induced following the TAE therapy, but no additional responses were observed as a result of OK432-stimulated DC transfer in the current study.

Serum levels of cytokines, chemokines and arginase activity

To screen for immunobiological responses induced following OK432-stimulated DC transfer, serum levels of cytokines and chemokines were measured simultaneously using the Bio-Plex multiplex suspension array system. The results were compared with the historical control patients treated with TAE without DC administration. Interestingly, serum con-





Fig. 3. Natural killer (NK) cell activity and intracellular cytokine production in peripheral blood mononuclear cells (PBMCs) of patients treated with OK432-stimulated dendritic cells (DCs) during transcatheter hepatic arterial embolization (TAE) therapy ($n = 13$). PBMCs were isolated before and 1 and 3 months after treatment and used for the analyses. Upper panel: NK cell cytotoxicity against K562 erythroleukaemia target cells was evaluated at the effector/target (E/T) cell ratios shown. NK cell activities were not changed following treatment. Middle and lower panels: PBMCs were stimulated with phorbol 12-myristate 13-acetate (PMA) and ionomycin, stained for CD4, CD8 and CD56 expression, permeabilized and stained for intracellular interferon (IFN)- γ and interleukin (IL)-4. Percentages of cytokine-positive cells were quantitated by flow cytometry. There were no significant changes in terms of cytokine production capacity in the CD4+, CD8+ and CD56+ subsets following the treatments. The data are given as means \pm standard deviation of the groups.

Fig. 4. Immune responses to human leucocyte antigen (HLA-DR⁻)-A24-restricted peptide epitopes derived from tumour antigens in HLA-A24-positive patients treated with OK432-stimulated DCs during transcatheter hepatic arterial embolization (TAE) therapy (numbers 2, 5, 6, 7, 10 and 11) and HLA-A24-positive historical controls treated with TAE without dendritic cell (DC) transfer (numbers t8, t14, t15, t19, t20 and t22). Peripheral blood mononuclear cells (PBMCs) were obtained before (open bars) and 1 month after the infusion (solid bars), pulsed with the peptides derived from squamous cell carcinoma antigen recognized by T cells 2 (SART2), SART3, multi-drug resistance protein 3 (MRP3), alpha-fetoprotein (AFP), human telomerase reverse transcriptase (hTERT) and interferon (IFN)- γ production was quantitated by enzyme-linked immunospot (ELISPOT). Negative controls consisted of a human immunodeficiency virus (HIV) envelope-derived peptide (HIVenv₅₈₄). Positive controls consisted of 10 ng/ml phorbol 12-myristate 13-acetate (PMA) or a cytomegalovirus (CMV) pp65-derived peptide (CMVpp65₃₂₈). The number of specific spots was determined by subtracting the number of spots in the absence of antigen from the number of spots in its presence. T lymphocyte responses to the peptide epitopes were induced following TAE therapy, but no additional responses were observed after DC transfer. Numbers denote specific spots beyond the upper limit of y-axis; n.d., not determined.



centrations of IL-9, IL-15 and TNF- α were greatly increased after OK432-stimulated DC infusion, in contrast to their reduction following TAE treatment alone (Fig. 5a). Furthermore, the chemokines eotaxin (CCL11) and MIP-1 β (CCL4) were induced markedly after DC transfer, although they were also decreased after TAE alone. These data indicate that transfer of OK432-stimulated DC during TAE therapy induced unique immune responses that may be mediated by the cytokines IL-9, IL-15 and TNF- α and the chemokines eotaxin and MIP-1 β .

In addition, serum arginase activity was reported to reflect numbers of myeloid-derived suppressor cells (MDSCs) that may inhibit T lymphocyte responses in cancer patients [36]. Therefore, serum arginase activity was measured after OK432-stimulated DC infusion, and it was found that it was

increased six- or sevenfold in patients treated with TAE. However, this increase was independent of the presence or absence of OK432-stimulated DC transfer (Fig. 5b). None the less, serum arginase activity was decreased again 4 weeks after treatment with both TAE and OK432-stimulated DC transfer but tended to be maintained at a high level in patients treated with TAE without DC transfer. However, these differences did not reach statistical significance ($P > 0.05$). Because arginase activity is known to be relatively high in liver and HCC cells [37], the influence of tissue injury was assessed biochemically by measuring serum levels of ALT and LDH activities. We did not observe ALT or LDH elevation, indicating that the increase of arginase activity was not due to tissue damage following treatment. Collectively, these results demonstrate that infusion of OK432-stimulated




Fig. 5. Cytokine and chemokine profiling and arginase activity in sera of patients treated with OK432-stimulated dendritic cells (DCs) during transcatheter hepatic arterial embolization (TAE) therapy (TAE + OK-DC; $n = 13$) and the historical controls treated with TAE without DC transfer (TAE; $n = 22$). (a) Serum samples were examined for their content of a validated panel of cytokines and chemokines using the Bioplex assay. Percentage changes in serum levels 2 weeks after the treatments were calculated as follows: [(post-treatment level – pretreatment level)/pretreatment level] $\times 100$. The data are means \pm standard error of the mean (s.e.m.) of the groups. * $P < 0.05$ when compared by the Mann–Whitney U -test. (b) Serum samples were tested for arginase activity by conversion of L-arginine to L-ornithine, and for alanine aminotransferase (ALT) and lactic dehydrogenase (LDH) activities. While there was a trend for the arginase activity in the TAE + OK-DC group to decrease 4 weeks after treatment, the difference did not reach statistical significance ($P > 0.05$). Percentage changes in serum levels 2 weeks after the treatments were calculated as follows: [(post-treatment level – pretreatment level)/pretreatment level] $\times 100$. The data indicate means \pm s.e.m. of the groups.

DCs during TAE treatment may reduce the immunosuppressive activities of MDSCs, and assist in developing a favourable environment for the induction of anti-tumour immunity.

Discussion

Although many novel strategies, including immunotherapies, have been developed in an attempt to suppress tumour recurrence after curative treatments for HCC, recurrence rates and survival times have not been improved significantly [38]. In the current study, we first established that OK432-stimulated DC administration during TAE therapy did not cause critical adverse events in patients with cirrhosis and HCC. Most importantly, DC transfer resulted in prolonged recurrence-free survival after combination therapy with TAE and OK432-stimulated DC administration. In terms of the immunomodulatory effects of DC transfer, although NK cell activity, intracellular cytokine production and T lymphocyte-mediated immune responses were not altered in PBMCs from treated patients, serum levels of IL-9, IL-15 and TNF- α and the chemokines eotaxin and MIP-1 β were enhanced markedly after DC transfer. In addition, serum levels of arginase activity were decreased following DC transfer. Collectively, this study demonstrated the feasibility, safety and beneficial anti-tumour effects of OK432-stimulated DC infusion into tumour tissues for patients with cirrhosis and HCC, suggesting the ability of an active immunotherapeutic strategy to reduce tumour recurrence after locoregional treatment of HCC.

DCs were stimulated with OK432 prior to infusion into tumour tissues through an arterial catheter. OK432 was reported to activate DCs through its binding to TLR-2 and -4 [16,39] that can be used for cancer therapy [33]. The current results indicate that OK432 stimulation of immature DCs

from HCC patients promoted their maturation processes while preserving antigen uptake capacity and enhancing tumoricidal activity, consistent with previous observations [16,19] and supporting the current strategy in which OK432-stimulated DCs were infused directly into tumour tissues. Because the tumoricidal activity of unstimulated DCs was not observed in *in vitro* experiments, OK432 stimulation obviously altered the cytotoxic properties of DCs. One of the mechanisms of DC killing was reported to be CD40/CD40 ligand interaction [19]. Further studies are needed to determine the killing mechanisms of DCs derived from HCC patients in a direct [TNF, TNF-related apoptosis inducing ligand (TRAIL), Fas ligand, nitric oxide (NO) and perforin/granzyme] and indirect (MHC-restricted) manner [40–43]. Although the main mechanism by which OK432-stimulated DCs prolonged the recurrence-free survival was not elucidated, the tumoricidal activity of mature DCs was implicated in *in vivo* enhancement of antigen presentation, co-stimulation and inflammatory cytokine production.

Very recent reports document injection of OK432-stimulated DCs into patients with cancer of the gastrointestinal tract or pancreas [44,45], but their anti-tumour effects were not defined clearly. The current study shows for the first time that OK432-stimulated DCs induce beneficial anti-tumour responses when transferred into tumour tissues during TAE therapy. The anti-tumour responses may have been enhanced as a result of optimal activation of the DCs with OK432 or combining infusion of stimulated DCs with TAE therapy. Inappropriately activated DCs may be unable to generate sufficient numbers of properly activated effector T lymphocytes [46]. As shown in Fig. 1, all these alterations could contribute to the further enhancement of anti-tumour effects compared to those in our previous study with immature DCs [20]. Furthermore, the tumour cell death-promoting therapies, e.g. chemotherapy [47] and TAE [48], can be expected to enhance the effects of therapeutic cancer vaccines by redressing the immunosuppressive tumour environment.

NK cell activity and intracellular cytokine responses in CD4⁺ and CD8⁺ T lymphocytes and CD56⁺ NK cell subsets in PBMCs were not changed significantly in patients treated with OK432-stimulated DCs. Furthermore, we did not observe tumour antigen-specific T lymphocyte responses associated clearly with DC administration. The data suggest therefore that the immune responses induced by the therapy applied here were not detectable systemically. Because cytotoxic T lymphocyte responses were enhanced in patients receiving $> 3 \times 10^7$ cells [49,50], the numbers of transferred OK432-stimulated DCs were apparently not sufficient to induce responses detectable in the peripheral blood, but were enough to exert beneficial anti-tumour effects. In addition, many studies have concluded that cytotoxic T lymphocyte responses rarely predict clinical outcomes of DC-based immunotherapies [51,52] and that in many cases, also including our own studies

[28,30], tumour-specific effector T lymphocytes co-exist with the tumours. Consistent with these observations, the current results suggest that cytotoxic T lymphocyte responses in PBMCs are not reliable predictors of beneficial anti-tumour effects in patients treated with the current OK432-stimulated DC strategy.

Serum levels of the cytokines IL-9, IL-15 and TNF- α and the chemokines eotaxin and MIP-1 β were increased following OK432-stimulated DC transfer, but decreased after TAE therapy without DC administration. IL-9 and IL-15 belong to the cytokine receptor common gamma chain (γ_c ; CD132) family, a member of the type I cytokine receptor family expressed on most lymphocyte populations [53]. IL-9 exerts pleiotropic activities on T and B lymphocytes, mast cells, monocytes and haematopoietic progenitors [54,55]. IL-15 and TNF- α are known to prime T lymphocytes and NK cells when secreted by DCs [56] and to induce anti-tumour immune responses [57]. Eotaxin is known to selectively recruit eosinophils also contributing to anti-tumour effects [58,59], and MIP-1 β is a chemoattractant for NK cells, monocytes and a variety of other immune cells [60]. In addition, serum levels of arginase tended to decrease after DC transfer. Because serum arginase activity reflects the numbers of MDSCs that inhibit T lymphocyte responses in cancer patients [36], the patients treated with OK432-stimulated DCs might have developed lower levels of suppressor cells. Collectively, the results suggest that infusion of OK432-stimulated DCs may orchestrate the immune environment in the whole body that could enhance beneficial anti-tumour effects, although the precise molecular and cellular mechanisms associated with the actions of these cytokines and chemokines were not defined clearly in the current analysis.

Acknowledgements

The authors thank Kazumi Fushimi and Mariko Katsuda for technical assistance. We also thank the patients for participating in this trial. This work was supported in part by research grants from the Ministry of Education, Culture, Sports, Science, and Technology of Japan, the Ministry of Health, Labour and Welfare of Japan and the Japanese Society of Gastroenterology.

Disclosure

The authors have declared that no conflict of interest exists.

References

- 1 Omata M, Tateishi R, Yoshida H, Shiina S. Treatment of hepatocellular carcinoma by percutaneous tumor ablation methods: ethanol injection therapy and radiofrequency ablation. *Gastroenterology* 2004; 127:S159–66.
- 2 Belghiti J. Resection and liver transplantation for HCC. *J Gastroenterol* 2009; 44 (Suppl. 19):132–5.

- 3 Nakamoto Y, Guidotti LG, Kuhlen CV, Fowler P, Chisari FV. Immune pathogenesis of hepatocellular carcinoma. *J Exp Med* 1998; **188**:341–50.
- 4 Ercolani G, Grazi GL, Ravaioli M *et al.* Liver resection for hepatocellular carcinoma on cirrhosis: univariate and multivariate analysis of risk factors for intrahepatic recurrence. *Ann Surg* 2003; **237**:536–43.
- 5 Shankaran V, Ikeda H, Bruce AT *et al.* IFN γ and lymphocytes prevent primary tumour development and shape tumour immunogenicity. *Nature* 2001; **410**:1107–11.
- 6 Vulink A, Radford KJ, Melief C, Hart DN. Dendritic cells in cancer immunotherapy. *Adv Cancer Res* 2008; **99**:363–407.
- 7 Banchereau J, Briere F, Caux C *et al.* Immunobiology of dendritic cells. *Annu Rev Immunol* 2000; **18**:767–811.
- 8 Lemos MP, Esquivel F, Scott P, Laufer TM. MHC class II expression restricted to CD8 α ⁺ and CD11b⁺ dendritic cells is sufficient for control of *Leishmania major*. *J Exp Med* 2004; **199**:725–30.
- 9 Ni K, O'Neill HC. The role of dendritic cells in T cell activation. *Immunol Cell Biol* 1997; **75**:223–30.
- 10 Andrews DM, Andoniou CE, Scalzo AA *et al.* Cross-talk between dendritic cells and natural killer cells in viral infection. *Mol Immunol* 2005; **42**:547–55.
- 11 Heiser A, Coleman D, Dannull J *et al.* Autologous dendritic cells transfected with prostate-specific antigen RNA stimulate CTL responses against metastatic prostate tumors. *J Clin Invest* 2002; **109**:409–17.
- 12 Banchereau J, Steinman RM. Dendritic cells and the control of immunity. *Nature* 1998; **392**:245–52.
- 13 Forster R, Schubel A, Breitfeld D *et al.* CCR7 coordinates the primary immune response by establishing functional microenvironments in secondary lymphoid organs. *Cell* 1999; **99**:23–33.
- 14 Martín-Fontecha A, Sebastiani S, Hopken UE *et al.* Regulation of dendritic cell migration to the draining lymph node: impact on T lymphocyte traffic and priming. *J Exp Med* 2003; **198**:615–21.
- 15 Ratzinger G, Stoltzner P, Ebner S *et al.* Matrix metalloproteinases 9 and 2 are necessary for the migration of Langerhans cells and dermal dendritic cells from human and murine skin. *J Immunol* 2002; **168**:4361–71.
- 16 Nakahara S, Tsunoda T, Baba T, Asabe S, Tahara H. Dendritic cells stimulated with a bacterial product, OK-432, efficiently induce cytotoxic T lymphocytes specific to tumor rejection peptide. *Cancer Res* 2003; **63**:4112–18.
- 17 Okamoto M, Oshikawa T, Tano T *et al.* Mechanism of anticancer host response induced by OK-432, a streptococcal preparation, mediated by phagocytosis and Toll-like receptor 4 signaling. *J Immunother* 2006; **29**:78–86.
- 18 Pasare C, Medzhitov R. Toll pathway-dependent blockade of CD4⁺CD25⁺ T cell-mediated suppression by dendritic cells. *Science* 2003; **299**:1033–6.
- 19 Hill KS, Errington F, Steele LP *et al.* OK432-activated human dendritic cells kill tumor cells via CD40/CD40 ligand interactions. *J Immunol* 2008; **181**:3108–15.
- 20 Nakamoto Y, Mizukoshi E, Tsuji H *et al.* Combined therapy of transcatheter hepatic arterial embolization with intratumoral dendritic cell infusion for hepatocellular carcinoma: clinical safety. *Clin Exp Immunol* 2007; **147**:296–305.
- 21 Steinman RM, Banchereau J. Taking dendritic cells into medicine. *Nature* 2007; **449**:419–26.
- 22 Tacken PJ, de Vries IJ, Torensma R, Figdor CG. Dendritic-cell immunotherapy: from *ex vivo* loading to *in vivo* targeting. *Nat Rev Immunol* 2007; **7**:790–802.
- 23 Makuuchi M. General rules for the clinical and pathological study of primary liver cancer, 2nd edn. Tokyo: Kanehara & Co., Ltd, 2003.
- 24 Veltri A, Moretto P, Doriguzzi A, Pagano E, Carrara G, Gandini G. Radiofrequency thermal ablation (RFA) after transarterial chemoembolization (TACE) as a combined therapy for unresectable non-early hepatocellular carcinoma (HCC). *Eur Radiol* 2006; **16**:661–9.
- 25 Dhodapkar MV, Steinman RM, Sapp M *et al.* Rapid generation of broad T-cell immunity in humans after a single injection of mature dendritic cells. *J Clin Invest* 1999; **104**:173–80.
- 26 Orange JS, Brodeur SR, Jain A *et al.* Deficient natural killer cell cytotoxicity in patients with IKK- γ /NEMO mutations. *J Clin Invest* 2002; **109**:1501–9.
- 27 Klausner RD, Donaldson JG, Lippincott-Schwartz J, Brefeldin A: insights into the control of membrane traffic and organelle structure. *J Cell Biol* 1992; **116**:1071–80.
- 28 Mizukoshi E, Nakamoto Y, Marukawa Y *et al.* Cytotoxic T cell responses to human telomerase reverse transcriptase in patients with hepatocellular carcinoma. *Hepatology* 2006; **43**:1284–94.
- 29 Mizukoshi E, Nakamoto Y, Tsuji H, Yamashita T, Kaneko S. Identification of alpha-fetoprotein-derived peptides recognized by cytotoxic T lymphocytes in HLA-A24⁺ patients with hepatocellular carcinoma. *Int J Cancer* 2006; **118**:1194–204.
- 30 Mizukoshi E, Honda M, Arai K, Yamashita T, Nakamoto Y, Kaneko S. Expression of multidrug resistance-associated protein 3 and cytotoxic T cell responses in patients with hepatocellular carcinoma. *J Hepatol* 2008; **49**:946–54.
- 31 Rodríguez PC, Quiceno DG, Zabaleta J *et al.* Arginase I production in the tumor microenvironment by mature myeloid cells inhibits T-cell receptor expression and antigen-specific T-cell responses. *Cancer Res* 2004; **64**:5839–49.
- 32 Itoh T, Ueda Y, Okugawa K *et al.* Streptococcal preparation OK432 promotes functional maturation of human monocyte-derived dendritic cells. *Cancer Immunol Immunother* 2003; **52**:207–14.
- 33 Kuroki H, Morisaki T, Matsumoto K *et al.* Streptococcal preparation OK-432: a new maturation factor of monocyte-derived dendritic cells for clinical use. *Cancer Immunol Immunother* 2003; **52**:561–8.
- 34 Gunn MD, Kyuwa S, Tam C *et al.* Mice lacking expression of secondary lymphoid organ chemokine have defects in lymphocyte homing and dendritic cell localization. *J Exp Med* 1999; **189**:451–60.
- 35 Imanishi T, Akaza T, Kimura A, Tokunaga K, Gojobori T. HLA 1991, Proceedings of the Eleventh International Histocompatibility Workshop and Conference. Tokyo: Oxford University Press, 1992.
- 36 Zea AH, Rodriguez PC, Atkins MB *et al.* Arginase-producing myeloid suppressor cells in renal cell carcinoma patients: a mechanism of tumor evasion. *Cancer Res* 2005; **65**:3044–8.
- 37 Chrzanowska A, Krawczyk M, Baranczyk-Kuzma A. Changes in arginase isoenzymes pattern in human hepatocellular carcinoma. *Biochem Biophys Res Commun* 2008; **377**:337–40.
- 38 Caldwell S, Park SH. The epidemiology of hepatocellular cancer: from the perspectives of public health problem to tumor biology. *J Gastroenterol* 2009; **44** (Suppl. 19):96–101.
- 39 Okamoto M, Oshikawa T, Tano T *et al.* Involvement of Toll-like

- receptor 4 signaling in interferon-gamma production and antitumor effect by streptococcal agent OK-432. *J Natl Cancer Inst* 2003; **95**:316–26.
- 40 Liu S, Yu Y, Zhang M, Wang W, Cao X. The involvement of TNF-alpha-related apoptosis-inducing ligand in the enhanced cytotoxicity of IFN-beta-stimulated human dendritic cells to tumor cells. *J Immunol* 2001; **166**:5407–15.
- 41 Lu G, Janjic BM, Janjic J, Whiteside TL, Storkus WJ, Vujanovic NL. Innate direct anticancer effector function of human immature dendritic cells. II. Role of TNE, lymphotoxin-alpha(1)beta(2), Fas ligand, and TNF-related apoptosis-inducing ligand. *J Immunol* 2002; **168**:1831–9.
- 42 Nicolas A, Cathelin D, Larmonier N *et al*. Dendritic cells trigger tumor cell death by a nitric oxide-dependent mechanism. *J Immunol* 2007; **179**:812–18.
- 43 Stary G, Bangert C, Tauber M, Strohal R, Kopp T, Stingl G. Tumoricidal activity of TLR7/8-activated inflammatory dendritic cells. *J Exp Med* 2007; **204**:1441–51.
- 44 West E, Morgan R, Scott K *et al*. Clinical grade OK432-activated dendritic cells: *in vitro* characterization and tracking during intralymphatic delivery. *J Immunother* 2009; **32**:66–78.
- 45 Hirooka Y, Itoh A, Kawashima H *et al*. A combination therapy of gemcitabine with immunotherapy for patients with inoperable locally advanced pancreatic cancer. *Pancreas* 2009; **38**:e69–74.
- 46 Melief CJ. Cancer immunotherapy by dendritic cells. *Immunity* 2008; **29**:372–83.
- 47 Zitvogel L, Apetoh L, Ghiringhelli F, Kroemer G. Immunological aspects of cancer chemotherapy. *Nat Rev Immunol* 2008; **8**:59–73.
- 48 Ayaru L, Pereira SP, Alisa A *et al*. Unmasking of alpha-fetoprotein-specific CD4(+) T cell responses in hepatocellular carcinoma patients undergoing embolization. *J Immunol* 2007; **178**:1914–22.
- 49 Thurner B, Haendle I, Roder C *et al*. Vaccination with mage-3A1 peptide-pulsed mature, monocyte-derived dendritic cells expands specific cytotoxic T cells and induces regression of some metastases in advanced stage IV melanoma. *J Exp Med* 1999; **190**:1669–78.
- 50 Banchereau J, Palucka AK, Dhodapkar M *et al*. Immune and clinical responses in patients with metastatic melanoma to CD34(+) progenitor-derived dendritic cell vaccine. *Cancer Res* 2001; **61**:6451–8.
- 51 Engell-Noerregaard L, Hansen TH, Andersen MH, Thor Straten P, Svane IM. Review of clinical studies on dendritic cell-based vaccination of patients with malignant melanoma: assessment of correlation between clinical response and vaccine parameters. *Cancer Immunol Immunother* 2009; **58**:1–14.
- 52 Itoh K, Yamada A, Mine T, Noguchi M. Recent advances in cancer vaccines: an overview. *Jpn J Clin Oncol* 2009; **39**:73–80.
- 53 Sugamura K, Asao H, Kondo M *et al*. The common gamma-chain for multiple cytokine receptors. *Adv Immunol* 1995; **59**:225–77.
- 54 Temann UA, Geba GP, Rankin JA, Flavell RA. Expression of interleukin 9 in the lungs of transgenic mice causes airway inflammation, mast cell hyperplasia, and bronchial hyperresponsiveness. *J Exp Med* 1998; **188**:1307–20.
- 55 McMillan SJ, Bishop B, Townsend MJ, McKenzie AN, Lloyd CM. The absence of interleukin 9 does not affect the development of allergen-induced pulmonary inflammation nor airway hyperreactivity. *J Exp Med* 2002; **195**:51–7.
- 56 de Saint-Vis B, Fugier-Vivier I, Massacrier C *et al*. The cytokine profile expressed by human dendritic cells is dependent on cell subtype and mode of activation. *J Immunol* 1998; **160**:1666–76.
- 57 Shanmugham LN, Petrarca C, Frydas S *et al*. IL-15 an immunoregulatory and anti-cancer cytokine. Recent advances. *J Exp Clin Cancer Res* 2006; **25**:529–36.
- 58 Kataoka S, Konishi Y, Nishio Y, Fujikawa-Adachi K, Tominaga A. Antitumor activity of eosinophils activated by IL-5 and eotaxin against hepatocellular carcinoma. *DNA Cell Biol* 2004; **23**:549–60.
- 59 Simson L, Ellyard JI, Dent LA *et al*. Regulation of carcinogenesis by IL-5 and CCL11: a potential role for eosinophils in tumor immune surveillance. *J Immunol* 2007; **178**:4222–9.
- 60 Bystry RS, Aluvihare V, Welch KA, Kallikourdis M, Betz AG. B cells and professional APCs recruit regulatory T cells via CCL4. *Nat Immunol* 2001; **2**:1126–32.