

- Ainsworth BE, et al. International physical activity questionnaire: 12-country reliability and validity. *Med Sci Sports Exerc.* 2003;35(8):1381–95.
20. Murase N, Katsumura T, Ueda C, Inoue S, Shimomitsu T. Validity and reliability of Japanese version of International Physical Activity Questionnaire. *J Health Welfare Stat.* 2002; 49(11):1–9 (in Japanese).
 21. Deng HB, Macfarlane DJ, Thomas GN, Lao XQ, Jiang CQ, Cheng KK, et al. Reliability and validity of the IPAQ-Chinese: the Guangzhou Biobank Cohort study. *Med Sci Sports Exerc.* 2008;40(2):303–7.
 22. Nelson ME, Rejeski WJ, Blair SN, Duncan PW, Judge JO, King AC, et al. Physical activity and public health in older adults: recommendation from the American College of Sports Medicine and the American Heart Association. *Circulation.* 2007;116(9): 1094–105.
 23. The Ministry of Health, Labour and Welfare of Japan. Tokyo: The National Health and Nutrition Survey [cited 2011 May 10]. Available from: http://www.mhlw.go.jp/bunya/kenkou/kenkou_eiyou_chousa.html.
 24. Fukuhara S, Suzukamo Y. Manual of the SF-8 Japanese version. Institute for Health Outcomes & Process Evaluation Research, Kyoto, 2004.
 25. Healy GN, Dunstan DW, Salmon J, Cerin E, Shaw JE, Zimmet PZ, et al. Objectively measured light-intensity physical activity is independently associated with 2-h plasma glucose. *Diabetes Care.* 2007;30(6):1384–9.
 26. Laurson KR, Eisenmann JC, Welk GJ, Wickel EE, Gentile DA, Walsh DA. Combined influence of physical activity and screen time recommendations on childhood overweight. *J Pediatr.* 2008;153(2):209–14.
 27. Eisenmann JC, Bartee RT, Smith DT, Welk GJ, Fu Q. Combined influence of physical activity and television viewing on the risk of overweight in US youth. *Int J Obes (Lond).* 2008;32(4): 613–8.
 28. Hamilton MT, Hamilton DG, Zderic TW. Role of low energy expenditure and sitting in obesity, metabolic syndrome, type 2 diabetes, and cardiovascular disease. *Diabetes.* 2007;56(11): 2655–67.
 29. Sugiyama T, Healy GN, Dunstan DW, Salmon J, Owen N. Is television viewing time a marker of a broader pattern of sedentary behavior? *Ann Behav Med.* 2008;35(2):245–50.
 30. Gorber SC, Tremblay M, Moher D, Gorber B. A comparison of direct vs. self-report measures for assessing height, weight and body mass index: a systematic review. *Obes Rev.* 2007;8(4):307–26.
 31. Spencer EA, Appleby PN, Davey GK, Key TJ. Validity of self-reported height and weight in 4808 EPIC-Oxford participants. *Public Health Nutr.* 2002;5(4):561–5.
 32. Lawlor DA, Bedford C, Taylor M, Ebrahim S. Agreement between measured and self-reported weight in older women. Results from the British Women's Heart and Health Study. *Age Ageing.* 2002;31(3):169–74.
 33. Gunnell D, Berney L, Holland P, Maynard M, Blane D, Frankel S, et al. How accurately are height, weight and leg length reported by the elderly, and how closely are they related to measurements recorded in childhood? *Int J Epidemiol.* 2000; 29(3):456–64.
 34. Inoue M, Toyokawa S, Miyoshi Y, Miyano Y, Suzuki T, Suyama Y, et al. Degree of agreement between weight perception and body mass index of Japanese workers: MY Health Up Study. *J Occup Health.* 2007;49(5):376–81.
 35. Wada K, Tamakoshi K, Tsunekawa T, Otsuka R, Zhang H, Murata C, et al. Validity of self-reported height and weight in a Japanese workplace population. *Int J Obes (Lond).* 2005;29(9): 1093–9.



SHORT PAPER

Open Access

Joint associations of physical activity and screen time with overweight among Japanese adults

Yung Liao^{1*}, Kazuhiro Harada^{2,3}, Ai Shibata³, Kaori Ishii³, Koichiro Oka³, Yoshio Nakamura³, Takemi Sugiyama⁴, Shigeru Inoue⁵ and Teruichi Shimomitsu⁵

Abstract

Background: Although both insufficient physical activity (PA) and high screen time (ST) are independent risk factors for obesity, how the combination of sufficient/insufficient PA and high/low ST could increase obesity risk among the adult population of Japan is not known. This study examined joint associations of PA and ST with overweight among Japanese adults.

Methods: An Internet-based survey collected data on height, weight, self-reported time spent in PA and ST, and sociodemographic variables from 2832 adults. Respondents were categorized into sufficient PA/low ST, sufficient PA/high ST, insufficient PA/low ST, or insufficient PA/high ST categories as per public PA guidelines and the median of ST. Logistic regression analysis examined the odds ratios (OR) of being overweight (body mass index, ≥ 25 kg/m²) according to the categories of PA and ST.

Results: In comparison with the sufficient PA/low ST category, participants in the insufficient PA/high ST category were significantly more likely overweight (OR, 1.48; 95% confidence interval [95%CI], 1.14, 1.93) after adjusting for sociodemographic variables. A significantly higher OR for overweight (including obesity) among insufficient PA/high ST category was also observed in men, but no significant association was found in women.

Conclusions: Both insufficient PA and prolonged ST contribute to overweight and obesity among Japanese adults. Public health initiatives addressing obesity in Japan need to consider both promoting PA and reducing ST, especially in men.

Background

Overweight and obesity increase the risk of developing chronic diseases including cardiovascular disease, hypertension, type 2 diabetes, and certain types of cancer [1,2]. Physically inactive lifestyles are considered to play important roles in the current obesity epidemic [3]. Research has consistently shown that physical activity (PA) is inversely associated with obesity measures [4,5]. Time spent in sitting (sedentary behavior) is also known to be associated with increased risk of obesity, independent of participation in PA [6,7].

Drawing on these research findings, an Australian study has examined the joint association of PA and sedentary behavior with obesity [8]. The study found that those who met PA guideline recommendations but

reported prolonged sedentary time and those with insufficient PA and lower sedentary time had similarly higher likelihood of being overweight compared with those who conducted sufficient PA and were low in sedentary time [8]. However, the combined effect of PA and sedentary behavior on obesity is not known in other countries. Japan offers a unique research opportunity in this context. Although the prevalence of obesity is relatively low in Japan compared with Western countries, it is increasing steadily [9]. In addition, partly because of easily available media-related technologies, television/video viewing and Internet use are highly prevalent and increasing among adults [10]. Since computer and Internet use has been found associated with adult overweight and obesity [11], it is of interest to examine the health impact of screen-based sedentary behavior in the presence (and absence) of PA. This study examined the joint associations of PA and screen-based sedentary behavior with overweight among Japanese adults.

* Correspondence: anthroliao@fuji.waseda.jp

¹Graduate School of Sport Sciences, Waseda University, Nakamura Laboratory 2-579-15 Mikajima Tokorozawa, Saitama 359-1192, Japan
Full list of author information is available at the end of the article



Methods

Participants

Data for this study came from an Internet-based cross-sectional survey in 2009. A total of 9418 adults were randomly selected from the database of a Japanese research service company (with approximately 264,000 registrants) and received invitation e-mails. Of these, 3000 individuals responded to the survey. Detailed methods and procedures have been reported elsewhere [12]. This study received prior approval from the Ethics Committee of Waseda University.

Outcome variable

The outcome variable of this study was body mass index (BMI) calculated from self-reported height and weight dichotomized into normal weight ($< 25 \text{ kg/m}^2$) or overweight (including obese, $\geq 25 \text{ kg/m}^2$), according to the criterion of Japanese National Health and Nutrition Survey [13].

Exposure variable

Exposure variable was calculated from levels of PA and screen time (ST). For PA the Short Version of International Physical Activity Questionnaire (IPAQ-SV) was used. Total time spent in vigorous-intensity PA, moderate-intensity PA, and walking was calculated and dichotomized into "sufficient PA" (≥ 150 minutes/week) or "insufficient PA" (< 150 minutes/week) based on public health guidelines [14]. Test-retest reliability and criterion validity of the Japanese version of IPAQ-SV have been validated [15]. For ST, participants reported their time spent in the following screen-related sedentary behaviors: watching television, videos, and DVDs; Internet use (except work related); and video game use. Reasonable validity and reliability were reported [16]. The sum of the time spent in these behaviors was dichotomized into "low ST" or "high ST" using the median (21 hours/week). According to the levels of PA and ST, participants were classified into the following four categories: sufficient PA/low ST; sufficient PA/high ST; insufficient PA/low ST; and insufficient PA/high ST.

Sociodemographic variables

Data on participants' sex, age (30-39; 40-49; 50-59 years), marital status (married; unmarried), educational level (junior high and high school degree; two-year college degree or equivalent; four-year college or higher degree), job status (full-time job; not full-time job), and household income (less than 5 million yen; 5-10 million yen; more than 10 million yen) were obtained from the research company.

Data analysis

Data for 2832 adults who provided complete information for the study variables were analyzed. Logistic

regression was conducted to estimate the odds ratios (ORs) of being overweight by PA/ST categories adjusted for sociodemographic variables. The sufficient PA/low ST category was the reference for this analysis. Regression analyses were performed for the whole sample, and separately for men and women, based on the gender differences have previously been found [8]. Analysis was conducted using SPSS 15.0; the level of significance was set at $p < 0.05$.

Results

Table 1 shows the sociodemographic characteristics of the total sample and the four PA/ST categories. Overall, 50% of the respondents were men and 74.5% were married. Of the total respondents, 33.6% were aged 30-39 years, 33.3% were aged 40-49 years, and 33.1% were aged 50-59 years. In addition, 47.6% of the participants had graduated from college or graduate school and 62.0% had full-time job, as well as 46.5% had 5-10 million yen income. The proportion of overweight and obese participants was 21.8% for the total sample. Correlation coefficient between PA and ST was -0.17, suggesting that these two behaviors were essentially independent from each other.

Table 2 shows ORs for being overweight or obese by combined categories of PA and ST for the total sample, for men, and for women, adjusting for sociodemographic factors. In the total sample, adults who were insufficient in PA and high in ST were 1.48 times more likely overweight compared with those who engaged in sufficient PA and were low in ST. The other categories (sufficient PA/high ST, insufficient PA/low ST) were not significantly different from the reference category. A similar pattern was observed in men. Compared with the reference category, men who engaged in insufficient PA and high ST were significantly more likely overweight. However, no significant association was found between the PA/ST categories and overweight in women.

Discussion

Japanese adults who engaged in insufficient PA and high ST were about 1.5 times more likely overweight than those with sufficient PA and low ST. Given that insufficient PA or high ST alone was not significantly associated with overweight, our findings suggest that it is the combination of lack of PA and prolonged ST that increases the risk of overweight and obesity in this sample of Japanese adults. This finding on the combined effect of PA and ST on overweight is consistent with a previous study that examined the joint association in the same manner [8]. However, our findings are different from that study in that we found the OR of being overweight was not significantly higher for sufficient PA/high ST and insufficient

Table 1 Sample characteristics by Physical Activity (PA) and Sitting Time (ST) Categories

	Total	Combined Categories of PA and ST				p-value
		Sufficient PA/Low ST	Sufficient PA/High ST	Insufficient PA/Low ST	Insufficient PA/High ST	
No. (%)	2832	905(32%)	656 (23%)	702 (25%)	569 (20%)	-
Sex						< 0.001
Male	50.0%	57.5%	44.7%	50.7%	43.4%	
Female	50.0%	42.5%	55.3%	49.3%	56.6%	
Age group						< 0.001
30-39 years	33.6%	35.6%	27.7%	38.0%	31.6%	
40-49 years	33.3%	29.8%	34.3%	36.8%	33.4%	
50-59 years	33.1%	34.6%	38.0%	25.2%	35.0%	
Marital status						< 0.001
Unmarried	25.5%	20.8%	31.3%	24.5%	27.6%	
Married	74.5%	79.2%	68.8%	75.5%	72.4%	
Educational level						< 0.001
Junior high/high school	26.4%	18.6%	33.5%	25.8%	31.5%	
Two-year college	26.0%	24.7%	25.0%	25.1%	30.4%	
Four-year college/graduate school	47.6%	56.7%	41.5%	49.1%	38.1%	
Job status						< 0.001
Full-time job	62.0%	68.3%	53.7%	69.8%	52.2%	
No full-time job	38.0%	31.7%	46.3%	30.2%	47.8%	
Household income (yen p.a)						< 0.001
< 5 million	38.0%	31.3%	41.6%	36.9%	45.9%	
5-10 million	46.5%	50.4%	42.8%	48.1%	42.7%	
> 10 million	15.5%	18.3%	15.6%	15.0%	11.4%	
BMI						0.182
Normal weight	78.2%	79.4%	79.1%	78.3%	74.9%	
Overweight	21.8%	20.6%	20.9%	21.7%	25.1%	
Mean BMI, kg/m ² (sd)	22.59 (3.53)	22.46 (3.25)	22.51 (3.49)	22.52 (3.50)	22.99 (4.02)	

Abbreviations: PA: physical activity; ST: screen time, p.a.: per annum.

PA/low ST categories. This is also inconsistent with previous studies that demonstrated associations of sedentary time with obesity measures independent of physical activity [6,7]. The inconsistency between this and previous studies may stem from behaviors that were not measured in this study such as nonscreen-based sedentary behaviors (e.g., during work and transport) and light-intensity physical activity. The latter has been shown associated with reduced metabolic risk

independent of moderate-to-vigorous physical activity [17]. It is possible that those in the sufficient PA/high ST category may be low in nonscreen-based sedentary behaviors (they may afford high ST in their leisure time due to less time commitment for work or transport), and those in the insufficient PA/low ST category may be high in light-intensity activity (they may have to cut PA and ST to perform duties such as household chores). Our findings suggest potentially different behavioral

Table 2 Adjusted Odds Ratios of Overweight/Obese by Physical Activity (PA) and Sitting Time (ST) Categories

	Being overweight (BMI, ≥ 25 kg/m ²)					
	Total (n = 2832)		Men (n = 1416)		Women (n = 1416)	
	OR (95% CI)	p-value	OR (95% CI)	p-value	OR (95% CI)	p-value
Sufficient PA/low ST	1.00 (ref.)	-	1.00 (ref.)	-	1.00 (ref.)	-
Sufficient PA/high ST	1.13 (0.87-1.46)	0.37	1.30 (0.94-1.79)	0.11	0.87 (0.55-1.40)	0.57
Insufficient PA/low ST	1.12 (0.87-1.43)	0.39	1.18 (0.88-1.59)	0.28	0.97 (0.61-1.55)	0.91
Insufficient PA/high ST	1.48 (1.14-1.93)	0.003	1.50 (1.08-2.09)	0.02	1.43 (0.92-2.23)	0.11

OR adjusted for sex (whole sample), age, marital status, educational level, job status, and household income.

mechanisms linking physical activity and sedentary behavior with metabolic risk between Japan and Western countries, where previous studies have been conducted. Further research using objective behavioral measures is warranted to explore such differences.

In this study, significant associations between combined PA/ST categories and likelihood of being overweight or obese were found in men but not in women. This pattern of sex difference diverges from the findings of studies conducted in Australia [18], Europe [19], and the United States [20], which have shown stronger associations between sedentary behavior and metabolic health risks in women. One possible explanation is that the prevalence of overweight or obesity is very low among women in this sample. It was 12.5%, which is even lower than the national prevalence for women reported in the Japanese National Health and Nutrition Survey (20%) [21]. It is possible that women in this sample, particularly those who are not very active, may pay close attention to diet so as to control their weight. Future studies should examine diet so as better to understand obesity risks among Japanese women.

Several limitations need to be considered. First, the study used a cross-sectional design; thus it is not possible to make causal inferences. Second, the utilization of IPAQ-SV may cause the overestimation of PA time due to recall bias [22,23]. Third, ST was not measured separately by weekday and weekend, which may contribute to an inaccurate estimation of ST. Fourth, as discussed above, potentially confounding behaviors such as light-intensity activity and diet were not assessed in the study. Finally, the study sample was extracted from the list held by an Internet survey company. Previous studies have indicated that respondents to Internet-based surveys are generally younger, better educated, have higher income and may have greater access to the Internet than respondents to traditional surveys [24,25]. Thus the findings obtained from our sample may not be representative for the entire adult population of Japan.

Regardless of the limitations, our findings suggest the importance of addressing both aspects of physical inactivity (insufficient PA and high ST) to reduce overweight and obesity at the population level. Future health promotion strategies addressing obesity in Japan should focus not only on increasing PA but also on reducing sedentary time, especially in men.

Acknowledgements

This study was supported by a Grant-in-Aid from the Ministry of Health, Labour, and Welfare of Japan (Comprehensive Research on Prevention of Cardiovascular Diseases and Other Lifestyle Related Diseases: H20-Junkankitou-Ippan-001), Grant-in-Aid for Scientific Research (C) 20500604 from the Japan Ministry of Education, Culture, Sports, Science, and Technology, Grants-in-Aid for Scientific Research (No. 20800054) from the Japan Society for the Promotion of Science, and Global COE Program "Sport

Sciences for the Promotion of Active Life" from the Japan Ministry of Education, Culture, Sports, Science, and Technology.

Author details

¹Graduate School of Sport Sciences, Waseda University, Nakamura Laboratory 2-579-15 Mikajima Tokorozawa, Saitama 359-1192, Japan. ²Japan Society for the Promotion of Science, Sumitomo-Ichibancho Bldg., 6 Ichibancho, Chiyoda-ku, Tokyo 102-8471, Japan. ³Faculty of Sport Sciences, Waseda University, 2-579-15 Mikajima Tokorozawa, Saitama 359-1192, Japan. ⁴Baker IDI Heart and Diabetes Institute, Level 4, 99 Commercial Road, Melbourne VIC 3004, Australia. ⁵Department of Preventive Medicine and Public Health, Tokyo Medical University, 6-1-1, Shinjuku-ku, Tokyo, 160-8402, Japan.

Authors' contributions

YL contributed to analysis and interpretation of data and drafted and revised the paper. KH participated in the study design, contributed to analysis and interpretation of data, and revised the paper. AS, KI, KO, SI conceived the study, participated in its design and coordination, and helped in drafting the manuscript. TSU, YN, TSH performed the sequence alignment and helped in drafting the manuscript. All the authors have read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 4 August 2011 Accepted: 30 November 2011
Published: 30 November 2011

References

1. Li Z, Bowerman S, Heber D: Health Ramifications of the Obesity Epidemic. *Surg Clin North Am* 2005, **85**(4):681-701.
2. Visscher TL, Seidell JC: The public health impact of obesity. *Annu Rev Public Health* 2001, **22**:355-75.
3. Kumanyika SK, Obarzanek E, Stettler N, Bell R, Field AE, Fortmann SP, Franklin BA, Gillman MW, Lewis CE, Poston WC, Stevens J, Hong Y, American Heart Association Council on Epidemiology and Prevention, Interdisciplinary Committee for Prevention: Population-based prevention of obesity: The need for comprehensive promotion of healthful eating, physical activity, and energy balance. *Circulation* 2008, **118**:428-464.
4. Dwyer T, Hosmer D, Hosmer T, Venn AJ, Blizzard CL, Granger RH, Cochrane JA, Blair SN, Shaw JE, Zimmet PZ, Dunstan D: The inverse relationship between number of steps per day and obesity in a population-based sample—the AusDiab study. *Int J Obes* 2007, **31**:797-804.
5. Lindstrom M: Means of transportation to work and overweight and obesity: a population-based study in southern Sweden. *Prev Med* 2008, **46**:22-28.
6. Healy GN, Wijndaele K, Dunstan DW, Shaw JE, Salmon J, Zimmet PZ, Owen N: Objectively measured sedentary time, physical activity, and metabolic risk: the Australian Diabetes, Obesity and Lifestyle Study (AusDiab). *Diabetes Care* 2008, **31**:369-371.
7. Jakes RW, Day NE, Khaw KT, Luben R, Oakes S, Welch A, Bingham S, Wareham NJ: Television viewing and low participation in vigorous recreation are independently associated with obesity and markers of cardiovascular disease risk: EPIC-Norfolk population-based study. *Eur J Clin Nutr* 2003, **57**:1089-1096.
8. Sugiyama T, Healy GN, Dunstan DW, Salmon J, Owen N: Joint associations of multiple leisure-time sedentary behaviours and physical activity with obesity in Australian adults. *Int J Behav Nutr Phys Act* 2008, **1**(5):35.
9. McCurry J: Japan battles with obesity. *Lancet* 2007, **369**(9560):451-2.
10. NHK: Survey on Time Use and Leisure Activities. Report from NHK. 2010.
11. Vandelandotte C, Sugiyama T, Gardiner P, Owen N: Associations of leisure-time Internet and computer use with overweight and obesity, physical activity and sedentary behaviors: cross-sectional study. *J Med Internet Res* 2009, **11**(3):e28.
12. Liao Y, Harada K, Shibata A, Ishii K, Oka K, Nakamura Y, Inoue S, Shimomitsu T: Perceived Environmental Factors Associated with Physical Activity among Normal-Weight and Overweight Japanese Men. *Int J Environ Res Public Health* 2011, **8**:931-943.
13. Ministry of Health, Labour and Welfare: *National Health and Nutrition Survey in 2009* Ministry of Health, Labour and Welfare: Tokyo, Japan; 2009.

14. Haskell WL, Lee IM, Pate RR, Powell KE, Blair SN, Franklin BA, Macera CA, Heath GW, Thompson PD, Bauman A: Physical activity and public health: updated recommendation for adults from the American College of Sports Medicine and the American Heart Association. *Med Sci Sports Exerc* 2007, **39**(8):1423-1434.
15. Murase N, Katsumura T, Ueda C, Inoue S, Shimomitsu T: International standardization of physical activity level: reliability and validity study of the Japanese version of the International Physical Activity Questionnaire (IPAQ) (Kosei no Shihyo). *J Health Welfare Statistics* 2003, **49**:1-9, (in Japanese).
16. Salmon J, Owen N, Crawford D, Bauman A, Sallis JF: Physical activity and sedentary behavior: a population-based study of barriers, enjoyment, and preference. *Health Psychol* 2003, **22**(2):178-88.
17. Healy GN, Dunstan DW, Salmon J, Cerin E, Shaw JE, Zimmet PZ, Owen N: Objectively measured light-intensity physical activity is independently associated with 2-h plasma glucose. *Diabetes Care* 2007, **30**(6):1384-9.
18. Healy GN, Dunstan DW, Salmon J, Shaw JE, Zimmet PZ, Owen N: Television time and continuous metabolic risk in physically active adults. *Med Sci Sports Exerc* 2008, **40**(4):639-45.
19. Bertrais S, Beyeme-Ondoua JP, Czernichow S, Galan P, Hercberg S, Oppert JM: Sedentary behaviors, physical activity, and metabolic syndrome in middle-aged French subjects. *Obes Res* 2005, **13**(5):936-44.
20. Ford ES, Kohl HW, Mokdad AH, Ajani UA: Sedentary behavior, physical activity, and the metabolic syndrome among U.S. adults. *Obes Res* 2005, **13**(3):608-14.
21. Handbook of Health and Welfare Statistics 2009, Ministry of Health, Labour and Welfare. [<http://www.mhlw.go.jp/english/database/db-hh/2-1.html>], (accessed on 21 July 2011).
22. Hallal PC, Gomez LF, Parra DC, Lobelo F, Mosquera J, Florindo AA, Reis RS, Pratt M, Sarmiento OL: Lessons learned after 10 years of IPAQ use in Brazil and Colombia. *J Phys Act Health* 2010, **7**(2):S259-64.
23. Rzewnicki R, Vanden Auweele Y, De Bourdeaudhuij I: Addressing overreporting on the International Physical Activity Questionnaire (IPAQ) telephone survey with a population sample. *Public Health Nutr* 2003, **6**(3):299-305.
24. Shibata A, Oka K, Harada K, Nakamura Y, Muraoka I: Psychological, social, and environmental factors to meeting physical activity recommendations among Japanese adults. *Int J Behav Nutr Phys Act* 2009, **28**(6):60.
25. Rhodes SD, Bowie DA, Hergenrather KC: Collecting behavioural data using the world wide web: considerations for researchers. *J Epidemiol Community Health* 2003, **57**(1):68-73.

doi:10.1186/1479-5868-8-131

Cite this article as: Liao et al.: Joint associations of physical activity and screen time with overweight among Japanese adults. *International Journal of Behavioral Nutrition and Physical Activity* 2011 **8**:131.

Submit your next manuscript to BioMed Central
and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



Sociodemographic Determinants of Pedometer-Determined Physical Activity Among Japanese Adults

Shigeru Inoue, MD, PhD, Yumiko Ohya, PhD, Yuko Odagiri, MD, PhD, Tomoko Takamiya, MD, PhD, Kenichi Suijo, MEd, Masamitsu Kamada, MEd, Shinpei Okada, MEd, Catrine Tudor-Locke, PhD, and Teruichi Shimomitsu, MD, PhD

Background: Although previous studies have reported physical activity and its sociodemographic determinants using self-report measures, there have been few studies using pedometers.

Purpose: To ascertain pedometer-determined physical activity and its sociodemographic determinants among community residents living in four Japanese cities.

Methods: A cross-sectional mail survey was conducted from February 2007 to January 2008 with a sample of 4000 residents (aged 20–69 years and 50% male) who were randomly selected from the registry of residential addresses. Complete responses for both questionnaire and pedometer were obtained from 790 residents (48.3±13.7 years, 46.7% male). Associations of 11 sociodemographic variables with steps per day were examined using multiple logistic regression analyses. Data were analyzed in 2010.

Results: Men averaged 8763±3497 steps/day and women averaged 8242±3277 steps/day. Further, 29.0% of men and 27.8% of women walked ≥10,000 steps/day. City of residence, good self-rated health, low educational attainment, and not owning a car were associated with taking ≥10,000 steps/day in men, whereas employed status and dog ownership were associated with walking ≥10,000 steps/day in women.

Conclusions: The results contribute to understanding of step-defined physical activity and its sociodemographic determinants. A diversity of step counts by sociodemographic variables clarifies specific populations among Japanese who are in need of intervention to promote physical activity. (Am J Prev Med 2011;40(5):566–571) © 2011 American Journal of Preventive Medicine

Introduction

Physical activity promotion is one of the priorities for chronic disease prevention.^{1,2} To develop effective intervention strategies, solid evidence of physical activity patterns among specific population groups is needed. Previous studies have reported the association between sociodemographic variables and physical activity.^{3–7} However, most of these studies (1) were conducted in Western countries and (2) used

self-reported physical activity, with a few exceptions.^{8,9} Thus, the generalizability of findings to countries of a different culture, such as Japan, is unclear.

Further, there remains the potential for information bias from self-reported physical activity.¹⁰ Step-counting devices provide an objective output and are becoming widespread as intervention tools.^{11,12} Therefore, data related to step-defined physical activity are increasingly important.

Thus, the objectives of the present study are to (1) report step-defined physical activity levels and (2) examine sociodemographic characteristics of people who achieve ≥10,000 steps/day¹³ using a randomly selected community sample from four Japanese cities.

Methods

Participants and Data Collection

This cross-sectional study was a part of larger project¹⁴ to investigate physical activity environment. Data were collected from February to March 2007 and from December 2007 to January 2008.

From the Department of Preventive Medicine and Public Health, Tokyo Medical University (Inoue, Ohya, Odagiri, Takamiya, Suijo, Shimomitsu), Tokyo; Physical Education and Medicine Research Center Unnan (Kamada), Unnan, Shimane; Physical Education and Medicine Research Foundation (Okada), Tomi, Nagano, Japan; and Walking Behavior Laboratory, Pennington Biomedical Research Center (Tudor-Locke), Baton Rouge, Louisiana

Address correspondence to: Shigeru Inoue, MD, PhD, Department of Preventive Medicine and Public Health, Tokyo Medical University, 6-1-1, Shinjuku, Shinjuku-ku, Tokyo 160-8402, Japan. E-mail: inoue@tokyo-med.ac.jp.

0749-3797/\$17.00

doi: 10.1016/j.amepre.2010.12.023

Four thousand residents (aged 20–69 years, 50% male) living in four Japanese cities (Koganei, Tsukuba, Shizuoka, and Kagoshima) were randomly selected from the registry of residential addresses, stratified by gender, age, and city of residence. Locations, areas, and population sizes of four cities are indicated in Appendix A (available online at www.ajpm-online.net). Koganei lies approximately at the center of Tokyo. Tsukuba is located 50 km northeast of Tokyo within commuting distance from Tokyo. Shizuoka and Kagoshima are the prefectural capitals in mid- and west Japan.

The study was conducted by mail. At first, participants were asked about sociodemographic characteristics and subsequently invited to wear an accelerometer. If participants consented to join, the accelerometer was mailed out. Of the targeted 4000 residents, 1508 responded to the questionnaire (37.7%). Of these, 886 offered to wear the accelerometer, but valid accelerometer data were obtained from 790 participants (overall response rate: 19.8%). All participants provided signed informed consent. The current study received prior approval from the Tokyo Medical University Ethics Committee. Further details of the survey were reported in a previous article.¹⁴

Assessment of Step Counts

Participants were asked to wear an accelerometer (Suzuken Life-recorder Ex 4 second version, Nagoya, Japan) for 7 consecutive days. This device has a step-counting function with a 35-day memory. Acceleration signals were used only for calculation of device use (wearing and nonwearing time). Previous studies^{15,16} have validated the step-counting feature of this device. The step data were systematically processed according to the following rules: (1) non-wear time was defined as no acceleration signal for ≥ 30 consecutive minutes; (2) a recorded day was deemed valid if it was worn at least 10 hours that day^{17,18}; and (3) to be included in the analysis the participant had to have 3 or more valid days of data.¹⁸ Finally, mean steps/day were calculated based on steps of valid days.

Sociodemographic Variables

Gender, age, and city of residence were obtained from the registry of residential addresses. Information on height, weight, self-rated health, education, employment, marital status, child (defined as junior high school students or younger; aged ≤ 15 years) in household, household motor vehicle, and dog ownership were obtained by questionnaire. BMI was calculated from self-reported weight and height.

Statistical Analyses

Multiple logistic regression analyses were conducted to examine the relationships between sociodemographic variables and steps/day. Steps/day were recoded into three categories, sedentary to low active (Se/LA), < 7500 steps/day; somewhat active (SA), 7500–9999 steps/day; and active to highly active (A/HA), $\geq 10,000$ steps/day.¹³ All 11 sociodemographic variables were included in the model. Cities of residence were included in the model as dummy variables. Following the analyses of the overall sample, stratified analyses by gender were conducted. The odds of higher step counts (SA and A/HA) compared with Se/LA by 11 sociodemographic attributes were calculated. Significance was considered to be $p < 0.05$. Analyses were conducted in 2010 using SPSS, version 17.0.

Results

Men accounted for 46.7% of participants. Mean age was 48.3 ± 13.7 years overall. Mean steps/day were 8763 ± 3497 steps/day in men and 8242 ± 3277 steps/day in women. The prevalence of taking $\geq 10,000$ steps/day was 29.0% in men and 27.8% in women. Further information about participants' characteristics is shown in Appendix B (available online at www.ajpm-online.net).

In multivariate analyses (Table 1), city of residence (Tsukuba or Koganei); good self-rated health; being employed; not having a household motor vehicle; and dog ownership were related to either SA or A/HA or both SA and A/HA. ORs (95% CI) of A/HA, that is, engaging in $\geq 10,000$ steps/day (vs Se/LA) were 1.80 (1.04, 3.12) for Tsukuba residents and 2.03 (1.17, 3.52) for Koganei residents compared with Kagoshima residents; 1.70 (1.18, 2.43) for people with good self-rated health; 1.67 (1.06, 2.63) for employed; 2.40 (1.28, 4.49) for nonowners of motor vehicles; and 1.77 (1.13, 2.76) for dog owners. According to stratified analyses by gender; city of residence (Tsukuba, Koganei, or Shizuoka); good self-rated health; lower educational attainment; and having no household motor vehicle were associated with more steps/day in men. High step-defined activity was associated with good self-rated health, being employed, and dog ownership in women.

Discussion

The present study showed the step-defined physical activity level of residents in four Japanese cities and its association with sociodemographic variables. Most previous studies conducted in Western countries have been based on self-report. Thus, the findings of the current study add new evidence from two perspectives. First, the present study was conducted in Japan and therefore extends understanding of physical activity determinants to include a distinct culture. Second, objective step data were collected. Step-defined physical activity is relevant to health promotion applications because of the potential of step counter to be widely used as an intervention tool.

Participants in the current study appear to be more active compared with national surveys in Japan¹⁹ and the U.S.²⁰ Japanese survey¹⁹ in 2007 reported that men and women took 7321 steps/day and 6267 steps/day, respectively. Differences in sampling and response rates may explain some of this discrepancy. The current sample did not include older adults (aged ≥ 70 years) and was primarily living in urban settings. In the 2005–2006 National Health and Nutrition Examination Survey²⁰ of the U.S., accelerometer-determined physical activity data treated to approximate pedome-

Table 1. ORs for active people, as determined by pedometer, by sociodemographic variables

	Overall (N=790)				Men (n=369)	
	SA (vs Se/LA)		A/HA (vs Se/LA)		SA (vs Se/LA)	
	7500-9999 (vs <7499)		≥10,000 (vs <7499)		7500-9999 (vs <7499)	
	OR (95% CI)	p-value ^a	OR (95% CI)	p-value ^a	OR (95% CI)	p-value ^a
Gender	—	—	—	—	—	—
Men	1.10 (0.77, 1.59)	0.595	1.07 (0.73, 1.55)	0.741	—	—
Women	1.00	—	1.00	—	1.00	—
Age (years)	—	—	—	—	—	—
20-39	1.46 (0.81, 2.63)	0.205	1.24 (0.69, 2.24)	0.474	2.31 (0.88, 6.10)	0.090
40-59	1.58 (0.96, 2.58)	0.069	1.14 (0.70, 1.85)	0.599	1.69 (0.80, 3.57)	0.172
60-69	1.00	—	1.00	—	1.00	—
City of residence	—	—	—	—	—	—
Tsukuba	1.40 (0.84, 2.34)	0.194	1.80 (1.04, 3.12)	0.035	2.69 (1.27, 5.71)	0.010
Koganei	1.59 (0.95, 2.67)	0.079	2.03 (1.17, 3.52)	0.012	2.16 (0.96, 4.87)	0.062
Shizuoka	0.98 (0.58, 1.63)	0.927	1.43 (0.84, 2.45)	0.187	1.20 (0.55, 2.63)	0.649
Kagoshima	1.00	—	1.00	—	1.00	—
BMI	—	—	—	—	—	—
<25	0.93 (0.60, 1.44)	0.737	1.08 (0.68, 1.72)	0.732	0.77 (0.43, 1.36)	0.362
≥25	1.00	—	1.00	—	1.00	—
Self-rated health	—	—	—	—	—	—
Good	1.52 (1.07, 2.16)	0.018	1.70 (1.18, 2.43)	0.004	1.38 (0.82, 2.34)	0.228
Fair or poor	1.00	—	1.00	—	1.00	—
Education (years)	—	—	—	—	—	—
≥13	1.04 (0.71, 1.51)	0.857	0.73 (0.50, 1.07)	0.103	0.95 (0.52, 1.71)	0.853
≤12	1.00	—	1.00	—	1.00	—
Employment status	—	—	—	—	—	—
Employed	1.55 (0.99, 2.43)	0.053	1.67 (1.06, 2.63)	0.028	2.15 (0.85, 5.48)	0.108
Not employed	1.00	—	1.00	—	1.00	—
Marital status	—	—	—	—	—	—
Married	1.53 (0.93, 2.52)	0.094	1.56 (0.94, 2.59)	0.089	1.94 (0.83, 4.55)	0.128
Not married	1.00	—	1.00	—	1.00	—
Child in household	—	—	—	—	—	—
Yes	0.91 (0.60, 1.38)	0.669	0.65 (0.41, 1.01)	0.056	1.04 (0.54, 1.98)	0.911
No	1.00	—	1.00	—	1.00	—
Household motor vehicle (no.)	—	—	—	—	—	—
0	1.88 (1.00, 3.56)	0.052	2.40 (1.28, 4.49)	0.006	4.24 (1.38, 13.01)	0.012
≥1	1.00	—	1.00	—	1.00	—
Dog ownership	—	—	—	—	—	—
Yes	1.65 (1.06, 2.55)	0.026	1.77 (1.13, 2.76)	0.012	1.40 (0.74, 2.67)	0.301
No	1.00	—	1.00	—	1.00	—

(continued on next page)

^ap-values were calculated by multiple logistic regression analyses adjusted for all other sociodemographic variables listed in the table.

A/HA, active and high active; SA, somewhat active; Se/LA, sedentary and low active

Table 1. (continued)

A/HA (vs Se/LA)		Women (n=421)			
		SA (vs Se/LA)		A/HA (vs Se/LA)	
≥10,000 (vs <7499)		7500-9999 (vs <7499)		≥10,000 (vs <7499)	
OR (95% CI)	p-value ^a	OR (95% CI)	p-value ^a	OR (95% CI)	p-value ^a
—	—	—	—	—	—
—	—	—	—	—	—
—	—	—	—	—	—
—	—	—	—	—	—
2.52 (0.94, 6.78)	0.067	1.02 (0.47, 2.23)	0.952	0.83 (0.38, 1.83)	0.649
1.37 (0.64, 2.89)	0.416	1.29 (0.65, 2.57)	0.468	1.03 (0.52, 2.03)	0.942
1.00	—	1.00	—	1.00	—
—	—	—	—	—	—
2.90 (1.26, 6.68)	0.013	0.77 (0.37, 1.60)	0.477	1.31 (0.62, 2.78)	0.481
3.22 (1.34, 7.73)	0.009	1.28 (0.63, 2.58)	0.493	1.70 (0.81, 3.55)	0.160
2.37 (1.06, 5.32)	0.036	0.85 (0.42, 1.71)	0.649	1.02 (0.49, 2.14)	0.960
1.00	—	1.00	—	1.00	—
—	—	—	—	—	—
1.06 (0.57, 1.98)	0.859	1.39 (0.65, 2.96)	0.397	1.01 (0.50, 2.04)	0.983
1.00	—	1.00	—	1.00	—
—	—	—	—	—	—
2.46 (1.41, 4.29)	0.001	1.81 (1.11, 2.94)	0.018	1.30 (0.80, 2.12)	0.294
1.00	—	1.00	—	1.00	—
—	—	—	—	—	—
0.46 (0.25, 0.84)	0.011	1.16 (0.69, 1.93)	0.582	1.00 (0.59, 1.68)	0.990
1.00	—	1.00	—	1.00	—
—	—	—	—	—	—
1.15 (0.49, 2.66)	0.749	1.43 (0.83, 2.45)	0.198	1.92 (1.09, 3.40)	0.024
1.00	—	1.00	—	1.00	—
—	—	—	—	—	—
2.15 (0.88, 5.24)	0.093	1.44 (0.75, 2.76)	0.272	1.48 (0.77, 2.86)	0.239
1.00	—	1.00	—	1.00	—
—	—	—	—	—	—
0.77 (0.38, 1.57)	0.475	0.78 (0.44, 1.37)	0.385	0.60 (0.34, 1.09)	0.092
1.00	—	1.00	—	1.00	—
—	—	—	—	—	—
5.47 (1.81, 16.51)	0.003	1.17 (0.52, 2.65)	0.706	1.47 (0.66, 3.29)	0.347
1.00	—	1.00	—	1.00	—
—	—	—	—	—	—
1.39 (0.71, 2.72)	0.341	1.78 (0.96, 3.29)	0.065	2.17 (1.18, 4.00)	0.013
1.00	—	1.00	—	1.00	—

ter-determined scaling indicated that American men took 7431 steps/day and women took 5756 steps/day. Although the differences of sampling strategy and of device should be considered, participants in the current study appear to be more active.

Gender and age were not significantly related to steps/day in multivariate analyses. In previous studies,^{3,4} men and younger adults were more active than women and older adults. In the present study, multivariate analyses included a wide range of sociodemographic variables. Thus, after adjustment for these variables, the effect of gender and age alone appears to be lessened. People in Tsukuba and Koganei were more active than those in Kagoshima. Both Tsukuba and Koganei are in the Greater Tokyo Area.

According to the Nationwide Person Trip Survey in Japan,²¹ 22.2% of commuters drive in metropolitan areas (including Koganei); 31.6% in suburban cities of metropolitan areas (including Tsukuba); and 63.0% in core local cities (including Shizuoka and Kagoshima). The differences among cities might be related to existing public transport networks. Koganei also has high walkability, characterized by high density and land use mix.^{22,23} Thus, the importance of environmental features was suggested.

Sociodemographic variables associated with steps/day differed by gender. High education level was related to lower steps/day in men. This is inconsistent with previous findings.^{3,4,6,7} However, many previous studies have focused on leisure-time physical activity. It is likely that the sociodemographic determinants of total physical activity are different. Physically demanding jobs among less-educated men may be one possible reason for this result. Employment status was an important factor for women. Women without jobs outside the home might have fewer opportunities to walk for transport. Association of motor vehicle ownership and steps/day suggested a car-reliant lifestyle among men. In contrast, women may feel an obligation to walk their dog and increase their activity as a result.

There were some limitations in the current study. First, the response rate might be considered low. If participants were reactive to wearing the accelerometer, they might have been active regardless of their sociodemographic characteristics, leading to an underestimation of the association between sociodemographic variables and steps/day. Second, the study setting was relatively urban because of the original purpose of this survey. Research including rural areas is also needed in the future.

In spite of these limitations, the present study contributes to understanding of step-defined physical activity and its sociodemographic determinants. A diversity of steps/day by sociodemographic variables clarifies specific

populations in need of intervention to promote physical activity.

This study was supported by a grant-in-aid from the Japan Ministry of Health, Labor, and Welfare (ID: H19-Junkankitou-Ippan-008 and H20-Junkankitou-Ippan-001) and a grant-in-aid for scientific research (C: 17590556) from the Japan Ministry of Education, Culture, Sports, Science, and Technology.

No financial disclosures were reported by the authors of this paper.

References

1. Physical Activity Guidelines Advisory Committee. Physical Activity Guidelines Advisory Committee report, 2008. Washington DC: USD-HHS, 2008.
2. Haskell WL, Lee IM, Pate RR, et al. Physical activity and public health: updated recommendation for adults from the American College of Sports Medicine and the American Heart Association. *Med Sci Sports Exerc* 2007;39(8):1423–34.
3. Sallis JF, Owen N. Physical activity and behavioral medicine. Thousand Oaks CA: Sage, 1999:110–34.
4. Trost SG, Owen N, Bauman AE, Sallis JF, Brown W. Correlates of adults' participation in physical activity: review and update. *Med Sci Sports Exerc* 2002;34(12):1996–2001.
5. Bergman P, Grjibovski AM, Hagstromer M, Bauman A, Sjostrom M. Adherence to physical activity recommendations and the influence of socio-demographic correlates—a population-based cross-sectional study. *BMC Public Health* 2008;8:367.
6. Bertrais S, Preziosi P, Mennen L, Galan P, Hercberg S, Oppert JM. Sociodemographic and geographic correlates of meeting current recommendations for physical activity in middle-aged French adults: the Supplementation en Vitamines et Mineraux Antioxydants (SUVIMAX) study. *Am J Public Health* 2004;94(9):1560–6.
7. Pan SY, Cameron C, Desmeules M, Morrison H, Craig CL, Jiang X. Individual, social, environmental, and physical environmental correlates with physical activity among Canadians: a cross-sectional study. *BMC Public Health* 2009;9:21.
8. Shibata A, Oka K, Nakamura Y, Muraoka I. Prevalence and demographic correlates of meeting the physical activity recommendation among Japanese adults. *J Phys Act Health* 2009;6(1):24–32.
9. Hagstromer M, Oja P, Sjostrom M. Physical activity and inactivity in an adult population assessed by accelerometry. *Med Sci Sports Exerc* 2007;39(9):1502–8.
10. Sallis JF, Saelens BE. Assessment of physical activity by self-report: status, limitations, and future directions. *Res Q Exerc Sport* 2000; 71(2S):S1–14.
11. Bravata DM, Smith-Spangler C, Sundaram V, et al. Using pedometers to increase physical activity and improve health: a systematic review. *JAMA* 2007;298(19):2296–304.
12. Richardson CR, Newton TL, Abraham JJ, Sen A, Jimbo M, Swartz AM. A meta-analysis of pedometer-based walking interventions and weight loss. *Ann Fam Med* 2008;6(1):69–77.
13. Tudor-Locke C, Bassett DR Jr. How many steps/day are enough? Preliminary pedometer indices for public health. *Sports Med* 2004;34(1):1–8.
14. Inoue S, Ohya Y, Odagiri Y, et al. Association between perceived neighborhood environment and walking among adults in 4 cities in Japan. *J Epidemiol* 2010;20(4):277–86.

15. Schneider PL, Crouter SE, Lukajic O, Bassett DR Jr. Accuracy and reliability of 10 pedometers for measuring steps over a 400-m walk. *Med Sci Sports Exerc* 2003;35(10):1779–84.
16. Schneider PL, Crouter SE, Bassett DR. Pedometer measures of free-living physical activity: comparison of 13 models. *Med Sci Sports Exerc* 2004;36(2):331–5.
17. Troiano RP, Berrigan D, Dodd KW, Masse LC, Tilert T, McDowell M. Physical activity in the U.S. measured by accelerometer. *Med Sci Sports Exerc* 2008;40(1):181–8.
18. Trost SG, McIver KL, Pate RR. Conducting accelerometer-based activity assessments in field-based research. *Med Sci Sports Exerc* 2005;37(11S):S531–43.
19. The Ministry of Health, Labour, and Welfare of Japan. The National Health and Nutrition Survey 2007, 2010. www.mhlw.go.jp/bunya/kenkou/eiyou09/.
20. Tudor-Locke C, Johnson WD, Katzmarzyk PT. Accelerometer-determined steps per day in U.S. adults. *Med Sci Sports Exerc* 2009;41(7):1384–91.
21. The Ministry of Land, Infrastructure and Transport of Japan. The report of the 4th Nationwide Person Trip Survey 2005, 2010, www.mlit.go.jp/crd/tosiko/zpt/index.html.
22. Owen N, Humpel N, Leslie E, Bauman A, Sallis JF. Understanding environmental influences on walking: review and research agenda. *Am J Prev Med* 2004;27(1):67–76.
23. Saelens BE, Handy SL. Built environment correlates of walking: a review. *Med Sci Sports Exerc* 2008;40(7S):S550–66.

Appendix

Supplementary data

Supplementary data associated with this article can be found, in the online version, at doi:10.1016/j.amepre.2010.12.023.

Have you seen the *AJPM* website lately?
Visit www.ajpm-online.net today!

Differences in Association of Walking for Recreation and for Transport With Maximum Walking Speed in an Elderly Japanese Community Population

Masamitsu Kamada, Jun Kitayuguchi, Kuninori Shiwaḡu, Shigeru Inoue, Shimpei Okada, and Yoshiteru Mutoh

Background: Physical activity contributes to maintaining functional ability later in life. Specific relationships between walking for particular purposes (eg, recreation or transport) and functional ability are not clear. It is useful for planning health promotion strategies to clarify whether walking time for recreation, or walking time for transport has the stronger relationship with maximum walking speed (MWS), a determinant of functional ability later in life in the elderly. **Methods:** A cross-sectional survey was conducted in 2007 using a sample of 372 community-dwelling elderly people aged 60 to 87 years in Mitoya Town, Unnan City, rural Japan. Associations with MWS were examined for self-reported weekly times of walking for recreation and for transport using multiple linear regression analyses. **Results:** Both in men and women, walking time for recreation was significantly associated with MWS after controlling for age, height, weight, hip and knee pain, and a number of chronic diseases (men: $\beta = 0.18$, $P = .024$; women: $\beta = 0.17$, $P < .01$). However, walking time for transport was not significantly associated with MWS (men: $\beta = -0.094$, $P = .24$; women: $\beta = -0.040$, $P = .50$). **Conclusions:** Walking for recreation may contribute to maintaining functional abilities such as MWS in the elderly.

Keywords: physical activity, physical fitness, public health, older adults, epidemiology

Physical activity plays a role in maintaining functional ability later in life.¹⁻⁴ Walking is the most common physical activity behavior of adults and the elderly.⁵⁻⁷ Previous studies have shown the health benefit of walking behaviors.^{1,3,4,8,9} However, there has been little study of the differences in the purposes of walking (eg, recreational vs. nonrecreational). If the intensities of walking for recreation and for transport are different, effects of those behaviors on health outcomes should be different, and taking such differences into account could be useful, in health education and the promotion of walking behaviors. Differences between specific associations of walking for recreation or for transport with health benefits, such as functional ability, have not been clearly identified.¹⁰

Maximum walking speed (MWS) is a good predictor of bone health¹¹ and functional dependence¹² in the elderly and believed to be susceptible to walking behavior.¹³ The purpose of this study was to clarify specific

relationships between walking for particular purposes and maximum walking speed in the elderly. We hypothesized that walking for recreation may have a greater influence on functional ability, such as maximum walking speed,¹¹ because this type of walking is assumed to be conducted with higher levels of intensity compared with walking for transport.

Methods

Study Location and Subjects

A cross-sectional survey was conducted in community-dwelling elderly in Mitoya Town (population 8241, area 82.7km²), Unnan City, Shimane Prefecture, in western rural Japan in October 2007. This study was a part of a Shimane study¹⁴ conducted concomitantly with an annual health examination. All adults aged 20 years and above then living in Mitoya were invited to participate in the study by direct mail, local public broadcasts, cable television, circulars called *kairanban*, and city newsletters. Figure 1 shows the flowchart of the recruitment. The following inclusion criteria were used for each subject: aged 60 years and over, community-dwelling, and a participant in a Shimane study conducted concomitantly with the subject's annual health examination. Exclusion criteria applied were those individuals in assisted living facilities, those requiring nursing care, and those who walked

Kamada and Kitayuguchi are with the Physical Education and Medicine Research Center UNNAN, Unnan, Japan. Shiwaḡu is with the Dept of Environmental and Preventive Medicine, Shimane University, Izumo, Japan. Inoue is with the Dept of Preventive Medicine and Public Health, Tokyo Medical University, Tokyo, Japan. Okada is with the Physical Education and Medicine Research Foundation, Tomi, Japan. Mutoh is with the Graduate School of Education, University of Tokyo, Tokyo, Japan.




Figure 1 — Subject recruitment.

using an assistive device. Data on 372 elderly people who qualified as subjects (132 men and 240 women) aged 60 to 87 years (70.5 ± 5.8) were used in the analyses.

Walking Behavior

We investigated average walking time for recreation (including for both exercise and leisure) and for transport (to or from work, shopping, to access public transportation, etc.) over a typical 7-day period (min/w), using a self-administered questionnaire and a face-to-face interview for confirmation. Subjects were asked to address only their walking activities lasting for at least 10 minutes at a time. This questionnaire and interview confirmation method had an acceptable 1-week test-retest reliability (Spearman's $\rho = 0.79$, $P < .01$) in 43 elderly people (22 men and 21 women) aged 75.7 ± 4.6 years in Kisuki Town, adjacent to Mitoya, in Unnan City. A validity test with a 1-week walking diary was also conducted in Kisuki with another group of 44 elderly people (12 men and 32 women) aged 72.7 ± 5.2 years. These subjects were provided with walking diaries and were asked to make daily records of their start and finish times and purpose of all walking activities continuing for at least 10 minutes. This validity test showed acceptable results of walking times for recreation (Spearman's $\rho = 0.49$, $P < .01$) and for transport (Spearman's $\rho = 0.38$, $P = .013$). Results of a validity test for total walking time with average daily step counts recorded by a uniaxial accelerometer (Life-corder, Suzuken Co. Ltd., Nagoya, Japan)^{15,16} were also acceptable (Spearman's $\rho = 0.41$, $P < .01$) in 93 elderly people (35 men and 58 women) aged 73.7 ± 5.0 years in Kisuki. The questionnaire used in this study is originally in Japanese and available online.¹⁷

Maximum Walking Speed (MWS)

Walking speed was measured once by trained examiners using digital stopwatches and a 10-m course set between the 2- and 12-m marks of a 14-m straight, flat, indoor walkway. We asked subjects to walk as fast as they safely could without running on the walkway course. The high reliability of walking (gait) speed measurements have been reported previously.¹³

Other Measurements

Subjects underwent anthropometric measurements and questionnaires with face-to-face interviews covering chronic musculoskeletal pain in the hip and knee,^{18,19} and chronic diseases (cerebrovascular disorder, diabetes, heart disease, liver disease, lung disease, hypertension).^{2,20}

Data and Statistical Analyses

Differences by gender were examined by *t* test for variables with normal distributions and Mann-Whitney test for variables with nonnormal distributions. Correlations with MWS were calculated by Pearson correlation for age, height, weight, and body mass index and Spearman's ρ for musculoskeletal pain, a number of chronic diseases, walking time for recreation, and walking time for transport with nonnormal distributions. The correlations between walking times for recreation and for transport were calculated by Spearman's ρ . Multiple linear regression models were used with MWS as the dependent variable and walking times for recreation and for transport as the independent variables separately by gender. Adjustments were made for selected variables whose *P* values were 0.10 and under in univariate analyses with

MWS based on age, height, weight, hip and knee pain, and/or a number of chronic diseases for model 1 and by all these potential confounding factors for model 2. Data on walking for recreation and for transport were simultaneously analyzed in multiple linear regression analyses. Significance was set at a level of $P < .05$. Analyses were conducted using SPSS 14.0J for Windows (SPSS Japan Inc., Tokyo, Japan, 2006)

This study was approved by the research ethics committee of the Physical Education and Medicine Research Center UNNAN. All the subjects took part in the study after signing informed consent forms.

Results

Subject characteristics are shown in Table 1. The mean values of MWS were 2.2 ± 0.4 m/s in men and 2.0 ± 0.4 m/s in women. The mean walking times for recreation and for transport were 66 ± 101 min/w and 26 ± 51 min/w, respectively, in men, and 47 ± 87 min/w and 35 ± 70 min/w, respectively, in women. The median values of walking times for recreation and for transport were 0 min/w for both men and women. About 40% of the subjects engaged in walking for both recreation and transport, while 37.9% of the subjects engaged in neither. Although

a direct comparison cannot be made, the prevalence of recreational walkers in our study was similar to national data of annual participation rates of walking for recreation in Japanese elderly (47.2% for 60–69 year-olds; 35.6% for 70+ year-olds).²¹ Women were more likely to be younger, have smaller height and weight, have hip and knee pain, and have lower MWS, compared with men.

MWS was significantly correlated with age and walking time for transport in men and with age, height, hip and knee pain, number of chronic diseases, and walking time for recreation in women (Table 2). The correlation between walking times for recreation and for transport was not significant (men: Spearman's $\rho = -0.048$, $P = .59$; women: Spearman's $\rho = 0.038$, $P = .56$, not shown in tables).

Table 3 shows the results of multiple linear regression analyses. Both in men and women, walking time for recreation was positively and significantly associated with MWS after controlling for potential confounding factors. However, walking time for transport was not significantly associated with MWS. Each model explained 22% of the variance of MWS in men and 25% in women. Results did not change much and the model R^2 values decreased somewhat when walking times were analyzed as categorical variables divided into nearly 3 equal parts

Table 1 Subject Characteristics (Shimane, Japan, 2007)

	Men (n = 132)	Women (n = 240)	<i>P</i> ^a
	Mean \pm SD or Number (%)	Mean \pm SD or Number (%)	
Age (years)	71.7 \pm 5.8	69.8 \pm 5.7	<0.01
60–64	17 (12.9)	48 (20.0)	
65–69	30 (22.7)	73 (30.4)	
70–74	41 (31.1)	69 (28.8)	
75–79	33 (25.0)	35 (14.6)	
80+	11 (8.3)	15 (6.3)	
Height (cm)	161.1 \pm 5.8	148.9 \pm 5.2	<0.01
Weight (kg)	58.3 \pm 8.6	50.1 \pm 7.0	<0.01
Body mass index (kg/m ²)	22.4 \pm 2.8	22.6 \pm 2.9	0.60
Hip and knee pain	63 (47.7)	154 (64.2)	<0.01
Number of chronic diseases ^b			0.08
0	56 (42.4)	118 (49.4)	
1	56 (42.4)	102 (42.7)	
2+	20 (15.2)	19 (7.9)	
Walking time for recreation (min/w)	66 \pm 101	47 \pm 87	0.06
Walking time for transport (min/w)	26 \pm 51	35 \pm 70	0.12
Maximum walking speed (m/s)	2.2 \pm 0.4	2.0 \pm 0.4	<0.01

Note. Sample sizes vary due to missing values.

^a Compares prevalences by gender using *t* test for age, height, weight, body mass index, and maximum walking speed and Mann-Whitney test for others.

^b Number of the following diseases: cerebrovascular disorder, diabetes, heart disease, liver disease, lung disease, hypertension.

Table 2 Correlations Between Maximum Walking Speed and the Measured Items (Shimane, Japan, 2007)

	Men (n = 132)		Women (n = 240)	
	r or ρ^a	P	r or ρ^a	P
Age (years)	-0.45	<0.001	-0.40	<0.001
Height (cm)	0.08	0.37	0.29	<0.001
Weight (kg)	0.05	0.61	0.06	0.37
Body mass index (kg/m ²)	0.00	0.97	-0.10	0.13
Hip and knee pain	0.082	0.36	-0.25	<0.001
Number of chronic diseases ^b	-0.17	0.058	-0.13	0.044
Walking time for recreation (min/w)	0.17	0.059	0.19	<0.01
Walking time for transport (min/w)	-0.18	0.040	-0.12	0.061

Note. Sample sizes vary due to missing values.

^a Pearson's correlation coefficients (*r*) for age, height, weight, and body mass index and Spearman's ρ for other variables with nonnormal distributions.

^b Number of the following diseases: cerebrovascular disorder, diabetes, heart disease, liver disease, lung disease, hypertension.

Table 3 Multiple Linear Regression Analyses for the Assessment of Associations Between Walking Behavior and Maximum Walking Speed (Shimane, Japan, 2007)

	Model 1 ^a		Model 2 ^a	
	β^b	P	β^b	P
Men (n = 132)				
Walking behavior				
Walking time for recreation (min/w)	0.16	0.046	0.18	0.024
Walking time for transport (min/w)	-0.12	0.14	-0.094	0.24
Covariates				
Age (years)	-0.43	<0.001	-0.46	<0.001
Height (cm)			0.035	0.71
Weight (kg)			-0.14	0.18
Hip and knee pain			0.10	0.21
Number of chronic diseases ^c	-0.033	0.68	-0.0037	0.97
Adjusted R ² of the model	0.22		0.22	
Women (n = 240)				
Walking behavior				
Walking time for recreation (min/w)	0.17	<0.01	0.17	<0.01
Walking time for transport (min/w)	-0.040	0.49	-0.040	0.50
Covariates				
Age (years)	-0.25	<0.001	-0.25	<0.001
Height (cm)	0.20	<0.01	0.19	<0.01
Weight (kg)			0.0088	0.89
Hip and knee pain	-0.22	<0.001	-0.22	<0.001
Number of chronic diseases ^c	-0.060	0.30	-0.062	0.30
Adjusted R ² of the model	0.25		0.25	

Note. Sample sizes vary due to missing values.

^a Model 1 is adjusted for covariates those P values were 0.10 and under in univariate correlation analyses with maximum walking speed (ie, age and chronic diseases for men and age, height, pain, and chronic diseases for women) and model 2 is adjusted for all covariates.

^b β is the standardized partial regression coefficient for maximum walking speed.

^c Number of the following diseases: cerebrovascular disorder, diabetes, heart disease, liver disease, lung disease, hypertension.

(ie, 0 min/w; >0 to <90 min/w; 90+ min/w), and body mass index was not significantly associated with MWS when it was analyzed in the models instead of height and weight (data not shown).

Discussion

Both in men and women, walking times for recreation had a significant relationship with MWS, while no significant relationship was observed between walking times for transport and MWS. Significant negative simple correlations of walking times for transport in men and a number of chronic diseases in women with MWS became insignificant in multiple linear regression analyses.

As we hypothesized, one possible interpretation of the observed stronger relationship of walking for recreation with MWS is the greater intensity involved with recreational walking in contrast to walking for transport. We believe that if the elderly walk intentionally to improve and maintain health, the intensity of such walking is greater than of those walks involving routine activities of daily living (eg, walking around home, to work, or for shopping).^{10,22} Our pilot study examining the intensity of walking for recreation and for transport by use of a walking diary and accelerometer in 16 Japanese elderly individuals showed a trend of greater intensity while walking for recreation than that of walking for transport (unpublished data). However, we believe it likely that in cases of elderly people engage in walking for relaxation,²³ the intensity of such walks are low, although this form of walking is deemed to be recreational. Hills et al conducted an experimental study on Australian adults and reported "walking for pleasure" (included in walking for recreation) was sufficient to improve cardiovascular fitness in obese, but not normal-weight, individuals.²⁴ There may be intra- and interpersonal variations in the intensities of walking behavior. Further investigation is needed to determine the intensities of walking for particular purposes.

Another possible cause of the lack of a significant relationship between walking time for transport and MWS is the lower standard deviation and the fewer instances of transport-walking than walking for recreation for this particular study population. In rural areas such as Unnan City, where the automobile is a dominant and necessary form of transportation, the interpersonal difference in walking times for transport is considered to be small. Thus, it is possible we were unable to detect any association between walking for transport and MWS because of insufficient statistical power.

The mean age of our subjects was less than 75 years. Shinkai et al suggested that MWS was most sensitive in predicting future functional dependence for those aged between 65 to 74 years, while usual walking speed was most sensitive for people aged ≥ 75 years.¹² Our observed stronger relationship of walking time for recreation with MWS, compared with walking time for transport, is also believed to be based on the fact that MWS is a relatively vigorous mobility index compared with other indices of

physical fitness (eg, usual walking speed) in the elderly. These results did not contradict the health benefits of active transport (ie, walking or bicycling for transport), especially in younger adults.²⁵

Previous study has shown that time spent in moderate to vigorous intensity physical activity significantly decreased with aging, even though the number of counted steps did not differ significantly.²⁶ Recent recommendations for physical activity in older adults have stated the importance of increased levels of aerobic activity over and above the routine activities of daily living which include light to moderate-intensity activity of less than 10 min duration.¹ Our present study also supports walking for recreation as an effective moderate physical activity for maintaining functional ability in the elderly. Both the promotion of walking for recreation and improvement of public transportation service to promote active transport is needed, especially in rural areas.²⁷

The determinants of walking behavior have been reported to vary with the purposes of walking,^{28,29} so it may be necessary to consider walking for recreation and for transport as separate behaviors, especially in the case of planning health promotion strategies and promoting these behaviors. Differences between the specific association of walking time for recreation and for transport with health benefits require further study.

In addition, the presence of hip or knee pain was associated to low MWS in women. Musculoskeletal conditions have been suggested as determinants of physical activity¹⁸ and functional ability¹⁹ in the elderly. Assistance in the care and treatment of musculoskeletal conditions may be of importance in promoting physical activity and maintaining functional ability in the elderly.

Limitations

This study had several limitations. First, the exclusion of other domains of physical activity,^{2,30} visual and mental health status,³¹ and/or socioeconomic status,³² may cause low model R^2 values (0.22 and 0.25) in multiple linear regression analyses. Assessment of how long the participants had been walking for recreation and for transport would also provide important insight into their effects on functional abilities. Second, our findings may not be attributable generally to populations of elderly people in other geographic locations, given that our subjects were restricted only to Japanese elderly in a rural town. Finally, we were unable to confirm any causality from this cross-sectional study. There is the possibility of reverse causation in the observed associations (eg, participants with higher MWS might be more capable of participating in recreational activities). Future prospective and intervention studies focusing on differences in health outcomes of walking for particular purposes would provide valuable knowledge to health professionals.

In spite of these limitations, the results of this study are notable in that they serve to advance our knowledge of differences between walking for recreation and for transport in association with maintained functional ability, which has not previously received much scrutiny.

Conclusions

Elderly people with longer walking times for recreation had faster MWS. However, walking time for transport was not significantly associated with MWS. Walking for recreation in addition to that for routine activities of daily living may contribute to the maintaining of functional ability in the elderly. Considering the several limitations of this study, further research is needed to confirm the differences in the health benefits associated with walking for recreation and those for transport.

Acknowledgments

We greatly appreciate the cooperation of the participants and staff members of the Shimane study. We also appreciate the generous assistance provided by Mr. Roger Matthews, Mr. Nicholas Zimmermann, and Ms. Atsuko Sato. This study was supported by Health and Labor Sciences Research Grants from the Japanese Ministry of Health, Labor and Welfare.

References

- Nelson ME, Rejeski WJ, Blair SN, et al. Physical activity and public health in older adults. Recommendation from the American College of Sports Medicine and the American Heart Association. *Circulation*. 2007;116:1094–1105.
- Brach JS, FitzGerald S, Newman AB, et al. Physical activity and functional status in community-dwelling older women: a 14-year prospective study. *Arch Intern Med*. 2003;163:2565–2571.
- DiPietro L. The epidemiology of physical activity and physical function in older people. *Med Sci Sports Exerc*. 1996;28:596–600.
- LaCroix AZ, Guralnik JM, Berkman LF, Wallace RB, Satterfield S. Maintaining mobility in late life. II. Smoking, alcohol consumption, physical activity, and body mass index. *Am J Epidemiol*. 1993;137:858–869.
- Sasakawa Sports Foundation. *Sports White Paper*. Tokyo, Sasakawa Sports Foundation, 2006. [in Japanese]
- Siegel PZ, Brackbill RM, Heath GW. The epidemiology of walking for exercise: implications for promoting activity among sedentary groups. *Am J Public Health*. 1995;85:706–710.
- Australian Bureau of Statistics. *Participation in sport and physical activities, Australia*. 2000. Available at: <http://www.abs.gov.au/AUSSTATS/abs@.nsf/DetailsPage/4177.01999-2000?OpenDocument>, accessed June 2009.
- Hayashi T, Tsumura K, Suematsu C, Okada K, Fujii S, Endo G. Walking to work and the risk for hypertension in men: the Osaka Health Survey. *Ann Intern Med*. 1999;131:21–26.
- Cummings SR, Nevitt MC, Browner WS, et al. Risk factors for hip fracture in white women. Study of Osteoporotic Fractures Research Group. *N Engl J Med*. 1995;332:767–773.
- Parise C, Sternfeld B, Samuels S, Tager IB. Brisk walking speed in older adults who walk for exercise. *J Am Geriatr Soc*. 2004;52:411–416.
- Sun W, Watanabe M, Tanimoto Y, et al. Assessment of the best gait parameter in relation to bone status in community-dwelling young-old and old-old women in Japan. *Arch Gerontol Geriatr*. 2009;49:156–161.
- Shinkai S, Watanabe S, Kumagai S, et al. Walking speed as a good predictor for the onset of functional dependence in a Japanese rural community population. *Age Ageing*. 2000;29:441–446.
- Bohannon RW. Comfortable and maximum walking speed of adults aged 20–79 years: reference values and determinants. *Age Ageing*. 1997;26:15–19.
- Wang T, Karino K, Yamasaki M, et al. Effects of G994T in the Lp-PLA2 gene on the plasma oxidized LDL level and carotid intima-media thickness in Japanese: the Shimane study. *Am J Hypertens*. 2009;22:742–747.
- Kumahara H, Schutz Y, Ayabe M, et al. The use of uniaxial accelerometry for the assessment of physical-activity-related energy expenditure: a validation study against whole-body indirect calorimetry. *Br J Nutr*. 2004;91:235–243.
- Rafamantanantsoa HH, Ebine N, Yoshioka M, et al. Validation of three alternative methods to measure total energy expenditure against the doubly labeled water method for older Japanese men. *J Nutr Sci Vitaminol (Tokyo)*. 2002;48:517–523.
- Physical Education and Medicine Research Center UNNAN. Shimane study questionnaire. Available at <http://user.kkm.ne.jp/shintai/kankoubutu/kankoubutu-index.html>, accessed January 2010. [in Japanese]
- Centers for Disease Control and Prevention. Arthritis as a potential barrier to physical activity among adults with heart disease—United States, 2005 and 2007. *MMWR Morb Mortal Wkly Rep*. 27 2009;58:165–169.
- Weaver GD, Kuo YF, Raji MA, et al. Pain and disability in older Mexican-American adults. *J Am Geriatr Soc*. 2009;57:992–999.
- Goldman SE, Stone KL, Ancoli-Israel S, et al. Poor sleep is associated with poorer physical performance and greater functional limitations in older women. *Sleep*. 2007;30:1317–1324.
- Cabinet Office, Government of Japan. *Public opinion poll on physical fitness and sports*. 2004. Available at: <http://www8.cao.go.jp/survey/h15/h15-sports/index.html>, accessed July 2009. [in Japanese]
- Murtagh EM, Boreham CA, Murphy MH. Speed and exercise intensity of recreational walkers. *Prev Med*. 2002;35:397–400.
- Clifton KJ, Livi AD. Gender differences in walking behavior, attitudes about walking, and perceptions of the environment in three Maryland communities. In: Transportation Research Board, National Research Council, United States, ed. *Research on women's issues in transportation, report of a conference*. Vol 2. 2005:79–88.
- Hills AP, Byrne NM, Wearing S, Armstrong T. Validation of the intensity of walking for pleasure in obese adults. *Prev Med*. 2006;42:47–50.
- Wen LM, Orr N, Millett C, Rissel C. Driving to work and overweight and obesity: findings from the 2003 New South Wales Health Survey, Australia. *Int J Obes (Lond)*. 2006;30(5):782–786.
- Ayabe M, Yahiro T, Yoshioka M, Higuchi H, Higaki Y, Tanaka H. Objectively measured age-related changes in the intensity distribution of daily physical activity in adults. *J Phys Act Health*. 2009;6:419–425.

27. Kamada M, Kitayuguchi J, Inoue S, Kamioka H, Mutoh Y, Shiwaku K. Environmental correlates of physical activity in driving and non-driving rural Japanese women. *Prev Med.* 2009;49:490-496.
28. Owen N, Humpel N, Leslie E, Bauman A, Sallis JF. Understanding environmental influences on walking: review and research agenda. *Am J Prev Med.* 2004;27:67-76.
29. Giles-Corti B, Donovan RJ. Socioeconomic status differences in recreational physical activity levels and real and perceived access to a supportive physical environment. *Prev Med.* 2002;35:601-611.
30. Park H, Kim KJ, Komatsu T, Park SK, Mutoh Y. Effect of combined exercise training on bone, body balance, and gait ability: a randomized controlled study in community-dwelling elderly women. *J Bone Miner Metab.* 2008;26:254-259.
31. Alexander NB, Goldberg A. Gait disorders: search for multiple causes. *Cleve Clin J Med.* 2005;72(7):586,589-590,592-584.
32. Thorpe RJ, Jr, Kasper JD, Szanton SL, Frick KD, Fried LP, Simonsick EM. Relationship of race and poverty to lower extremity function and decline: findings from the Women's Health and Aging Study. *Soc Sci Med.* 2008;66:811-821.

Article

Perceived Environmental Factors Associated with Physical Activity among Normal-weight and Overweight Japanese Men

Yung Liao ^{1,*}, Kazuhiro Harada ^{1,2}, Ai Shibata ³, Kaori Ishii ³, Koichiro Oka ³,
Yoshio Nakamura ³, Shigeru Inoue ⁴ and Teruichi Shimomitsu ⁴

¹ Graduate School of Sport Sciences, Waseda University, Nakamura Lab. 2-579-15 Mikajima Tokorozawa, Saitama 359-1192, Japan; E-Mail: harada-ka@fuji.waseda.jp (K.H.)

² Japan Society for the Promotion of Science, Sumitomo-Ichibancho Bldg., 6 Ichibancho, Chiyoda-ku, Tokyo 102-8471, Japan

³ Faculty of Sport Sciences, Waseda University, 2-579-15 Mikajima Tokorozawa, Saitama 359-1192, Japan; E-Mails: aishibata@aoni.waseda.jp (A.S.); ishiikaori@aoni.waseda.jp (K.I.); koka@waseda.jp (K.O.); nakamura@waseda.jp (Y.N.)

⁴ Department of Preventive Medicine and Public Health, Tokyo Medical University, 6-1-1, Shinjuku-ku, Tokyo 160-8402, Japan; E-Mails: inoue@tokyo-med.ac.jp (S.I.); tshimo@tokyo-med.ac.jp (T.S.)

* Author to whom correspondence should be addressed; E-Mail: anthroliao@fuji.waseda.jp; Tel.: +81-4-2947-7189; Fax: +81-4-2947-7189.

Received: 26 January 2011; in revised form: 3 March 2011 / Accepted: 22 March 2011 /

Published: 28 March 2011

Abstract: Although it is crucial to examine the environmental correlates of physical activity (PA) for developing more effective interventions for overweight populations, limited studies have investigated differences in the environmental correlates on body mass index (BMI). The purpose of the present study was to examine the perceived environmental correlates of PA among normal-weight and overweight Japanese men. Data were analyzed for 1,420 men (aged 44.4 ± 8.3 years), who responded to an internet-based cross-sectional survey of answering the short version of the International Physical Activity Questionnaire and its Environment Module. Binary logistic regression analyses were utilized to examine the environmental factors associated with meeting the PA recommendation (150 minutes/week) between the normal-weight and overweight men. After adjusting for socio-demographic variables, common and different environmental