

3. 睡眠呼吸障害と合併症—特に糖尿病との関連での最近の話題—

- tive sleep apnea and incident coronary heart disease and heart failure : The Sleep Heart Health Study. *Circulation* 2010 ; 122 : 352-360.
- 6) Yaggi HK et al : Obstructive sleep apnea as a risk factor for stroke and death. *N Engl J Med* 2005 ; 353 : 2034-2041.
- 7) Redline S et al : Obstructive sleep apnea-hypopnea and incident stroke : The Sleep Heart Health Study. *Am J Respir Crit Care Med* 2010 ; 182 : 269-277.
- 8) Meslier N et al : Impaired glucose-insulin metabolism in males with obstructive sleep apnea syndrome. *Eur Respir J* 2003 ; 22 : 156-160.
- 9) Resnick HE et al : Diabetes and sleep disturbances : findings from the Sleep Heart Health Study. *Diabetes Care* 2003 ; 26 : 702-709.
- 10) Shaw JE et al : Sleep-disordered breathing and type 2 diabetes : A report from the International Diabetes Federation Taskforce on Epidemiology and Prevention. *Diabetes Res Clin Pract* 2008 ; 81 : 2-12.
- 11) Spiegel K et al : Impact of sleep debt on metabolic and endocrine function. *Lancet* 1999 ; 354 : 1435-1439.
- 12) Punjabi NM and Beamer BA : Alterations in glucose disposal in sleep-disordered breathing. *Am J Respir Crit Care Med* 2009 ; 179 : 235-240.
- 13) Tasali E et al : Slow-wave sleep and the risk of type 2 diabetes in humans. *PNAS* 2008 ; 105 : 1044-1049.
- 14) Stamatakis KA and Punjabi NM : Effects of sleep fragmentation on glucose metabolism in normal subjects. *Chest* 2010 ; 137 : 95-101.
- 15) Louis M and Punjabi NM : Effects of acute intermittent hypoxia on glucose metabolism in awake healthy volunteers. *J Appl Physiol* 2009 ; 106 : 1538-1544.
- 16) Punjabi NM et al : Sleep-disordered breathing, glucose intolerance, and insulin resistance : The Sleep Heart Health Study. *Am J Epidemiol* 2004 ; 160 : 521-530.
- 17) Seicean S et al : Sleep-disordered breathing and impaired glucose metabolism in normal-weight and overweight/obese individuals : the Sleep Heart Health Study. *Diabetes Care* 2008 ; 31 : 1001-1006.
- 18) Reichmuth KJ et al : Association of sleep apnea and type II diabetes : a population-based study. *Am J Respir Care Med* 2005 ; 172 : 1590-1595.
- 19) Botros N et al : Obstructive sleep apnea as a risk factor for type 2 diabetes. *Am J Med* 2009 ; 122 : 1122-1127.
- 20) Marshall NS et al : Is sleep apnea an independent risk factor for prevalent and incident diabetes in the Busselton Health Study? *J Clin Sleep Med* 2009 ; 5 : 15-20.
- 21) Shin C et al : Association of habitual snoring with glucose and insulin metabolism in nonobese Korean adult men. *Am J Respir Crit Care Med* 2005 ; 171 : 287-291.
- 22) Muraki I et al : Nocturnal intermittent hypoxia and the development of type 2 diabetes : the Circulatory Risk in Communities Study(CIRCS). *Diabetologia* 2010 ; 53 : 481-488.

