



Traditional risk factor management for stroke: a never-ending challenge for health behaviors of diet and physical activity

Yoshihiro Kokubo

Purpose of review

Recently, many guidelines have given new evidence on the risk factors for stroke. In this review, I refer to the most important guidelines for primary prevention of stroke and hypertension, especially focused on diet and physical activity.

Recent findings

The health behavior recommendations in recent guidelines for the primary prevention of stroke are virtually identical, and the same recommendations appear in the recent guidelines for the management of hypertension, especially with respect to diet and physical activity. The recommended health behaviors consist of weight reduction, reduction of salt intake, increase in fruit and vegetable intake, decrease in saturated and total fat intake (increase in fish intake), physical activity, and moderation of alcohol consumption. Fruits and vegetables have high levels of potassium, antioxidants, phytochemicals, and dietary fiber, and thus are also considered preventive of cardiovascular disease and its risk factors. It was found that individuals with many of these health behaviors have been shown to have a lowered risk of stroke.

Summary

The health behaviors, especially those related to diet and physical activity, appearing in recent guidelines for the management of hypertension are also important for the primary prevention of stroke, and appear in recent stroke guidelines.

Keywords

diet, guidelines, health behavior, physical activity, primary prevention of stroke

INTRODUCTION

Stroke is one of the leading causes of mortality and morbidity worldwide [1]. Figure 1 shows the schema of the progression from lifestyle behaviors to the onset of cardiovascular disease (CVD) as follows:

Environmental factor (lifestyle) → risk factors → cardiovascular disease

Environmental and genetic factors are the key factors for the primary stage of preventive CVD. Environmental factors include diet, physical activity, smoking, drinking, mental condition, and socioeconomic factors. Cardiovascular risk factors, such as hypertension, diabetes, and dyslipidemia, are the key factors for the secondary stage of preventive CVD. It is essential to consider these factors in order to prevent CVD, irrespective of whether lifestyle behaviors and risk factors were improved in the early stage.

Hypertension is the strongest risk factor for CVD worldwide [2,3]. The total population-attributable fractions of high-normal blood pressure and hypertension for CVD were approximately 50% in men and 30% in women [4]. When high blood pressure levels and other risk factors, such as chronic kidney disease [5] or diabetes mellitus [6^a], are combined, the risk of cardiovascular disease becomes much

Department of Preventive Cardiology, National Cerebral and Cardiovascular Center, Suita, Japan

Correspondence to Yoshihiro Kokubo, MD, PhD, FAHA, FACC, FESC, Department of Preventive Cardiology, National Cerebral and Cardiovascular Center, 5-7-1, Fujishiro-dai, Suita, Osaka 565-8565, Japan.
Tel: +81 6 6833 5012 x2186; fax: +81 6 6833 5300; e-mail: ykokubo@hsp.ncvc.go.jp

Curr Opin Neurol 2012, 25:11–17

DOI:10.1097/WCO.0b013e32834eb58e

KEY POINTS

- Reduced intake of sodium and increased intake of potassium.
- Higher consumption of fruits, vegetables, low-fat dairy products and foods reduced in saturated fat (rich in fish and n-3 polyunsaturated fat).
- Increasing physical activity.
- Lifestyle recommendations for primary prevention of stroke are similar to those for preventing hypertension.

higher. The prevention of hypertension is the best way to prevent primary strokes. According to the 2007 European Society of Cardiology/European Society of Hypertension (ESC/ESH) Guidelines for the management of arterial hypertension [7], the Seventh Report of the Joint National Committee (JNC 7) [8], and the Japanese Society of Hypertension Guidelines for the Management of Hypertension (JSH 2009) [9], the factors that are effective for the prevention of hypertension are weight reduction, reduction of salt intake, increase in fruit and vegetable intake, decrease in saturated and total fat intake, physical activity, and moderation of alcohol consumption (Table 1). These factors are also considered as preventive for stroke (Table 2).

The Guidelines for the Primary Prevention of Stroke from the American Heart Association/American Stroke Association state that several aspects of diet can contribute to elevated blood pressure: excess salt intake, weight, drinking, and low potassium intake [10¹¹]. These guidelines also recommend a DASH (Dietary Approaches to Stop Hypertension)-style diet, which emphasizes consumption of fruits, vegetables, and low-fat dairy products. The European Stroke Organization (ESO) guidelines similarly recommend

the following measures in order to reduce the risk of stroke: moderate drinking, regular physical activity, maintenance of appropriate weight, and a diet low in salt and saturated fat and rich in fruits, vegetables, and fiber [11]. However, antioxidant vitamin supplements and hormone replacement therapy are not recommended for the primary prevention of stroke in the ESO guidelines [11].

This review will focus on the prevention of stroke events through dietary modifications and increased physical activity, which are never-ending challenges for health behaviors of diet and physical activity.

DIET

Diet is one of the important lifestyle factors for the prevention of stroke, because we habitually eat three times a day. As mentioned in the Introduction, the following dietary factors have been related to stroke prevention: reduction of salt intake, increase in fruit and vegetable intake, and decrease in saturated and total fat intake.

Reduction of salt intake

In the INTERSALT (International Study of Salt and Blood Pressure) study, it is estimated that with a 100 mmol lower daily sodium intake the average decrease in blood pressure from age 25 to 55 would be by 9.0 mmHg for systolic and 4.5 mmHg for diastolic [12]. In Japan, the incidence of stroke is higher than in other countries, partly because both the salt intake and the frequency of hypertension are higher in Japan than in other countries. In a large Japanese general population, sodium intake was increasing associated with mortality from stroke [hazard ratio 1.6, 95% confidence interval (CI) 1.2–2.0], ischemic stroke (hazard ratio 2.0, 95% CI 1.4–2.9), and CVD (hazard ratio 1.4, 95% CI 1.2–1.7) [13].




FIGURE 1. Schema of the progression from lifestyle changes to the incidence of cardiovascular disease. CHD, coronary heart disease; CVD, cardiovascular disease.

Table 1. Lifestyle recommendations to prevent hypertension in various guidelines

	JNC 7 [8]	ESC/ESH [7]	JSH 2009 [9]
Weight reduction	Maintain normal body weight	Weight reduction (and weight stabilization)	Maintaining appropriate body weight: BMI <25kg/m ²
Adopt DASH eating plan	Consume a diet rich in fruits, vegetables, and low-fat dairy products with a reduced content of saturated and total fat	Increase in fruit and vegetable intake and decrease in saturated and total fat intake	Increased intake of fruits and vegetables. Reduce intake of cholesterol and saturated fatty acids. Increased intake of fish (fish oil)
Dietary sodium reduction	Reduce dietary sodium intake to no more than 100 mmol per day	Reduction of salt intake	Salt restriction to <6 g/day
Physical activity	Engage in regular aerobic physical activity such as brisk walking (at least 30 min per day)	Physical exercise	Exercise: In hypertensive patients with no cardiovascular disease, exercise, which is primarily moderate aerobic exercise, should be performed periodically (for 30 min daily if possible)
Moderation of alcohol consumption	Limit consumption to no more than 2 drinks per day in men and to no more than 1 drink per day in women and lighter weight men	Reduction of excessive alcohol intake	Restriction of alcohol intake: 20–30 ml/day in men and 10–20 ml/day in women as ethanol
Stop smoking	For overall cardiovascular risk reduction, stop smoking	Smoking cessation	Quitting smoking

Table 2. Lifestyle recommendations to prevent primary stroke in various guidelines

	Guidelines for the primary prevention of stroke: AHA/ASA Guideline [10 ^{aaa}]	The European Stroke Initiative recommendations for stroke management [11]
Weight reduction	Among overweight and obese persons, weight reduction is reasonable as a means of reducing risk of stroke	Individuals with an elevated body mass index are recommended to take a weight-reducing diet
Diet and nutrition	Reduced intake of sodium and increased intake of potassium as indicated in the report Dietary Guidelines for Americans are recommended to lower blood pressure A DASH-style diet, which emphasizes consumption of fruits, vegetables, and low-fat dairy products and is reduced in saturated fat, also lowers blood pressure and is recommended A diet that is rich in fruits and vegetables and thereby high in potassium is beneficial and may lower risk of stroke	A diet low in salt is recommended A diet low in saturated fat, high in fruit and vegetables and rich in fibre is recommended
Physical activity	Increased physical activity is recommended because it is associated with a reduction in risk of stroke Adults should engage in at least 150 min (2 h and 30 min) per week of moderate intensity or 75 min (1 h and 15 min) per week of vigorous intensity aerobic physical activity	Regular physical activity is recommended

DASH, Dietary Approaches to Stop Hypertension.

Increase in fruit and vegetable intake

Cohort studies have shown that high fruit and vegetable intake reduces stroke incidence and mortality. Two meta-analysis studies reported that fruit and vegetable consumption decreases the risk of stroke [14,15]. The risk reductions of stroke from meta-analysis of seven cohorts were 11% (95% CI 7–15%) for each additional portion per day of fruit and 5% (3–8%) for fruit and vegetables [14]. Those of nine cohorts were 26% (21–31%) for more than five servings per day of fruit and vegetables and 11% (3–17%) for three to five servings per day compared to less than three servings per day [15]. In participants in a large Japanese cohort study, the inverse association between fruit intake and risk was observed only among nonsmokers [16]. Smoking, which causes oxidative stress, diminishes the healthy effects of fruits and vegetables. The mechanism by which high intake of fruits and vegetables prevents stroke may involve the high levels of potassium, antioxidants, phytochemicals, and dietary fiber in these foods, since each of these components has been shown to individually prevent cardiovascular risk factors.

Potassium excretion has been negatively correlated with blood pressure in individuals. A 50 mmol/day lower urinary excretion of potassium has been associated with an average blood pressure increase of 3.4 mmHg systolic or 1.9 mmHg diastolic [12]. A meta-analysis of 33 randomized controlled trials of potassium supplementation showed that the supplementation significantly reduced systolic and diastolic blood pressure. The systolic and diastolic blood pressures were decreased by 4.0 and 2.5 mmHg, respectively, in hypertensive patients, whereas they were decreased by 1.8 and 1.0 mmHg, respectively, in normotensive individuals.

In the US general population, higher potassium intake was associated with lower mortality risk (hazard ratio 0.80, 95% CI 0.67–0.94, per 1000 mg/day of potassium) [17^a]. In a pooled meta-analysis of 15 cohorts including 247 510 participants at baseline for 5–19 years of follow-up, a 42 mmol/day higher potassium intake was associated with a 21% lower risk of stroke (hazard ratio 0.79, 95% CI 0.68–0.90) [18^a].

Fruits and vegetables are rich sources of antioxidants that may help prevent CVD, including vitamins C and E, carotenoids, polyphenols, and flavonoids. Cohort participants in the top quartiles of baseline plasma vitamin C concentrations had 41–61% lower risk than did those in the bottom quartiles [19–21]. In a Rotterdam cohort study, compared with participants in the lowest tertile of vitamin C intake, those in the second and third tertiles of vitamin C intake had a significantly lower

risk of ischemic stroke (hazard ratios 0.69 and 0.66, 95% CIs 0.49–0.98 and 0.46–0.93, respectively) [22]. In another study, a high serum vitamin A concentration was found to have a beneficial effect on early outcome in ischemic stroke [23]. Dietary supplementation with antioxidants including vitamins C, E, and beta-carotene was tested in a meta-analysis of randomized controlled trials, and no significant association with secondary CVD prevention was observed [24].

Flavonoids are a large group of polyphenolic compounds abundant in vegetables, fruits, tea, and red wine, and may contribute to the protective effect of these foods. There are five subclasses of flavonoids, that is, flavonols, flavones, flavanones, flavan-3-ols and anthocyanidins, which have been estimated to contribute to daily dietary intake and thus potentially have effects on health [25]. In a cohort study, Finnish men in the highest quartile of flavonol and flavan-3-ol intake had a relative risk of 0.55 (95% CI 0.31–0.99) and 0.59 (0.30–1.14) for ischemic stroke, respectively, as compared with the lowest quartile [26]. In a Japanese cohort study, green tea consumption of five or more cups per day was associated with a relative risk of 0.88 (95% CI 0.79–0.98) for ischemic stroke compared with less than one cup per day [27]. Recent cohort studies have shown that coffee consumption is protective of stroke. The relative risks of total stroke in Swedish women were 1.00 (reference), 0.78 (95% CI 0.66–0.91), 0.75 (95% CI 0.64–0.88), and 0.77 (95% CI 0.63–0.92) for coffee consumption of less than 1 cup/day, 1–2 cups/day, 3–4 cups/day, and at least 5 cups/day, respectively [28].

Many studies on the intake of soy, a food rich in isoflavones, have reported an impact on decreasing plasma cholesterol levels [29,30]. However, only a few studies exist on the impact of soy intake on CVD, because the average soy intake in Japanese is 10–70 times higher than that in Western people [31,32]. A prospective study of Dutch women did not support the idea that dietary isoflavones lowered the risk of CVD [32]. The quantity of isoflavones consumed by Dutch women, however, was quite small [32]. A prospective study of Japanese men and women showed that soy intake was weakly and inversely associated with total mortality but not mortality due to CVD [33]. Recently, a Japanese community-based prospective study with 40 462 participants has shown that high consumption of soy and isoflavones was associated with a reduced risk of incidence and mortality of cerebral and myocardial infarction among women, particularly postmenopausal women [34]. This study suggests that the consumption of dietary isoflavones and soy intake may be beneficial to

postmenopausal women for the prevention of ischemic CVD.

Population cohort studies have revealed dietary fiber, consisting of water-soluble and water-insoluble fiber, to be inversely associated with the risks of coronary heart disease (CHD) [35,36²⁰] and of stroke [37]. Water-soluble fiber may result in an improvement of glycemic control and a lowering of triglyceride levels [38], as well as a cholesterol-lowering effect [39], especially on low-density lipoprotein cholesterol. Insoluble fiber may slow the intestinal absorption of foods and reduce the levels of clotting factors [40], fibrinolysis [41], coagulation [42], and inflammatory markers [43]. Recently, a relatively large Japanese population-based prospective study has shown higher dietary intakes of total and insoluble fiber to be associated with reduced risk of total strokes, cerebral infarction and intracerebral hemorrhage in women [44²¹]. The inverse associations between dietary total fiber intake and cardiovascular diseases were statistically significant only for non-smoking men and women, but not for smoking men and women.

Fish and n-3 polyunsaturated fat (decrease in saturated and total fat intake)

In two large Japanese community samples, dietary intake of fish and n-3 polyunsaturated fatty acid was inversely associated with incidence of CHD [45] and mortality of CVD [46]. However, the association between intake of fat and stroke is controversial. A meta-analysis of randomized controlled trials showed no effects of n-3 fatty acids on cardiovascular events [47]. A large cohort study in men in the US did not show an increase in the risk of ischemic stroke according to dietary intakes of total fat, animal fat, saturated fat, and vegetable fat [48]. The estimated level of association between dairy intake and stroke events pooled from seven prospective studies was not statistically significant [49²²]. In a Japanese cohort, a high consumption of animal fat and cholesterol was associated with a reduced risk of cerebral infarction death, whereas those relationships have been diminished in Western counties [50], where the animal product intake is higher than in Japan.

Interestingly, an increased concentration in serum of n-3 polyunsaturated fatty acids, especially docosahexaenoic acid, a marker of fish or fish oil consumption, may protect against atrial fibrillation according to a prospective population-based study [51,52]. Atrial fibrillation is a strong risk factor for cerebral infarction [53]. Therefore, an increased concentration of n-3 polyunsaturated fatty acids may be of benefit for the prevention

of cerebral infarction, especially cardio-embolism infarction.

PHYSICAL ACTIVITY

The 2008 Physical Activity Guidelines for Americans include an extensive review of the literature and concludes that individuals engaged in physical activity have a 25–30% lower risk of stroke than inactive individuals [54²³,55]. In another study, moderate activity was associated with a lower risk of stroke compared with inactivity (hazard ratio 0.80, 95% CI 0.74–0.86) [56]. This association was mediated through beneficial effects on body weight, blood pressure, serum total cholesterol levels, and glucose levels. The 2008 Physical Activity Guidelines for Americans recommended that adults should engage in at least 150 min per week of moderate-intensity or 75 min per week of vigorous-intensity aerobic physical activity, or an equivalent combination of moderate and vigorous-intensity aerobic activity [10²⁴,54²³].

According to the Physical Activity Guideline for Older Adults [57²⁵], to promote and maintain health, older adults should participate in moderate-intensity aerobic activity for at least 30 min on 5 days of the week, or vigorous-intensity aerobic activity for at least 20 min on 3 days of the week [58]. Even in older adults with chronic illnesses or disabilities, significant health benefits can be obtained by daily nonstrenuous physical activity, such as stretching exercises [55].

Physical activity in leisure time (2–5 h per week) has been significantly associated with a reduced severity of ischemic stroke at admission [59]. Japanese individuals who reported the highest level of physical activity (i.e. walking 1 h/day) had lower mortality of stroke than did those in the second lowest physical activity category (i.e. walking 0.5 h/day) (hazard ratio 0.71, 95% CI 0.54–0.94). [60]

COMBINED IMPACT OF HEALTH BEHAVIORS

In the above mentioned guidelines for the primary prevention of stroke, there is a study on health behaviors for the prevention of stroke, with a focus on diet and physical activity [61]. This prospective population study (20 040 men and women aged 40–79 at baseline) reported on the potential combined impact of four health behaviors, that is current nonsmoking, physical inactivity, moderate alcohol intake, and plasma concentration of vitamin C at least 50 $\mu\text{mol/l}$ (indicating fruit and vegetable intake of at least five servings a day), on incidence of stroke. The relative risks for stroke were 1.15 (95% CI 0.89–

1.49) for three health behaviors, 1.58 (1.22–2.05) for two, 2.18 (1.63–2.92) for one, and 2.31 (1.33–4.02) for none ($P < 0.001$ for trend), compared with people with all four health behaviors.

CONCLUSION

The health behavior recommendations in guidelines for the primary prevention of stroke are highly similar, and the same recommendations appear in the guidelines for the management of hypertension, especially with respect to diet and physical activity. The recommended health behaviors consist of weight reduction, reduction of salt intake, increase in fruit and vegetable intake, decrease in saturated and total fat intake, physical activity, and moderation of alcohol consumption. Individuals with many of these health behaviors were found to be at lower risk of stroke.

Acknowledgements

The present study was supported by the Intramural Research Fund of the National Cerebral and Cardiovascular Center (22-4-5) and by a grant (no. 23390178) from the Ministry of Education, Science, and Culture of Japan.

Conflicts of interest

There are no conflicts of interest.

REFERENCES AND RECOMMENDED READING

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
- of outstanding interest

Additional references related to this topic can also be found in the Current World Literature section in this issue (pp. 96–97).

1. Feigin VL, Lawes CM, Bennett DA, Anderson CS. Stroke epidemiology: a review of population-based studies of incidence, prevalence, and case-fatality in the late 20th century. *Lancet Neurol* 2003; 2:43–53.
2. Kubo M, Hata J, Doi Y, *et al.* Secular trends in the incidence of and risk factors for ischemic stroke and its subtypes in Japanese population. *Circulation* 2008; 118:2672–2678.
3. Sjol A, Thomsen KK, Schroll M. Secular trends in blood pressure levels in Denmark 1964–1991. *Int J Epidemiol* 1998; 27:614–622.
4. Kokubo Y, Kamide K, Okamura T, *et al.* Impact of high-normal blood pressure on the risk of cardiovascular disease in a Japanese urban cohort: the Suita study. *Hypertension* 2008; 52:652–659.
5. Kokubo Y, Nakamura S, Okamura T, *et al.* The relationship between blood pressure category and incidence of stroke and myocardial infarction in an urban Japanese population with and without chronic kidney disease: the Suita study. *Stroke* 2009; 40:2674–2679.
6. Kokubo Y, Okamura T, Watanabe M, *et al.* The combined impact of blood pressure category and glucose abnormality on the incidence of cardiovascular diseases in a Japanese urban cohort: the Suita study. *Hypertens Res* 2010; 33:1238–1243.
- This cohort study showed that the high-normal blood pressure patients in all glucose categories and the normal blood pressure patients with impaired fasting glucose showed increased risk of cardiovascular disease in this Japanese population.
7. Mancia G, De Backer G, Dominiczak A, *et al.* Guidelines for the management of arterial hypertension: the task force for the management of arterial hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). *J Hypertens* 2007; 25:1105–1187.
8. Chobanian AV, Bakris GL, Black HR, *et al.* Seventh report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure. *Hypertension* 2003; 42:1206–1252.
9. Ogihara T, Kikuchi K, Matsuoka H, *et al.* The Japanese Society of Hypertension Guidelines for the Management of Hypertension (JSH 2009). *Hypertens Res* 2009; 32:3–107.
10. Goldstein LB, Bushnell CD, Adams RJ, *et al.* Guidelines for the primary prevention of stroke: a guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke* 2011; 42:517–584.
- This guideline provides an overview of the evidence on established and emerging risk factors for stroke to provide evidence-based recommendations for the reduction of risk of a first stroke. Extensive evidence identifies a variety of specific factors that increase the risk of a first stroke and that provide strategies for reducing that risk.
11. Guidelines for management of ischaemic stroke and transient ischaemic attack 2008. *Cerebrovasc Dis* 2008; 25:457–507.
12. Intersalt: an international study of electrolyte excretion and blood pressure. Results for 24 h urinary sodium and potassium excretion. Intersalt Cooperative Research Group. *Br Med J* 1988; 297:319–328.
13. Umesawa M, Iso H, Date C, *et al.* Relations between dietary sodium and potassium intakes and mortality from cardiovascular disease: the Japan Collaborative Cohort Study for Evaluation of Cancer Risks. *Am J Clin Nutr* 2008; 88:195–202.
14. He FJ, Nowson CA, MacGregor GA. Fruit and vegetable consumption and stroke: meta-analysis of cohort studies. *Lancet* 2006; 367:320–326.
15. Dauchet L, Amouyel P, Dallongeville J. Fruit and vegetable consumption and risk of stroke: a meta-analysis of cohort studies. *Neurology* 2005; 65:1193–1197.
16. Takachi R, Inoue M, Ishihara J, *et al.* Fruit and vegetable intake and risk of total cancer and cardiovascular disease: Japan Public Health Center-Based Prospective Study. *Am J Epidemiol* 2008; 167:59–70.
17. Yang Q, Liu T, Kuklina EV, *et al.* Sodium and potassium intake and mortality among US adults: prospective data from the Third National Health and Nutrition Examination Survey. *Arch Intern Med* 2011; 171:1183–1191.
- The Third National Health and Nutrition Examination Survey Linked Mortality File (1988–2006), a prospective cohort study showed that a higher sodium–potassium ratio is associated with significantly increased risk of cardiovascular disease and all-cause mortality, and higher sodium intake is associated with increased total mortality in the general US population.
18. D'Elia L, Barba G, Cappuccio FP, Strazzullo P. Potassium intake, stroke, and cardiovascular disease: a meta-analysis of prospective studies. *J Am Coll Cardiol* 2011; 57:1210–1219.
- A meta-analysis of prospective studies showed that higher dietary potassium intake is associated with lower rates of stroke and might also reduce the risk of coronary heart disease and cardiovascular disease.
19. Myint PK, Luben RN, Welch AA, *et al.* Plasma vitamin C concentrations predict risk of incident stroke over 10 y in 20 649 participants of the European Prospective Investigation into Cancer Norfolk prospective population study. *Am J Clin Nutr* 2008; 87:64–69.
20. Kurl S, Tuomainen TP, Laukkanen JA, *et al.* Plasma vitamin C modifies the association between hypertension and risk of stroke. *Stroke* 2002; 33:1568–1573.
21. Yokoyama T, Date C, Kokubo Y, *et al.* Serum vitamin C concentration was inversely associated with subsequent 20-year incidence of stroke in a Japanese rural community. The Shibata study. *Stroke* 2000; 31:2287–2294.
22. Voko Z, Hollander M, Hofman A, *et al.* Dietary antioxidants and the risk of ischemic stroke: the Rotterdam Study. *Neurology* 2003; 61:1273–1275.
23. De Keyser J, De Klippel N, Merckx H, *et al.* Serum concentrations of vitamins A and E and early outcome after ischaemic stroke. *Lancet* 1992; 339:1562–1565.
24. Bleyers J, Miller ER 3rd, Pastor-Barriuso R, *et al.* Vitamin-mineral supplementation and the progression of atherosclerosis: a meta-analysis of randomized controlled trials. *Am J Clin Nutr* 2006; 84:880–887 (quiz 954–955).
25. Dwyer JT, Peterson JJ. Measuring flavonoid intake: need for advanced tools. *Public Health Nutr* 2002; 5:925–930.
26. Mursu J, Voutilainen S, Nurmi T, *et al.* Flavonoid intake and the risk of ischaemic stroke and CVD mortality in middle-aged Finnish men: the Kuopio Ischaemic Heart Disease Risk Factor Study. *Br J Nutr* 2008; 100:890–895.
27. Kuriyama S, Shimazu T, Ohmori K, *et al.* Green tea consumption and mortality due to cardiovascular disease, cancer, and all causes in Japan: the Ohsaki study. *J Am Med Assoc* 2006; 296:1255–1265.
28. Larsson SC, Virtamo J, Wolk A. Coffee consumption and risk of stroke in women. *Stroke* 2011; 42:908–912.
29. Anderson JW, Johnstone BM, Cook-Newell ME. Meta-analysis of the effects of soy protein intake on serum lipids. *N Engl J Med* 1995; 333:276–282.
30. Zhan S, Ho SC. Meta-analysis of the effects of soy protein containing isoflavones on the lipid profile. *Am J Clin Nutr* 2005; 81:397–408.
31. Erdman JW Jr. AHA Science Advisory: soy protein and cardiovascular disease: a statement for healthcare professionals from the Nutrition Committee of the AHA. *Circulation* 2000; 102:2555–2559.
32. van der Schouw YT, Kreijkamp-Kaspers S, Peeters PH, *et al.* Prospective study on usual dietary phytoestrogen intake and cardiovascular disease risk in Western women. *Circulation* 2005; 111:465–471.
33. Nagata C, Takatsuka N, Shimizu H. Soy and fish oil intake and mortality in a Japanese community. *Am J Epidemiol* 2002; 156:824–831.

34. Kokubo Y, Iso H, Ishihara J, *et al.* Association of dietary intake of soy, beans, and isoflavones with risk of cerebral and myocardial infarctions in Japanese populations: the Japan Public Health Center-based (JPHC) study cohort I. *Circulation* 2007; 116:2553–2562.
35. Bazzano LA, He J, Ogden LG, *et al.* Dietary fiber intake and reduced risk of coronary heart disease in US men and women: the National Health and Nutrition Examination Survey I Epidemiologic Follow-up Study. *Arch Intern Med* 2003; 163:1897–1904.
36. Mente A, de Koning L, Shannon HS, Anand SS. A systematic review of the evidence supporting a causal link between dietary factors and coronary heart disease. *Arch Intern Med* 2009; 169:659–669.
- A systematic search of MEDLINE for prospective cohort studies or randomized trials showed a valid association of a limited number of dietary factors and dietary patterns with coronary heart disease.
37. Oh K, Hu FB, Cho E, *et al.* Carbohydrate intake, glycemic index, glycemic load, and dietary fiber in relation to risk of stroke in women. *Am J Epidemiol* 2005; 161:161–169.
38. Anderson JW, Tietjen-Clark J. Dietary fiber: hyperlipidemia, hypertension, and coronary heart disease. *Am J Gastroenterol* 1986; 81:907–919.
39. Brown L, Rosner B, Willett WW, Sacks FM. Cholesterol-lowering effects of dietary fiber: a meta-analysis. *Am J Clin Nutr* 1999; 69:30–42.
40. Marckmann P, Sandstrom B, Jespersen J. Effects of total fat content and fatty acid composition in diet on factor VII coagulant activity and blood lipids. *Atherosclerosis* 1990; 80:227–233.
41. Pereira MA, Pins JJ. Dietary fiber and cardiovascular disease: experimental and epidemiologic advances. *Curr Atheroscler Rep* 2000; 2:494–502.
42. Jenkins DJ, Axelsen M, Kendall CW, *et al.* Dietary fibre, lente carbohydrates and the insulin-resistant diseases. *Br J Nutr* 2000; 83 (Suppl 1):S157–S163.
43. Ajani UA, Ford ES, Mokdad AH. Dietary fiber and C-reactive protein: findings from National Health and Nutrition Examination Survey Data. *J Nutr* 2004; 134:1181–1185.
44. Kokubo Y, Iso H, Saito I, *et al.* Dietary fiber intake and risk of cardiovascular disease in the Japanese population: the Japan Public Health Center-based study cohort. *Eur J Clin Nutr* 2011; 65:1233–1241.
- A cohort study showed that higher total dietary fiber was associated with reduced risk of cardiovascular disease in Japanese nonsmokers.
45. Iso H, Kobayashi M, Ishihara J, *et al.* Intake of fish and n3 fatty acids and risk of coronary heart disease among Japanese: the Japan Public Health Center-Based (JPHC) Study Cohort I. *Circulation* 2006; 113:195–202.
46. Yamagishi K, Iso H, Date C, *et al.* Fish, omega-3 polyunsaturated fatty acids, and mortality from cardiovascular diseases in a nationwide community-based cohort of Japanese men and women: the JACC (Japan Collaborative Cohort Study for Evaluation of Cancer Risk) Study. *J Am Coll Cardiol* 2008; 52:988–996.
47. Hooper L, Thompson RL, Harrison RA, *et al.* Risks and benefits of omega 3 fats for mortality, cardiovascular disease, and cancer: systematic review. *Br Med J* 2006; 332:752–760.
48. He K, Merchant A, Rimm EB, *et al.* Dietary fat intake and risk of stroke in male US healthcare professionals: 14 year prospective cohort study. *Br Med J* 2003; 327:777–782.
49. Soedamah-Muthu SS, Ding EL, Al-Delaimy WK, *et al.* Milk and dairy consumption and incidence of cardiovascular diseases and all-cause mortality: dose-response meta-analysis of prospective cohort studies. *Am J Clin Nutr* 2011; 93:158–171.
- This dose-response meta-analysis of prospective studies indicates that milk intake is not associated with total mortality but may be inversely associated with overall cardiovascular disease risk; however, these findings are based on limited numbers.
50. Sauvaget C, Nagano J, Hayashi M, Yamada M. Animal protein, animal fat, and cholesterol intakes and risk of cerebral infarction mortality in the adult health study. *Stroke* 2004; 35:1531–1537.
51. Virtanen JK, Mursu J, Voutilainen S, Tuomainen TP. Serum long-chain n-3 polyunsaturated fatty acids and risk of hospital diagnosis of atrial fibrillation in men. *Circulation* 2009; 120:2315–2321.
52. Brouwer IA, Heeringa J, Geleijnse JM, *et al.* Intake of very long-chain n-3 fatty acids from fish and incidence of atrial fibrillation. The Rotterdam Study. *Am Heart J* 2006; 151:857–862.
53. Nakayama T, Date C, Yokoyama T, *et al.* A 15.5-year follow-up study of stroke in a Japanese provincial city. The Shibata Study. *Stroke* 1997; 28:45–52.
54. Carlson SA, Fulton JE, Schoenborn CA, Loustalot F. Trend and prevalence estimates based on the 2008 Physical Activity Guidelines for Americans. *Am J Prev Med* 2010; 39:305–313.
- In the 2008 Physical Activity Guidelines among US adults, little progress has been made during the past 10 years in increasing physical activity levels. There is much room for improvement in achieving recommended levels of physical activity among Americans, particularly among relatively inactive subgroups.
55. Physical Activity Guidelines Advisory Committee report, 2008. To the Secretary of Health and Human Services. Part A: executive summary. *Nutr Rev* 2009; 67:114–120.
56. Lee CD, Folsom AR, Blair SN. Physical activity and stroke risk: a meta-analysis. *Stroke* 2003; 34:2475–2481.
57. Elsayy B, Higgins KE. Physical activity guidelines for older adults. *Am Fam Physician* 2010; 81:55–59.
- This article provides information and guidance on the amount of physical activity recommended to maintain health and fitness for older adults in the US.
58. Nelson ME, Rejeski WJ, Blair SN, *et al.* Physical activity and public health in older adults: recommendation from the American College of Sports Medicine and the American Heart Association. *Med Sci Sports Exerc* 2007; 39:1435–1445.
59. Deplanque D, Masse I, Lefebvre C, *et al.* Prior TIA, lipid-lowering drug use, and physical activity decrease ischemic stroke severity. *Neurology* 2006; 67:1403–1410.
60. Noda H, Iso H, Toyoshima H, *et al.* Walking and sports participation and mortality from coronary heart disease and stroke. *J Am Coll Cardiol* 2005; 46:1761–1767.
61. Myint PK, Luben RN, Wareham NJ, *et al.* Combined effect of health behaviours and risk of first ever stroke in 20 040 men and women over 11 years' follow-up in Norfolk cohort of European Prospective Investigation of Cancer (EPIC Norfolk): prospective population study. *Br Med J* 2009; 338:b349.

Associations of impaired glucose metabolism and dyslipidemia with cardiovascular diseases: what have we learned from Japanese cohort studies for individualized prevention and treatment?

Yoshihiro Kokubo

Received: 12 January 2011 / Accepted: 10 March 2011 / Published online: 2 April 2011
© European Association for Predictive, Preventive and Personalised Medicine 2011

Abstract Metabolic disorder is a modifiable risk factor for cardiovascular diseases (CVD), and lifestyle modification is the key to improving metabolic disorder. Diabetes mellitus has been shown to be a risk factor for coronary heart disease (CHD) and ischemic stroke in both Western and Japanese populations. An association between impaired fasting glucose and pre-hypertension found in an urban Japanese population emphasized the combined risk of CVD. Mean total cholesterol levels in Japan have been increasing in the last three decades. The Japanese evidence for the positive association of total cholesterol with CHD is similar to that in the West. Higher low-density lipoprotein cholesterol (LDL-C) levels pose an increased risk of CHD and atherothrombotic infarction, whereas lower LDL-C levels may pose an increased risk of intracerebral hemorrhage in Japan. Overall, the studies reviewed here show that impaired glucose metabolism and dyslipidemia are emerging risk factors for CVD in the Japanese population.

Keywords Impaired glucose metabolism · Dyslipidemia · Lifestyle · Predictors for cardiovascular disease · Cohort study · Japanese population

Introduction

Metabolic syndrome comprises a cluster of components of impaired glucose metabolism, abdominal fat accumulation, dyslipidemia, and elevated blood pressure [1]. Each component has been shown to be an independent risk factor for cardiovascular diseases (CVD) in Japanese community cohort studies: impaired fasting glucose [2]; abdominal obesity [3, 4]; low-density lipoprotein cholesterol [5, 6]; and high-normal blood pressure [7–9].

Recently, a new streamlined definition of metabolic syndrome for global populations has been introduced [10]. In this definition, abdominal obesity is just one of the possible components for metabolic syndrome. However, in the Japanese definition of metabolic syndrome, abdominal obesity is an essential component [11]. The definition may mislead a prevention of CVD for non-obesity metabolic disorder, which is a modifiable risk factor for CVD. Changes in lifestyle are the key to improving metabolic disorder, including diabetes mellitus (DM) and dyslipidemia, and thereby to reducing the risk of cardiovascular disease. In this review paper, the focus is on the relation of metabolic symptoms, namely impaired glucose metabolism and dyslipidemia to CVD in the Japanese population.

Impaired glucose metabolism: trends, combination with blood pressure elevation, and individualized preventive approaches

Diabetes mellitus has become a major public health problem [12, 13] as well as a risk factor for mortality [12] and CVD

Y. Kokubo (✉)
Department of Preventive Cardiology,
National Cerebral and Cardiovascular Center,
5-7-1, Fujishiro-dai,
Suita, Osaka 565–8565 Japan
e-mail: ykokubo@hsp.ncvc.go.jp

[14–16]. In Japan, the frequencies of hyperglycemia for 1961, 1974, and 1989 show a trend of increasing incidence: 12.1%, 13.8%, and 31.9% in men and 4.8%, 8.1%, and 27.2% in women, respectively [17].

Previous meta-analysis studies have shown that diabetes is a risk factor for ischemic stroke [18] and coronary artery disease [15, 18, 19]. The Framingham Offspring cohort study showed a positive relationship between impaired fasting glucose and coronary heart disease in women and between diabetes and coronary heart disease in men and women [20]. These results are compatible with those of previous cohort studies in Japan, as described below and summarized in Table 1.

The Hisayama Study indicated that diabetes as defined by the glucose tolerance test was a risk factor for ischemic stroke in men (hazard ratio 2.54; 95% confidence intervals 1.40 to 4.63) and women (hazard ratio 2.02; 95% confidence intervals 1.07 to 3.81) and coronary heart

disease in women (hazard ratio 3.46; 95% confidence intervals 1.59 to 7.54) [21]; however, impaired glucose tolerance and impaired fasting glucose were not risk factors for ischemic stroke or coronary heart disease.

In a study of five communities in Japan, diabetes defined by non-fasting glucose levels was a risk factor for non-embolic ischemic stroke in men (hazard ratio 1.8; 95% confidence intervals 1.0 to 3.2) and women (hazard ratio 2.2; 95% confidence intervals 1.2 to 4.0) [22]. This result is similar to that of lacunar infarction. The risk was observed in both non-hypertensive subjects and obese subjects. The positive association was particularly strong in hypertensive subjects with higher skin-fold thickness values (hazard ratio 1.9 and 95% confidence intervals 1.0 to 3.7 for borderline diabetes; hazard ratio 4.9 and 95% confidence intervals 2.5 to 9.5 for diabetes).

A positive association between diabetes and CVD mortality was also observed in a Japanese general population. The

Table 1 Association between blood glucose categories and cardiovascular diseases in Japanese cohort studies

Study name	Number	Sex	Follow-up	End point	Results	Reference
The Suita Study	5321	MF	11.5	Stroke	DM, HR=2.08	[2]
				CHD	IFG, HR=1.46; DM, HR=2.28	
		M		Stroke	DM, HR=1.78	
		F		Stroke	DM, HR=2.66	
		CHD		IFG, HR=1.83; DM, HR=4.32		
The Hisayama Study	2421	M	14	CI	DM, HR=2.15; 2hPG, HR=2.71	[21]
		F		CI	DM, HR=2.10; 2hPG, HR=2.19	
		CHD		DM, HR=3.83; 2hPG, HR=4.44		
Five Japanese communities	4287	M	17	CI	DM, HR=1.8	[22]
		Lacunar infarction		DM, HR=2.1		
	6295	F		CI	DM, HR=2.2	
	Lacunar infarction	DM, HR=2.4				
JPHC study	31,192	MF	12.9	CHD	Borderline HR=1.5	[25]
					DM, HR=2.38	
					DM with higher BMI, HR=2.2	
NIPPON DATA80	9444	MF	17.3	All causes mortality	CBG \geq 11.1 mmol/L, HR=1.63	[23]
				CVD mortality	5.22 mmol/L \leq CBG<7.77 mmol/L, HR=1.22	
				7.77 mmol/L \leq CBG<11.1 mmol/L, HR=1.46		
				CBG \geq 11.1 mmol/L, HR=1.82		
				CHD mortality	7.77 mmol/L \leq CBG<11.1 mmol/L, HR=2.43	
The Funagata Diabetes Study	2534	MF	5.7	All cause mortality	ADA 2007: DM, HR=2.11	[24]
				CVD mortality	WHO 1985: IGT, HR=2.3; DM, HR=3.54	
				ADA 2007: DM, HR=3.17		

M men; F women; CHD coronary heart disease; CI cerebral infarction; CVD cardiovascular diseases; DM diabetes mellitus; IFG impaired fasting glucose; 2hPG 2 h post-loaded glucose levels; HR hazard ratio; BMI body mass index; CBG casual blood glucose; ADA American Diabetes Association; WHO World Health Organization.

NIPPON DATA 80 Study indicated that high and borderline-high casual blood glucose groups (11.1 mmol/L and 7.77 to 11.1 mmol/L, respectively) had increased risks of coronary heart disease mortality [23]. Similar results were observed for both CVD and all-cause mortality. The Funagata Diabetes Study showed that diabetes defined by both the WHO criteria (1985) and ADA recommendations (1997) were risk factors for all-cause mortality and cardiovascular mortality [24]. However, impaired fasting glucose was not a risk factor for all-cause mortality and cardiovascular mortality. These Japanese cohort studies for mortality risks indicated that diabetes is a risk factor for all-cause and CVD mortality. However, impaired fasting glucose and impaired glucose tolerance may not be risk factors for all-cause or CVD mortality. Further investigations of larger cohort studies are required to clarify these matters.

Recently, the Suita Study showed that impaired fasting glucose is a risk factor for the incidence of cardiovascular disease (hazard ratio 1.49; 95% confidence intervals 1.02 to 2.16) or coronary heart disease (hazard ratio 1.83; 95% confidence intervals 1.01 to 3.32) in women, and that diabetes is a risk factor for stroke in both men (hazard ratio 1.78; 95% confidence intervals 1.00 to 3.12) and women (hazard ratio 2.66; 95% confidence intervals 1.22 to 5.80) and for coronary heart disease in women (hazard ratio 1.78; 95% confidence intervals 1.00 to 3.12) [2]. In addition, compared with normoglycemic and optimal blood pressure Japanese subjects, increased risks of CVD were observed in normoglycemic subjects with high-normal blood pressure or hypertension, impaired fasting glucose subjects with normal or higher blood pressure, and diabetic subjects regardless of blood pressure category (Fig. 1: P -value for interaction=0.046). These two borderline categories may augment the risk of CVD.




Fig. 1 Multivariable-adjusted hazard ratios of cardiovascular diseases according to the combination of blood pressure and glucose categories. Multivariable-adjusted hazard ratios were adjusting for age, sex, body mass index, smoking, drinking, and hyperlipidemia. *BP* blood pressure; *NGT* normal glycaemic tolerance; *IFG* impaired fasting glucose; *DM* diabetes mellitus; *: $P < 0.05$ (compared with the optimal blood pressure and normoglycemic group)

In a collaborative meta-analysis of 102 prospective studies from around the world, increased risks of coronary heart disease were observed in unknown diabetic subjects (i.e., subjects with diabetes who are not aware of their disease) with 5.6 to <6.1 mmol/L and 6.1 to <7 mmol/L (hazard ratios [95% confidence intervals]: 1.11 [1.04 to 1.18] and 1.17 [1.08 to 1.26], respectively) [18]. Quite recently, the Japan Public Health Center-based prospective (JPHC) study consisting of a 12.9-year of follow-up for 31,192 individuals aged 40–69 years was observed that diabetes mellitus and hyperglycemia carried an increased risk for coronary heart disease in a Japanese general population [25]. Based on this study and the Suita study [2, 25], both diabetes mellitus and impaired fasting glucose may be a risk factor for the incidence of coronary heart disease in Japan.

In the JPHC study's systematic review of the evidence for pre-diabetes, impaired fasting glucose and impaired glucose tolerance were shown to be associated with modest increases in the risk for CVD [25]. Overall relative risks (95% confidence intervals) for the association between two categories of impaired fasting glucose (100 to 125 mg/dL and 110 to 125 mg/dL) and CVD were 1.18 (1.09 to 1.28) and 1.20 (1.12 to 1.28), respectively. Meanwhile, the risks of CVD for impaired fasting glucose and impaired glucose tolerance were 1.10 (0.98 to 1.23) and 1.20 (1.07 to 1.34), respectively. In two Japanese cohort studies using the oral glucose tolerance test, impaired fasting glucose was not found to pose a risk of CVD [21, 24]. However, in one of these cohort studies, impaired glucose tolerance was observed as a risk of CVD mortality [24]. Larger Japanese cohort studies are required to establish the association between oral glucose tolerance test results and CVD.

In order to prevent CVD, a person with impaired glucose metabolism should eliminate or reduce other cardiovascular risk factors, such as high blood pressure, obesity, smoking, and excess drinking. The Suita study showed that the combination of impaired fasting glucose and high blood pressure greatly increased the risk of cardiovascular disease [2]. Recently, the frequencies of obesity in men and elderly women have increased in Japan [26]. Obesity is a risk factor for impaired glucose metabolism; weight loss reduces the risk of both impaired glucose metabolism [27] and CVD [3].

The average smoking rate in men worldwide has rapidly decreased in recent years, and in 2005 had dropped by 45.5% since its peak. However, the rate in Japan is still higher than that in Western countries [28]. And while, the smoking rate in women is generally lower than that in men. In Japan, the rate for younger women aged 20 to 39 years has increased to around 17 to 20% [28]. Smoking itself is a risk factor for diabetes mellitus [29]. Smoking cessation reduces the risk of both diabetes mellitus [30] and CVD

[31]. Excess drinking is a risk factor for increasing blood pressure [32] and stroke [33]. In order to reduce the risk of CVD, subjects with a combination of two borderline risks, high-normal blood pressure and hyperglycemia, should modify their lifestyles, aiming at cessation of smoking and even moderate drinking, as well as weight reduction if there are overweight or obese.

Dyslipidemia: risks, trends, and individual treatment approaches

According to Japan's National Survey on Circulatory Disorders in 1980, 1990, and 2000, mean levels of total cholesterol increased from 186 mg/dL and 191 mg/dL in 1980 to 200 mg/dL and 208 mg/dL in 2000 among adult men and women, respectively. Frequencies of hypercholesterolemia (total cholesterol levels of 220 mg/dL or more) increased 15% and 19% in 1980 to 27% and 35% in 2000 among men and women, respectively [34]. The difference in mean total cholesterol levels between Japanese and American men was approximately 40 mg/dL in the 1980s. However, this difference had diminished to 15 mg/dL by 1990 and 2000. No changes in mean total cholesterol levels in Japanese or American men aged 60 years and over were observed from 1990 to 2000 [28].

Total cholesterol

A previous meta-analytic study has shown that total cholesterol was positively associated with ischemic heart disease in both middle and old age and at all blood pressure categories [35]. In contrast, the association between total cholesterol and the risk of stroke remains unclear. The Multiple Risk Factor Intervention trial has shown that the risk of death from intracerebral hemorrhage was three times higher in men with serum cholesterol levels under 160 mg/dL than in those with higher cholesterol levels, whereas a positive association was observed between the serum cholesterol level and death from ischemic stroke [36]. A meta-analysis of 45 prospective cohort studies showed no association between blood cholesterol levels and stroke except in those under 45 years of age when screened [37]. The inconsistency in results may be due to the different etiologic origins of stroke. Dyslipidemia may be important for some subtypes of stroke but not for others, because stroke is a heterogeneous disease of various etiologic origins. In American men, an inverse association was observed between the total cholesterol levels and the incidence of intracerebral hemorrhage, while a positive association was observed between total cholesterol levels and the incidence of cerebral infarction [36].

Japanese cohort studies have shown evidence of the positive association of total cholesterol with coronary heart disease similar to that in Western studies. NIPPON DATA80 showed associations between total cholesterol and risk of all-cause mortality (hazard ratio 1.19 and 1.36; 95% confidence intervals, 1.03 to 1.37 and 1.05 to 1.77) in both the lowest (<160 mg/dl) and highest (\geq 260 mg/dl) total cholesterol groups, respectively [38]. In addition, the hazard ratios of coronary heart disease mortality were 1.4 in the 200 to 219 mg/dL total cholesterol group, 1.7 in the 220 to 239 mg/dL group, 1.8 in the 240 to 259 mg/dL groups, and 3.8 in the 260 mg/dL or above group, compared with the 160 to 179 mg/dL total cholesterol (healthy control) group [38].

LDL cholesterol and non-HDL cholesterol

The Suita Study showed a positive association between serum low-density lipoprotein (LDL) cholesterol and non-high-density lipoprotein (non-HDL) cholesterol levels and increased incidence of myocardial infarction, but not with any type of stroke [6]. The hazard ratio for myocardial infarction was highest in the top quintile of LDL cholesterol (hazard ratio, 3.03; 95% confidence intervals, 1.32 to 6.96) when men and women were combined. The hazard ratio for myocardial infarction was also highest in the top quintile of non-HDL cholesterol (hazard ratio, 2.97; 95% confidence intervals, 1.26 to 6.97). There was no substantial difference in the predictive value for the incidence of myocardial infarction between LDL cholesterol and non-HDL cholesterol. LDL cholesterol can be calculated from fasting blood sample and measurement of total cholesterol, HDL cholesterol, and triglyceride levels from fasting blood samples, according to the Friedewald formula [39]. The formula is not applicable for serum triglyceride levels equal to or greater than \geq 400 mg/dL. However, non-HDL cholesterol levels can be easily calculated by routine measured parameters, total and HDL cholesterol without the effect of non-fasting status or hypertriglyceridemia (\geq 400 mg/dL).

The Hisayama Study showed that the positive association between LDL cholesterol levels and risk of atherothrombotic infarction (P for trend=0.02) and coronary heart disease (P for trend=0.03) remained significant after multivariable adjustment. In the Ibaraki Prefectural Health Study, which was a very large sample consisting of 30,802 men and 60,417 women, high LDL cholesterol levels were associated with an increased risk of mortality from coronary heart disease in men, but not in women [40]. The same study showed that lower LDL cholesterol levels are associated with elevated risk of mortality from intracerebral hemorrhage, but are not associated with increasing risk of cerebral infarction [41]. These Japanese cohort studies suggest that higher levels

of LDL cholesterol pose an increased risk of coronary heart disease and possibly atherothrombotic infarction, whereas lower levels of LDL cholesterol may increase the risk of intracerebral hemorrhage.

Recently, the lectin-like oxidized LDL receptor 1 (LOX-1) has been implicated in atherothrombotic diseases [42]. Activation of LOX-1 in humans can be evaluated by use of the LOX index, obtained by multiplying the circulating concentration of LOX-1 ligands containing apolipoprotein B by that of the soluble form of LOX-1. In the Suita Study, higher LOX index values were associated with an increased risk of coronary heart disease [43].

HDL cholesterol

Lower HDL cholesterol predicts coronary heart disease mortality and occurrence of new coronary heart disease events [44]. Elevated total cholesterol was not found to be associated with coronary heart disease mortality in older men, but may be a risk factor for coronary heart disease in older women [44]. The Israeli Ischemic Heart Disease Study showed an independent negative association between HDL cholesterol and ischemic stroke mortality during 21 years of follow-up [45].

Among the Japanese cohort studies, the Oyabe Study demonstrated that lower HDL cholesterol levels were related significantly and independently to an increased risk of all-stroke incidence and ischemic stroke incidence [46]. And in a combined prospective cohort study of 13 urban industrial companies, coronary heart disease incidence was inversely related to HDL cholesterol in Japanese men [47].

Triglyceride

Based on combined data from prospective studies, a high serum triglyceride level is a risk factor for CVD for both men (hazard ratio, 1.32; 95% confidence intervals, 1.26 to 1.39) and women (hazard ratio, 1.76; 95% confidence intervals, 1.50 to 2.07) in the general population, independent of HDL cholesterol [48]. In 26 prospective studies in Asian and Pacific populations, serum triglycerides were an independent predictor of coronary heart disease and stroke risk [49]. In 29 prospective meta-analytic studies in the West, triglyceride levels were moderately associated with coronary heart disease [50].

In Japanese cohort studies, two studies have provided evidence regarding the association between serum triglyceride levels and coronary heart disease and ischemic stroke. A cohort study of four rural communities showed that a high serum triglyceride level was a risk factor for coronary heart disease; even after adjustment for HDL cholesterol levels the significant association remained, although 80% of the baseline participants were non-fasting [51]. The Suita Study has also

shown that the risk for ischemic stroke was highest in participants with high triglycerides alone, and that a combination of high serum levels of triglyceride and non-HDL cholesterol was associated with an increased risk of myocardial infarction [52]. High serum levels of triglyceride and non-HDL cholesterol are both important targets for the prevention of CVD, which requires evidence-based guidelines for management in the primary care setting.

Concluding remarks and outlook

This paper reviews the associations of impaired glucose metabolism and dyslipidemia with CVD in Japanese cohort studies. Diabetes mellitus is a risk factor for coronary heart disease and ischemic stroke. Impaired fasting glucose and high-normal blood pressure were shown to be independent risk factors for CVD and coronary heart disease in an urban cohort. The combination of these two borderline categories may increase the risk for CVD. Impaired glucose tolerance has not been observed as a risk factor for the incidence of CVD in Japan. The Japanese evidence for the positive association of total cholesterol with coronary heart disease is similar to that of previous Western studies. Associations with all-cause mortality were observed for both the lower and higher levels of cholesterol: Higher levels of LDL cholesterol have been shown to increase the risk of coronary heart disease and atherothrombotic infarction, whereas lower levels of LDL cholesterol may increase the risk of intracerebral hemorrhage in Japan, as elsewhere. HDL cholesterol levels were inversely related with ischemic stroke. Positive associations between serum triglyceride levels and coronary heart disease and ischemic stroke have also been observed in Japanese populations. The Japanese diet has been rapidly changing in recent decades, as reflected in many of its health indicators such as cholesterol levels, and both impaired glucose metabolism and dyslipidemia are emerging as important risk factors for CVD in the Japanese population. In order to reduce the risk of CVD, subjects with metabolic disorder should reduce other cardiovascular risk factors and improve their lifestyle.

References

1. Eckel RH, Grundy SM, Zimmet PZ. The metabolic syndrome. *Lancet*. 2005;365:1415–28.
2. Kokubo Y, Okamura T, Watanabe M, et al. The combined impact of blood pressure category and glucose abnormality on the incidence of cardiovascular diseases in a Japanese urban cohort: the Suita Study. *Hypertens Res*. 2010;33:1238–43.
3. Furukawa Y, Kokubo Y, Okamura T, et al. The relationship between waist circumference and the risk of stroke and myocar-

- dial infarction in a Japanese urban cohort: the Suita study. *Stroke*. 2010;41:550–3.
4. Doi Y, Ninomiya T, Hata J, et al. Proposed criteria for metabolic syndrome in Japanese based on prospective evidence: the Hisayama study. *Stroke*. 2009;40:1187–94.
 5. Executive Summary of the Third Report of the National Cholesterol Education Program (NCEP). Expert panel on detection, evaluation, and treatment of high blood cholesterol in adults (Adult Treatment Panel III). *JAMA*. 2001;285:2486–97.
 6. Okamura T, Kokubo Y, Watanabe M, et al. Low-density lipoprotein cholesterol and non-high-density lipoprotein cholesterol and the incidence of cardiovascular disease in an urban Japanese cohort study: The Suita study. *Atherosclerosis*. 2009;203:587–92.
 7. Kokubo Y, Kamide K, Okamura T, et al. Impact of high-normal blood pressure on the risk of cardiovascular disease in a Japanese urban cohort: the Suita study. *Hypertension*. 2008;52:652–9.
 8. Ikeda A, Iso H, Yamagishi K, Inoue M, Tsugane S. Blood pressure and the risk of stroke, cardiovascular disease, and all-cause mortality among Japanese: the JPHC Study. *Am J Hypertens*. 2009;22:273–80.
 9. Kokubo Y, Kamide K. High-normal blood pressure and the risk of cardiovascular disease. *Circ J*. 2009;73:1381–5.
 10. Alberti KG, Eckel RH, Grundy SM, et al. Harmonizing the metabolic syndrome: a joint interim statement of the international diabetes federation task force on epidemiology and prevention; national heart, lung, and blood institute; american heart association; world heart federation; international atherosclerosis society; and international association for the study of obesity. *Circulation*. 2009;120:1640–5.
 11. Kokubo Y, Okamura T, Yoshimasa Y, et al. Impact of metabolic syndrome components on the incidence of cardiovascular disease in a general urban Japanese population: the Suita study. *Hypertens Res*. 2008;31:2027–35.
 12. Garcia MJ, McNamara PM, Gordon T, Kannel WB. Morbidity and mortality in diabetics in the Framingham population. Sixteen year follow-up study. *Diabetes*. 1974;23:105–11.
 13. Manuel DG, Schultz SE. Health-related quality of life and health-adjusted life expectancy of people with diabetes in Ontario, Canada, 1996–1997. *Diabetes Care*. 2004;27:407–14.
 14. Fox CS, Coady S, Sorlie PD, et al. Increasing cardiovascular disease burden due to diabetes mellitus: the framingham heart study. *Circulation*. 2007;115:1544–50.
 15. Lee WL, Cheung AM, Cape D, Zinman B. Impact of diabetes on coronary artery disease in women and men: a meta-analysis of prospective studies. *Diabetes Care*. 2000;23:962–8.
 16. Folsom AR, Rasmussen ML, Chambless LE, et al. Prospective associations of fasting insulin, body fat distribution, and diabetes with risk of ischemic stroke. The Atherosclerosis Risk in Communities (ARIC) study investigators. *Diabetes Care*. 1999;22:1077–83.
 17. Kubo M, Kiyohara Y, Kato I, et al. Trends in the incidence, mortality, and survival rate of cardiovascular disease in a Japanese community: the Hisayama study. *Stroke*. 2003;34:2349–54.
 18. Sarwar N, Gao P, Seshasai SR, et al. Diabetes mellitus, fasting blood glucose concentration, and risk of vascular disease: a collaborative meta-analysis of 102 prospective studies. *Lancet*. 2010;375:2215–22.
 19. Sarwar N, Aspelund T, Eiriksdottir G, et al. Markers of dysglycaemia and risk of coronary heart disease in people without diabetes: Reykjavik prospective study and systematic review. *PLoS Med*. 2010;7:e1000278.
 20. Levitzky YS, Pencina MJ, D'Agostino RB, et al. Impact of impaired fasting glucose on cardiovascular disease: the framingham heart study. *J Am Coll Cardiol*. 2008;51:264–70.
 21. Doi Y, Ninomiya T, Hata J, et al. Impact of glucose tolerance status on development of ischemic stroke and coronary heart disease in a general Japanese population: the Hisayama study. *Stroke*. 2010;41:203–9.
 22. Iso H, Imano H, Kitamura A, et al. Type 2 diabetes and risk of non-embolic ischaemic stroke in Japanese men and women. *Diabetologia*. 2004;47:2137–44.
 23. Kadowaki S, Okamura T, Hozawa A, et al. Relationship of elevated casual blood glucose level with coronary heart disease, cardiovascular disease and all-cause mortality in a representative sample of the Japanese population. NIPPON DATA80. *Diabetologia*. 2008;51:575–82.
 24. Tomiyama M, Eguchi H, Manaka H, Igarashi K, Kato T, Sekikawa A. Impaired glucose tolerance is a risk factor for cardiovascular disease, but not impaired fasting glucose. The Funagata Diabetes Study. *Diabetes Care*. 1999;22:920–4.
 25. Saito I, Kokubo Y, Yamagishi K, Iso H, Inoue M, Tsugane S. Diabetes and the risk of coronary heart disease in the general Japanese population: the Japan Public Health Center-based prospective (JPHC) study. *Atherosclerosis*. 2011. doi:10.1016/j.atherosclerosis.2011.01.021.
 26. Yoshiike N, Seino F, Tajima S, et al. Twenty-year changes in the prevalence of overweight in Japanese adults: the national nutrition survey 1976–95. *Obes Rev*. 2002;3:183–90.
 27. Lechleitner M. Obesity and the metabolic syndrome in the elderly—a mini-review. *Gerontology*. 2008;54:253–9.
 28. Ueshima H. Explanation for the Japanese paradox: prevention of increase in coronary heart disease and reduction in stroke. *J Atheroscler Thromb*. 2007;14:278–86.
 29. Willi C, Bodenmann P, Ghali WA, Faris PD, Cornuz J. Active smoking and the risk of type 2 diabetes: a systematic review and meta-analysis. *JAMA*. 2007;298:2654–64.
 30. Yeh HC, Duncan BB, Schmidt MI, Wang NY, Brancati FL. Smoking, smoking cessation, and risk for type 2 diabetes mellitus: a cohort study. *Ann Intern Med*. 2010;152:10–7.
 31. Honjo K, Iso H, Tsugane S, et al. The effects of smoking and smoking cessation on mortality from cardiovascular disease among Japanese: pooled analysis of three large-scale cohort studies in Japan. *Tob Control*. 2010;19:50–7.
 32. Beilin LJ, Puddey IB, Burke V. Alcohol and hypertension—kill or cure? *J Hum Hypertens*. 1996;10 Suppl 2:S1–5.
 33. Iso H, Baba S, Mannami T, et al. Alcohol consumption and risk of stroke among middle-aged men: the JPHC study cohort I. *Stroke*. 2004;35:1124–9.
 34. Iso H. Changes in coronary heart disease risk among Japanese. *Circulation*. 2008;118:2725–9.
 35. Lewington S, Whitlock G, Clarke R, et al. Blood cholesterol and vascular mortality by age, sex, and blood pressure: a meta-analysis of individual data from 61 prospective studies with 55,000 vascular deaths. *Lancet*. 2007;370:1829–39.
 36. Iso H, Jacobs Jr DR, Wentworth D, Neaton JD, Cohen JD. Serum cholesterol levels and six-year mortality from stroke in 350,977 men screened for the multiple risk factor intervention trial. *N Engl J Med*. 1989;320:904–10.
 37. Cholesterol, diastolic blood pressure, and stroke: 13,000 strokes in 450,000 people in 45 prospective cohorts. Prospective studies collaboration. *Lancet*. 1995;346:1647–53.
 38. Okamura T, Tanaka H, Miyamatsu N, et al. The relationship between serum total cholesterol and all-cause or cause-specific mortality in a 17.3-year study of a Japanese cohort. *Atherosclerosis*. 2007;190:216–23.
 39. Friedewald WT, Levy RI, Fredrickson DS. Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *Clin Chem*. 1972;18:499–502.
 40. Noda H, Iso H, Irie F, Sairenchi T, Ohtaka E, Ohta H. Gender difference of association between LDL cholesterol concentra-

- tions and mortality from coronary heart disease amongst Japanese: the Ibaraki prefectural health study. *J Intern Med*. 2010;267:576–87.
41. Noda H, Iso H, Irie F, et al. Low-density lipoprotein cholesterol concentrations and death due to intraparenchymal hemorrhage: the Ibaraki prefectural health study. *Circulation*. 2009;119:2136–45.
 42. Sawamura T, Kume N, Aoyama T, et al. An endothelial receptor for oxidized low-density lipoprotein. *Nature*. 1997;386:73–7.
 43. Inoue N, Okamura T, Kokubo Y, et al. LOX index, a novel predictive biochemical marker for coronary heart disease and stroke. *Clin Chem*. 2010;56:550–8.
 44. Corti MC, Guralnik JM, Salive ME, et al. HDL cholesterol predicts coronary heart disease mortality in older persons. *JAMA*. 1995;274:539–44.
 45. Goldbourt U, Yaari S, Medalie JH. Isolated low HDL cholesterol as a risk factor for coronary heart disease mortality. A 21-year follow-up of 8000 men. *Arterioscler Thromb Vasc Biol*. 1997;17:107–13.
 46. Soyama Y, Miura K, Morikawa Y, et al. High-density lipoprotein cholesterol and risk of stroke in Japanese men and women: the Oyabe Study. *Stroke*. 2003;34:863–8.
 47. Kitamura A, Iso H, Naito Y, et al. High-density lipoprotein cholesterol and premature coronary heart disease in urban Japanese men. *Circulation*. 1994;89:2533–9.
 48. Hokanson JE, Austin MA. Plasma triglyceride level is a risk factor for cardiovascular disease independent of high-density lipoprotein cholesterol level: a meta-analysis of population-based prospective studies. *J Cardiovasc Risk*. 1996;3:213–9.
 49. Patel A, Barzi F, Jamrozik K, et al. Serum triglycerides as a risk factor for cardiovascular diseases in the Asia-Pacific region. *Circulation*. 2004;110:2678–86.
 50. Sarwar N, Danesh J, Eiriksdottir G, et al. Triglycerides and the risk of coronary heart disease: 10,158 incident cases among 262,525 participants in 29 Western prospective studies. *Circulation*. 2007;115:450–8.
 51. Iso H, Naito Y, Sato S, et al. Serum triglycerides and risk of coronary heart disease among Japanese men and women. *Am J Epidemiol*. 2001;153:490–9.
 52. Okamura T, Kokubo Y, Watanabe M, et al. Triglycerides and non-high-density lipoprotein cholesterol and the incidence of cardiovascular disease in an urban Japanese cohort: the Suita study. *Atherosclerosis*. 2010;209:290–4.

