

Table 1. Baseline characteristics by BMI^a category in 13 764 men aged 40–64 years

	BMI (kg/m ²)							P value ^b
	<18.5	18.5–20.9	21.0–22.9	23.0–24.9	25.0–27.4	27.5–29.9	≥30.0	
No. of subjects	310	2159	3591	3852	2637	903	312	
Mean age (years) (SD ^a)	54.2 (7.7)	53.0 (8.0)	53.1 (7.8)	53.2 (7.6)	52.8 (7.6)	52.1 (7.4)	52.0 (7.6)	<0.0001
Mean weight (kg) (SD)	48.9 (4.8)	54.6 (4.6)	59.4 (4.7)	64.6 (5.2)	70.3 (5.8)	77.4 (6.3)	86.9 (14.8)	<0.0001
Mean height (cm) (SD)	166.3 (7.6)	164.9 (6.6)	164.0 (6.3)	164.1 (6.2)	164.1 (6.5)	164.8 (6.5)	163.6 (19.1)	<0.0001
Mean BMI (kg/m ²) (SD)	17.6 (0.9)	20.1 (0.7)	22.1 (0.5)	24.0 (0.6)	26.1 (0.7)	28.5 (0.7)	32.4 (4.7)	<0.0001
Weight change since age 20 years (%)								
≤−10.0 kg	24.4	8.5	4.0	2.8	1.5	2.5	0.7	<0.0001
−9.9 to −5.0 kg	26.8	21.9	14.2	8.0	4.5	3.3	3.0	
−4.9 to +4.9 kg	43.5	61.2	57.2	36.1	15.5	6.6	4.3	
+5.0 to +9.9 kg	4.4	7.1	19.9	32.9	28.1	12.3	7.2	
≥+10.0 kg	1.0	1.3	4.6	20.2	50.4	75.2	84.9	
Education (%)								
Junior high school or less	57.2	56.6	55.8	55.1	52.3	54.2	55.8	NS ^a
High school	35.7	35.9	36.8	37.2	39.0	37.3	35.6	
College/university or higher	7.1	7.5	7.5	7.7	8.7	8.6	8.6	
Marital status (%)								
Married	84.3	87.0	88.8	89.7	88.6	90.4	82.4	0.0072
Unmarried	15.7	13.0	11.2	10.3	11.4	9.6	17.6	
Smoking status (%)								
Never smoker	13.1	13.1	18.2	21.5	24.9	24.9	23.6	<0.0001
Past smoker	16.8	14.1	17.2	19.4	22.9	21.2	19.5	
Current smoker, 1–19 cigarettes/day	23.4	26.6	20.9	19.0	14.7	13.3	13.8	
Current smoker, ≥20 cigarettes/day	46.7	46.3	43.7	40.2	37.5	40.7	43.1	
Alcohol drinking (%)								
Never drinker	20.7	14.7	14.3	14.5	14.7	15.2	19.7	<0.0001
Past drinker	14.4	7.3	7.3	5.9	7.0	5.8	8.9	
Current drinker	64.9	78.0	78.4	79.7	78.3	79.1	71.5	
Time spent walking (%)								
≥1 hour/day	44.3	55.9	53.8	53.0	49.1	46.1	45.3	<0.0001
<1 hour/day	55.8	44.1	46.2	47.0	50.9	53.9	54.7	
Sports and physical exercise (%)								
≥5 hours/week	2.4	5.8	5.4	5.8	4.5	4.4	5.9	0.0367
3–4 hours/week	3.8	4.1	4.7	5.2	5.0	4.8	3.0	
1–2 hours/week	11.9	12.1	14.2	14.1	14.3	15.8	12.9	
<1 hour/week	81.9	78.1	75.7	75.0	76.2	75.0	78.2	
History of hypertension (%)								
Yes	10.0	12.6	14.6	17.2	21.2	26.8	25.3	<0.0001
No	90.0	87.4	85.4	82.8	78.8	73.2	74.7	
History of diabetes (%)								
Yes	6.1	4.5	5.9	5.9	6.6	7.3	9.6	0.0015
No	93.9	95.6	94.2	94.1	93.4	92.7	90.4	
History of kidney disease (%)								
Yes	4.5	4.3	3.3	3.0	3.5	4.7	2.6	NS
No	95.5	95.7	96.7	97.0	96.6	95.4	97.4	
History of liver disease (%)								
Yes	6.1	6.4	6.2	7.0	7.4	7.6	8.0	NS
No	93.9	93.7	93.8	93.0	92.6	92.4	92.0	

^aBMI, body mass index; SD, standard deviation; NS, not significant.

^bP values were calculated by using the chi-square test (for categorical variables) or ANOVA (for continuous variables).

23.0 ≤ BMI ≤ 24.9 kg/m² (model 4). The HRs were similar to model 1 HRs: the model 4 multivariate HRs in underweight men were 1.18 (0.88–1.60) in middle-aged men and 1.42 (1.26–1.76) in elderly men, in obese men they were 1.64 (1.13–2.38) in middle-aged men and 1.25 (0.87–1.80) in elderly men, in underweight women they were 1.38 (0.94–1.99) in middle-aged women and 1.43 (1.19–1.71) in elderly women, and in obese women they were 1.41 (0.92–2.16) in middle-aged women and 1.25 (0.95–1.64) in elderly women.

DISCUSSION

The present results indicate that the mortality risk associated with underweight and obesity might be dependent upon sex and age group. We noted significant increased risks of mortality only in middle-aged obese men and elderly underweight men. In women, there was no significant excess risk of mortality with age in the obese, and no significant increased risk of mortality, irrespective of age group, in the underweight.

Table 2. Baseline characteristics by BMI^a category in 7274 men aged 65–79 years

	BMI (kg/m ²)							P value ^b
	<18.5	18.5–20.9	21.0–22.9	23.0–24.9	25.0–27.4	27.5–29.9	≥30.0	
No. of subjects	422	1518	2026	1805	1089	310	104	
Mean age (years) (SD ^a)	71.5 (4.2)	70.4 (4.2)	70.2 (4.0)	69.9 (4.0)	69.6 (3.9)	69.5 (3.8)	69.9 (4.3)	<0.0001
Mean weight (kg) (SD)	47.1 (4.9)	52.0 (4.4)	56.5 (4.7)	61.6 (5.0)	66.7 (5.4)	72.4 (6.5)	81.0 (21.6)	<0.0001
Mean height (cm) (SD)	164.5 (8.0)	161.1 (6.4)	160.1 (6.5)	160.4 (6.3)	159.9 (6.2)	159.6 (7.2)	153.8 (13.7)	<0.0001
Mean BMI (kg/m ²) (SD)	17.4 (1.0)	20.0 (0.7)	22.0 (0.6)	23.9 (0.6)	26.0 (0.7)	28.4 (0.6)	34.3 (8.1)	<0.0001
Weight change since age 20 years (%)								
≤−10.0 kg	55.0	29.9	13.5	6.9	3.1	3.1	4.0	<0.0001
−9.9 to −5.0 kg	28.9	37.1	30.5	16.6	8.4	5.9	5.0	
−4.9 to +4.9 kg	15.4	29.9	43.6	43.0	26.1	13.8	12.0	
+5.0 to +9.9 kg	0.8	2.3	9.7	22.1	29.1	19.3	8.0	
≥+10.0 kg	0.0	0.8	2.7	11.4	33.4	57.9	71.0	
Education (%)								
Junior high school or less	69.4	73.9	75.4	72.1	73.2	72.9	69.6	0.0107
High school	20.9	19.4	18.5	19.2	20.9	22.4	24.5	
College/university or higher	9.8	6.7	6.1	8.8	6.0	4.7	5.9	
Marital status (%)								
Married	89.8	90.3	90.0	89.2	91.7	91.4	89.6	NS ^a
Unmarried	10.2	9.7	10.0	10.9	8.3	8.6	10.4	
Smoking status (%)								
Never smoker	12.7	14.0	14.6	18.5	18.6	23.4	29.4	<0.0001
Past smoker	32.7	31.7	34.9	38.9	43.1	42.3	35.9	
Current smoker, 1–19 cigarettes/day	34.8	30.7	29.2	24.5	21.0	15.7	18.5	
Current smoker, ≥20 cigarettes/day	19.8	23.6	21.4	18.1	17.3	18.5	16.3	
Alcohol drinking (%)								
Never drinker	21.3	19.3	19.9	19.2	18.4	20.1	20.8	<0.0001
Past drinker	24.0	18.1	15.0	14.0	14.9	12.3	18.8	
Current drinker	54.8	62.6	65.1	66.7	66.7	67.6	60.4	
Time spent walking (%)								
≥1 hour/day	36.9	47.1	46.0	43.0	37.6	41.7	34.8	<0.0001
<1 hour/day	63.1	52.9	54.0	57.0	62.4	58.3	65.2	
Sports and physical exercise (%)								
≥5 hours/week	16.2	15.4	17.0	17.4	15.3	15.4	10.7	NS
3–4 hours/week	9.2	9.8	10.4	10.4	11.4	8.8	7.1	
1–2 hours/week	14.8	17.0	18.9	17.6	20.4	24.2	26.2	
<1 hour/week	59.9	57.9	53.7	54.5	52.9	51.7	56.0	
History of hypertension (%)								
Yes	21.6	27.8	32.8	38.2	43.3	48.4	52.9	<0.0001
No	78.4	72.2	67.2	61.8	56.7	51.6	47.1	
History of diabetes (%)								
Yes	6.2	6.7	9.2	9.7	9.7	12.9	11.5	0.0010
No	93.8	93.3	90.8	90.3	90.3	87.1	88.5	
History of kidney disease (%)								
Yes	5.0	4.0	3.4	4.5	3.5	2.9	3.9	NS
No	95.0	96.1	96.6	95.5	96.5	97.1	96.2	
History of liver disease (%)								
Yes	8.3	7.7	5.8	5.7	6.5	7.4	11.5	0.0284
No	91.7	92.3	94.2	94.4	93.5	92.6	88.5	

^aBMI, body mass index; SD, standard deviation; NS, not significant.

^bP values were calculated by using the chi-square test (for categorical variables) or ANOVA (for continuous variables).

We considered several important confounding factors: cigarette smoking, alcohol consumption, and physical activity are major confounding factors associated with both BMI and mortality.^{1–15,17,18} We also considered education level and marital status as potential confounding factors, as in past studies.^{1,3,4,6,7,9,11,17} Furthermore, the presence of subclinical disease or a history of illness could induce weight loss and increase the risk of death.^{1–4,8,9,11,14,15,17,18} To eliminate any effect of medical history, we excluded participants with a history of cancer, myocardial infarction, or stroke, and adjusted

for weight change since age 20 years, history of kidney disease, and history of liver disease, in multivariate analysis.

Multivariate adjustment attenuated the HR estimates associated with a BMI of 27.5–29.9 or ≥30.0 kg/m² in women, but not in men. No single covariate resulted in significant attenuation, although an increase in body weight of 5 kg or more since age 20 years, current drinking, and ≥1 hour physical activity per week attenuated hazard ratios. In contrast, a decrease in body weight of 5 kg or less since age 20 years, past drinking, being unmarried, <1 hour spent

Table 3. Baseline characteristics by BMI^a category in 14 457 women aged 40–64 years

	BMI (kg/m ²)							P value ^b
	<18.5	18.5–20.9	21.0–22.9	23.0–24.9	25.0–27.4	27.5–29.9	≥30.0	
No. of subjects	425	2135	3521	3770	2890	1227	489	
Mean age (years) (SD ^b)	54.1 (7.7)	53.5 (7.7)	53.9 (7.5)	54.4 (7.2)	55.6 (6.9)	55.4 (6.8)	54.9 (7.0)	<0.0001
Mean weight (kg) (SD)	42.1 (4.1)	47.3 (3.7)	51.4 (3.7)	55.9 (4.0)	60.4 (4.4)	65.6 (4.9)	73.0 (11.0)	<0.0001
Mean height (cm) (SD)	154.8 (7.7)	153.6 (5.4)	152.7 (5.2)	152.6 (5.2)	152.0 (5.2)	151.5 (5.5)	149.6 (8.6)	<0.0001
Mean BMI (kg/m ²) (SD)	17.5 (0.9)	20.0 (0.7)	22.0 (0.6)	24.0 (0.6)	26.1 (0.7)	28.5 (0.7)	32.6 (4.6)	<0.0001
Weight change since age 20 years (%)								
≤−10.0 kg	19.1	8.2	3.2	1.2	0.4	0.9	0.9	<0.0001
−9.9 to −5.0 kg	31.4	22.5	14.0	5.6	2.6	1.3	1.5	
−4.9 to +4.9 kg	46.8	58.6	52.7	35.6	15.0	7.1	4.3	
+5.0 to +9.9 kg	2.2	9.5	24.3	36.9	33.7	17.4	6.1	
≥+10.0 kg	0.5	1.3	5.8	20.8	48.3	73.4	87.2	
Education (%)								
Junior high school or less	49.4	42.8	45.7	49.2	55.1	58.8	63.9	<0.0001
High school	40.2	45.0	43.3	41.6	37.8	34.1	31.1	
College/university or higher	10.4	12.2	11.0	9.3	7.0	7.1	5.0	
Marital status (%)								
Married	74.4	81.7	83.6	84.8	84.1	84.1	81.1	<0.0001
Unmarried	25.6	18.3	16.4	15.2	15.9	15.9	18.9	
Smoking status (%)								
Never smoker	79.4	82.8	88.8	90.1	89.1	88.0	87.9	<0.0001
Past smoker	2.3	2.5	1.8	2.0	2.3	2.4	2.1	
Current smoker, 1–19 cigarettes/day	12.4	9.6	6.4	5.3	6.2	5.7	5.1	
Current smoker, ≥20 cigarettes/day	5.9	5.1	3.0	2.6	2.5	3.8	4.9	
Alcohol drinking (%)								
Never drinker	66.9	64.7	68.4	68.0	68.2	69.2	64.8	0.0002
Past drinker	6.4	5.1	3.4	3.6	4.4	5.6	8.5	
Current drinker	26.7	30.2	28.3	28.3	27.4	25.2	26.7	
Time spent walking (%)								
≥1 hour/day	41.0	47.5	46.8	47.9	45.2	39.9	39.4	<0.0001
<1 hour/day	59.0	52.5	53.2	52.1	54.8	60.1	60.6	
Sports and physical exercise (%)								
≥5 hours/week	3.6	3.5	4.4	3.7	3.9	3.0	3.6	NS ^a
3–4 hours/week	4.1	4.3	4.9	5.1	5.1	4.5	3.4	
1–2 hours/week	14.0	14.6	14.1	14.7	16.4	14.7	11.7	
<1 hour/week	78.4	77.6	76.6	76.5	74.6	77.8	81.4	
History of hypertension (%)								
Yes	10.6	11.0	15.2	20.7	28.8	35.6	41.3	<0.0001
No	89.4	89.0	84.8	79.3	71.2	64.4	58.7	
History of diabetes (%)								
Yes	3.3	3.4	3.2	3.5	4.5	4.5	6.1	0.0043
No	96.7	96.6	96.9	96.6	95.5	95.5	93.9	
History of kidney disease (%)								
Yes	6.4	5.3	3.9	3.2	2.7	3.9	4.9	<0.0001
No	93.7	94.7	96.1	96.8	97.3	96.1	95.1	
History of liver disease (%)								
Yes	5.2	3.2	3.6	3.9	3.7	4.5	5.7	NS
No	94.8	96.8	96.4	96.1	96.3	95.5	94.3	

^aBMI, body mass index; SD, standard deviation; NS, not significant.

^bP values were calculated by using the chi-square test (for categorical variables) or ANOVA (for continuous variables).

walking per day, and histories of kidney disease and liver disease significantly increased HRs in men.

Almost all previous studies agree that the excess risk of mortality due to obesity decreases with age,^{1–14,17,18} and our results accord with this. In underweight adults, the results of past studies have been inconsistent.^{1–15} Our results are in agreement with 2 of 14 studies of men,^{5,13} and 4 of 13 studies of women.^{3,4,10,14}

In Japan, Matsuo et al reported the effect of age on the association between BMI and all-cause mortality.¹⁰

Their findings agree with ours, except for underweight men. They adjusted only for age, alcohol intake, and smoking status in multivariate analysis; however, physical activity and socioeconomic status have also been identified as confounding factors for the risk of all-cause mortality.^{1–15,17,18} Although their result differ from ours for underweight men, our study was more careful in adjusting for physical activity, socioeconomic status, weight change since age 20 years, marital status, and histories of kidney disease and liver disease.

Table 4. Baseline characteristics by BMI^a category in 8477 women aged 65–79 years

	BMI (kg/m ²)							P value ^b
	<18.5	18.5–20.9	21.0–22.9	23.0–24.9	25.0–27.4	27.5–29.9	≥30.0	
No. of subjects	503	1383	1977	1906	1666	702	340	
Mean age (years) (SD ^a)	72.0 (4.3)	70.9 (4.3)	70.4 (4.2)	70.0 (4.0)	70.0 (4.0)	70.0 (4.1)	70.0 (4.0)	<0.0001
Mean weight (kg) (SD)	39.6 (4.5)	44.6 (3.7)	48.8 (3.7)	53.6 (4.0)	58.1 (4.7)	62.5 (5.2)	68.7 (12.0)	<0.0001
Mean height (cm) (SD)	151.9 (8.8)	149.5 (5.8)	148.8 (5.3)	149.3 (5.3)	149.0 (5.6)	147.8 (5.9)	144.6 (10.3)	<0.0001
Mean BMI (kg/m ²) (SD)	17.2 (1.2)	19.9 (0.7)	22.0 (0.6)	24.0 (0.6)	26.1 (0.7)	28.6 (0.7)	33.0 (5.6)	<0.0001
Weight change since age 20 years (%)								
≤−10.0 kg	41.9	22.7	12.0	4.5	3.3	1.5	1.3	<0.0001
−9.9 to −5.0 kg	33.3	35.1	25.4	16.1	8.7	4.5	2.6	
−4.9 to +4.9 kg	22.7	36.9	46.3	39.0	21.6	15.3	6.2	
+5.0 to +9.9 kg	1.8	4.9	12.7	26.6	29.9	22.7	13.1	
≥+10.0 kg	0.2	0.3	3.6	13.8	36.5	56.0	76.8	
Education (%)								
Junior high school or less	65.7	68.8	68.4	67.1	72.6	75.2	82.1	<0.0001
High school	28.6	26.4	25.0	26.2	22.1	19.2	14.7	
College/university or higher	5.7	4.8	6.6	6.7	5.2	5.6	3.3	
Marital status (%)								
Married	59.8	61.7	62.9	62.8	63.9	65.5	62.0	NS ^a
Unmarried	40.2	38.4	37.1	37.2	36.1	34.5	38.0	
Smoking status (%)								
Never smoker	84.1	90.0	90.0	91.1	92.3	91.4	89.8	0.0016
Past smoker	4.1	2.6	3.0	3.6	2.9	3.0	5.1	
Current smoker, 1–19 cigarettes/day	10.5	6.3	6.1	4.8	3.9	4.7	3.9	
Current smoker, ≥20 cigarettes/day	1.3	1.1	1.0	0.5	0.8	1.0	1.2	
Alcohol drinking (%)								
Never drinker	82.0	81.1	81.8	81.4	82.0	78.3	80.4	NS
Past drinker	4.6	4.5	4.9	4.4	3.4	4.8	5.8	
Current drinker	13.5	14.4	13.3	14.2	14.6	16.9	13.8	
Time spent walking (%)								
≥1 hour/day	34.3	40.4	39.8	38.7	35.5	34.7	28.5	0.0002
<1 hour/day	65.7	59.6	60.2	61.3	64.5	65.3	71.5	
Sports and physical exercise (%)								
≥5 hours/week	4.6	8.4	6.9	9.2	8.5	7.2	9.7	0.0003
3–4 hours/week	6.7	7.6	8.5	8.1	8.1	7.8	6.6	
1–2 hours/week	12.7	14.9	19.3	19.4	18.2	18.1	13.1	
<1 hour/week	76.0	69.0	65.3	63.3	65.2	66.9	70.7	
History of hypertension (%)								
Yes	24.7	29.9	35.0	39.8	45.8	50.7	54.7	<0.0001
No	75.4	70.1	65.0	60.2	54.2	49.3	45.3	
History of diabetes (%)								
Yes	5.8	6.1	8.7	7.5	8.2	9.0	12.9	0.0004
No	94.2	93.9	91.4	92.6	91.8	91.0	87.1	
History of kidney disease (%)								
Yes	4.4	4.3	4.5	4.0	5.0	4.4	2.1	NS
No	95.6	95.7	95.5	96.0	95.0	95.6	97.9	
History of liver disease (%)								
Yes	4.0	4.8	5.5	5.0	3.8	4.6	6.2	NS
No	96.0	95.2	94.5	95.0	96.2	95.4	93.8	

^aBMI, body mass index; SD, standard deviation; NS, not significant.

^bP values were calculated by using the chi-square test (for categorical variables) or ANOVA (for continuous variables).

Development of measures to address underweight has been slower than for obesity. However, Grabowski et al and Sergi et al showed that a low BMI in elderly adults was a predictor of mortality.^{26,27} Okoro et al found that underweight was associated with subsequent disability in elderly adults.²⁸ Our study also found that underweight is associated with a high mortality risk in elderly men and women, irrespective of age group.

A major strength of the present study was that the participants were recruited from the general Japanese

population. According to the Global Database on Body Mass Index of the WHO, the prevalence of underweight participants is higher in Japan (10%–20%) than in Western populations (0%–5%). Therefore, the Japanese population is one of the best in which to examine the excess risk of mortality due to underweight.

Several limitations of our study should be considered. First, although BMI has been accepted as satisfactory index of underweight and obesity, it cannot be used to identify distributions of fat and muscle tissue. Second, we used self-

Table 5. HRs^a and 95% CIs^a of all-cause mortality in 21 038 men by BMI^a category, stratified by age group

	BMI						
	<18.5	18.5–20.9	21.0–22.9	23.0–24.9	25.0–27.4	27.5–29.9	≥30.0
Total							
No.	732	3677	5617	5657	3726	1213	416
Person-years	6282	35 339	55 681	57 157	37 954	12 484	4162
No. of deaths	270	805	1004	861	513	165	67
Mortality rate ^b	43.0	22.8	18.0	15.1	13.5	13.2	16.1
Age-smoking-adjusted HRs	2.78	1.49	1.19	1.00	0.90	0.90	1.11
	(2.42–3.18)	(1.35–1.64)	(1.09–1.22)	(reference)	(0.81–1.00)	(0.76–1.06)	(0.87–1.43)
Multivariate HRs1 ^c	1.42	1.10	1.04	1.00	1.01	1.10	1.44
	(1.23–1.65)	(0.99–1.22)	(0.95–1.14)	(reference)	(0.90–1.13)	(0.92–1.31)	(1.11–1.87)
Multivariate HRs2 ^d	1.52	1.14	1.06	1.00	1.00	1.06	1.39
	(1.31–1.76)	(1.03–1.26)	(0.96–1.16)	(reference)	(0.89–1.12)	(0.89–1.26)	(1.07–1.81)
Multivariate HRs3 ^e	1.35	1.06	1.01	1.00	0.99	1.05	1.42
	(1.15–1.59)	(0.95–1.19)	(0.92–1.12)	(reference)	(0.88–1.12)	(0.87–1.26)	(1.07–1.88)
40–64 y							
No.	310	2159	3591	3852	2637	903	312
Person-years	3053	21 992	36 885	40 026	27 421	9425	3221
No. of deaths	47	224	340	305	211	75	33
Mortality rate ^b	15.4	10.2	9.2	7.6	7.7	8.0	10.2
Age-smoking-adjusted HRs	1.76	1.25	1.17	1.00	1.07	1.17	1.54
	(1.29–2.39)	(1.05–1.49)	(1.01–1.37)	(reference)	(0.90–1.28)	(0.91–1.51)	(1.08–2.21)
Multivariate HRs1 ^c	1.26	1.07	1.11	1.00	1.14	1.27	1.71
	(0.92–1.73)	(0.89–1.28)	(0.95–1.30)	(reference)	(0.95–1.37)	(0.97–1.66)	(1.17–2.50)
Multivariate HRs2 ^d	1.32	1.09	1.12	1.00	1.12	1.22	1.64
	(0.96–1.82)	(0.91–1.31)	(0.96–1.32)	(reference)	(0.93–1.35)	(0.93–1.60)	(1.12–2.40)
Multivariate HRs3 ^e	1.24	1.11	1.13	1.00	1.16	1.20	1.62
	(0.87–1.78)	(0.92–1.35)	(0.95–1.34)	(reference)	(0.96–1.41)	(0.90–1.61)	(1.07–2.45)
65–79 y							
No.	422	1518	2026	1805	1089	310	104
Person-years	3229	13 347	18 796	17 131	10 533	3059	941
No. of deaths	223	581	664	556	302	90	34
Mortality rate ^b	69.1	43.5	35.3	32.5	28.7	29.4	36.1
Age-smoking-adjusted HRs	1.88	1.26	1.06	1.00	0.91	0.96	1.21
	(1.61–2.20)	(1.12–1.41)	(0.94–1.18)	(reference)	(0.79–1.05)	(0.77–1.19)	(0.86–1.71)
Multivariate HRs1 ^c	1.49	1.11	1.01	1.00	0.94	1.01	1.25
	(1.26–1.76)	(0.98–1.26)	(0.90–1.14)	(reference)	(0.81–1.09)	(0.80–1.27)	(0.87–1.80)
Multivariate HRs2 ^d	1.59	1.16	1.03	1.00	0.93	0.97	1.23
	(1.35–1.89)	(1.03–1.32)	(0.92–1.16)	(reference)	(0.80–1.07)	(0.77–1.23)	(0.86–1.76)
Multivariate HRs3 ^e	1.48	1.09	0.98	1.00	0.88	0.91	1.22
	(1.23–1.78)	(0.96–1.25)	(0.87–1.11)	(reference)	(0.76–1.03)	(0.71–1.17)	(0.83–1.79)

^aHR, hazard ratio; CI, confidence interval; BMI, body mass index.

^bMortality rate was defined as number of deaths per 1000 person-years.

^cMultivariate HRs1 were adjusted for age in 5-year categories; weight change since age 20 years (loss of 10.0 kg or more, loss of 5.0–9.9 kg, change of less than ±5.0 kg, gain of 5.0–9.9 kg, or gain of 10.0 kg or more); education (junior high school or less, high school, or college/university or higher); marital status (married or unmarried); cigarette smoking (never smoker, past smoker, current smoker consuming 1–19 cigarettes per day, or current smoker consuming at least 20 cigarettes per day); alcohol drinking (never drinker, past drinker, or current drinker); time spent walking per day (less than 1 hour, or 1 hour or longer); sports and physical exercise time per week (less than 1 hour, 1–2 hours, 3–4 hours, or 5 hours or longer); history of kidney disease (yes or no); history of liver disease (yes or no).

^dMultivariate HRs2 were further adjusted for history of hypertension (yes or no) and history of diabetes (yes or no).

^eMultivariate HRs3 excluded from multivariate HRs2 the 473 men who died within the 2 years after baseline.

reported BMI at baseline. Niedhammer et al showed that there is a systematic bias in self-reported weight and height.²⁹ However, we previously evaluated the validity of self-reported BMI, and demonstrated a high correlation and appropriate agreement between self-reported BMI and measured BMI in a subsample of 14 883 participants ($r = 0.88$, $\kappa = 0.72$). We consider this bias to be a nondifferential misclassification that is not dependent upon all-cause death. This misclassification weakens the true association toward the

null. Third, as a result of stratification by age group, there was a possibility of beta error because of inadequate numbers of participants and events. Finally, there is a possibility of residual confounding by physical activity.

In summary, obesity increases mortality risk in middle-aged men, whereas underweight, rather than obesity, is associated with high mortality risk in elderly men. In women, obesity increases mortality risk in middle age, and underweight increases mortality risk irrespective of age. Although there

Table 6. HRs^a and 95% CIs^a of all-cause mortality in 22 934 women by BMI^a category, stratified by age group

	BMI						
	<18.5	18.5–20.9	21.0–22.9	23.0–24.9	25.0–27.4	27.5–29.9	≥30.0
Total							
No.	928	3518	5498	5676	4556	1929	829
Person-years	9011	34 782	55 716	57 537	46 281	19 477	8313
No. of deaths	174	371	451	415	357	159	95
Mortality rate ^b	19.3	10.7	8.1	7.2	7.7	8.2	11.4
Age-smoking-adjusted HRs	2.66	1.48	1.12	1.00	1.07	1.13	1.59
	(2.23–3.18)	(1.28–1.70)	(0.98–1.28)	(reference)	(0.93–1.23)	(0.94–1.36)	(1.27–1.99)
Multivariate HRs1 ^c	1.49	1.15	0.99	1.00	1.03	1.07	1.33
	(1.24–1.80)	(0.99–1.33)	(0.87–1.14)	(reference)	(0.89–1.19)	(0.89–1.30)	(1.05–1.69)
Multivariate HRs2 ^d	1.58	1.19	1.01	1.00	1.00	1.04	1.24
	(1.31–1.91)	(1.03–1.38)	(0.88–1.15)	(reference)	(0.87–1.16)	(0.85–1.26)	(0.97–1.57)
Multivariate HRs3 ^e	1.44	1.18	1.02	1.00	1.06	1.09	1.37
	(1.17–1.78)	(1.01–1.38)	(0.88–1.18)	(reference)	(0.91–1.24)	(0.88–1.34)	(1.07–1.77)
40–64 y							
No.	425	2135	3521	3770	2890	1227	489
Person-years	4416	21 274	35 734	38 262	29 435	12 484	4999
No. of deaths	32	92	137	128	104	49	28
Mortality rate ^b	7.2	4.3	3.8	3.3	3.5	3.9	5.6
Age-smoking-adjusted HRs	2.10	1.30	1.16	1.00	0.99	1.11	1.59
	(1.43–3.10)	(0.99–1.70)	(0.91–1.48)	(reference)	(0.76–1.28)	(0.80–1.54)	(1.06–2.39)
Multivariate HRs1 ^c	1.46	1.10	1.09	1.00	1.00	1.06	1.47
	(0.96–2.22)	(0.82–1.47)	(0.85–1.40)	(reference)	(0.77–1.31)	(0.75–1.51)	(0.94–2.27)
Multivariate HRs2 ^d	1.55	1.14	1.10	1.00	0.98	1.01	1.38
	(1.02–2.36)	(0.85–1.52)	(0.86–1.41)	(reference)	(0.75–1.28)	(0.71–1.43)	(0.89–2.14)
Multivariate HRs3 ^e	1.78	1.36	1.21	1.00	1.02	0.99	1.32
	(1.13–2.81)	(1.00–1.86)	(0.92–1.59)	(reference)	(0.76–1.36)	(0.68–1.45)	(0.82–2.15)
65–79 y							
No.	503	1383	1977	1906	1666	702	340
Person-years	4595	13 508	19 982	19 275	16 845	6994	3314
No. of deaths	142	279	314	287	253	110	67
Mortality rate ^b	30.9	20.7	15.7	14.9	15.0	15.7	20.2
Age-smoking-adjusted HRs	1.67	1.26	1.00	1.00	1.02	1.07	1.33
	(1.37–2.05)	(1.07–1.48)	(0.85–1.17)	(reference)	(0.86–1.21)	(0.86–1.34)	(1.02–1.73)
Multivariate HRs1 ^c	1.47	1.14	0.95	1.00	1.04	1.07	1.26
	(1.19–1.82)	(0.96–1.36)	(0.81–1.12)	(reference)	(0.87–1.24)	(0.85–1.35)	(0.95–1.68)
Multivariate HRs2 ^d	1.56	1.19	0.96	1.00	1.01	1.04	1.17
	(1.26–1.93)	(1.00–1.41)	(0.82–1.13)	(reference)	(0.85–1.21)	(0.83–1.31)	(0.88–1.55)
Multivariate HRs3 ^e	1.45	1.17	0.96	1.00	1.04	1.07	1.24
	(1.15–1.83)	(0.97–1.40)	(0.81–1.15)	(reference)	(0.86–1.25)	(0.83–1.37)	(0.92–1.68)

^aHR, hazard ratio; CI, confidence interval; BMI, body mass index.

^bMortality rate was defined as number of deaths per 1000 person-years.

^cMultivariate HRs1 were adjusted for age in 5-year categories; weight change since age 20 years (loss of 10.0 kg or more, loss of 5.0–9.9 kg, change of less than ±5.0 kg, gain of 5.0–9.9 kg, or gain of 10.0 kg or more); education (junior high school or less, high school, or college/university or higher); marital status (married or unmarried); cigarette smoking (never smoker, past smoker, current smoker consuming 1–19 cigarettes per day, or current smoker consuming at least 20 cigarettes per day); alcohol drinking (never drinker, past drinker, or current drinker); time spent walking per day (less than 1 hour, or 1 hour or longer); sports and physical exercise time per week (less than 1 hour, 1–2 hours, 3–4 hours, or 5 hours or longer); history of kidney disease (yes or no); history of liver disease (yes or no).

^dMultivariate HRs2 were further adjusted for history of hypertension (yes or no) and history of diabetes (yes or no).

^eMultivariate HRs3 excluded from multivariate HRs2 the 266 women who died within the 2 years after baseline.

was no significant interaction by age group or sex, the mortality risks associated with underweight and obesity may nevertheless be dependent on sex and age group.

ACKNOWLEDGMENTS


This study was supported by a Health Sciences Research Grant for Health Services (H18-Choju-Ippan-014, H19-Seisaku-Ippan-026, H18-Junkankitou [Seisyu]-Ippan-012,

H20-Junkankitou [Seisyu]-Ippan-013), Ministry of Health, Labour and Welfare, Japan. Masato Nagai is a recipient of a Research Fellowship of the Japan Society for the Promotion of Science for Young Scientists.

REFERENCES

1. Stevens J, Cai J, Pamuk ER, Williamson DF, Thun MJ, Wood JL. The effect of age on the association between body-mass

- index and mortality. *N Engl J Med.* 1998;338:1–7.
2. Jee SH, Sull JW, Park J, Lee SY, Ohrr H, Guallar E, et al. Body-mass index and mortality in Korean men and women. *N Engl J Med.* 2006;355:779–87.
 3. Calle EE, Thun MJ, Petrelli JM, Rodriguez C, Heath CW Jr. Body-mass index and mortality in a prospective cohort of U.S. adults. *N Engl J Med.* 1999;341:1097–105.
 4. Adams KF, Schatzkin A, Harris TB, Kipnis V, Mouw T, Ballard-Barbash R, et al. Overweight, obesity, and mortality in a large prospective cohort of persons 50 to 71 years old. *N Engl J Med.* 2006;355:763–78.
 5. Flegal KM, Graubard BI, Williamson DF, Gail MH. Excess deaths associated with underweight, overweight, and obesity. *JAMA.* 2005;293:1861–7.
 6. Gu D, He J, Duan X, Reynolds K, Wu X, Chen J, et al. Body weight and mortality among men and women in China. *JAMA.* 2006;295:776–83.
 7. Pednekar MS, Hakama M, Hebert JR, Gupta PC. Association of body mass index with all- and cause-specific mortality: findings from a prospective cohort study in Mumbai (Bombay), India. *Int J Epidemiol.* 2008;37:524–35.
 8. Rissanen A, Heliövaara M, Knekt P, Aromaa A, Reunanen A, Maatela J. Weight and mortality in Finnish men. *J Clin Epidemiol.* 1989;42:781–9.
 9. Dorn JM, Schisterman EF, Winkelstein W Jr, Trevisan M. Body mass index and mortality in a general population sample of men and women. The Buffalo Health Study. *Am J Epidemiol.* 1997;146:919–31.
 10. Matsuo T, Sairenchi T, Iso H, Irie F, Tanaka K, Fukasawa N, et al. Age- and gender-specific BMI in terms of the lowest mortality in Japanese general population. *Obesity (Silver Spring).* 2008;16:2348–55.
 11. Pischon T, Boeing H, Hoffmann K, Bergmann M, Schulze MB, Overvad K, et al. General and abdominal adiposity and risk of death in Europe. *N Engl J Med.* 2008;359:2105–20.
 12. Klenk J, Nagel G, Ulmer H, Strasak A, Concin H, Diem G, et al; The VHM&PP Study Group. Body mass index and mortality: results of a cohort of 184,697 adults in Austria. *Eur J Epidemiol.* 2009;24:83–91.
 13. Janssen I. Morbidity and mortality risk associated with an overweight BMI in older men and women. *Obesity (Silver Spring).* 2007;15:1827–40.
 14. Rissanen A, Knekt P, Heliövaara M, Aromaa A, Reunanen A, Maatela J. Weight and mortality in Finnish women. *J Clin Epidemiol.* 1991;44:787–95.
 15. Ajani UA, Lotufo PA, Gaziano JM, Lee IM, Spelsberg A, Buring JE, et al. Body mass index and mortality among US male physicians. *Ann Epidemiol.* 2004;14:731–9.
 16. Bender R, Jöckel KH, Trautner C, Spraul M, Berger M. Effect of age on excess mortality in obesity. *JAMA.* 1999;281:1498–504.
 17. Baik I, Ascherio A, Rimm EB, Giovannucci E, Spiegelman D, Stampfer MJ, et al. Adiposity and mortality in men. *Am J Epidemiol.* 2000;152:264–71.
 18. Prospective Studies Collaboration, Whitlock G, Lewington S, Sherliker P, Clarke R, Emberson J, Halsey J, et al. Body-mass index and cause-specific mortality in 900 000 adults: collaborative analyses of 57 prospective studies. *Lancet.* 2009;373:1083–96.
 19. Low S, Chin MC, Ma S, Heng D, Deurenberg-Yap M. Rationale for redefining obesity in Asians. *Ann Acad Med Singapore.* 2009;38:66–9.
 20. Tsuji I, Nishino Y, Ohkubo T, Kuwahara A, Ogawa K, Watanabe Y, et al. A prospective cohort study on National Health Insurance beneficiaries in Ohsaki, Miyagi Prefecture, Japan: study design, profiles of the subjects and medical cost during the first year. *J Epidemiol.* 1998;8:258–63.
 21. Kuriyama S, Shimazu T, Ohmori K, Kikuchi N, Nakaya N, Nishino Y, et al. Green tea consumption and mortality due to cardiovascular disease, cancer, and all causes in Japan: the Ohsaki study. *JAMA.* 2006;296:1255–65.
 22. Funada S, Shimazu T, Kakizaki M, Kuriyama S, Sato Y, Matsuda-Ohmori K, et al. Body mass index and cardiovascular disease mortality in Japan: the Ohsaki Study. *Prev Med.* 2008;47:66–70.
 23. World Health Organization. The use and interpretation of anthropometry: report of a WHO expert committee. *World Health Organ Tech Rep Ser.* 1995;854:312–409.
 24. SAS Institute Inc., 2004 SAS Institute Inc. SAS/STAT 9.1 User's Guide. Cary, NC: SAS Institute Inc; 2004.
 25. World Health Organization. Men, ageing and health. Achieving health across the life span. Geneva: World Health Organization; 2001.
 26. Grabowski DC, Ellis JE. High body mass index does not predict mortality in older people: analysis of the Longitudinal Study of Aging. *J Am Geriatr Soc.* 2001;49:968–79.
 27. Sergi G, Perissinotto E, Pisent C, Buja A, Maggi S, Coin A, et al; ILSA Working Group. An adequate threshold for body mass index to detect underweight condition in elderly persons: the Italian Longitudinal Study on Aging (ILSA). *J Gerontol A Biol Sci Med Sci.* 2005;60:866–71.
 28. Okoro CA, Hootman JM, Strine TW, Balluz LS, Mokdad AH. Disability, arthritis, and body weight among adults 45 years and older. *Obes Res.* 2004;12:854–61.
 29. Niedhammer I, Bugel I, Bonenfant S, Goldberg M, Leclerc A. Validity of self-reported weight and height in the French GAZEL cohort. *Int J Obes Relat Metab Disord.* 2000;24:1111–8.



Participation in health check-ups and mortality using propensity score matched cohort analyses

Atsushi Hozawa^{a,b,*}, Shinichi Kuriyama^a, Ikue Watanabe^a, Masako Kakizaki^a, Kaori Ohmori-Matsuda^a, Toshimasa Sone^a, Masato Nagai^a, Yumi Sugawara^a, Akemi Nitta^a, Qiang Li^a, Takayoshi Ohkubo^c, Yoshitaka Murakami^d, Ichiro Tsuji^a

^a Division of Epidemiology, Department of Public Health and Forensic Medicine, Tohoku University Graduate School of Medicine, Sendai, Japan

^b Department of Public Health, Yamagata University Graduate School of Medical Science, Yamagata, Japan

^c Department of Health Science, Shiga University of Medical Science, Otsu, Japan

^d Department of Medical Statistics, Shiga University of Medical Science, Otsu, Japan

ARTICLE INFO

Available online 7 September 2010

Keywords:

Mass screening
Lifestyle
Health check-ups
Prospective studies
Cardiovascular disease
Mortality

ABSTRACT

Objective. All Japanese aged ≥ 40 years are eligible for free annual health check-ups including blood pressure and cholesterol measurements. It is well known that health check-up screenees are more likely to have healthy lifestyles and better health conditions than non-screenees. Therefore, controlling these factors is required to investigate whether screenees have a lower mortality risk than non-screenees independent of their lifestyles or health conditions.

Methods. We followed 48,775 Japanese National Health Insurance beneficiaries aged 40–79 years since 1994 for 11 years. We used Cox proportional hazard models adjusted for possible confounding factors. We also performed propensity for use of the health check-up matched cohort analyses.

Results. Compared to non-screenees, multiple-adjusted hazard ratios (95% confidence intervals) for all-cause and cardiovascular disease mortality among screenees were 0.74 (0.62–0.88) and 0.65 (0.44–0.95) for men and 0.69 (0.52–0.91) and 0.61 (0.36–1.04) for women, respectively. These relations were also observed when we used propensity matched cohort analyses.

Conclusion. This is the first study to show that mortality rates are lower among screenees than non-screenees in Japanese health check-ups when propensity matched cohort analyses were used for adjusting confounding factors. Further prospective studies, including randomized controlled trials, are required to confirm whether screening lowers mortality.

© 2010 Elsevier Inc. All rights reserved.

Introduction

Japanese have the longest life span in the world, and this has increased remarkably (Statistics and Information Department, Minister's Secretariat, Ministry of Health, Labour and Welfare, 2007). The reduction of stroke mortality in Japan is one explanation for the current status of Japanese longevity (Statistics and Information Department, Minister's Secretariat, Ministry of Health, Labour and Welfare, 2008).

One strategy to reduce stroke in Japan has been the national health check-up system. The Health Services for the Elderly Act 1982 provides for six health services including general health checks (Tatara et al., 1991). These services are inexpensive or free for all who reside anywhere in Japan and hold a resident card, who are aged ≥ 40 years, and who do not have access to other health examinations in the

workplace. Thus, all Japanese adults aged ≥ 40 years can attend annual health check-ups. Height, weight, blood pressure (BP), lipids, glucose, liver function, and renal function are measured during these health check-ups (Tatara et al., 1991). Specific cancer screenings are not included in the health check-up examination. After the check-ups, depending on the results, the screenees can receive health education in a group setting or ask for individual health counseling (Tatara et al., 1991).

The U.S. Preventive Task Force (USPSTF) recommends regular monitoring for high BP, lipid disorders, and obesity in adults (U.S. Preventive Services Task Force, 2009), and these risk factors are included in the national health check system. Thus, this Japanese system of health check-up coupled with follow-up treatment of abnormalities should contribute to lowering stroke mortality among Japanese.

However, estimating the effect of health check-ups on mortality is difficult because no randomized trial exists, and participants who attend annual health check-ups are more likely to have healthy lifestyles or health conditions than those who do not.

Although previous studies did not adjust for these lifestyles or health conditions (Iwasaki et al., 2006; Lannerstad et al., 1997;

* Corresponding author. Department of Public Health, Yamagata University Graduate School of Medical Science, 2-2-2 Iidamachi, 990-9585, Yamagata, Japan.
E-mail address: hozawa-thk@umin.ac.jp (A. Hozawa).

Wilhelmsen et al., 1986), recent Japanese studies reported the relationship between health check-up and risk of mortality adjusted for possible confounding factors (Khan et al., 2004; Ikeda et al., 2005). Khan et al. (2004) investigated 3185 men and women for about 15 years and found an inverse relationship between screening and all-cause mortality. Ikeda et al. (2005) also investigated 68,825 men and women for 10 years and reported an inverse relationship between participation in screening and all-cause and cause-specific mortality only in women. However, neither of the studies adjusted for information on diet, self-rated health, nor physical function, which should also be different between participants who used health check-up (screenees) and those who did not (non-screenees).

The Ohsaki National Health Insurance (NHI) Cohort Study investigated lifestyles of the general population with very high response rate and determined whether participants used the annual health check system in 1995 (Hozawa et al., 2009; Kuriyama et al., 2006; Ohmori-Matsuda et al., 2007; Sone et al., 2008; Tsuji et al., 1998). Because the Ohsaki study had several detailed lifestyle information than the previous report, such as diet, physical function, or self-rated health, we could control for these important confounding factors.

Recently, propensity scores have been widely used in cardiovascular research (D'Agostino, 2007). This approach was used to reduce bias in observational studies. Therefore, in this study, we also used this approach to investigate the relationship between health check-up and mortality.

The present study compares general and specific mortality rates among screenees and non-screenees, after 11 years of follow-up, controlling lifestyle, and other risk factors.

Methods

Study cohort

Details of the Ohsaki NHI Cohort Study have been described elsewhere (Hozawa et al., 2009; Kuriyama et al., 2006; Ohmori-Matsuda et al., 2007; Sone et al., 2008; Tsuji et al., 1998). In brief, we sent a self-administered questionnaire between October and December 1994 to all NHI beneficiaries aged 40–79 years living in the catchment area of the Ohsaki Public Health Center. The NHI in Japan is used by farmers, self-employed individuals,

pensioners, and their dependents. Because these populations usually do not have access to any other health examinations, they could all be considered targets of health check-ups. Of the 54,996 eligible individuals mailed the questionnaire, 52,029 (95%) responded. To ascertain the date of and reason for withdrawal from the NHI, we started the prospective collection of NHI withdrawal history files on January 1, 1995. We excluded 774 participants who had withdrawn from the NHI before the baseline questionnaire survey. Thus, 51,255 participants ultimately formed the study cohort. Among the participants of the Ohsaki NHI Cohort Study, 16,515 (32.2%) had undergone an annual health check between April and December 1995. In Japan, calendar year started in April, and health check-ups were usually carried out from April to December. To exclude bias from participants who died before being able to attend a health check, we further excluded 2480 participants who died or moved from the area before the health check ended on December 6, 1995 (Fig. 1). Thus, our study participants comprised of 48,775 men and women (men, 23,451; women, 25,324). We defined 15,985 of them as screenees and 32,790 as non-screenees of the health check in 1995.

The ethics committee of Tohoku University School of Medicine reviewed and approved the study protocol. We considered that returning signed, self-administered questionnaires implied written consent to participate in this study.

Exposure data

The questionnaire included items about cigarette smoking, alcohol consumption, self-reported weight and height, histories of diseases, participation in sports or exercise, time spent walking, sense that life is worth living (*ikigai*) (Sone et al., 2008), self-rated health, physical function, the frequency of recent average consumption of 4 beverages (green tea, oolong tea, black tea, and coffee) and 36 food items, and cancer screening within the previous 5 years. Validation of the questionnaire for time spent walking (Tsubono et al., 2002) or the food frequency questionnaire (Ogawa et al., 2003) has already been reported. The physical function status was assessed using the six-item measure of the Medical Outcomes Study (MOS) Short-form General Health Survey (Stewart et al., 1988): able to perform vigorous activity (MOS scores of 5 or 6); capable of moderate but not vigorous activity (MOS scores of 2–4); and capable of only low physical activity (MOS scores of 0 or 1). We defined MOS 0 or 1 as limited physical function.

Follow-up

The end points among the cohort of 48,775 were all-cause mortality and cause-specific mortality from December 6, 1995, to December 31, 2006. The




Fig. 1. Flowchart of the study participants.

details of follow-up and certification of death causes have been described in previous reports (Hozawa et al., 2009; Kuriyama et al., 2006; Ohmori-Matsuda et al., 2007; Sone et al., 2008).

Statistical analysis

To compare the baseline characteristics, we used the Student's *t*-test and the chi-squared test for continuous and categorical variables, respectively. We used the Cox proportional hazards regression analysis with age as the time scale, using left and right censoring, to calculate the hazard ratios (HRs) and 95% confidence intervals (CIs) of all-cause and cause-specific mortality (Korn et al., 1997). All data were analyzed using the SAS statistical software, version 9.1 (SAS Institute Inc., Cary, NC, USA). Non-screenees were considered as the reference category. Because smoking habit or alcohol consumption is largely different between men and women in Japan, we separately analyzed men and women in all analyses. One exception was the propensity score matched cohort analysis. Because the pairs were matched both by sex and propensity score, we showed men only, women only, and men and women combined information in this analysis.

We used two models to estimate the difference in mortality between screenees and non-screenees. Model 1 used the following lifestyles and conditions as potential confounders: body mass index (BMI); participation in sports or exercise; time spent walking; sense of life worth living (*ikigai*); self-rated health; physical function; history of any extant diseases; smoking status; alcohol consumption; daily consumption of meat, fish, green and yellow vegetables, and green tea; job status; education; residential area; and history of screening for lung, gastric, colon, breast, and uterine cancer. To minimize the possibility that participants did not attend health check-ups because of illness or moving restriction, we used Model 2: excluding participants with any extant diseases, those with limited physical function, and those who died within the first 3 years of follow-up.

For the propensity score matched cohort analyses, we used participants who did not have any extant diseases, those without limited physical function, and those who did not die within the first 3 years of follow-up ($N=7732$ for men and $N=7435$ for women). Among the participants, we calculated the propensity score, which was determined without regard to outcome, using multivariable logistic regression analysis using possible confounding factors in Model 2 together with age categories at baseline (40–49, 50–59, 60–69, and 70–79). *C* statistics of these models were 0.79 for men and 0.78 for women. Then, we made propensity score categories for every 5% (Table 1). The number of pairs was defined as the minimum of the number of screenees and non-screenees in each propensity score category. We randomly selected the participants from each category. Finally, we sorted screenees and non-screenees according to the propensity score. As a result, 1800 and 2087 propensity matched pairs (1:1) for men and women were selected. The relation between the use of the health check-up and all-cause and cause-specific mortality was calculated using the Cox proportional hazard model stratified on the propensity score matched pairs. All reported *p* values are 2-tailed.

Results

Mean ages (standard deviation, SD) of screenees and non-screenees were 61.7 (10.0) and 59.6 (10.7) years for men and 61.4 (8.9) and 62.0 (10.5) years for women, respectively. Mean BMIs (SD) of screenees and non-screenees were 23.4 (2.9) and 23.3 (3.2) kg/m² for men and 23.8 (3.2) and 23.8 (3.7) kg/m² for women, respectively. The proportions who never smoked, who often consumed green and yellow vegetables, and who had been screened for cancer were much higher among screenees than non-screenees (Table 2). The proportions of participants who considered that life was worth living, who had unlimited physical functions, or who had good/excellent self-rated health were also higher among screenees than non-screenees. Table 2 also shows the proportion with a history of severe diseases such as stroke, myocardial infarction, or cancer was higher among non-screenees than screenees. Thus, screenees were in better health than non-screenees, probably because, as we have been able to check using other analyses, they tended to have healthier lifestyles.

During 11 years of follow-up, the all-cause mortality rate (per 1000 person-years) was lower among screenees (men, 15.8; women, 5.7)

Table 1
Selection of pair for propensity matched cohort analyses. The Ohsaki Cohort Study.

Propensity score	Men			Women		
	Non-screenees	Screenees	Pair selected	Non-screenees	Screenees	Pair selected
0–0.049	423	17	17	74	3	3
0.05–0.099	1093	83	83	445	37	37
0.1–0.149	994	148	148	492	76	76
0.15–0.199	759	154	154	508	117	117
0.2–0.249	508	157	157	555	155	155
0.25–0.299	348	125	125	465	152	152
0.3–0.349	264	108	108	391	166	166
0.35–0.399	192	143	143	270	146	146
0.4–0.449	177	136	136	211	197	197
0.45–0.499	175	149	149	199	203	199
0.5–0.549	152	163	152	176	213	176
0.55–0.599	129	181	129	178	199	178
0.6–0.649	108	183	108	126	235	126
0.65–0.699	84	187	84	120	217	120
0.7–0.749	69	161	69	89	267	89
0.75–0.799	29	94	29	78	298	78
0.8–0.849	9	26	9	57	233	57
0.85–0.899	0	2	0	15	70	15
0.9–	0	2	0	0	2	0
Total	5513	2219	1800	4449	2986	2087

We used participants who did not have any extant diseases, those without limited physical function, and those who did not die within the first 3 years of follow-up ($N=7732$ for men and $N=7435$ for women).

Number of pair was defined as the minimum of the number of screenees and non-screenees in each propensity score category.

than that among non-screenees (men, 23.8; women, 14.0). This tendency was similar for cause-specific mortality. Both male and female screenees had a reduced multiple-adjusted HR of death due to all-causes, CVD, cancer, and other causes (Model 1) (Table 3). For men, if we used more detailed categories of current smoking (≥ 20 cigarettes/day and < 20 cigarettes/day) and current drinking (≥ 69 g of ethanol/day, 46–68.9 g of ethanol /day, 23–45.9 g of ethanol/day, and 0–22.9 g of ethanol/day), the HRs were unchanged (data not shown).

When we excluded participants with any extant diseases, those with restricted movement, and those who died during the first 3 years of follow-up (Model 2), this approach did not alter the point estimate of the relative risk substantially.

For the propensity matched cohort analyses, significant differences were not found in the characteristics between screenees and non-screenees at baseline when the propensity matched cohort analyses were made (data not shown). Screenees consistently showed a reduced HR of all-cause and cause-specific mortality when we used propensity matched cohort analysis (Table 4). When we combined men and women in the same model, screenees had a significantly reduced HR of death due to all-cause and cause-specific mortality, except for cancer mortality.

Discussion

We found that screenees of a health check-up in 1995 had healthier lifestyles than non-screenees. We also found that all-cause and cause-specific mortality risks were lower among screenees than non-screenees. The difference in mortality between screenees and non-screenees persisted, despite attenuation when adjusted for lifestyle or when participants with extant diseases were excluded. Furthermore, except for cancer mortality, propensity matched analyses also showed significantly lower all-cause and cause-specific mortality risks among screenees than non-screenees, even though propensity matched screenees and non-screenees had identical baseline characteristics.

The strengths of this study are the large sample cohort and the high (95%) response rate to the questionnaire. Thus, we considered our screenees and non-screenees to be representatives of the target

Table 2
Baseline characteristics of participants who used health check-up in 1995 and those who did not. The Ohsaki Cohort Study, 1994.

		Men			Women		
		Screenees	Non-screenees	P	Screenees	Non-screenees	P
N		6814	16,637		9171	16,153	
Age category	40–49 years (%)	16.9	24.4	<0.001	12.9	17.3	<0.001
	50–59 years (%)	20.1	22.1		26.3	21.6	
	60–69 years (%)	42.3	36.0		45.2	36.1	
	70–79 years (%)	20.8	17.5		15.7	25.1	
Body mass index	<18.5 kg/m ² (%)	2.9	4.1	<0.001	2.9	5.0	<0.001
	18.5–21.9 kg/m ² (%)	28.4	30.1		25.7	26.6	
	22–24.9 kg/m ² (%)	43.5	40.4		40.0	36.1	
	25–29.9 kg/m ² (%)	23.7	23.2		28.8	27.9	
	≥30 kg/m ² (%)	1.6	2.2		2.5	4.4	
Smoking	Current (%)	48.2	57.8	<0.001	4.9	10.5	<0.001
	Former (%)	30.5	24.8		1.7	3.2	
	Never (%)	21.3	17.4		93.4	86.3	
Alcohol drinking	Current (%)	76.1	70.3	<0.001	22.0	22.7	<0.001
	Former (%)	8.3	12.8		3.1	5.4	
	Never (%)	15.6	17.0		75.0	71.9	
Engaging in sports or exercise	≥1 hour/week (%)	29.9	35.3	<0.001	30.2	25.5	<0.001
Time spent walking	≥1 hour/day (%)	48.0	48.4	<0.001	44.2	41.4	<0.001
Sense of life worth living	Yes (%)	68.1	58.4	<0.001	60.3	53.7	<0.001
Self-rated health	Excellent (%)	9.2	10.7	<0.001	7.9	7.5	<0.001
	Good (%)	64.2	53.4		60.0	49.7	
	Fair (%)	13.6	14.9		14.1	15.8	
	Poor (%)	10.5	14.3		14.8	18.9	
	Bad (%)	2.5	6.7		3.4	8.1	
Physical function	Unlimited (%)	93.6	86.6	<0.001	92.3	85.4	<0.001
Meat consumption	≥3–4 times/week (%)	26.1	27.3	<0.001	29.9	26.0	<0.001
Fish	Almost everyday (%)	39.1	35.3	<0.001	41.9	35.3	<0.001
Green and yellow vegetables	Almost everyday (%)	36.3	28.9	<0.001	48.3	39.2	<0.001
Green tea	≥5 cups/day (%)	30.5	25.5	<0.001	34.3	33.4	<0.001
Education	Until 15 years of age (%)	59.6	63.5	<0.001	55.1	60.2	<0.001
	Until 16–18 years of age (%)	32.1	29.3		36.5	32.2	
	Until ≥19 years of age (%)	8.3	7.2		8.3	7.6	
Job status	Working (%)	57.2	56.4	<0.001	35.1	30.8	<0.001
Living area	City (%)	14.8	27.3	<0.001	15.5	30.2	<0.001
Cancer screening/lung	≥1/5 years (%)	92.5	77.6	<0.001	90.8	77.0	<0.001
Cancer screening/gastric	≥1/5 years (%)	80.6	49.8	<0.001	77.3	48.6	<0.001
Cancer screening/colon	≥1/5 years (%)	57.3	28.6	<0.001	53.3	27.4	<0.001
Cancer screening/uterus	≥1/5 years (%)	–	–		71.4	43.4	<0.001
Cancer screening/breast	≥1/5 years (%)	–	–		62.3	34.9	<0.001
History of diseases	Stroke (%)	1.7	3.4	<0.001	0.7	2.3	<0.001
	Hypertension (%)	24.4	24.6	0.73	24.6	31.0	<0.001
	Myocardial infarction (%)	2.7	3.6	<0.001	1.8	2.8	<0.001
	Kidney diseases (%)	3.7	3.8	0.64	3.6	4.6	<0.001
	Liver diseases (%)	7.0	7.0	0.85	4.1	4.6	0.08
	Cholecystitis or cholelithiasis (%)	5.8	4.9	0.003	6.6	7.6	0.004
	Diabetes (%)	6.4	7.9	<0.001	3.5	7.1	<0.001
	Gastric ulcer (%)	23.5	18.6	<0.001	12.3	10.2	<0.001
	Tuberculosis (%)	5.7	4.3	<0.001	3.1	3.0	0.60
	Hearing problem (%)	5.6	4.7	0.003	3.1	4.1	<0.001
	Cataract (%)	5.7	4.7	0.002	11.2	11.8	0.14
	Arthritis (%)	6.6	6.1	0.13	12.0	13.0	0.02
	Osteoporosis (%)	1.1	1.2	0.55	6.9	7.0	0.71
	Cancer (%)	2.5	3.1	0.01	3.3	4.0	<0.01
	Blood transfusion (%)	10.8	11.6	0.09	11.8	13.5	<0.01

population in this area. Although it is unknown whether this information could be directly applicable for participants who had other insurance for employees and their dependents or who lived in other countries or other areas in Japan, our information might be important for other settings.

Several important confounding factors should be considered when comparing mortality rates between screenees and non-screenees of health check-ups. Firstly, screenees had healthier habits on important lifestyle factors such as smoking or green and yellow vegetable consumption. Secondly, we found that participants with a history of diseases were less likely to undergo the health check-up. This may be because they were feeling too bad to attend or they often visited a physician anyway. Thirdly, screenees were more active, had better self-rated health, and had a greater sense that life is worth living. These characteristics are associated with lower mortality (Sone et al.,

2008; Idler and Benyamini, 1997). However, although we considered them as confounding factors, these factors could not fully account for the difference in mortality. Finally, our result did not change substantially when we used propensity matched cohort analyses to minimize the self-selection bias. Therefore, we concluded that the difference in mortality rates between screenees and non-screenees was observed, even though we used detailed lifestyle information, extant diseases, or propensity for participating in health check-up. However, as with all prospective cohort studies, unknown confounding factors might exist. Although the effect of unknown and unmeasured confounding factors on mortality might be reduced by using propensity matched cohort analyses, only randomized controlled trials can provide an unbiased estimate of the effect.

A beneficial effect of health checks on mortality rates might be attributable to early detection of risk factors and early intervention

Table 3

Relation between using a health check-up in 1995 and cause-specific mortality. The Ohsaki Cohort Study, 1995–2006.

		All-cause mortality		CVD mortality		Cancer mortality		Mortality due to other causes	
		Screenees	Non-screenees	Screenees	Non-screenees	Screenees	Non-screenees	Screenees	Non-screenees
Men	<i>n</i>	6814	16,637	6814	16,637	6814	16,637	6814	16,637
	Number of deaths	1052	3589	261	1043	418	1219	373	1327
	Mortality rate (per 1000 person-years)	15.8	23.8	3.9	6.9	6.3	8.1	5.6	8.8
	Hazard ratio (95% CI) Model 1	0.72 (0.67–0.77)	1	0.68 (0.58–0.79)	1	0.75 (0.66–0.84)	1	0.71 (0.63–0.81)	1
	Hazard ratio (95% CI) Model 2	0.73 (0.61–0.87)	1	0.64 (0.43–0.94)	1	0.76 (0.58–0.997)	1	0.77 (0.57–1.04)	1
Women	<i>n</i>	9171	16,153	9171	16,153	9171	16,153	9171	16,153
	Number of deaths	519	2125	174	758	179	574	166	793
	Mortality rate (per 1000 person-years)	5.7	14.0	1.9	5.0	2.0	3.8	1.8	5.2
	Hazard ratio (95% CI) Model 1	0.66 (0.59–0.73)	1	0.70 (0.58–0.84)	1	0.66 (0.55–0.79)	1	0.62 (0.51–0.74)	1
	Hazard ratio (95% CI) Model 2	0.66 (0.50–0.86)	1	0.60 (0.36–1.006)	1	0.64 (0.41–0.98)	1	0.70 (0.43–1.14)	1

n: number of participants; CVD: cardiovascular diseases; CI: confidence interval; Model 1: we used Cox proportional hazards model with age as the time scale. In the model, we adjusted for body mass index (calculated as weight in kilograms divided by height in meters squared; <18.5, 18.5–21.9, 22.0–24.9, 25.0–29.9, and ≥30); participation in sports or exercise (<1 hour/week and ≥1 hour/week); time spent walking (<1 or ≥1 hour/day); sense of life worth living (*ikigai*); self-rated health; physical function; history of any extant diseases; smoking status (never, former, and current); alcohol consumption (never, former, and current); daily consumption of meat, fish, green and yellow vegetables, and green tea; job status; education (until 15 years of age, between 16 and 18 years of age, and ≥19 years of age); residential area; and history of screening for lung, gastric, colon, breast, and uterine cancer. Model 2: excluding participants with any extant diseases, those with limited physical function, and those who died within the first 3 years of follow-up.

including treatment. Risk factors such as high BP, abnormal liver function, and excessive alcohol consumption can be identified earlier among screenees of health check-ups. When risk factors are identified, participants are advised to visit a physician or to alter their lifestyle. If participants follow such advice, risk factors might be better controlled and lead to a reduction in future mortality rates. Although the health check-ups mainly screen risk factors for CVD, this process is also applicable to non-CVD mortality. Changes in smoking status or alcohol consumption could reduce rates of not only CVD diseases but also of cancer, liver, and respiratory diseases. Education about decreasing salt intake to lower BP might contribute to reducing salt-related diseases such as gastric cancer and kidney disease. However, our study did not collect detailed information about whether screenees with abnormal findings determined through health check-ups actually visited clinicians, changed their lifestyles, or controlled their risk factors appropriately through medication or lifestyle modification. This is a study limitation.

We had another limitation in this study. Except for several questionnaires, we generally relied on the self-reported questionnaire. We also rely on the data from secondary sources, such as death certificate. Usage of this information might yield uncontrolled errors and imprecision (Doria-Rose et al., 2010).

In conclusion, we found that the mortality risk was lower among screenees than non-screenees when we adjusted for possible confounding factors, such as lifestyle and extant diseases. This is also true when we used propensity matched cohort analyses. However, this study could not clarify whether the health check processes actually lead to a decreased risk of mortality. Further studies including randomized controlled trials are required to confirm our findings, but such trials could not be performed in Japan, since the national health check-up service is available to all. Therefore, we believe that the present findings represent the best available information regarding the relationship between the Japanese health check system and mortality.

Conflict of interest statement

The authors declare that there are no conflicts of interest.

Acknowledgments

This study was supported by Health Sciences Research Grants for Health Services (H19-Seisaku-Ippan-026 and H20-Junkankitou (Seisyu)-Ippan-013), Ministry of Health, Labour and Welfare, Japan.

This study is not related with any industry.

Table 4

Relation between using health check-up in 1995 and all-cause and cause-specific mortality using propensity for undergoing health check-up matched cohort. The Ohsaki Cohort Study, 1995–2006.

		All-cause mortality		CVD mortality		Cancer mortality		Mortality due to other causes	
		Screenees	Non-screenees	Screenees	Non-screenees	Screenees	Non-screenees	Screenees	Non-screenees
Men	<i>n</i> of participants	1800	1800	1800	1800	1800	1800	1800	1800
	Number of deaths	137	189	26	43	59	76	52	70
	HR (95% CI)	0.70 (0.56–0.88)	1	0.63 (0.38–1.03)	1	0.79 (0.55–1.12)	1	0.67 (0.46–0.97)	1
Women	<i>n</i> of participants	2087	2087	2087	2087	2087	2087	2087	2089
	Number of deaths	72	99	18	27	29	40	25	32
	HR (95% CI)	0.73 (0.53–0.99)	1	0.68 (0.37–1.26)	1	0.73 (0.45–1.17)	1	0.77 (0.45–1.32)	1
Men and women combined	<i>n</i> of participants	3887	3887	3887	3887	3887	3887	3887	3887
	Number of deaths	209	288	44	70	88	116	77	102
	HR (95% CI)	0.71 (0.59–0.86)	1	0.65 (0.44–0.95)	1	0.76 (0.58–1.01)	1	0.70 (0.51–0.95)	1

n: number; HR: hazard ratio; CI: confidence interval.

We thank Yoshiko Nakata, Mika Wagatsuma, and Tomoko Muroi from the Division of Epidemiology, Department of Public Health and Forensic Medicine, Tohoku University Graduate School of Medicine, Sendai, Japan, for their research assistance. We also thank for Professor Aaron R. Folsom for his valuable comments on this paper.

References

- D'Agostino Jr., R.B., 2007. Propensity scores in cardiovascular research. *Circulation* 115, 2340–2343.
- Doria-Rose, V.P., Marcus, P.M., Miller, A.B., et al., 2010. Does the source of death information affect cancer screening efficacy results? A study of the use of mortality review versus death certificates in four randomized trials. *Clin. Trials* 7, 69–77.
- Hozawa, A., Kuriyama, S., Kakizaki, M., Ohmori-Matsuda, K., Ohkubo, T., Tsuji, I., 2009. Attributable risk fraction of prehypertension on cardiovascular disease mortality in the Japanese population: the Ohsaki study. *Am. J. Hypertens.* 22, 267–272.
- Idler, E.L., Benyamini, Y., 1997. Self-rated health and mortality: a review of twenty-seven community studies. *J. Health Soc. Behav.* 38, 21–37.
- Ikeda, A., Iso, H., Toyoshima, H., Fujino, Y., et al., 2005. The relationships between interest for and participation in health screening and risk of mortality: the Japan Collaborative Cohort Study. *Prev. Med.* 41, 767–771.
- Iwasaki, M., Yamamoto, S., Otani, T., et al., 2006. Japan Public Health Center-based Prospective Study Group. Generalizability of relative risk estimates from a well-defined population to a general population. *Eur. J. Epidemiol.* 21, 253–262.
- Khan, M.M., Goto, R., Sonoda, T., et al., 2004. Impact of health education and screening over all-cause mortality in Japan: evidence from a cohort study during 1984–2002. *Prev. Med.* 38, 786–792.
- Korn, E.L., Graubard, B.I., Midthune, D., 1997. Time-to-event analysis of longitudinal follow-up for a survey: choice of the time-scale. *Am. J. Epidemiol.* 145, 72–80.
- Kuriyama, S., Shimazu, T., Ohmori, K., et al., 2006. Green tea consumption and mortality due to cardiovascular disease, cancer, and all causes in Japan: the Ohsaki study. *JAMA* 296, 1255–1265.
- Lannerstad, O., Sternby, N.H., Isacson, S.O., Lindgren, G., Lindell, S.E., 1997. Effects of a health screening on mortality and causes of death in middle-aged men. A prospective study from 1970 to 1974 of men in Malmö, born 1914. *Scand. J. Soc. Med.* 5, 137–140.
- Ogawa, K., Tsubono, Y., Nishino, Y., et al., 2003. Validation of a food-frequency questionnaire for cohort studies in rural Japan. *Public Health Nutr.* 6, 147–157.
- Ohmori-Matsuda, K., Kuriyama, S., Hozawa, A., Nakaya, N., Shimazu, T., Tsuji, I., 2007. The joint impact of cardiovascular risk factors upon medical costs. *Prev. Med.* 44, 349–355.
- Sone, T., Nakaya, N., Ohmori, K., et al., 2008. Sense of life worth living (*ikigai*) and mortality in Japan: Ohsaki study. *Psychosom. Med.* 70, 709–715.
- Statistics and Information Department, Minister's Secretariat, Ministry of Health, Labour and Welfare, 2007. The 20th Life Tables. Health and Welfare Statistics Association, Tokyo, Japan. (in Japanese).
- Statistics and Information Department, Minister's Secretariat, Ministry of Health, Labour and Welfare, 2008. Vital Statistics of Japan. Health and Welfare Statistics Association, Tokyo, Japan. (in Japanese).
- Stewart, A.L., Hays, R.D., Ware, J.E., 1988. The MOS Short-form General Health Survey. Reliability and validity in a patient population. *Med. Care* 26, 724–735.
- Tatara, K., Shinsho, F., Suzuki, M., Takatorige, T., Nakanishi, N., Kuroda, K., 1991. Relation between use of health check ups starting in middle age and demand for inpatient care by elderly people in Japan. *BMJ* 302, 615–618.
- Tsubono, Y., Tsuji, I., Fujita, K., et al., 2002. Validation of walking questionnaire for population-based prospective studies in Japan: comparison with pedometer. *J. Epidemiol.* 12, 305–309.
- Tsuji, I., Nishino, Y., Ohkubo, T., et al., 1998. A prospective cohort study on National Health Insurance beneficiaries in Ohsaki, Miyagi Prefecture, Japan: study design, profiles of the subjects and medical cost during the first year. *J. Epidemiol.* 8, 258–263.
- U.S. Preventive Services Task Force (USPSTF), <http://www.ahrq.gov/clinic/uspstfix.htm>. Final access at Jan 9, 2009.
- Wilhelmsen, L., Berglund, G., Elmfeldt, D., et al., 1986. The multifactor primary prevention trial in Göteborg, Sweden. *Eur. Heart J.* 7, 279–288.

Impact of walking on life expectancy and lifetime medical expenditure: the Ohsaki Cohort Study

Masato Nagai,¹ Shinichi Kuriyama,^{1,2} Masako Kakizaki,¹ Kaori Ohmori-Matsuda,¹ Toshimasa Sone,¹ Atsushi Hozawa,^{1,4} Miyuki Kawado,⁵ Shuji Hashimoto,⁵ Ichiro Tsuji¹

To cite: Nagai M, Kuriyama S, Kakizaki M, *et al*. Impact of walking on life expectancy and lifetime medical expenditure: the Ohsaki Cohort Study. *BMJ Open* 2011;1:e000240. doi:10.1136/bmjopen-2011-000240

► Prepublication history for this paper is available online. To view these files please visit the journal online (<http://bmjopen.bmj.com>).

Received 28 June 2011
Accepted 25 August 2011

This final article is available for use under the terms of the Creative Commons Attribution Non-Commercial 2.0 Licence; see <http://bmjopen.bmj.com>

ABSTRACT

Objective: People who spend a longer time walking have lower demands for medical care. However, in view of their longer life expectancy, it is unclear whether their lifetime medical expenditure increases or decreases. The present study examined the association between time spent walking, life expectancy and lifetime medical expenditure.

Method: The authors followed up 27 738 participants aged 40–79 years and prospectively collected data on their medical expenditure and survival covering a 13-year-period. Participants were classified into those walking <1 and ≥1 h per day. The authors constructed life tables and estimated the life expectancy and lifetime medical expenditure from 40 years of age using estimate of multiaadjusted mortality and medical expenditure using a Poisson regression model and linear regression model, respectively.

Results: Participants who walked ≥1 h per day have a longer life expectancy from 40 years of age than participants who walked <1 h per day. The multiaadjusted life expectancy for those who walked ≥1 h per day was 44.81 years, significantly lower by 1.38 years in men ($p=0.0073$) in men and 57.78 years in women, non-significantly lower by 1.16 years in women ($p=0.2351$). In addition to their longer life expectancy, participants who walked ≥1 h per day required a lower lifetime medical expenditure from 40 years of age than participants who walked <1 h per day. The multiaadjusted lifetime medical expenditure for those who walked ≥1 h per day was £99 423.6, significantly lower by 7.6% in men ($p=0.0048$) and £128 161.2, non-significantly lower by 2.7% in women ($p=0.2559$).

Discussion: Increased longevity resulting from a healthier lifestyle does not necessarily translate into an increased amount of medical expenditure throughout life. Encouraging people to walk may extend life expectancy and decrease lifetime medical expenditure, especially for men.

INTRODUCTION

Previous studies have agreed that a higher level of physical activity extends life expect-

ARTICLE SUMMARY

Article focus

- Medical expenditure per month was reduced when the amount of time spent walking was increased.
- Walking is associated with a decreased risk of mortality.
- In view of the increased life expectancy of those who walk longer, it is unclear whether lifetime medical expenditure increases or decreases as a result.

Key messages

- Lifetime medical expenditure from the age of 40 years for men and women who walked ≥1 h per day was reduced by 7.6% and 2.7%, respectively, in comparison with those who walked <1 h per day.
- Years of life added as a result of a healthy lifestyle did not necessarily translate into an increased amount of lifetime medical expenditure.

Strengths and limitations of this study

- This is the first study to investigate the association between walking, life expectancy and lifetime medical expenditure.
- We assessed walking using a simple questionnaire, in which we asked the participants to report only the time spent walking, and did not ask about walking pace, distance walked or any distinction between walking for exercise and other reasons.

ancy.^{1–4} Walking is part of a physically active lifestyle. Previous studies have indicated that a longer time spent walking,^{5–13} walking pace^{8 14–16} and a longer distance walked^{17 18} are significantly associated with a decreased risk of mortality.

We previously reported that medical expenditure per month was significantly reduced among those who spent a longer time walking, based on the same cohort dataset as that used here.¹⁹ Similar findings have been reported worldwide.^{20 21} However,

For numbered affiliations see end of article.

Correspondence to

Dr Masato Nagai;
m-nagai@med.tohoku.ac.jp

Walking and life expectancy and lifetime medical cost

in view of the increased life expectancy of those who walk for a longer time, it is unclear whether lifetime medical expenditure increases or decreases as a result. In other words, the question to be answered here is whether a lifelong healthy lifestyle eventually increases lifetime medical expenditure because of extended life expectancy.

So far, only one study has examined the association between physical activity and lifetime costs.⁴ This revealed that people with a high physical activity level tended to live longer than people with a lower physical activity level, and that the former had lower lifetime medical costs than the latter. However, that study was only a simulation analysis based on assumed variations in the health and economic effects of active and sedentary lifestyles.

The objective of the present study was to examine the association between walking, life expectancy and lifetime medical expenditure using actual individual data derived from a population-based 13-year prospective observation period. The population comprised 27 738 Japanese adults aged 40–79 years living in the community who were free of any functional limitations or chronic conditions interfering with physical activity, with an accrued total of 285 342 person-years. This cohort study has been monitoring survival and medical care utilisation, and its costs, for all participants.^{19–22} Using this dataset, we constructed a life table to estimate life expectancy and lifetime medical expenditure according to the time spent walking.^{24–25}

MATERIALS AND METHODS

Study cohort

The present data were derived from the Ohsaki National Health Insurance (NHI) Cohort Study.^{19–22–25} We conducted a self-administered questionnaire survey of various lifestyle habits between October and December 1994 for all NHI beneficiaries aged 40–79 years who lived in the catchment area of Ohsaki Public Health Center, Miyagi Prefecture, northeastern Japan. Out of 54 996 eligible individuals, 52 029 (95%) responded.

We excluded 776 participants who had withdrawn from the NHI before 1 January 1995 because their cost data were not available. Thus, the remaining 51 253 participants formed the study cohort. The study protocol was approved by the Ethics Committee of Tohoku University School of Medicine. Participants who had returned the self-administered questionnaires and signed them were considered to have consented to participate.

For the present analysis, we excluded participants who had functional limitation or chronic conditions interfering with physical activity. Physical function status was assessed using the six-item measure of the Medical Outcomes Study Short-form General Health Survey.²⁶ Participants were excluded if they stated on the Medical Outcomes Study questionnaire that they were unable to perform moderate or vigorous activities ($n=15\,916$). We

excluded participants who reported severe bodily pain ($n=949$), or any history of stroke ($n=474$), myocardial infarction ($n=585$) or arthritis ($n=2176$). We also excluded those who died within the first year ($n=174$) or did not provide complete responses in the walking status questionnaire ($n=3241$). Thus, a total of 27 738 participants (15 521 men and 12 217 women) remained. These participants were apparently healthy enough to walk for as long as they wished.

Time spent walking

The self-administered questionnaire included items on time spent walking. Time spent walking was assessed through the subject's response to the question, 'About how much time do you walk per day on average?' The participants were asked to choose one of three answers: '1 h or more,' '30 min–1 h' or '30 min or less.' In Japan, the Ministry of Health, Labour and Welfare recommended to walk ≥ 1 h per day in Exercise and Physical Activity Reference for Health Promotion 2006. Then, we divided the participants into two groups according to the time spent walking daily: < 1 h and ≥ 1 h. We had previously evaluated and reported the validity of self-reported time spent walking.^{5–19–27} This validation study had indicated that self-reported walking time was reasonably reproducible and sufficiently valid for studying the health effects of walking.

Health-insurance system in Japan

Details of the Japanese NHI system have been described previously.^{22–28–29} Briefly, everyone living in Japan is required to enrol in a health-insurance system. The NHI covers 35% of the Japanese population, mainly farmers, self employed or retired people. The NHI covers almost all medical treatment, including diagnostic tests, medication, surgery, supplies and materials, physicians and other personnel costs, inpatient care and most dental treatment. It covers treatment by physicians and nurses but not that by other professionals such as home health aides. Payment to medical providers is made on a fee-for-service basis, where the price of each service is determined by a uniform national fee schedule.

When a participant withdraws from the NHI system because of death, emigration or employment, the withdrawal date and its reason are coded in the NHI withdrawal history files. We recorded any mortality or migration by reviewing the NHI withdrawal history files and collected data on the death of participants by reviewing the death certificates filed at Ohsaki Public Health Center. We thus followed up the participants and prospectively collected data on medical care utilisation and its costs for all individuals in the cohort from 1 January 1995 to 31 December 2007. Study participants (16.3%) were lost to follow-up, so their vital status was unknown.

Statistical analysis

Using the Ohsaki NHI cohort database, we estimated mortality and medical expenditure for individual age

groups, and for the categories of time spent walking, for both men and women. We divided age into the following groups: 40–44, 45–49, 50–54, 55–59, 60–64, 65–69, 70–74, 75–79, 80–84 and ≥ 85 years. The multiadjusted mortalities for each age category were estimated from a Poisson regression model based on person-years and the number of deaths from 1996 until 2007. The dependent variable was mortality, and the independent variables were age groups, categories of time spent walking and the following covariates: smoking status (current and past smoker, or never smoker), alcohol consumption (current drinker consuming 1–499 g/week, current drinker consuming ≥ 450 g/week, or never and past drinker), body mass index (BMI: < 21 kg/m², 21–24.9 kg/m² or ≥ 25 kg/m²), self-rated health (good or not good), sports and physical activity (≥ 3 h/week or < 3 h/week), history of hypertension disease (presence or absence), history of diabetes mellitus (presence or absence), history of cancer (presence or absence), history of liver disease (presence or absence) and history of kidney disease (presence or absence).¹⁹ The data on all covariates were obtained from a self-administered questionnaire. We estimated the mortality of participants aged ≥ 85 years by multiplying the estimated mortality for the 85–89-year age group by the ratio of mortality for the same age group relative to the mortality for ≥ 85 years from complete life tables for the year 2000, as there were few person-years for participants aged over 90 years in our dataset.³⁰

Because medical expenditure increases before death, we separately calculated medical expenditure for participants who survived through the index year and for those who died. The multiadjusted medical expenditure per year for survivors and decedents, respectively, was estimated for each of the age groups and the categories of time spent walking using a linear regression model adjusted for the above covariates.

The estimates of multiadjusted mortality and medical expenditure for each age group were used for estimating life expectancy and lifetime medical expenditure from 40 years of age. Life expectancy was calculated using Chiang's analytical method on the basis of the latest published complete life tables of Japan for the year 2000.^{23–30} Lifetime medical expenditure was estimated from the sum obtained by multiplying the static population in life table by the medical expenditure for survivors and the number of deaths in the life table by the increased medical expenditure owing to death, which was calculated by subtracting the medical expenditure in year of survivors from that in the period of 1 year before death. That is, the life expectancy (e_x) and lifetime medical expenditure (M_x) for each age group (x) were estimated using the numbers of survivors (l_x), deaths (d_x), static population (L_x), multiadjusted medical expenditure for survivors (a_y) and multiadjusted medical expenditure for the deceased (b_y) as follows:

\sum is the sum of $y \geq x$

$$e_x = \frac{\sum L_y}{l_x}$$

$$M_x = \frac{\sum (L_y a_y + d_y b_y)}{l_x}$$

The 95% CIs were estimated using a Monte Carlo simulation with 100 000 replicates based on a Poisson regression model and linear regression model. All analyses were used the SAS V.9.1 statistical software package.

We used a purchasing-power parity rate (British pounds to Japanese yen) of £1.00=¥140.

RESULTS

After 13 years of follow-up, we observed 2936 deaths (2193 men and 743 women) among the 27738 participants (15521 men and 12217 women). The mean medical expenditure per year for survivors who walked ≥ 1 h per day was £1714.2 in men and £1621.4 in women, significantly lower than for those who walked < 1 h per day (men: £2064.3, $p < 0.0001$; women: £1878.6, $p < 0.0001$). Also, the mean medical expenditure in the year of death for participants who walked ≥ 1 h per day was £16878.6 in men and £17464.3 in women, which was not significantly different from those who walked < 1 h per day (men: £16650.0, $p = 0.7315$; women: £17742.9, $p = 0.8330$).

Baseline characteristics in terms of categories for time spent walking

Table 1 shows the baseline characteristics of the study participants according to the categories of time spent walking for men and women, respectively.

As compared with those who walked < 1 h per day, participants who walked ≥ 1 h per day were less likely to be smokers and obese. Self-reported histories of hypertension, diabetes mellitus and liver disease were all significantly less prevalent in those who walked ≥ 1 h per day.

Mortality in terms of categories for time spent walking

Figure 1A (for men) and Figure 1B (for women) show the multiadjusted mortality (per 1000) in each of the age groups according to the categories of time spent walking.

In men in each age group, the multiadjusted mortality was lower in participants who walked ≥ 1 h per day than in those who walked < 1 h per day. In women in all age groups except for the aged 40–44 and 45–49 year groups, the multiadjusted mortality was lower in participants who walked ≥ 1 h per day than in those who walked < 1 h per day.

Walking and life expectancy and lifetime medical cost

Table 1 Baseline characteristics by time-spent-walking categories in 27 738 participants

	Men			Women		
	Time spent walking		p Value*	Time spent walking		p Value*
	<1 h	≥1 h		<1 h	≥1 h	
No of subjects	7363	8158	0.2784	6303	5914	0.0714
Mean (SD) age	57.4 (10.6)	57.2 (10.2)		57.9 (10.1)	57.6 (9.7)	
Smoking status (%)						
Current and past smoker	81.9	79.8	0.0011	12.6	9.9	<0.0001
Never smoker	18.1	20.2		87.5	90.1	
Alcohol drinking (%)						
Never and past drinker	26.5	25.1	0.1117	74.7	75.4	0.1380
Current drinker,	61.3	62.2		24.3	23.9	
1–449 g/week						
Current drinker,	12.2	12.8		1.1	0.7	
≥450 g/week						
Body mass index (%)						
<21 kg/m ²	18.5	20.5	<0.0001	18.6	19.4	0.0019
21–24.9 kg/m ²	53.3	55.8		50.7	52.9	
≥25 kg/m ²	28.3	23.7		30.7	27.7	
Self-rated health (%)						
Good	73.1	79.3	<0.0001	72.5	78.2	<0.0001
Not good	26.9	20.7		27.5	21.8	
Sports and physical activity (%)						
≥3 h/week	13.2	18.7	<0.0001	10.7	16.6	<0.0001
<3 h/week	86.8	81.3		89.3	83.4	
History of hypertension (%)						
Presence	21.5	18.6	<0.0001	24.2	20.1	<0.0001
Absence	78.5	81.4		75.8	79.9	
History of diabetes mellitus (%)						
Presence	7.4	4.8	<0.0001	5.1	3.2	<0.0001
Absence	92.6	95.2		94.9	96.8	
History of cancer (%)						
Presence	2.1	1.8	0.2563	2.8	2.8	0.9032
Absence	97.9	98.2		97.2	97.2	
History of liver disease (%)						
Presence	6.9	5.7	0.0013	3.7	2.9	0.0223
Absence	93.1	94.4		96.3	97.1	
History of kidney disease (%)						
Presence	2.9	2.7	0.3493	3.6	2.7	0.0049
Absence	97.1	97.3		96.4	97.3	

*p Values were calculated using the χ^2 test.

Table 2 shows the mortality ratio with 95% CIs according to the categories of time spent walking. In men, the multiadjusted mortality ratio for participants who walked ≥1 h per day was significantly lower than that for participants who walked <1 h per day (0.90, 95% CI 0.82 to 0.98, $p=0.0153$). In women, the multiadjusted mortality ratio for participants who walked ≥1 h per day was non-significantly lower than that for participants who walked <1 h per day (0.95, 95% CI 0.82 to 1.10, $p=0.4693$).

Life expectancy and lifetime medical expenditure in terms of time spent walking

Table 3 shows the life expectancy and lifetime medical expenditure from 40 years of age with 95% CIs according to the categories of time spent walking.

In men, the multiadjusted life expectancy of those who walked ≥1 h per day was 44.81 years (95% CI 43.66 to

45.94), which was significantly longer by 1.38 years ($p=0.0073$) than for those who walked <1 h per day (43.43 years; 95% CI 42.39 to 44.41). In women, the same results were observed, although the differences did not reach statistical significance.

In spite of their longer life expectancy, their lifetime medical expenditure from 40 years of age was significantly lower in men and non-significantly lower in women. The multiadjusted lifetime medical expenditure for participants who walked ≥1 h per day was £99 423.6 (95% CI 92 515.9 to 106 694.7), significantly lower by 7.6% ($p=0.0048$) than for those who walked <1 h per day (£107 544.2; 95% CI 101 234.0 to 114 044.6). In women, the multiadjusted lifetime medical expenditure for participants who walked ≥1 h per day was £128 161.2 (95% CI 111 335.0 to 148 494.7), non-significantly lower by 2.7% ($p=0.2559$) than for

Walking and life expectancy and lifetime medical cost




Figure 1 Multiadjusted mortality by time-spent-walking categories in each age group in (A) men and (B) women.

those who walked <1 h per day (£131 766.8; 95% CI 115 902.4 to 150 714.3).

DISCUSSION

The present results indicate that the multiadjusted lifetime medical expenditure from the age of 40 years for those who walked ≥ 1 h per day was significantly lower by 7.6% in men and non-significantly lower by 2.7% in women than for those who walked <1 h per day. This decrease in lifetime medical expenditure was observed in spite of a longer life expectancy (1.38 years for men and 1.16 years for women) among those who walked ≥ 1 h per day. Thus, a healthy lifestyle not only extended longevity but also decreased the amount of lifetime medical expenditure, especially men.

We observed statistically significant differences for men but not for women. Although the differences did not reach statistical significance, the same results were observed for women. The reason why the impact of walking was smaller in women than in men was unknown. In women, other factors such as obesity and postmenopausal change might have a stronger impact on life expectancy and lifetime medical expenditure than walking.

Comparison with other studies

Four studies have addressed the association between physical activity level and life expectancy.¹⁻⁴ Additionally, studies investigating associations between time spent walking,⁵⁻¹³ or distance walked^{17 18} and mortality have consistently shown that participants who have a higher

physical activity have a longer life expectancy and lower mortality than participants who have a lower physical activity. Only one study has reported the association between physical activity and lifetime medical expenditure.⁴ In a simulation study, Keeler *et al* demonstrated that if participants with a sedentary lifestyle had exercised regularly, the additional exercise would have increased their life expectancy by 300 days and saved £1900 in lifetime costs. Even though there is a difference between physical activity and walking, this result is consistent with the present finding that participants who walked ≥ 1 h per day lived longer and had a lower lifetime medical expenditure.

We previously calculated life expectancy and lifetime medical expenditure in relation to smoking and BMI from age 40 years using the same dataset as that for the present study.²⁴ The results indicated that lifetime medical expenditure was lower for smokers than for non-smokers, reflecting the 3.5-year shorter life expectancy of smokers.²⁴ On the other hand, lifetime medical expenditure was higher for participants who were obese ($BMI \geq 30.0$) than for those who were of normal weight ($18.5 \leq BMI < 25.0$), even though the former lived 2 years less than the latter (Nagai M, Kuriyama S, Kakizaki M, *et al*. Impact of obesity, overweight and underweight on life expectancy and lifetime medical expenditures. *Popul Health Metr*; under review). In fact, both smokers and obese participants had a shorter life expectancy than non-smokers and normal-weight participants, whereas their lifetime medical expenditure was conversely

Table 2 Mortality ratio for time-spent-walking categories in 27 738 participants

Time spent walking	Univariate		Multiadjusted*	
	Mortality ratio (95% CI)	p Value	Mortality ratio (95% CI)	p Value
Men	<1 h	1.00	1.00	
	≥ 1 h	0.87 (0.80 to 0.94)	0.90 (0.82 to 0.98)	0.0153
Women	<1 h	1.00	1.00	
	≥ 1 h	0.89 (0.77 to 1.03)	0.95 (0.82 to 1.10)	0.4693

*Adjusted for age groups, smoking status, alcohol drinking, body mass index, self-rated health, sports and physical activity, and history of hypertension, diabetes mellitus, cancer, liver disease and kidney disease.

Walking and life expectancy and lifetime medical cost

Table 3 Life expectancy and lifetime medical expenditure at age 40 years for time-spent-walking categories in 27 738 participants

	Time spent walking	Univariate			Multiadjusted*		
		Estimate	95% CI	p Value	Estimate	95% CI	p Value
Men	Life expectancy (years) at age 40 years						
	<1 h	42.41	41.45 to 43.26	0.0004	43.43	42.39 to 44.41	0.0073
	≥1 h	44.19	43.15 to 45.19		44.81	43.66 to 45.94	
	Lifetime medical expenditure (£) at age 40 years						
<1 h	107 023.2	101 093.6 to 113 066.3	<0.0001	107 544.2	101 234.0 to 114 044.6	0.0048	
≥1 h	94 402.1	87 812.3 to 101 248.0		99 423.6	92 515.9 to 106 694.7		
Women	Life expectancy (years) at age 40 years						
	<1 h	52.25	49.79 to 54.92	0.0569	56.62	53.17 to 60.62	0.2351
	≥1 h	54.25	51.38 to 57.48		57.78	54.02 to 62.22	
	Lifetime medical expenditure (£) at age 40 years						
<1 h	123 553.0	111 619.5 to 137 549.6	0.0644	131 766.8	115 902.4 to 150 714.3	0.2559	
≥1 h	115 896.0	102 406.6 to 131 792.1		128 161.2	111 335.0 to 148 494.7		

*Adjusted for age groups, smoking status, alcohol drinking, body mass index, self-rated health, sports and physical activity, and history of hypertension, diabetes mellitus, cancer, liver disease and kidney disease as same as table 2.

increased. These differences could be explained by the impact of these risk factors on quality of life. Using prospective data for 16 176 adult Caucasians in the USA,³¹ Reuser *et al* estimated life expectancy and years of life with and without activities of daily-living disability in relation to smoking and BMI, respectively. The results indicated that smoking decreased both life expectancy and years of life with activities of daily-living disability, whereas obesity decreased the former but increased the latter, thus leading to the conclusion that 'smoking kills, and obesity disables'. Reuser's conclusion is concordant with the impact of walking on life expectancy and lifetime medical expenditure. Physical activity decreases not only mortality but also disability,^{5–13 17 18 32} leading to lower medical expenditure.³³ For instance, walking has been significantly associated with a lower risk of cardiovascular disease,⁷ stroke,³⁴ coronary heart disease,^{35 36} type 2 diabetes³⁷ and hypertension.³⁸ Consequently, walking also reduces expenditure on medication needed for these conditions.³⁹

Strengths and limitations

This is the first study to have investigated the association between walking, life expectancy and lifetime medical expenditure. A major strength of this study was that we collected individual data on medical expenditure based on a cohort study of survival and medical-care utilisation, and its cost, for all participants,^{19 22 24 28} and the NHI covers almost all medical treatment in Japan. Second, we conducted a 13-year prospective observation of 27 738 Japanese adults aged 40–79 years living in the community who were free of any functional limitations or chronic conditions interfering with physical activity, with an accrued total of 285 342 person-years. Third, in order to reduce bias or reverse causation in that people did not walk because of functional limitations that also required

medical expenditure, we excluded participants who, at the baseline, reported limited physical function or conditions interfering with physical activity. Additionally, to control for confounders, we also included various covariates in our Poisson regression model and linear regression model.

On the other hand, several limitations should also be considered. First, we assessed walking using a simple questionnaire in which we asked the participants to report only the time spent walking and did not ask about walking pace, distance walked or any distinction between walking for exercise and other reasons. Second, a longer time spent walking may be a reflection of performing more vigorous activity, making it difficult to distinguish the impact of walking from other types of physical activity. However, the present result did not change after multivariate adjustment.

Conclusions and policy implication

In summary, lifetime medical expenditure was shown to be decreased in participants who walked ≥1 h per day, despite the fact that they lived longer. Increased longevity resulting from a healthier lifestyle did not necessarily translate into an increased amount of medical expenditure throughout life. However, in the present study, around 50% of study participants walk <1 h per day. To increase their walking time, the recommendation of walking with a pedometer may be useful.⁴⁰ An increase in walking time at the population level would bring about a tremendous change in people's health and medical cost. A campaign to encourage people to walk for longer and a program to make walking environments safer and more pleasant should be implemented. This intervention may extend life expectancy without apparently increasing lifetime medical expenditure, especially for men.

Walking and life expectancy and lifetime medical cost

Author affiliations

¹Division of Epidemiology, Department of Public Health and Forensic Medicine, Tohoku University Graduate School of Medicine, Sendai, Japan

²Department of Molecular Epidemiology, Genome Research Center, Tohoku University Graduate School of Medicine, Sendai, Japan

³Department of Environment and Genome Research Center, Tohoku University Graduate School of Medicine, Sendai, Japan

⁴Department of Public Health, Yamagata University Graduate School of Medical Science, Yamagata, Japan

⁵Department of Hygiene, Fujita Health University School of Medicine, Toyoake, Japan

Correction notice The "To cite: ..." information and running footer in this article have been updated with the correct volume number (volume 1).

Funding This study was supported by a Health Sciences Research Grant for Health Services (H21-Choju-Ippan-001, H20-Junkankitou (Seisyu)-Ippan-013, H22- Junkankitou (Seisyu)-Ippan-012, H23- Junkankitou (Seisyu)-Ippan-005), Ministry of Health, Labour and Welfare, Japan. MN is a recipient of a Research Fellowships of the Japan Society for the Promotion of Science for Young Scientists.

Competing interests None.

Patient consent Obtained.

Ethical approval Ethics approval was provided by the Ethics Committee of Tohoku University School of Medicine.

Contributors All authors contributed to the design of the study. MN, SK, MK, KO-M, TS and IT carried out the data collection. MN, SK, AH, MK and SH carried out the data analysis. MN, MK, KO-M, TS, AH, MK and SH wrote the report. SK and IT carried out a critical revision of the manuscript. All authors approved the final version of the report for submission.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement No additional data available.

REFERENCES

1. Franco OH, de Laet C, Peeters A, *et al*. Effects of physical activity on life expectancy with cardiovascular disease. *Arch Intern Med* 2005;165:2355–60.
2. Hatziaendreu EI, Koplan JP, Weinstein MC, *et al*. A cost-effectiveness analysis of exercise as a health promotion activity. *Am J Public Health* 1988;78:1417–21.
3. Jonker JT, De Laet C, Franco OH, *et al*. Physical activity and life expectancy with and without diabetes: life table analysis of the Framingham Heart Study. *Diabetes Care* 2006;29:38–43.
4. Keeler EB, Manning WG, Newhouse JP, *et al*. The external costs of a sedentary life-style. *Am J Public Health* 1989;79:975–81.
5. Fujita K, Takahashi H, Miura C, *et al*. Walking and mortality in Japan: the Miyagi Cohort Study. *J Epidemiol* 2004;14(Suppl 1): S26–32.
6. Kujala UM, Kaprio J, Sarna S, *et al*. Relationship of leisure-time physical activity and mortality: the Finnish twin cohort. *JAMA* 1998;279:440–4.
7. LaCroix AZ, Leveille SG, Hecht JA, *et al*. Does walking decrease the risk of cardiovascular disease hospitalizations and death in older adults? *J Am Geriatr Soc* 1996;44:113–20.
8. Lee IM, Rexrode KM, Cook NR, *et al*. Physical activity and coronary heart disease in women: is 'no pain, no gain' passe? *JAMA* 2001;285:1447–54.
9. Morioka S. [A cohort study on the relationship between lifestyles and total mortality] (in Japanese). *Nippon Kosho Eisei Zasshi* 1996;43:469–78.
10. Seki N. [Relationships between walking hours, sleeping hours, meaningfulness of life (ikigai) and mortality in the elderly: prospective cohort study] (in Japanese). *Nippon Eiseigaku Zasshi* 2001;56:535–40.
11. Stessman J, Maaravi Y, Hammerman-Rozenberg R, *et al*. The effects of physical activity on mortality in the Jerusalem 70-Year-Olds Longitudinal Study. *J Am Geriatr Soc* 2000;48:499–504.
12. Wannamethee SG, Shaper AG, Walker M. Physical activity and mortality in older men with diagnosed coronary heart disease. *Circulation* 2000;102:1358–63.
13. Weller I, Corey P. The impact of excluding non-leisure energy expenditure on the relation between physical activity and mortality in women. *Epidemiology* 1998;9:632–5.
14. Dumurgier J, Elbaz A, Ducimetière P, *et al*. Slow walking speed and cardiovascular death in well functioning older adults: prospective cohort study. *BMJ* 2009;339:b4460.
15. Tanasescu M, Leitzmann MF, Rimm EB, *et al*. Exercise type and intensity in relation to coronary heart disease in men. *JAMA* 2002;288:1994–2000.
16. Cooper R, Kuh D, Hardy R. Objectively measured physical capability levels and mortality: systematic review and meta-analysis. *BMJ* 2010;341:c4467.
17. Hakim AA, Petrovitch H, Burchfiel CM, *et al*. Effects of walking on mortality among nonsmoking retired men. *N Engl J Med* 1998;338:94–9.
18. Lee IM, Paffenbarger RS Jr. Associations of light, moderate, and vigorous intensity physical activity with longevity. The Harvard Alumni Health Study. *Am J Epidemiol* 2000;151:293–9.
19. Tsuji I, Takahashi K, Nishino Y, *et al*. Impact of walking upon medical care expenditure in Japan: the Ohsaki Cohort Study. *Int J Epidemiol* 2003;32:809–14.
20. Jones TF, Eaton CB. Cost-benefit analysis of walking to prevent coronary heart disease. *Arch Fam Med* 1994;3:703–10.
21. Perkins AJ, Clark DO. Assessing the association of walking with health services use and costs among socioeconomically disadvantaged older adults. *Prev Med* 2001;32:492–501.
22. Tsuji I, Nishino Y, Ohkubo T, *et al*. A prospective cohort study on National Health Insurance beneficiaries in Ohsaki, Miyagi Prefecture, Japan: study design, profiles of the subjects and medical cost during the first year. *J Epidemiol* 1998;8:258–63.
23. Chiang CL. *The Life Table and Its Applications*. Malabar: Robert E Krieger Publishing Company, 1984.
24. Hayashida K, Imanaka Y, Murakami G, *et al*. Difference in lifetime medical expenditures between male smokers and non-smokers. *Health Policy* 2010;94:84–9.
25. Kuriyama S, Shimazu T, Ohmori K, *et al*. Green tea consumption and mortality due to cardiovascular disease, cancer, and all causes in Japan: the Ohsaki study. *JAMA* 2006;296:1255–65.
26. Stewart AL, Hays RD, Ware JE Jr. The MOS short-form general health survey. Reliability and validity in a patient population. *Med Care* 1988;26:724–35.
27. Tsubono Y, Tsuji I, Fujita K, *et al*. Validation of walking questionnaire for population-based prospective studies in Japan: comparison with pedometer. *J Epidemiol* 2002;12:305–9.
28. Kuriyama S, Tsuji I, Ohkubo T, *et al*. Medical care expenditure associated with body mass index in Japan: the Ohsaki Study. *Int J Obes Relat Metab Disord* 2002;26:1069–74.
29. Nakamura K, Okamura T, Kanda H, *et al*. Medical costs of obese Japanese: a 10-year follow-up study of National Health Insurance in Shiga, Japan. *Eur J Public Health* 2007;17:424–9.
30. Statistics and Information Department Minister's Secretariat, Ministry of Health, Labour and Welfare of Japan. *The 20th Life Tables*. Tokyo: Health and Welfare Statistics Association, Japan, 2000.
31. Reuser M, Bonneux LG, Willekens FJ. Smoking kills, obesity disables: a multistate approach of the US Health and Retirement Survey. *Obesity (Silver Spring)* 2009;17:783–9.
32. Clark DO. The effect of walking on lower body disability among older blacks and whites. *Am J Public Health* 1996;86:57–61.
33. Mitra S, Findley PA, Sambamoorthi U. Health care expenditures of living with a disability: total expenditures, out-of-pocket expenses, and burden, 1996 to 2004. *Arch Phys Med Rehabil* 2009;90:1532–40.
34. Hu FB, Stampfer MJ, Colditz GA, *et al*. Physical activity and risk of stroke in women. *JAMA* 2000;283:2961–7.
35. Hakim AA, Curb JD, Petrovitch H, *et al*. Effects of walking on coronary heart disease in elderly men: the Honolulu Heart Program. *Circulation* 1999;100:9–13.
36. Manson JE, Hu FB, Rich-Edwards JW, *et al*. A prospective study of walking as compared with vigorous exercise in the prevention of coronary heart disease in women. *N Engl J Med* 1999;341:650–8.
37. Hu FB, Sigal RJ, Rich-Edwards JW, *et al*. Walking compared with vigorous physical activity and risk of type 2 diabetes in women: a prospective study. *JAMA* 1999;282:1433–9.
38. Hayashi T, Tsumura K, Suematsu C, *et al*. Walking to work and the risk for hypertension in men: the Osaka Health Survey. *Ann Intern Med* 1999;131:21–6.
39. Williams PT. Reduced diabetic, hypertensive, and cholesterol medication use with walking. *Med Sci Sports Exerc* 2008;40:433–43.
40. Snyder A, Colvin B, Gammack JK, *et al*. Pedometer use increases daily steps and functional status in older adults. *J Am Med Dir Assoc*. Published Online First: 7 October 2010. doi:10.1016/j.jamda.2010.06.007.