

14. 梶谷展生, 四方賢一:糖尿病性細小血管症の発症・進展の分子メカニズム
Microinflammationの関与. 日本臨床 68(9), 2010
15. 四方賢一、槇野博史:慢性合併症の臨床 2(腎症)1. 腎症の成因. 糖尿病学の進歩 2010, 日本糖尿病学会編
16. 四方賢一、南條輝志男、羽田勝計、高木均、中村二郎:病期から見た糖尿病合併症対策. 糖尿病学の進歩 2010, 日本糖尿病学会編
17. 四方賢一:腎不全期 糖尿病最新の治療, 南江堂, 2010
18. 宮本聡、四方賢一:脂質異常症・セミナー 実地診療で遭遇する問題点の対応 腎疾患患者に合併する脂質異常症—管理の意義と方法—MedicalPractice 27(3), 2010.
19. 小寺亮、四方賢一:DPP-4 阻害薬の登場で薬物療法はどう変わるか. 月刊薬事 52(3), 2010.
20. 四方賢一、槇野博史:全身疾患と腎障害 1. 糖尿病性腎症. メディカルノート腎臓がわかる腎・尿路疾患水・電解質代謝異常 p. 78-82, 2010
21. 梶谷展生、宮本聡、小寺亮、四方賢一、槇野博史:糖尿病性腎症の理解と予防 外来看護最前線生活習慣病&外来がん看護 日総研, 2009
22. 小寺亮、四方賢一、槇野博史:糖尿病が引き起こす細小血管症③糖尿病性腎症 からだの科学 261号糖尿病のすべて, 2009
23. 小寺亮、四方賢一、槇野博史:腎臓専門医での診断とかかりつけ医への逆紹介の要点 4. 糖尿病腎症の確定診断はどのように行うのですか? かかりつけ医と専門医のためのCKD診療ガイド, 2009
24. 小寺亮、四方賢一:糖尿病性腎症 第24回免疫抑制薬の臨床応用実践論『炎症と免疫』 17(6), 2009
25. 四方賢一、宮本聡:糖尿病性腎症の成因としてのmicroinflammation. 月刊糖尿病 1(5)p43-49, 2009
- ガイドライン
1. 日本腎臓学会・日本医学放射線学会・日本循環器学会編「腎障害患者におけるヨード造影剤使用に関するガイドライン」2012 日本腎臓学会 「CKD診療ガイド2012」
- 2)海外
- 口頭発表 20 件
- 原著論文による発表 35 件
- それ以外(レビュー等)の発表 5 件
- そのうち主なもの
- 原著論文発表
1. Horio M, Yasuda Y, Kaimori J, Ichimaru N, Isaka Y, Takahara S, Nishi S, Uchida K, Takeda A, Hattori R, Kitada H, Tsuruya K, Imai E, Takahashi K, Watanabe T, Matsuo S. Performance of the Japanese GFR equation in potential kidney donors. Clin Exp Nephrol. in press. 2012
2. Suzuki Y, Ito Y, Mizuno M, Kinashi H, Sawai A, Noda Y, Mizuno T, Shimizu H, Fujita Y, Matsui K, Maruyama S, Imai E, Matsuo S, Takei Y. Transforming growth factor- β induces vascular endothelial growth factor-C expression leading to lymphangiogenesis in rat unilateral ureteral obstruction. Kidney Int. in press. 2012
3. Toda S, Ito Y, Mizuno M, Suzuki Y, Ito I, Hiramatsu H, Ozaki T, Tsuboi N, Sato W, Maruyama S, Imai E, Matsuo S. Asymptomatic diverticulosis identified by computed tomography is not a risk factor for enteric peritonitis. Nephrol Dial Transplant. in press. 2011
4. Shinzawa M, Yamamoto R, Nagasawa Y, Shoji T, Obi Y, Namba T, Kitamura H, Kaneko T,

- Okada N, Iwatani H, Yamauchi A, Tsubakihara Y, Imai E, Isaka Y, Rakugi H. Gene polymorphisms contributing to hypertension in immunoglobulin A nephropathy. *Clin Exp Nephrol.* in press. 2011
5. Kawada N, Moriyama T, Kitamura H, Yamamoto R, Furumatsu Y, Matsui I, Takabatake Y, Nagasawa Y, Imai E, Wilcox CS, Rakugi H, Isaka Y. Towards developing new strategies to reduce the adverse side-effects of nonsteroidal anti-inflammatory drugs. *Clin Exp Nephrol.* in press. 2011
 6. Imai E, Yasuda Y, Horio M, Shibata K, Kato S, Mizutani Y, Imai J, Hayashi M, Kamiya H, Oiso Y, Murohara T, Maruyama S, Matsuo S. Validation of the equations for estimating daily sodium excretion from spot urine in patients with chronic kidney disease. *Clin Exp Nephrol.* 15(6):861-7. 2011
 7. Suzuki Y, Mizuno M, Nakashima R, Hiramatsu H, Toda S, Sato W, Tsuboi N, Ito I, Maruyama S, Imai E, Matsuo S, Ito Y. A case of perforative peritonitis caused by a piece of bamboo in a patient on peritoneal dialysis. *Clin Exp Nephrol.* 15(6):962-5. 2011
 8. Iwatani H, Iio K, Nagasawa Y, Yamamoto R, Horii A, Okuzaki D, Inohara H, Nojima H, Imai E, Rakugi H, Isaka Y. Microarray analysis of tonsils of IgA nephropathy patients. *Adv Otorhinolaryngol.* 72:75-8. 2011
 9. Horio M, Imai E, Yasuda Y, Watanabe T, Matsuo S. Performance of serum cystatin C versus serum creatinine as a marker of glomerular filtration rate as measured by inulin renal clearance. *Clin Exp Nephrol.* 15(6):868-76. 2011
 10. Mizuno M, Ito Y, Hayasaki T, Suzuki Y, Hiramatsu H, Toda S, Mizuno T, Tatematsu M, Ozaki T, Yasuda Y, Sato W, Tsuboi N, Ito I, Maruyama S, Imai E, Matsuo S. A case of acute renal failure caused by cholesterol embolization after carotid artery stenting that was improved by peritoneal dialysis. *Intern Med.* 50(16):1719-23. 2011
 11. Li PK, Chow KM, Matsuo S, Yang CW, Jha V, Becker G, Chen N, Sharma SK, Chittinandana A, Chowdhury S, Harris DC, Hooi LS, Imai E, Kim S, Kim SG, Langham R, Padilla BS, Teo BW, Togtokh A, Walker RG, Wang HY, Tsukamoto Y; Asian Forum for Chronic Kidney Disease Initiatives. Asian chronic kidney disease best practice recommendations: positional statements for early detection of chronic kidney disease from Asian Forum for Chronic Kidney Disease Initiatives (AFCKDI). *Nephrology (Carlton).* 16(7):633-41. 2011
 12. Yamamoto R, Nagasawa Y, Shoji T, Katakami N, Ohtoshi K, Hayaishi-Okano R, Yamasaki Y, Yamauchi A, Tsubakihara Y, Imai E, Rakugi H, Isaka Y. A candidate gene approach to genetic contributors to the development of IgA nephropathy. *Nephrol Dial Transplant.* in press. 2011
 13. Mizuno M, Ito Y, Tanaka A, Suzuki Y, Hiramatsu H, Watanabe M, Tsuruta Y, Matsuoka T, Ito I, Tamai H, Kasuga H, Shimizu H, Kurata H, Inaguma D, Hiramatsu T, Horie M, Naruse T, Maruyama S, Imai E, Yuzawa Y, Matsuo S. Peritonitis is still an important factor for withdrawal from peritoneal dialysis therapy in the Tokai area of Japan. *Clin Exp Nephrol.* 15(5):727-37. 2011

14. Nakamori A, Ando Y, Matsuda H, Kimura T, Minami H, Imai E, Yura T. Influence of proteinuria on renal Doppler sonographic measurements in chronic kidney disease and in diabetes mellitus. *J Clin Ultrasound.* 39(9):506-11. 2011
15. Takabatake Y, Li XK, Mizui M, Miyasato K, Matsui I, Kawada N, Imai E, Hünig T, Takahara S, Wada T, Furuichi K, Rakugi H, Isaka Y. A superagonistic monoclonal antibody for CD28 ameliorates crescentic glomerulonephritis in wistar-kyoto rats. *Mol Med.* 17(7-8):686-96. 2011
16. Mizuno M, Ito Y, Masuda T, Toda S, Hiramatsu H, Suzuki Y, Ozaki T, Yasuda Y, Ito I, Tsuboi N, Sato W, Maruyama S, Imai E, Matsuo S. A case of fulminant peritonitis caused by *Streptococcus mitis* in a patient on peritoneal dialysis. *Intern Med.* 50(5):471-4. 2011
17. Akizawa T, Makino H, Matsuo S, Watanabe T, Imai E, Nitta K, Ohashi Y, Hishida A; Chronic Kidney Disease Japan Cohort Study Group. Management of anemia in chronic kidney disease patients: baseline findings from Chronic Kidney Disease Japan Cohort Study. *Clin Exp Nephrol.* 15(2):248-57. 2011
18. Imai E, Yasuda Y, Horio M, Shibata K, Kato S, Mizutani Y, Imai J, Hayashi M, Kamiya H, Oiso Y, Murohara T, Maruyama S, Matsuo S.: Validation of the equations for estimating daily sodium excretion from spot urine in patients with chronic kidney disease. *Clin Exp Nephrol.* 2011 Dec;15(6):861-7.
19. Horio M, Imai E, Yasuda Y, Watanabe T, Matsuo S. Modification of the CKD Epidemiology Collaboration (CKD-EPI) Equation for Japanese: Accuracy and Use for Population Estimates. *Am J Kidney Dis.* 56: 32-38, 2010.
20. Horio M, Yasuda Y, Takahara S, Watanabe T, Matsuo S. Comparison of simple method and standard method for inulin renal clearance. *Clin Exp Nephrol* 14(5):427-430, 2010.
21. Kakuta Y, Okumi M, Ichimaru N, Abe T, Nonomura N, Okuyama A, Kojima Y, Isaka Y, Takahara S, Imai E, Horio M. Utility of the Japanese GFR estimation equation for evaluating potential donor kidney function. *Clin Exp Nephrol.* 14: 63-37, 2010
22. Horio M, Imai E, Yasuda Y, Watanabe T, Matsuo S. Performance of serum cystatin C versus serum creatinine as a marker of glomerular filtration rate as measured by inulin renal clearance. *Clin Exp Nephrol.* 2011, 15:868-876
23. Gansvoort RT, Matsushita K, van der Verde M, Astor BC, et al. Lower estimated GFR and higher albuminuria are associated with adverse kidney outcomes in both general and high-risk population: A meta-analysis of general and high-risk population cohorts. *Kidney Int* 80: 93-104, 2011
24. Iseki K, Chiho Iseki, and Kozen Kinjo. C-reactive protein is a predictor for developing proteinuria in a screened cohort
25. Nakayama M, Sato T, Sato H, Yamaguchi Y, Obara K, Kurihara I, Sato K, Hotta O, Seino J, Miyata M, Takeuchi K, Nakayama K, Matsushima M, Otaka T, Kinoshita Y, Taguma Y, and Ito S. Different clinical outcomes for cardiovascular events and mortality in chronic kidney disease according to underlying renal disease: the Gonryo study. *Clin. Exp Nephrol* 14: 333-339,

- 2010.
26. Nakayama M, Sato T, Miyazaki M, Matsushima M, Sato H, Taguma Y and Ito S. Increased risk of cardiovascular events and mortality among non-diabetic chronic kidney disease patients with hypertensive nephropathy: the Gonryo study. *Hypertens Res* 34: 1106-1110, 2011.
 27. Ogawa D, Asanuma M, Miyazaki I, Tachibana H, Wada J, Sogawa N, Sugaya T, Kitamura S, Maeshima Y, Shikata K, Makino H. High glucose increases metallothionein expression in renal proximal tubular epithelial cells. *Exp Diabetes Res*. 2011. 534872, 2011.
 28. Watanabe N, Shikata K, Shikata Y, Sarai K, Omori K, Kodera R, Sato C, Wada J, Makino H. Involvement of MAPKs in ICAM-1 expression in glomerular endothelial cells in diabetic nephropathy. *Acta Med Okayama* 65:247-257, 2011.
 29. Sasaki M, Shikata K, Okada S, Miyamoto S, Nishishita S, Kataoka HU, Sato C, Wada J, Ogawa D, Makino H. The macrophage is a key factor in renal injuries caused by glomerular hyperfiltration. *Acta Med Okayama*. 2011 Apr; 65:81-89, 2011.
 30. Matsushita Y, Ogawa D, Wada J, Yamamoto N, Shikata K, Sato C, Tachibana H, Toyota N, Makino H. Activation of peroxisome proliferator-activated receptor delta inhibits streptozotocin-induced diabetic nephropathy through anti-inflammatory mechanisms in mice. *Diabetes*. 60:960-968, 2011.
 31. Kodera R, Shikata K, Kataoka HU, Takatsuka T, Miyamoto S, Sasaki M, Kajitani N, Nishishita S, Sarai K, Hirota D, Sato C, Ogawa D, Makino H. Glucagon-like peptide-1 receptor agonist ameliorates renal injury through its anti-inflammatory action without lowering blood glucose level in a rat model of type 1 diabetes. *Diabetologia*. 54:965-978, 2011.
 32. Sato C, Shikata K, Hirota D, Sasaki M, Nishishita S, Miyamoto S, Kodera R, Ogawa D, Tone A, Kataoka HU, Wada J, Kajitani N, Makino H. P-selectin glycoprotein ligand-1 deficiency is protective against obesity-related insulin resistance. *Diabetes*. 60:189-199, 2011.
 33. Kido Y, Ogawa D, Shikata K, Sasaki M, Nagase R, Okada S, Usui Kataoka HU, Wada J, Makino H. Intercellular adhesion molecule-1 plays a critical role in glomerulosclerosis after subtotal nephrectomy. *Clin Exp Nephrol*. 15:212-219, 2011.
 34. Miyatake N, Shikata K, Makino H, Numata T. Decreasing Abdominal Circumference Is Associated with Improving Estimated Glomerular Filtration Rate (eGFR) with Lifestyle Modification in Japanese Men: A Pilot Study. 1. *Acta Med Okayama*. 65:363-367, 2011.
 35. Tone A, Shikata K, Nakagawa K, Hashimoto M, Makino H. Renoprotective effects of clarithromycin via reduction of urinary MCP-1 levels in type 2 diabetic patients. *Clin Exp Nephrol*. 15:79-85, 2011.
 36. Miyatake N, Shikata K, Makino H, Numata T. The Relation between estimated glomerular filtration rate (eGFR) and proteinuria in Okayama prefecture, Japan. *Environ Health Prev Med* 16: 191-195, 2011.
 37. Miyatake N, Shikata K, Makino H, Numata T. Comparison of ventilatory threshold between subjects with and without proteinuria in Japanese. *Health* 3:

- 394-399, 2011.
38. Miyatake N, Shikata K, Makino H, Numata T: Decreasing serum uric acid levels might be associated with improving estimated glomerular filtration rate (eGFR) in Japanese men. *Health 3*: 498-503, 2011.
 39. Miyatake N, Nishii K, Numata T: Relationship between work style and cigarette smoking in Japanese workers. *Health 3*: 537-541, 2011.
 40. Miyatake N, Shikata K, Makino H, Numata T: The relation between estimated glomerular filtration rate (eGFR) and coffee consumption in the Japanese. *Health 3*: 549-552, 2011.
 41. Miyatake N, Shikata K, Makino H, Numata T: Comparison of muscle strength between subjects with and without proteinuria. *Health 3*: 698-702, 2011.
 42. Shikata K, Haneda M, Koya D, Suzuki Y, Tomino Y, Yamada K, Maeda S, Kawakami N, Uzu T, Nishimura M, Sato C, Ogawa D, Makino H; DNETT-Japan Study Group. Diabetic Nephropathy Remission and Regression Team Trial in Japan (DNETT-Japan): Rationale and study design. *Diabetes Res Clin Pract.* 2010 Feb;87(2):228-232.
 43. Miyatake N, Shikata K, Makino H, Numata T: Relationship between estimated glomerular filtration rate (eGFR) and metabolic syndrome in Japanese. *Acta Med Okayama* 64: 203-206, 2010.
 44. Miyatake N, Shikata K, Makino H, Numata T: Relation between estimated glomerular filtration rate and pulse wave velocity in Japanese. *Intern Med* 49: 1315-1320, 2010.
 45. Miyatake N, Shikata K, Makino H, Numata T: Decreasing systolic blood pressure is associated with improving estimated glomerular filtration rate (eGFR) with lifestyle modification in Japanese healthy women. *Acta Med Okayama* 64: 339-343, 2010.
 46. Kajitani N, Shikata K, Nakamura A, Nakatou T, Hiramatsu M, Makino H : Microinflammation is a common risk factor for progression of nephropathy and atherosclerosis in Japanese patients with type 2 diabetes. *Diabetes Res Clin Pract* 88: 171-176, 2010.
 47. Tone A, Iseda I, Higuchi C, Tsukamoto K, Katayama A, Matsushita Y, Hida K, Wada J, Shikata K. Comparison of insulin detemir and insulin glargine on glycemic variability in patients with type 1 and type 2 diabetes. *Exp Clin Endocrinol Diabetes.* 18:320-324. 2010.
 48. 総説: Iseki K. Editorials. Role of chronic kidney disease in cardiovascular disease: are we different from others? *Clin Exp Nephrol* 15:450-455, 2011
 49. Iseki K. Role of urinalysis in the diagnosis of chronic kidney disease (CKD). *JMAJ* 54: 27-30, 2011
 50. Ito S. Cardiorenal connection in chronic kidney disease *Clin. Exp Nephrol* DOI 10.1007/s10157-011-0493-2 Published online; 01 November 2011.
 51. Matsuo S, Yasuda Y, Imai E, Horio M. : Current status of estimated glomerular filtration rate (eGFR) equations for Asians and an approach to create a common eGFR equation. *Nephrology (Carlton).* 2010 Jun;15 Suppl 2:45-8
 52. Stevens LA, Claybon MA, Schmid CH, Chen J, Horio M, Imai E, Nelson RG, Van Deventer M, Wang HY, Zuo L, Zhang YL, Levey AS. : Evaluation of the Chronic Kidney Disease Epidemiology Collaboration equation for

- estimating the glomerular filtration rate in multiple ethnicities. *Kidney Int.* 79(5):555-62, 2010
53. Horio M, Yasuda Y, Takahara S, Imai E, Watanabe T, Matsuo S.: Comparison of a simple and a standard method for inulin renal clearance. *Clin Exp Nephrol.* 14(5):427-30, 2010
 54. Yamamoto R, Nagasawa Y, Shoji T, Iwatani H, Hamano T, Kawada N, Inoue K, Uehata T, Kaneko T, Okada N, Moriyama T, Horio M, Yamauchi A, Tsubakihara Y, Imai E, Rakugi H, Isaka Y.: Cigarette smoking and progression of IgA nephropathy. *Am J Kidney Dis.* 56(2):313-24, 2010
 55. Hosoya T, Okada H, Horio M, Ohno I, Tamura S, Sugimoto H, Tsushima Y, Hayashi H, Fukuda K, Korogi Y, Yoshifumi N.: Guidelines for administering gadolinium-based contrast agents to patients with renal dysfunction--second edition. *Nippon Jinzo Gakkai Shi.* 51(7):839-42, 2009
 56. Iseki K, Horio M, Imai E, Matsuo S, and Yamagata K. Geographic difference in prevalence of chronic kidney disease among Japanese screened subjects: Ibaraki vs. Okinawa. *Clin Exp Nephrol* 13:44-49, 2009
 57. Imai E, Horio M, Watanabe T, Iseki K, Yamagata K, Hara S, Ura N, Kiyohara Y, Moriyama T, Ando Y, Fujimoto S, Konta T, Yokoyama H, Makino H, Hishida A, and Matsuo S. Prevalence of chronic kidney disease (CKD) in Japanese general population. *Clin Exp Nephrol* 13: 621-630, 2009
 58. Yamagata K, Makino H, Akizawa T, Iseki K, Itoh S, Kimura K, Koya D, Narita I, Mitarai T, Miyazaki M, Tsubakihara Y, Watanabe T, Wada T, Sakai O, and Advisory Committee for FROM-J. Design and methods of a strategic outcome study for chronic kidney disease - Frontier of Renal Outcome Modifications in Japan (FROM-J) -*Clin Exp Nephrol* 14; 144-151, 2010
 59. Iseki K. Renal outcomes in chronic kidney disease. *Nephrology* 15: S273-S276, 2010
 60. Iseki K, Chiho Iseki, and Kozen Kinjo. C-reactive protein is a predictor for developing proteinuria in a screened cohort. *Nephron Clin Pract* 117: C51-C56, 2011
 61. Kohagura K, Tomiyama N, Kinjo K, Takishita S, Iseki K. Prevalence of anemia according to stage of chronic kidney disease in a large screening cohort of Japanese. *Clin Exp Nephrology* 13: 614-620, 2009
 62. Horio M, Imai E, Yasuda Y, Hishida A, Matsuo S on behalf of the collaborators for developing the Japanese equation for estimating GFR. Simple sampling strategy for measuring inulin renal clearance. *Clin Exp Nephrol* 13: 50-54, 2009.
 63. Iseki K, Horio M, Imai E, Matsuo S, Yamagata K.: Geographic difference in the prevalence of chronic kidney disease among Japanese screened subjects: Ibaraki versus Okinawa. *Clin Exp Nephrol.* 13(1):44-9. 2009
 64. Nakamura H, Kanemura T, Takara C, Tsukayama A, Tohyama K, Matsumoto T, Iseki K: A retrospective analysis of 4,000 patients with OSA in Okinawa, Japan. *Sleep and Biological Rhythms* 7: 103-112, 2009
 65. Koya D, Haneda M, Inomata S, Suzuki Y, Suzuki D, Makino H, Shikata K, Murakami Y, Tomino Y, Yamada K, Araki SI, Kashiwagi A, Kikkawa R; Low-Protein Diet Study Group. Long-term effect of modification of

- dietary protein intake on the p
progression of diabetic nephropathy: a
randomised controlled trial.
Diabetologia. 52:2037-2045、 2009
66. Sarai K, Shikata K, Shikata Y, Omori K,
Watanabe N, Sasaki M, Nishishita S, Wada
J, Goda N, Kataoka N, Makino H:
Endothelial barrier protection of FTY720
under hyperglycemic condition:
Involvement of focal adhesion kinase,
small GTPases and adherens junction
proteins. *Am J Physiol Cell Physiol*, 2009
67. Matsuo S, Imai E, Horio M, Yasuda Y, Tomita
K, Nitta K, Yamagata K, Tomino Y, Yokoyama
H, Hishida A. on behalf of the
collaborators developing the Japanese
equation for estimating GFR. Revised
Equations for Estimating Glomerular
Filtration Rate (GFR) from Serum
Creatinine in Japan. *Am J Kidney Dis*.
53:982-992, 2009.
68. Tsuda H, Isaka Y, Takahara S, Horio M.
Discrepancy between serum levels of low
molecular weight proteins in acute kidney
injury model rats with bilateral ureteral
obstruction and bilateral nephrectomy
Clin Exp Nephrol 13: 567-570, 2009.
69. Tanaka H, Hamano T, Fujii N, Tomida K,
Matsui I, Mikami S, Nagasawa Y, Ito T,
Moriyama T, Horio M, Imai E, Isaka Y,
Rakugi H. The impact of diabetes mellitus
on vitamin D metabolism in predialysis
patients. *Bone*. 45: 949-955, 2009.
70. Imai E, Horio M, Watanabe T, Iseki K,
Yamagata K, Hara S, Ura N, Kiyohara Y,
Moriyama T, Ando Y, Fujimoto S, Konta T,
Yokoyama H, Makino H, Hishida A, Matsuo S.
Prevalence of chronic kidney disease in
the Japanese general population. *Clin Exp
Nephrol*. 13 : 621-630, 2009.
71. Kakuta Y, Okumi M, Ichimaru N, Abe T,
Nonomura N, Okuyama A, Kojima Y, Isaka Y,
Takahara S, Imai E, Horio M. Utility of the
Japanese GFR estimation equation for
evaluating potential donor kidney
function. *Clin Exp Nephrol*.
14(1):63-7. 2009
- 著書
1. Rayner H, Imai E: Approach to Renal
Replacement Therapy, “COMPREHENSIVE
CLINICAL NEPHROLOGY” , 4th Edition edited
by Floege J, Johnson RJ, and Feehally J
Elsevier Inc. p 1019-1030, 2010
- 学会発表
1. Ambulatory Blood Pressure Monitoring
(ABPM) in the CKD Japan Cohort (CKD-JAC)
Study. Enyu Imai, Satoshi Iimuro, Kosaku
Nitta, Tsuyoshi Watanabe, Seiichi
Matsuo, , Tadao Akizawa, Hirofumi Makino,
Yasuo Ohashi, Akira Hishida The American
Society of Nephrology 44th Annual
Meeting(Philadelphia) Nov 10, 2011
2. Erythropoietin Responses in Chronic
Kidney Disease: A Cross-Sectional
Analysis from the CKD-JAC Study. Tadao
Akizawa, Takayuki Hamano, Enyu Imai,
Seiichi Matsuo, Hirofumi Makino,
Tsuyoshi Watanabe, Kosaku Nitta, Yasuo
Ohashi, Akira Hishida The American
Society of Nephrology 44th Annual
Meeting(Philadelphia) Nov 10, 2011
3. Low Serum Cultured Adipose-Derived
Mesenchymal Stem Cells, but Not
Bone-Marrow Derived Mesenchymal Stem
Cells, Ameliorate Rat Crescentic
Glomerulonephritis by Functional
Polarization of Macrophages into
Immunoregulatory Phenotype. Kazuhiro

- Furuhashi, Naotake Tsuboi, PhD, Hangsoo Kim, Takayuki Katsuno, Takenori Ozaki, Waichi Sato, Enyu Imai, Seiichi Matsuo, Shoichi Maruyama The American Society of Nephrology 44th Annual Meeting(Philadelphia) Nov 10, 2011
4. The Relationship between Minnesota Code Findings and CKD: Result from the CKD-JAC Study. Satoshi Iimuro, Takeo Okada, Enyu Imai, Kosaku Nitta, Tsuyoshi Watanabe, Seiichi Matsuo, Tadao Akizawa, Hirofumi Makino, Yasuo Ohashi, Akira Hishida The American Society of Nephrology 44th Annual Meeting(Philadelphia) Nov 10, 2011
 5. Prognostic Value of Geriatric Nutritional Risk Index for Cardiovascular and All-Cause Mortality in End-Stage Renal Disease Patients on Chronic Hemodialysis. Kaoru Yasuda, Shoichi Maruyama, Hirotake Kasuga, Tomoki Kosugi, Naotake Tsuboi, Waichi Sato, Yasuhiko Ito, Enyu Imai, Seiichi Matsuo The American Society of Nephrology 44th Annual Meeting(Philadelphia) Nov 10, 2011
 6. Association of Adipokines with Cardiovascular Mortality in Patients on Hemodialysis. Shoichi Maruyama, Kaoru Yasuda, Hirotake Kasuga, Yoshinari Yasuda, Tomoki Kosugi, Waichi Sato, Naotake Tsuboi, Yasuhiko Ito, Enyu Imai, Seiichi Matsuo The American Society of Nephrology 44th Annual Meeting(Philadelphia) Nov 10, 2011
 7. Low Serum Cultured Adipose Derived Stem Cells Ameliorate Zymosan Induced Severe Rat Peritonitis Model. Hangsoo Kim, Masashi Mizuno, Kazuhiro Furuhashi, Takayuki Katsuno, Kaoru Yasuda, Takenori Ozaki, Waichi Sato, Naotake Tsuboi, Yasuhiko Ito, Enyu Imai, Shoichi Maruyama, Seiichi Matsuo The American Society of Nephrology 44th Annual Meeting(Philadelphia) Nov 10, 2011
 8. The Structure of Health-Related QOL Varies According to Underlying Diseases of CKD: Results from the CKD-JAC Study. Yasuo Ohashi, Satoshi Iimuro, Tadao Akizawa, Enyu Imai, Seiichi Matsuo, Tsuyoshi Watanabe, Kosaku Nitta, Hirofumi Makino, Akira Hishida The American Society of Nephrology 44th Annual Meeting(Philadelphia) Nov 10, 2011
 9. Clinical Impact of the Examination of the Ocular Fundus for CKD Patients without Atherosclerosis Risk Factors. Yoshinari Yasuda, Kiyoshi Shibata, Sadao Suzuki, Sawako Kato, Shoichi Maruyama, Enyu Imai, Seiichi Matsuo The American Society of Nephrology 44th Annual Meeting(Philadelphia) Nov 10, 2011
 10. A Comparison of the Acute Kidney Injury Network and Kidney Disease: Improving Global Outcomes Criteria for Acute Kidney Injury in Critically Ill Patients. Hibiki Shinjo, Waichi Sato, Tomoki Kosugi, Hiroki Hayashi, Shoichi Maruyama, Enyu Imai, Yukio Yuzawa, Seiichi Matsuo The American Society of Nephrology 44th Annual Meeting(Philadelphia) Nov 10, 2011
 11. Lymphangiogenesis Develops during Peritoneal Fibrosis in Peritoneal Dialysis Patients and Rat Peritonitis Model. Hiroshi Kinashi, Yasuhiko Ito, MD, Masashi Mizuno, Yasuhiro Suzuki, Fumiko Nagura, Waichi Sato, Naotake Tsuboi, Shoichi Maruyama, Enyu Imai, Yoshifumi Takei, Seiichi Matsuo The American Society of Nephrology 44th Annual Meeting(Philadelphia) Nov 10, 2011
 12. Association of Cardiac Valve

- Calcification and C-Reactive Protein (CRP) with Cardiovascular and All-Cause Mortality in End-Stage Renal Disease Patients – 10-Year Follow-Up Study from Induction of Hemodialysis Therapy. Yasuhiko Ito, Hirotake Kasuga, Masashi Mizuno, Keiko Kimura, Shoichi Maruyama, Enyu Imai, Seiichi Matsuo The American Society of Nephrology 44th Annual Meeting (Philadelphia) Nov 10, 2011
13. A Candidate Gene Approach to Genetic Contributors to Development of IgA Nephropathy. Ryohei Yamamoto, Yasuyuki Nagasawa, Matsuya Shoji, Tetsuya Kaneko, Kazunori Inoue, Hirotatsu Iwatani, Enyu Imai, Atsushi Yamauchi, Yoshiharu Tsubakihara, Hiromi Rakugi, Yoshitaka Isaka The American Society of Nephrology 44th Annual Meeting (Philadelphia) Nov 10, 2011
 14. Evaluation of Diverticular Diseases by Computed Tomography Is Useful for Selection of Peritoneal Dialysis Modality and Asymptomatic Diverticulosis Is Not a Risk Factor for Peritonitis. Yasuhiko Ito, Susumu Toda, Masashi Mizuno, Yasuhiro Suzuki, Isao Ito, Hideki Hiramatsu, Takenori Ozaki, Waichi Sato, Naotake Tsuboi, Shoichi Maruyama, Enyu Imai, Seiichi Matsuo The American Society of Nephrology 44th Annual Meeting (Philadelphia) Nov 10, 2011
 15. Imai E: Regional variation in GFR. Detection of CKD: International perspectives. World Congress of Nephrology April 10, 2011, Vancouver.
 16. Imai E: Comparisons between Japan and Taiwan based on CKD/ESRD risk factors. “2011 Kidney Health Forum” January 15–16, Zhunan Town, Taiwan, R. O. C.
 17. Imai E: Natural History of CKD and Cohort Study The First International Symposium on the Non-Communicable Disease (NCDs). Osong City, Korea. March 16, 2011.
 18. Imai E: Anemia in renal disease: diagnosis and management. the 5th AFCKDI Guangzhou P.R. China March 18, 2011
 19. Horio M, Simple sampling strategy for measuring inulin renal clearance. The 3rd Research Forum on Chronic Kidney Disease. Nagoya, Feb 7, 2009.
 20. Horio M, Imai E, Yasuda Y, Watanabe T, Matsuo S. Modification of the CKD-EPI equation for Japanese; Accuracy and use for population estimates. The American Society of Nephrology 43th Annual Meeting Denver Nov 18, 2010
 21. Horio M, GFR equations from serum cystatin-C alone and in combination with serum creatinine. The 5th Research Forum on Chronic Kidney Disease. Nagoya, Feb 12, 2011.
 22. Yasuda Y, Horio M, Chen HC, Lee HY, Imai E, Matsuo S. Validation of MDRD and Japanese eGFR Equation among Taiwanese and Korean: Approach To Set the Fundamental Scheme for eGFR Evaluation in Asia. The American Society of Nephrology 44th Annual Meeting Philadelphia Nov 11, 2011
 23. Iseki K Changes in serum uric acid have reciprocal effect on eGFR change: a 10-year follow-up study of community-based screening in Okinawa, Japan. 44th Annual Meeting Philadelphia Nov 11, 2011
 24. Iseki K, Iseki C, Watanabe T. Relation between glomerular filtration rate (GFR) and medical expense among the screened subjects of the Japan Health Insurance

- Association. 44th Annual Meeting
Philadelphia Nov 11, 2011
25. Yamamoto T, Nakayama M, Miyazaki M, Sato H, Sato T, Ito S. Rapid Progression of Chronic Kidney Disease Is Associated with Cardiovascular Mortality in the Japanese Population: The Gonryo Study 44th Annual Meeting Philadelphia Nov 11, 2011
 26. Kodera R, Kataoka H, Shikata K, Makino H. • Exendin-4 ameliorates renal injuries through anti-inflammatory effects in type 1 diabetic rats. Asia Islet Biology and Incretin Symposium (AIBIS). August 1-2, 2009
 27. Miyamoto S, Shikata K, Sasaki M, Nishishita S, Sato C, Kataoka H, Kodera R, Hirota D, Kajitani N, Makino H. Cholecystokinin ameliorates diabetic renal injuries via anti-inflammatory effects. 45th EASD Annual Meeting, September 29-October 2, 2009
 28. Kodera R, Kataoka H, Shikata K, Sato C, Sasaki M, Nishishita S, Miyamoto S, Hirota D, Kajitani N, Ogawa D, Makino H. • Exendin-4 ameliorates renal injuries through anti-oxidative and anti-inflammatory effects in type 1 diabetic rats. 45th EASD Annual Meeting. September 30-October 2, 2009.
 29. Kodera R, Kataoka H, Shikata K, Sato C, Sasaki M, Nishishita S, Miyamoto S, Hirota D, Kajitani N, Ogawa D, Makino H. Exendin-4 ameliorates renal injuries through anti-oxidative and anti-inflammatory effects in type 1 diabetic rats. 46th EASD Annual Meeting Stockholm, Sweden, September 21-24, 2010
 30. Ogawa D, Matsushita Y, Shikata K, Wada J, Sato C, Makino H. PPAR δ -mediated anti-inflammatory mechanisms inhibit streptozotocin-induced diabetic nephropathy. Keystone Symposia, Banff, Alberta, Canada. February 12-17, 2010
 31. Ogawa D, Matsushita Y, Shikata K, Wada J, Sato C, Makino H. PPAR δ -mediated anti-inflammatory mechanisms inhibit streptozotocin-induced diabetic nephropathy. ISN-NEXUS Kyoto in 2010. Kyoto, April 15-18, 2010
 32. Ogawa D, Matsushita Y, Shikata K, Wada J, Sato C, Makino H. PPAR δ -mediated anti-inflammatory mechanisms inhibit streptozotocin-induced diabetic nephropathy. The 2nd Scientific Meeting of the Asian Association for the Study of Diabetes. May 29, 2010.
 33. Kajitani N, Shikata K, Kodera R, Miyamoto S, Hirota D, Sato C, Wada J, Ogawa D, and Makino H. Helmsartan attenuates diabetic nephropathy through anti-oxidative and anti-inflammatory actions via activation of peroxisome proliferator-activated receptor- γ . 47th Annual Meeting of the European Association for the Study of Diabetes Lisbon Portugal, 12-16 September 2011.
 34. Ogawa D, Wada J, Matsushita Y, Nakatsuka A, Murakami K, Kanzaki M, Shikata K, Makino H. Activation of Peroxisome Proliferator-activated Receptor δ Inhibits Streptozotocin-induced Diabetic Nephropathy Through Anti-inflammatory Mechanisms in Mice. Keystone Symposia (Keystone, Colorado, USA) 2011. 1. 12-1. 17
 35. Tachibana H, Ogawa D, Matsushita Y, Yamamoto N, Toyota N, Sato C, Wada J, Shikata K, Makino H. Activation of Peroxisome Proliferator-activated

Receptor δ Inhibits
Streptozotocin-induced Diabetic
Nephropathy Through Anti-inflammatory
Mechanisms in Mice. American Diabetes
Association 71st Scientific Sessions (San
Diego, California, USA) 2011.6.24-6.28

36. 4th AFCKDI. Iseki K. Pan Asian CKD
registry: Background and Japanese CKD
registry. 2010,6/4 Seoul
37. APCN2010. Iseki K. Chronic kidney
disease: from early prediction to current
facts. 2010,6/5 Seoul
38. KDIGO Controversies Conference: CKD and
CVD. 2010,10/29 London

8 知的所有権の出願・取得状況（予定を含む）

- 1 特許取得 なし
- 2 実用新案登録 なし
- 3 その他 なし

厚生労働科学研究費補助金（腎疾患対策研究事業）

班 員 名 簿

課題		氏名	所属
	代表者	今井 圓裕	名古屋大学腎臓内科
CKD早期発見に必要な腎機能推算式の開発	分担者	堀尾 勝	大阪大学医学系研究科保健学専攻機能診断科学講座
	協力者	猪阪 善隆	大阪大学腎臓内科
	協力者	安田 宜成	名古屋大学腎臓内科
腎不全発症率の地域格差の研究	分担者	井関 邦敏	琉球大学医学部附属病院血液浄化療法部
	協力者	山縣 邦弘	筑波大学腎臓内科
	協力者	今田 恒夫	山形大学腎臓内科
	協力者	鶴屋 和彦	九州大学腎疾患治療部
	協力者	渡辺 毅	福島県立医科大学第三内科
	協力者	藤元 昭一	宮崎大学第一内科
CKD患者におけるCVDの危険因子の同定	分担者	伊藤 貞嘉	東北大学大学院医学系研究科腎・高血圧・内分泌学
	協力者	宮崎 真理子	東北大学大学院医学系研究科腎・高血圧・内分泌学
	協力者	佐藤 博	東北大学大学院薬学研究科臨床薬学分野
	協力者	佐藤 壽伸	仙台社会保険病院
	協力者	中山 昌明	福島県立医科大学腎臓高血圧・糖尿病内分泌代謝内科学
	協力者	竹内 和久	宏人会 中央クリニック
	協力者	田熊 淑男	仙台社会保険病院
	協力者	秋保 直樹	仙台市立病院
	協力者	小原 克也	東北公済病院
	協力者	栗原 功	東北公済病院 宮城野分院
	協力者	山口 裕二	仙台赤十字病院
	協力者	清野 仁	仙台医療センター
	協力者	太田 耕造	大崎市民病院
	協力者	木下 康道	石巻赤十字病院
	協力者	佐藤 和人	石巻赤十字病院
	協力者	大高 徹也	公立総合刈田病院
生活習慣病のCKD発症への危険因子の同定	分担者	四方賢一	岡山大学病院 新医療研究開発センター
	協力者	和田 淳	岡山大学大学院医歯薬学総合研究科 腎・免疫・内分泌代謝内科学
	協力者	梶谷展生	岡山大学大学院医歯薬学総合研究科 腎・免疫・内分泌代謝内科学
	協力者	利根惇仁	国立岡山医療センター 糖尿病内科
	協力者	肥田和之	国立岡山医療センター 糖尿病内科
	協力者	石井啓太	公立学校共済組合中国中央病院 内科
	協力者	平田教至	公立学校共済組合中国中央病院 内科
	協力者	宮武伸行	香川大学医学部衛生学
	協力者	中村明彦	おさふねクリニック
腎硬化症のリスク因子の解明	分担者	富田 公夫	熊本大学大学院生命科学研究部 腎臓内科
	協力者	白石 直樹	熊本大学大学院生命科学研究部 腎臓内科
高齢者への薬剤投与とCKD対策が及ぼす効果の評価	分担者	細谷 龍男	東京慈恵会医科大学腎臓・高血圧内科
	協力者	大野 岩男	東京慈恵会医科大学腎臓・高血圧内科
	協力者	宇都宮 保典	東京慈恵会医科大学腎臓・高血圧内科
	協力者	宮崎 陽一	東京慈恵会医科大学腎臓・高血圧内科
	協力者	川村 哲也	東京慈恵会医科大学腎臓・高血圧内科
	協力者	小倉 誠	東京慈恵会医科大学腎臓・高血圧内科
	協力者	池田 雅人	東京慈恵会医科大学腎臓・高血圧内科

厚生労働科学研究費補助金（腎疾患対策研究事業）

研究成果の刊行物・別刷

研究成果の刊行に関する一覧表

書籍

著者氏名	論文タイトル名	書籍全体の 編集者名	書 籍 名	出版社 名	出版地	出版年	ページ
梶谷展生, 四方賢一	糖尿病性細小血管症(第2版)-発症 ・進展制御の最前線- II. 糖尿病性細小血管症の発症 ・進展の分子メカニズム Microinflammationの関与.		日本臨床	日本臨床社	東京	61-64	2010
廣田大昌, 四方賢一	特集 糖尿病合併症とその治療に関するエポックメイキングトピックスの展開 腎症の寛解は可能か? : 最近のエビデンスより.		糖尿病診療マスター	医学書院	東京	403-406	2010
廣田大昌, 四方賢一	特集: 糖尿病と慢性腎臓病(CKD) 糖尿病におけるCKDの成因と治療.		プラクティス	医歯薬出版社	東京	272-280	2010
小寺亮, 四方賢一	特集 腎疾患治療法開発の最前線~糖尿病性腎症治療薬開発の動向~.		細胞	ニュー・サイエンス社	東京	147-150	2010
小寺亮, 四方賢一	新たなステージへ進む糖尿病診療 各論: DPP-4阻害薬の登場で薬物療法はどう変わるか		月刊薬事	じほう	東京	345-349	2010
小寺亮, 四方賢一	糖尿病性細小血管症-III 糖尿病性腎症の病理 (Pathology of diabetic nephropathy).		日本臨床	日本臨床社	東京	370-374	2010
四方賢一	VIII 糖尿病腎症 4. 腎不全期.	難波光義、相澤徹、池上博司、西村理明、浜口朋也、宮川潤一郎、森豊	糖尿病最新の治療	南江堂	東京	2010-2012	2010

四方賢一	新しい時代の糖尿病治療 ～インクレチン製剤への期待～.		山県郡医師会会報	山県郡医師会	広島	46-47	2010
四方賢一、槇野博史	シンポジウム：慢性合併症の臨床2（腎症）1 腎症の成因.	日本糖尿病学会	糖尿病学の進歩（第44集）	診断と治療社	東京		2010
四方賢一、槇野博史	糖尿病腎症の形態的アプローチ.		糖尿病ナビゲーター 第2版	メディカルレビュー社	大阪		2010
四方賢一	病期から見た合併症対策 発言1 腎症の病期に応じた対応のポイント.	堀田饒、清野裕、門脇孝、柏木厚典、中村二郎	対糖尿病合併症対策のイノベーション 成因から管理、治療まで.	時事通信出版局	東京	196-203	2010
南条輝志男、羽田勝計、四方賢一、高木均、中村二郎	病期から見た合併症対策 総合討論.	堀田饒、清野裕、門脇孝、柏木厚典、中村二郎	対糖尿病合併症対策のイノベーション 成因から管理、治療まで.	時事通信出版局	東京	222-235	2010
松下裕一、四方賢一	糖尿病性腎症：晩期糖尿病の管理と注意点.	槇野博史、秋澤忠男	腎疾患・透析 最新の治療 2011-2013	南江堂	東京	2011	179-183
今井圓裕、安田宜成、松尾清一	CKD運動の意義とCKD患者の動向		臨牀透析	日本メディカルセンター	東京	2011	9-16
今井圓裕、松尾清一	慢性腎症の抗原		腎臓	日本腎臓財団	東京	2011	1-310
今井圓裕	慢性腎症		医学と薬学	自然科学社	東京	2011	149-155
今井圓裕、友雅司	第55回日本透析医学会・第53回日本腎臓学会共催シンポジウムより		日本透析医学会雑誌	日本透析医学会	東京	2011	123-136
今井圓裕	『透析導入基準』		progress in Medicine	ライフ・サイエンス	東京	2011	389-394
今井圓裕	疫学と疾患としての重要性		腎と透析	東京医学社	東京	2011	236-239
今井圓裕	海外の腎性貧血ガイドライン		医学のあゆみ	医歯薬出版	東京	2011	301-302

今井圓裕	ステロイドパルス療法		総合臨牀	永井書店	大阪	2011	1320-1328
------	------------	--	------	------	----	------	-----------

雑誌

発表者氏名	論文タイトル名	発表誌名	巻号	ページ	出版年
Miyazawa N, Abe M, Souma T, Tanemoto M, Abe T, Nakayama M, Ito S.	Methylglyoxal augments intracellular oxidative stress in human aortic endothelial cells.	Free Radic Res.	44(1)	101-7	2010
Nakayama M, Kabayama S, Nakano H, Zhu WJ, Terawaki H, Nakayama K, Katoh K, Satoh T, Ito S.	Biological effects of electrolyzed water in hemodialysis.	Nephron Clin Pract.	c9-15	112(1)	2009
Guo Q, Mori T, Jiang Y, Hu C, Osaki Y, Yoneki Y, Sun Y, Hosoya T, Kawamata A, Ogawa S, Nakayama M, Miyata T, Ito S.	Methylglyoxal contributes to the development of insulin resistance and salt sensitivity in Sprague-Dawley rats.	J Hypertens.	27(8)	1664-71	2009
Nakayama K, Nakayama M, Terawaki H, Murata Y, Sato T, Kohno M, Ito S.	Carbonated soft drinks and carbonyl stress burden	J Toxicol Sci.	34(6)	699-702	2009

<p>Toyohara T, Suzuki T, Morimoto R, Akiyama Y, Souma T, Shiwaku HO, Takeuchi Y, Mishima E, Abe M, Tanemoto M, Masuda S, Kawano H, Maemura K, Nakayama M, Sato H, Mikkaichi T, Yamaguchi H, Fukui S, Fukumoto Y, Shimokawa H, Inui K, Terasaki T, Goto J, Ito S, Hishinuma T, Rubera I, Tauc M, Fujii- Kuriyama Y, Yabuuchi H, Moriyama Y, Soga T, Abe T.</p>	<p>SLC04C1 transporter eliminates uremic toxins and attenuates hypertension and renal inflammation.</p>	<p>J Am Soc Nephrol.</p>	<p>20(12)</p>	<p>2546-55</p>	<p>2009</p>
<p>Matsuo S, Imai E, Horio M, Yasuda Y, Tomita K, Nitta K, Yamagata K, Tomino Y, Yokoyama H, Hishida A. on behalf of the collaborators developing the Japanese equation for estimating GFR</p>	<p>Revised Equations for Estimating Glomerular Filtration Rate (GFR) from Serum Creatinine in Japan</p>	<p>Am J Kidney Dis</p>	<p>53</p>	<p>982-992</p>	<p>2009</p>
<p>Tsuda H, Isaka Y, Takahara S, Horio M</p>	<p>Discrepancy between serum levels of low molecular weight proteins in acute kidney injury model rats with bilateral ureteral obstruction and bilateral nephrectomy</p>	<p>Clin Exp Nephrol</p>	<p>13</p>	<p>567-570</p>	<p>2009</p>

Tanaka H, Hamano T, Fujii N, Tomida K, Matsui I, Mikami S, Nagasawa Y, Ito T, Moriyama T, Horio M, Imai E, Isaka Y, Rakugi H	The impact of diabetes mellitus on vitamin D metabolism in predialysis patients	Bone	45	949-955	2009
Imai E, Horio M, Watanabe T, Iseki K, Yamagata K, Hara S, Ura N, Kiyohara Y, Moriyama T, Ando Y, Fujimoto S, Konta T, Yokoyama H, Makino H, Hishida A, Matsuo S	Prevalence of chronic kidney disease in the Japanese general population	Clin Exp Nephrol	13	621-630	2009
Kakuta Y, Okumi M, Ichimaru N, Abe T, Nonomura N, Okuyama A, Kojima Y, Isaka Y, Takahara S, Imai E, Horio M	Utility of the Japanese GFR estimation equation for evaluating potential donor kidney function	Clin Exp Nephrol			2009
Shikata K, Haneda M, Makino H, et al	DNETT-Japan Study Group. Diabetic Nephropathy Remission and Regression Team Trial in Japan (DNETT-Japan): Rationale and study design.	Diabetes Res Clin Pract.	In press		
Koya D, Haneda M, Inomata S, Shikata K, Murakami Y, Kashiwagi A, Kikkawa R, Et al	Long-term effect of modification of dietary protein intake on the progression of diabetic nephropathy: a randomised controlled trial.	Diabetologia.	52	2037-2045.	2009
Sarai K, Shikata K, Shikata Y, Omori K, Watanabe N, Sasaki M, Nishishita S, Wada J, Goda N, Kataoka N, Makino H.	Endothelial barrier protection by FTY720 under hyperglycemic condition: involvement of focal adhesion kinase, small GTPases, and adherens junction proteins.	Am J Physiol Cell Physiol.	945-954.	945-954.	2009

Horio M, Imai E, Yasuda Y, Watanabe T, Matsuo S.	Modification of the CKD Epidemiology Collaboration (CKD-EPI) Equation for Japanese: Accuracy and Use for Population Estimates.	Am J Kidney Dis.	56(1)	32-38	2010
Stevens LA, Claybon MA, Schmid CH, Chen J, Horio M, Imai E, Nelson RG, Van Deventer M, Wang HY, Zuo L, Zhang YL, Levey AS.	Evaluation of the Chronic Kidney Disease Epidemiology Collaboration equation for estimating the glomerular filtration rate in multiple ethnicities.	Kidney Int.	in press		2010
Horio M, Yasuda Y, Takahara S, Imai E, Watanabe T, Matsuo S.	Comparison of a simple and a standard method for inulin renal clearance.	Clin Exp Nephrol.	14(5)	427-430	2010
Matsuo S, Yasuda Y, Imai E, Horio M.	Current status of estimated glomerular filtration rate (eGFR) equations for Asians and an approach to create a common eGFR equation.	Nephrology (Carlton).	Suppl 2	45-48	2010
Yamamoto R, Nagasawa Y, Shoji T, Iwatani H, Hamano T, Kawada N, Inoue K, Uehata T, Kaneko T, Okada N, Moriyama T, Horio M, Yamauchi A, Tsubakihara Y, Imai E, Rakugi H, Isaka Y.	Cigarette smoking and progression of IgA nephropathy.	Am J Kidney Dis.	56(2)	313-324	2010
Hosoya T, Okada H, Horio M, Ohno I, Tamura S, Sugimoto H, Tsushima Y, Hayashi H, Fukuda K, Korogi Y, Yoshifumi N.	Guidelines for administering gadolinium-based contrast agents to patients with renal dysfunction--second edition.	Nippon Jinzo Gakkai Shi.	51(7)	839-842	2009
Kakuta Y, Okumi M, Ichimaru N, Abe T, Nonomura N, Okuyama A, Kojima Y, Isaka Y, Takahara S, Imai E, Horio M.	Utility of the Japanese GFR estimation equation for evaluating potential donor kidney function.	Clin Exp Nephrol.	14(1)	63-67	2010