

CHS: Michael Shlipak, Mark J Sarnak, Ronit Katz, and Linda P Fried; **HUNT:** Stein Hallan, Stian Lydersen, and Jostein Holmen; **KP-Hawaii Cohort:** Brian J Lee; **MESA:** Michael Shlipak, Mark J Sarnak, Ronit Katz, and Linda P Fried; **MR-FIT:** Areef Ishani, James Neaton, and Ken Svendsen; **OGHMA-OKID 83 and 93:** Kunitoshi Iseki; **ONTARGET:** Johannes FE Mann, Salim Yusuf, Koon K. Teo, and Peggy Gao; **Pima Indian:** Robert G Nelson and William C Knowler; and **TRANSCEND:** Johannes FE Mann, Salim Yusuf, Koon K Teo, and Peggy Gao.

CKD Prognosis Consortium Analytic Team: Brad C Astor, Priscilla Auguste, Josef Coresh, Ron T Gansevoort, Paul E de Jong, Kunihiro Matsushita, Marije van der Velde, Kasper Veldhuis, Yaping Wang, and Mark Woodward.

CKD Prognosis Consortium Administration Staff: Laura Camarata and Beverly Thomas.

National Kidney Foundation Staff: Tom Manley.

SUPPLEMENTARY MATERIAL

Table S1. Acronyms/abbreviations for individual studies.

Table S2. Incidence rate for end-stage renal disease.

Table S3. Incidence rate for acute kidney injury.

Table S4. Incidence rate for progressive chronic kidney disease.

Table S5. Distribution of subjects for analysis of incident end-stage renal disease.

Table S6. Distribution of incident end-stage renal disease events.

Table S7. Distribution of incident acute kidney injury events.

Table S8. Distribution of incident progressive chronic kidney disease events.

Table S9. Statistical significance for interaction between eGFR and age, and between eGFR and albuminuria for end-stage renal disease, acute kidney injury, and progressive chronic kidney disease.

Table S10. Hazard ratios for incident end-stage renal disease.

Table S11. Hazard ratios for incident acute kidney injury.

Table S12. Odds ratios for incident progressive chronic kidney disease.

Figure S1. Pooled adjusted hazard ratios for acute kidney injury according to spline eGFR and albumin-to-creatinine ratio adjusted for each other and for age, sex, and cardiovascular risk factors.

Figure S2. Pooled adjusted hazard ratios for progressive chronic kidney disease according to spline eGFR and albumin-to-creatinine ratio adjusted for each other and for age, sex, and cardiovascular risk factors.

Figure S3. Pooled adjusted hazard ratios for end-stage renal disease according to eGFR and albuminuria for four groups (general population cohorts with albumin-to-creatinine ratio data, general population cohorts with dipstick data, high-risk cohorts with albumin-to-creatinine ratio data, and high-risk cohorts with dipstick data).

Supplementary material is linked to the online version of the paper at <http://www.nature.com/ki>

REFERENCES

- Eckardt KU, Berns JS, Rocco MV *et al.* Definition and classification of CKD: the debate should be about patient prognosis—a position statement from KDOQI and KDIGO. *Am J Kidney Dis* 2009; **53**: 915–920.
- Levey AS, de Jong PE, Coresh J *et al.* Chronic kidney disease - definition, classification and prognosis: a KDIGO controversies conference reaches a consensus. *Kidney Int* 2010; **375**: 2073–2081.
- National Kidney Foundation. K/DOQI clinical practice guidelines for chronic kidney disease: evaluation, classification, and stratification. *Am J Kidney Dis* 2002; **39**: S1–266.
- Levey AS, Coresh J, Balk E *et al.* NKF practice guidelines for CKD: evaluation, classification and stratification. *Arch Int Med* 2003; **139**: 137–147.
- Levey AS, Eckardt KU, Tsukamoto Y *et al.* Definition and classification of chronic kidney disease: a position statement from Kidney Disease: Improving Global Outcomes (KDIGO). *Kidney Int* 2005; **67**: 2089–2100.
- Levey AS, Atkins R, Coresh J *et al.* Chronic kidney disease as a global public health problem: approaches and initiatives - a position statement from Kidney Disease Improving Global Outcomes. *Kidney Int* 2007; **72**: 247–259.
- Gansevoort RT, de Jong PE. The case for using albuminuria in staging chronic kidney disease. *J Am Soc Nephrol* 2009; **20**: 465–468.
- Glasscock RJ, Winearls C. An epidemic of chronic kidney disease: fact or fiction? *Nephrol Dial Transpl* 2008; **23**: 1117–1123.
- Ikizler TA. CKD classification: time to move beyond KDOQI. *J Am Soc Nephrol* 2009; **20**: 929–930.
- Wetzels JF, Willems HL, den Heijer M. Age- and gender-specific reference values of estimated glomerular filtration rate in a Caucasian population: results of the Nijmegen Biomedical Study. *Kidney Int* 2008; **73**: 657–658.
- Winearls CG, Glasscock RJ. Dissecting and refining the staging of chronic kidney disease. *Kidney Int* 2009; **75**: 1009–1014.
- Ruggenti P, Perna A, Mosconi L *et al.* Urinary protein excretion rate is the best independent predictor of ESRF in non-diabetic proteinuric chronic nephropathies. 'Gruppo Italiano di Studi Epidemiologici in Nefrologia' (GISEN). *Kidney Int* 1998; **53**: 1209–1216.
- Keane WF, Zhang Z, Lyle PA, *et al.* Brenner BM, RENAAL Study Investigators. Risk scores for predicting outcomes in patients with type 2 diabetes and nephropathy: the RENAAL study. *Clin J Am Soc Nephrol* 2006; **1**: 761–767.
- Jafar TH, Stark PC, Schmid CH, *et al.* Levey AS, AIPRD Study Group. Progression of chronic kidney disease: the role of blood pressure control, proteinuria, and angiotensin-converting enzyme inhibition: a patient-level meta-analysis. *Ann Intern Med* 2003; **139**: 244–252.
- Wen CP, Cheng TY, Tsai MK *et al.* All-cause mortality attributable to chronic kidney disease: a prospective cohort study based on 462 293 adults in Taiwan. *Lancet* 2008; **371**: 2173–2182.
- Go AS, Chertow GM, Fan D *et al.* Chronic kidney disease and the risks of death, cardiovascular events, and hospitalization. *N Engl J Med* 2004; **351**: 1296–1305.
- Hallan S, Astor B, Romundstad S *et al.* Association of kidney function and albuminuria with cardiovascular mortality in older vs younger individuals: the HUNT II Study. *Arch Intern Med* 2007; **167**: 2490–2496.
- Astor BC, Hallan SI, Miller III ER *et al.* Glomerular filtration rate, albuminuria, and risk of cardiovascular and all-cause mortality in the US population. *Am J Epidemiol* 2008; **167**: 1226–1234.
- Brantsma AH, Bakker SJ, Hillege HL, *et al.* Gansevoort RT, PREVEND Study Group. Cardiovascular and renal outcome in subjects with K/DOQI stage 1–3 chronic kidney disease: the importance of urinary albumin excretion. *Nephrol Dial Transplant* 2008; **23**: 3851–3858.
- Hemmelgarn BR, Manns BJ, Lloyd A *et al.* Relation between kidney function, proteinuria, and adverse outcomes. *JAMA* 2010; **303**: 423–429.
- van der Velde M, Halbesma N, de Charro FT *et al.* Screening for albuminuria identifies individuals at increased renal risk. *J Am Soc Nephrol* 2009; **20**: 852–862.
- Hallan SI, Ritz E, Lydersen S *et al.* Combining GFR and albuminuria to classify CKD improves prediction of ESRD. *J Am Soc Nephrol* 2009; **20**: 1069–1077.
- Coca SG. Long-term outcomes of acute kidney injury. *Curr Opin Nephrol Hypertens* 2010; **19**: 266–272.
- Goldberg R, Dennen P. Long-term outcomes of acute kidney injury. *Adv Chronic Kidney Dis* 2008; **15**: 297–307.
- The Chronic Kidney Disease Prognosis Consortium. Association of estimated glomerular filtration rate and albuminuria with all-cause and cardiovascular mortality: a collaborative meta-analysis of general population cohorts. *Lancet* 2010; **375**: 2073–2081.
- The Chronic Kidney Disease Prognosis Consortium. Association of estimated glomerular filtration rate and albuminuria with all-cause and cardiovascular mortality: a collaborative meta-analysis of high risk cohorts. *Kidney Int.* (submitted).
- The Chronic Kidney Disease Prognosis Consortium. Association of estimated glomerular filtration rate and albuminuria with mortality and end-stage renal disease: a collaborative meta-analysis of kidney disease cohorts. *Kidney Int.* (submitted).
- Iseki K, Kinjo K, Iseki C *et al.* Relationship between predicted creatinine clearance and proteinuria and the risk of developing ESRD in Okinawa, Japan. *Am J Kidney Dis* 2004; **44**: 806–814.
- Ishani A, Grandits GA, Grimm RH *et al.* Association of single measurements of dipstick proteinuria, estimated glomerular filtration rate, and hematocrit with 25-year incidence of end-stage renal disease in

- the multiple risk factor intervention trial. *J Am Soc Nephrol* 2006; **17**: 1444–1452.
30. Imai E, Horio M, Yamagata K *et al.* Slower decline of glomerular filtration rate in the Japanese general population: a longitudinal 10-year follow-up study. *Hypertens Res* 2008; **31**: 433–441.
31. Ninomiya T, Perkovic V, de Galan BE, *et al.*, for the ADVANCE Collaborative Group. Albuminuria and kidney function independently predict cardiovascular and renal outcomes in diabetes. *J Am Soc Nephrol* 2009; **20**: 1813–1821.
32. Kshirsagar AV, Bang H, Bomback AS *et al.* A simple algorithm to predict incident kidney disease. *Arch Intern Med* 2008; **168**: 2466–2473.
33. Crowe E, Halpin D, Stevens P, on behalf of the Guideline Development G. Early identification and management of chronic kidney disease: summary of NICE guidance. *BMJ* 2008; **337**: 812–815.
34. O'Hare AM, Choi AI, Bertenthal D *et al.* Age affects outcomes in chronic kidney disease. *J Am Soc Nephrol* 2007; **18**: 2758–2765.
35. O'Hare AM, Bertenthal D, Covinsky KE *et al.* Mortality risk stratification in chronic kidney disease: one size for all ages? *J Am Soc Nephrol* 2006; **17**: 846–853.
36. Matsushita K, Selvin E, Bash LD *et al.* Change in estimated GFR associates with coronary heart disease and mortality. *J Am Soc Nephrol* 2009; **20**: 2617–2624.
37. White SL, Polkinghorne KR, Atkins RC *et al.* Comparison of the prevalence and mortality risk of CKD in Australia using the CKD Epidemiology collaboration (CKD-EPI) and modification of diet in renal disease (MDRD) Study GFR estimating equations: the AusDiab (Australian Diabetes, Obesity and Lifestyle) Study. *Am J Kidney Dis* 2010; **55**: 660–670.
38. Shankar A, Klein R, Klein BEK. The association among smoking, heavy drinking and chronic kidney disease. *Am J Epidemiol* 2006; **164**: 263–271.
39. Shlipak MG, Katz R, Kestenbaum B *et al.* Rate of kidney function decline in older adults: a comparison using creatinine and cystatin C. *Am J Nephrol* 2009; **30**: 171–178.
40. Hallan SJ, Ritz E, Lydersen S *et al.* Combining GFR and albuminuria to classify CKD improves prediction of ESRD. *J Am Soc Nephrol* 2009; **20**: 1069–1077.
41. Bui AL, Katz R, Kestenbaum B *et al.* Cystatin C and carotid intima-media thickness in asymptomatic adults: the Multi-Ethnic Study of Atherosclerosis (MESA). *Am J Kidney Dis* 2009; **53**: 389–398.
42. Iseki K, Ikemiya Y, Iseki C *et al.* Proteinuria and the risk of developing end-stage renal disease. *Kidney Int* 2003; **63**: 1468–1474.
43. Tonelli M, Jose P, Curhan G, *et al.*, Cholesterol and Recurrent Events (CARE) Trial Investigators. Proteinuria, impaired kidney function, and adverse outcomes in people with coronary disease: analysis of a previously conducted randomised trial. *BMJ* 2006; **332**: 1426–1431.
44. Mann JF, Schmieder RE, McQueen M, *et al.*, ONTARGET investigators. Renal outcomes with telmisartan, ramipril, or both, in people at high vascular risk (the ONTARGET study): a multicentre, randomised, double-blind, controlled trial. *Lancet* 2008; **372**: 547–553.
45. Pavkov ME, Knowler WC, Hanson RL *et al.* Predictive power of sequential measures of albuminuria for progression to ESRD or death in Pima Indians with type 2 diabetes. *Am J Kidney Dis* 2008; **51**: 759–766.
46. Mann JFE, Schmieder RE, Dyal L *et al.* Effects of telmisartan on renal outcomes. *Ann Int Med* 2009; **151**: 1–10.
47. Lee BJ, Forbes K. The role of specialists in managing the health of populations with chronic illness: the example of chronic kidney disease. *BMJ* 2009; **339**: 800–802.
48. Levey AS, Coresh J, Greene T *et al.* Using standardized serum creatinine values in the modification of diet in renal disease study equation for estimating glomerular filtration rate. *Ann Intern Med* 2006; **145**: 247–254.
49. Konta T, Hao Z, Takasaki S *et al.* Clinical utility of trace proteinuria for microalbuminuria screening in the general population. *Clin Exp Nephrol* 2007; **11**: 51–55.
50. Woodward M. *Epidemiology: Study Design and Data Analysis*. 2nd edn. Chapman & Hall/CRC: Boca Raton, 2005.

Significance of past history of renal failure for the detection of high-risk individuals for cardiovascular and end-stage renal disease: analysis of data from a nationwide health checkup

Kazunobu Ichikawa · Tsuneo Konta · Ami Ikeda · Shouichi Fujimoto · Kunitoshi Iseki · Toshiaki Moriyama · Kunihiro Yamagata · Kazuhiko Tsuruya · Hideaki Yoshida · Koichi Asahi · Issei Kurahashi · Yasuo Ohashi · Tsuyoshi Watanabe

Received: 10 May 2011 / Accepted: 18 July 2011 / Published online: 5 August 2011
© Japanese Society of Nephrology 2011

Abstract

Background Clinical information regarding risk factors may be helpful in the detection of various diseases. This cross-sectional study examined the characteristics of subjects with past history of renal failure, and assessed whether this information would be useful for the efficient detection of high-risk individuals for chronic kidney disease (CKD) and cardiovascular disease (CVD) at health checkup.

Methods This study utilized data from a nationwide health checkup, “The Specific Health Check and Guidance in Japan,” and data for 250,130 adult subjects were analyzed. Subjects with self-reported history of renal failure and receiving dialysis therapy were defined as having a history of renal failure.

Results Among total participants, there were 1,400 (0.6%) with a history of renal failure. The prevalence of a history of renal failure was higher in subjects with CKD than in those

without CKD (1.5 vs. 0.3%, $P < 0.001$) and increased with progression of the stage of CKD (0.9–43.5%). Subjects with a history of renal failure had a reduced estimated glomerular filtration rate (44.6 ± 20.3 ml/min/1.73 m²) and a higher prevalence of CKD (50.5%) and CVD (31.9%), compared with subjects with hypertension, diabetes or metabolic syndrome. Multivariate logistic regression analysis showed an independent association between a history of CVD and renal failure (odds ratio 3.68, 95% confidence interval 3.26–4.15), after adjustment for confounding factors.

Conclusions A history of renal failure was strongly associated with advanced CKD and CVD. Information regarding history of renal failure could be utilized to efficiently detect high-risk individuals at health checkup.

Keywords Chronic kidney disease · Epidemiology · Past history

K. Ichikawa · T. Konta (✉) · A. Ikeda
Department of Cardiology, Pulmonology,
and Nephrology, Yamagata University School of Medicine,
2-2-2 Iida-Nishi, Yamagata 990-9585, Japan
e-mail: kkonta@med.id.yamagata-u.ac.jp

S. Fujimoto
Dialysis Division, University of Miyazaki Hospital,
Miyazaki, Japan

K. Iseki · T. Moriyama · K. Yamagata · K. Tsuruya ·
H. Yoshida · K. Asahi · T. Watanabe
Steering Committee for the “Research on the Positioning of
Chronic Kidney Disease (CKD) in Specific Health Check and
Guidance in Japan”, Otawara, Japan

I. Kurahashi · Y. Ohashi
Department of Biostatistics/Epidemiology and Preventive Health
Sciences, School of Health Sciences and Nursing,
University of Tokyo, Tokyo, Japan

Introduction

Chronic kidney disease (CKD) is a risk factor for cardiovascular disease (CVD) as well as end-stage renal disease; therefore, early detection and treatment of CKD is important [1]. Local health checks have been performed annually for the Japanese general population to facilitate the early detection of high-risk subjects for CKD and CVD. The main target of the recent health checkup “Specific Health Check and Guidance in Japan” was metabolic syndrome, which is associated with cardiovascular and kidney disease.

In mass screening programs, it is necessary from a cost-benefit perspective to efficiently detect high-risk subjects. Expensive investigations such as direct measurement of albuminuria may be acceptable in a high-risk population such as diabetic patients [2], but not in a low-risk population

such as the general population [3]. Therefore, for low-risk populations, it is desirable to reduce the number of subjects screened, by consideration of the characteristics of participants, including age and comorbidities [3]. For this purpose, previous studies have utilized information such as a family history of CKD [4, 5], and past history of hypertension and diabetes [6] for the detection of patients with CKD.

A question on past history of renal failure was included in the “Specific Health Check and Guidance in Japan” survey. This information on the history of renal failure may be useful for detecting subjects at high-risk of end-stage renal disease and CVD. Therefore, to clarify the characteristics of subjects with a history of renal failure, and to assess whether this information could be used for the efficient detection of high-risk individuals at health checkup, a cross-sectional study was conducted using data from the nationwide annual health check of the Japanese general population.

Methods

Study population

This study was part of the ongoing “Research on the Positioning of CKD in Specific Health Check and Guidance in Japan” program. The Specific Health Check and Guidance program is an annual health check in which all inhabitants over the age of 20 years who are covered by national insurance in Japan are invited to participate. Data from the nationwide database was obtained for 13 prefectures (Yamagata, Miyagi, Fukushima, Niigata, Tokyo, Kanagawa, Ibaraki, Osaka, Okayama, Kochi, Fukuoka, Miyazaki and Okinawa) that agreed with the study aims. In 2008 and 2009, data were collected on 278,017 men and 383,586 women (a total of 676,905 individuals, aged 20–101 years), who took part in the health checks. The study was approved by the institutional ethics committee.

Of the 676,905 participants, 426,775 were excluded from the present analysis because essential data, including information on proteinuria and serum creatinine levels, was incomplete. Therefore, data for 101,147 males and 148,983 females (a total of 250,130 individuals, aged 20–88 years) were used in the final statistical analyses. Among the 250,130 participants, there were 112,002 hypertensive subjects (44.8%), 27,820 subjects with metabolic syndrome (11.2%), 23,403 subjects with diabetes (9.4%), 138,535 subjects with dyslipidemia (55.4%) and 45,845 subjects with CKD (18.3%).

Measurements

Subjects used a self-report questionnaire to document their medical history, current medications, smoking habit

(smoker or non-smoker) and alcohol intake (drinker or non-drinker). To assess their past history of renal failure, all participants were asked “Have you ever been told that you have renal failure or have you received dialysis therapy before?” If the answer was “yes,” the subject was considered to have a past history of renal failure. The past history of renal failure mainly consists of past history of AKI and CKD stage 4 or 5. However, the patients on maintenance dialysis (stage 5D) were not included. Systolic blood pressure (SBP) and diastolic blood pressure (DBP) were determined using a standard sphygmomanometer or an automated device, with subjects in the sitting position after a sufficient period of rest. Hypertension was defined as a SBP ≥ 140 mmHg or a DBP ≥ 90 mmHg, or use of anti-hypertensive medication. Body mass index (BMI) was calculated as weight (kg) divided by height squared (m^2). For both men and women, obesity was defined as BMI ≥ 25.0 kg/ m^2 . Blood samples were obtained after overnight fasting. Plasma glucose levels were measured using the hexokinase enzymatic reference method. Subjects with diabetes were identified either by self-reported physical diagnosis or by a fasting plasma glucose concentration ≥ 126 mg/dl or a HbA1c value $\geq 6.5\%$. Triglyceride and low-density lipoprotein cholesterol (LDL-C) concentrations were measured by enzymatic methods. High-density lipoprotein cholesterol (HDL-C) concentrations were measured directly. Dyslipidemia was defined as triglyceride concentrations ≥ 150 mg/dl or LDL-C concentrations ≥ 140 mg/dl, or HDL-C concentrations < 40 mg/dl or use of anti-lipidemic medication. Metabolic syndrome was defined according to the Japanese criteria for high waist circumference (≥ 85 cm in men and ≥ 90 cm in women) plus any two of the following: (1) blood pressure $\geq 130/85$ mmHg and/or use of anti-hypertensive medication; (2) fasting plasma glucose ≥ 110 mg/dl and/or use of anti-diabetic medication; and (3) triglyceride concentrations ≥ 150 mg/dl and/or HDL-C concentrations < 40 mg/dl [7].

Urinalysis by the dipstick method was performed on a single spot urine specimen collected in the early morning after overnight fasting. The results of the urinalysis were recorded as (–), trace, (1+), (2+) or (3+). Positivity for proteinuria was defined as (1+) or greater. Serum creatinine was measured using an enzymatic method, and the estimated glomerular filtration rate (eGFR) was calculated using the Japanese equation for eGFR [8]. We assumed the subjects with the presence of proteinuria and/or renal insufficiency (eGFR < 60 ml/min/ 1.73 m^2) as having CKD. CKD was further categorized into five stages: stage 1, eGFR ≥ 90 ml/min/ 1.73 m^2 with proteinuria; stage 2, eGFR 60–89 with proteinuria; stage 3, eGFR 30–59; stage 4, eGFR 15–29; stage 5, eGFR < 15 [9]. To investigate in detail the relationship between stage of CKD and history of

renal failure, stage 3 was further divided into stages 3A (eGFR 45–59) and 3B (eGFR 30–44).

Statistical analyses

Unpaired *t* tests were used to compare mean values, and chi-square tests were used to evaluate differences in proportions. The trend in proportions was examined by chi-square test. To examine the factors associated with CVD (stroke and heart disease), unadjusted and adjusted multivariate logistic regression analyses were performed, including age, gender, alcohol consumption, smoking, hypertension, diabetes, metabolic syndrome, proteinuria, renal insufficiency and past history of renal failure as independent variables. Continuous data are expressed as mean ± SD. All statistical analyses were performed using JMP version 8 software (SAS Institute Inc., Cary, NC). Differences were regarded as significant when *P* values were <0.05.

Results

Characteristics of subjects in the general population with a past history of renal failure

Among the 250,130 participants, there were 650 men and 750 women with a past history of renal failure (a total of

1,400 subjects, 0.6% of all participants, 1.5% of the subjects with CKD). Subjects with a history of renal failure were more likely to be older, to be male, obese, and have CKD, hypertension, diabetes, metabolic syndrome, hematuria (≥1+), proteinuria (≥1+), a past history of CVD (stroke and heart disease), to have received treatment for hypertension, diabetes or dyslipidemia, and less likely to be smokers compared with the 248,730 subjects without a history of renal failure (Table 1). The subjects with a history of renal failure were mostly aged over 40 years (mean age 64.5 years); as percentages of all participants, 0.1% were <40 years, 0.5% were 40–49 years, 0.5% were 50–59 years, 0.5% were 60–69 years and 0.7% were ≥70 years.

The association between CKD and a past history of renal failure

The prevalence of a history of renal failure was higher in the CKD population than in the population without CKD (1.5 vs. 0.3%, *P* < 0.001) and increased with progression of the stage of CKD: 0.9% in stage 1, 1.0% in stage 2, 1.3% in stage 3 (0.9% in stage 3A and 5.4% in stage 3B), and 23.2% in stage 4 and 43.5% in stage 5 (*P* for trend <0.001) (Fig. 1).

The distribution of eGFR in subjects with a history of renal failure and subjects with other common disorders was

Table 1 The characteristics of participants with and without past history of renal failure

	Past history of renal failure		<i>P</i> value
	(+)	(–)	
Number (%)	1,400 (0.6%)	248,730 (99.4%)	
Age (years)	64.5 ± 8.2	63.6 ± 8.7	<0.001
Male (%)	46.4	40.4	<0.001
Serum creatinine (mg/dl)	1.16 ± 1.27	0.72 ± 0.22	<0.001
eGFR (ml/min/1.73 m ²)	60.6 ± 23.6	75.2 ± 16.1	<0.001
CKD (%)	50.6	18.2	<0.001
Hypertension (%)	60.8	44.7	<0.001
Diabetes (%)	15.4	9.3	<0.001
Dyslipidemia (%)	57.4	55.4	0.123
Drinker (%)	37.6	45.4	<0.001
Smoker (%)	13.1	13.7	0.515
Obesity (%)	29.5	25.5	<0.001
Metabolic syndrome (%)	13.7	11.2	0.002
Hematuria ≥1+ (%)	27.9	17.1	<0.001
Proteinuria ≥1+ (%)	25.6	5.5	<0.001
Past history of CVD (%)	31.9	9.0	<0.001
Past history of stroke (%)	19.6	3.5	<0.001
Past history of heart disease (%)	24.9	6.1	<0.001
Treatment in hypertension (%)	49.3	29.5	<0.001
Treatment in diabetes (%)	11.5	5.2	<0.001
Treatment in dyslipidaemia (%)	23.4	17.2	<0.001

Mean ± SD
eGFR estimated glomerular filtration rate, *CKD* chronic kidney disease, *CVD* cardiovascular disease (stroke and heart disease)




Fig. 1 The prevalence of a history of renal failure according to stage of CKD. CKD chronic kidney disease




Fig. 3 The proportion of CKD stages. MetS metabolic syndrome




Fig. 2 The distribution of eGFR in subjects with a history of renal failure and subjects with other common disorders. eGFR estimated glomerular filtration rate, MetS metabolic syndrome

examined. The eGFR distribution was shifted to the low range in subjects with a history of renal failure, as compared to those with hypertension, metabolic syndrome or diabetes (Fig. 2). The prevalence of proteinuria ($\geq 1+$) was higher in subjects with a history of renal failure (25.6%), as compared with the total population (5.5%), those with metabolic syndrome (13.0%), those with hypertension (8.5%) and those with diabetes (14.4%). Furthermore, high grade proteinuria ($\geq 2+$) was frequently observed in subjects with a history of renal failure.

The prevalence of CKD in subjects with various disorders was investigated. Subjects with a history of renal failure had a higher prevalence of CKD (50.5%) as compared with the total population (18.3%), those with metabolic syndrome (22.6%), those with hypertension (23.8%) and those with diabetes (26.9%) (Fig. 3). Although the prevalence of early stages of CKD (stages 1–3A) was

similar between subjects with different disorders (20.0–26.7%), the prevalence of advanced CKD (stages 3B–5) was much higher in subjects with a history of renal failure (23.8%), compared to those with other disorders (2.6–3.1%).

In addition, the significance of a history of renal failure among the population with CKD was investigated. Among the total of 45,845 subjects with CKD, there were 406 men and 302 women with a history of renal failure (a total of 708 subjects). As compared with the 45,137 subjects without a history of renal failure, subjects with a history of renal failure were more likely to be male, to have a lower eGFR (44.6 ± 20.3 vs. 58.4 ± 14.1 ml/min/1.73 m², $P < 0.001$) and a higher prevalence of hypertension, diabetes, hematuria ($\geq 1+$), proteinuria ($\geq 1+$), a history of CVD (stroke and heart disease), to have received treatment for hypertension, diabetes or dyslipidemia, and were less likely to consume alcohol (Table 2). The proportion of subjects with optimal blood pressure control for CKD ($<130/80$ mmHg) did not differ significantly between those with or without a history of renal failure (34.1 vs. 34.6%, $P = 0.833$).

The association between a past history of cardiovascular disease and renal failure

Finally, the relationship between a past history of CVD (stroke and heart disease) and renal failure was examined. The subjects with a history of renal failure had a higher prevalence of CVD (31.9%) as compared with the total population (9.1%), those with metabolic syndrome (12.1%), those with hypertension (13.7%) and those with diabetes (14.8%) (Fig. 4). Logistic regression analyses showed that a past history of CVD was independently associated with proteinuria, renal insufficiency (eGFR

Table 2 The characteristics of participants with and without past history of renal failure among CKD population

	Past history of renal failure		P value
	(+)	(-)	
Number (%)	708 (1.5%)	45,137 (98.5%)	
Age (years)	66.3 ± 6.7	66.3 ± 6.9	0.83
Male (%)	57.3	51.8	0.004
Systolic BP (mmHg)	132.4 ± 18.5	131.9 ± 17.7	0.454
Diastolic BP (mmHg)	76.8 ± 11.0	77.6 ± 10.8	0.079
Serum creatinine (mg/dl)	1.63 ± 1.65	0.93 ± 0.38	<0.001
eGFR (ml/min/1.73 m ²)	44.6 ± 20.3	58.4 ± 14.1	<0.001
Hypertension (%)	75	57.8	<0.001
Diabetes (%)	19.2	13.6	<0.001
Dyslipidemia (%)	59.2	61.7	0.168
Drinker (%)	34.6	45.7	<0.001
Smoker (%)	10.7	12.8	0.111
Obesity (%)	31.8	32.7	0.595
Metabolic syndrome (%)	17.1	17.4	0.841
Hematuria ≥1+ (%)	35.9	20.5	<0.001
Proteinuria ≥1+ (%)	50.7	30.2	<0.001
BP <130/80 (mmHg)	34.1	34.6	0.833
Past history of CVD (%)	29.2	13.6	<0.001
Past history of stroke (%)	12.9	5.7	<0.001
Past history of heart disease (%)	21.5	9.2	<0.001
Treatment in hypertension (%)	66	42	<0.001
Treatment in diabetes (%)	15.1	8.2	<0.001
Treatment in dyslipidemia (%)	26.1	21.4	0.003

Mean ± SD

CKD chronic kidney disease, BP blood pressure, eGFR estimated glomerular filtration rate, CVD cardiovascular disease




Fig. 4 The prevalence of a history of cardiovascular disease. CVD cardiovascular disease (stroke and heart disease), MetS metabolic syndrome

<60 ml/min/1.73 m²) and a history of renal failure, both in unadjusted and adjusted models. In the adjusted model that included possible confounders, the odds ratio (OR) for CVD and a history of renal failure [OR 3.68, 95% confidence interval (CI) 3.26–4.15] was greater than those for CVD and proteinuria (OR 1.16, 95% CI 1.10–1.22) or

CVD and eGFR <60 ml/min/1.73 m² (OR 1.32, 95% CI 1.27–1.36) (Table 3).

Discussion

This study showed, for the first time, that the proportion of subjects who had a past history of renal failure was low at health checkup; however, they had a high prevalence of advanced stages of CKD, a history of cardiovascular events and risk factors for atherosclerosis.

Several studies reported that the prevalence of a previous diagnosis of renal failure in the general population, irrespective of severity and time course (acute or chronic), was 2.0% in the United States [10] and 3.3% in Italy [11]. Other studies documented a higher prevalence of family history of kidney disease (17–24%) [4] or end-stage renal disease (9.5%) [5]. The present study, which focused on severe kidney disease (renal failure or requirement for dialysis therapy), showed that the prevalence (0.6%) was much less than those reported in previous studies and was closer to the prevalence of acute kidney disease, as identified in the International Classification of Diseases (ICD) [12]. Although the prevalence of a history of kidney disease varies, depending on the criteria used, our finding

Table 3 Factors related with the past history of cardiovascular diseases

	Unadjusted		Adjusted ^a	
	OR (95% CI)	P value	OR (95% CI)	P value
Proteinuria ($\geq 1+$)	1.75 (1.67–1.84)	<0.001	1.16 (1.10–1.22)	<0.001
eGFR <60 (ml/min/1.73 m ²)	1.87 (1.81–1.93)	<0.001	1.32 (1.27–1.36)	<0.001
Past history of renal failure	4.74 (4.23–5.31)	<0.001	3.68 (3.26–4.15)	<0.001

OR odds ratio, eGFR estimated glomerular filtration rate

^a Adjusted with age, gender, drinking, smoking, hypertension, diabetes and metabolic syndrome

showed that the number of subjects with a past history of renal failure, who might require further examination for advanced CKD and CVD, is small and could be easily identified at health checkup.

Subjects with a history of renal failure were not only likely to have CKD, but also to have CVD and multiple risk factors for atherosclerosis, including male gender, hypertension and metabolic syndrome. Therefore, these subjects should be regarded as a high-risk population both for end-stage renal disease and CVD. However, a history of renal failure was not emphasized in health checkups in Japan, and blood pressure was optimally controlled (<130/80 mmHg) [13] in only one-third of these subjects. An Italian study reported that the evaluation and treatment of kidney disease by general physicians was likely to be suboptimal [14, 15]. These findings suggest that the management of a majority of subjects with CKD may be inadequate, and that aggressive evaluation and treatment of kidney disease should be encouraged.

Of note, a history of renal failure was independently associated with cardiovascular events, with a greater odds ratio than those for other renal indices such as proteinuria and renal insufficiency. Due to the cross-sectional nature of this study, it is not possible to infer a causal relationship; however, there is a possibility that renal failure might directly affect the development of cardiovascular events, or vice versa. The causal relationship between these factors should be clarified by a longitudinal investigation.

This study had several strengths. First, we focused on information regarding a past history of renal failure that was readily obtained at health checkup and shows a strong association with advanced CKD and CVD. Currently, the measurement of serum creatinine is not mandatory at health checkup, mainly due to the low cost-effectiveness. Consequently, serum creatinine is not determined in more than half of participants. This causes advanced CKD subjects without urine abnormalities to be overlooked. Although the past history of renal failure cannot cover the whole population of CKD and CVD, it could be a help to find occult subjects with advanced stages of CKD without extra effort and cost. Second, we utilized a large nationwide database that reflects the

current health status of the entire Japanese population. This enabled us to analyze the characteristics of the target population in detail and to obtain useful clinical information for the detection of individuals at high-risk of CKD and CVD. In previous reports the incidence rate of end-stage renal disease among individuals with AKI was 2.6–2.8 per 100 person-years [16, 17]. In a referred CKD cohort 44.9% of CKD patients had at least one AKI episode [18]. Consistent with these reports that acute kidney injury is the risk factor of end-stage renal disease, half of the subjects with a history of renal failure had CKD in this study. However, it is of interest that the other half of them showed no sign of CKD. This indicates that the population with a history of renal failure is heterogeneous and should be treated individually.

This study also had some limitations that should be noted. First, the single measurements of clinical parameters may have led to some misclassification of comorbidities, including CKD. Such misclassification would probably have been non-differential and would have biased the relationship toward null. Therefore, there is a possibility that the strength of the observed relationship between a history of renal failure and various parameters may have been underestimated. Second, there was no detailed information available on the cause and clinical course of kidney disease. The observed relationship may be affected by the etiology and treatment of kidney disease. Third, there may have been a selection bias in that the study population was comprised of relatively healthy local inhabitants. Therefore, caution is required in generalizing these findings to other populations.

In conclusion this study showed that a past history of renal failure was strongly associated with current CKD and CVD. Information on past history of renal failure could be utilized for the efficient detection of high-risk individuals for end-stage renal disease and CVD at health checkup.

Acknowledgments This work was supported by Health and Labour Sciences Research Grants for “Research on the Positioning of Chronic Kidney Disease (CKD) in Specific Health Check and Guidance in Japan” (20230601), the Ministry of Health, Labour and Welfare of Japan.

Conflict of interest None of the authors has a conflict of interest to declare.

References

- Go AS, Chertow GM, Fan D, McCulloch CE, Hsu CY. Chronic kidney disease and the risks of death, cardiovascular events, and hospitalization. *N Engl J Med*. 2004;351:1296–305.
- Kiberd BA, Jindal KK. Screening to prevent renal failure in insulin dependent diabetic patients: an economic evaluation. *BMJ*. 1995;311:1595–9.
- Boulware LE, Jaar BG, Tarver-Carr ME, Brancati FL, Powe NR. Screening for proteinuria in US adults: a cost-effectiveness analysis. *JAMA*. 2003;290:3101–14.
- Jolly SE, Burrows NR, Chen SC, Li S, Jurkovitz CT, Narva AS, et al. Racial and ethnic differences in albuminuria in individuals with estimated GFR greater than 60 mL/min/1.73 m²: results from the Kidney Early Evaluation Program (KEEP). *Am J Kidney Dis*. 2010;55:S15–22.
- McClellan W, Speckman R, McClure L, Howard V, Campbell RC, Cushman M, et al. Prevalence and characteristics of a family history of end-stage renal disease among adults in the United States population: Reasons for Geographic and Racial Differences in Stroke (REGARDS) renal cohort study. *J Am Soc Nephrol*. 2007;18:1344–52.
- Brown WW, Peters RM, Ohmit SE, Keane WF, Collins A, Chen SC, et al. Early detection of kidney disease in community settings: the Kidney Early Evaluation Program (KEEP). *Am J Kidney Dis*. 2003;42:22–35.
- Committee to Evaluate Diagnostic Standards for Metabolic Syndrome. Definition and the diagnostic standard for metabolic syndrome (in Japanese). *Nippon Naika Gakkai Zasshi*. 2005;94:794–809.
- Matsuo S, Imai E, Horio M, Yasuda Y, Tomita K, Nitta K, et al. Revised equations for estimated GFR from serum creatinine in Japan. *Am J Kidney Dis*. 2009;53:982–92.
- Levey AS, Eckardt KU, Tsukamoto Y, Levin A, Coresh J, Rossert J, et al. Definition and classification of chronic kidney disease: a position statement from Kidney Disease: Improving Global Outcomes (KDIGO). *Kidney Int*. 2005;67:2089–100.
- Coresh J, Byrd-Holt D, Astor BC, Briggs JP, Eggers PW, Lacher DA, et al. Chronic kidney disease awareness, prevalence, and trends among U.S. adults, 1999 to 2000. *J Am Soc Nephrol*. 2005;16:180–8.
- Cirillo M, Laurenzi M, Mancini M, Zanchetti A, Lombardi C, De Santo NG. Low glomerular filtration in the population: prevalence, associated disorders, and awareness. *Kidney Int*. 2006;70:800–6.
- Grams ME, Astor BC, Bash LD, Matsushita K, Wang Y, Coresh J. Albuminuria and estimated glomerular filtration rate independently associate with acute kidney injury. *J Am Soc Nephrol*. 2010;21:1757–64.
- Japanese Society of Nephrology. Evidence-based practice guideline for the treatment of CKD. *Clin Exp Nephrol*. 2009;13:537–66.
- Minutolo R, De Nicola L, Mazzaglia G, Postorino M, Cricelli C, Mantovani LG, et al. Detection and awareness of moderate to advanced CKD by primary care practitioners: a cross-sectional study from Italy. *Am J Kidney Dis*. 2008;52:444–53.
- Muntner P, Anderson A, Charleston J, Chen Z, Ford V, Makos G, et al. Hypertension awareness, treatment, and control in adults with CKD: results from the Chronic Renal Insufficiency Cohort (CRIC) Study. *Am J Kidney Dis*. 2010;55:441–51.
- Ishani A, Xue JL, Himmelfarb J, Eggers PW, Kimmel PL, Molitoris BA, et al. Acute kidney injury increases risk of ESRD among elderly. *J Am Soc Nephrol*. 2009;20:223–8.
- Wald R, Quinn RR, Luo J, Li P, Scales DC, Mamdani MM, et al. Chronic dialysis and death among survivors of acute kidney injury requiring dialysis. *JAMA*. 2009;302:1179–85.
- Lafrance JP, Djurdjev O, Levin A. Incidence and outcomes of acute kidney injury in a referred chronic kidney disease cohort. *Nephrol Dial Transplant*. 2010;25:2203–9.

see commentary on page 229

Association between prehypertension and chronic kidney disease in the Japanese general population

Yuichiro Yano¹, Shouichi Fujimoto², Yuji Sato¹, Tsuneo Konta³, Kunitoshi Iseki⁴, Toshiki Moriyama⁴, Kunihiro Yamagata⁴, Kazuhiko Tsuruya⁴, Hideaki Yoshida⁴, Koichi Asahi⁴, Issei Kurahashi⁵, Yasuo Ohashi⁵ and Tsuyoshi Watanabe⁴

¹Circulatory and Body Fluid Regulation, Department of Internal Medicine, Faculty of Medicine, University of Miyazaki, Miyazaki, Japan; ²Dialysis Division, University of Miyazaki Hospital, Miyazaki, Japan; ³Department of Cardiology, Pulmonology, and Nephrology, Yamagata University School of Medicine, Yamagata, Japan; ⁴Steering Committee for the 'Research on the positioning of chronic kidney disease (CKD) in Specific Health Check and Guidance in Japan', Japan and ⁵Department of Biostatistics/Epidemiology and Preventive Health Sciences, School of Health Sciences and Nursing, University of Tokyo, Tokyo, Japan

The increased prevalence of chronic kidney disease (CKD) is a consequence of the accumulation of risk factors, one of which is hypertension. Here we assessed the prevalence of CKD according to blood pressure among 232,025 patients in a Japanese nationwide database with a focus on the prevalence and risk factors of CKD in prehypertension. Patients were stratified by blood pressure and included 75,474 with optimal blood pressure (less than 120/80 mm Hg); 59,194 with prehypertension and a normal blood pressure (120–129/80–84 mm Hg) or 46,547 patients with high-normal blood pressure (130–139/85–89 mm Hg); and 50,810 with hypertension (over 140/90 mm Hg without anti-hypertensive drugs). CKD was defined as an estimated glomerular filtration rate of stage 3 or lower or having proteinuria greater than 1+ by a dipstick method. The prevalence of CKD among patients with optimal blood pressure, prehypertension having normal or high-normal blood pressure, and hypertension was 13.9, 15.6, 18.1, and 20.7% in men, and 10.9, 11.6, 12.9, and 15.0% in women, with a significant difference between genders at each strata of blood pressure. In men, but not in women, whose blood pressure was high-normal, the CKD risk was significantly greater (odds ratio 1.11) than those with optimal blood pressure. Obesity (body mass index over 25) was significantly associated with an increased risk of CKD in both men and women (odds ratio 1.43 and 1.26, respectively), and there was an additive effect of obesity and pre-hypertension on CKD risk in men compared with men with optimal blood pressure. Thus, the prevalence of CKD increased with the severity of blood pressure. Prehypertension with high-normal

blood pressure, particularly in conjunction with obesity, was found to be an independent risk factor of CKD in men.

Kidney International (2012) **81**, 293–299; doi:10.1038/ki.2011.346; published online 28 September 2011

KEYWORDS: chronic kidney disease; high-normal blood pressure; obesity; prehypertension

Chronic kidney disease (CKD) is now recognized as a major global public health problem.^{1,2} It is increasingly apparent that CKD is associated with increased risk of not only progression to renal failure but also excess cardiovascular morbidity and mortality in a manner independent of other known risk factors.^{1,2}

CKD affects 10–15% of the adult population worldwide.^{3,4} A recent Japanese survey demonstrated that the prevalence of CKD increased significantly in men, but not in women, from the 1970s to the 2000s in the general population.⁵ The reasons are not well understood, but it is likely that the increased prevalence of CKD is a consequence of the accumulation of risk factors, such as hypertension or metabolic abnormalities including diabetes, dyslipidemia, and obesity, over the last three decades.⁵ Furthermore, Japan is known to have a high incidence of end-stage renal disease, and the number of patients undergoing dialysis has been increasing.^{6,7} The incidence and prevalence of end-stage renal disease are higher in men than in women in Japan.^{8,9} Individuals with CKD have reduced life expectancy, and the social burden of CKD with or without end-stage renal disease is becoming greater. Accordingly, it should be a public health priority to identify CKD-prone high-risk subjects in the general population and to treat risk factors in the initial phase of CKD in order to prevent and delay the progression to renal failure. Such efforts would also help to prevent cardiovascular diseases.

Correspondence: Shouichi Fujimoto, Dialysis Division, University of Miyazaki Hospital, Kihara 5200, Kiyotake, Miyazaki 889-1692, Japan.
E-mail: fujimos@fc.miyazaki-u.ac.jp

Received 10 April 2011; revised 2 August 2011; accepted 9 August 2011; published online 28 September 2011

Hypertension is well established as both a cause and consequence of CKD.^{10–12} In Asian countries in particular, high blood pressure (BP) is the strongest risk factor for renal outcome.¹⁰ A previous study in Japan demonstrated that there was a linear continuous association between BP and incidence of end-stage renal disease; even in subjects without hypertension (i.e., even in subjects with prehypertension: systolic BP/diastolic BP, 120–139/80–89 mm Hg), there was a greater risk of future development of end-stage renal disease compared with the risk in subjects with optimal BP (<120/80 mm Hg).¹¹ Given the evidence that the risk of end-stage renal disease is increased throughout the BP range, understanding the burden of CKD in subjects with prehypertension could help in promoting prevention and screening efforts for both CKD and prehypertension.¹³ Recently, the National Health and Nutrition Examination Survey in the United States demonstrated that the prevalence of CKD among those with prehypertension was 17.3%, compared with 13.4% in those with optimal BP.¹⁴ However, there has been no comparable analysis of a nationwide database in Japan.

Accordingly, in the present study, we examined the prevalence of CKD within BP classification using a large nationwide database of subjects recruited from the national health checkup system in Japan. In addition, we examined some clinical characteristics other than BP that are prone to increase risk of CKD.

RESULTS

Patient characteristics

By reviewing the data from the national health checkup program in Japan, we identified 346,942 subjects for whom all the clinical data required for the present analysis were available. A total of 84,854 subjects with a history

of treatment with anti-hypertensive medications, 12,771 subjects with a previous history of cardiovascular diseases, and 17,049 subjects with both were excluded from the present analysis. Moreover, 243 subjects with CKD stage 5 (estimated glomerular filtration rate (eGFR) <15 ml/min per 1.73 m²) were excluded. Table 1 shows the clinical characteristics of all subjects included in the present study (*n* = 232,025, left column) or the clinical characteristics according to gender difference (right column).

BP classification

Among the study subjects, 75,474 subjects (32.5%) had optimal BP, 105,741 subjects (45.6%) had prehypertension (normal BP: 59,194 subjects, 25.5%; high-normal BP: 46,547 subjects, 20.1%), and 50,810 subjects (21.9%) had hypertension. As the prevalence of such BP classification differed between men and women, the clinical characteristics according to BP classification were described by gender (Table 2). In accordance with the severity of BP classification, significant increases of age and body mass index, and significant decrease in the prevalence of current smoking, were observed. Information about glucose and lipid parameters could be obtained in some subjects, although not all: according to the severity of BP classification, there were significant differences in the glucose and lipid parameters (Supplementary Table S1 online).

CKD and BP classification

A total of 32,692 subjects (14.1%) were diagnosed with CKD, and 8751 subjects (3.8%) had proteinuria ($\geq 1+$). There was a gender difference in the prevalence of CKD (17.0% in men versus 12.2% in women; *P* < 0.001); accordingly, we determined the relationship between prevalence of CKD and BP classification separately for each gender (Table 2).

Table 1 | Characteristics of the study population overall (left column) or by gender (right column)

	Total subjects (<i>n</i> =232,025)	Gender difference		<i>P</i> -value
		Women (<i>n</i> =142,293)	Men (<i>n</i> =89,732)	
Age, years	61.8 ± 9.4	62.0 ± 9.1	61.4 ± 9.9	<0.001
Men, <i>n</i> (%)	89,732 (38.7)	—	89,732 (100)	<0.001
Body mass index, kg/m ²	22.6 ± 3.2	22.2 ± 3.2	23.4 ± 3.0	<0.001
Obesity, <i>n</i> (%)	58,061 (25.0)	29,358 (20.6)	28,703 (32.0)	<0.001
Current smoker, <i>n</i> (%)	36,058 (15.5)	9912 (7.0)	26,146 (29.1)	<0.001
Daily drinker, <i>n</i> (%)	50,495 (21.8)	12,471 (8.8)	38,024 (42.4)	<0.001
eGFR, ml/min per 1.73m ²	76.9 ± 16.0	76.9 ± 15.9	76.8 ± 16.3	0.57
CKD, <i>n</i> (%)	32,692 (14.1)	17,409 (12.2)	15,283 (17.0)	<0.001
Stage 1 and 2, <i>n</i> (%)	7041 (3.0)	3232 (2.3)	3809 (4.2)	<0.001
Stage 3, <i>n</i> (%)	25,547 (11.0)	14,117 (9.9)	11,430 (12.7)	
Stage 4, <i>n</i> (%)	104 (0.04)	60 (0.04)	44 (0.05)	
Proteinuria ($\geq 1+$), <i>n</i> (%)	8751 (3.8)	3948 (2.8)	4803 (5.4)	<0.001
BP measurement				
Systolic BP, mm Hg	126 ± 17	124 ± 17	128 ± 17	<0.001
Diastolic BP, mm Hg	75 ± 11	73 ± 10	77 ± 11	<0.001

Abbreviations: BP, blood pressure; CKD, chronic kidney disease; eGFR, estimated glomerular filtration rate.

Data are expressed as the means ± SD or percentage. *P*-values were obtained by an unpaired *t*-test or χ^2 -test between women and men. Statistical significance was defined as *P* < 0.05. Obesity was defined as body mass index (BMI) ≥ 25 kg/m², and CKD was defined as eGFR <60 ml/min per 1.73 m² and/or presence of proteinuria ($\geq 1+$). The proteinuria number in each column includes all stage 1/2 patients plus a few in stage 3/4.

Table 2 | Patient characteristics and BP values according to the BP classification by gender

	Women (n=142,293)				P-value	Men (n=89,732)				P-value
	Optimal BP (n=51,715)	Prehypertension with normal BP (n=36,182)	Prehypertension with high-normal BP (n=27,348)	Hypertension (n=27,048)		Optimal BP (n=23,759)	Prehypertension with normal BP (n=23,012)	Prehypertension with high-normal BP (n=19,199)	Hypertension (n=23,762)	
Age, years	58.8 ± 10.2	62.7 ± 8.4	64.4 ± 7.5	64.8 ± 7.2	<0.001	59.0 ± 10.7	61.0 ± 10.1	62.9 ± 9.3	63.0 ± 8.8	<0.001
Body mass index, kg/m ²	21.4 ± 2.9	22.2 ± 3.1	22.7 ± 3.2	23.2 ± 3.5	<0.001	22.5 ± 2.8	23.3 ± 2.9	23.6 ± 3.0	24.0 ± 3.1	<0.001
Obesity, n (%)	6775 (13.1)	7349 (20.3)	6863 (25.1)	8371 (30.9)	<0.001	5256 (22.1)	7168 (31.1)	6689 (34.8)	9590 (40.4)	<0.001
Current smoker, n (%)	4852 (9.4)	2234 (6.2)	1488 (5.4)	1338 (4.9)	<0.001	7953 (33.5)	6562 (28.5)	5071 (26.4)	6560 (27.6)	<0.001
Daily drinker, n (%)	4594 (8.9)	3120 (8.6)	2350 (8.6)	2407 (8.9)	0.33	8059 (33.9)	9428 (41.0)	8713 (45.4)	11,824 (49.8)	<0.001
eGFR, ml/min per 1.73m ²	77.8 ± 15.9	76.9 ± 15.9	76.1 ± 15.7	75.8 ± 15.8	<0.001	78.1 ± 16.5	77.0 ± 16.1	76.1 ± 16.0	76.0 ± 16.4	<0.001
CKD, n (%)	5619 (10.9)	4204 (11.6)	3540 (12.9)	4046 (15.0)	<0.001	3303 (13.9)	3582 (15.6)	3475 (18.1)	4923 (20.7)	<0.001
Stage 1 and 2, n (%)	864 (1.7)	672 (1.9)	650 (2.4)	1046 (3.9)	<0.001	729 (3.1)	799 (3.5)	814 (4.2)	1467 (6.2)	<0.001
Stage 3, n (%)	4774 (9.2)	3516 (9.7)	2874 (10.5)	2983 (11.0)	<0.001	2565 (10.8)	2775 (12.1)	2652 (13.8)	3438 (14.5)	<0.001
Stage 4, n (%)	11 (0.02)	16 (0.04)	16 (0.05)	17 (0.06)	<0.001	9 (0.03)	8 (0.03)	9 (0.04)	18 (0.07)	<0.001
Proteinuria (≥1+), n (%)	1040 (2.0)	812 (2.2)	796 (2.9)	1300 (4.8)	<0.001	872 (3.7)	1003 (4.4)	1013 (5.3)	1915 (8.1)	<0.001
BP measurement										
Systolic BP, mm Hg	107 ± 8	123 ± 4	133 ± 4	149 ± 12	<0.001	109 ± 7	123 ± 4	132 ± 4	148 ± 13	<0.001
Diastolic BP, mm Hg	65 ± 7	73 ± 7	77 ± 7	85 ± 10	<0.001	67 ± 7	75 ± 6	79 ± 7	88 ± 10	<0.001

Abbreviations: BP, blood pressure; CKD, chronic kidney disease; eGFR, estimated glomerular filtration rate. Data are expressed as the means ± SD or percentage. Obesity was defined as body mass index (BMI) ≥ 25 kg/m², and CKD was defined as eGFR < 60 ml/min per 1.73 m² and/or presence of proteinuria (≥ 1+). The proteinuria number in each column includes all stage 1/2 patients plus a few in stage 3/4.




Figure 1 | Prevalence of chronic kidney disease according to the blood pressure (BP) classification in women (white bar) and men (black bar). The gender difference in the prevalence of chronic kidney disease increased in accordance with the severity of BP classification. Chronic kidney disease was defined as estimated glomerular filtration rate (eGFR) < 60 ml/min per 1.73 m² and/or the presence of proteinuria (≥ 1+).

The prevalence of CKD and/or proteinuria (≥ 1+) paralleled the severity of BP classification in both genders (Figure 1). The gender difference of CKD became greater and more prominent with increasing severity of BP classification.

Using multiple logistic regression analysis, the odds ratio for the presence of CKD was estimated. Hypertension was significantly associated with CKD in both genders. In contrast, only in men, but not in women, prehypertension with high-normal BP was significantly associated with an increased risk of CKD even after adjustment for confounders, such as age, obesity, current smoking, and daily drinking

(Table 3). We also reanalyzed the results of Table 3 after adjusting for serum glucose, triglyceride, high-density lipoprotein, and low-density lipoprotein levels: these factors had no influence on the association between prehypertension with high-normal BP and CKD in men (data not shown).

Lifestyle factors, obesity, and CKD

Obesity was positively associated with CKD in both genders, and eGFR was significantly decreased in the subjects with obesity compared with those without obesity (76.1 ± 16.2 versus 77.1 ± 16.0 ml/min per 1.73 m²; P < 0.001). When we reanalyzed the risk of CKD conferred by obesity in either the subjects with low eGFR (< 60 ml/min per 1.73 m²) or the subjects with proteinuria (≥ 1+), the conclusion remained unchanged (data not shown). In contrast, daily drinking was inversely associated with CKD in both genders. Additional analysis of the subgroup of subjects for whom daily alcohol intake data were available (n = 70,416 men and n = 75,416 women) revealed that the inverse association between daily drinking and CKD was consistent regardless of the amount of daily intake (≥ 23 g of ethanol or < 23 g of ethanol) in men (odds ratio (95% confidence interval, CI): 0.77 (0.73–0.80) and 0.89 (0.84–0.95), respectively; both P < 0.001); in women, the inverse association between daily drinking and CKD was found only in those with a daily intake of < 23 g of ethanol (odds ratio (95% CI): 0.91 (0.84–0.99); P = 0.03).

In women, current smoking status was positively associated with CKD. In contrast, among men, current smoking was inversely associated with CKD; that is, male current smokers had a significantly higher level of eGFR than current non-smokers (mean (95% CI) of eGFR: 79.0 (78.8–79.2) versus 75.9 (75.8–76.1) ml/min per 1.73 m²; P < 0.001). In contrast, there was no significant difference in eGFR between female current smokers and non-smokers (mean (95% CI) of eGFR: 77.0 (76.7–77.3) versus 76.9 (76.8–77.0) ml/min per 1.73 m²; P = 0.45). When we reanalyzed the association of current smoking with the presence

Table 3 | Odds ratio (95% confidence interval) for CKD by gender

	Women (n=142,293)		Men (n=89,732)	
	Odds ratio (95% confidence interval)	P-value	Odds ratio (95% confidence interval)	P-value
Age, 10 years	1.39 (1.37:1.42)	<0.001	1.82 (1.78:1.87)	<0.001
Obesity (0=no, 1=yes)	1.26 (1.22:1.31)	<0.001	1.43 (1.38:1.49)	<0.001
Current smoker (0=no, 1=yes)	1.34 (1.26:1.43)	<0.001	0.90 (0.86:0.94)	<0.001
Daily drinker (0=no, 1=yes)	0.92 (0.86:0.98)	0.006	0.78 (0.76:0.81)	<0.001
BP classification^a				
Optimal BP	1 (Reference)		1 (Reference)	
Prehypertension with normal BP	0.95 (0.91:1.00)	0.03	1.01 (0.96:1.07)	0.60
Prehypertension with high-normal BP	1.02 (0.97:1.06)	0.54	1.11 (1.05:1.17)	<0.001
Hypertension	1.17 (1.12:1.23)	<0.001	1.32 (1.25:1.38)	<0.001

Abbreviations: BP, blood pressure; CKD, chronic kidney disease.

Obesity was defined as body mass index (BMI) ≥ 25 kg/m². BP classification was defined as follows: optimal BP, systolic blood pressure (SBP) < 120 mm Hg and diastolic blood pressure (DBP) < 80 mm Hg; prehypertension with normal BP, SBP 120–129 mm Hg and/or 80–84 mm Hg; prehypertension with high-normal BP, SBP 130–139 mm Hg and/or DBP 85–89 mm Hg; hypertension, SBP ≥ 140 mm Hg and/or DBP ≥ 90 mm Hg. Statistical significance was defined as $P < 0.05$.

^aBP classification: Odds ratio was adjusted for age, obesity, current smoking, and daily drinking.

of proteinuria, there was a positive association between current smoking and proteinuria in both genders (odds ratio (95% CI): 1.47 (1.38–1.56) in men and odds ratio (95% CI): 1.89 (1.15–3.11) in women; both $P < 0.001$).

Effect of obesity on the association between CKD and BP classification

Among subjects without hypertension ($n = 181,215$), the risk of CKD conferred by prehypertension with high-normal BP increased when these conditions were accompanied by obesity (≥ 25 kg/m²) in men (Figure 2a), but not in women (Figure 2b). Accordingly, we examined whether or not there was an interaction between obesity and prehypertension with high-normal BP on CKD risk among subjects without hypertension. Using a multivariable logistic regression analysis, we showed that there was an additive effect, but not a synergistic one, of obesity and prehypertension with high-normal BP on CKD risk in men (data not shown). Furthermore, we also examined whether there was an interaction between obesity and hypertension ($\geq 140/90$ mm Hg) on CKD risk among all subjects ($n = 232,025$). The results showed that there was no synergistic interaction in either gender (data not shown).

DISCUSSION

Prehypertension and CKD

In this nationwide study of 232,025 Japanese aged 20 years or older, we have demonstrated the prevalence of CKD across the diagnostic spectrum of BP classification. In the present study, the prevalence of CKD was 17.0% in men and 12.2% in women. The prevalence was lower than a previous Japanese report,⁵ because the present study excluded treated hypertensive patients. In particular, we focused on the prevalence of CKD among subjects with prehypertension (16.7% in men and 12.2% in women). The prevalence of CKD among subjects with prehypertension with high-normal BP was greater in men than in women (18.1% versus 12.9%), and prehypertension with high-normal BP was an




Figure 2 | Logistic regression analysis of chronic kidney disease risk among subjects without hypertension. The odds ratio (95% confidence interval) of chronic kidney disease risk in subjects with or without obesity and/or prehypertension with high-normal blood pressure (BP) is shown in men (a) and women (b). The analysis was adjusted for age, current smoking, and daily drinking. Chronic kidney disease was defined as estimated glomerular filtration rate (eGFR) < 60 ml/min per 1.73 m² and/or the presence of proteinuria ($\geq 1+$). ^{*} $P < 0.001$ versus group (1) and [†] $P < 0.001$ versus group (3).

independent risk factor for CKD in men, but not in women, even after adjustment for confounders.

Evidence is accumulating that prehypertension, and particularly a high-normal BP range, is associated with a variety of cardiovascular diseases and cardiovascular-associated and all-cause mortality;^{15–17} however, information about the association of prehypertension with CKD is scarce in Japan.¹⁸ Much as in other previous reports worldwide,^{14–16} older age, higher prevalence of men, and obesity or obesity-related metabolic abnormalities were more prevalent in subjects with prehypertension than those with optimal BP (Table 2). These characteristics could partly explain the cardiovascular risk of prehypertension;^{15–17} however, our data show that the association between CKD and prehypertension with high-normal BP in men is independent of these confounders.

The increased risk of CKD among prehypertensive subjects with high-normal BP was recognized only in men; this means that the parallel increase of CKD in accordance with the level of severity of BP begins at an earlier phase in men than in women. This gender difference cannot be fully explained by the gender differences in metabolic factors or BP itself. It is speculated that it may be related to gender-specific differences in glomerular structure, hemodynamic condition, activity of local cytokines and hormones, gene expression, and/or the effects of sex hormones on kidney cells.^{9,19}

As shown in several previous reports,^{10–12} hypertension ($\geq 140/90$ mm Hg) is a clear risk factor for CKD in both the genders. In the present study, we excluded 84,854 subjects who had been treated with anti-hypertensive medication, and included 50,810 subjects who had never been treated with anti-hypertensive medication. This exclusion rate suggests that about a quarter of hypertensive subjects have not been treated for their condition. This proportion is substantially improved as compared with a previous report,²⁰ but more health promotion to increase awareness and treatment of hypertension is still considered necessary.

Obesity, BP, and CKD

Obesity is an independent risk factor for CKD both in men and women (Table 3). Intriguingly, our data indicate that the risk of CKD conferred by prehypertension with high-normal BP in men increased when these conditions were accompanied by obesity (Figure 2a). There was an additive effect of obesity and prehypertension with high-normal BP on CKD risk in men.

Obesity-associated glucose and lipid abnormalities could partly explain the increased risk of CKD in obesity.^{21,22} However, our data show that the increased risk of CKD conferred by obesity was independent of these confounders, although there was some lack of data on glucose and lipid parameters. There remain several other possible explanations for the risk of obesity. First, unmeasured obesity-associated factors, such as insulin resistance, inflammatory and oxidative stress, and abnormal adipocytokine production, may be involved in the increased risk of CKD in obesity.^{22,23} Second, obesity has a fairly consistent effect on renal

hemodynamics, suggestive of glomerular hypertension.^{24,25} At an early phase, obesity is associated with an elevated GFR with a less pronounced increase, or even a decrease, in effective renal plasma flow, resulting in an increased filtration rate. This alteration, that is, a predominant decrease in afferent rather than efferent glomerular tone in obese subjects, may confer enhanced renal susceptibility toward damage when BP increase is superimposed.^{24,25} Obesity-induced hyperfiltration, if continued for a certain period, can cause a decline in GFR, which may be one of the reasons why our data showed that obese subjects had a lower eGFR than nonobese subjects in both genders.

Lifestyle factors and CKD

Lifestyle factors, such as smoking and drinking, are also important contributors to CKD.²⁶ In the present study, an inverse association between CKD and current smoking was found (Table 3), despite the fact that several previous studies have identified smoking as an important risk factor in the promotion and progression of renal dysfunction in healthy subjects or those with complications.^{27,28} Our study is a cross-sectional study, and thus there may have been artifacts due to the observation of sick subjects after they have changed their lifestyles. However, the effects of smoking on eGFR are still controversial.^{27,29,30} In fact, we observed that male current smokers had a higher eGFR than male non-smokers, whereas no such association was found in women. On the other hand, our present results agree with previous reports;^{27,29} in that we found a positive association between smoking and proteinuria in both genders, suggesting the possibility that smoking causes endothelial dysfunction, partly through an inflammatory or oxidative pathway.^{28,29} It was also unexpected that there was an inverse association between the BP increase and the prevalence of current smoking (Table 2); this may have been attributable to one of the following: (1) Some of the smokers in the hypertensive group may have had knowledge that they were hypertensive, and may have ceased to smoke on the advice of their physicians. (2) There may have been a so-called survival effect, as smokers who develop hypertension were more likely to have died and thus not to have been included in the cross-sectional study. (3) Daytime BP under daily activity would likely be more elevated in smokers compared with non-smokers, even when there is no difference in the clinical or office BP between them (i.e., masked hypertension is more prevalent in smokers).³¹

Evidence on the association between CKD and alcohol intake has been scarce. We found that subjects with a daily drinking habit had a lower likelihood of CKD compared with those who had no alcohol intake. We could not assess the kinds or total amount of alcohol; therefore, to discuss this issue is beyond the scope of the present research. Further investigation with prospective or lifestyle interventional studies, such as smoking cessation studies, are warranted to better elucidate the impact of smoking or drinking on renal outcomes.

Several limitations of our study should be mentioned. First, we cannot infer a cause–effect relationship based on our cross-sectional data. Second, only a single measurement of serum creatinine, as well as only a single assessment of proteinuria, is not fully accurate, and thus there may be an underestimation of the true association between CKD and BP level. Third, subjects who participated in the present survey were generally healthy individuals who were interested in their health; therefore, the prevalence of prehypertension/hypertension or CKD may have been underestimated. Finally, little is known about the cost-effectiveness of screening male subjects with prehypertension and high-normal BP range for CKD; therefore, an additional study is needed to identify the most appropriate populations to undergo CKD screening.

CONCLUSION

Using a nationwide Japanese database, we show an increased prevalence of CKD across the diagnostic spectrum of hypertension. Among men, even in the state of prehypertension, high-normal BP, particularly when in conjunction with obesity, was an independent risk factor for CKD. Considering the fact that the prevalence of CKD and the incidence of end-stage renal disease are increasing in Japanese men,^{5,8,9} these data have important clinical implications; as CKD is often asymptomatic but progressive, more attention must be paid to men and women with hypertension or obesity and to men even with high-normal BP for the early detection and prevention of CKD, or to delay the progression to renal failure.

MATERIALS AND METHODS

Study population

The methods of the study are detailed in the Supplementary Information section online. Briefly, based on a recent survey that showed that obesity and metabolic syndrome are not uncommon in Japan (<http://www-bm.mhlw.go.jp/houdou/2008/04/h0430-2.html>), the Japanese government started a new health-care strategy that targeted early diagnosis and intervention for metabolic syndrome from 2008 (Specific Health Checkups and Guidance System (Tokutei-Kensin)). In this new health-care system, people diagnosed with metabolic syndrome are obligated to receive repeated lifestyle guidance over a 6-month period after an annual health examination.

Thirteen of the prefectures participating in this nationwide project (Yamagata, Miyagi, Fukushima, Niigata, Tokyo, Kanagawa, Ibaraki, Osaka, Okayama, Kochi, Fukuoka, Miyazaki, and Okinawa) agreed on our study purpose and were included in the present analysis. The population surveyed included a total of 346,942 subjects (41% ($n=141,938$) were men) above 20 years of age, for whom all the data necessary for our research purposes were available—namely, information about age, gender, BP, body mass index, habitual smoking or drinking, use of anti-hypertensive drugs, previous history of cardiovascular diseases (i.e., cardiac disease and stroke), and data about the serum creatinine level and dipstick urine test for proteinuria. This study was granted ethics approval from the respective institutional review boards. Data were sent to an independent data center called the NPO Japan Clinical Research Support Unit, and verified by trained staff.

Baseline measurement

At the baseline examination, all subjects completed a self-administered questionnaire about lifestyle factors (current smoking status, daily drinking), and provided medical information on treatment with anti-hypertensive drugs and a previous history of cardiac disease or stroke. The study physicians performed a physical examination of each subject and rechecked their medical history to improve the precision of the information.

According to the recommendations of the Ministry of Health, Labor and Welfare (<http://www.mhlw.go.jp/bunya/shakaihoshou/iryouseido01/info03a.html>), BP was measured by trained observers using a standard sphygmomanometer or an automated device on the right arm after resting for 5 min in a seated position with the legs not crossed. Conversation and alcohol/caffeine consumption should also be avoided before measurement. Subjects were classified according to their BP level as follows: optimal BP (systolic BP/diastolic BP < 120/80 mm Hg), prehypertension³² that comprises normal BP (systolic BP 120–129 mm Hg, diastolic BP 80–84 mm Hg or both) and high-normal BP (systolic BP 130–139 mm Hg, diastolic BP 85–89 mm Hg or both), and treated or untreated hypertension (systolic BP/diastolic BP \geq 140/90 mm Hg or usage of anti-hypertensive medication).³³

Body height and weight were measured in light clothing without shoes, and the body mass index was calculated (kg/m^2). According to the Japan Society for the Study of Obesity,³⁴ obesity was defined as a body mass index \geq 25 kg/m^2 .

Blood samples were collected after an overnight fast and were assayed within 24 h. For the purpose of our study, there were no missing data on the serum creatinine level, but there was a substantial lack of data on the glucose and lipid parameters (Supplementary Table S1 online). Freshly voided urine samples were tested by the dipstick methods in all subjects. Proteinuria was defined as 1+ or more.

Definition of CKD

Serum creatinine was assayed by an enzymatic method. eGFR was derived using the following equation:

$$\text{eGFR (ml/min per } 1.73 \text{ m}^2) = 194 \times \text{age (years)}^{-0.287} \times \text{serum creatinine (mg/dl)}^{-1.094} \text{ (if women } \times 0.739).^{35}$$

Details about this equation are also shown in the Supplementary Information section. CKD was defined as either the presence of proteinuria or eGFR < 60 ml/min per 1.73 m^2 . The clinical stages of CKD were classified according to the recommendations of the National Kidney Foundation Kidney Disease Outcomes Quality Initiative guidelines³⁶: Stage 1 or 2 (eGFR \geq 60 ml/min per 1.73 m^2 and the presence of proteinuria), Stage 3 (eGFR 30–59 ml/min per 1.73 m^2), Stage 4 (eGFR 15–29 ml/min per 1.73 m^2), and Stage 5 (eGFR < 15 ml/min per 1.73 m^2).

Statistical analysis

All statistical analyses were performed with the SPSS version 18.0J software (SPSS, Chicago, IL). The differences of patient characteristics and BP values according to the BP classification were assessed using analysis of variance, and categorical parameters were compared with the χ^2 -test. As there is a significant gender difference in the prevalence of CKD, we examined the association between CKD and the severity of BP classification separately in men and women. The odds ratio and 95% CI of each BP classification group (optimal BP group (reference) versus prehypertension with normal BP, prehypertension with high-normal BP, and untreated hypertension group) were calculated for the presence of CKD by multiple

logistic regression analysis. Finally, we used a multivariable logistic regression analysis to examine the effect of obesity on the association between CKD and BP classification, as well as whether or not there was an interaction between obesity and prehypertension with high-normal BP on CKD risk. Statistical significance was defined as $P < 0.05$.

DISCLOSURE

All the authors declared no competing interests.

ACKNOWLEDGMENTS

This work was supported by the Health and Labor Sciences Research Grants for 'Research on the positioning of CKD in Specific Health Check and Guidance in Japan' (20230601), the Ministry of Health, Labor, and Welfare of Japan. We acknowledge the contributions of the staff members, who collected data and instructed subjects with metabolic syndrome, at the regional screening centers providing data for this study: Yamagata, Miyagi Fukushima, Ibaraki, Tokyo, Kanagawa, Niigata, Osaka, Okayama, Kochi, Fukuoka, Miyazaki, and Okinawa. Follow-up screenings are ongoing.

SUPPLEMENTARY MATERIAL

Table S1. Glucose and lipid parameters according the BP classification by gender.

Supplementary material is linked to the online version of the paper at <http://www.nature.com/ki>

REFERENCES

- Levey AS, Atkins R, Coresh J et al. Chronic kidney disease as a global public health problem: approaches and initiatives—a position statement from Kidney Disease Improving Global Outcomes. *Kidney Int* 2007; **72**: 247–259.
- Schiffrin EL, Lipman ML, Mann JF. Chronic kidney disease: effects on the cardiovascular system. *Circulation* 2007; **116**: 85–97.
- Imai E, Horio M, Watanabe T et al. Prevalence of chronic kidney disease in the Japanese general population. *Clin Exp Nephrol* 2009; **13**: 621–630.
- Hall YN, Hsu CY, Iribarren C et al. The conundrum of increased burden of end-stage renal disease in Asians. *Kidney Int* 2005; **68**: 2310–2316.
- Nagata M, Ninomiya T, Doi Y et al. Trends in the prevalence of chronic kidney disease and its risk factors in a general Japanese population: the Hisayama Study. *Nephrol Dial Transplant* 2010; **25**: 2557–2564.
- Nakai S, Masakane I, Akiba T et al. Overview of regular dialysis treatment in Japan as of 31 December 2006. *Ther Apher Dial* 2008; **12**: 428–456.
- Imai E, Matsuo S, Makino H et al. Chronic Kidney Disease Japan Cohort study: baseline characteristics and factors associated with causative diseases and renal function. *Clin Exp Nephrol* 2010; **14**: 558–570.
- Iseki K, Iseki C, Ikemiya Y et al. Risk of developing end-stage renal disease in a cohort of mass screening. *Kidney Int* 1996; **49**: 800–805.
- Iseki K. Gender differences in chronic kidney disease. *Kidney Int* 2008; **74**: 415–417.
- O'Seaghdha CM, Perkovic V, Lam TH et al. Blood pressure is a major risk factor for renal death: an analysis of 560 352 participants from the Asia-Pacific region. *Hypertension* 2009; **54**: 509–515.
- Iseki K, Ikemiya Y, Kinjo K et al. Body mass index and the risk of development of end-stage renal disease in a screened cohort. *Kidney Int* 2004; **65**: 1870–1876.
- Yamagata K, Ishida K, Sairenchi T et al. Risk factors for chronic kidney disease in a community-based population: a 10-year follow-up study. *Kidney Int* 2007; **71**: 159–166.
- Kiberd B. The chronic kidney disease epidemic: stepping back and looking forward. *J Am Soc Nephrol* 2006; **17**: 2967–2973.
- Crews DC, Plantinga LC, Miller 3rd ER et al. Prevalence of chronic kidney disease in persons with undiagnosed or prehypertension in the United States. *Hypertension* 2010; **55**: 1102–1109.
- Pimenta E, Oparil S. Medscape. Prehypertension: epidemiology, consequences and treatment. *Nat Rev Nephrol* 2001; **6**: 21–30.
- Ishikawa Y, Ishikawa J, Ishikawa S et al. Prehypertension and the risk for cardiovascular disease in the Japanese general population: the Jichi Medical School Cohort Study. *J Hypertens* 2010; **28**: 1630–1637.
- Kalaitzidis RG, Bakris GL. Prehypertension: is it relevant for nephrologists? *Kidney Int* 2010; **77**: 194–200.
- Ninomiya T, Kubo M, Doi Y et al. Prehypertension increases the risk for renal arteriosclerosis in autopsies: the Hisayama Study. *J Am Soc Nephrol* 2007; **18**: 2135–2142.
- Iseki K, Iseki C, Ikemiya Y et al. Risk of developing end-stage renal disease in a cohort of mass screening. *Kidney Int* 1996; **49**: 800–805.
- Tanaka T, Okamura T, Yamagata Z et al. Awareness and treatment of hypertension and hypercholesterolemia in Japanese workers: the High-risk and Population Strategy for Occupational Health Promotion (HIPOP-OHP) study. *Hypertens Res* 2007; **30**: 921–928.
- Wahba IM, Mak RH. Obesity and obesity-initiated metabolic syndrome: mechanistic links to chronic kidney disease. *Clin J Am Soc Nephrol* 2007; **2**: 550–562.
- Guarnieri G, Zanetti M, Vinci P et al. Metabolic syndrome and chronic kidney disease. *J Ren Nutr* 2010; **20**(5 Suppl): S19–S23.
- Wisse BE. The inflammatory syndrome: the role of adipose tissue cytokines in metabolic disorders linked to obesity. *J Am Soc Nephrol* 2004; **15**: 2792–2800.
- Ribstein J, du Cailar G, Mimran A. Combined renal effects of overweight and hypertension. *Hypertension* 1995; **26**: 610–615.
- Krikken JA, Bakker SJ, Navis GJ. Role of renal haemodynamics in the renal risks of overweight. *Nephrol Dial Transplant* 2009; **24**: 1708–1711.
- Appel LJ. Lifestyle modification as a means to prevent and treat high blood pressure. *J Am Soc Nephrol* 2003; **14**(7 Suppl 2): S99–S102.
- Ishizaka N, Ishizaka Y, Toda E et al. Association between cigarette smoking and chronic kidney disease in Japanese men. *Hypertens Res* 2008; **31**: 485–492.
- Orth SR. Effects of smoking on systemic and intrarenal hemodynamics: influence on renal function. *J Am Soc Nephrol* 2004; **15**(Suppl 1): S58–S63.
- Sauriasari R, Sakano N, Wang DH et al. C-reactive protein is associated with cigarette smoking-induced hyperfiltration and proteinuria in an apparently healthy population. *Hypertens Res* 2010; **33**: 1129–1136.
- Shankar A, Klein R, Klein BE. The association among smoking, heavy drinking, and chronic kidney disease. *Am J Epidemiol* 2006; **164**: 263–271.
- Mann SJ, James GD, Wang RS et al. Elevation of ambulatory systolic blood pressure in hypertensive smokers. A case-control study. *JAMA* 1991; **265**: 2226–2228.
- Chobanian AV, Bakris GL, Black HR et al. The seventh report of the joint national committee on prevention, detection, evaluation, and treatment of high blood pressure: the JNC 7 report. *JAMA* 2003; **289**: 2560–2572.
- Ogihara T, Kikuchi K, Matsuoka H et al. The Japanese society of hypertension guidelines for the management of hypertension (JSH 2009). *Hypertens Res* 2009; **32**: 3–107.
- Examination Committee of Criteria for 'Obesity Disease' in Japan; Japan Society for the Study of Obesity. New criteria for 'obesity disease' in Japan. *Circ J* 2002; **66**: 987–992.
- Matsuo S, Imai E, Horio M et al. Revised equations for estimated GFR from serum creatinine in Japan. *Am J Kidney Dis* 2009; **53**: 982–992.
- National Kidney Foundation. K/DOQI clinical practice guidelines for chronic kidney disease: evaluation, classification, and stratification. *Am J Kidney Dis* 2002; **39**(2 Suppl 1): S1–S266.

Risk factor profiles based on estimated glomerular filtration rate and dipstick proteinuria among participants of the Specific Health Check and Guidance System in Japan 2008

Kunitoshi Iseki · Koichi Asahi · Toshiki Moriyama · Kunihiro Yamagata · Kazuhiko Tsuruya · Hideaki Yoshida · Shoichi Fujimoto · Tsuneo Konta · Issei Kurahashi · Yasuo Ohashi · Tsuyoshi Watanabe

Received: 30 June 2011 / Accepted: 12 October 2011
© Japanese Society of Nephrology 2011

Abstract

Background Estimated glomerular filtration rate (eGFR) and albuminuria (proteinuria) are both important determinants of the risk of cardiovascular disease (CVD), end-stage renal disease (ESRD), and mortality. Few studies, however, have examined the risk factor profiles based on eGFR and proteinuria among the general population.

Methods Data of the newly developed nationwide screening program of the Specific Health Check-up and Guidance System (Tokutei-Kensin) initiated in 2008 were used in this study. The aim of this screening, targeting people 40–74 years of age, was to detect those with metabolic syndrome and to offer those services regarding lifestyle modifications that will lead to the reduction of diabetes mellitus (DM) and DM-related ESRD. Individual records of 580,000 participants in 69 cities and towns and 3 union cohorts throughout Japan were anonymously provided and included in the present study.

Results Details of 332,174 participants (57.3% of the total) with both serum creatinine and dipstick urine test

data were analyzed. Mean (SD) age was 63.6 (8.3) years and 40.6% were men. The mean (SD) eGFR was 75.0 (16.2) ml/min/1.73 m² and 5.4% had proteinuria. The prevalence of chronic kidney disease (CKD) stage 3, 4, and 5 was 14.2%, 0.2%, and 0.07%, respectively. The prevalence of DM, hypertension, and history of stroke and heart disease was correlated with the combination of eGFR and degree of proteinuria.

Conclusion The findings of the present study indicate that CKD and risk factors for CVD are quite common among middle-aged Japanese. CKD classification based on eGFR and proteinuria may be useful for predicting CVD, mortality rate, and ESRD in the Japanese population.

Keywords eGFR · CKD · Screening · Proteinuria · Epidemiology

Introduction

Chronic kidney disease (CKD) is a common condition and is a risk factor for developing cardiovascular disease (CVD) and end-stage renal disease (ESRD) [1]. Both the prevalence and incidence of treated ESRD are very high in Japan [2]. Furthermore, the incidence and prevalence continue to increase, despite several preventive strategies aimed at early detection and treatment of CKD. Japan has a long history of universal screening, a program that might facilitate the early detection of CKD [3]. A higher mean age at the start of dialysis can be interpreted as delaying the progression of CKD, but it may also simply reflect the increase in the elderly population and longevity. Dipstick proteinuria is a strong predictor of developing ESRD in a setting of community-based screening [4]. Delayed visits to the nephrology clinic result in an inevitable initiation of

K. Iseki (✉)
Dialysis Unit, University Hospital of the Ryukyus, 207 Uehara,
Nishihara, Okinawa 903-0215, Japan
e-mail: chihokun@med.u-ryukyu.ac.jp

K. Iseki · K. Asahi · T. Moriyama · K. Yamagata ·
K. Tsuruya · H. Yoshida · S. Fujimoto ·
T. Konta · T. Watanabe
Steering Committee for “Research on the Positioning of Chronic
Kidney Disease (CKD) in the Specific Health Check and
Guidance System of Japan, Tokyo, Japan

I. Kurahashi · Y. Ohashi
Department of Biostatistics/Epidemiology and Preventive Health
Sciences, School of Health Sciences and Nursing,
University of Tokyo, Tokyo, Japan

dialysis with a short duration of follow-ups [5, 6]. Such ‘late referral’ negatively impacts survival after dialysis is initiated. Preliminary results of the Japanese Society Dialysis Therapy support the notion that the longer the duration of pre-ESRD treatment, the better the survival. Because CKD remains asymptomatic until the late stages, effective strategies for the early detection and treatment of CKD are necessary.

The increasing prevalence of obesity and diabetes mellitus (DM) has become the leading cause of ESRD. A specific nationwide health check-up and guidance system, called Tokutei-Kenshin, was initiated in April 2008 in Japan (The Ministry of Health, Labour and Welfare; <http://www-admin.mhlw.go.jp>). The aim of this project is to detect metabolic syndrome and if confirmed, to provide individual instruction to modify lifestyle and the necessary treatment. The target population comprises Japanese citizens between the ages of 40–74 years. Data on the prevalence of risk factors for developing CKD, ESRD, and CVD are limited to the Japanese population. In the present study, we examined the demographics of participants of the newly developed screening system in Japan. Risk factor profiles were examined according to the new CKD classification based on the combination of estimated glomerular filtration rate (eGFR) and dipstick proteinuria findings [7]. Results of dipstick proteinuria were categorized into three groups: (–) and (±), 1+, and ≥2+. The present study provides the baseline characteristics for the future outcome study as the unique identification number was set by the government.

Methods

Individual records for 580,000 participants in 12 communities or prefectures were anonymously provided and included in this analysis. Among these participants, subjects with data for both serum creatinine and dipstick proteinuria were selected for this study. A test was mandatory for dipstick proteinuria, but not for serum creatinine. Therefore, rates of measurement of serum creatinine differ among cohorts or prefectures. Databases included in this study were from Yamagata, Miyagi, Fukushima, Ibaraki, Tokyo, Kanagawa, Niigata, Osaka, Okayama, Kochi, Fukuoka, Miyazaki and Okinawa, and ethical approval was obtained from the respective institute review boards. Data were sent to a data center called the NPO Japan Clinical Research Support Unit to be verified. Outliers were deleted through winsorization and accounted for 0.01–0.1% of the total. Eligible participants visited a pre-assigned clinic or hospital and responded to a questionnaire regarding past history of stroke, cardiac disease, kidney disease, lifestyle habits such as smoking, alcohol intake, walking, etc., and medications for

hypertension, DM, and dyslipidemia. Screening participants are eligible for public support for the standard health checks, such as measurement of height, weight, waist circumference, blood pressure, fasting blood glucose, hemoglobin A1c, triglyceride, serum high-density lipoprotein (HDL) cholesterol, low-density lipoprotein (LDL) cholesterol, glutamyl oxaloacetic transaminase, glutamate pyruvate transaminase, gamma-glutamyl transpeptidase, hemoglobin, uric acid, serum creatinine, dipstick urine test for proteinuria, hematuria, and glucosuria. Proteinuria was coded as (–), (±), (1+), (2+), and (3+). Serum creatinine was measured using the enzymatic method. Glomerular filtration rate was calculated using the formula of the Japanese Society of Nephrology [8]. Reference levels for triglyceride, HDL cholesterol, LDL cholesterol, uric acid, fasting blood glucose, and hemoglobin A1c were set at 150, 40, 7, 110 mg/dl, and 6.1%, respectively. Blood pressure was measured in all cohorts using a standard sphygmomanometer. Hypertension was defined as ≥140/90 mmHg or on antihypertensive medication. DM was defined as hemoglobin A1c ≥6.1% or on medication for DM. Obesity was defined as body mass index (BMI) ≥25 kg/m².

Statistical analysis

Data were analyzed with SAS/STAT software (version 6.03, SAS Institute, Tokyo, Japan). Student’s *t* test and the Chi-squared test were performed to compare the significance of discrete variables. A *P* value of less than 0.05 was considered statistically significant in all analyses.

Results

Demographics of the screened subjects are summarized in Table 1. The prevalence of CKD (i.e., eGFR <60 ml/min/1.73 m²) was as high as 14.2%. Compared to national statistics, smoking rates were lower in both men and women than in the general population. Those with low eGFR comprised 14.2% and proteinuria was distributed as follows: negative and ± 94.55%, 1+ 3.75%, and ≥2+ 1.7%.

The prevalence of obesity, DM, and hypertension is summarized based on the results of eGFR and proteinuria in Table 2. The prevalence of obesity, DM, and hypertension increased in relation to the degree of proteinuria in each eGFR group. Higher levels of proteinuria together with lower levels of eGFR were associated with an increased prevalence of hypertension (Fig. 1).

History of stroke, heart disease, and CVD (either stroke or heart disease) is summarized in Table 3. The prevalence of CVD was highest (25.2%) in those with proteinuria of (1+) and an eGFR of 15–29 ml/min/1.73 m², and the prevalence was lowest (6.1%) in those negative and ± for

Table 1 Demographics of the screened cohorts. Screening was performed during April 1, 2008 to March 31, 2009

Number of participants	332,174		
Men (%)	134,751 (40.6)		
Age (years)	63.6 (8.3), 40–74		
Body height (cm)	157.2 (8.5)	Men 164.6 (6.3) [#]	Women 152.2 (5.7)
Body weight (kg)	57.6 (10.5)	Men 64.5 (9.5) [#]	Women 52.8 (8.3)
Body mass index (kg/m ²)	23.2 (3.3)	Men 23.8 (3.1) [#]	Women 22.8 (3.5)
Waist circumference (cm)	84.1 (9.2)	Men 85.7 (8.3) [#]	Women 83 (9.5)
Systolic blood pressure (mmHg)	128.9 (17.4)		
Diastolic blood pressure (mmHg)	76.3 (10.7)		
Fasting blood glucose (mg/dl)	98.2 (21.5)		
Hemoglobin A1c (%)	5.3 (0.7)		
Triglyceride (mg/dl)	122.5 (84.0)		
HDL cholesterol (mg/dl)	62.1 (16.3)		
LDL cholesterol (mg/dl)	125.9 (30.6)		
Hemoglobin (g/dl)	13.6 (1.4)		
Serum creatinine (mg/dl)	0.7 (0.2)	Men 0.8 (0.3) [#]	Women 0.6 (0.2)
eGFR (ml/min/1.73 m ²)	75.0 (16.2)		
<15	240 (0.07%)		
15–29	655 (0.20%)		
30–44	4,300 (1.29%)		
45–59	42,975 (12.94%)		
60–89	225,081 (67.76%)		
≥90	58,923 (17.74%)		
Serum uric acid (mg/dl)	5.2 (1.4)	Men 6.0 (1.3) [#]	Women 4.7 (1.1)
Glucosuria ^a	2.30%		
Proteinuria ^a	5.40%		
Hematuria ^a	7.50%		
Past history (%)			
Stroke	3.30		
Cardiac disease	6.00		
Kidney disease	0.70		
Medication (%)			
Anti-hypertensive drugs	28.80		
Lipid lowering drugs	15.80		

Table 1 continued

Insulin or hypoglycemic drugs	5.20		
Lifestyle (%)			
Smoking	13.50	Men 25.2% [#]	Women 5.5%
Drinking	44.30	Men 65.2% [#]	Women 30.0%

Data are mean (SD)

^a Positive urine test denote ≥1+ by dipstick[#] *P* < 0.01 (vs women)

proteinuria and having an eGFR ≥ 90 ml/min/1.73 m². The combination of higher levels of proteinuria and lower levels of eGFR was associated with an increased prevalence of a history of CVD (Fig. 2).

Mean (SD) levels of BMI and smoking rate are summarized in Table 4. Both BMI and smoking rate were higher in men than in women. The smoking rate tended to decrease in the lower eGFR category.

Discussion

The target population of this screening in Japan comprised participants from 40 to 74 years of age, and the expected turnout was approximately 58 million. In the 2008 screening, the actual participation rate remained low, 20–30%, probably because of the lack of preparation for implementing this new system. The total number of participants in the present study was approximately 0.58 million; therefore, our analysis included at least 1% of the target population in Japan.

The results revealed the current health status among the general Japanese population. The proportion of the population comprising elderly people is high in Japan and its rate of increase is currently the highest in the world. The proportion of those with a low GFR (<60 ml/min/1.73 m²), regardless of proteinuria, increases with aging. Fortunately, the rate of decline of the GFR in the Japanese is relatively low, 0.36 ml/min/1.73 m²/year [9]. Elderly people are at risk for CVD and death. Effective strategies to establish a health check and guidance system are necessary to better accommodate the future burden of medical and social costs due to the aging population. Based on the findings of the present study, we propose that a cost–benefit analysis be performed on programs designed for the early detection and treatment of CKD, including education regarding lifestyle modification.

CVD is a recently recognized risk factor for CKD and ESRD [7]. The prevalence of CVD in CKD stage 5 is

Table 2 Prevalence of obesity, DM, and hypertension based on the combination of eGFR and proteinuria

eGFR	Proteinuria	Number (%)	Obesity (%)	DM (%)	Hypertension (%)
<15	Minus, ±	101 (0.03)	22.8	10.2	50.5 [#]
	1+	35 (0.01)	28.6	17.6	88.6*
	≥2+	103 (0.03)	31.1	32.0*	92.2*
15–29	Minus, ±	251 (0.08)	29.1	25.2*	81.1*
	1+	119 (0.04)	31.1	22.7*	85.5*
	≥2+	285 (0.09)	40.3*	38.4*	91.9*
30–44	Minus, ±	3,194 (0.96)	35.8*	13.9*	68.2*
	1+	504 (0.15)	43.8*	24.9*	83.3*
	≥2+	579 (0.17)	46.5*	33.9*	85.9*
45–59	Minus, ±	39,265 (11.82)	30.7*	9.3*	52.7*
	1+	2,408 (0.72)	42.2*	19.6*	71.6*
	≥2+	1,326 (0.40)	49.5*	31.1*	80.2*
60–89	Minus, ±	214,768 (64.66)	25.4*	8.4*	43.3*
	1+	7,579 (2.28)	38.5*	19.3*	62.1*
	≥2+	2,703 (0.81)	46.5*	31.1*	72.0*
≥90	Minus, ±	56,495 (17.01)	24.3	10.6	39.8
	1+	1,812 (0.55)	37.9*	26.5*	57.4*
	≥2+	647 (0.19)	46.5*	36.4*	72.0*

Total number of participants was 332,174. Parentheses are the percentage to the total participants in each column. Obesity, BMI ≥25 kg/m²; DM, HbA1c ≥6.1% or on treatment; hypertension, 140/90 mmHg or on treatment

* $P < 0.0001$, [#] $P < 0.05$ (vs. reference value of eGFR ≥90 and proteinuria minus or ±)

Prevalence of Hypertension




Fig. 1 Prevalence of hypertension by the combination of eGFR and proteinuria. Prevalence of hypertension was significantly ($P < 0.0001$) higher in every column except those with eGFR 15–29 and proteinuria minus or (±) ($P < 0.05$) when compared to the reference value of eGFR ≥90 and proteinuria minus or (±)

approximately 25%; similar to the prevalent dialysis population. Ethnic variations in CVD incidence and subtype are well described in the general population [10, 11]. The stroke mortality rate is high in Japan; however, in the present study, the prevalence of stroke was lower than that of cardiac disease (Table 1). The reasons for this finding are not clear, but many people with stroke are unable to participate in this type of screening program.

Metabolic syndrome is an important risk factor for developing CKD [12], and for DM and hypertension, which are the main causes of ESRD [13]. We previously reported

Table 3 Prevalence of history of stroke and heart disease based on the combination of eGFR and proteinuria

eGFR	Proteinuria	Stroke	Heart Disease	CVD
<15	Minus, ±	4.0	6.9	10.9*
	1+	5.7	17.1*	22.9
	≥2+	8.7*	12.6	19.4 [#]
15–29	Minus, ±	13.9*	15.9*	25.1*
	1+	15.1*	15.1*	25.2*
	≥2+	11.6*	13.7*	22.5*
30–44	Minus, ±	8.6*	13.1*	19.2*
	1+	9.9*	16.1*	22.4*
	≥2+	10.5*	16.4*	23.7*
45–59	Minus, ±	4.8*	8.5*	12.3*
	1+	6.9*	11.7*	16.2*
	≥2+	8.3*	13.1*	19.2*
60–89	Minus, ±	3.0*	5.6*	8.1*
	1+	4.7*	7.3*	11.1*
	≥2+	5.8*	9.3*	13.8*
≥90	Minus, ±	2.4	4.1	6.1
	1+	3.5 [†]	6.0*	8.8*
	≥2+	4.5 [†]	4.6	8.5 [†]

Total number of participants was 332,174. Cardiovascular disease (CVD) denotes stroke and/or heart disease

* $P < 0.0001$, [†] $P < 0.02$, [#] $P < 0.05$ (vs reference value of eGFR ≥90 and proteinuria minus or ±)

the significance of obesity in the risk for ESRD. Recent societal changes in lifestyle related to motorized transportation and high-calorie intake may have contributed to the