

## References

1. Prall F, Nollau P, Neumaier M, Haubeck HD, Drzeniek U, Helmchen U, Loning T, Wagener C: CD66a (BGP), an adhesion molecule of the carcinoembryonic antigen family, is expressed in epithelium, endothelium, and myeloid cells in a wide range of normal human tissues. *J Histochem Cytochem* 1996, 44:35–41
2. Kammerer R, Hahn S, Singer BB, Luo JS, von Kleist S: Biliary glycoprotein (CD66a), a cell adhesion molecule of the immunoglobulin superfamily, on human lymphocytes: structure, expression and involvement in T cell activation. *Eur J Immunol* 1998, 28:3664–3674
3. Gray-Owen SD, Blumberg RS: CEACAM1: contact-dependent control of immunity. *Nat Rev Immunol* 2006, 6:433–446
4. Hammarstrom S: The carcinoembryonic antigen (CEA) family: structures, suggested functions and expression in normal and malignant tissues. *Semin Cancer Biol* 1999, 9:67–81
5. Izzi L, Turbide C, Houde C, Kunath T, Beauchemin N: Cis-determinants in the cytoplasmic domain of CEACAM1 responsible for its tumour inhibitory function. *Oncogene* 1999, 18:5563–5572
6. Morales VM, Christ A, Watt SM, Kim HS, Johnson KW, Utku N, Texeira AM, Mizoguchi A, Mizoguchi E, Russell GJ, Russell SE, Bhan AK, Freeman GJ, Blumberg RS: Regulation of human intestinal intraepithelial lymphocyte cytolytic function by biliary glycoprotein (CD66a). *J Immunol* 1999, 163:1363–1370
7. Nakajima A, Iijima H, Neurath MF, Nagaishi T, Nieuwenhuis EES, Raychowdhury R, Glickman J, Blau DM, Russell S, Holmes KV, Blumberg RS: Activation-induced expression of carcinoembryonic antigen-cell adhesion molecule 1 regulates mouse T lymphocyte function. *J Immunol* 2002, 168:1028–1035
8. Boulton IC, Gray-Owen SD: Neisserial binding to CEACAM1 arrests the activation and proliferation of CD4<sup>+</sup> T lymphocytes. *Nat Immunol* 2002, 3:229–236
9. Markel G, Wolf D, Hanna J, Gazit R, Goldman-Wohl D, Lavy Y, Yagel S, Mandelboim O: Pivotal role of CEACAM1 protein in the inhibition of activated decidual lymphocyte functions. *J Clin Inv* 2002, 110:943–953
10. Chen D, Iijima H, Nagaishi T, Nakajima A, Russell S, Raychowdhury R, Morales V, Rudd CE, Utku N, Blumberg RS: Carcinoembryonic antigen-related cellular adhesion molecule 1 isoforms alternatively inhibit and costimulated human T cell function. *J Immunol* 2004, 172:3535–3543
11. Chen CJ, Shively JE: The cell-cell adhesion molecule carcinoembryonic antigen-related cellular adhesion molecule 1 inhibits IL-2 production and proliferation in human T cells by association with Src homology protein-1 and down-regulates IL-2 receptor. *J Immunol* 2004, 172:3544–3552
12. Iijima H, Neurath MF, Nagaishi T, Glickman JN, Nieuwenhuis EE, Nakajima A, Chen D, Fuss IJ, Utku N, Lewicki DN, Becker C, Gallagher TM, Holmes KV, Blumberg RS: Specific regulation of T helper cell 1-mediated murine colitis by CEACAM1. *J Exp Med* 2004, 199:471–482
13. Wekerle H: Experimental autoimmune encephalomyelitis as a model of immune-mediated CNS disease. *Curr Opin Neurobiol* 1993, 3:779–784
14. Steinman L: Assessment of animal models for MS and demyelinating disease in the design of rational therapy. *Neuron* 1999, 24:511–514
15. Hemmer B, Archelos JJ, Hartung HP: New concepts in the immunopathogenesis of multiple sclerosis. *Nat Rev Neurosci* 2002, 3:291–301
16. Sospedra M, Martin R: Immunology of multiple sclerosis. *Annu Rev Immunol* 2005, 23:683–747
17. Willenborg DO, Fordham S, Bernard CC, Cowden WB, Ramshaw IA: IFN- $\gamma$  plays a critical down-regulatory role in the induction and effector phase of myelin oligodendrocyte glycoprotein-induced autoimmune encephalomyelitis. *J Immunol* 1996, 157:3223–3227
18. Krakowski M, Owens T: Interferon- $\gamma$  confers resistance to experimental allergic encephalomyelitis. *Eur J Immunol* 1996, 26:1641–1646
19. Willenborg DO, Fordham SA, Staykova MA, Ramshaw IA, Cowden WB: IFN- $\gamma$  is critical to the control of murine autoimmune encephalomyelitis and regulates both in the periphery and in the target tissue: a possible role for nitric oxide. *J Immunol* 1999, 163:5278–5286
20. Tran EH, Prince EN, Owens T: IFN- $\gamma$  shapes immune invasion of the central nervous system via regulation of chemokines. *J Immunol* 2000, 164:2759–2768
21. Bettelli E, Sullivan B, Szabo SJ, Sobel RA, Glimcher LH, Kuchroo VK: Loss of T-bet, but not STAT1, prevents the development of experimental autoimmune encephalomyelitis. *J Exp Med* 2004, 200:79–87
22. Cua DJ: Interleukin-23 rather than interleukin-12 is the critical cytokine for autoimmune inflammation of the brain. *Nature* 2003, 421:744–748
23. Steinman L: A brief history of T(H)17, the first major revision in the T(H)1/T(H)2 hypothesis of T cell mediated tissue damage. *Nat Med* 2007, 13:139–145
24. Komiya Y, Nakae S, Matsuki T, Nambu A, Ishigame H, Kakuta S, Sudo K, Iwakura Y: IL-17 plays an important role in the development of experimental autoimmune encephalomyelitis. *J Immunol* 2006, 177:566–73
25. Chitinis T: Effect of targeted disruption of STAT4 and STAT6 on the induction of experimental autoimmune encephalomyelitis. *J Clin Invest* 2001, 108:739–747
26. Korn T: Myelin-specific regulatory T-cells accumulate in the central nervous system, but fail to suppress pathogenic effector T-cells at the peak of autoimmune inflammation. *Nat Med* 2007, 13:423–431
27. Bettelli E, Oukka M, Kuchroo VK: TH-17 cells in the circle of immunity and autoimmunity. *Nat Immunol* 2007, 8:345–350
28. Stromnes IM, Cerretti LM, Liggitt D, Harris RA, Goverman JM: Differential regulation of central nervous system autoimmunity by Th1 and Th17 cells. *Nat Med* 2007, 14:337–342
29. Kroenke MA, Carlson TJ, Andjelkovic AV, Segal BM: IL-12- and IL-23-modulated T cells induce distinct types of EAE based on histology. CNS chemokine profile, and response to cytokine inhibition. *J Exp Med* 2008, 205:1535–1541
30. Bendala A, Savage PB, Teyton L: The biology of NKT cells. *2007*;25:297–336
31. Kronenberg M: Toward an understanding of NKT cell biology: progress and paradoxes. *Annu Rev Immunol* 2005, 23:877–900
32. Miyake S, Yamamura T: Therapeutic potential of CD1d-restricted invariant natural killer T cell-based treatment for autoimmune diseases. *Int Rev Immunol* 2007, 26:73–94
33. Kuprina NI, Baranov VN, Yazova AK, Rudinskaya TD, Escobedo M, Cordier J, Gleiverman AS, Goussev AI: The antigen of bile canaliculi of the mouse hepatocyte: identification and ultrastructural localization. *Histochemistry* 1990, 94:179–186
34. Gallagher TM: A role for naturally occurring variation of the murine coronavirus spike protein in stabilizing association with the cellular receptor. *J Virol* 1997, 71:3129–3137
35. Miyamoto K, Miyake S, Mizuno M, Oka N, Kusunoki S, Yamamura T: Selective COX-2 inhibitor celecoxib prevents experimental autoimmune encephalomyelitis through COX-2-independent pathway. *Brain* 2006, 129:1984–1992
36. Kammerer R, Stober D, Singer BB, Obrink B, Reimann J: Carcinoembryonic antigen-related cell adhesion molecule 1 on murine dendritic cells in a potent regulator or T cell stimulation. *J Immunol* 2001, 166:6537–6544
37. Korn T, Bettelli E, Gao W, Awasthi A, Jager A, Strom TB, Oukka M, Kuchroo VK: IL-21 initiates an alternative pathway to induce proinflammatory T<sub>H</sub>17 cells. *Nature* 2007, 448:484–487
38. Mars LT, Araujo L, Kerschen P, Diem S, Bourgeois E, Van LP, Carrie N, Dy M, Liblau RS, Herbelin A: Invariant NKT cells inhibit development of the Th<sub>17</sub> lineage. *Proc Natl Acad Sci USA* 2009, 106:6238–6243
39. Yokote H, Miyake S, Croxford JL, Oki S, Mizusawa H, Yamamura T: NKT cell-dependent amelioration of a mouse model of multiple sclerosis by altering gut flora. *Am J Pathol* 2008, 173:1714–1723
40. Gray-Owen SD, Lorenzen DR, Hude A, Meyer TF, Dehio C: Differential Opa specificities for CD66 receptors influence tissue interactions and cellular response to *Neisseria Gonorrhoeae*. *Mol Microbiol* 1997, 26:971–980
41. Virji M, Makepeace K, Ferguson DJP, Watt SM: Carcinoembryonic antigens (CD66) on epithelial cells and neutrophils are receptors for Opa proteins of pathogenic neisseriae. *Mol Microbiol* 1996, 22:941–950
42. Hill DJ, Toleman MA, Evans DJ, Villullas S, Van AL, Virji M: The variable P5 proteins of typeable and non-typeable *Haemophilus influenzae* target human CEACAM1. *Mol Microbiol* 2001, 39:850–862

43. Toleman M, Aho E, Virji M: Expression of pathogen-like Opa adhesions in commensal *Neisseria*: genetic and functional analysis. *Cell Microbiol* 2001, 3:33–44
44. Virji M, Evams D, Griffith J, Hill D, Serino L, Hadfield A, Watt SM: Carcinoembryonic antigens are targeted by diverse strains of typable and non-typable *Haemophilus influenzae*. *Mol Microbiol* 2000, 36:784–795
45. Berger CN, Billker O, Meyer TF, Servin AL, Kansau I: Differential recognition of members of the carcinoembryonic antigen family by Afa/Dr adhesins of diffusely adhering *Escherichia coli* (Afa/Dr DAEC). *Mol Microbiol* 2004, 52:963–983
46. Miyamoto K, Miyake S, Yamamura T: A synthetic glycolipid prevents autoimmune encephalomyelitis by inducing TH2 bias of natural killer T cells. *Nature* 2001, 413:531–534

# Ghrelin: Friend or Foe for Neuroinflammation

SACHIKO MIYAKE AND TAKASHI YAMAMURA

**Abstract:** Ghrelin is a recently identified gastric hormone that displays strong growth hormone (GH) releasing activity mediated by the GH secretagogue receptor (GHS-R). While this unique endogenous peptide participates in the regulation of energy homeostasis, increases food intake, and decreases energy expenditure, its ability to modulate immune regulation is another important feature. Here we discuss the effect of ghrelin on the immune system. Ghrelin was initially reported as an immune enhancing factor. More recently, however, the immunosuppressive effects of ghrelin have been found in several animal models including bowel disease, arthritis, and sepsis and endotoxemia. We recently demonstrated that exogenous administration of ghrelin suppressed experimental autoimmune encephalomyelitis, an animal model of multiple sclerosis in association with the reduction of pro-inflammatory cytokines in microglia. These results shed light on the new role of ghrelin in the regulation of disorders that pro-inflammatory cytokines contribute to the pathogenesis such as neuroinflammatory and mental diseases.

Malnourished people are more susceptible to

---

*Sachiko Miyake, M.D., Ph.D., and Takashi Yamamura, M.D., Ph.D., is at the Department of Immunology, National Institute of Neuroscience, NCNP, Tokyo 187-8502, Japan.  
Corresponding author: Dr. Sachiko Miyake (Sachiko Miyake: miyake@ncnp.go.jp).*

pathogens, yet it is said, though mostly anecdotal, that people with autoimmune diseases such as multiple sclerosis or rheumatoid arthritis may claim that their symptoms can be reduced by fasting. Although little was known about the mechanistic link between starvation and immunity, recent studies have shed light on the immunomodulatory potency of a range of feeding regulatory hormones such as ghrelin, leptin, and neuropeptide Y (NPY). For example, serum leptin is decreased after acute starvation in parallel with immunosuppression or Th2 bias, whereas exogenous leptin would correct the altered Th1/Th2 balance towards Th1 (Lord et al., 1998; Sanna et al., 2003). In contrast, endogenous NPY is increased after starvation. Exogenous NPY has been reported to shift the Th1/Th2 balance towards Th2 and can ameliorate the severity of EAE (Bedoui et al., 2003). More recent study using NPY1 receptor-deficient mice has shown that NPY has the bimodal role in altering immune responses (Wheway et al., 2005). Namely, a new role of NPY to promote APC activation has emerged in addition to its role for downregulating Th1-responses. Interestingly, both peptide hormones are linked to ghrelin, a focus of this short review, in an endocrine feedback system. Ghrelin level increased after starvation, and can potently stimulate the release of NPY in the central nervous system (CNS) (Cowley et al., 2003). Moreover, ghrelin shows antagonistic effects against leptin (Kalra et al., 2005).

## The effect of ghrelin on immune system

Ghrelin is a gastric hormone first identified in the rat stomach in 1999 as a mediator of growth hormone (GH) release (Kojima et al., 1999). The bio-

logical effect of ghrelin is mediated by a G protein-coupled receptor called GH secretagogue receptor (GHS-R). Although ghrelin is predominantly secreted from mucosal endocrine cells of stomach, ghrelin and GHS-R are widely distributed in various organs. Furthermore, it is measurable in the systemic circulation, indicating its hormonal nature. In consistent with its broad expression, ghrelin does not only stimulate GH release, but also increases food intake, regulates energy homeostasis and decreases energy expenditure by lowering the catabolism of fat (Nakazato et al., 2001; Wu et al., 2004). Because of its orexigenic and adipogenic character, ghrelin may be potentially useful for the treatment of anorexia and cachexia (Nagaya et al., 2005). Ghrelin and the GHS-R were detected in immune cells and lymphoid tissues. Initially, ghrelin was reported as an immune enhancing factor. Koo et al. reported that ghrelin induced increase of levels of peripheral blood lymphocytes and thymic cellularity and differentiation (Koo et al., 2001). In addition, they showed ghrelin treatment reduced tumor initiation and subsequent metastases by the increase of the number of cytotoxic lymphocytes. More recently, however, the immunosuppressive effects of ghrelin have been reported. Anti-inflammatory functions of ghrelin were first reported against T cells and macrophages in vitro (Dixit et al., 2004). And then, the immunosuppressive activity of ghrelin in vivo was shown in several animal models including bowel disease (Gonzalez-Rey et al., 2006), arthritis (Chorny et al., 2008; Granado et al., 2005) and sepsis and endotoxemia (Chorny et al., 2008; Li et al., 2004; Wu et al., 2007). In these reports, the anti-inflammatory effect of ghrelin is focused on the suppression of pro-inflammatory cytokines such as IL-1, IL-6, and TNF- $\alpha$ . Li et al. demonstrated that ghrelin attenuated TNF- $\alpha$ -induced nuclear translocation of NF- $\kappa$ B, indicating that blockade for activation of the transcription factor NF- $\kappa$ B could be a potential mechanism whereby ghrelin modulates inflammatory responses (Li et al., 2004).

### **Ghrelin suppresses experimental autoimmune encephalomyelitis**

In consistent with these observations, we recently demonstrated that exogenous administration of ghrelin suppressed experimental autoimmune encephalomyelitis (EAE), an animal model of mul-

tipple sclerosis (Theil et al., 2009). Since ghrelin is the most potent NPY releasing hormone and NPY suppresses EAE by a Th2 bias, we examined whether ghrelin affects the Th1/Th2 balance in ways that are similar to NPY. It was also interesting to clarify whether its potential mechanism of EAE suppression is primarily mediated through immune cells or secondarily through NPY release. Both the proliferative response and the cytokine production including IFN- $\gamma$ , IL-17, and IL-4 by encephalitogenic antigen-primed T cells did not significantly alter between ghrelin- and sham-treated mice. Based upon these observations, we concluded that the suppression of EAE mediated by ghrelin is independent on the NPY effect. Furthermore, we revealed that transfer of T cells from ghrelin-treated mice to normal mice produced a similar disease compared to transfer of T cells from control mice, suggesting that other cells rather than T cells are main targets of ghrelin in EAE. Interestingly, the expression of pro-inflammatory cytokines was reduced in spinal cords, particularly in microglia, of ghrelin-treated mice. In fact, LPS-stimulated production of pro-inflammatory cytokines by microglia isolated from naive mice was reduced when cultured in the presence of ghrelin in vitro, indicating the direct effect of ghrelin on microglia. Although monocytes have been reported to be mainly affected by ghrelin treatment in animal models of sepsis or arthritis, microglia seem to be an important target in EAE. Microglia are derived from hematopoietic cells in central nervous system. Microglia is a double-edged sword which has both pro-inflammatory and anti-inflammatory functions (Block et al., 2007). Microglia play an important role in keeping homeostasis of CNS as a scavenger cells in normal conditions. However, microglia are important sources of inflammatory cytokines in pathogenic conditions such as inflammation. More recently, the involvement of pro-inflammatory cytokines has gained attention not only in inflammation but also in psychiatric disorders such as depression and neurodegenerative diseases including Parkinson's disease, amyotrophic lateral sclerosis (ALS) and Alzheimer's disease. Therefore, to understand the regulatory mechanisms of pro-inflammatory cytokine production in microglia is critical to control a variety of neurological and mental diseases.

### **Future directions**

The discrepancy between initial studies to show the immune-enhancing effect of ghrelin and recent studies to show anti-inflammatory function of ghrelin still remains to be explained. In both cases, however, the effects of ghrelin on the immune system seem to be beneficial to our body, as the initial study demonstrated anti-tumor effect and the later studies showed suppression of excessive immune reactions such as sepsis or autoimmune diseases including arthritis and encephalomyelitis. Therefore ghrelin might play a protective role for our body either by enhancing or inhibiting immunity depending on the specific situations.

Ghrelin is currently under the clinical trial for cardiac failure, chronic obstructive pulmonary disease, and anorexia nervosa. Proinflammatory cytokines are upregulated in patients with cognitive heart failure, particularly those with cardiac cachexia, and have been implicated in the pathophysiology of these diseases. For these patients, anti-inflammatory effect may be beneficial in addition of orexigenic effect. In addition, our findings to suppress microglial activation highlight the potential of ghrelin to treat disorders that proinflammatory cytokines contribute to the pathogenesis such as neuroinflammatory diseases and depression.

## References

- Bedoui S, Miyake S, Lin Y, Miyamoto K, Oki S, Kawamura N, Beck-Sickinger A, von Horsten S, Yamamura T. Neuropeptide Y (NPY) suppresses experimental autoimmune encephalomyelitis: NPY1 receptor-specific inhibition of autoreactive Th1 responses in vivo. *J Immunol* 171:3451-3458, 2003.
- Block ML, Zecca L, Hong J-S. Microglia-mediated neurotoxicity: uncovering the molecular mechanisms. *Nat Rev Neurosci* 8:57-69, 2007.
- Chorny A, Anderson P, Gonzalez-Rey E, Delgado M. Ghrelin protects against experimental sepsis by inhibiting high-mobility group box 1 release and by killing bacteria. *J Immunol* 180:8369-8377, 2008.
- Cowley MA, Smith RG, Diano S, Tschop M, Pronchuk N, Grove KL, Strasburger CJ, Bidlingmaier M, Esterman M, Heiman ML, Garcia-Segura LM, Nillni EA, Mendez P, Low MJ, Sotonyi P, Friedman JM, Liu H, Pinto S, Colmers WF, Cone RD, Horvath TL. The distribution and mechanism of action of ghrelin in the CNS demonstrates a novel hypothalamic circuit regulating energy homeostasis. *Neuron* 37:649-661, 2003.
- Dixit VD, Schaffer E M, Pyle R S, Collins GD, Sakthivel SK, Palaniappan R, Lillard Jr JW, Taub DD. Ghrelin inhibits leptin- and activation-induced proinflammatory cytokine expression by human monocytes and T cells. *J Clin Invest* 114:57-66, 2004.
- Gonzalez-Rey E., Chorny A, Delgado M. Therapeutic action of ghrelin in a mouse model of colitis. *Gastroenterology* 130:1707-1720, 2006.
- Granado M, Priego T, Martin AI, Villanua A, Lopez-Caldron A. Anti-inflammatory effect of the ghrelin agonist growth hormone-releasing peptide-2 (GHRP-2) in arthritic rats. *Am J Physiol Endocrinol Metab* 288:E486-E492, 2005.
- Kalra SP, Ueno N, Kalra PS. Stimulation of appetite by ghrelin is regulated by leptin restraint: peripheral and central sites of action. *J Nutr* 135:1331-1335, 2005.
- Kojima M, Hosoda H, Date Y, Nakazato M, Matsuo H, Kangawa K. Ghrelin is a growth-hormone-releasing acylated peptide from stomach. *Nature* 402:656-660, 1999.
- Koo GC, Huang C, Camacho R, Trainor C, Blake JT, Sirotina-Meisher A, Schleim KD, Wu T-J, Cheng K, Nargund R, Mckissick G. Immune enhancing effect of a growth hormone secretagogue. *J Immunol* 16:4195-4201, 2001.
- Li WG, Gavrilu D, Liu X, Wang L, Gunnlaugsson S, Stoll LL, McCormick ML, Sigmund CD, Tang C, Weintraub NL. Ghrelin inhibits proinflammatory responses and nuclear factor- $\kappa$ B activation in human endothelial cells. *Circulation* 109:2221-2226, 2004.
- Lord GM, Matarese G, Howard JK, Baker RJ, Bloon SR, Lechler RI. Leptin modulates the T-cell immune response and reverses starvation-induced immunosuppression. *Nature* 394:897-901, 1998.
- Nagaya N, Itoh T, Murakami S, Oya H, Uemasu M, Miyatake K, Kangawa K. Treatment of cachexia

- with ghrelin in patients with COPD. *Chest* 128:1187-1193, 2005.
- Nakazato M, Murakami N, Date Y, Kojima M, Matsuo H, Kangawa K, Matsukura S. A role for ghrelin in the central regulation of feeding. *Nature* 409:194-198, 2001.
- Sanna V, Di Giacomo A, La Cava A, Lechler I, Fontana S, Zappacosta S, Matarese G. Leptin surge precedes onset of autoimmune encephalomyelitis and correlates with development of pathogenic T cell responses. *J Clin Invest* 111:241-250, 2003.
- Theil M-M, Miyake S, Mizuno M, Tomi C, Croxford JL, Hosoda H, Theil J, von Horsten S, Yokote H, Chiba A, Lin Y, Oki S, Akamizu T, Kangawa K, Yammaura T. Suppression of experimental autoimmune encephalomyelitis by ghrelin. *J Immunol* 183:2859-2866, 2009.
- Wassem T, Duxbury M, Ito H, Ashley SW, Robinson MK. Exogenous ghrelin modulates release of pro- and anti-inflammatory cytokines in LPS-stimulated macrophages through distinct signaling pathways. *Surgery* 143:334-342, 2008.
- Wheway J, Mackay CR, Newton RA, Sainsbury A, Boey D, Herzog H, Mackay F. A fundamental bimodal role for neuropeptide Y1 receptor in the immune system. *J Exp Med* 202(11):1527-1538, 2005.
- Wu JT, Kral JG. Ghrelin: integrative neuroendocrine peptide in health and disease. *Annu Surg* 239:464-474, 2004.
- Wu R, Dong W, Cui X, Zhou M, Simms HH, Ravikumar TS, Wang P. Ghrelin down-regulates proinflammatory cytokines in sepsis through activation of the vagus nerve. *Ann Surg* 245:480-486, 2007.

# Synthetic Retinoid AM80 Inhibits Th17 Cells and Ameliorates Experimental Autoimmune Encephalomyelitis

Christian Klemann,\* Benjamin J.E. Raveney,\*  
Anna K. Klemann,\* Tomoko Ozawa,\*  
Stephan von Hörsten,<sup>†</sup> Koichi Shudo,<sup>‡</sup> Shinji Oki,\*  
and Takashi Yamamura\*

From the Department of Immunology,\* National Institute of Neuroscience, NCNP, Tokyo, Japan; the Department of Experimental Therapy,<sup>†</sup> Franz-Penzoldt Center, Erlangen, Germany; and the Research Foundation IISUU Laboratory,<sup>‡</sup> Tokyo, Japan

**Recent evidence suggests that interleukin-17-producing CD4<sup>+</sup> T cells (Th17 cells) are the dominant pathogenic cellular component in autoimmune inflammatory diseases, including multiple sclerosis. It has recently been demonstrated that *all-trans* retinoic acid can suppress Th17 differentiation and promote the generation of Foxp3<sup>+</sup> regulatory T cells via retinoic acid receptor signals. Here, we investigated the effects of AM80, a synthetic retinoid with enhanced biological properties to *all-trans* retinoic acid, on Th17 differentiation and function and evaluated its therapeutic potential in experimental autoimmune encephalomyelitis (EAE), an animal model of multiple sclerosis. AM80 treatment was more effective than *all-trans* retinoic acid in inhibiting Th17 differentiation *in vitro*. Oral administration of AM80 was protective for the early development of EAE and the down-modulation of Th17 differentiation and effector functions *in vivo*. Moreover, AM80 inhibited interleukin-17 production by splenic memory T cells, *in vitro*-differentiated Th17 cells, and central nervous system-infiltrating effector T cells. Accordingly, AM80 was effective when administered therapeutically after the onset of EAE. Continuous AM80 treatment, however, was ineffective at inhibiting late EAE symptoms despite the maintained suppression of ROR $\gamma$ t and interleukin-17 expression levels by central nervous system-infiltrating T cells. We reveal that continuous AM80 treatment also led to the suppression of interleukin-10 production by a distinct T cell subset that expressed both Foxp3 and ROR $\gamma$ t. These findings sug-**

**gest that retinoid signaling regulates both inflammatory Th17 cells and Th17-like regulatory cells. (Am J Pathol 2009, 174:2234–2245; DOI: 10.2353/ajpath.2009.081084)**

Multiple sclerosis (MS) is a chronic inflammatory autoimmune disease affecting the central nervous system (CNS).<sup>1</sup> Previous studies of experimental autoimmune encephalomyelitis (EAE), a murine model of MS,<sup>2,3</sup> indicated that autoimmune responses were initiated by a subset of myelin-specific CD4<sup>+</sup> T cells secreting the inflammatory cytokine interferon (IFN)- $\gamma$ , termed Th1 cells.<sup>4,5</sup> More recently, the identification of an additional subset of differentiated inflammatory helper T cells secreting large amount of the cytokine interleukin (IL)-17 (Th17 cells) have allowed new insights into the pathology of a range of inflammatory autoimmune diseases.<sup>6,7</sup> Indeed, the presence of such Th17 cells among CNS-infiltrating leukocytes has been demonstrated in EAE animals.<sup>7</sup> Furthermore, induction of EAE in IL-17-deficient mice leads to less severe disease<sup>8</sup> and mice that lack IL-23, a cytokine required for Th17 cell survival,<sup>9</sup> are resistant to EAE.<sup>6</sup> Critically, increased levels of IL-17 are detected in MS patients<sup>10</sup> and the presence of IL-17-secreting T cells has been shown to link with acute CNS lesions in MS.<sup>11</sup>

Differentiation of naïve T cells into Th17 cells *in vitro* requires culture with a combination of IL-6, an inflammatory cytokine elaborated by innate immune cells subsequent to ligation of pathogen-associated molecular pattern receptors, and transforming growth factor (TGF)- $\beta$ ,<sup>12–14</sup> a cyto-

Supported in part by Health and Labor Sciences Research Grants on Intractable Diseases from the Ministry of Health, Labor and Welfare of Japan.

Accepted for publication March 5, 2009.

Supplemental material for this article can be found on <http://ajp.amjpathol.org>.

Address reprint requests to Takashi Yamamura, M.D., Ph.D., Director, Department of Immunology, or Shinji Oki, Ph.D., Section Chief, Department of Immunology, National Institute of Neuroscience, NCNP, 4-1-1 Ogawahigashi, Kodaira, Tokyo, 187-8502, Japan. E-mail: [yamamura@ncnp.go.jp](mailto:yamamura@ncnp.go.jp) and [soki@ncnp.go.jp](mailto:soki@ncnp.go.jp).

kine classically regarded as anti-inflammatory and also associated with the differentiation of regulatory T cells.<sup>15</sup> The requirement for IL-6 and TGF- $\beta$  in Th17 differentiation has also been demonstrated *in vivo*<sup>12,16</sup> Phenotypically, Th17 cells express the retinoic acid (RA)-related orphan receptor  $\gamma$ t (ROR $\gamma$ t) in a Stat3-dependent manner, and produce high levels of many inflammatory cytokines, including IL-17, IL-6, tumor necrosis factor (TNF)- $\alpha$ , IL-21, and IL-22.<sup>7,17–19</sup> Interestingly, naïve T cells receiving TGF- $\beta$  signaling alone are induced to become a CD4<sup>+</sup>CD25<sup>+</sup> regulatory T cell population (Treg cells).<sup>13</sup> Treg cells are capable of suppressing inflammation, mediating self-tolerance, and produce suppressive cytokines such as TGF- $\beta$  and/or IL-10.<sup>20</sup> The generation of Treg cells requires the forkhead/winged-helix transcription factor Foxp3 and its functional impairment leads to autoimmunity.<sup>21,22</sup> Foxp3<sup>+</sup> Treg cells in the CNS have been shown to ameliorate EAE via the production of IL-10,<sup>23</sup> which has recently been associated with restraining Th17-mediated pathology in EAE.<sup>24</sup> Furthermore, IL-10-producing ROR $\gamma$ t<sup>+</sup>Foxp3<sup>+</sup> T cells have been identified *in vivo*, suggesting the existence of a regulatory Th17-like cell type.<sup>25</sup>

RA, the active metabolite of vitamin A, has multiple effects on cell differentiation and survival through ligation to the two families receptors: retinoic acid receptors (RAR) and retinoid X receptor (RXR), each of which has multiple isoforms.<sup>26</sup> Recently, *all-trans* retinoic acid (ATRA) has been reported to suppress the differentiation of Th17 cells through ligation to the RAR- $\alpha$ ,<sup>27,28</sup> accompanied by a down-regulation of ROR $\gamma$ t and reciprocal induction of Treg cells expressing Foxp3.<sup>27,29</sup> Possible mechanisms of action of ATRA for the suppression of Th17 cell function have been reported to be a result of reduced expression of IL-6 receptor and IL-23 receptor as well as enhanced TGF- $\beta$  signaling in a Smad3-dependent manner.<sup>30</sup>

RA has been shown to ameliorate EAE.<sup>31,32</sup> But as these studies were carried through before the discovery of Th17 cells, the amelioration was attributed to suppression of Th1 cells. In addition, the therapeutic application of RA to date has been limited by instability and poor bioavailability of this compound as well as by its non-selective binding to a broad range of retinoid receptors, which conceivably leads to unexpected side effects.<sup>26,33,34</sup> To circumvent these potential problems in the clinical use of RAR agonists, a variety of synthetic RAR agonists with improved biological properties *in vivo* have been developed. One of these synthetic retinoids, AM80, is already available as medication under the trade name Tamibarotene for human diseases such as acute promyelocytic leukemia (APL) and psoriasis.<sup>35–37</sup> AM80 is specific for the RAR $\alpha/\beta$  and characterized by a higher stability, fewer potential side effects, and superior bioavailability compared with ATRA.<sup>35–38</sup> Therefore, we may open up new therapeutic avenues for treating Th17-mediated autoimmune diseases by testing AM80 in an immunoregulatory context.

In this study, we demonstrate for the first time that AM80 inhibits Th17 differentiation *in vitro* with a higher potency than ATRA. Treatment with AM80 ameliorates EAE and inhibits both the differentiation of Th17 cells and

the effector function of Th17 cells *in vivo* without generating general immunosuppression. In addition, AM80 proved to be effective in rescue from acute EAE when administered after the onset of the disease. Interestingly, continuous AM80 treatment failed to improve chronic disease despite of apparent suppression of T cell expression of IL-17 and ROR $\gamma$ t. We demonstrate that continuous AM80 treatment results in the suppression of IL-10 production by a unique subset of T cells, which is identified as T cells that co-expresses ROR $\gamma$ t and Foxp3. We conclude that treatment with the synthetic retinoid AM80 is a considerable intervention strategy for the acute phase of Th17-mediated autoimmune diseases such as MS.

## Materials and Methods

### Animals and EAE Induction

C57BL/6J (B6) mice (CLEA Laboratory Animal Corp., Tokyo, Japan) were maintained in specific pathogen-free conditions in accordance with institutional guidelines (National Institute of Neuroscience, NCNP, Tokyo, Japan). This study used female mice at 8 to 10 weeks of age. For EAE induction mice were injected subcutaneously with 100  $\mu$ g of myelin oligodendrocyte glycoprotein (MOG) amino acids 35–55 (MOG<sub>35–55</sub> peptide MEVGWYRSPFSRVVHLYRNGK)<sup>39</sup> and 1 mg of heat-killed *Mycobacterium tuberculosis* H37RA emulsified in complete Freund's adjuvant (Difco, Lawrence, KS). 200 ng of pertussis toxin (List Biological Laboratories) was injected intraperitoneally on days 0 and 2 after immunization. Some groups of mice also received 3 mg/kg AM80 in 0.5% carboxymethylcellulose (CMC) solution (WAKO Chemicals, Osaka, Japan) by oral gavage.

EAE was clinically scored daily (0, no clinical signs; 1, partial tail paralysis; 2, flaccid tail; 3, partial hindlimb paralysis; 4, total hindlimb paralysis; 5, Hind- and foreleg paralysis).<sup>39</sup> Disease was also assessed using histological examination of CNS tissue as previously described.<sup>40</sup> Briefly, animals were perfused with 20 ml of cold phosphate-buffered saline, and CNS tissue was excised and fixed in formal saline. Paraffin-embedded sections were prepared and stained with either Luxol fast blue or H&E and photomicrographs acquired with a light microscope (Eclipse E800M, Nikon, Japan).

### Cell Isolation and Purification

CNS-infiltrating lymphocytes were isolated from spinal cords and brains as previously described.<sup>39</sup> Briefly, tissue was cut into small pieces and digested for 40 minutes at 37°C in media (GIBCO, Auckland, New Zealand) supplemented with 1.4 mg/ml collagenase H (Roche, Mannheim, Germany) and 100  $\mu$ g/ml DNase I (Roche). Resulting tissue homogenates were forced through a 70- $\mu$ m cell strainer and leukocytes were enriched using a discontinuous 40%/80% Percoll density gradient centrifugation. Leukocytes were collected from the interface and where indicated cell numbers for leukocytes and/or T cell subsets per mouse were counted with an improved



Neubauer counting chamber and via flow cytometry with reference to a cell number curve as previously described.<sup>41</sup> T cells were purified from splenocytes, draining lymph nodes and CNS infiltrates using a pan T cell MACS isolation kit with an AutoMACS separator according to manufacturer's instructions (Miltenyi Biotech, Bergisch Gladbach, Germany). Where required, naïve CD4<sup>+</sup>CD44<sup>-</sup>CD25<sup>-</sup>CD62L<sup>high</sup> T cells or memory CD4<sup>+</sup>CD44<sup>+</sup>CD25<sup>-</sup>CD62L<sup>low</sup> T cells were further sorted using a fluorescence-activated cell sorter ARIA (BD Cytometry System, Franklin Lakes, NJ).

### Cell Culture

RPMI 1640 medium (GIBCO) supplemented with 10% fetal calf serum, 2 mmol/L L-glutamine, 100 U/ml penicillin-streptomycin, and 50  $\mu$ mol/L 2-mercaptoethanol (Invitrogen, Carlsbad, CA) was used for all cultures. Cells were activated with immobilized anti-CD3 monoclonal antibody (mAb) (2C-11; 2  $\mu$ g/ml) and soluble anti-CD28 mAb (BD PharMingen) or, when measuring recall responses of secondary lymphoid tissue 10 days after immunization, with MOG<sub>35-55</sub> peptide (0–100  $\mu$ mol/L). Where indicated, cells were cultured under Th17 polarizing conditions: 2 ng/ml TGF- $\beta$ , 20 ng/ml IL-6 (Pepro-Tech, London, UK), anti-IFN- $\gamma$  mAb (HB170; 10  $\mu$ g/ml), and anti-IL-4 mAb (HB188; 10  $\mu$ g/ml). Into some cultures ATRA (Sigma-Aldrich, Steinheim, Germany) or AM80 dissolved in dimethyl sulfoxide (Sigma) was added at the required concentrations. Supernatants were harvested at 72 or 96 hours for cytokine measurement and proliferation was determined by incubating with [<sup>3</sup>H]thymidine (1  $\mu$ Ci/well) for the final 12 hours of culture assessing incorporation of radioactivity with a beta 1205 counter (Pharmacia Biotech, Freiburg, Germany). In some experiments, to measure differentiation of T cells, after 96 hours of activation cells were rested for 48 hours before restimulation with anti-CD3 for a further 96 hours.

### Cytokine Measurement

IL-17 was assessed via a mouse IL-17 DuoSet (R&D Systems). All other cytokines were assessed by cytometric bead array using a mouse inflammation kit or a mouse Th1/Th2 cytokine kit (BD Biosciences).

### Intracellular Staining for Cytokines and Transcription Factors

To stain cells for production of cytokines, cells were restimulated with 50 ng/ml phorbol 12-myristate 13-acetate and 500 ng/ml ionomycin (Sigma-Aldrich) for 4 to 6 hours before surface staining in the presence of monensin-containing GolgiPlug (BD Biosciences). A LIVE/DEAD fixable dead cell stain kit (Invitrogen) was used to exclude dead cells. Intracellular staining was then performed using a Cytotfix/cytoperm kit (BD PharMingen), according to manufacturer's instructions. To stain intracellular transcription factors, we used an anti-mouse/rat

Foxp3 staining set (eBioscience, San Diego, CA) with anti-ROR $\gamma$ t antibody (BioLegend, San Diego, CA) and visualized with a secondary PE-conjugated goat anti-rabbit antibody (Invitrogen).

### RNA Extraction and Quantitative Reverse Transcription-Polymerase Chain Reaction (RT-PCR)

Total RNA was extracted from cell populations using an RNeasy Mini Kit (Qiagen, Maryland) according to the manufacturer's instructions. cDNA was prepared using a first-strand cDNA Kit (Takara, Otsu, Shiga, Japan). Quantitative real-time PCR was performed using a LightCycler-FastStart DNA Master SYBR Green I kit (Roche Diagnostics) with a LightCycler real-time PCR machine (Roche). Gene expression values were normalized to the expression of the GAPDH housekeeping gene. Primers used in this study were GAPDH fw: 5'-AACGACCCCTTCATTGAC-3' rv: 5'-TCCACATACTCAGCAC-3', RORc fw: 5'-TGTCCCTGGGCTACCCTACTG-3' rv: 5'-GTGCAGGAGTAGGCCACATT-3', mFOXP3 fw: 5'-TTCTCACAACAAGGCCACTTG-3' rv: 5'-CCCAGGAAAGACAGCAACCCT-3', mT-bet fw: 5'-GC-CAGGGAACCGCTTATATG-3' rv: 5'-GACGATCATCTGGGTACATTGT-3', mIL-10 fw: 5'-CATGGGTCTTGGGAGAGAA-3' rv: 5'-CATTCCCAGAGGAATTGCAT-3'.


### Statistics

EAE clinical scores for groups of mice are presented as the mean group clinical score  $\pm$  SEM, and statistical differences were analyzed by two-way analysis of variance (analysis of variance) for repeated measures and significance calculated with a Bonferroni post-test. Cytokine secretion data were analyzed with a two-tailed Student's *t*-test or with one-way analysis of variance with Bonferroni's multiple comparison test. Unless otherwise stated, *P* < 0.05 was considered significant and indicated on plots by asterisks.

## Results

### RAR Agonists AM80 Inhibit Th17 Cell Differentiation in Vitro

The synthetic retinoid AM80 is already available as medication for human diseases such as APL and psoriasis and characterized by superior pharmacological properties compared with ATRA. As previous studies demonstrated that Th17 cells are the dominant pathogenic cellular component in autoimmune inflammatory diseases, such as MS, and ATRA modulates Th17 differentiation,<sup>27-29</sup> we examined whether AM80 could be used to treat autoimmune diseases. To this end, whole splenocytes or purified naïve T cells were stimulated under a range of Th17-inducing conditions (ie, with either TGF- $\beta$  plus IL-6, or IL-23) with or without either AM80 or ATRA and were assessed for their IL-17 production. Addition of retinoids to the splenocyte culture significantly reduced




**Figure 1.** AM80 is a potent inhibitor for Th17 cell differentiation *in vitro*. Whole splenocytes were stimulated with soluble anti-CD3 with retinoids added at a range of concentrations. **A:** Cells were cultured in the presence of IL-23 (broken line), TGF-β and IL-6 (solid line), or without the addition of cytokines (closed triangle) for 3 days and IL-17 production assessed by enzyme-linked immunosorbent assay (ELISA). **B:** Intracellular cytokine staining of IL-17 and IFN-γ among Tcr<sup>+</sup>CD4<sup>+</sup> cells with or without 10 nmol/L retinoid treatment assessed after 3 days of culture. Data depicted in **A** and **B** are representative of three independent experiments. In **C**, CD4<sup>+</sup>CD44<sup>-</sup>CD25<sup>-</sup>CD62L<sup>high</sup> naïve T cells were stimulated under Th17-priming conditions with retinoids at a range of concentrations for 96 hours. Cells were rested for 48 hours before restimulation and IL-17 production was measured in culture supernatants after further 96 hours of stimulation. \**P* < 0.001. RNA from these cells was analyzed by quantitative RT-PCR for the transcription factors RORγt and Foxp3 (**D**). Data depicted in **C** and **D** are representative of four similar experiments.

the amount of IL-17 in supernatants (Figure 1A). Accumulation of IL-17-producing T cells after restimulation of the culture was also suppressed by AM80 or ATRA with minimal effect on the development of IFN-γ-producing T cells (Figure 1B). Interestingly, AM80 appeared to be more effective than ATRA at inhibiting IL-17 production especially at low doses. Furthermore, both AM80 and ATRA inhibited Th17 cell differentiation of naïve T cells, as revealed by the reductions in IL-17 secretion (Figure 1C). Importantly, the effect of retinoid treatment on naïve T cell differentiation is not merely due to a suppression of T cell activation or an increase in cell death, as such treatment did not reduce proliferation, or total live cell number in the cultures (data not shown). Also, treatment with AM80 or ATRA led to reduced expression of RORγt, a key Th17 cell-specific transcription factor, as compared with untreated controls (Figure 1D). Such reductions in Th17 phenotype were accompanied by the reciprocal increase of Foxp3 expression, indicative of a regulatory T cell phenotype (Figure 1D). AM80 was also more effective at modulating the transcription factors specific for Th17 cells as compared with ATRA (Figure 1D). In addition, no up-regulation in T-bet and GATA-3 was observed, indicating that the inhibition of Th17 differentiation by retinoid treatment was not a result of an altered Th1/Th2 phenotype. These results suggest that AM80 is a superior modulator of *in vitro* Th17 differentiation as compared with ATRA.

### AM80 Treatment Ameliorates Acute Autoimmune Inflammation

EAE, the murine model of MS,<sup>3</sup> is an autoimmune disease in which Th17 cells play an important pathogenic role.<sup>42</sup> At high doses ATRA can delay onset of this disease putatively via mechanisms that affect the development of Th17 cell function.<sup>30</sup> As AM80 is more effective in inhibiting Th17 development *in vitro*, we tested whether or not administration of this compound could modulate EAE. Thus, EAE was induced in B6 mice, and some groups of mice received AM80 orally every other day from the day of immunization. The onset of clinical disease was delayed and maximal clinical score was significantly reduced in animals treated with AM80 as compared with control mice treated with vehicle alone (Figure 2A). Histological examination of spinal cords at the peak of clinical disease (day 15) showed that AM80 treatment led to a marked reduction in cellular infiltrates into spinal cord and maintained normal myelin structure (Figure 2B). Flow cytometric analysis confirmed that AM80 treatment led to reduced numbers of infiltrating cells in the brains and spinal cords and this difference was particularly apparent when T cell numbers were compared (Figure 2C). As some T cell infiltration was still observed in the CNS of AM80-treated mice, we examined the functional properties of such cells to ascertain whether or not AM80 modulated T cell differentiation *in vivo*. The expression of




**Figure 2.** AM80 treatment ameliorates EAE with reduction of IL-17 production *in vivo*. EAE was induced in B6 mice by immunization with MOG<sub>35-55</sub>. Groups of mice received either CMC or 3 mg/kg AM80 in CMC orally every other day from day 0. Mice were scored daily for clinical disease (A) analysis of variance for repeated measures shows significant differences from day 11 to 18 (\**P* < 0.001). On day 15 after immunization, groups of mice were sacrificed and spinal cords were examined histologically. Representative sections are shown in B. H&E staining (upper panels), Luxol fast blue staining (lower panels). Scale bar = 200 μm. Leukocytes and T cells were purified from the CNS at day 15 after immunization and cell numbers evaluated by hemocytometer (C). Quantitative RT-PCR was used to measure an array of transcription factors and cytokines (D, upper row) in which error bars represent duplicated PCR of the same samples. In the lower row, CNS-infiltrating T cells were restimulated with immobilized anti-CD3 antibody and supernatants were measured after 72 hours for the presence of a range of cytokines and chemokines using ELISA or CBA. Error bars represent measurements from duplicate wells. Data are representative of at least two independent experiments.

Th17-specific genes including RORγt, IL-17 and IL-23 receptor in T cells isolated from CNS was reduced in the group treated with AM80 (Figure 2D). In contrast, the expression of Foxp3 and T-bet (specific for Treg cells and Th1 cells respectively) were not elevated by AM80 treatment. Flow cytometric analysis confirmed that fewer IL-17-producing cells but similar numbers of Foxp3-positive cells were present among the CNS infiltrating T cells in AM80-treated mice as compared with vehicle treated controls (data not shown). Also, we note that AM80 treatment did not affect the expression of activation markers (including CD62L, CD44 and CD25) by CNS-infiltrating T cells (data not shown). Furthermore, on anti-CD3 mAb stimulation, CNS-infiltrating T cells isolated from animals

treated with AM80 produced significantly reduced levels of pro-inflammatory cytokines and chemokines (Figure 2D). Myeloid cell and T cell phenotyping via flow cytometric analysis revealed no significant differences between treatment groups (data not shown). Taken together, these results indicate that AM80 treatment decreases the number of T cells infiltrating the CNS during EAE and also lowers their IL-17 producing capacity.

We have performed adoptive transfer experiments to determine whether AM80 ameliorates EAE through a direct effect on T cells or not. Draining lymph node cells isolated from immunized mice with oral administration of AM80 or vehicle were restimulated with MOG peptide and cultured for one week without retinoid treatment.



**Figure 3.** RAR agonist AM80 suppresses Th17 differentiation without inhibiting typical proliferative responses *in vivo*. B6 mice were immunized with MOG<sub>35-55</sub> in CFA and vehicle (CMC) or AM80 (3 mg/kg in CMC) were administered orally every other day starting from day 0 until day 8. Draining lymph node cells were isolated at day 10 and restimulated with MOG peptide at various concentrations. After 72 hours, antigen-specific IL-17 production was assessed by ELISA (A). In B, TNF- $\alpha$  and IFN- $\gamma$  production after restimulation with 100  $\mu$ mol/L MOG were assessed by CBA. C: Intracellular cytokine staining of draining lymph node cells in the presence of 10  $\mu$ mol/L MOG<sub>35-55</sub> after 96 hours of culture shows reduced percentages of IL-17<sup>+</sup> and IL-17<sup>+</sup>/IFN- $\gamma$ <sup>+</sup> double producing cells. Plots are gated on TCR<sup>+</sup>CD4<sup>+</sup> lymphocytes. D: Expression of ROR $\gamma$ t in T cells obtained from C was assessed by quantitative RT-PCR. E: Cell proliferation was assessed either by [<sup>3</sup>H]thymidine incorporation or cell number evaluation by fluorescence-activated cell sorting. Data in A, B, and E are representative of four independent experiments with three to five mice per group and C and D depict results from two experiments with three mice per group.


After transfer of MOG-reactive T cell blasts into naive mice, AM80-treated T cells caused only minimal disease compared with a significant disease development after transfer of vehicle-treated T cells (see Supplemental Figure S1 at <http://ajp.amjpathol.org>). In addition, when encephalitogenic T cells isolated from vehicle-treated animals were incubated *ex vivo* with AM80 for 2 hours before adoptive transfer into naive mice, no disease developed (see Supplemental Figure S1 at <http://ajp.amjpathol.org>). Taken together, the suppressive capacity of AM80 is attributed, at least in part, to a direct inhibitory effect on encephalitogenic T cells *in vivo*.

### AM80 Suppresses Antigen-Specific IL-17 Production of T Cells

Since AM80 treatment suppressed the onset of clinical EAE and also inhibited inflammatory cellular responses in the target organ, we set out to elucidate the cellular and molecular mechanism by which AM80 suppresses EAE development by examining antigen-specific effector T cells responses with or without AM80 treatment. MOG-specific production of pro-inflammatory IL-17 by draining lymph node cells was dramatically reduced after *in vivo* treatment with AM80 (Figure 3A). In addition, production of other proinflammatory cytokines such as IFN- $\gamma$  and TNF- $\alpha$  was also significantly reduced (Figure 3B). The

reduction in IL-17 secretion following retinoid treatment could result from decreased frequency of IL-17 producing T cells, or decreased production of cytokines by each T cell. Therefore, we examined cytokine production among draining lymph node T cells on a per cell basis using flow cytometric intracellular cytokine staining. Also, Th17 differentiation was estimated by quantifying ROR $\gamma$ t expression among draining lymph node, using quantitative RT-PCR. As shown in Figure 3C, the IL-17-positive population in TCR<sup>+</sup>CD4<sup>+</sup> subset was reduced after treatment with AM80, indicating that the treatment resulted in a lower frequency of IL-17 producing T cells developing in draining lymph node. In line with these findings, isolated T cells from animals that had received AM80 treatment expressed lower levels of ROR $\gamma$ t (Figure 3D). Interestingly, unlike the effect of AM80 *in vitro*, such reductions in ROR $\gamma$ t expression were not associated with an increase in Foxp3 expression (data not shown).

It is conceivable that AM80 may protect from EAE by generating systemic immunosuppression. Indeed, a previous study using relatively high doses of the broad spectrum RAR agonist ATRA to treat EAE was unable to rule out this mechanism of action as administration of ATRA resulted in a suppression of peripheral proliferative T cell responses.<sup>30</sup> With our treatment approach, however, cellular proliferation as measured by [<sup>3</sup>H]thymidine incorporation and increase in cell numbers determined by fluo-



**Figure 4.** AM80 suppresses IL-17 production by differentiated Th17 cells and ameliorates EAE in therapeutic intervention. **A:** CD4<sup>+</sup>CD44<sup>+</sup>CD25<sup>+</sup>CD62L<sup>low</sup> memory T cells were stimulated by plate-bound anti-CD3 antibody for 4 days in the presence of IL-23. Dose-dependent decrease of IL-17 production by RAR agonists was shown in left panel. Intracellular cytokine staining of TCR<sup>+</sup>CD4<sup>+</sup> cells shows decreased percentage of IL-17<sup>+</sup> memory T cells after treatment with 100 nmol/L AM80 (**B**). Graphs are representative of two independent experiments. **C:** CNS-infiltrating T cells from mice with severe EAE (score 3–4) were isolated and restimulated with immobilized anti-CD3 in the presence of RAR agonists (100 nmol/L) for 3 days. Supernatants were analyzed for IL-17 production by ELISA and cells were subjected to quantitative RT-PCR for RORγt expression. Data are a representative of four similar experiments. **D:** Clinical EAE scores of animals treated daily from day 13 on (arrow) either with vehicle (CMC) or AM80. Displayed scores are representative of two experiments with *n* = 6.

rescence-activated cell sorting following re-stimulation with antigen did not differ between treated and control groups (Figure 3E). Collectively, these data suggest that rather acting as a systemic immunosuppressive agent, AM80 acts as an immunomodulatory agent for antigen-specific T cell responses, primarily affecting Th17 cell function *in vivo*.


#### AM80 Treatment Ameliorates Ongoing Inflammatory Responses by Suppressing IL-17 Production from Differentiated Th17 Effector Cells

As we wished to evaluate the therapeutic potential of AM80 for MS treatment, we tested whether or not AM80 had an inhibitory effect on Th17 effector function in addition to its effect on Th17 differentiation. To this end, we stimulated memory cells in the presence of IL-23, a cytokine that promotes Th17 memory cell function and survival,<sup>9</sup> and in the presence of increasing doses of AM80 or ATRA. AM80 and ATRA suppressed IL-17 production by memory T cells responding to anti-CD3 mAb activation (Figure 4A). Intracellular cytokine staining also demonstrated that those retinoids inhibit IL-17 production by differentiated Th17 cells (AM80, Figure 4B; ATRA, data not shown). Furthermore, we also confirmed that IL-17 production by T cells differentiated *in vitro* by a combination of TGF-β and IL-6, was effectively inhibited in the presence of AM80 after restimulation with immobilized anti-CD3 mAb (data not shown). As Th17 cells are shown to have an unstable phenotype when differentiated *ex vivo*,<sup>43</sup> we examined how retinoids affect the function of Th17 cells that have stably differentiated *in vivo*. We stimulated CNS-infiltrating T cells isolated from mice at peak

EAE, which consist of a high proportion of Th17 cells,<sup>42</sup> in the presence of AM80 or ATRA. Both RAR agonists successfully suppressed IL-17 production by those CNS infiltrating T cells concomitant with significant reduction of RORγt expression (Figure 4C). We further tested whether or not this effect was sufficient to modulate established EAE. Rescue treatment with AM80 starting after the onset of disease scores significantly suppressed the increase of disease scores that was observed in control vehicle-treated EAE mice (Figure 4D). The maximal disease score was reduced from 3.4 ± 0.39 in untreated EAE mice to 2.58 ± 0.47 in AM80-treated animals. Taken together, these data indicate the retinoid treatment is effective at inhibiting the function of activated Th17 cells.

#### Continuous AM80 Treatment Alters Cytokine Profile in the Chronic Phase of Disease


Our data have demonstrated that treatment with AM80 can both protect and rescue from EAE, associated with a suppression of pathogenic Th17 cell differentiation and function. However, when observed at later time points after EAE induction, the disease scores of both groups became alike (Figures 2A and 4D). Accordingly at such time points, there is no clear differences between treatment groups in terms of cellular infiltrates in CNS tissue as observed by histology (Figure 5A) or flow cytometry (Figure 5B). We then investigated the effector properties of CNS-infiltrating T cells derived from either vehicle-treated or AM80-treated animals with equivalent disease scores at day 25 after EAE induction. CNS-infiltrating T cells derived from AM80-treated animals contained strongly decreased levels of mRNA transcripts for RORγt, IL-23 receptor and IL-17 (Figure 5C). In addition, CNS-



**Figure 5.** Continuous AM80 treatment is less effective on EAE suppression with a differentially modulated cytokine profile. **A:** The H&E-stained sections and the Luxol fast blue-stained sections in EAE mice treated with or without AM80 are shown. The control animal (score 2) and AM80-treated animal (score 2) at day 25 were subjected to histological examination. Scale bar = 200  $\mu$ m. **B:** Total leukocyte numbers and isolated T cell numbers from spinal cords were evaluated in animals that had received CMC or AM80 at day 25 after immunization. The **upper row** of **C** depicts purified T cells from **B** that were subjected to quantitative RT-PCR. Error bars represent duplicated PCR of the same samples. In the **lower row** of **C**, CNS-infiltrating T cells were restimulated with immobilized anti-CD3 mAb and cytokine or chemokine production in culture supernatants were assessed by ELISA or CBA after 72 hours. Error bars represent measurements from duplicate wells (\* $P < 0.001$ ). **D:** IL-10 production by stimulated CNS-infiltrating T cells derived from CMC-treated or AM80-treated animals were assessed by either quantitative RT-PCR or CBA. CNS-infiltrating T cells were isolated at day 15 or day 25 after EAE induction. **B** is a representative of four independent experiments; **A**, **C**, and **D** of two with six animals per group.

infiltrating T cells of AM80-treated animals produce significantly lower amounts of proinflammatory cytokines (IL-17, IFN- $\gamma$ , and IL-6) and chemokines (CCL2) on restimulation *in vitro* (Figure 5C). Furthermore, although we have observed disease development until day 45, there were no further differences between treatment groups over this time period (data not shown). Collectively, these data suggest that inhibition of Th17 cell function alone is not sufficient to protect mice from later

onset of the disease. Foxp3 expression in CNS infiltrating T by quantitative RT-PCR (Figure 5C) or intracellular Foxp3 staining (data not shown) showed no apparent differences between treatment groups, albeit a trend toward lower levels of Foxp3 expression in brain infiltrating T cells of AM80 treated animals could be observed. Next, we investigated several regulatory cytokines and found that levels of IL-10 were reduced in CNS-infiltrating T cell of animals that had received AM80 (Figure 5D). Interest-



**Figure 6.** Continuous AM80 treatment impairs IL-10 production by T cells, which are identified as RORγt<sup>+</sup>/FOXP3<sup>+</sup> population. **A:** Naive T cells were cultured under neutral (no priming) or Th17-priming conditions in the presence of retinoids (100 nmol/L). After 96 hours, expression of IL-10 mRNA was assessed by quantitative RT-PCR. For further analysis, cells were rested for 48

ingly, we observed a more profound inhibitory effect of AM80 on IL-10 production by CNS-infiltrating T cells at the later time point after EAE induction (day 25) compared with those in earlier phases of the disease (day 15). Despite simultaneous suppression of IL-17 production, continuous AM80 treatment may deplete the host immune system of its intrinsic T cell production of IL-10, leading to a possible loss of the protective function of retinoids.

### Continuous Treatment with RAR Agonists Abrogate IL-10 Production by T Cells, Which Are Identified To Be RORγt<sup>+</sup>Foxp3<sup>+</sup> Double Positive

As continuous AM80 treatment suppressed the production of IL-10 by CNS-infiltrating T cells, we further investigated the effect of AM80 on the production of IL-10 *in vitro*. As shown in Figure 6A, we detected higher levels of IL-10 transcripts among effector T cells activated under Th17 priming conditions as compared with those primed under neutral conditions, supporting a previous study in which demonstrate increase IL-10 secretion in Th17 cultures on TGF-β and IL-6 signaling.<sup>24</sup> Addition of RAR agonists during the primary culture confers reduced expression of IL-10 transcripts, even when cells underwent a secondary stimulation in the absence of retinoids, pointing toward a stable phenotype. To assess the effect of retinoids on differentiated Th17 cells, T cells were primed under Th17 priming conditions without retinoids, but restimulation in the presence of retinoids decreased the expression of IL-10 (Figure 6B). In addition, *ex vivo* restimulation of CNS-infiltrating T cells in the presence of retinoids also reduced IL-10 expression (Figure 6C). Taken together, treatment with either AM80 or ATRA inhibits the production of IL-10 by Th17 cells, which have differentiated *in vitro* or *in vivo*.

Regulatory T cells have previously been indicated as main source of IL-10-producing T cell subsets.<sup>44</sup> In Figure 6D, we confirm that Foxp3<sup>+</sup> cells are the source of IL-10 in whole splenocyte stimulated in the presence of

and restimulated with anti-CD3 antibody for another 96 hours and examined for expression of IL-10 transcript. **B:** Naive T cells stimulated under Th17 priming conditions without RAR agonists were then restimulated as described above. RAR agonists (100 nmol/L) were added during the secondary stimulation and assessed for their expression of IL-10 mRNA by quantitative RT-PCR. Results are representative of two independent experiments. **C:** CNS-infiltrating T cells were isolated from EAE animals (score 3) and restimulated with immobilized anti-CD3 mAb in the presence of AM80 or ATRA (100 nmol/L) for 72 hours. The amount of IL-10 in culture supernatants was examined by ELISA. **D:** Whole splenocytes were cultured for 96 hours with TGF-β and IL-6 and RAR agonists (10 nmol/L) and CBA analysis were performed for analyzing the levels of IL-10 production in the culture supernatants. Cells were then subjected to intracellular cytokine staining by fluorescence-activated cell sorting. The histogram gated on CD45<sup>hi</sup>Sb<sup>+</sup>TCR<sup>+</sup>CD4<sup>+</sup> lymphocytes displays the comparative population of IL-10<sup>+</sup> cells by Foxp3<sup>+</sup> (black line) and Foxp3<sup>-</sup> (gray line) cell populations. Broken line represents the data stained with isotype control antibody. **E:** CD45<sup>hi</sup>Sb<sup>+</sup>TCR<sup>+</sup>CD4<sup>+</sup> lymphocytes derived from **D** were analyzed for their expression of RORγt and Foxp3. Comparative dot plots were shown with or without AM80 treatment (10 nmol/L). Addition of AM80 decreased the number of RORγt<sup>+</sup> cells, including the RORγt<sup>+</sup>FOXP3<sup>+</sup> cells. **F:** The levels of IL-10 producing cells from the quadrants labeled in **E** are shown.

TGF- $\beta$  and IL-6. Interestingly, addition of retinoids expanded the number of Foxp3<sup>+</sup> cells (data not shown), but abolished their production of IL-10 (Figure 6D). Recently, a further subset of IL-10 producing regulatory T cells that express of ROR $\gamma$ t and Foxp3 simultaneously have been highlighted.<sup>25</sup> Therefore, we hypothesized that such cells represent the major population of IL-10-secreting T cells in Th17 differentiation cultures and thus-it is these cells on which retinoids act to abolish IL-10 production. In fact, Foxp3<sup>+</sup> cells could be successfully divided into ROR $\gamma$ t<sup>+</sup> and ROR $\gamma$ t<sup>-</sup> populations (Figure 6E). Addition of AM80 to cultures reduced the proportion of ROR $\gamma$ t<sup>+</sup> and ROR $\gamma$ t<sup>+</sup>Foxp3<sup>+</sup> double positive cells, but increased the proportion of Foxp3<sup>+</sup>ROR $\gamma$ t<sup>-</sup> cells (Figure 6E), at the same time as abolishing IL-10 production (Figure 6D). Intracellular IL-10 staining revealed that the ROR $\gamma$ t<sup>+</sup>Foxp3<sup>+</sup> double positive population was the predominant source of IL-10 (Figure 6F). Taken together, these data suggests that retinoid treatment reduces the production of IL-10 by inhibiting the effector function of a distinct ROR $\gamma$ t<sup>+</sup>Foxp3<sup>+</sup> population, which might serve as a regulatory T cell subset.

## Discussion

ATRA, the physiologically active metabolite of vitamin A, inhibits the differentiation of Th17 cells and reciprocally promotes the generation of Treg cells *in vitro*.<sup>27,29</sup> In this study, we demonstrate for the first time that the synthetic RAR agonist AM80 inhibits the differentiation and effector function of Th17 cells more effectively than ATRA. Oral administration of AM80 attenuates antigen-specific Th17 cell differentiation, thus such treatment is able to reduce disease in the acute phase of EAE.

ATRA suppresses Th17 differentiation and effector function via RAR $\alpha$  signaling,<sup>27,28</sup> but ATRA can also bind to RAR $\beta$  and RAR $\gamma$ , which can form a variety of homo- and heterodimers with three RXR receptors.<sup>26,34</sup> Non-selective receptor binding is thought to be a major cause of the side effects associated with the administration of ATRA and other pan-RAR agonists in the clinic. AM80 is a synthetic RAR agonist that has high affinity to the RAR $\alpha/\beta$  and is currently available as medication for human diseases such as APL and psoriasis.<sup>35-37,45</sup> In addition to greater specificity for RAR $\alpha$ , AM80 offers several other advantages over ATRA as a therapeutic agent: it has a lower toxicity, higher stability, fewer potential side effects, and a superior bioavailability. Also, we demonstrate that a lower dose of AM80 is required to inhibit IL-17 production by T cells than similar treatment with ATRA. Interestingly, at very low doses, AM80 treatment reduced IL-17 production dramatically, while ROR $\gamma$ t expression was only slightly reduced. Although ROR $\gamma$ t expression is a pre-requisite for Th17 development,<sup>17</sup> recent studies have demonstrated that Foxp3 can physically interact with ROR $\gamma$ t inhibiting IL-17 production when both are co-expressed.<sup>25,46,47</sup> It is conceivable that the disconnection between ROR $\gamma$ t expression and IL-17 secretion following low dose AM80 treatment we observed may result from concomitant Foxp3 up-regulation.

However, at higher doses of AM80, the inhibition of IL-17 production was associated with a reduction in ROR $\gamma$ t expression; thus there may be multiple mechanisms of action in operation.

A recent study demonstrated that intraperitoneal injection of high doses of ATRA protected animals from EAE induction and that this protection was associated with reduced IL-17 and IFN- $\gamma$  production.<sup>30</sup> However, such treatment was also found to reduce proliferative T cells responses following antigen restimulation *ex vivo*,<sup>30</sup> suggesting that, in this study, ATRA may ameliorate EAE by generating systemic immune-suppression. We have observed similar immunosuppression by ATRA and other retinoids at a high dose (data not shown). In our study, we treated with AM80 at a dose 5–10 times lower than the dose that ATRA has been previously tested at and we were able to administer the retinoid orally. Our treatment regimen also suppressed Th17 cell differentiation and IL-17 production, but antigen-specific T cell proliferation was not altered. Thus, we were able to target pathogenic Th17 cells specifically, without inducing general immunosuppression.

We and others have found that there is no induction of regulatory T cells when treating inflammatory diseases with RAR agonists,<sup>29,30</sup> and it has been speculated that this may be due to a lack of TGF- $\beta$  *in vivo*.<sup>29</sup> An alternative hypothesis is that Treg generation is inhibited by the strong induction of inflammatory cytokines, including IL-6, TNF- $\alpha$ , and IL-1.<sup>30</sup> Thus, under both of these suggestions, it is likely that AM80 suppresses EAE by inhibiting the generation and activity of Th17 cells.

AM80 treatment inhibited acute EAE in mice, but continuous administration of AM80 did not suppress chronic inflammation. Interestingly, T cells isolated from the CNS tissue of AM80-treated mice during the chronic phase of the disease continued to express only low levels of ROR $\gamma$ t, IL-23 receptor, and IL-17. This was in marked contrast to vehicle-treated control mice, which, despite analogous clinical disease scores, had CNS-infiltrating T cells that expressed high levels of Th17-related factors. Thus, we suggest that inhibition of Th17 cell function alone is not sufficient to protect mice from chronic CNS inflammation.

In fact, we found that treatment with RAR-agonists suppressed T cell production of IL-10 at late stage disease. Recently, a unique T cell subset that co-expresses ROR $\gamma$ t and Foxp3, and predominantly secretes regulatory IL-10 has been identified *in vivo*.<sup>23</sup> Intriguingly, we show that RAR agonists not only suppress pathogenic Th17 cells, but also suppress IL-10 production by these ROR $\gamma$ t+Foxp3<sup>+</sup> T cells. Korn *et al* propose a model of sequential infiltration by different subsets of differentiated CD4<sup>+</sup> T cells during organ-specific autoimmunities such as EAE. In this model, Th17 cells mediate the acute phase of disease, while Th1 cells are more prominent in the chronic phase, and at much later phase of disease, there is a moderate up-regulation of IL-10 production.<sup>48,49</sup> In addition, previous studies indicate that IL-10 is a key cytokine for the suppression of T cell-mediated autoimmune inflammation in the CNS.<sup>23,49</sup> Therefore, the residual expression of IL-17 or IFN- $\gamma$  at later time points



with continuous AM80 treatment may cause moderate progression of EAE development under the condition with reduced IL-10 production *in vivo*. We have little information about how those two subsets of IL-17-producing inflammatory ROR $\gamma$ t+ T cells and IL-10-producing immunoregulatory ROR $\gamma$ t+Foxp3+ T cells developed during immunological responses. Although we demonstrate here that the suppressive effect of AM80 on EAE is most likely through modulation of T cell production of both IL-17 and IL-10 *in vivo*, we don't exclude the possible involvement other factors, as EAE/MS are complex diseases that do not depend only on a IL-17 versus IL-10 dichotomy. For example, ongoing neurodegeneration<sup>50</sup> may also contribute significantly to the observed phenotype especially at the later phase of disease. As Th1 cells also play a significant role in the induction or EAE/MS, we should also point out that retinoic acid has been previously shown to exert direct effects on T cells, suppressing Th1 development and enhancing Th2 development via retinoic acid receptors.<sup>51</sup> This indicates that AM80 has multiple beneficial effects for disease protection through oral administration. The immunomodulatory effect of RAR agonists on those Th17-like regulatory T cells in inflammatory autoimmune diseases is to be considered if conducting *in vivo* attenuation of Th17 cells for treatment of autoimmune diseases with retinoids. At this point in time, we advocate that the use of AM80 on inflammatory autoimmune diseases should target the acute phases of Th17-mediated pathogenesis.

In addition, we found that T cells that had infiltrated the spinal cord produced much lower amounts of CCL2 (MCP-1) as compared with brain-infiltrating T cells. CCL2 plays a crucial role in the progression of EAE<sup>52,53</sup> and previously we have shown that human Th17 cells express the corresponding receptor, CCR2.<sup>54</sup> Our data showing reductions in CCL2 in the spinal cord may represent previously reported differences in inflammatory cell populations at different CNS sites.<sup>55</sup>

In summary, we demonstrate that oral treatment with AM80 effectively inhibits IL-17 production without generating systemic immunosuppression. In addition, AM80 treatment protected mice from the development of acute EAE and rescued mice with established EAE from acute autoimmune inflammation. Collectively, these data advocate AM80 as a potent therapeutic agent against acute Th17-mediated autoimmune diseases including MS.

## References

1. Sospedra M, Martin R: Immunology of multiple sclerosis. *Annu Rev Immunol* 2005, 23:683-747
2. Kuchroo VK, Anderson AC, Waldner H, Munder M, Bettelli E, Nicholson LB: T cell response in experimental autoimmune encephalomyelitis (EAE): role of self and cross-reactive antigens in shaping, tuning, and regulating the autopathogenic T cell repertoire. *Annu Rev Immunol* 2002, 20:101-123
3. Gold R, Lington C, Lassmann H: Understanding pathogenesis and therapy of multiple sclerosis via animal models: 70 years of merits and culprits in experimental autoimmune encephalomyelitis research. *Brain* 2006, 129:1953-1971
4. Murphy KM, Ouyang W, Farrar JD, Yang J, Ranganath S, Asnagli H, Afkarian M, Murphy TL: Signaling and transcription in T helper development. *Annu Rev Immunol* 2000, 18:451-494
5. Zamvil SS, Steinman L: The T lymphocyte in experimental allergic encephalomyelitis. *Annu Rev Immunol* 1990, 8:579-621
6. Cua DJ, Sherlock J, Chen Y, Murphy CA, Joyce B, Seymour B, Luciani L, To W, Kwan S, Churakova T, Zurawski S, Wiekowski M, Lira SA, Gorman D, Kastelein RA, Sedgwick JD: Interleukin-23 rather than interleukin-12 is the critical cytokine for autoimmune inflammation of the brain. *Nature* 2003, 421:744-748
7. Langrish CL, Chen Y, Blumenschein WM, Mattson J, Basham B, Sedgwick JD, McClanahan T, Kastelein RA, Cua DJ: IL-23 drives a pathogenic T cell population that induces autoimmune inflammation. *J Exp Med* 2005, 201:233-240
8. Komiyama Y, Nakae S, Matsuki T, Nambu A, Ishigame H, Kakuta S, Sudo K, Iwakura Y: IL-17 plays an important role in the development of experimental autoimmune encephalomyelitis. *J Immunol* 2006, 177:566-573
9. McGeachy MJ, Cua DJ: The link between IL-23 and Th17 cell-mediated immune pathologies. *Semin Immunol* 2007, 19:372-376
10. Matuszewski D, Kivisakk P, He B, Kostulas N, Ozenci V, Fredrikson S, Link H: Interleukin-17 mRNA expression in blood and CSF mononuclear cells is augmented in multiple sclerosis. *Mult Scler* 1999, 5:101-104
11. Tzartos JS, Friese MA, Craner MJ, Palace J, Newcombe J, Esiri MM, Fugger L: Interleukin-17 production in central nervous system-infiltrating T cells and glial cells is associated with active disease in multiple sclerosis. *Am J Pathol* 2008, 172:146-155
12. Veldhoen M, Hocking RJ, Atkins CJ, Locksley RM, Stockinger B: TGFbeta in the context of an inflammatory cytokine milieu supports de novo differentiation of IL-17-producing T cells. *Immunity* 2006, 24:179-189
13. Bettelli E, Carrier Y, Gao W, Korn T, Strom TB, Oukka M, Weiner HL, Kuchroo VK: Reciprocal developmental pathways for the generation of pathogenic effector TH17 and regulatory T cells. *Nature* 2006, 441:235-238
14. Mangan PR, Harrington LE, O'Quinn DB, Helms WS, Bullard DC, Elson CO, Hatton RD, Wahl SM, Schoeb TR, Weaver CT: Transforming growth factor-beta induces development of the T(H)17 lineage. *Nature* 2006, 441:231-234
15. Shevach EM, Davidson TS, Huter EN, Dipaolo RA, Andersson J: Role of TGF-beta in the induction of Foxp3 expression and T regulatory cell function. *J Clin Immunol* 2008, 28:640-646
16. Korn T, Bettelli E, Gao W, Awasthi A, Jager A, Strom TB, Oukka M, Kuchroo VK: IL-21 initiates an alternative pathway to induce proinflammatory T(H)17 cells. *Nature* 2007, 448:484-487
17. Ivanov II, McKenzie BS, Zhou L, Tadokoro CE, Lepelley A, Lafaille JJ, Cua DJ, Littman DR: The orphan nuclear receptor RORgamma directs the differentiation program of proinflammatory IL-17+ T helper cells. *Cell* 2006, 126:1121-1133
18. Harris TJ, Grosso JF, Yen HR, Xin H, Kortylewski M, Albesiano E, Hipkiss EL, Getnet D, Goldberg MV, Maris CH, Housseau F, Yu H, Pardoll DM, Drake CG: Cutting edge: an *in vivo* requirement for STAT3 signaling in TH17 development and TH17-dependent autoimmunity. *J Immunol* 2007, 179:4313-4317
19. Liang SC, Tan XY, Luxenberg DP, Karim R, Dunussi-Joannopoulos K, Collins M, Fouser LA: Interleukin (IL)-22 and IL-17 are coexpressed by Th17 cells and cooperatively enhance expression of antimicrobial peptides. *J Exp Med* 2006, 203:2271-2279
20. Sakaguchi S, Ono M, Setoguchi R, Yagi H, Hori S, Fehervari Z, Shimizu J, Takahashi T, Nomura T: Foxp3+ CD25+ CD4+ natural regulatory T cells in dominant self-tolerance and autoimmune disease. *Immunol Rev* 2006, 212:8-27
21. Fontenot JD, Rasmussen JP, Williams LM, Dooley JL, Farr AG, Rudensky AY: Regulatory T cell lineage specification by the forkhead transcription factor foxp3. *Immunity* 2005, 22:329-341
22. Brunkow ME, Jeffery EW, Hjerrild KA, Paeppe B, Clark LB, Yasayko SA, Wilkinson JE, Galas D, Ziegler SF, Ramsdell F: Disruption of a new forkhead/winged-helix protein, scurf, results in the fatal lymphoproliferative disorder of the scurfy mouse. *Nat Genet* 2001, 27:68-73
23. Zhang X, Koldzic DN, Izikson L, Reddy J, Nazareno RF, Sakaguchi S, Kuchroo VK, Weiner HL: IL-10 is involved in the suppression of experimental autoimmune encephalomyelitis by CD25+CD4+ regulatory T cells. *Int Immunol* 2004, 16:249-256

24. McGeachy MJ, Bak-Jensen KS, Chen Y, Tato CM, Blumenschein W, McClanahan T, Cua DJ: TGF-beta and IL-6 drive the production of IL-17 and IL-10 by T cells and restrain T(H)-17 cell-mediated pathology. *Nat Immunol* 2007, 8:1390-1397
25. Lochner M, Peduto L, Cherrier M, Sawa S, Langa F, Varona R, Riethmacher D, Si-Tahar M, Di Santo JP, Eberl G: In vivo equilibrium of proinflammatory IL-17+ and regulatory IL-10+ Foxp3+ ROR-gamma t+ T cells. *J Exp Med* 2008, 205:1381-1393
26. Mark M, Ghyselincx NB, Chambon P: Function of retinoid nuclear receptors: lessons from genetic and pharmacological dissections of the retinoic acid signaling pathway during mouse embryogenesis. *Annu Rev Pharmacol Toxicol* 2006, 46:451-480
27. Schambach F, Schupp M, Lazar MA, Reiner SL: Activation of retinoic acid receptor-alpha favours regulatory T cell induction at the expense of IL-17-secreting T helper cell differentiation. *Eur J Immunol* 2007, 37:2396-2399
28. Elias KM, Laurence A, Davidson TS, Stephens G, Kanno Y, Shevach EM, O'Shea JJ: Retinoic acid inhibits Th17 polarization and enhances FoxP3 expression through a Stat-3/Stat-5 independent signaling pathway. *Blood* 2008, 111:1013-1020
29. Mucida D, Park Y, Kim G, Turovskaya O, Scott I, Kronenberg M, Cheroutre H: Reciprocal TH17 and regulatory T cell differentiation mediated by retinoic acid. *Science* 2007, 317:256-260
30. Xiao S, Jin H, Korn T, Liu SM, Oukka M, Lim B, Kuchroo VK: Retinoic acid increases Foxp3+ regulatory T cells and inhibits development of Th17 cells by enhancing TGF-beta-driven Smad3 signaling and inhibiting IL-6 and IL-23 receptor expression. *J Immunol* 2008, 181:2277-2284
31. Massaccesi L, Abbamondi AL, Giorgi C, Sarlo F, Lolli F, Amaducci L: Suppression of experimental allergic encephalomyelitis by retinoic acid. *J Neurol Sci* 1987, 80:55-64
32. Racke MK, Burnett D, Pak SH, Albert PS, Cannella B, Raine CS, McFarlin DE, Scott DE: Retinoid treatment of experimental allergic encephalomyelitis. IL-4 production correlates with improved disease course. *J Immunol* 1995, 154:450-458
33. Patatanian E, Thompson DF: Retinoic acid syndrome: a review. *J Clin Pharm Ther* 2008, 33:331-338
34. Lefebvre P, Martin PJ, Flajollet S, Dedieu S, Billaut X, Lefebvre B: Transcriptional activities of retinoic acid receptors. *Vitam Horm* 2005, 70:199-264
35. Tobita T, Takeshita A, Kitamura K, Ohnishi K, Yanagi M, Hiraoka A, Karasuno T, Takeuchi M, Miyawaki S, Ueda R, Naoe T, Ohno R: Treatment with a new synthetic retinoid, Am80, of acute promyelocytic leukemia relapsed from complete remission induced by all-trans retinoic acid. *Blood* 1997, 90:967-973
36. Miwako I, Kagechika H: Tamibarotene. *Drugs Today (Barc)* 2007, 43:563-568
37. Ohnishi K: PML-RARalpha inhibitors (ATRA, tamibarotene, arsenic trioxide) for acute promyelocytic leukemia. *Int J Clin Oncol* 2007, 12:313-317
38. Kagechika H: Novel synthetic retinoids and separation of the pleiotropic retinoid activities. *Curr Med Chem* 2002, 9:591-608
39. Doi Y, Oki S, Ozawa T, Hohjoh H, Miyake S, Yamamura T: Orphan nuclear receptor NR4A2 expressed in T cells from multiple sclerosis mediates production of inflammatory cytokines. *Proc Natl Acad Sci USA* 2008, 105:8381-8386
40. Croxford JL, Miyake S, Huang YY, Shimamura M, Yamamura T: Invariant V(alpha)19i T cells regulate autoimmune inflammation. *Nat Immunol* 2006, 7:987-994
41. Raveney BJ, Richards C, Akinin ML, Copland DA, Burton BR, Kerr E, Nicholson LB, Williams NA, Dick AD: The B subunit of Escherichia coli heat-labile enterotoxin inhibits Th1 but not Th17 cell responses in established experimental autoimmune uveoretinitis. *Invest Ophthalmol Vis Sci* 2008, 49:4008-4017
42. Aranami T, Yamamura T: Th17 Cells and autoimmune encephalomyelitis (EAE/MS). *Allergol Int* 2008, 57:115-120
43. Lexberg MH, Taubner A, Forster A, Albrecht I, Richter A, Kamradt T, Radbruch A, Chang HD: Th memory for interleukin-17 expression is stable in vivo. *Eur J Immunol* 2008, 38:2654-2664
44. Zhou X, Schmidtke P, Zepp F, Meyer CU: Boosting interleukin-10 production: therapeutic effects and mechanisms. *Curr Drug Targets Immune Endocr Metabol Disord* 2005, 5:465-475
45. Takeuchi M: Clinical experience with a new synthetic retinoid, tami-barotene (Am-80) for relapsed or refractory acute promyelocytic leukemia. *Gan To Kagaku Ryoho* 2006, 33:397-401
46. Zhou L, Lopes JE, Chong MM, Ivanov II, Min R, Victora GD, Shen Y, Du J, Rubtsov YP, Rudensky AY, Ziegler SF, Littman DR: TGF-beta-induced Foxp3 inhibits T(H)17 cell differentiation by antagonizing RORgamma function. *Nature* 2008, 453:236-240
47. Ichihama K, Yoshida H, Wakabayashi Y, Chinen T, Saeki K, Nakaya M, Takaesu G, Hori S, Yoshimura A, Kobayashi T: Foxp3 inhibits RORgamma-mediated IL-17A mRNA transcription through direct interaction with RORgamma. *J Biol Chem* 2008, 283:17003-17008
48. Korn T, Oukka M, Kuchroo V, Bettelli E: Th17 cells: effector T cells with inflammatory properties. *Semin Immunol* 2007, 19:362-371
49. Korn T, Reddy J, Gao W, Bettelli E, Awasthi A, Petersen TR, Backstrom BT, Sobel RA, Wucherpfennig KW, Strom TB, Oukka M, Kuchroo VK: Myelin-specific regulatory T cells accumulate in the CNS but fail to control autoimmune inflammation. *Nat Med* 2007, 13:423-431
50. Herrero-Herranz EPL, Gold R, Linker RA: Pattern of axonal injury in murine myelin oligodendrocyte glycoprotein induced experimental autoimmune encephalomyelitis: implications for multiple sclerosis. *Neurobiol Dis* 2008, 30:162-173
51. Iwata M, Eshima Y, Kagechika H: Retinoic acids exert direct effects on T cells to suppress Th1 development and enhance Th2 development via retinoic acid receptors. *Int Immunol* 2003, 15:1017-1025
52. Izikson L, Klein RS, Charo IF, Weiner HL, Luster AD: Resistance to experimental autoimmune encephalomyelitis in mice lacking the CC chemokine receptor (CCR)2. *J Exp Med* 2000, 192:1075-1080
53. Dogan RN, Elhofy A, Karpus WJ: Production of CCL2 by central nervous system cells regulates development of murine experimental autoimmune encephalomyelitis through the recruitment of TNF- and iNOS-expressing macrophages and myeloid dendritic cells. *J Immunol* 2008, 180:7376-7384
54. Sato W, Aranami T, Yamamura T: Cutting edge: human Th17 cells are identified as bearing CCR2+CCR5- phenotype. *J Immunol* 2007, 178:7525-7529
55. Stromnes IM, Cerretti LM, Liggitt D, Harris RA, Goverman JM: Differential regulation of central nervous system autoimmunity by T(H)1 and T(H)17 cells. *Nat Med* 2008, 14:337-342

# Suppression of Experimental Autoimmune Encephalomyelitis by Ghrelin<sup>1</sup>

Michael-Mark Theil,<sup>\*†</sup> Sachiko Miyake,<sup>2\*</sup> Miho Mizuno,<sup>\*</sup> Chiharu Tomi,<sup>\*</sup> J. Ludovic Croxford,<sup>\*</sup> Hiroshi Hosoda,<sup>‡</sup> Julia Theil,<sup>\*†</sup> Stephan von Hörsten,<sup>†§</sup> Hiroaki Yokote,<sup>\*</sup> Asako Chiba,<sup>\*</sup> Youwei Lin,<sup>\*</sup> Shinji Oki,<sup>\*</sup> Takashi Akamizu,<sup>¶</sup> Kenji Kangawa,<sup>¶</sup> and Takashi Yamamura<sup>2\*</sup>

Ghrelin is a recently identified gastric hormone that displays strong growth hormone-releasing activity mediated by the growth hormone secretagogue receptor. While this unique endogenous peptide participates in the regulation of energy homeostasis, increases food intake, and decreases energy expenditure, its ability to inhibit the production of proinflammatory cytokines *in vitro* indicates its role in the regulation of inflammatory process *in vivo*. Here we examine the effect of exogenous ghrelin on the development of experimental autoimmune encephalomyelitis (EAE), a representative model of multiple sclerosis. In the C57BL/6 mouse model of EAE induced by sensitization to myelin oligodendrocyte glycoprotein 35–55 peptide, we found that alternate-day s.c. injections of ghrelin (5  $\mu$ g/kg/day) from day 1 to 35 significantly reduced the clinical severity of EAE. The suppression of EAE was accompanied by reduced mRNA levels of proinflammatory cytokines such as TNF- $\alpha$ , IL-1 $\beta$ , and IL-6 in the spinal cord cellular infiltrates and microglia from ghrelin-treated mice at the peak of disease, suggesting the role of ghrelin as an antiinflammatory hormone. Consistently, ghrelin significantly suppressed the production of proinflammatory cytokines in LPS-stimulated microglia *in vitro*. These results shed light on the new role of ghrelin in the regulation of inflammation with possible implications for management of human diseases. *The Journal of Immunology*, 2009, 183: 2859–2866.

Small synthetic compounds, referred to as growth hormone (GH)<sup>3</sup> secretagogues (GHS), have been known to stimulate GH release, working through a G protein-coupled receptor called GHS receptor (GHS-R) (1–3). It is now established that a new endogenous peptide, ghrelin, discovered in rat gastric extracts, is an endogenous ligand for GHS-R and is involved in the regulation of GH release. Ghrelin is a 28-aa polypeptide with an essential *n*-octanoyl modification on serine at position 3 (4). Although ghrelin is predominantly secreted from mucosal endocrine cells of stomach, it is widely distributed in various organs, including lymphoid tissues (5, 6). Furthermore, it is measurable in the systemic circulation, indicating its hormonal nature (7).

Ghrelin does not only stimulate GH release, but it also increases food intake, regulates energy homeostasis, and decreases energy expenditure by lowering the catabolism of fat (4, 8, 9). Because of its orexigenic and adipogenic character, ghrelin may be potentially useful for the treatment of anorexia and cachexia (10, 11). Although the precise mechanisms remain to be clarified, the orexigenic activities of ghrelin may be mediated by another feeding regulatory hormone neuropeptide Y (NPY) via stimulation of Y1 and Y5 receptors (12). Furthermore, the antagonistic effect of ghrelin on leptin-induced decrease of food intake seems to be mediated by ghrelin-induced release of NPY and subsequent stimulation of the Y1 receptor (13).

Ghrelin has been shown to exhibit antiinflammatory functions against T cells and macrophages *in vitro* (14–16). The potential activity of ghrelin as antiinflammatory reagent *in vivo* was shown in several animal models, including bowel disease (17), arthritis (16, 18), sepsis, and endotoxemia (16, 19, 20). Here we report that s.c. injections of ghrelin could significantly attenuate the clinical severity of the representative model of experimental autoimmune encephalomyelitis (EAE) induced in C57BL/6 (B6) mice by sensitization against myelin oligodendrocyte glycoprotein (MOG)<sub>35–55</sub> peptide. Furthermore, we demonstrate that *in vivo* treatment with ghrelin significantly suppressed the mRNA levels of the proinflammatory cytokines TNF- $\alpha$ , IL-1 $\beta$ , and IL-6 in microglia and infiltrating T cells derived from the spinal cords of ghrelin-treated mice. Finally, we confirm that LPS-stimulated microglia and monocytes produced lower amounts of proinflammatory cytokines when they were pretreated with ghrelin *in vitro*. In conclusion, the present study indicates the potential use of ghrelin as an antiinflammatory drug to control human CNS pathology.

## Materials and Methods

### Mice and reagents

We used female B6 mice (CLEA Japan) between 6 and 10 wk of age in specific pathogen-free conditions. Animal care and use were in accordance

\*Department of Immunology, National Institute of Neuroscience, National Centre of Neurology and Psychiatry, Tokyo, Japan; <sup>†</sup>Department of Functional and Applied Anatomy, Hannover Medical School, Hannover, Germany; <sup>‡</sup>Department of Biochemistry, National Cardiovascular Center Research Institute, Fujishirodai, Suita, Osaka, Japan; <sup>§</sup>Experimental Therapy, Franz-Penzoldt-Center, Friedrich-Alexander-University Erlangen-Nürnberg, Erlangen, Germany; and <sup>¶</sup>Ghrelin Research Project, Translational Research Center, Kyoto University Faculty of Medicine, Kyoto, Japan

Received for publication October 14, 2008. Accepted for publication June 9, 2009.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

<sup>1</sup> This work was supported by the grants from the Ministry of Health, Labor, and Welfare of Japan, the Program for Promotion of Fundamental Studies in Health Sciences of the National Institute of Biomedical Innovation (NIBIO), and the Ministry of Education, Science, Culture, Sports, and Technology of Japan.

<sup>2</sup> Address correspondence and reprint requests to Dr. Sachiko Miyake or Dr. Takashi Yamamura, Department of Immunology, National Institute of Neuroscience, National Centre of Neurology and Psychiatry, 4-1-1 Ogawahigashi, Kodaira, Tokyo 187-8502. E-mail address: miyake@ncnp.go.jp or yamamura@ncnp.go.jp

<sup>3</sup> Abbreviations used in this paper: GH, growth hormone; EAE, experimental autoimmune encephalomyelitis; GHS, growth hormone secretagogue; GHS-R, growth hormone secretagogue receptor; LN, lymph node; MOG, myelin oligodendrocyte glycoprotein; NPY, neuropeptide Y.

Copyright © 2009 by The American Association of Immunologists, Inc. 0022-1767/09/\$2.00

Table I. Amino acid sequence of mouse ghrelin and des-acyl ghrelin

Peptide	Amino Acid Sequence <sup>a</sup>	Ser <sup>3</sup> acylation	Reference
Ghrelin	GSS <u>FLS</u> PEHQKAQQRKESKKPPAKLQPR	<i>n</i> -Octanoic acid	(4)
Des-acyl ghrelin	GSS <u>FLS</u> PEHQKAQQRKESKKPPAKLQPR		(7)

<sup>a</sup> The underlined letter S represents the third serine (Ser<sup>3</sup>).

with institutional guidelines. Animal experiments were approved by our institutional review committee. Rat MOG<sub>35-55</sub> (amino acid sequence MEVGWYRSPFSRVVHLYRNGK) was synthesized at Toray Research Center (Tokyo, Japan). Ghrelin and des-acyl ghrelin (Table I) were synthesized as previously described (4, 7).

#### Immunization and clinical assessment of EAE

We immunized mice ( $n = 5-15$  per group) s.c. in the tail base with 100  $\mu$ g of MOG<sub>35-55</sub>-peptide dissolved in 0.1 ml of PBS and 0.1 ml of CFA containing 1 mg of *M. tuberculosis* H37Ra (Difco). Shortly after immunization and 48 h later, the mice were injected i.p. with 200 ng of pertussis toxin (List Biological Laboratories). Clinical scores of EAE were daily assigned as follows: 0, normal; 1, weakness of the tail and/or paralysis of the distal half of the tail; 2, loss of tail tonicity; 3, partial hind limb paralysis; 4, complete hind limb paralysis; 5, forelimb paralysis or moribund; 6, death. The cumulative scores were calculated for individual mice by summing up the daily scores.

#### Administration of ghrelin and des-acyl ghrelin

For EAE treatment, we s.c. injected ghrelin and des-acyl ghrelin diluted in 0.9% saline. In the first series of experiments, mice were injected with ghrelin or des-acyl ghrelin at doses of 0.5, 5, or 50  $\mu$ g/kg every other day for 35 days. Sham-treated animals were injected with 0.9% saline (standard protocol). In the next experiment, we injected the mice with 5  $\mu$ g/kg ghrelin every day from day 1 to 10 (induction phase treatment) or from day 11 to 20 (effector phase treatment) and in-between with 0.9% saline. The controls were injected every day from day 1 to 20 with 0.9% saline (alternative protocol).

#### Assessment of histological EAE

To evaluate the histological manifestations of EAE, we treated mice with 5  $\mu$ g/kg ghrelin or 0.9% saline following the standard protocol and sacrificed them on day 17 postimmunization. The spinal cords were removed and fixed in buffered formalin. They were embedded in paraffin, sectioned, and stained with H&E and Luxol fast blue for histopathological analysis.

#### Flow cytometry and isolation of mononuclear cells from the CNS

B6 mice were challenged for EAE, treated following the standard protocol with 5  $\mu$ g/kg ghrelin or 0.9% saline and sacrificed on day 17 postimmunization. We removed spleen, lymph nodes (LN), and thymus as well as spinal cord from the ghrelin- and saline-treated mice for flow cytometer analysis. Single-cell suspensions were prepared according to standard methods. The spinal cord cell suspensions were centrifuged at  $200 \times g$  for 10 min and resuspended in 4 ml of 70% isotonic Percoll (Amersham Biosciences)/PBS and overlaid by equal volumes of 37% and 30% isotonic Percoll. The gradient was centrifuged at  $500 \times g$  for 15 min and the mononuclear cells were harvested from the 37%–70% interface, washed, and counted. The cells were stained for 5 min with anti-Fc $\gamma$  III/II mAb (BD Pharmingen), washed, and labeled with the following mAbs for surface phenotype analysis: FITC-CD4 mAb, FITC-CD19 mAb, PE-CD8a mAb, PE-NK1.1 mAb, PE-CD25 mAb, allophycocyanin-FoxP3, and PerCP-Cy5.5-CD3e mAb (BD Pharmingen) and FITC-F4/80 mAb (Dainihon Seiyaku). The cytofluorometric analysis was performed using a FACSCalibur operated by CellQuest software (BD Biosciences).

#### Cytokine and cell proliferation assay

MOG<sub>35-55</sub>-immunized B6 mice were treated s.c. with 5  $\mu$ g/kg/day of ghrelin or 0.9% saline every day from day 1 to 10. The LN cells were collected on day 11 after immunization and suspended in our standard lymphocyte culture medium (RPMI 1640 supplemented with  $5 \times 10^{-5}$  M 2-ME, 2 mM L-glutamine, 100 U/ml penicillin/streptomycin) added with 1% syngeneic mouse serum. The cells were cultured in 96-well round-bottom plates at  $1 \times 10^6$ /well for 72 h in the presence of 100  $\mu$ g/ml

MOG<sub>35-55</sub>. Levels of IFN- $\gamma$ , IL-17, and IL-4 in the supernatant were determined by using a sandwich ELISA. Proliferative responses were measured using a Beta-1205 counter (Pharmacia) to detect the incorporation of [<sup>3</sup>H]thymidine (1  $\mu$ Ci/well) for the final 16 h of culture.

#### Evaluation of encephalitogenic T cell induction in B6 mice treated with ghrelin

To evaluate whether in vivo ghrelin treatment may affect the induction of encephalitogenic T cells after immunization with MOG<sub>35-55</sub>, we evaluated the ability of the lymphoid cells from ghrelin- or saline-treated mice to passively transfer EAE into naive recipients. Donor B6 mice were immunized with MOG<sub>35-55</sub> and treated every day from day 1 to 10 with 5  $\mu$ g/kg/day of ghrelin or 0.9% saline. We removed spleens and LN from the donor mice on day 11 and prepared lymphoid cell suspensions. The lymphoid cells were stimulated with MOG<sub>35-55</sub> (33  $\mu$ g/ml) in the standard medium added with FCS (10%) for 96 h and then we isolated the CD4<sup>+</sup> T cells for cell transfer by depletion of CD8<sup>+</sup>, CD19<sup>+</sup>, and NK1.1<sup>+</sup> cells. In brief, the MOG<sub>35-55</sub>-stimulated total lymphoid cells were labeled with PE-CD8a mAb, PE-NK1.1 mAb, and PE-CD19 mAb (BD Pharmingen) for 30 min, washed, and incubated with anti-PE microbeads (Miltenyi Biotec) for 15 min. Using autoMACS (Miltenyi Biotec), we isolated CD4<sup>+</sup> T cells (CD8<sup>-</sup>, CD19<sup>-</sup>, and NK1.1<sup>-</sup> fraction) as a pass-through and suspended the cells in PBS. We injected  $1.0 \times 10^7$  of the cells into the peritoneal cavity of syngeneic recipient mice that had been X-irradiated (550 rad) shortly before. We also injected 200 ng of pertussis toxin i.p. on the same day and 48 h later.

#### Reverse transcription and real-time PCR

To analyze the mechanism of ghrelin effects in vivo, we extracted total RNA from spinal cord, spleen, thymus, and LN samples using the RNeasy Mini Kit (Qiagen). The RNA was subjected to reverse transcription with the Advantage RT-for-PCR kit (BD Biosciences). Real-time PCR was conducted in the LightCycler quantitative PCR system (Roche Molecular Biochemicals) by using the LightCycler-FastStart DNS Master SYBR Green I kit (Roche Molecular Biochemicals). We followed the manufacturer's specification using 4 mM MgCl<sub>2</sub> and 1 pM primers. The primers used are as follows: TNF- $\alpha$ , CTGTGAAGGGAATGGGTGTT (sense) and GGTCACTGTCCCAGCATCTT (antisense); IL-1 $\beta$ , TGAAATGCCACCTTTTGACA (sense) and GTAGCTGCCACAGCTTCTCC (antisense); IL-6, TTCCATCCAGTTGCCTT-CCT (sense) and CAGAATTGCCATTGCAACA (antisense); TGF- $\beta$ , TGCGCTTGACAGA-GATTAAAA (sense) and GCTGAATCGAAAGCCCTGTA (antisense); and HPRT, GTTGGG-TACAGGCCAGACTTTGTGTTG (sense) and GAGGGTAGGCTGGCCTATAGGCT (antisense). Values are presented as the relative amount of transcript of each sample normalized to the housekeeping gene hypoxanthine phosphoribosyltransferase (HPRT).

#### In vitro effect of ghrelin on RAW 264.7 monocytes treated with LPS

To examine the effect of ghrelin on monocytes, RAW 264.7 monocytes (American Type Culture Collection) were suspended in the standard culture medium supplemented with 10% FCS and cultured in 96-well flat bottom plates at  $1 \times 10^5$ /well overnight. Various concentrations of ghrelin ( $10^{-6}$  M,  $10^{-8}$  M,  $10^{-10}$  M) were added to the culture and 1 h later the cells were stimulated with LPS (Sigma-Aldrich) at various doses (0.1, 1, 10  $\mu$ g/ml). After 2 h of incubation at 37°C, supernatants were collected and the levels of TNF- $\alpha$  and IL-6 were detected by using a sandwich ELISA.

#### Isolation of microglial cells from the CNS

The spinal cords were incubated with 35 mg/ml Liberase Blendzyme 3 (Roche Molecular Biochemicals) and 0.1 mg/ml DNaseI (Roche Molecular Biochemicals) in RPMI 1640 medium at 37°C for 30 min. Mononuclear cells were isolated on 30%–80% discontinuous Percoll gradients and were stained with FITC-CD11b mAb, PE-CD45 mAb, and allophycocyanin-CD3 mAb (BD Pharmingen). CD11b<sup>high</sup>CD45<sup>high</sup> macrophage cells, CD11b<sup>int</sup>CD45<sup>int</sup> microglial cells, and CD3<sup>+</sup> T cells were isolated using