

10. Wanner A: **Mucociliary clearance in the trachea.** *Clin Chest Med* 1986, **7**:247-258.
11. Kiyono H, Fukuyama S: **NALT- versus Peyer's-patch-mediated mucosal immunity.** *Nat Rev Immunol* 2004, **4**:699-710.
12. Wu HY, Weiner HL: **Oral tolerance.** *Immunol Res* 2003, **28**:265-284.
13. Mora JR, Iwata M, Eksteen B, Song SY, Junt T, Senman B, Otipoby KL, Yokota A, Takeuchi H, Ricciardi-Castagnoli P *et al.*: **Generation of gut-homing IgA-secreting B cells by intestinal dendritic cells.** *Science* 2006, **314**:1157-1160.
14. Cerutti A: **The regulation of IgA class switching.** *Nat Rev Immunol* 2008, **8**:421-434.
15. Mestecky J, McGhee JR: **Immunoglobulin A (IgA): molecular and cellular interactions involved in IgA biosynthesis and immune response.** *Adv Immunol* 1987, **40**:153-245.
16. Gohda M, Kunisawa J, Miura F, Kagiya Y, Kurashima Y, Higuchi M, Ishikawa I, Ogahara I, Kiyono H: **Sphingosine 1-phosphate regulates the egress of IgA plasmablasts from Peyer's patches for intestinal IgA responses.** *J Immunol* 2008, **180**:5335-5343.
17. Kunkel EJ, Kim CH, Lazarus NH, Vierra MA, Soler D, Bowman EP, Butcher EC: **CCR10 expression is a common feature of circulating and mucosal epithelial tissue IgA Ab-secreting cells.** *J Clin Invest* 2003, **111**:1001-1010.
18. Mestecky J: **The common mucosal immune system and current strategies for induction of immune responses in external secretions.** *J Clin Immunol* 1987, **7**:265-276.
19. Mestecky J, Michalek SM, Moldoveanu Z, Russell MW: **Routes of immunization and antigen delivery systems for optimal mucosal immune responses in humans.** *Behring Inst Mitt* 1997:33-43.
20. Quiding-Jarbrink M, Nordstrom I, Granstrom G, Kilander A, Jertborn M, Butcher EC, Lazarovits AI, Holmgren J, Czerkinsky C: **Differential expression of tissue-specific adhesion molecules on human circulating antibody-forming cells after systemic, enteric, and nasal immunizations. A molecular basis for the compartmentalization of effector B cell responses.** *J Clin Invest* 1997, **99**:1281-1286.
21. Ogra PL: **An introductory overview.** In *Mucosal Vaccines*. Edited by Kiyono H, Ogra PL, McGhee JR. Academic Press, Inc.; 1996: 3-14.
22. Holmgren J, Czerkinsky C: **Mucosal immunity and vaccines.** *Nat Med* 2005, **11**:S45-S53.
23. Cha HR, Ko HJ, Kim ED, Chang SY, Seo SU, Cuburu N, Ryu S, Kim S, Kweon MN: **Mucosa-associated epithelial chemokine/CCL28 expression in the uterus attracts CCR10+ IgA plasma cells following mucosal vaccination via estrogen control.** *J Immunol* 2011, **187**:3044-3052.
- This study demonstrates that intranasal vaccination can induce the migration of antigen-specific IgA-secreting cells into the uterus in a CCL28-dependent manner.
24. Kunisawa J, Nochi T, Kiyono H: **Immunological commonalities and distinctions between airway and digestive immunity.** *Trends Immunol* 2008, **29**:505-513.
25. Johansen FE, Braathen R, Brandtzaeg P: **Role of J chain in secretory immunoglobulin formation.** *Scand J Immunol* 2000, **52**:240-248.
26. Kaetzel CS: **The polymeric immunoglobulin receptor: bridging innate and adaptive immune responses at mucosal surfaces.** *Immunol Rev* 2005, **206**:83-99.
27. Russell MWaK M: **Biological activities.** In *Mucosal Immunology*, edn 3. Edited by Mestecky J, Lamm ME, Strober W, Bienenstock J, McGhee JR, Mayer L. Elsevier; 2005:267-289.
28. Mebius RE: **Organogenesis of lymphoid tissues.** *Nat Rev Immunol* 2003, **3**:292-303.
29. Heritage PL, Brook MA, Underdown BJ, McDermott MR: **Intranasal immunization with polymer-grafted microparticles activates the nasal-associated lymphoid tissue and draining lymph nodes.** *Immunology* 1998, **93**:249-256.
30. Niess JH, Brand S, Gu X, Landsman L, Jung S, McCormick BA, Vyas JM, Boes M, Ploegh HL, Fox JG *et al.*: **CX3CR1-mediated dendritic cell access to the intestinal lumen and bacterial clearance.** *Science* 2005, **307**:254-258.
31. Sung SS, Fu SM, Rose CE Jr, Gaskin F, Ju ST, Beaty SR: **A major lung CD103 (alphaE)-beta7 integrin-positive epithelial dendritic cell population expressing Langerin and tight junction proteins.** *J Immunol* 2006, **176**:2161-2172.
32. Owen RL: **Sequential uptake of horseradish peroxidase by lymphoid follicle epithelium of Peyer's patches in the normal unobstructed mouse intestine: an ultrastructural study.** *Gastroenterology* 1977, **72**:440-451.
33. Park HS, Francis KP, Yu J, Cleary PP: **Membranous cells in nasal-associated lymphoid tissue: a portal of entry for the respiratory mucosal pathogen group A streptococcus.** *J Immunol* 2003, **171**:2532-2537.
34. Siebers A, Finlay BB: **M cells and the pathogenesis of mucosal and systemic infections.** *Trends Microbiol* 1996, **4**:22-29.
35. Wu Y, Wang X, Csencsits KL, Haddad A, Walters N, Pascual DW: **M cell-targeted DNA vaccination.** *Proc Natl Acad Sci U S A* 2001, **98**:9318-9323.
36. Kim DY, Sato A, Fukuyama S, Sagara H, Nagatake T, Kong IG, Goda K, Nochi T, Kunisawa J, Sato S *et al.*: **The airway antigen sampling system: respiratory M cells as an alternative gateway for inhaled antigens.** *J Immunol* 2011, **186**:4253-4262.
- This study demonstrates the presence of novel airway antigen sampling sites—namely respiratory M cells—in the murine nasal passages. The respiratory M cells act independently of nasopharynx-associated lymphoid tissue.
37. Fukuyama S, Hiroi T, Yokota Y, Rennert PD, Yanagita M, Kinoshita N, Terawaki S, Shikina T, Yamamoto M, Kurono Y *et al.*: **Initiation of NALT organogenesis is independent of the IL-7R, LTbetaR, and NIK signaling pathways but requires the Id2 gene and CD3(-)CD4(+)CD45(+) cells.** *Immunity* 2002, **17**:31-40.
38. Teitelbaum R, Schubert W, Gunther L, Kress Y, Macaluso F, Pollard JW, McMurray DN, Bloom BR: **The M cell as a portal of entry to the lung for the bacterial pathogen *Mycobacterium tuberculosis*.** *Immunity* 1999, **10**:641-650.
39. Shinya K, Ebina M, Yamada S, Ono M, Kasai N, Kawaoka Y: **Avian flu: influenza virus receptors in the human airway.** *Nature* 2006, **440**:435-436.
40. Randall TD, Carragher DM, Rangel-Moreno J: **Development of secondary lymphoid organs.** *Annu Rev Immunol* 2008, **26**:627-650.
41. Sun Z, Unutmaz D, Zou YR, Sunshine MJ, Pierani A, Brenner-Morton S, Mebius RE, Littman DR: **Requirement for RORgamma in thymocyte survival and lymphoid organ development.** *Science* 2000, **288**:2369-2373.
42. Yokota Y, Mansouri A, Mori S, Sugawara S, Adachi S, Nishikawa S, Gruss P: **Development of peripheral lymphoid organs and natural killer cells depends on the helix-loop-helix inhibitor Id2.** *Nature* 1999, **397**:702-706.
43. Yoshida H, Honda K, Shinkura R, Adachi S, Nishikawa S, Maki K, Ikuta K, Nishikawa SI: **IL-7 receptor alpha+ CD3(-) cells in the embryonic intestine induces the organizing center of Peyer's patches.** *Int Immunol* 1999, **11**:643-655.
44. Ansel KM, Ngo VN, Hyman PL, Luther SA, Forster R, Sedgwick JD, Browning JL, Lipp M, Cyster JG: **A chemokine-driven positive feedback loop organizes lymphoid follicles.** *Nature* 2000, **406**:309-314.
45. Finke D, Acha-Orbea H, Mattis A, Lipp M, Kraehenbuhl J: **CD4+CD3- cells induce Peyer's patch development: role of alpha4beta1 integrin activation by CXCR5.** *Immunity* 2002, **17**:363-373.
46. Fukuyama S, Nagatake T, Kim DY, Takamura K, Park EJ, Kaisho T, Tanaka N, Kurono Y, Kiyono H: **Cutting edge: uniqueness of lymphoid chemokine requirement for the initiation and**

8 Viral pathogenesis

- maturation of nasopharynx-associated lymphoid tissue organogenesis. *J Immunol* 2006, **177**:4276-4280.**
47. Honda K, Nakano H, Yoshida H, Nishikawa S, Rennert P, Ikuta K, Tamechika M, Yamaguchi K, Fukumoto T, Chiba T *et al.*: **Molecular basis for hematopoietic/mesenchymal interaction during initiation of Peyer's patch organogenesis. *J Exp Med* 2001, **193**:621-630.**
48. Yoshida H, Kawamoto H, Santee SM, Hashi H, Honda K, Nishikawa S, Ware CF, Katsura Y, Nishikawa SI: **Expression of alpha(4)beta(7) integrin defines a distinct pathway of lymphoid progenitors committed to T cells, fetal intestinal lymphotoxin producer, NK, and dendritic cells. *J Immunol* 2001, **167**:2511-2521.**
49. Yoshida H, Naito A, Inoue J, Satoh M, Santee-Cooper SM, Ware CF, Togawa A, Nishikawa S: **Different cytokines induce surface lymphotoxin-alpha on IL-7 receptor-alpha cells that differentially engender lymph nodes and Peyer's patches. *Immunity* 2002, **17**:823-833.**
50. Weih F, Caamano J: **Regulation of secondary lymphoid organ development by the nuclear factor-kappaB signal transduction pathway. *Immunol Rev* 2003, **195**:91-105.**
51. Harmsen A, Kusser K, Hartson L, Tighe M, Sunshine MJ, Sedgwick JD, Choi Y, Littman DR, Randall TD: **Cutting edge: organogenesis of nasal-associated lymphoid tissue (NALT) occurs independently of lymphotoxin-alpha (LT alpha) and retinoic acid receptor-related orphan receptor-gamma, but the organization of NALT is LT alpha dependent. *J Immunol* 2002, **168**:986-990.**
52. Bienenstock J, McDermott MR: **Bronchus- and nasal-associated lymphoid tissues. *Immunol Rev* 2005, **206**:22-31.**
53. Tschernig T, Pabst R: **Bronchus-associated lymphoid tissue (BALT) is not present in the normal adult lung but in different diseases. *Pathobiology* 2000, **68**:1-8.**
54. Tango M, Suzuki E, Gejyo F, Ushiki T: **The presence of specialized epithelial cells on the bronchus-associated lymphoid tissue (BALT) in the mouse. *Arch Histol Cytol* 2000, **63**:81-89.**
55. Moyron-Quiroz JE, Rangel-Moreno J, Kusser K, Hartson L, Sprague F, Goodrich S, Woodland DL, Lund FE, Randall TD: **Role of inducible bronchus associated lymphoid tissue (iBALT) in respiratory immunity. *Nat Med* 2004, **10**:927-934.**
56. Rangel-Moreno J, Carragher DM, de la Luz Garcia-Hernandez M, Hwang JY, Kusser K, Hartson L, Kolls JK, Khader SA, Randall TD: **The development of inducible bronchus-associated lymphoid tissue depends on IL-17. *Nat Immunol* 2011, **12**:639-646.**
- This study demonstrates that, in iBALT formation, IL-17 produced by CD4⁺ T cells promotes lymphotoxin- α -independent expression of CXCL13, which is important for lymphoid follicle formation.
57. Zhu M, Fu Y: **Proinflammatory IL-17 induces iBALT development. *Cell Mol Immunol* 2012, **9**:101-102.**
58. Takeuchi O, Akira S: **Innate immunity to virus infection. *Immunol Rev* 2009, **227**:75-86.**
59. Takeuchi O, Akira S: **Pattern recognition receptors and inflammation. *Cell* 2010, **140**:805-820.**
- This is a useful review for understanding the role of innate pattern-recognition receptors and their signaling pathways in the control of inflammatory responses.
60. Horimoto T, Kawaoka Y: **Influenza: lessons from past pandemics, warnings from current incidents. *Nat Rev Microbiol* 2005, **3**:591-600.**
61. Diebold SS, Kaisho T, Hemmi H, Akira S, Reis e Sousa C: **Innate antiviral responses by means of TLR7-mediated recognition of single-stranded RNA. *Science* 2004, **303**:1529-1531.**
62. Heil F, Hemmi H, Hochrein H, Ampenberger F, Kirschning C, Akira S, Lipford G, Wagner H, Bauer S: **Species-specific recognition of single-stranded RNA via toll-like receptor 7 and 8. *Science* 2004, **303**:1526-1529.**
63. Kato H, Sato S, Yoneyama M, Yamamoto M, Uematsu S, Matsui K, Uematsu T, Takeda K, Fujita T, Takeuchi O *et al.*: **Cell type-specific involvement of RIG-I in antiviral response. *Immunity* 2005, **23**:19-28.**
64. Kato H, Takeuchi O, Sato S, Yoneyama M, Yamamoto M, Matsui K, Uematsu S, Jung A, Kawai T, Ishii KJ *et al.*: **Differential roles of MDA5 and RIG-I helicases in the recognition of RNA viruses. *Nature* 2006, **441**:101-105.**
65. Ichinohe T, Pang IK, Iwasaki A: **Influenza virus activates inflammasomes via its intracellular M2 ion channel. *Nat Immunol* 2010, **11**:404-410.**
- This study demonstrates the mechanism by which influenza virus activates the NLRP3 inflammasome pathway.
66. Gazit R, Gruda R, Elboim M, Arnon TI, Katz G, Achdout H, Hanna J, Qimron U, Landau G, Greenbaum E *et al.*: **Lethal influenza infection in the absence of the natural killer cell receptor gene Ncr1. *Nat Immunol* 2006, **7**:517-523.**
67. Mandelboim O, Lieberman N, Lev M, Paul L, Arnon TI, Bushkin Y, Davis DM, Strominger JL, Yewdell JW, Porgador A: **Recognition of haemagglutinins on virus-infected cells by NKp46 activates lysis by human NK cells. *Nature* 2001, **409**:1055-1060.**
68. Huntington ND, Voshchenrich CA, Di Santo JP: **Developmental pathways that generate natural-killer-cell diversity in mice and humans. *Nat Rev Immunol* 2007, **7**:703-714.**
69. Nogusa S, Ritz BW, Kassim SH, Jennings SR, Gardner EM: **Characterization of age-related changes in natural killer cells during primary influenza infection in mice. *Mech Ageing Dev* 2008, **129**:223-230.**
70. Stein-Streilein J, Guffee J, Fan W: **Locally and systemically derived natural killer cells participate in defense against intranasally inoculated influenza virus. *Reg Immunol* 1988, **1**:100-105.**

Reintroduction of H5N1 highly pathogenic avian influenza virus by migratory water birds, causing poultry outbreaks in the 2010–2011 winter season in Japan

Yoshihiro Sakoda,^{1†} Hiroshi Ito,^{2,3†} Yuko Uchida,^{4†} Masatoshi Okamoto,¹ Naoki Yamamoto,¹ Kosuke Soda,^{1,3} Naoki Nomura,¹ Saya Kuribayashi,¹ Shintaro Shichinohe,¹ Yuji Sunden,⁵ Takashi Umemura,⁵ Tatsufumi Usui,^{3,6} Hiroichi Ozaki,^{3,7} Tsuyoshi Yamaguchi,^{3,6} Toshiyuki Murase,^{3,7} Toshihiro Ito,^{2,3} Takehiko Saito,⁴ Ayato Takada⁸ and Hiroshi Kida^{1,8,9}

Correspondence

Hiroshi Kida
kida@vetmed.hokudai.ac.jp

¹Laboratory of Microbiology, Department of Disease Control, Graduate School of Veterinary Medicine, Hokkaido University, Sapporo 060-0818, Japan

²Laboratory of Veterinary Public Health, Faculty of Agriculture, Tottori University, Tottori 680-8553, Japan

³Avian Zoonosis Research Center, Faculty of Agriculture, Tottori University, Tottori 680-8553, Japan

⁴Research Team for Zoonotic Diseases, National Institute of Animal Health, Tsukuba 305-0856, Japan

⁵Laboratory of Comparative Pathology, Department of Veterinary Clinical Sciences, Graduate School of Veterinary Medicine, Hokkaido University, Sapporo 060-0818, Japan

⁶Laboratory of Veterinary Hygiene, Faculty of Agriculture, Tottori University, Tottori 680-8553, Japan

⁷Laboratory of Veterinary Microbiology, Faculty of Agriculture, Tottori University, Tottori 680-8553, Japan

⁸Research Center for Zoonosis Control, Hokkaido University, Sapporo 001-0020, Japan

⁹Japan Science and Technology Agency (JST), 4-1-8 Honcho, Kawaguchi 332-0012, Japan

H5N1 highly pathogenic avian influenza virus (HPAIV) was reintroduced and caused outbreaks in chickens in the 2010–2011 winter season in Japan, which had been free from highly pathogenic avian influenza (HPAI) since 2007 when HPAI outbreaks occurred and were controlled. On 14 October 2010 at Lake Ohnuma, Wakkanai, the northernmost part of Hokkaido, Japan, H5N1 HPAIVs were isolated from faecal samples of ducks flying from their nesting lakes in Siberia. Since then, in Japan, H5N1 HPAIVs have been isolated from 63 wild birds in 17 prefectures and caused HPAI outbreaks in 24 chicken farms in nine prefectures by the end of March in 2011. Each of these isolates was genetically closely related to the HPAIV isolates at Lake Ohnuma, and those in China, Mongolia, Russia and Korea, belonging to genetic clade 2.3.2.1. In addition, these isolates were genetically classified into three groups, suggesting that the viruses were transmitted by migratory water birds through at least three different routes from their northern territory to Japan. These isolates were antigenic variants, which is consistent with selection in poultry under the immunological pressure induced by vaccination. To prevent the perpetuation of viruses in the lakes where water birds nest in summer in Siberia, prompt eradication of HPAIVs in poultry is urgently needed in Asian countries where HPAI has not been controlled.

Received 14 September 2011

Accepted 20 November 2011

†These authors contributed equally to this work.

Supplementary figures and a supplementary table with the GenBank/EMBL/DDBJ accession numbers of the nucleotide sequences of the representative H5N1 isolates determined in this study are available with the online version of this paper.

INTRODUCTION

Avian influenza caused by infection with H5N1 highly pathogenic avian influenza virus (HPAIV) has spread in poultry in more than 60 countries in Eurasia and Africa since 1996, when the first outbreak occurred at a goose farm in Guangdong province in China (Smith *et al.*, 2006; Xu *et al.*, 1999). H5N1 HPAIV infections have become endemic in several countries and cause accidental transmissions to humans. H5N1 viruses are thus now recognized as one of the most likely candidates for the next pandemic (Li *et al.*, 2004; Peiris *et al.*, 2007). The widespread presence of H5N1 HPAIVs in poultry, especially in domestic free-range-reared ducks, has inevitably resulted in the water-borne transmission of viruses to wild-bird populations, as domestic ducks and geese infected with HPAIV shed progeny virus with their faeces into ponds on farms where migratory water birds visit. In the past, such infections were restricted to wild birds found dead in the vicinity of infected poultry farms, but it is now a concern that infections in wild birds in which HPAIV has caused mild clinical signs (e.g. ducks) could result in the spread of viruses to large areas (Kim *et al.*, 2009; Smith *et al.*, 2009). Infection with HPAIVs in many wild-bird species at two water-bird parks in Hong Kong was reported in 2002 (Ellis *et al.*, 2004), and more significant outbreaks in wild water birds occurred at Lake Qinghai in western China, and at the Khunt and Erkhel Lakes in Mongolia in 2005 (Chen *et al.*, 2005; Sakoda *et al.*, 2010). H5N1 HPAIV infections in poultry and wild birds have now spread in Asia, Europe and Africa, and it has been suggested that the H5N1 virus could spread by migratory water birds to the west and south, as genetically closely related H5N1 viruses (clade 2.2) have been isolated in several countries since 2005 (Monne *et al.*, 2008; Salzberg *et al.*, 2007; Starick *et al.*, 2008).

In Japan, the outbreaks caused by H5N1 HPAIVs occurred in chicken farms in 2004 (Mase *et al.*, 2005) and 2007. The H5N1 HPAIV isolates in 2004 and 2007 were classified genetically into clades 2.5 and 2.2, respectively. Both outbreaks were controlled by the culling of chickens on the farms where the outbreaks occurred (four farms in each year), intensive surveillance and improved biosecurity measures. In addition, H5N1 HPAIVs were isolated from jungle crows, mountain hawk eagles and whooper swans in 2004, 2007 and 2008, respectively (Shivakoti *et al.*, 2010; Tanimura *et al.*, 2006; Uchida *et al.*, 2008). Since then, it has been confirmed that Japan was free from HPAIV infection in poultry and wild birds by intensive surveillance.

H5N1 viruses of clade 2.3.2 were first isolated from ducks, geese and other mammals in China and Vietnam in 2005 (Chen *et al.*, 2006; Robertson *et al.*, 2006). In intensive surveillance studies in China, viruses belonging to clade 2.3.2 have been characterized as the dominant isolates in poultry and wild birds (Ellis *et al.*, 2009; Jiang *et al.*, 2010; Kou *et al.*, 2009; Smith *et al.*, 2009). In the updated unified nomenclature of H5 HPAIVs, recent H5N1 isolates

belonging to clade 2.3.2 were defined as clade 2.3.2.1 (WHO/OIE/FAO H5N1 Evolution Working Group, 2012). H5N1 HPAIVs of clade 2.3.2.1 were isolated from migratory water birds in Japan in 2008, in China in 2009, in Mongolia in 2009 and 2010, in Russia in 2009 and 2010, and in Korea in 2010 and 2011 (Kwon *et al.*, 2011; Li *et al.*, 2011; Sakoda *et al.*, 2010; Sharshov *et al.*, 2010; Uchida *et al.*, 2008). In addition, the infections of chickens and wild birds with HPAIVs belonging to clade 2.3.2.1 have now spread to Europe (Reid *et al.*, 2011). These H5N1 HPAIVs were isolated only from migratory water birds on the way back to their northern territory, and not from those flying to the south from their nesting lakes in Siberia in autumn, suggesting that H5N1 HPAIVs had not dominantly perpetuated at their nesting lakes in Siberia until 2009 (Sakoda *et al.*, 2010; Yamamoto *et al.*, 2011).

On 14 October 2010 at Lake Ohnuma, Wakkanai, in the northernmost part of Hokkaido, Japan, H5N1 HPAIVs were isolated from faecal samples from ducks flying from their nesting lakes in Siberia (Kajihara *et al.*, 2011). Since then, in Japan, H5N1 HPAIVs have been isolated from 63 wild birds, and caused HPAI outbreaks in 24 chicken farms by the end of March. The aim of the present study was to characterize genetically and antigenically the H5N1 viruses isolated from wild birds and chickens in Japan.

RESULTS

Isolation and identification of H5N1 HPAIVs from wild birds and chickens

In the intensive surveillance of HPAIV infection in poultry and wild birds, H5N1 HPAIV had not been isolated from migratory water birds that flew from their nesting lakes in Siberia to Japan until the 2009–2010 winter season (data not shown). In the 2010–2011 winter season, 5591 dead wild birds of ~100 different species were found in Japan. After the isolation of H5N1 HPAIVs from faecal samples of ducks at Lake Ohnuma, Hokkaido (Kajihara *et al.*, 2011), H5N1 viruses were isolated from 63 dead wild birds (63 isolates) and chickens from 24 farms (24 isolates) in Japan (Fig. 1 and Table 1). The multiple basic amino acid sequence (RERRRKR/G), which is a marker of HPAIVs (OIE, 2011), was found at the cleavage site of the deduced amino acid sequence of the haemagglutinin (HA) of all 87 isolates. The pathogenicity of four representative isolates, A/duck/Fukushima/2/2011 (H5N1), A/whooper swan/Hokkaido/4/2011 (H5N1), A/peregrine falcon/Tochigi/15/2011 (H5N1) and A/peregrine falcon/Aomori/7/2011 (H5N1), to chickens was evaluated using an intravenous pathogenicity index (IVPI) test. All chickens inoculated with each virus died within 3 days of inoculation, and IVPI scores ranged from 2.80 to 2.98, which were categorized as HPAIV in chickens. The nucleotide sequences of the representative H5N1 isolates obtained in the present study have been registered in GenBank and are given in Supplementary Table S1 (available in JGV Online).




Fig. 1. H5N1 HPAIV infections in wild birds and chickens in the 2010–2011 winter season in Japan. (a) Geographical location of Japan in Asia and migration routes of wild water birds from Siberia in autumn (arrows). (b) On 14 October 2010 at Lake Ohnuma, Wakkanai, Hokkaido, Japan (red star), H5N1 HPAIVs were isolated from faecal samples from ducks that had flown from their nesting lakes in Siberia (Kajihara *et al.*, 2011). H5N1 HPAIVs were isolated from 63 wild birds in 17 prefectures (red circles) and chickens of 24 farms in nine prefectures (blue circles) by the end of March 2011. Occurrences at different geographical location are indicated by the star or circles, and subsequent cases at the same place are omitted.

Phylogenetic analysis of the H5N1 isolates

For the phylogenetic analysis of HA genes, 30 isolates were selected from 63 isolates of wild birds, and three isolates were also selected from 24 isolates of chickens. The HA genes of the representative 33 H5N1 isolates were analysed by the neighbour-joining method along with those of other HPAIVs isolated recently in Asia (Fig. 2a). The HA genes of the isolates in the 2010–2011 winter season in Japan were closely related to the isolates from poultry or wild birds in China, Mongolia, Russia and Korea in 2009–2011, and were classified into clade 2.3.2.1. These isolates in Japan were divided into three groups (A–C) based on the results of phylogenetic analysis (Fig. 2b and Table 1). This classification by the neighbour-joining method was supported by analyses using maximum-likelihood and maximum-parsimony methods with 1000 bootstrap replicates (data not shown). In particular, A/duck/Hokkaido/WZ83/2010 (H5N1), the first isolate at Lake Ohnuma, Wakkanai, Hokkaido, in October 2010, indicated with an asterisk in Fig. 2(b), was classified into group C, and not group A containing subsequent isolates from Hokkaido (A/pintail/Hokkaido/1/2011, A/greater scaup/Hokkaido/2/2011, A/whooper swan/Hokkaido/3/2011, A/whooper swan/Hokkaido/4/2011, A/whooper swan/Hokkaido/6/2011, A/whooper swan/Hokkaido/13-21/2011, A/whooper swan/Hokkaido/13-27/2011, A/greater scaup/Hokkaido/28/2011 and A/whooper swan/Hokkaido/A13/2011) and Fukushima (A/tufted duck/Fukushima/2/2011, A/tufted duck/Fukushima/4/2011, A/tufted duck/Fukushima/5/2011, A/tufted duck/Fukushima/7/2011, A/tufted

duck/Fukushima/16/2011 and A/tundra swan/Fukushima/207/2011). All occurrences in Hokkaido after January 2011 were only in the eastern Kushiro area, 350 km south-east of Lake Ohnuma, Wakkanai (Fig. 1b). The cases in the Kushiro area in Hokkaido started in mid-January 2011, and ended in mid-February 2011 (Table 1). The isolates from wild birds in this area were genetically closely related to each other and were classified into group A (Fig. 2b). In group B, all viruses were isolated only from western areas (Aichi, Kyoto, Hyogo, Tokushima and Shimane). In group C, viruses were isolated from the whole of the country (Hokkaido, Aomori, Tochigi, Aichi, Mie, Tottori, Yamaguchi, Kochi, Oita, Nagasaki, Miyazaki and Kagoshima). In addition, A/mandarin duck/Kochi/3901C005/2011 (H5N1) isolated in Kochi Prefecture, in south-western Japan, belonging to group C, had the highest nucleotide identity of the HA gene with A/mallard duck/Korea/W401/2011 (H5N1) and A/mandarin duck/Korea/K10-515/2011 (H5N1) isolated in Korea in the 2010–2011 winter season (Kwon *et al.*, 2011).

To assess the genetic relationship of the HPAIVs in gene segments other than the HA, the nucleotide sequences of a representative 30 H5N1 isolates were analysed and compared with those of other H5N1 HPAIVs (see Supplementary Figs S1–S7, available in JGV Online). These viruses were isolates from wild birds and were used for the phylogenetic tree analysis of the HA gene. The genes of these isolates were closely related to each other, and no genetic reassortment with other previous HPAIVs was identified. Each of the polymerase subunit (PB2 and PB1), nucleoprotein (NP),

Table 1. Cases of infection with H5N1 HPAIVs in Japan in the 2010–2011 winter season

Information about the cases from chicken farms is underlined. NT, Not tested.

Area	Prefecture	Date of reports	Species of birds*	Genetic subgroup of representative isolates†
Hokkaido	Hokkaido	14 Oct 2010‡, 12, 17, 18, 19 and 28 Jan 2011, 3, 7 and 17 Feb 2011	Duck (2)‡, whooper swan (6), greater scaup (2), pintail (1)	A, C‡
Honshu	Aomori	10 Mar (2011)	Peregrine falcon (1)	C
	Fukushima	4, 5, 7, 10 and 23 Jan 2011, 10 Feb 2011	Tufted duck (5), tundra swan (1)	A
	Tochigi	14 Feb 2011, 25 Mar 2011	Peregrine falcon (1), goshawk (1)	C
	Chiba	<u>12 and 16 Mar 2011</u>	<u>Chicken (2)</u>	NT
	Aichi	17 Feb 2011 <u>27 Jan 2011, 14 Feb 2011</u>	Peregrine falcon (1) <u>Chicken (2)</u>	B, C
	Toyama	16 Dec 2010	Mute swan (1)	NT
	Mie	<u>15 and 26 Feb 2011</u>	<u>Chicken (2)</u>	C
	Wakayama	<u>15 Feb 2011</u>	<u>Chicken (1)</u>	NT
	Kyoto	16 Feb 2011	Peregrine falcon (1)	B
	Nara	<u>28 Feb 2011</u>	<u>Chicken (1)</u>	NT
	Hyogo	12 and 25 Jan 2011, 11 and 22 Feb 2011	Common pochard (1), little grebe (1), mute swan (1), great crested grebe (1)	B
	Tottori	4 Dec 2010, 19 and 24 Jan 2011, 1, 3 and 6 Feb 2011	Tundra swan (1), black-headed gull (1), tufted duck (2), common pochard (1), peregrine falcon (1)	C
	Shimane	14 Jan 2011, 1 and 8 Feb 2011 <u>29 Nov 2010</u>	Tufted duck (4), common pochard (1) <u>Chicken (1)</u>	B
	Shikoku	Yamaguchi	6 and 9 Feb 2011	Tufted duck (1), black swan (1)
Tokushima		8 Feb 2011	Ural owl (1)	B
Kochi		26 Jan 2011	Mandarin duck (1)	C
Kyushu	Nagasaki	31 Jan 2011, 4 and 12 Feb 2011	Mandarin duck (3), peregrine falcon (1)	C
	Oita	7, 8, 9 and 15 Feb 2011 <u>2 Feb 2011</u>	Mandarin duck (4), grey heron (1) <u>Chicken (1)</u>	C
	Miyazaki	1, 2, 8, 11, 14, 15 and 18 Feb 2011 <u>22, 24, 27, 28, 29 and 30 Jan 2011, 1, 4, 5, 6, 7 and 17 Feb 2011, 5 Mar 2011</u>	Mandarin duck (3), peregrine falcon (3), little grebe (1) <u>Chicken (13)</u>	C
	Kagoshima	19, 20, 21 and 24 Dec 2010, 13 Feb 2011 <u>26 Jan 2011</u>	Hooded crane (7) <u>Chicken (1)</u>	C

*The number of dead wild birds or outbreaks in chicken farms is shown in parentheses.

†Based on the phylogenetic tree of the HA gene shown in Fig. 1.

‡Viruses were isolated from faecal samples (Kajihara *et al.*, 2011).

neuraminidase (NA) and matrix (M) genes of the isolates was divided into three genetic groups, corresponding to the classification of the HA genes (groups A–C), although a few isolates were not divided into these groups (Supplementary Figs S1–S5). Because the sequence identities of the polymerase subunit (PA) and non-structural (NS) genes were so high, the genes of these isolates were not classified completely into groups A, B and C (Supplementary Figs S6 and S7).

Antigenic analysis of the HA of the H5N1 HPAIV isolates

The HAs of H5N1 isolates were analysed antigenically using a panel of mAbs recognizing six different epitopes on the HA

of A/duck/Pennsylvania/10218/84 (H5N2) (Okamatsu *et al.*, 2010; Soda *et al.*, 2008; Yamamoto *et al.*, 2011) (Table 2). Each of the non-pathogenic avian influenza viruses (NPAIVs) isolated from migratory ducks in Mongolia and Hokkaido in 2000–2010 bound to all the mAbs used in the present study. Each of the H5N1 HPAIVs isolated before 2005 [A/Hong Kong/483/1997 (H5N1), A/Vietnam/1194/2004 (H5N1), A/chicken/Yamaguchi/7/2004 (H5N1) and A/whooper swan/Mongolia/3/2005 (H5N1)] bound to most mAbs; however, each of the H5N1 viruses belonging to genetic clade 2.3.2.1, including two strains isolated in the present study and A/duck/Hokkaido/WZ83/2010 (H5N1) isolated at Lake Ohnuma, Wakkanai, bound only to mAb D101/1 (Table 2).




Fig. 2. Phylogenetic trees of HA genes of the isolates in the 2010–2011 winter season in Japan. (a) Phylogenetic tree of H5 avian influenza viruses. The unified nomenclature of the A/goose/Guangdong/1/1996 lineage of Eurasian HPAIVs was based on the homology of the HA gene and classified into ten distinct clades (clades 0–9) containing second-order (or third-order) clades proposed by the WHO/OIE/FAO H5N1 Evolution Working Group (2008, 2009). Recently, a new classification was proposed by the same group (WHO/OIE/FAO H5N1 Evolution Working Group, 2012) and 2.3.2.1 is one of the new nomenclature systems. The H5N1 HPAIVs isolated in this study were classified into clade 2.3.2.1 with other recent isolates in Asia from 2007 onward. A/mallard/Hokkaido/24/09 (H5N1) is indicated as representative strain of NPAIV isolated from water birds and its HA gene was classified into the Eurasian lineage (Yamamoto *et al.*, 2011). The HA genes of A/chicken/Pennsylvania/1/1983 (H5N2) and A/chicken/Ibaraki/1/2005 (H5N2) belong to the North American lineage. The individual clade of each isolate is shown in parentheses. (b) Phylogenetic trees of the HA genes of H5N1 HPAIVs including the isolates in the 2010–2011 winter season in Japan. To assess the genetic relationships among H5 avian influenza virus isolates, the nucleotide sequences of the HA gene of each isolate in the present study were compared with those of recent isolates in Asia in 2007–2011 belonging to genetic clade 2.3.2.1. Phylogenetic trees were constructed by the neighbour-joining method and bootstrap testing (1000 replicates). The phylogenetic tree was rooted to A/whooper swan/Hokkaido/1/2008 (H5N1). The HA genes of the recent isolates in this study (shaded) were divided into three genetic groups (A–C). A/duck/Hokkaido/WZ83/2010 (H5N1) HPAIV, isolated from faecal samples on 14 October 2010 at Lake Ohnuma, Hokkaido, Japan (Kajihara *et al.*, 2011), is indicated with an asterisk. Isolates from Korea, Russia, Mongolia, China, Laos and Vietnam in 2007–2011 are underlined. Horizontal distances (bars) are proportional to the minimum number of nucleotide differences required to join nodes and sequences. HA and NA subtypes have been left out of the names of the H5N1 viruses. Ck, Chicken.

Table 2. Antigenic analyses of H5 influenza viruses

Virus*	Clade	mAb†						Polyclonal antibody HI titre‡		
		I (88)	II (145)	III (157)	IV (168)		V (169)	VI (205)	Mal/ Hok/09 (H5N1)	Ws/ Hok/08 (H5N1)
		D101/1	A310/39	64/1	B9/5	B220/1	B59/5	25/2		
NPAIV										
Dk/Pennsylvania/ 10218/1984 (H5N2)	–	+	+	+	+	+	+	+	1280	80
Dk/Mongolia/54/2001 (H5N2)	–	+	+	+	+	+	+	+	640	80
Dk/Hokkaido/167/2007 (H5N3)	–	+	+	+	+	+	+	+	1280	160
Dk/Hokkaido/WZ21/ 2008 (H5N2)	–	+	+	+	+	+	+	+	2560	80
Mal/Hokkaido/24/2009 (H5N1)	–	+	+	+	+	+	+	+	<u>1280</u>	160
Dk/Hokkaido/101/2010 (H5N2)	–	+	+	+	+	+	+	+	640	80
HPAIV										
Hong Kong/483/1997 (H5N1)	0	–	+	+	+	+	+	+	1280	320
Vietnam/1194/2004 (H5N1)	1	+	+	+	+	+	–	+	640	640
Ck/Yamaguchi/7/2004 (H5N1)	2.5	–	+	+	+	+	–	+	1280	1280
Ws/Mongolia/3/2005 (H5N1)	2.2	+	–	+	+	+	–	+	320	640
Ws/Hokkaido/1/2008 (H5N1)	2.3.2.1	+	–	–	–	–	–	–	40	<u>1280</u>
Ws/Mongolia/6/2009 (H5N1)	2.3.2.1	+	–	–	–	–	–	–	80	1280
Ws/Mongolia/1/2010 (H5N1)	2.3.2.1	+	–	–	–	–	–	–	80	640
Dk/Hokkaido/WZ83/ 2010 (H5N1)	2.3.2.1	+	–	–	–	–	–	–	40	320
Ws/Hokkaido/4/2011 (H5N1)	2.3.2.1	+	–	–	–	–	–	–	40	320
Pf/Aomori/7/2011 (H5N1)	2.3.2.1	+	–	–	–	–	–	–	40	320

*Viruses indicated in bold are isolates from the 2010–2011 winter season in Japan. Dk, Duck; Mal, mallard; Ck, chicken; Ws, whooper swan; Pf, peregrine falcon.

†Reactivity of mAbs against the HA of A/duck/Pennsylvania/10218/1984 (H5N2) to the representative H5 viruses (I–VI) were compared in fluorescent antibody methods. The location of amino acid substitutions in antigenic variants selected in the presence of the respective mAbs (Soda *et al.*, 2008) is indicated in parentheses.

‡Haemagglutination inhibition (HI) titres of hyperimmunized polyclonal antibodies against representative H5 viruses were measured. Homologous combination between virus and polyclonal antibody is indicated by underlining.

These H5N1 isolates were also analysed antigenically using hyperimmunized chicken antisera to A/mallard/Hokkaido/24/2009 (H5N1) and A/whooper swan/Hokkaido/1/2008 (H5N1) (Table 2). A/mallard/Hokkaido/24/2009 (H5N1) was isolated from a faecal sample, and the antigenicity and pathogenicity of this isolate in chickens were similar to those of other H5 NPAIVs isolated from migratory ducks

(Yamamoto *et al.*, 2011). The reactivity of the present H5N1 isolates in Japan with the antiserum to A/mallard/Hokkaido/24/2009 (H5N1) was quite low. In contrast, the reactivity of these H5N1 isolates with antiserum to A/whooper swan/Hokkaido/1/2008 (H5N1) was comparatively high. These results indicated that the HAs of H5N1 isolates in the 2010–2011 winter season in Japan were

antigenically distinct from H5 NPAIVs and HPAIVs isolated before 2005.

DISCUSSION

In October 2010, H5N1 viruses were isolated from faecal samples of ducks at Lake Ohnuma, Wakkanai, Hokkaido, on their way south from their nesting lakes in Siberia (Kajihara *et al.*, 2011). Since then, nationwide H5N1 HPAIV infections in wild birds and chickens have occurred in Japan, and 63 and 24 isolates were identified from wild birds and chickens, respectively. The present results indicate that the viruses isolated from wild birds and chickens from November 2010 onward were genetically related to the isolates from migratory ducks at Lake Ohnuma, Wakkanai, in October 2010. In Hokkaido, H5N1 viruses were isolated in two areas, Wakkanai and Kushiro (Fig. 1b). A/duck/Hokkaido/WZ83/2010 (H5N1), the first isolate at Lake Ohnuma, Wakkanai, was identified as a member of genetic group C, and not group A containing subsequent isolates in Kushiro in January and February 2011. Based on the genetic analysis, A/duck/Hokkaido/WZ83/2010 (H5N1) was closely related to A/tundra swan/Tottori/12-002/2010 (H5N1) belonging to group C. The isolates of group C were detected over the whole of the country, and some isolates of group C had the highest nucleotide identity with that of wild ducks in Korea (Kwon *et al.*, 2011). By contrast, the isolates of group B were detected only in the western area. Wild water birds start their migration from their nesting lakes in the northern territory to the south in the middle of August. The migratory routes of water birds are from Siberia to northern Japan via the Kamchatka Peninsula or Sakhalin Island, and to southern Japan via the Korean Peninsula or the coast of north-eastern China (Fig. 1a). Our results indicated that the viruses circulating in different populations of wild migratory birds at their nesting lakes in Siberia in summer were transmitted through at least three different routes via China, Korea and Russia to Japan in the 2010–2011 winter season. Further virus spread then occurred in wild birds at the resting lakes of birds in Japan by water-borne transmission or predation of carcasses. Taken together, these results raise the possibility that H5N1 HPAIVs were perpetuated at the nesting lakes in Siberia before the migration of water birds to Japan.

Concerning the origin of these H5N1 viruses, the HA genes of isolates from chickens and wild birds in China (Jiang *et al.*, 2010; Li *et al.*, 2011) and from wild birds in Mongolia and Russia in 2009 and 2010 (Sakoda *et al.*, 2010; Sharshov *et al.*, 2010) were closely related to those of the present isolates in Japan. The isolates in Laos in 2010 were released recently in the public database (GenBank accession no. CY098351), although epidemiological information is not available. The season of isolation of these viruses from wild birds in China, Mongolia and Russia in 2009 was May to July, the period when migratory water birds return to their nesting lakes in Siberia. As Japan and Mongolia are located

on the flyways of migratory water birds that fly from their nesting lakes in Siberia to the south in autumn, intensive surveillance of avian influenza has been performed in Hokkaido, Japan and Mongolia every year since 1996. No HPAIV was found in a total of 634 virus isolates from 13 740 faecal samples of migratory water birds until 2009 (Sakoda *et al.*, 2010; Yamamoto *et al.*, 2011). These results suggested that the origin of the viruses isolated from wild birds in China, Mongolia and Russia in 2009 was poultry in China, and that these viruses did not perpetuate at their nesting areas in Siberia until 2009. The isolation of H5N1 HPAIVs in the spring of 2010 in Mongolia and Russia demonstrated that virus spread from poultry to wild birds occurred again in China and that H5N1 HPAIVs have circulated in wild water birds since last summer at their nesting lakes in Siberia. These viruses have been maintained in wild migratory bird populations and were brought to Japan in the 2010–2011 winter season. To clarify whether H5N1 HPAIV has been perpetuated dominantly at their nesting lakes in Siberia and viruses are brought by migratory birds from Siberia to the south in autumn, intensive surveillance of avian influenza in migratory birds should be strengthened.

HPAIVs are not under immunological selection pressure in the non-vaccinated chicken populations as HPAIV causes acute infection and death in chickens. The generation of escape mutants against H5 HPAIV was first observed in the follow-up phase of H5N2 HPAIV outbreaks in Mexico in the 1990s (Lee *et al.*, 2004). As vaccine use for poultry has increased in several countries, antigenic variants have been selected in H5N1 HPAIVs under immunological selection pressure (Cattoli *et al.*, 2011; Chen, 2009; Grund *et al.*, 2011). The present results support the findings that H5N1 viruses belonging to clade 2.3.2.1 were antigenically distinct from other HPAIVs and NPAIVs of H5 subtype (Okamatsu *et al.*, 2010; Smith *et al.*, 2009). The vaccination was applied based on the optimistic expectation of preventing H5N1 influenza virus infection in poultry and humans; however, several countries using vaccines against H5 HPAIV have not yet eliminated viruses in poultry because the efficacy of the vaccine against HPAI is limited to suppress virus replication, and does not confer immunity to prevent infection with the virus. It is reasonable to argue that vaccination of poultry results in the selection of antigenic variants and that the vaccine does not confer immunity against antigenic variants for humans and animals. To stop infection with H5 HPAIV in poultry, thorough culling of infected birds must be carried out worldwide.

In the 2010–2011 winter season in Japan, outbreaks of H5N1 HPAIV infection in chicken farms were sporadic, except in Miyazaki Prefecture (13 cases), although a large number of infections in wild birds occurred and the natural environment was contaminated with H5N1 HPAIVs throughout the country. In Japan, each of the outbreaks in poultry was controlled by culling, intensive surveillance, improved biosecurity measures and compensation, without the use of a vaccine, and ended in March 2011. H5N1

HPAIV strains have persisted throughout the world for more than 15 years, and antigenic variants have been selected because some countries use vaccines for the control of HPAIV infection. In the chickens vaccinated against HPAIV, it is hard to find infected birds because they do not show clinical signs, despite shedding of the virus. As a result, HPAIV has returned to migratory water birds from domestic poultry, and many feral water birds have died on the way back to their northern territory in Siberia in spring. Some migratory water birds infected with the virus must have returned to their nesting lakes in Siberia and then disseminated the virus to other birds though water-borne transmission at their nesting lakes. To prevent the perpetuation of HPAIVs among migratory water birds at their nesting lakes in Siberia, HPAIVs should be contained within poultry in Asia. Thus, we strongly recommend that a stamping-out strategy is the only way to achieve prompt eradication of H5N1 HPAIV and that vaccination may be an optional tool for the control of HPAI in addition to the stamping-out policy. Otherwise, disasters will occur every year throughout Asian countries.

METHODS

Viruses. The H5N1 viruses isolated in the present study and the reference H5 viruses shown in Table 2 were propagated in 10-day-old embryonated chicken eggs. As reference strains, H5 NPAIVs isolated from the faecal material of migratory ducks (Yamamoto *et al.*, 2011) and the H5N1 HPAIVs shown in Table 2 (Kajihara *et al.*, 2011; Mase *et al.*, 2005; Muramoto *et al.*, 2006; Okamatsu *et al.*, 2010; Sakoda *et al.*, 2010; Suarez *et al.*, 1998) were used for antigenic analyses.

Isolation and identification of viruses. Virus isolation was carried out from faecal samples, tracheal and cloacal swabs, or homogenates of the tissues of wild birds and chickens throughout the year. Faecal samples were mixed with transport medium containing minimum essential medium (Nissui), 10 000 U penicillin G (Meiji Seika) ml⁻¹, 10 mg streptomycin (Meiji Seika) ml⁻¹, 0.3 mg gentamicin (Merck) ml⁻¹, 250 U nystatin (Sigma) ml⁻¹ and 0.5% BSA fraction V (Roche) to yield a 10–20% suspension. Tracheal and cloacal swabs were mixed with 2 ml transport medium. Organ tissue was homogenized with transport medium to yield a 10% suspension. Samples from wild birds and chickens were inoculated into the allantoic cavities of 10-day-old embryonated chicken eggs, and the subtypes of the HA and NA of influenza virus isolates were identified by haemagglutination inhibition (HI) and neuraminidase inhibition tests, respectively, according to a standard protocol (OIE, 2011).

H5N1 HPAIVs were isolated from 17 species of dead or diseased wild birds found at the waterside of their resting areas and in the gardens of private houses from November 2010 to March 2011 (Table 1): whooper swan (*Cygnus cygnus*), greater scaup (*Aythya marila*), pintail (*Anas acuta*), peregrine falcon (*Falco peregrinus*), tufted duck (*Aythya fuligula*), mute swan (*Cygnus olor*), common pochard (*Aythya ferina*), little grebe (*Tachybaptus ruficollis*), great crested grebe (*Podiceps cristatus*), tundra swan (*Cygnus columbianus*), black-headed gull (*Larus ridibundus*), black swan (*Cygnus atratus*), ural owl (*Strix uralensis*), mandarin duck (*Aix galericulata*), grey heron (*Ardea cinerea*), hooded crane (*Grus monacha*) and goshawk (*Accipiter gentilis*).

Experimental infection of chickens with H5N1 isolates. To assess the pathogenicity of the representative H5N1 virus isolates, A/duck/Fukushima/2/2011 (H5N1), A/whooper swan/Hokkaido/4/2011

(H5N1), A/peregrine falcon/Tochigi/15/2011 (H5N1) and A/peregrine falcon/Aomori/7/2011 (H5N1) were inoculated intravenously into 4–6-week-old chickens (*Gallus gallus*) for an IVPI test according to a standard protocol (OIE, 2011). Each bird was housed in a self-contained isolator unit (Tokuiwa Kagaku) at a Biosafety Level 3 facility at Hokkaido University, Japan.

Sequencing and phylogenetic analysis. For the genetic analysis, 30 isolates were selected from 63 isolates of wild birds, and three isolates were also selected from 24 isolates of chickens. Viral RNA was extracted from the allantoic fluid of embryonated chicken eggs using TRIzol LS reagent (Invitrogen) and reverse transcribed with the Uni12 primer (Hoffmann *et al.*, 2001) and Moloney murine leukemia virus reverse transcriptase (Invitrogen). The full-length or partial sequence of each gene segment was amplified by PCR with gene-specific primer sets reported previously (Hoffmann *et al.*, 2001) or designed exclusively in the present study. The sequences of primers designed in the present study were: PB2-826F: 5'-GTTAGGAGG-CAACAGTATCAG-3', PB2-2135R: 5'-TCATTGATGCTCAATGCC-GG-3', PB1-547F: 5'-ACACATTTCCAGAGAAAGAG-3', PB1-2128R: 5'-TCCACCATGCTAGAAATCCC-3', PA-38F: 5'-GTGCGACAATG-CTTCAATCC-3', PA-1372R: 5'-CCTGCAATGGGATACTTCCGC-3', NP-57F: 5'-TGGAACTGGTGGAGAACGC-3', NP-1456R: 5'-TTGTCTCCGAAGAAATAAGA-3', M-19F: 5'-GTCGAAACGTAC-GTTCTCTC-3', M-853R: 5'-GAATCCACAATATCAAGTGCAAG-3' and NS-848R: 5'-TCATTAATAAGCTGGAACG-3'. Direct sequencing of each gene segment was performed using a 3130 or 3500 Genetic Analyzer (Applied Biosystems). To assess the genetic relationship among influenza virus isolates, nt 34–1019 (986 bp) of HA, nt 197–1206 (1010 bp) of NA, nt 1017–1929 (913 bp) of PB2, nt 1064–1657 (594 bp) of PB1, nt 269–1218 (950 bp) of PA, nt 760–1329 (570 bp) of NP, nt 97–771 (675 bp) of M and nt 73–750 (678 bp) of NS of isolates in the present study were compared with those of other recent H5N1 isolates in Asia. For the NA and internal genes, reference strains of each genotype according to a previous report (Duan *et al.*, 2008) were included. Phylogenetic trees were constructed by the neighbour-joining method (Saitou & Nei, 1987) using MEGA 5 software (<http://www.megasoftware.net/>).

Antigenic analysis. The antigenic properties of the representative H5 viruses [A/duck/Hokkaido/WZ83/2010 (H5N1), A/whooper swan/Hokkaido/4/2011 (H5N1) and A/peregrine falcon/Aomori/7/2011 (H5N1)] were compared with those of the reference H5 viruses by a fluorescent antibody method using mAbs against H5 HA (Soda *et al.*, 2008). Madin–Darby canine kidney cells infected with H5 influenza viruses were fixed with cold 100% acetone at 8 h post-inoculation. The reactivity patterns of the H5 viruses with mAbs were investigated with an FITC-conjugated goat anti-mouse IgG (MP Biomedicals) using a fluorescence microscope (Axiovert 200; Carl Zeiss).

The antigenic properties of the representative H5 viruses were also assessed using hyperimmunized chicken antisera against A/mallard/Hokkaido/24/2009 (H5N1) and A/whooper swan/Hokkaido/1/2008 (H5N1) by an HI test according to a standard protocol (OIE, 2011). HI titres were expressed as the reciprocals of the highest serum dilutions that showed complete HI.

ACKNOWLEDGEMENTS

We deeply appreciate the kind cooperation of the Ministry of Environment and Ministry of Agriculture, Forestry and Fisheries, Government of Japan. This study was supported by Strategic Funds for the Promotion of Science and Technology (2011–2013), Japan. The present work was supported in part by the Japan Initiative for Global Research Network on Infectious Diseases (J-GRID) and Japan Science and Technology Agency Basic Research Programs.

REFERENCES

- Cattoli, G., Milani, A., Temperton, N., Zecchin, B., Buratin, A., Molesti, E., Aly, M. M., Arafa, A. & Capua, I. (2011). Antigenic drift in H5N1 avian influenza virus in poultry is driven by mutations in major antigenic sites of the hemagglutinin molecule analogous to those for human influenza virus. *J Virol* **85**, 8718–8724.
- Chen, H. (2009). Avian influenza vaccination: the experience in China. *Rev Sci Tech* **28**, 267–274.
- Chen, H., Smith, G. J., Zhang, S. Y., Qin, K., Wang, J., Li, K. S., Webster, R. G., Peiris, J. S. & Guan, Y. (2005). Avian flu: H5N1 virus outbreak in migratory waterfowl. *Nature* **436**, 191–192.
- Chen, H., Smith, G. J., Li, K. S., Wang, J., Fan, X. H., Rayner, J. M., Vijaykrishna, D., Zhang, J. X., Zhang, L. J. & other authors (2006). Establishment of multiple sublineages of H5N1 influenza virus in Asia: implications for pandemic control. *Proc Natl Acad Sci U S A* **103**, 2845–2850.
- Duan, L., Bahl, J., Smith, G. J., Wang, J., Vijaykrishna, D., Zhang, L. J., Zhang, J. X., Li, K. S., Fan, X. H. & Cheung, C. (2008). The development and genetic diversity of H5N1 influenza virus in China, 1996–2006. *Virology* **380**, 243–254.
- Ellis, T. M., Bousfield, R. B., Bissett, L. A., Dyrting, K. C., Luk, G. S., Tsim, S. T., Sturm-Ramirez, K., Webster, R. G., Guan, Y. & Malik Peiris, J. S. (2004). Investigation of outbreaks of highly pathogenic H5N1 avian influenza in waterfowl and wild birds in Hong Kong in late 2002. *Avian Pathol* **33**, 492–505.
- Ellis, T. M., Dyrting, K. C., Wong, C. W., Chadwick, B., Chan, C., Chiang, M., Li, C., Li, P., Smith, G. J. & other authors (2009). Analysis of H5N1 avian influenza infections from wild bird surveillance in Hong Kong from January 2006 to October 2007. *Avian Pathol* **38**, 107–119.
- Grund, C., Abdelwhab, S. M., Arafa, A. S., Ziller, M., Hassan, M. K., Aly, M. M., Hafez, H. M., Harder, T. C. & Beer, M. (2011). Highly pathogenic avian influenza virus H5N1 from Egypt escapes vaccine-induced immunity but confers clinical protection against a heterologous clade 2.2.1 Egyptian isolate. *Vaccine* **29**, 5567–5573.
- Hoffmann, E., Stech, J., Guan, Y., Webster, R. G. & Perez, D. R. (2001). Universal primer set for the full-length amplification of all influenza A viruses. *Arch Virol* **146**, 2275–2289.
- Jiang, W.-M., Liu, S., Chen, J., Hou, G.-Y., Li, J.-P., Cao, Y.-F., Zhuang, Q.-Y., Li, Y., Huang, B.-X. & Chen, J.-M. (2010). Molecular epidemiological surveys of H5 subtype highly pathogenic avian influenza viruses in poultry in China during 2007–2009. *J Gen Virol* **91**, 2491–2496.
- Kajihara, M., Matsuno, K., Simulundu, E., Muramatsu, M., Noyori, O., Manzoor, R., Nakayama, E., Igarashi, M., Tomabeche, D. & other authors (2011). An H5N1 highly pathogenic avian influenza virus that invaded Japan through waterfowl migration. *Jpn J Vet Res* **59**, 89–100.
- Kim, J. K., Negovetich, N. J., Forrest, H. L. & Webster, R. G. (2009). Ducks: the “Trojan horses” of H5N1 influenza. *Influenza Other Respir Viruses* **3**, 121–128.
- Kou, Z., Li, Y., Yin, Z., Guo, S., Wang, M., Gao, X., Li, P., Tang, L., Jiang, P. & other authors (2009). The survey of H5N1 flu virus in wild birds in 14 Provinces of China from 2004 to 2007. *PLoS ONE* **4**, e6926.
- Kwon, H.-I., Song, M.-S., Pascua, P. N., Baek, Y. H., Lee, J. H., Hong, S.-P., Rho, J.-B., Kim, J.-K., Poo, H. & other authors (2011). Genetic characterization and pathogenicity assessment of highly pathogenic H5N1 avian influenza viruses isolated from migratory wild birds in 2011, South Korea. *Virus Res* **160**, 305–315.
- Lee, C.-W., Senne, D. A. & Suarez, D. L. (2004). Effect of vaccine use in the evolution of Mexican lineage H5N2 avian influenza virus. *J Virol* **78**, 8372–8381.
- Li, K. S., Guan, Y., Wang, J., Smith, G. J., Xu, K. M., Duan, L., Rahardjo, A. P., Puthavathana, P., Buranathai, C. & other authors (2004). Genesis of a highly pathogenic and potentially pandemic H5N1 influenza virus in eastern Asia. *Nature* **430**, 209–213.
- Li, Y., Liu, L., Zhang, Y., Duan, Z., Tian, G., Zeng, X., Shi, J., Zhang, L. & Chen, H. (2011). New avian influenza virus (H5N1) in wild birds, Qinghai, China. *Emerg Infect Dis* **17**, 265–267.
- Mase, M., Tsukamoto, K., Imada, T., Imai, K., Tanimura, N., Nakamura, K., Yamamoto, Y., Hitomi, T., Kira, T. & other authors (2005). Characterization of H5N1 influenza A viruses isolated during the 2003–2004 influenza outbreaks in Japan. *Virology* **332**, 167–176.
- Monne, I., Fusaro, A., Al-Blawi, M. H., Ismail, M. M., Khan, O. A., Dauphin, G., Tripodi, A., Salviato, A., Marangon, S. & other authors (2008). Co-circulation of two sublineages of HPAI H5N1 virus in the Kingdom of Saudi Arabia with unique molecular signatures suggesting separate introductions into the commercial poultry and falconry sectors. *J Gen Virol* **89**, 2691–2697.
- Muramoto, Y., Le, T. Q., Phuong, L. S., Nguyen, T., Nguyen, T. H., Sakai-Tagawa, Y., Iwatsuki-Horimoto, K., Horimoto, T., Kida, H. & Kawaoka, Y. (2006). Molecular characterization of the hemagglutinin and neuraminidase genes of H5N1 influenza A viruses isolated from poultry in Vietnam from 2004 to 2005. *J Vet Med Sci* **68**, 527–531.
- OIE (2011). Avian influenza. In *Manual of Diagnostic Tests and Vaccines for Terrestrial Animals* 2011, Chapter 2.3.4. Paris: The World Organization for Animal Health http://www.oie.int/eng/normes/mmanual/A_summry.htm
- Okamatsu, M., Tanaka, T., Yamamoto, N., Sakoda, Y., Sasaki, T., Tsuda, Y., Isoda, N., Kokumai, N., Takada, A. & other authors (2010). Antigenic, genetic, and pathogenic characterization of H5N1 highly pathogenic avian influenza viruses isolated from dead whooper swans (*Cygnus cygnus*) found in northern Japan in 2008. *Virus Genes* **41**, 351–357.
- Peiris, J. S., de Jong, M. D. & Guan, Y. (2007). Avian influenza virus (H5N1): a threat to human health. *Clin Microbiol Rev* **20**, 243–267.
- Reid, S. M., Shell, W. M., Barboi, G., Onita, I., Turcitu, M., Cioranu, R., Marinova-Petkova, A., Goujgoulova, G., Webby, R. J. & other authors (2011). First reported incursion of highly pathogenic notifiable avian influenza A H5N1 viruses from clade 2.3.2 into European poultry. *Transbound Emerg Dis* **58**, 76–78.
- Robertson, S. I., Bell, D. J., Smith, G. J., Nicholls, J. M., Chan, K. H., Nguyen, D. T., Tran, P. Q., Streicher, U., Poon, L. L. & other authors (2006). Avian influenza H5N1 in viverrids: implications for wildlife health and conservation. *Proc Biol Sci* **273**, 1729–1732.
- Saitou, N. & Nei, M. (1987). The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* **4**, 406–425.
- Sakoda, Y., Sugar, S., Batchluun, D., Erdene-Ochir, T. O., Okamatsu, M., Isoda, N., Soda, K., Takakuwa, H., Tsuda, Y. & Yamamoto, N. (2010). Characterization of H5N1 highly pathogenic avian influenza virus strains isolated from migratory waterfowl in Mongolia on the way back from the southern Asia to their northern territory. *Virology* **406**, 88–94.
- Salzberg, S. L., Kingsford, C., Cattoli, G., Spiro, D. J., Janies, D. A., Aly, M. M., Brown, I. H., Couacy-Hymann, E., De Mia, G. M. & other authors (2007). Genome analysis linking recent European and African influenza (H5N1) viruses. *Emerg Infect Dis* **13**, 713–718.
- Sharshov, K., Silko, N., Sousloparov, I., Zaykovskaya, A., Shestopalov, A. & Drozdov, I. (2010). Avian influenza (H5N1) outbreak among wild birds, Russia, 2009. *Emerg Infect Dis* **16**, 349–351.
- Shivakoti, S., Ito, H., Otsuki, K. & Ito, T. (2010). Characterization of H5N1 highly pathogenic avian influenza virus isolated from a mountain hawk eagle in Japan. *J Vet Med Sci* **72**, 459–463.
- Smith, G. J., Fan, X. H., Wang, J., Li, K. S., Qin, K., Zhang, J. X., Vijaykrishna, D., Cheung, C. L., Huang, K. & other authors (2006).

Emergence and predominance of an H5N1 influenza variant in China. *Proc Natl Acad Sci U S A* **103**, 16936–16941.

Smith, G. J., Vijaykrishna, D., Ellis, T. M., Dyrting, K. C., Leung, Y. H., Bahl, J., Wong, C. W., Kai, H., Chow, M. K. & other authors (2009). Characterization of avian influenza viruses A (H5N1) from wild birds, Hong Kong, 2004–2008. *Emerg Infect Dis* **15**, 402–407.

Soda, K., Ozaki, H., Sakoda, Y., Isoda, N., Haraguchi, Y., Sakabe, S., Kuboki, N., Kishida, N., Takada, A. & Kida, H. (2008). Antigenic and genetic analysis of H5 influenza viruses isolated from water birds for the purpose of vaccine use. *Arch Virol* **153**, 2041–2048.

Starick, E., Beer, M., Hoffmann, B., Staubach, C., Werner, O., Globig, A., Strebelow, G., Grund, C., Durban, M. & other authors (2008). Phylogenetic analyses of highly pathogenic avian influenza virus isolates from Germany in 2006 and 2007 suggest at least three separate introductions of H5N1 virus. *Vet Microbiol* **128**, 243–252.

Suarez, D. L., Perdue, M. L., Cox, N., Rowe, T., Bender, C., Huang, J. & Swayne, D. E. (1998). Comparisons of highly virulent H5N1 influenza A viruses isolated from humans and chickens from Hong Kong. *J Virol* **72**, 6678–6688.

Tanimura, N., Tsukamoto, K., Okamatsu, M., Mase, M., Imada, T., Nakamura, K., Kubo, M., Yamaguchi, S., Irishio, W. & other authors (2006). Pathology of fatal highly pathogenic H5N1 avian influenza virus infection in large-billed crows (*Corvus macrorhynchos*) during the 2004 outbreak in Japan. *Vet Pathol* **43**, 500–509.

Uchida, Y., Mase, M., Yoneda, K., Kimura, A., Obara, T., Kumagai, S., Saito, T., Yamamoto, Y., Nakamura, K. & other authors (2008). Highly pathogenic avian influenza virus (H5N1) isolated from whooper swans, Japan. *Emerg Infect Dis* **14**, 1427–1429.

WHO/OIE/FAO H5N1 Evolution Working Group (2008). Toward a unified nomenclature system for highly pathogenic avian influenza virus (H5N1). *Emerg Infect Dis* **14**, e1.

WHO/OIE/FAO H5N1 Evolution Working Group (2009). Continuing progress towards a unified nomenclature for the highly pathogenic H5N1 avian influenza viruses: divergence of clade 2.2 viruses. *Influenza Other Respir Viruses* **3**, 59–62.

WHO/OIE/FAO H5N1 Evolution Working Group (2012). Continued evolution of highly pathogenic avian influenza A (H5N1): updated nomenclature. *Influenza Other Respir Viruses* **6**, 1–5.

Xu, X., Subbarao, K., Cox, N. J. & Guo, Y. (1999). Genetic characterization of the pathogenic influenza A/Goose/Guangdong/1/96 (H5N1) virus: similarity of its hemagglutinin gene to those of H5N1 viruses from the 1997 outbreaks in Hong Kong. *Virology* **261**, 15–19.

Yamamoto, N., Sakoda, Y., Motoshima, M., Yoshino, F., Soda, K., Okamatsu, M. & Kida, H. (2011). Characterization of a non-pathogenic H5N1 influenza virus isolated from a migratory duck flying from Siberia in Hokkaido, Japan, in October 2009. *Virology* **418**, 65.

An H9N2 Influenza Virus Vaccine Prepared from a Non-Pathogenic Isolate from a Migratory Duck Confers Protective Immunity in Mice against Challenge with an H9N2 Virus Isolated from a Girl in Hong Kong

Naoki NOMURA¹), Yoshihiro SAKODA¹), Kosuke SODA¹), Masatoshi OKAMATSU¹) and Hiroshi KIDA^{1,2,3)*}

¹Laboratory of Microbiology, Department of Disease Control, Graduate School of Veterinary Medicine, Hokkaido University, Kita 18 Nishi 9, Kita-ku, Sapporo 060-0818, Japan

²Research Center for Zoonosis Control, Hokkaido University, Sapporo 001-0020, Japan

³Japan Science and Technology Agency Basic Research Programs, 4-1-8 Honcho, Kawaguchi, Saitama 332-0012, Japan

(Received 20 October 2011; Accepted 13 November 2011; Published online in J-STAGE 25 November 2011)

ABSTRACT H9N2 influenza viruses circulate in wild birds and poultry in Eurasian countries, and have been isolated from pigs and humans in China. H9N2 viruses isolated from birds, pigs and humans have been classified into three sublineages based on antigenic and genetic features. Chicken antisera to H9N2 viruses of the Korean sublineage reacted with viruses of different sublineages by the hemagglutination-inhibition test. A test vaccine prepared from a non-pathogenic A/duck/Hokkaido/49/1998 (H9N2) strain of the Korean sublineage, obtained from our influenza virus library, induced immunity in mice to reduce the impact of disease caused by the challenge with A/Hong Kong/1073/1999 (H9N2), which is of a different sublineage. The present results indicate that an inactivated whole virus vaccine prepared from a non-pathogenic influenza virus from the library could be used as an emergency vaccine during the early stage of a pandemic caused by H9N2 infection.

KEY WORDS: antigenicity, H9N2 influenza virus, vaccine.

doi: 10.1292/jvms.11-0471; *J. Vet. Med. Sci.* 74(4): 441–447, 2012

Avian influenza viruses of various subtypes are circulating in poultry worldwide [1, 18, 19, 21, 29, 38]. In particular, H9N2 influenza virus is prevailing poultry populations in Eurasian countries [9–11, 24]. Since H9N2 viruses were isolated from quails in Hong Kong in 1988, they have become prevalent in live bird markets and poultry farms in Asia [6, 32]. The wide spread of H9N2 virus have been greatly concerned not only in the poultry industry but also for public health [6, 38]. The hemagglutinin (HA) genes of Eurasian H9N2 viruses have been phylogenetically divided into G1, Y280, and Korean sublineages [10]. H9N2 viruses do not substantially cause severe disease in poultry, but co-infection with bacteria such as *Staphylococcus aureus*, *Haemophilus paragallinarum*, or attenuated coronavirus vaccine exacerbates the disease [13, 22]. H9N2 viruses were also isolated from domestic pigs in China [38] and Korea, and from humans with febrile respiratory illness in Hong Kong in 1998, 1999, 2003, 2008, and 2009 [4, 5, 23, 31]. It has therefore been postulated that H9N2 virus has the potential to cause pandemic influenza in humans.

In the present study, as the preparedness for pandemic influenza, H9 virus strains from the influenza virus library in our laboratory [19] were analyzed antigenically and phylogenetically to select a strain suitable for a vaccine. A/duck/Hokkaido/49/1998 (H9N2) was selected and

an inactivated whole virus vaccine was prepared. The efficacy of the vaccine against challenge with A/Hong Kong/1073/1999 (H9N2) was assessed in mice.

MATERIALS AND METHODS

Viruses: A/duck/Hong Kong/Y280/1997 (H9N2), A/chicken/Hong Kong/G9/1997 (H9N2), A/quail/Hong Kong/G1/1997 (H9N2), A/chicken/Hong Kong/FY20/1999 (H9N2), A/silkie chicken/Hong Kong/SF43/1999 (H9N2), and A/quail/Hong Kong/A17/1999 (H9N2) were provided by Dr. K. F. Shortridge (The University of Hong Kong, China). A/ostrich/South Africa/9508103/1995 (H9N2) and A/chicken/Pakistan/2/1999 (H9N2) were provided by Dr. I. H. Brown (Animal Health and Veterinary Laboratories Agency, Weybridge, U.K.). A/Hong Kong/1073/1999 (H9N2) (HK/1073/99), which was isolated from a 4-year-old girl in Hong Kong in 1999 [23], was provided by Dr. A. J. Hay (MRC National Institute for Medical Research, U.K.). H9N2 influenza virus strains isolated from birds and mammals, and A/duck/Hokkaido/49/1998 (H9N2) (Dk/Hok/49/98) [25] are listed in Table 1. The viruses were grown in 10-day-old embryonated chicken eggs and infectious allantoic fluids were stored at –80°C until use.

Phylogenetic analysis: Viral RNAs were extracted from the allantoic fluids of chicken embryos infected with viruses using TRIzol LS Reagent (Invitrogen, Carlsbad, CA, U.S.A.) and reverse-transcribed using the Uni12 primer [14] and M-MLV reverse transcriptase (Invitrogen). The cDNA was amplified by using the Takara Ex Taq (Takara Bio, Inc., Shiga, Japan). The first cycle of the amplifica-

*CORRESPONDENCE TO: KIDA, H., Department of Disease Control, Graduate School of Veterinary Medicine, Hokkaido University, Kita 18 Nishi 9, Kita-ku, Sapporo 060-0818, Japan.
e-mail: kida@vetmed.hokudai.ac.jp

Table 1. H9N2 viruses used in this study

Sublineage	Virus	HA gene ^{a)}
Y280	A chicken Hebei 1/1996	AF536693
	A duck Hong Kong Y280/1997	AF156376
	A chicken/Hong Kong G9/1997	AF156373
	A duck Hong Kong W213/1998	AB432938
	A swine/Hong Kong/10/1998	AF222811
	A chicken/Hong Kong/FY20/1999	AF222611
	A silkie chicken Hong Kong SF43/1999	AF186268
Korean	A ostrich/South Africa/9508103/1995	AF218102
	A duck/Hokkaido/31/1997	AB125927
	A duck/Hokkaido/49/1998	AB125928
	A duck/Hokkaido/9/1999	AB125929
	A duck/Hokkaido/26/1999	AB125930
	A duck/Hokkaido/13/2000	AB125931
	A duck/Hokkaido/HY57/2005	AB455035
	A duck/Mongolia/564/2005	AB538969 ^{b)}
	A duck/Hokkaido/294/2006	AB538967 ^{b)}
	A duck/Hokkaido/W299/2006	AB538968 ^{b)}
	A duck/Hokkaido/238/2008	AB485600 ^{b)}
GI	A quail Hong Kong/GI/1997	AF156378
	A Hong Kong/1073/1999	AJ404626
	A chicken Pakistan 2/1999	AJ291392
	A quail Hong Kong/A17/1999	AF222606
North American	A/turkey/Wisconsin/1/1966	D90305

a) GenBank/EMBL/DBJ Accession No. b) The HA gene sequence was submitted to the GenBank/EMBL/DBJ databases in this study.

tion program consisted of a 5 min period at 94°C and was followed by 30 cycles with the following conditions: 98°C for 10 sec, 55°C for 30 sec, and 72°C for 1 min. The last cycle was done at 72°C for 10 min. Polymerase chain reaction amplification of the viral genes was performed using a PTC-200 thermal cycler (BIO-RAD, Hercules, CA, U.S.A.). The primers used for HA gene amplification were H9-101F (5'-GGCCACCAGTCAACAACTC-3') [24] and H9-134IR (5'-GTTTACATTCGCATCATGCTC-3'). Direct sequencing of the HA gene was performed using a CEQ 2000XL autosequencer (Beckman Coulter, Fullerton, CA, U.S.A.). For phylogenetic analysis, sequence data obtained for the genes together with those from public databases were analyzed using the neighbor-joining method [33] using MEGA 5.0 software (<http://www.megasoftware.net/>).

Antigenic analysis: Antigenic characterization of H9N2 influenza viruses was done by hemagglutination-inhibition (HI) test [35]. Hyperimmunized chicken antisera against seven H9N2 viruses were prepared [20]. Briefly, the sera were serially two-fold diluted with phosphate buffered saline (PBS) in 96-well microplates. The diluted sera were mixed with 8 hemagglutinin units of virus antigen and incubated at room temperature for 30 min. Chicken red blood cells (0.5%) were added to the antigen-serum dilution mixtures and incubated at room temperature for 30 min. HI titers were expressed as reciprocals of the highest serum dilutions that showed complete HI.

Virus replication and pathogenicity in embryonated chicken eggs: Viruses were inoculated into 10-day-old embryonated chicken eggs and incubated for 48 hr at 35°C. HA titers and 50% egg infectious dose (EID₅₀) were measured every 12 hr post-inoculation. Pathogenicity of Dk/Hok/49/98 against embryonated chicken eggs was evaluated by the mean death time as described Abenes *et al.* [2].

Vaccine preparation: Dk/Hok/49/98 and HK/1073/99 were injected into the allantoic cavities of 10-day-old embryonated chicken eggs and propagated at 35°C for 48 hr. The viruses in the allantoic fluids (512 HA for Dk/Hok/49/98 and 1,024 HA for HK/1073/99) were purified by differential centrifugation and sedimentation through a sucrose gradient [14]. The protein concentration was measured using the BCA Protein Assay Reagent (Thermo Fisher Scientific K. K., Waltham, MA, U.S.A.). The purified virus was inactivated with 0.1% formalin at 4°C for 7 days. The HA content was standardized as described [28]. Proteins of purified viruses were separated by sodium dodecyl sulfate-polyacrylamide gel electrophoresis on a 15% gel (BIO-RAD) and stained with Coomassie brilliant blue. The gel image was captured and analyzed by LumiVisionPRO (AISIN, Aichi, Japan), and the ratio of HA protein to total protein was calculated. On the basis of this method, concentration of HA protein was 14.7 µg in 50 µg of vaccine.

Challenge with HK/1073/99 into mice vaccinated once or twice: Inactivated Dk/Hok/49/98 or HK/1073/99 vaccines were injected once intraperitoneally into 4-week-old female BALB/c mice (Japan SLC, Inc., Shizuoka, Japan). PBS was injected into control mice. Three weeks later, 10 mice in each group were challenged intranasally with 30 µl of 10^{6.5} EID₅₀ of HK/1073/99 under anesthesia. Mixture of tiletamine hydrochloride (20 mg/kg) (United States Pharmacopeia, Rockville, MA, U.S.A.), zolazepam hydrochloride (20 mg/kg) (United States Pharmacopeia), and xylazine (20 mg/kg) (Bayer HealthCare, Osaka, Japan) was injected intraperitoneally into mice for anesthesia.

Inactivated Dk/Hok/49/98 vaccines were also injected twice intraperitoneally into 4-week-old female BALB/c mice. Two weeks later, the vaccine was again intraperitoneally injected into the mice. One week after the second vaccination, 10 mice in each group were challenged intranasally with 30 µl of 10^{6.5} EID₅₀ of HK/1073/99 under anesthesia.

On day 3 post-infection, five mice in each group were sacrificed and the lungs were homogenized to make a 10% (w/v) suspension with minimal essential medium (Nissui, Tokyo, Japan) with antibiotics (penicillin G potassium, streptomycin sulfate, gentamicin sulfate, and nystatin) and 0.5% Bovine Serum Albumin Fraction V (Roche, Basel, Switzerland). The virus titers of the supernatants of the lung tissue homogenates were calculated in 10-day-old embryonated chicken eggs and expressed as the EID₅₀/g of tissue.

In neutralization (NT) tests, titers were determined as the reciprocal of that maximum antibody dilution that completely prevented cytopathic effect caused by 100 plaque forming units of virus using MDCK cells.

RESULTS

Phylogenetic analysis of the HA genes of H9N2 influenza viruses: The HA genes of 22 H9N2 viruses were sequenced and phylogenetically analyzed by the neighbor-joining method. All of the HA genes were classified into the Eurasian lineage, and further classified into the Korean (n=11), Y280 (n=7), and G1 (n=4) sublineages (Fig. 1). The H9N2 viruses of the Korean and Y280 sublineages were isolated from water birds, poultry, pigs, and humans in East Asian countries, and those of the G1 sublineage were isolated from poultry in west Asian countries (Fig. 1).

Antigenicity of the H9N2 influenza viruses: H9N2 influenza viruses were antigenically analyzed by HI test (Table 2). Antisera against H9N2 viruses of the Y280 sublineage reacted slightly with H9N2 viruses of the G1 and Korean sublineages. Antisera against H9N2 viruses of the G1 sublineage reacted more with H9N2 viruses of the Y280 sublineage than those of the Korean sublineage. On the other hand, antisera against H9N2 viruses of the Korean sublineage reacted with H9N2 viruses of all sublineages. This result suggested that the H9N2 vaccine strain should be selected from the viruses of the Korean sublineage.

Selection of H9N2 vaccine strain: To select an H9N2 vaccine strain, four H9N2 viruses, Dk/Hok/49/98, A/duck/Hokkaido/13/2000 (H9N2) (Dk/Hok/13/00), A/duck/Hokkaido/9/1999 (H9N2) (Dk/Hok/9/99), and A/duck/Hokkaido/26/1999 (H9N2) (Dk/Hok/26/99), were selected from HI isolates of the Korean sublineage, and their replication and pathogenicity in embryonated chicken eggs were assessed. HA titers of Dk/Hok/49/98, Dk/Hok/13/00, Dk/Hok/9/99, and Dk/Hok/26/99 were 512, 512, 256, and 128, respectively. Virus titers were $10^{9.7}$, $10^{8.3}$, $10^{8.3}$, and $10^{7.3}$ EID₅₀/ml, respectively, indicating that Dk/Hok/49/98 replicated efficiently in 10-day-old embryonated chicken eggs. Pathogenicity of Dk/Hok/49/98 in the embryonated chicken eggs was determined by mean death time and that of Dk/Hok/49/98 was 91.8 hr, indicating that Dk/Hok/49/98 had low pathogenicity in chicken embryos. This virus was selected as a candidate H9N2 vaccine strain.

Protective efficacy of the test vaccine in mice against H9N2 virus challenge: To assess the efficacy of the vaccine against H9N2 virus infection, HK/1073/99 was intranasally inoculated into mice that had previously been vaccinated once with inactivated HK/1073/99 or Dk/Hok/49/98. Immunogenicity of the inactivated vaccine was assessed by NT test, and virus titers in the lungs were measured to assess protective immunity induced by the vaccine (Table 3). Serum antibodies were detected in mice injected with 50, 10, and 2 μ g protein of HK/1073/99 vaccine. The virus titers in the lungs were $<10^{1.5}$ – $10^{3.7}$ EID₅₀/g in mice injected with 50 or 10 μ g protein of HK/1073/99 vaccine, and $10^{4.7}$ – $10^{6.8}$ EID₅₀/g in the 2 and 0.4 μ g vaccine groups, and in the PBS control group (Table 3). A reduction in body weight was observed in mice injected with 10, 2, and 0.4 μ g protein, and in the control group from day 2 post-infection, reaching up to 10% body weight loss at days 3–4 post-infection, compared with in the mice that received 50 μ g of protein




Fig. 1. Phylogenetic tree of the HA genes of H9N2 influenza viruses. Nucleotides 163–1,048 (886 bases) of the HA genes were used for the analysis. Horizontal distances are proportional to the minimum number of nucleotide differences required to join nodes and sequences. Numbers at the nodes indicate confidence levels in a bootstrap analysis with 1,000 replicates. Viruses stocked in our laboratory are highlighted in gray. Representative viruses in each sublineage are underlined. Abbreviations: Sw, swine; Ck, chicken; Sck, silky chicken; Dk, duck; Osr, ostrich; Qa, quail; Ty, turkey; HK, Hong Kong; Hb, Hebei; S.Af, South Africa; Hok, Hokkaido; Pak, Pakistan; and Wis, Wisconsin.

(Fig. 2A).

We also tested the efficacy of vaccination with Dk/Hok/49/98 on protection against subsequent intranasal infection with HK/1073/99. Serum antibodies were slightly detected in mice injected with Dk/Hok/49/98 vaccine containing 50 and 10 μ g protein (Table 3). The virus titers in the lungs of mice injected with Dk/Hok/49/98 vaccine containing 50 and 10 μ g protein were $10^{4.3}$ – $10^{5.3}$ EID₅₀/g. In the mice injected with 2 and 0.4 μ g protein, the virus titers in the lungs of mice were similar to those of non-vaccinated control mice (Table 3). Although, a reduction in body weight was observed in mice at all doses of the Dk/Hok/49/98 vaccine, slight significant difference was observed in mice injected with 50 μ g protein, compared with in mice injected with PBS (Fig. 2B).

Table 2. The cross-reactivity of H9N2 viruses with antisera by HI test

Sublineage	Virus ^{b)}	Antisera ^{a)}							
		Y280		Korean		G1		North American	
		Ck HK G9/97	Dk/HK/Y280/97	Dk HK W213/98	Dk Hok/49/98	Dk/Hok/13/00	Qa HK G1/97	Ty Wis 1/66	
Y280	Ck Hb 1/96	10,240	10,240	2,560	2,560	2,560	2,560	40	
	Ck HK G9/97	<u>40,960</u>	10,240	2,560	1,280	2,560	2,560	40	
	Dk HK Y280/97	20,480	<u>20,480</u>	2,560	2,560	2,560	5,120	320	
	Sw HK 10/98	2,560	10,240	640	640	1,280	1,280	40	
	Dk HK W213/98	40,960	20,480	<u>2,560</u>	1,280	2,560	2,560	80	
	Ck HK FY20/99	10,240	10,240	2,560	2,560	2,560	5,120	160	
Korean	Dk Hok 31/97	640	640	640	1,280	2,560	160	640	
	Dk Hok 49/98	320	320	160	<u>2,560</u>	2,560	160	640	
	Dk Hok 9/99	640	640	160	2,560	2,560	80	320	
	Dk Hok 26/99	320	640	160	2,560	2,560	40	640	
	Dk Hok 13/00	640	640	160	1,280	<u>2,560</u>	80	640	
	Dk Mon 564/03	320	320	160	1,280	2,560	160	320	
	Dk Hok HY57/05	640	320	320	1,280	2,560	80	320	
	Dk Hok W299/06	640	320	640	1,280	2,560	80	640	
	Dk Hok 238/08	640	640	640	1,280	2,560	80	640	
G1	Qa HK G1/97	640	1,280	320	1,280	1,280	<u>5,120</u>	320	
	Ck Pak 2/99	1,280	1,280	640	640	640	640	80	
	HK 1073/99	1,280	320	160	1,280	80	1,280	320	
North American	Ty Wis 1/66	80	20	20	320	320	<20	<u>640</u>	

a) Homologous reactions are underlined. b) This panel showed the representative strains of each sublineage.

Table 3. Neutralizing antibody titers before challenge and virus titers of the lungs after challenge in mice vaccinated once

Vaccine	Dose of vaccine	NT titer to		Virus titer ^{a)} (log ₁₀ EID ₅₀ /g)
		HK/1073/99	Dk/Hok/49/98	
HK/1073/99	50 µg	320, 320, 160, 320, 640	ND	<1.5, <1.5, <1.5, <1.5, <1.5
	10 µg	80, 20, 80, 80, 40	ND	1.8, 3.7, 2.5, 2.3, 3.0
	2 µg	40, 20, 20, 20, 20	ND	4.7, 5.5, 5.5, 5.3, 6.0
	0.4 µg	<10, <10, <10, <10, <10	ND	5.7, 6.0, 5.0, 5.8, 5.5
	PBS	<10, <10, <10, <10, <10	ND	6.5, 6.2, 6.8, 6.5, 6.8
Dk Hok 49/98	50 µg	40, 40, 40, 20, 40	160, 80, 80, 80, 80	4.3, 4.8, 4.5, 4.5, 4.8
	10 µg	<10, <10, <10, <10, <10	80, 80, 80, 80, 20	4.8, 5.0, 5.2, 4.7, 5.3
	2 µg	<10, <10, <10, <10, <10	<10, <10, <10, <10, <10	6.5, 5.8, 6.0, 6.3, 6.0
	0.4 µg	<10, <10, <10, <10, <10	<10, <10, <10, <10, <10	6.3, 5.8, 6.0, 6.5, 6.3
	PBS	<10, <10, <10, <10, <10	<10, <10, <10, <10, <10	6.2, 6.0, 6.5, 5.8, 6.5

Each of vaccine was injected intraperitoneally with 10 mice. Serum samples were collected 3 weeks after the vaccination. Mice were challenged with 10^{6.0} EID₅₀ of HK/1073/99 intranasally. a) The lung samples were collected at 3 d.p.c. and virus titers were measured.

In the mice injected twice with Dk/Hok/49/98 vaccine on days 0 and 14, serum antibodies were detected in mice in the 50, 10, and 2 µg groups at one week after the second injection (Table 4). The virus titers in the lungs were <10^{1.5}–10^{3.8} EID₅₀/g in mice injected with 50, 10, and 2 µg protein, and 10^{3.3}–10^{6.5} EID₅₀/g in the other vaccinated mice (Table 4). A reduction in body weight was observed in mice injected with 2 and 0.4 µg protein, and in control group, reaching up to 10% weight loss from days 4–6 post-infection, compared with in mice injected with 50 and 10 µg protein (Fig. 3). These results suggest that the repeat administration of the test vaccine confers immunity, and

prevents body weight loss and decreases virus replication, after infection of mice with H9 influenza virus.

DISCUSSION

H9N2 viruses of each of the three sublineages, G1, Y280, and Korean, were recently isolated from wildbirds and poultry worldwide [3, 6, 27]. H9N2 viruses were also isolated from pigs and humans in China [4, 5, 37] and Korea, suggesting that these viruses have the potential to cause pandemic influenza in humans. H9N2 viruses isolated from pigs in China and Korea were classified into the Y280 and




Fig. 2. Changes in body weight in mice vaccinated once following challenge with HK/1073/99. Five vaccinated mice of each group injected with HK/1073/99 vaccine (A), and with Dk/Hok/49/98 (B) were inoculated intranasally with HK/1073/99 and body weight was monitored for 14 days. Data are shown as mean body weight changes in each group with corresponding standard deviation (SD). Asterisks indicate that body weights were not significantly ($P < 0.05$) decreased than PBS injected group.




Fig. 3. Changes in body weight in mice vaccinated twice with Dk/Hok/49/98 vaccine following challenge with HK/1073/99. Five mice from each Dk/Hok/49/98 vaccine group were inoculated intranasally with HK/1073/99 and body weight was monitored for 14 days. Data are shown as mean body weight changes in each group with corresponding SD. Asterisks indicate that body weights were not significantly ($P < 0.05$) decreased than PBS injected group.

Korean sublineages, while H9N2 viruses isolated from humans in China was classified into the G1 and Y280 sublineages [4, 5, 23, 31]. It is suggested that H9N2 viruses isolated from pigs and humans are antigenically distinct among viruses of the Korean, Y280, and G1 sublineages [5, 23, 31, 37]. Therefore, it is important that any H9N2 influenza virus vaccine to be used for pandemic influenza can broadly cross-react with antisera against all sublineage viruses. In the present study, Dk/Hok/49/98 was selected from the Korean sublineage, since antisera to the virus cross-reacted with all sublineages virus. Furthermore, Dk/Hok/49/98 replicated efficiently in embryonated chicken eggs and was non-pathogenic in chicken embryos. Recently, H9N2 viruses were isolated from pigs and humans in China [4, 5, 23, 31], it is necessary to analyze the antigenicity of these H9 isolates and evaluate the efficacy of test vaccine against

Table 4. Neutralizing antibody titers before challenge and virus titers of the lungs after challenge in mice vaccinated twice

Vaccine	Dose of vaccine	NT titer to		Virus titer ^{a)} (logEID ₅₀ /g)
		HK/1073/99	Dk/Hok/49/98	
Dk/Hok/49/98	50 µg	320, 320, 320, 160, 160	1,280, 640, 640, 320, 640	<1.5, <1.5, <1.5, <1.5, <1.5
	10 µg	40, <10, 20, 20, 40	320, 160, 320, 160, 160	2.0, 2.5, 2.0, 2.3, 2.5
	2 µg	<10, <10, <10, <10, <10	80, 160, 80, 80, 80	3.8, 3.5, 3.8, 3.5, 3.3
	0.4 µg	<10, <10, <10, <10, <10	<10, <10, <10, <10, <10	5.3, 5.8, 5.8, 6.0, 5.5
	PBS	<10, <10, <10, <10, <10	<10, <10, <10, <10, <10	5.8, 6.5, 6.0, 6.3, 5.8

Each of vaccine was injected intraperitoneally twice with 10 mice. Serum samples were collected 2 weeks after the second vaccination. Mice were challenged with 10^{6.0} EID₅₀ of HK/1073/99 intranasally. a) The lung samples were collected at 3 d.p.c. and virus titers were measured.

them. Taken together, it is important to carry out surveillance of avian influenza consecutively and to analyze the isolates antigenically and phylogenetically.

In the present study, it was suggested that the test whole particle vaccine has the potency against challenge with H9N2 virus of different sublineage in mice. It was already reported that whole particle vaccine induced strong immune responses and H5N1 whole particle vaccine induced protective immunity against antigenically distinct challenge virus [12, 26]. Although the efficacy of the test vaccine observed slightly in mice vaccinated once due to the antigenic difference between Dk/Hok/49/98 and HK/1073/99, it was clear that the vaccine induced protective immunity in mice injected twice, indicating the usefulness for the preparedness of the pandemic.

The current cycle of seasonal influenza vaccine production requires detailed planning up to 6 months before vaccine manufacture [7]. In the case of influenza pandemic in 2009, it also took 5 months to have an H1N1 vaccine available [8, 36]. To prepare for the emergence of pandemic influenza in birds and mammals including humans, we have carried out global surveillance of avian influenza [15, 30, 34, 39]. Avian influenza viruses of 144 combinations of HA and NA subtypes have been stocked for use in vaccine and diagnosis. Since the viruses stocked in our influenza virus library were already assessed the pathogenicity and replication in embryonated chicken eggs, we can exclude those tests to select a vaccine strain and prepare a vaccine rapidly [16, 17, 19]. The present results indicate that the inactivated whole virus vaccine prepared from an influenza virus from the library could be used as an emergency vaccine during the early stage of a pandemic caused by H9N2 influenza virus infection.

ACKNOWLEDGMENTS. We thank Dr. Ian Brown, Dr. Kennedy F. Shortridge, and Dr. Alan Hay for providing H9N2 influenza viruses. This study was supported by the Program of Founding Research Centers for Emerging and Reemerging Infectious Diseases of Ministry of Education, Culture, Sports, Science and Technology of Japan. This work was also supported by Japan Science and Technology Agency Basic Research Programs.

REFERENCES

1. Abbas, M. A., Spackman, E., Swayne, D. E., Ahmed, Z., Sarmiento, L., Siddique, N., Nacem, K., Hameed, A. and Rehmani, S. 2010. Sequence and phylogenetic analysis of H7N3 avian influenza viruses isolated from poultry in Pakistan 1995–2004. *Evol. J.* **7**: 137–137. [Medline] [CrossRef]
2. Abenes, G. B., Okazaki, K., Fukushi, H., Kida, H., Honda, E., Yagyu, K., Tsuji, M., Sato, H., Omo, E., Yanagawa, R. and Yamachi, N. 1982. Isolation of ortho- and paramyxoviruses from feral birds in Hokkaido, Japan—1980 and 1981. *Nihon Jinguaku Zasshi* **44**: 703–708. [Medline] [CrossRef]
3. Banet-Noach, C., Perk, S., Simanov, L., Grebenyuk, N., Rozenblut, E., Pokamunski, S., Pirak, M., Tendler, Y. and Panshin, A. 2007. H9N2 influenza viruses from Israeli poultry: a five-year outbreak. *Avian Dis.* **51**: 290–296. [Medline] [CrossRef]
4. Butt, K. M., Smith, G. J., Chen, H., Zhang, L. J., Leung, Y. H., Xu, K. M., Lim, W., Webster, R. G., Yuen, K. Y., Peiris, J. S. and Guan, Y. 2005. Human infection with an avian H9N2 influenza A virus in Hong Kong in 2003. *J. Clin. Microbiol.* **43**: 5760–5767. [Medline] [CrossRef]
5. Cheng, V. C., Chan, J. F., Wen, X., Wu, W. L., Que, T. L., Chen, H., Chan, K. H. and Yuen, K. Y. 2011. Infection of immunocompromised patients by avian H9N2 influenza A virus. *J. Infect.* **62**: 394–399. [Medline] [CrossRef]
6. Ge, F. F., Zhou, J. P., Liu, J., Wang, J., Zhang, W. Y., Sheng, L. P., Xu, F., Ju, H. B., Sun, Q. Y. and Liu, P. H. 2009. Genetic evolution of H9 subtype influenza viruses from live poultry markets in Shanghai, China. *J. Clin. Microbiol.* **47**: 3294–3300. [Medline] [CrossRef]
7. Gerdtl, C. 2003. The annual production cycle for influenza vaccine. *Vaccine* **21**: 1776–1779. [Medline] [CrossRef]
8. Girard, M. P., Tam, J. S., Assossou, O. M. and Kieny, M. P. 2010. The 2009 A (H1N1) influenza virus pandemic: a review. *Vaccine* **28**: 4895–4902. [Medline] [CrossRef]
9. Guan, Y., Shortridge, K. F., Krauss, S. and Webster, R. G. 1999. Molecular characterization of H9N2 influenza viruses: were they the donors of the "internal" genes of H5N1 viruses in Hong Kong? *Proc. Natl. Acad. Sci. U.S.A.* **96**: 9363–9367. [Medline] [CrossRef]
10. Guan, Y., Shortridge, K. F., Krauss, S., Chin, P. S., Dyrting, K. C., Ellis, T. M., Webster, R. G. and Peiris, M. 2000. H9N2 influenza viruses possessing H5N1-like internal genomes continue to circulate in poultry in southeastern China. *J. Virol.* **74**: 9372–9380. [Medline] [CrossRef]
11. Guo, Y. J., Krauss, S., Senne, D. A., Mo, F. P., Lo, K. S., Xiong, X. P., Norwood, M., Shortridge, K. F., Webster, R. G. and Guan, Y. 2000. Characterization of the pathogenicity of members of the newly established H9N2 influenza virus lineages in Asia. *Virology* **267**: 279–288. [Medline] [CrossRef]
12. Hagenaars, N., Mastroianni, E., Glansbeek, H., Heldens, J., van den Bosh, H., Schijns, V., Betheder, D., Vromans, H. and Jiskoot, W. 2008. Head-to-head comparison of four nonadjuvanted inactivated cell culture-derived influenza vaccines: effect of composition, spatial organization and immunization route on the immunogenicity in a murine challenge model. *Vaccine* **26**: 6555–6563. [Medline] [CrossRef]
13. Haghghat-Jahromi, M., Asasi, K., Nili, H., Dadras, H. and Shooshtari, A. H. 2008. Coinfection of avian influenza virus (H9N2 subtype) with infectious bronchitis live vaccine. *Arch. Virol.* **153**: 651–655. [Medline] [CrossRef]
14. Hoffmann, E., Stehm, J., Guan, Y., Webster, R. G. and Perez, D. R. 2001. Universal primer set for the full-length amplification of all influenza A viruses. *Arch. Virol.* **146**: 2275–2289. [Medline] [CrossRef]
15. Ito, T., Okazaki, K., Kawaoka, Y., Takada, A., Webster, R. G. and Kida, H. 1995. Perpetuation of influenza A viruses in Alaskan waterfowl reservoirs. *Arch. Virol.* **140**: 1163–1172. [Medline] [CrossRef]
16. Itoh, Y., Ozaki, H., Ishigaki, H., Sakoda, Y., Nagata, T., Soda, K., Isoda, N., Miyake, T., Ishida, H., Okamoto, K., Nakayama, M., Tsuchiya, H., Torii, R., Kida, H. and Ogasawara, K. 2010. Subcutaneous inoculation of a whole virus particle vaccine prepared from a non-pathogenic virus library induces protective immunity against H7N7 highly pathogenic avian influenza virus in cynomolgus macaques. *Vaccine* **28**: 780–789. [Medline] [CrossRef]
17. Itoh, Y., Ozaki, H., Tsuchiya, H., Okamoto, K., Torii, R., Sa-

- koda, Y., Kawaoka, Y., Ogasawara, K. and Kida, H. 2008. A vaccine prepared from a non-pathogenic H5N1 avian influenza virus strain confers protective immunity against highly pathogenic avian influenza virus infection in cynomolgus macaques. *Vaccine* **26**: 562–572. [Medline] [CrossRef]
18. Jeong, O. M., Kim, Y. J., Choi, J. G., Kang, H. M., Kim, M. C., Kwon, J. H. and Lee, Y. J. 2011. Genetic characterization of H1 avian influenza viruses isolated from migratory birds and domestic ducks in Korea. *Virus Genes* **42**: 55–63. [Medline] [CrossRef]
 19. Kida, H. and Sakoda, Y. 2006. Library of influenza virus strains for vaccine and diagnostic use against highly pathogenic avian influenza and human pandemics. *Dev. Biol. (Basel)* **124**: 69–72. [Medline]
 20. Kida, H. and Yamagawa, R. 1979. Isolation and characterization of influenza A viruses from wild free-flying ducks in Hokkaido, Japan. *Zentralbl. Bakteriol. [Orig. A]* **244**: 135–143. [Medline]
 21. Kim, H. R., Lee, Y. J., Lee, K. K., Oem, J. K., Kim, S. H., Lee, M. H., Lee, O. S. and Park, C. K. 2010. Genetic relatedness of H6 subtype avian influenza viruses isolated from wild birds and domestic ducks in Korea and their pathogenicity in animals. *J. Gen. Virol.* **91**: 208–219. [Medline] [CrossRef]
 22. Kishida, N., Sakoda, Y., Ito, M., Sunaga, Y. and Kida, H. 2004. Co-infection of *Staphylococcus aureus* or *Haemophilus paragallinarum* exacerbates H9N2 influenza A virus infection in chickens. *Arch. Virol.* **149**: 2095–2104. [Medline] [CrossRef]
 23. Lin, Y. P., Shaw, M., Gregory, V., Cameron, K., Lim, W., Klimov, A., Subbarao, K., Guan, Y., Krauss, S., Shortridge, K., Webster, R., Cox, N. and Hay, A. 2000. Avian-to-human transmission of H9N2 subtype influenza A viruses: relationship between H9N2 and H5N1 human isolates. *Proc. Natl. Acad. Sci. U.S.A.* **97**: 9654–9658. [Medline] [CrossRef]
 24. Liu, J., Okazaki, K., Ozaki, H., Sakoda, Y., Wu, Q., Chen, F. and Kida, H. 2003. H9N2 influenza viruses prevalent in poultry in China are phylogenetically distinct from A quail Hong Kong G1/97 presumed to be the donor of the internal protein genes of the H5N1 Hong Kong/97 virus. *Avian Pathol.* **32**: 551–560. [Medline] [CrossRef]
 25. Liu, J. H., Okazaki, K., Shi, W. M., Wu, Q. M., Mweene, A. S. and Kida, H. 2003. Phylogenetic analysis of neuraminidase gene of H9N2 influenza viruses prevalent in chickens in China during 1995–2002. *Virus Genes* **27**: 197–202. [Medline] [CrossRef]
 26. Lu, X., Edwards, L. E., Desheva, J. A., Nguyen, D. C., Rekstin, A., Stephenson, L., Szretter, K., Cox, N. J., Rudenko, L. G., Klimov, A. and Katz, J. M. 2006. Cross-protective immunity in mice induced by live-attenuated or inactivated vaccines against highly pathogenic influenza A (H5N1) viruses. *Vaccine* **24**: 6588–6593. [Medline] [CrossRef]
 27. Nagarajan, S., Rajukumar, K., Tosh, C., Ramaswamy, V., Purohit, K., Saxena, G., Behera, P., Pattinark, B., Pradhan, H. K. and Dubey, S. C. 2009. Isolation and pathotyping of H9N2 avian influenza viruses in Indian poultry. *Vet. Microbiol.* **133**: 154–163. [Medline] [CrossRef]
 28. Ninomiya, A., Imai, M., Tashiro, M. and Odagiri, T. 2007. Inactivated influenza H5N1 whole-virus vaccine with aluminum adjuvant induces homologous and heterologous protective immunities against lethal challenge with highly pathogenic H5N1 avian influenza viruses in a mouse model. *Vaccine* **25**: 3554–3560. [Medline] [CrossRef]
 29. Nguyen, D. C., Uyekci, I. M., Jadhao, S., Maines, T., Shaw, M., Matsuoka, Y., Smith, C., Rowe, T., Lu, X., Hall, H., Xu, X., Balish, A., Klimov, A., Tumpey, T. M., Swayne, D. E., Huynh, L. P., Nghiem, H. K., Nguyen, H. H., Hoang, L. T., Cox, N. J. and Katz, J. M. 2005. Isolation and characterization of avian influenza viruses, including highly pathogenic H5N1, from poultry in five bird markets in Hanoi, Vietnam, in 2001. *J. Virol.* **79**: 4201–4212. [Medline] [CrossRef]
 30. Okazaki, K., Takada, A., Ito, T., Imai, M., Takakuwa, H., Hatta, M., Ozaki, H., Ianzaki, T., Nagano, T., Ninomiya, A., Demenev, V. A., Tyapirganov, M. M., Karatayeva, T. D., Yamaikova, S. S., Lvov, D. K. and Kida, H. 2000. Precursor genes of future pandemic influenza viruses are perpetuated in ducks nesting in Siberia. *Arch. Virol.* **145**: 885–893. [Medline] [CrossRef]
 31. Peiris, M., Yuen, K. Y., Leung, C. W., Chan, K. H., Ip, P. L., Lai, R. W., Orr, W. K. and Shortridge, K. F. 1999. Human infection with influenza H9N2. *Lancet* **354**: 916–917. [Medline] [CrossRef]
 32. Perez, D. R., Lim, W., Seiler, J. P., Yi, G., Peiris, M., Shortridge, K. F. and Webster, R. G. 2003. Role of quail in the interspecies transmission of H9 influenza A viruses: molecular changes on HA that correspond to adaptation from ducks to chickens. *J. Virol.* **77**: 3148–3156. [Medline] [CrossRef]
 33. Saitou, N. and Nei, M. 1987. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol. Biol. Evol.* **4**: 406–425. [Medline]
 34. Sakoda, Y., Sugar, S., Batchluun, D., Erdene-Ochir, T. O., Okamatsu, M., Isoda, N., Soda, K., Takakuwa, H., Tsuda, Y., Yamamoto, N., Kishida, N., Matsuno, K., Nakayama, E., Kajihara, M., Yokoyama, A., Takada, A., Sodnomdarjaa, R. and Kida, H. 2010. Characterization of H5N1 highly pathogenic avian influenza virus strains isolated from migratory waterfowl in Mongolia on the way back from the southern Asia to their northern territory. *Virology* **406**: 88–94. [Medline] [CrossRef]
 35. Sever, J. L. 1962. Application of a microtechnique to viral serological investigations. *J. Immunol.* **88**: 320–329. [Medline]
 36. Stone, R. 2009. Swine flu outbreak: China first to vaccinate against novel H1N1 virus. *Science* **325**: 1482–1483. [Medline] [CrossRef]
 37. Xu, C., Fan, W., Wei, R. and Zhao, H. 2004. Isolation and identification of swine influenza recombinant A Swine Shandong/1/2003(H9N2) virus. *Microbes Infect.* **6**: 919–925. [Medline] [CrossRef]
 38. Xu, K. M., Smith, G. J., Bahl, J., Duan, F., Tai, H., Vijaykrishna, D., Wang, J., Zhang, J. X., Li, K. S., Fan, X. H., Webster, R. G., Chen, H., Peiris, J. S. and Guan, Y. 2007. The genesis and evolution of H9N2 influenza viruses in poultry from southern China, 2000 to 2005. *J. Virol.* **81**: 10389–10401. [Medline] [CrossRef]
 39. Yamamoto, N., Sakoda, Y., Motoshima, M., Yoshino, F., Soda, K., Okamatsu, M. and Kida, H. 2011. Characterization of a non-pathogenic H5N1 influenza virus isolated from a migratory duck flying from Siberia in Hokkaido, Japan, in October 2009. *Virol. J.* **8**: 65. [Medline] [CrossRef]

Characterization of Neutralizing Antibodies in Adults After Intranasal Vaccination With an Inactivated Influenza Vaccine

Akira Ainai,^{1,2} Shin-ichi Tamura,² Tadaki Suzuki,² Ryo Ito,^{2,3} Hideki Asanuma,¹ Takeshi Tanimoto,⁴ Yasuyuki Gomi,⁴ Sadao Manabe,⁴ Toyokazu Ishikawa,⁴ Yoshinobu Okuno,⁴ Takato Odagiri,¹ Masato Tashiro,¹ Tetsutaro Sata,² Takeshi Kurata,² and Hideki Hasegawa^{1,2*}

¹Influenza Virus Research Center, National Institute of Infectious Diseases, Musashimurayama, Tokyo, Japan

²Department of Pathology, National Institute of Infectious Diseases, Musashimurayama, Tokyo, Japan

³Department of Biological Science and Technology, Tokyo University of Science, Noda, Chiba, Japan

⁴The Research Foundation for Microbial Diseases of Osaka University, Kanonji, Kagawa, Japan

The levels and properties of neutralizing antibodies in nasal wash and serum collected from five healthy adults were examined after intranasal administration of an A/Uruguay/716/2007 (H3N2) split vaccine (45 μ g hemagglutinin (HA) per dose; five doses, with an interval of 3 weeks between each dose). Prior to the assays, nasal wash samples were concentrated so that the total amount of antibodies was equivalent to about 1/10 of that found in the natural nasal mucus. Vaccination induced virus-specific neutralizing antibody responses, which increased with the number of vaccine doses given. Neutralizing antibodies were produced more efficiently in the nasal passages than in the serum: A \geq 4-fold increase in nasal neutralization titres was observed after the second vaccination in four out of five subjects, whereas a rise in serum neutralization titres was observed only after the fifth vaccination. Nasal and serum neutralizing antibodies were mainly found in the polymeric IgA and monomeric IgG fractions, respectively, after gel filtration. Taken together, these results suggest that intranasal administration of an inactivated split vaccine induces high levels of nasal neutralizing antibodies (primarily polymeric IgA) and low levels of serum neutralizing antibodies (primarily monomeric IgG). *J. Med. Virol.* 84:336–344, 2012. © 2011 Wiley Periodicals, Inc.

KEY WORDS: influenza; vaccine; neutralizing antibody

INTRODUCTION

To prevent influenza, protective immunity must be induced in advance by administration of a vaccine.

Currently available inactivated vaccines, detergent disrupted split-viruses, or purified glycoproteins (surface antigen vaccines) are given via parenteral injection [Murphy and Webster, 1996]. Parenteral vaccination, that is, vaccination via the non-mucosal route, induces serum IgG antibodies, which are highly protective against homologous virus infection, but less effective against heterologous virus infection. Thus, intramuscular vaccination of seasonal influenza vaccine would be less effective in protecting against a heterologous virus epidemic.

A large number of studies show that the protective immunity induced by influenza virus infection is mainly mediated by secretory IgA (S-IgA) and IgG antibodies within the respiratory tract. S-IgA is carried to the mucus by transepithelial transport, while serum IgG is transported from the serum to the mucus by diffusion [Murphy and Clements, 1989; Brandtzig et al., 1994; Murphy, 1994; Asahi et al., 2002; Asahi-Ozaki et al., 2004]. S-IgA in the upper respiratory tract prevents viral infection, while IgG supports S-IgA-mediated protection by neutralizing newly-generated viruses [Ito et al., 2003; Renegar et al., 2004]. IgG is the main antibody involved in anti-viral protection in the lungs [Rampal et al., 1979; Palladino et al., 1995; Renegar et al., 1998; Ito et al., 2003]. Also, polymeric S-IgA neutralizes viruses more effectively than monomeric IgA or IgG [Taylor

Grant sponsor: Japanese Ministry of Health, Labor, and Welfare.

*Correspondence to: Hideki Hasegawa, MD, PhD, Influenza Virus Research Center and Department of Pathology, National Institute of Infectious Diseases, 4-7-1 Gakuen, Musashimurayama, Tokyo, 208-011 Japan. E-mail: hasegawa@nih.go.jp

Received 3 October 2011

DOI 10.1002/jmv.22273

Published online in Wiley Online Library
(wileyonlinelibrary.com).