

(Becton Dickinson, San Jose, CA, USA) and data were analyzed using CellQuest software (Becton Dickinson).

In vitro phagocytosis assays

Inflammatory macrophages were injected into the peritoneal cavity with 4% Brewer's thioglycolate (Difco). Peritoneal exudate cells were harvested 4 days later by peritoneal lavage with complete medium (RPMI containing 5% FBS (Thermo Scientific HyClone, South Logan, UT, USA) 50 mM 2-ME, 2 mM L-glutamine, 100 U/mL penicillin and 100 µg/mL streptomycin). Cells (2×10^5) were plated in 48-well plates, and non-adherent cells were removed after 2 h. The macrophage monolayers were cultured overnight in complete medium. CFSE-labeled parasitized erythrocytes (2×10^6) were then added to the wells. The plates were incubated for 2 h at 37°C. Adherent cells were then detached and analyzed by flow cytometry to assess phagocytosis of labeled cells. Resident splenic macrophages were also used.

Because the ratio of ring-infected erythrocytes differed in each preparation, the clearance of CFSE labeled ring-infected erythrocytes was adjusted according to the following:

Clearance rate of ring-infected erythrocytes = clearance rate of erythrocytes in Percoll pellet \times ratio of ring-infected erythrocytes to the total erythrocytes in the pellet.

Measurement of the in vivo clearance of fluorescence-labeled erythrocytes

CFSE-labeled erythrocytes (1×10^7) were injected intravenously into mice. Blood was taken from the mice and the percentage of CFSE-positive erythrocytes estimated by flow cytometry. In some experiments, mice were injected intraperitoneally (i.p.) with 500 µL of CGN at a concentration of 2 mg/mL in PBS once every 2 days. This treatment was started 1 day before infection and continued until the end of each experiment.

Determination of annexin binding to erythrocytes

Suspensions of Py-infected erythrocytes were stained with APC-annexin (BD biosciences) and the DNA dye Syto 16 (Invitrogen, Carlsbad, CA, USA) to detect PS and parasite DNA, respectively. For annexin V binding, erythrocytes were incubated with annexin V for 20 min at room temperature in annexin-binding buffer (140 mM NaCl, 10 mM HEPES, 5 mM glucose, 5 mM CaCl₂, pH 7.4). Syto16 (final concentration of 20 nM) was then added to the suspensions followed by incubation for 20 min at room temperature. Cells were then analyzed by flow cytometry.

Measurement of intracellular Ca²⁺

Intracellular Ca²⁺ measurement was performed as previously described [35] with minor modifications. Packed erythrocytes

(2 µL in 2 mL of Ringer's solution (1% Hct)) were loaded with Fluo-4/AM (Invitrogen) by addition of 2 µL of a Fluo-4/AM stock solution (2.0 mM in DMSO). The cells were incubated at 37°C for 15 min with vigorous shaking in a dark room followed by incubation with an additional 2 µM of Fluo-4/AM and 0.2 µg/mL Hoechst 33342 (Molecular Probes) for another 25 min. Cells were then washed twice with Ringer's solution containing 0.5% BSA (Sigma) and once with Ringer's solution alone. As a positive control, erythrocytes were stimulated with 1 µM ionomycin for 3 min prior to analysis to increase intracellular Ca²⁺ activity. Thin blood films were prepared and slides were analyzed using a fluorescence microscope (Keyence, Osaka, Japan).

Statistical analysis

Differences between groups were analyzed for statistical significance using the Mann–Whitney or Wilcoxon tests. For the survival curves, Kaplan–Meier plots and χ^2 tests were used. A *p* value < 0.05 was considered to be statistically significant.

Acknowledgements: The authors thank Mr. T. Matsumoto, (Keyence Co. Ltd., Osaka, Japan) for technical support in using fluorescence microscopy and the members of the Department of Parasitology, Institutes of Tropical Medicine, Nagasaki University, for support in completing additional experiments. This work was supported by the Ministry of Education, Culture, Sport, Science, and Technology of Japan (Grants 21022036, 20390121).

Conflict of interest: The authors declare no financial or commercial conflict of interest.

References

- 1 WHO/UNICEF/UNU. *Iron deficiency anemia Assessment, Prevention and Control. A Guide for Program Managers.* World Health Organization, Geneva 2001.
- 2 de Silva, A., Atukorala, S., Weerasinghe, I. and Ahluwalia, N., Iron supplementation improves iron status and reduces morbidity in children with or without upper respiratory tract infection: a randomized controlled study in Colombo Sri Lanka. *Am. J. Clin. Nutr.* 2003. 77: 234–241.
- 3 World malaria report. World Health Organization, 2009.
- 4 Nyakeriga, A. M., Troye-Blomberg, M., Dorfman, J. R., Alexander, N. D., Bäck, R., Kortok, M., Chemtai, A. K. et al., Iron Deficiency and malaria among children living on the coast of Kenya. *J. Infect. Dis.* 2004. 190: 439–447.
- 5 Oppenheimer, S. J., Gibson, F. D., Macfarlane, S. B., Moody, J. B., Harrison, C., Spencer, A. and Bunari, O., Iron supplementation increases prevalence and effects of malaria: report on clinical studies in Papua New Guinea. *Trans. R. Soc. Trop. Med. Hyg.* 1986. 80: 603–8012.

- 6 Sazawal, S., Black, R. E., Ramsan, M., Chwaya, H. M., Stoltzfus, R. J., Dutta, A., Dhingra, U. et al., Effect of routine prophylactic supplementation with iron and folic acid on admission to hospital and mortality in preschool children in a high malaria transmission setting: community-based, randomized, placebo-controlled trial. *Lancet* 2006. 367: 133–143.
- 7 Verhoef, H., West, C. E., Nzyuko, S. M., de Vogel, S., van der Valk, R., Wanga, M. A., Kuijsten, A. et al., Intermittent administration of iron and sulfadoxine-pyrimethamine to control anemia in Kenyan children: a randomized controlled trial. *Lancet* 2002. 360: 908–914.
- 8 Langhorne, J., The immune response to the blood stages of *Plasmodium* in animal models. *Immunol. Lett.* 1994. 41: 99–102.
- 9 Hisaeda, H., Maekawa, Y., Iwakawa, D., Okada, H., Himeno, K., Kishihara, K., Tsukumo, S. et al., Escape of malaria parasites from host immunity requires CD4⁺CD25⁺ regulatory T cells. *Nat. Med.* 2004. 10: 29–30.
- 10 Hisaeda, H., Tetsutani, K., Imai, T., Moriya, C., Tu, L., Hamano, S., Duan, X. et al., Malaria parasites require TLR9 signaling for immune evasion by activating regulatory T cells. *J. Immunol.* 2008. 180: 2496–2503.
- 11 Coban, C., Ishii, K. J., Kawai, T., Hemmi, H., Sato, S., Uematsu, S., Takeuchi, O. et al., Toll-like receptor 9 mediates innate immune activation by the malaria pigment hemozoin. *J. Exp. Med.* 2005. 201: 19–25.
- 12 Bastos, K. R., Barboza, R., Elias, R. M., Sardinha, L. R., Grisotto, M. G., Marinho, C. R., Amarante-Mendes, G. P. et al., Impaired macrophage responses may contribute to exacerbation of blood-stage *Plasmodium chabaudi chabaudi* malaria in interleukin-12-deficient mice. *J. Interferon Cytokine Res.* 2002. 22: 1191–1199.
- 13 Koka, S., Föller, M., Lamprecht, G., Boini, K. M., Lang, C., Huber, S. M. and Lang, F., Iron deficiency influences the course of malaria in *Plasmodium berghei* infected mice. *Biochem. Biophys. Res. Commun.* 2007. 357: 608–614.
- 14 Bosman, G. J., Willekens, F. L. and Were, J. M., Erythrocyte aging: a more than superficial resemblance to apoptosis? *Cell. Physiol. Biochem.* 2005. 16: 1–8.
- 15 Hargreaves, J., Yoeli, M. and Nussenzweig, R. S., Immunological studies in rodent malaria I: Protective immunity induced in mice by mild strains of *Plasmodium berghei yoelii* against a virulent and fatal line of this plasmodium. *Ann. Trop. Med. Parasitol.* 1975. 69: 289–299.
- 16 Leida, M. N., Mahoney, J. R. and Eaton, J. W., Intraerythrocytic plasmodial calcium metabolism. *Biochem. Biophys. Res. Commun.* 1981. 103: 402–406.
- 17 Goma, J., Renia, L., Miltgen, F. and Mazier, D., Effect of iron deficiency on the hepatic development of *Plasmodium yoelii*. *Parasite* 1995. 2: 351–356.
- 18 Lewis-Hughes, P. H. and Howell, M. J., In vitro culture of *Plasmodium yoelii* blood stage. *Inter. J. Parasitol.* 1984. 14: 447–451.
- 19 Luzzi, G. A., Torii, M., Aikawa, M. and Pasvol, G., Unrestricted growth of *Plasmodium falciparum* in microcytic erythrocytes in iron deficiency and thalassaemia. *Br. J. Haematol.* 1990. 74: 519–524.
- 20 Fadok, V. A., Bratton, D. L., Rose, D. M., Pearson, A., Ezekewitz, R. A. and Henson, P. M., A receptor for phosphatidylserine-specific clearance of apoptotic cells. *Nature* 2000. 405: 85–90.
- 21 López-Revuelta, A., Sánchez-Gallego, J. I., García-Montero, A. C., Hernández-Hernández, A., Sánchez-Yagüe, J. and Llanillo, M., Membrane cholesterol in the regulation of aminophospholipid asymmetry and phagocytosis in oxidized erythrocytes. *Free Radic. Biol. Med.* 2007. 42: 1106–1118.
- 22 Miki, Y., Tazawa, T., Hirano, K., Matsushima, H., Kumamoto, S., Hamasaki, N., Yamaguchi, T. et al., Clearance of oxidized erythrocytes by macrophage: Involvement of caspases in the generation of clearance signal at band 3 glycoprotein. *Biochem. Biophys. Res. Commun.* 2007. 363: 57–62.
- 23 Orrenius, S., Zhivotovsky, B. and Nicotera, P., Regulation of cell death: The calcium-apoptosis link. *Nat. Rev. Mol. Cell. Biol.* 2003. 7: 552–565.
- 24 Mandal, D., Mazumder, A., Das, P., Kundu, M., Basu, J., Fas-, caspase 8-, and caspase 3-dependent signaling regulates the activity of the aminophospholipid translocase and phosphatidylserine externalization in human erythrocytes. *J. Biol. Chem.* 2005. 280: 39460–39467.
- 25 Föller, M., Huber, S. M., Lang, F., Erythrocyte programmed cell death. *IUBMB Life* 2008. 60: 661–668.
- 26 Huang, C. J., Gurlo, T., Haataja, L., Costes, S., Daval, M., Ryazantsev, S., Wu, X. et al., Calcium-activated calpain-2 is a mediator of beta cell dysfunction and apoptosis in type 2 diabetes. *J. Biol. Chem.* 2010. 285: 339–348.
- 27 Cook, J. D. and Lynch, S. R., The liabilities of iron deficiency. *Blood* 1986. 68: 803–809.
- 28 Kleinclauss, F., Perruche, S., Masson, E., de Carvalho Bittencourt, M., Biichle, S., Remy-Martin, J. P., Ferrand, C. et al., Intravenous apoptotic spleen cell infusion induces a TGF-beta-dependent regulatory T-cell expansion. *Cell Death Differ.* 2006. 13: 41–52.
- 29 Ayi, K., Turrini, F., Piga, A. and Arese, P., Enhanced phagocytosis of ring-parasitized mutant erythrocytes: a common mechanism that may explain protection against falciparum malaria in sickle trait and beta-thalassemia trait. *Blood* 2004. 104: 3364–3371.
- 30 Cappadoro, M., Giribaldi, G., O'Brien, E., Turrini, F., Mannu, F., Ulliers, D. and Simula, G., Early phagocytosis of glucose-6-phosphate dehydrogenase (G6PD)-deficient erythrocytes parasitized by *Plasmodium falciparum* may explain malaria protection in G6PD deficiency. *Blood* 1998. 92: 2527–2534.
- 31 Min-Oo, G. and Gros, P., Erythrocyte variants and the nature of their malaria protective effect. *Cell. Microbiol.* 2005. 92: 753–763.
- 32 Harvey, P. W., Bell, R. G., Nesheim, M. C., Iron deficiency protects inbred mice against infection with *Plasmodium chabaudi*. *Infect. Immun.* 1985. 50: 932–934.
- 33 DeMaeyer, E. M., Dallman, P., Gurney, J. M., Hallberg, L., Sood, S. K. and Srikantia, S. G., Preventing and controlling iron deficiency anaemia through primary health care. World Health Organization, Geneva 1989.
- 34 Urban, B. C., Ferguson, D. J., Pain, A., Willcox, N., Plebanski, M., Austyn, J. M. and Roberts, D. J., *Plasmodium falciparum*-infected erythrocytes modulate the maturation of dendritic cells. *Nature* 1999. 400: 73–77.
- 35 Kempe, D. S. and Lang, F., Enhanced programmed cell death of iron-deficiency erythrocytes. *FASEB J.* 2005. 20: 368–370.

Abbreviations: APT: aminophospholipid translocase · CGN: carrageenan · IDA: iron deficiency anemia · PS: phosphatidylserine · Py: *Plasmodium yoelii*

Full correspondence: Dr. Hajime Hisaeda, Department of Parasitology, Graduate School of Medicine, Gunma University, 3-39-22 Showa-machi, Maebashi, Gunma 371-8511, Japan
Fax: +81-27-220-8025
e-mail: hisa@med.gunma-u.ac.jp

Received: 10/9/2010
Revised: 2/2/2011
Accepted: 23/2/2011
Accepted article online: 1/3/2011

A novel C-type lectin identified by EST analysis in tissue migratory larvae of *Ascaris suum*

Ayako Yoshida · Eiji Nagayasu · Yoichiro Horii · Haruhiko Maruyama

Received: 19 July 2011 / Accepted: 30 September 2011
© Springer-Verlag 2011

Abstract C-type lectins (CTLs) are a group of proteins which bind to carbohydrate epitopes in the presence of Ca^{2+} , which have been described in a wide range of species. In this study, a cDNA sequence coding a putative CTL has been identified from the cDNA library constructed from the pig round worm *Ascaris suum* lung L3 (LL3) larvae, which was designated as *A. suum* C-type lectin-1 (As-CTL-1). The 510 nucleotide open reading frame of As-CTL-1 cDNA encoded the predicted 169 amino acid protein including a putative signal peptide of 23 residues and C-type lectin/C-type lectin-like domain (CLECT) at residue 26 to 167. As-CTL-1 was most similar to *Toxocara canis* C-type lectin-1 and 4 (*Tc*-CTL-1 and 4), and highly homologous to nematode CTLs and mammalian CTLs as well, such as human C-type lectin domain family 4 member G (CLECG4). In addition, As-CTL-1 was strongly expressed in tissue migrating LL3 and the L4 larvae, which were developmental larvae stages within the mammalian host. These results suggest that *A. suum* larvae might utilize As-CTL-1 to avoid

pathogen recognition mechanisms in mammalian hosts due to its similarity to host immune cell receptors.

Introduction

C-type lectins (CTLs) constitute a large family of proteins that binds carbohydrate moieties in a Ca^{2+} -dependent manner (Drickamer 1988, 1996). They are characterized by a conserved C-type lectin/C-type lectin-like domain (CLECT) which shares Ca^{2+} - and carbohydrate-binding motifs. CLECT also contains at least four critical cysteine residues which form a two-loop structure by disulphide bonds. It is well known that CTLs are widely expressed among metazoan organisms (Drickamer and Fadden 2002; Zelensky and Gready 2005). In vertebrates, CTL represents a very large family that is classified into 17 groups (Drickamer and Fadden 2002), many of which are known as pattern-recognition receptors implicated in the recognition of pathogens by innate immunity (Weis et al. 1998). In addition, some evidence have indicated that CTLs play an important role in immune homeostasis by endogenous 'self' ligand recognition (García-Vallejo and van Kooyk 2009), and they themselves have a bactericidal activation (Cash et al. 2006).

Ascaris suum, a common round worm in pigs, is infective to a wide range of hosts, including humans, mice, cattle and chickens. When embryonated eggs are ingested by a definitive swine host, larvae hatch in the small intestine, penetrate the intestinal mucosa, and migrate through the liver and lungs, before finally reaching the intestine, where they sexually mature and produce eggs (Dold and Holland 2011). In contrast, it is generally considered that larvae, which are reached the lungs following liver migration, disperse into various tissues and

A. Yoshida (✉) · E. Nagayasu · H. Maruyama
Department of Infectious Diseases, Division of Parasitology,
Faculty of Medicine, University of Miyazaki,
5200 Kihara, Kiyotake,
Miyazaki 889-1692, Japan
e-mail: kukuri@med.miyazaki-u.ac.jp

Y. Horii
Laboratory of Veterinary Parasitic Diseases,
Department of Veterinary Sciences, Faculty of Agriculture,
University of Miyazaki,
Miyazaki, Japan

organs without further development in non-swine host (Slotved et al. 1998; Crompton 2001), although *A. suum* has been reported to develop into adult stage infrequently in human hosts (Anderson 1995; Nejsum et al. 2005; Arizono et al. 2010). However, it has not been fully explained how they discriminate pigs and other animals and what kind of interaction is involved between host and parasite during the lung phase of migration.

Expressed sequencing tag (EST) analysis is a powerful tool for profiling the gene expression pattern in a particular parasite population. Although publicly available EST databases of *A. suum* already exist in NEMBASE4 (<http://www.nematodes.org/nembase4/index.shtml>), they were constructed from adult worm, intestinal L4 larvae, newly hatched infective larvae (iL3) and egg embryos. EST analysis of tissue-migrating larvae has not yet been performed. Therefore, we explored cDNA of *A. suum* lung L3 (LL3) collected from infected rabbit lungs in order to examine what kind of biological processes were activated in tissue-migrating larvae. As one of the most frequently occurring clones, we identified a cDNA sequence for a putative *A. suum* CTL that showed specific expression during internal larval stages in mammalian hosts.

Materials and methods

Parasites and infection

Adult *A. suum* were collected from infected pigs at a local abattoir in Japan. Eggs were freed from the uterine tissue by incubating uteri in 0.1 N NaOH. After washing with distilled water, eggs were suspended and stirred in 0.1 N H₂SO₄ and cultured at 27°C for 3–4 weeks. Infective L3 larvae (iL3) were mechanically hatched from eggs and isolated free from egg shell contaminants (Takamiya et al. 1993). For the preparation of lung L3 larvae, male Japanese white rabbits (Kyudo, Kumamoto, Japan) were orally inoculated with 1.5×10^5 embryonated eggs. Six days after infection, the lungs were removed and cut into 5-mm cubes using scissors. The cubes were wrapped with Kimwipe papers and incubated in phosphate-buffered saline (PBS) at 37°C for 1.5 h, and then emerging worms were collected. Culture driven-L4 larvae (cL4) were obtained from cultures of LL3 in vitro (Islam et al. 2006).

RNA isolation and cDNA library construction

Total RNA of LL3 was isolated with TRIzol Reagent (Invitrogen, Carlsbad, CA), followed by purification of poly (A)⁺ RNA with GenElute™ mRNA Miniprep Kit (Sigma, St. Louis, MO). A cDNA library was constructed using the SMART cDNA Library Construction Kit

(Clontech, Mountain View, CA). The reverse transcription step was performed using MMLV Reverse Transcriptase with the SMART IV oligonucleotide primer and the CDS III/3' PCR primer provided in the kit. The double-stranded cDNA (ds-cDNA) was synthesized by long distance PCR with the 5' PCR primer and the CDS III/3' PCR primer using the Advantage 2 PCR kit (Clontech). The ds-cDNA was treated with proteinase K and then digested by *Sfi*I. After size fractionation, cDNA was cloned into pDNR-LIB vector, and transformed into *Escherichia coli* ElectroMAX™ DH10B™ cells (Invitrogen, Carlsbad, CA).

EST sequencing, processing and analysis

Plasmid DNA of the 2,024 randomly selected clones was extracted and single-pass sequenced from the 5'-end using sequencing primer (5'-GCATACATTATACGAAGTTATCAGTCG-3'). The sequencing was conducted on an ABI Prism 3130xl Genetic Analyzer (Applied Biosystems, Carlsbad, CA), using ABI Prism Big-Dye Terminator v3.1 Cycle Sequencing Kit (Applied Biosystems). EST sequences were clustered using SEQUENCHER (Gene Codes Corporation, Ann Arbor, MI), with a minimum sequence overlap length cut-off of 30 bases and an identity threshold of 90%, for the removal of flanking vector and adaptor sequences, followed by assembly. These contigs and singletons were subjected to BLASTN and BLASTX programs (*E* value of $\leq 1 \times 10^{-5}$) at the National Center for Biotechnology Information (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>). A protein sequence motif was identified using the InterProScan at The European Bioinformatics Institute (Zdobnov and Apweiler 2001). Alignment of ESTs was conducted by using GENETYX-WIN software (Genetyx Corporation, Tokyo, Japan).

Real-time PCR analysis

Total RNA from iL3, LL3, cL4 and adult worm tissues (head, muscle, intestine, uterus, ovary and testis) was extracted with TRIzol reagent. After treatment with DNaseI (Ambion Inc., Austin, TX), cDNA was generated from 250 ng of total RNA using PrimeScript® 1st strand cDNA Synthesis Kit (Takara Bio Inc., Shiga, Japan). Primer sets for amplification were as follows: As CTL-1 (sense, 5'-CCACCATGTTCTCGACCGTTGCT-3'; antisense, 5'-ATTCCTCCTACTGGCGCTCCT-3') and 18S ribosomal RNA gene (sense, 5'-ATCGGTTCGCGTAGGGTGGCT-3'; antisense, 5'-AAGCCGCAGGCTCCACTCCT-3'). Real-time PCR was then performed with an ABI Prism 7000 Sequence Detection Systems (Applied Biosystems) and a GoTaq® qPCR Master Mix (Promega, Madison, WI). Relative quantification was assessed by normalizing the amount of the target transcript to the 18S ribosomal RNA gene.

Results and discussion

In paratenic hosts such as humans, larvae of *A. suum* penetrate the mucosal epithelium but thereafter remain developmentally arrested in the migratory tissue phase (Crompton 2001). To understand biological events taking place in the arrested larvae, we carried out EST analysis in a cDNA library of migrating L3 larvae (LL3) collected from infected rabbit lungs (LL3). As a result from 5' ends single-pass sequencing of 2,024 clones, 1,650 ESTs were yielded. Upon clustering, these ESTs were represented by 279 distinct gene products, which consist of 78 contigs and 201 singletons.

The consensus sequences of contigs and singletons were compared against NCBI BLAST databases in BLASTX analysis, revealing a novel CTL sequence referred to as *A. suum* C-type lectin-1 (As-CTL-1). We focused on this molecule, because CTL might contribute to the establishment of successful parasitism in nematodes (Loukas et al. 1999; Urwin et al. 2002). Using RT-PCR, the full-length cDNA corresponding to As-CTL-1 was successfully amplified from LL3 RNA (data not shown). As-CTL-1 is 710 nucleotides (GenBank accession no. HQ025087), which encoded the protein of 169 amino acids including the putative signal peptide of 23 residues and C-type lectin/C-type lectin-like domain (CLECT) at residue 26 to 167. The four cysteine residues at positions 62, 136, 154 and 166, which are required to form the CLECT internal disulfide bridge formations (Zelensky and Gready 2003), were completely conserved in As-CTL-1, but the WIGL and WND motifs conserved in the classical CTLs (Zelensky and Gready 2003) were replaced by WLAL and WDD. According to carbohydrate specificity, CTLs are categorized into mannose/GlcNAc - or galactose/GalNAc -recognizing lectins (Weis et al. 1992; Iobst and Drickamer 1994). These differences suggest substitutions at the key substrate binding residues. As-CTL-1 had QPD, which was found in galactose/GalNAc-binding CTLs. Thus, although the Ca²⁺-dependent carbohydrate binding activity of As-CTL-1 was not assessed in this study, it is most likely a galactose-binding CTL from its amino acid motifs.

Subsequent sequence analysis showed that the amino acid sequence of CLECT in As-CTL-1 was found to have 33% identity to canine roundworm *Toxocara canis* C-type lectin-1 (*Tc*-CTL-1) and 38% identity to *Tc*-CTL-4. It has been reported that the free-living nematode *Caenorhabditis elegans* has more than 270 genes encoding CLECT of CTLs in their genome (Schulenburg et al. 2008). Interestingly, As-CTL-1 showed greater identity with human CTL domain family 4 member G (CLECG4; 28% identity) than with the *C. elegans* homologue (clec-149; 24% identity). Considering the eukaryotic phylogeny, it is reasonable that several CTLs from parasitic nematodes, such as *Ancylostoma ceylanicum*

(AceCTL-1), *Necator americanus* (NaCTL-2), *Heligmosomoides polygyrus* (*Hp*-CTL-1) and *Nippostrongylus brasiliensis* (*Nb*-CTL-1 and 2) CTLs, share greater identity with *C. elegans* CTLs than mammalian CTLs (Brown et al. 2006; Daub et al. 2000; Harcus et al. 2009). On the other hand, *Tc*-CTL-1, *Tc*-CTL-4 and NaCTL-1, as well as As-CTL-1, appear to be much closer to homologues in mammalian CTLs than those in *C. elegans* (Loukas et al. 1999, 2000; Daub et al. 2000), raising speculations about the role they might play in the adaptation to parasitism.

The expression of As-CTL-1 was evaluated by real-time PCR in different developmental stages, including mechanically hatched iL3, LL3, cL4 and adult (Fig. 1). The mRNA for As-CTL-1 was scarcely detected in iL3 and tissues from adult worms, including the head, muscle, intestine, uterus, ovary and testis, whereas LL3 and cL4 larvae showed strong expression of As-CTL-1 transcript. After arriving at the jejunum, L3 larvae developed to L4 stage larvae in definitive swine host in vivo. These results indicate that the expression of As-CTL-1 is up-regulated through the tissue migrating stage and intestinal larval stage.

Although the physiological function of CTLs remains unclear, a number of nematode CTLs identified so far act as a pathogen recognition molecule or an antibacterial protein in immune responses to protect the worm itself against microbial infection (O'Rourke et al. 2006; Schulenburg et al. 2008). However, considering that the expression of As-CTL-1 is confined during tissue migration, it would not be very likely that As-CTL-1 is employed for the recognition of microbes in *A. suum*, because the worms may not encounter hazardous bacteria on the migrating route, which would be maintained clean by the host immunity. Instead, greater sequence identity that As-CTL-1 shares with mammalian CTLs than *C. elegans* proteins, seems to suggest that As-CTL-1 acts as a




Fig. 1 Comparison of As-CTL-1 mRNA expression in developmental stages. Real-time PCR was performed with mechanically hatched iL3, LL3, cL4 and various adult worm tissues (head, muscle, intestine, uterus, ovary and testis). Relative expression of the As-CTL-1 mRNA was assessed by normalizing to 18S rRNA expression. Data were expressed as a ratio to As-CTL-1 gene expression in LL3

competitor for host cell receptors, possibly interfering host immune response to the worms (Loukas and Maizels 2000; Loukas and Prociv 2001). As-CTL-1 might be able to bind to ligands for mammalian CTL, such as NK receptors and/or macrophage/dendritic cell receptors (Osorio and Reis e Sousa 2010; Sun and Lanier 2011). In the present study, As-CTL-1 showed high similarity to *Tc*-CTLs and specific expression in LL3 and cL4, both of which were exposed to attack by host immune responses. Considering these findings, *A. suum* larvae might interfere with host inflammation processes by As-CTL-1 to avoid protective immune responses in infected animals during tissue migration. Further study on functional aspects of this molecule will identify novel dimensions of the host–parasite relationship and the significance of tissue migration in ascarid infection.

Acknowledgements We thank Hisako Kyan and Naomi Takeda for providing *A. suum* adult worm. This work was supported by grants from the Ministry of Education, Culture, Sports, Science and Technology of Japan (Grant-in-Aid for Scientific Research C 21590466, Grant-in-Aid for Scientific Research on Priority Areas 'Matrix of Infection Phenomena' 21022041), the Ministry of Health, Labour and Welfare (H23-Shinko-Ippan-014, H23-Kokui-Shitei-003, H22-Seisakusouyaku-Ippan-003) and grants from the University of Miyazaki (Support Program for Female Researchers M41D1123 and Integrated Research Project for Human and Veterinary Medicine).

References

- Anderson TJC (1995) *Ascaris* infections in humans from North America: molecular evidence for cross-infection. *Parasitology* 110(2):215–219
- Arizono N, Yoshimura Y, Tohzaka N, Yamada M, Tegoshi T, Onishi K, Uchikawa R (2010) Ascariasis in Japan: is pig-derived *Ascaris* infecting humans? *Jpn J Infect Dis* 63(6):447–448
- Brown AC, Harrison LM, Kapulkin W, Jones BF, Sinha A, Savage A, Villalon N, Cappello M (2006) Molecular cloning and characterization of a C-type lectin from *Ancylostoma ceylanicum*: evidence for a role in hookworm reproductive physiology. *Mol Biochem Parasitol* 151(2):141–147
- Cash HL, Whitham CV, Behrendt CL, Hooper LV (2006) Symbiotic bacteria direct expression of an intestinal bactericidal lectin. *Science* 313(5790):1126–1130
- Crompton DW (2001) *Ascaris* and ascariasis. *Adv Parasitol* 48:285–375
- Daub J, Loukas A, Pritchard DI, Blaxter M (2000) A survey of genes expressed in adults of the human hookworm, *Necator americanus*. *Parasitology* 120(2):171–184
- Dold C, Holland CV (2011) *Ascaris* and ascariasis. *Microbes Infect* 13(7):624–631
- Drickamer K (1988) Two distinct classes of carbohydrate-recognition domains in animal lectins. *J Biol Chem* 263(20):9557–9560
- Drickamer K (1996) Ca(2+)-dependent sugar recognition by animal lectins. *Biochem Soc Trans* 24(1):146–150
- Drickamer K, Fadden AJ (2002) Genomic analysis of C-type lectins. *Biochem Soc Symp* 69:59–72
- García-Vallejo JJ, van Kooyk Y (2009) Endogenous ligands for C-type lectin receptors: the true regulators of immune homeostasis. *Immunol Rev* 230(1):22–37
- Harcus Y, Nicoll G, Murray J, Filbey K, Gomez-Escobar N, Maizels RM (2009) C-type lectins from the nematode parasites *Heligmosomoides polygyrus* and *Nippostrongylus brasiliensis*. *Parasitol Int* 58(4):461–470
- Iobst ST, Drickamer K (1994) Binding of sugar ligands to Ca(2+)-dependent animal lectins: II. Generation of high-affinity galactose binding by site-directed mutagenesis. *J Biol Chem* 269(22):15512–15519
- Islam MK, Miyoshi T, Yamada M, Alim MA, Huang X, Motobu M, Tsuji N (2006) Effect of piperazine (diethylenediamine) on the moulting, proteome express and pyrophosphate activity of *Ascaris suum* lung-stage larvae. *Acta Trop* 99:208–217
- Loukas A, Mullin NP, Tetteh KK, Moens L, Maizels RM (1999) A novel C-type lectin secreted by a tissue-dwelling parasitic nematode. *Curr Biol* 9(15):825–828
- Loukas A, Maizels RM (2000) Helminth C-type lectins and host–parasite interactions. *Parasitol Today* 16(8):333–339
- Loukas A, Doedens A, Hintz M, Maizels RM (2000) Identification of a new C-type lectin, TES-70, secreted by infective larvae of *Toxocara canis*, which binds to host ligands. *Parasitology* 121(5):545–554
- Loukas A, Prociv P (2001) Immune responses in hookworm infections. *Clin Microbiol Rev* 14:689–703
- Nejsum P, Parker ED Jr, Frydenberg J, Roepstorff A, Boes J, Haque R, Astrup I, Prag J, Skov Sørensen UB (2005) Ascariasis is a zoonosis in Denmark. *J Clin Microbiol* 43(3):1142–1148
- O'Rourke D, Baban D, Demidova M, Mott R, Hodgkin J (2006) Genomic clusters, putative pathogen recognition molecules, and antimicrobial genes are induced by infection of *C. elegans* with *M. nematophilum*. *Genome Res* 16(8):1005–1016
- Osorio F, Reis e Sousa C (2010) Myeloid C-type lectin receptors in pathogen recognition and host defense. *Immunity* 34(5):651–664
- Schulenburg H, Hoepfner MP, Weiner J 3rd, Bornberg-Bauer E (2008) Specificity of the innate immune system and diversity of C-type lectin domain (CTLCD) proteins in the nematode *Caenorhabditis elegans*. *Immunobiology* 213(3–4):237–250
- Slotved HC, Eriksen L, Murrell KD, Nansen P (1998) Early *Ascaris suum* migration in mice as a model for pigs. *J Parasitol* 84(1):16–18
- Sun JC, Lanier LL (2011) NK cell development, homeostasis and function: parallels with CD8+ T cells. *Nat Rev Immunol* 11:645–657
- Takamiya S, Kita K, Wang H, Weinstein PP, Hiraishi A, Oya H, Aoki T (1993) Developmental changes in the respiratory chain of *Ascaris* mitochondria. *Biochim Biophys Acta* 1141:65–74
- Urwin PE, Lilley CJ, Atkinson HJ (2002) Ingestion of double-stranded RNA by preparasitic juvenile cyst nematodes leads to RNA interference. *Mol Plant Microbe Interact* 15(8):747–752
- Weis WI, Drickamer K, Hendrickson WA (1992) Structure of a C-type mannose-binding protein complexed with an oligosaccharide. *Nature* 360(6400):127–134
- Weis WI, Taylor ME, Drickamer K (1998) The C-type lectin superfamily in the immune system. *Immunol Rev* 163:19–34
- Zdobnov EM, Apweiler R (2001) InterProScan – an integration platform for the signature-recognition methods in InterPro. *Bioinformatics* 17:847–848
- Zelensky AN, Gready JE (2003) Comparative analysis of structural properties of the C-type-lectin-like domain (CTLCD). *Proteins* 52:466–477
- Zelensky AN, Gready JE (2005) The C-type lectin-like domain superfamily. *FEBS J* 272(24):6179–6217

Short Communication

Eosinophilic Pneumonia Due to Visceral Larva Migrans Possibly Caused by *Ascaris suum*: a Case Report and Review of Recent Literatures

Koichi Izumikawa^{1,2*}, Yoshihisa Kohno¹, Kinichi Izumikawa¹, Kohei Hara¹, Hiroko Hayashi³, Haruhiko Maruyama⁴, and Shigeru Kohno²

¹Department of Internal Medicine, Izumikawa Hospital, Nagasaki 859-1504;

²Department of Molecular Microbiology and Immunology and

³Department of Investigative Pathology, Nagasaki University

Graduate School of Biomedical Sciences, Nagasaki 852-8501; and

⁴Division of Parasitology, Department of Infectious Diseases, Faculty of Medicine, University of Miyazaki, Miyazaki 889-1692, Japan

(Received February 12, 2011. Accepted July 11, 2011)

SUMMARY: We report the case of a 62-year-old man who developed eosinophilic pneumonia due to visceral larva migrans (VLM) that was possibly caused by *Ascaris suum*. The patient, a resident of the middle Kyushu area who was fond of eating raw porcine liver, complained of dry cough without dyspnea. The chest radiography showed a migration of infiltrative shadow. Transbronchial lung biopsy of the right middle lobe revealed massive infiltration of eosinophils. The multi-dot enzyme-linked immunosorbent assay (ELISA) and microtiter plate ELISA showed positive results for *A. suum*; therefore, the patient was diagnosed with VLM caused by *A. suum*. The patient was administered albendazole (600 mg/day) for 28 days; he recovered successfully with no adverse effects except mild liver dysfunction. Several cases of VLM caused by *A. suum* have been reported in Japan, with a majority of the cases being reported in Kyushu. Careful history taking of the patient's area of residence and dietary habit is essential for the diagnosis of this parasitic disease with underestimated prevalence.

Visceral larva migrans (VLM) caused by *Ascaris suum* is a major parasitic infection that especially affects people living in southern Kyushu, Japan, which has a prominent livestock industry (1). *A. suum* infects pigs, and 30% of all the pigs in southern Kyushu are infected (2). Humans are usually infected when they eat the raw liver or meat of infected cattle or chicken or fresh vegetables grown in soil fertilized with porcine excrement contaminated with *A. suum* eggs. In humans, the larvae of *A. suum* migrate to various organs and cause a wide variety of nonspecific symptoms such as general malaise, cough, liver dysfunction, hyper-eosinophilia with hepatomegaly and/or pneumonia (3,4). Here, we report a case of eosinophilic pneumonia resulting from VLM that could have been possibly caused by *A. suum*, and present a review of the recent literature on VLM.

A 62-year-old man living in Shimabara, Nagasaki Prefecture, Kyushu, Japan, was referred and admitted to Izumikawa Hospital because he had dry cough and chest radiography had shown an infiltration shadow in both lungs. All in one cold and flu capsules prescribed in a previous clinic had not been effective. The patient

complained of dry cough without dyspnea at the time of admission. He had no remarkable underlying diseases, although he possessed a unique dietary habit such as eating raw porcine, chicken, and cattle livers.

At the time of admission, the vital signs of the patient were as follows: body temperature, 36.8°C; heart rate, 72 beats/min (regular rhythm); respiratory rate, 16 breaths/min; and blood pressure, 110/60 mmHg. Auscultation revealed no abnormal pulmonary crackles or heart sounds. The patient showed no clinical signs of lymphadenopathy, hepatosplenomegaly, and pretibial edema.

Chest radiography showed an infiltrative shadow in both the middle and lower lung fields (Fig. 1A), and computed tomography (CT) showed consolidation with ground-glass opacity in both lung fields (Fig. 1B and 1C). The laboratory test results were as follows: leukocytes count, $9.5 \times 10^3/\mu\text{L}$; eosinophil count, 2,175/ μL (22.9%); C-reactive protein (CRP) concentration, 0.4 mg/dL; and IgE level, 333.2 IU/mL. All other results were within the normal range. The results of the arterial blood gas (ABG) analysis at room air were as follows; pH, 7.404; PaO₂, 72.6 Torr; and PaCO₂, 40.8 Torr. Routine microbiological tests revealed no causative bacteria. Bronchoscopy was performed and the bronchoalveolar lavage (BAL) fluid was analyzed; the results of the cell count analysis were as follows: alveolar macrophages, 2%; eosinophils, 93%; lymphocytes, 3%; and basophils, 2%. No microorganisms, including fungi and mycobacteria, could be isolated from the BAL fluid in routine microbiological tests. Although the polymerase

*Corresponding author: Mailing address: Department of Molecular Microbiology and Immunology, Nagasaki University Graduate School of Biomedical Sciences, 1-7-1 Sakamoto, Nagasaki 852-8501, Japan. Tel: +81(95)819-7273, Fax: +81(95)849-7285, E-mail: koizumik@nagasaki-u.ac.jp




Fig. 1. Chest X-ray films on admission. (A) Chest X-ray film showing infiltrative shadow in both middle and lower lung fields. (B) and (C) CT scan images showing consolidation with ground-glass opacity at right upper lobe and left lower lobe.

chain reaction test for mycobacteria showed positive results for *Mycobacterium intracellulare*, an 8-week culture of the BAL fluid sample showed negative growth. Transbronchial lung biopsy of the right middle lobe (B₄) revealed massive infiltration of eosinophils in the parenchyma and that of alveolar macrophages in the alveoli. Eosinophilic pneumonia was diagnosed on the basis of the results of this pathological analysis (Fig. 2). Multi-dot enzyme-linked immunosorbent assay (multi-dot ELISA) was performed for detecting anti-parasitic antibodies in the patient's serum (5). A serum sample of the patient showed positive results for *Dirofilaria immitis*, *A. suum*, and *Gnathostoma doloresi* but negative results for *Toxocara canis* (the test was not performed for *T. cati*). A microtiter plate-ELISA for the semi-quantitative measurement of the antibodies for the three parasites (6) was performed, and the strongest reaction was observed for *A. suum* antibodies. Since the patient had a history of eating raw porcine liver, we diagnosed his condition as eosinophilic pneumonia due to VLM that was possibly caused by *A. suum*.

The patient received no treatment during an observation period of 17 days after the diagnosis of VLM, and the clinical symptoms and signs such as cough and hyper eosinophilia persisted. Chest radiography performed on day 17 after the admission showed that the infiltrative shadow in the right middle and lower lung




Fig. 2. Pathology of transbronchial lung biopsy from the right middle lobe (right B₄) demonstrates severe eosinophil infiltrations in lung parenchyma. HE stain, $\times 40$.

fields had migrated, and CT showed new consolidation with ground-glass opacity in the right upper and lower lobes of the lungs (Fig. 3A, 3B, and 3C). The patient was administered albendazole (600 mg/day) for 28 days. The clinical symptoms resolved completely, and the eosinophil count decreased to 390/ μ L. The infiltra-




Fig. 3. Chest X-ray films before albendazole treatment (17 days after admission). (A) Chest X-ray film showing infiltrative shadow in right middle and lower lung fields and migrated from the time of admission. (B) and (C) CT scan images showing new consolidation with ground-glass opacity at right upper and lower lobes.

tive shadow disappeared completely in 28 days. No adverse effects except mild liver dysfunction were noted during the 28 days. No recurrence was observed after discharge.

VLM was first described by Beaver et al. in 1952 and is mainly caused by *T. canis* and *T. cati* (3,7). Humans become infected when they ingest *Toxocara* eggs. *A. suum* is also a known cause of VLM (8), especially in Kyushu, Japan, because the residents of this region eat the raw meat and liver of cattle, poultry, and horse or fresh vegetables cultivated using organic fertilizers (9).

VLM as a zoonosis has emerged as a clinical concern because of an increase in the number of dogs and cats kept as pets in Japan. People who have a habit of eating the raw meat of wild animals are at risk of infection with parasitic worms. The current trend of eating fresh vegetables as a part of a healthy lifestyle also increases the risk of infection with parasitic worms since fresh vegetables may be contaminated with them.

A definitive diagnosis of VLM is possible only if the larvae of *Toxocara* or *Ascaris* are found in the patient's body; however, detecting these larvae is quite difficult and not practical. To date, no suitable or applicable molecular methods are available for accurately detecting the genomic DNA of parasites. The multi-dot ELISA method (5), performed using the patient's serum sample, is a useful and convenient tool for diagnosing

VLM. Although it is a simple method, cross-reactions among the parasite antigens have been observed. Therefore, a definite diagnosis of VLM cannot be made unless the larvae or DNA of the causative organism, such as *Toxocara* or *Ascaris*, is detected in the patient's body. Information such as the patients' area of residence and their dietary habits should be obtained and carefully evaluated by attending physicians.

An outbreak of VLM caused by *A. suum* in Japan was reported by Maruyama et al. (6) in 1996, with a total of 17 patients with pronounced eosinophilia and high antibody titers against the *A. suum* antigen. A review of recent literature on *A. suum* cases showed that at least 9 cases of VLM in Japan were reported in various journals after 1996 (Table 1). All the patients, except 2, were infected in the Kyushu island, and possible sources of infection were the raw meat of chicken, boar, deer, and cattle (5 patients); vegetables (2 patients); raw meat of poultry or horse, raw liver of cattle or vegetables cultivated using organic fertilizer (1 patient); and unknown (1 patient). Because the eggs of *A. suum* could not be detected in these possible sources, the apparent source and route of infection were not confirmed. Almost all the patients showed high levels of serum IgE and hyper eosinophilia. Albendazole and ivermectin were administered and were effective in 7 patients and 1 patient, respectively. Only one patient was diagnosed

Table 1. Summary of cases of visceral larva migrans due to *Ascaris suum* in Japan

Study (ref. no.)	Year	Age (y)	Sex	Place	Case	Possible source of <i>A. suum</i>	Eosinophil (μ l)	Serum IgE (IU/ml)	Treatment
Matsushita et al. (12)	1997	70	Female	Miyazaki, Kyushu	Eosinophilic pneumonia, + intrahepatic lesion	Raw chicken	9,440	7,022	Albendazole
Takeyama et al. (13)	1997	56	Female	Kyushu	Eosinophilic colitis	N.A.	7,872	10,960	Prednisolone
Matsuyama et al. (14)	1998	46	Male	Kagoshima, Kyushu	Eosinophilic pneumonia	Fresh vegetables cultivated using pig manure	9,188	3,190	Ivermectin
Arimura et al. (15)	2001	26	Male	Miyazaki, Kyushu	Pulmonary nodule	Raw boar, deer meat	750	926	Albendazole
Arimura et al. (15)	2001	57	Male	Miyazaki, Kyushu	Pulmonary nodule	Raw chicken, turkey	342	832	Albendazole
Sakakibara et al. (16)	2002	32	Male	Aichi, Honsyu	Eosinophilic pneumonia, + intrahepatic lesion	Fresh vegetables cultivated using organic fertilizer, raw meat of cattle liver, poultry meat, horsemeat	10,773	20,284	Albendazole
Sakurai et al. (17)	2003	25	Female	Tokyo, Honsyu	Eosinophilic pneumonia	Raw liver of cow	7,290	98	Albendazole
Tokojima et al. (18)	2004	50	Male	Kagoshima, Kyushu	Eosinophilic pneumonia	Vegetables	445	1,208	Albendazole
Hirakawa et al. (19)	2009	64	Male	Kagoshima, Kyushu	Eosinophilic pneumonia	Raw chicken liver	4,223	279	Albendazole

All cases are diagnosed with multi-dot enzyme-linked immunosorbent assay. Outcome of all cases are cured. N.A., not available.

with eosinophilic colitis and was administered prednisolone but not albendazole.



In the present case, the following clinical signs were consistent with those of VLM: (i) remarkable eosinophilia and high IgE levels, (ii) positive results in the multi-dot ELISA and the strongest reaction for the antibody for *A. suum*, in microtiter plate-ELISA, (iii) migration of the pulmonary infiltrates, and (iv) eosinophilic pneumonia, as diagnosed on the basis of the results of BAL fluid analysis and transbronchial lung biopsy. Although there is little evidence in favor of any treatment modality for VLM caused by *A. suum*, administration of albendazole or ivermectin for 2 to 3 weeks is recommended (10,11). The patient was administered albendazole (600 mg/day) for 28 days, and he showed no remarkable adverse effects except mild impairment of liver function (a common adverse effect of albendazole). Although most of the cases of VLM caused by *A. suum* are not fatal, VLM could sometimes become life threatening if a large number of *A. suum* eggs are ingested (8). It is important for clinicians to consider VLM caused by *A. suum* in case a patient presents hypereosinophilia, high IgE levels, and a migrating pneumonia shadow in addition to various nonspecific symptoms. Careful history taking of the patients' area of residence and dietary habit is essential for the diagnosis of this parasitic disease with underestimated prevalence. Furthermore, although VLM caused by *A. suum* is most prevalent in Kyushu, a couple of VLM cases have been reported in the Honshu region as well (Table 1). Owing to advances in mass-transportation of fresh vegetables and meat and improvement of the related logistics, cases of such originally localized parasitic infections are now being detected in other areas of Japan and even in other countries.

Conflict of interest None to declare.

REFERENCES

1. Maruyama, H., Nawa, Y., Noda, S., et al. (1996): An outbreak of visceral larva migrans due to *Ascaris suum* in Kyushu, Japan. *Lancet*, 347, 1766-1767.
2. Nawa, Y., Maruyama, H. and Noda, S. (1996): Visceral larva migrans due to *Ascaris suum* in Kyushu. *Infect. Agents Surveillance Rep.*, 17, 191-192 (in Japanese).
3. Beaver, P.C., Snyder, C.H., Carrera, G.M., et al. (1952): Chronic eosinophilia due to visceral larva migrans; report of three cases. *Pediatrics*, 9, 7-19.
4. van Knapen, F., Buijs, J., Kortbeek, L.M., et al. (1992): Larva migrans syndrome: toxocara, ascaris, or both? *Lancet*, 340, 550-551.
5. Itoh, M. and Sato, S. (1990): Multi-dot enzyme-linked immunosorbent assay for serodiagnosis of trematodiasis. *Southeast Asian J. Trop. Med. Public Health*, 21, 471-474.
6. Maruyama, H., Noda, S., Choi, W.-Y., et al. (1997): Fine binding specificities to *Ascaris suum* and *Ascaris lumbricoides* antigens of the sera from patients of probable visceral larva migrans due to *Ascaris suum*. *Parasitol. Int.*, 46, 181-188.
7. Ogilvie, B. and Savigny, D. (1982): *Immune Response to Nematodes*. 2nd ed. Blackwell, Oxford.
8. Phills, J.A., Harrold, A.J., Whiteman, G.V., et al. (1972): Pulmonary infiltrates, asthma and eosinophilia due to *Ascaris suum* infestation in man. *N. Engl. J. Med.*, 286, 965-970.
9. Inatomi, Y., Murakami, T., Tokunaga, M., et al. (1999): Encephalopathy caused by visceral larva migrans due to *Ascaris suum*. *J. Neurol. Sci.*, 164, 195-199.
10. Caumes, E., Carriere, J., Datry, A., et al. (1993): A randomized trial of ivermectin versus albendazole for the treatment of cutaneous larva migrans. *Am. J. Trop. Med. Hyg.*, 49, 641-644.
11. Bhatia, V. and Sarin, S.K. (1994): Hepatic visceral larva migrans: evolution of the lesion, diagnosis, and role of high-dose albendazole therapy. *Am. J. Gastroenterol.*, 89, 624-627.
12. Matsushita, R., Tahara, Y., Yamamoto, S., et al. (1997): A case of visceral larva migrans (VLM) with multiple intra-hepatic nodular lesions due to *Ascaris suum*. *Liver (Kanzou)*, 38, 730-734 (in Japanese).
13. Takeyama, Y., Kamimura, S., Suzumiya, J., et al. (1997): Case report: eosinophilic colitis with high antibody titre against *Ascaris*

- suum*. J. Gastroenterol. Hepatol., 12, 204–206.
14. Matsuyama, W., Mizoguchi, A., Iwami, F., et al. (1998): A case of pulmonary infiltration with eosinophilia caused by *Ascaris suum*. J. Jpn. Respir. Soc., 36, 208–212 (in Japanese).
 15. Arimura, Y., Mukae, H., Yanagi, S., et al. (2001): Two cases of visceral larva migrans due to *Ascaris suum* showing a migratory nodular shadow. J. Jpn. Respir. Soc., 39, 716–720 (in Japanese).
 16. Sakakibara, A., Baba, K., Niwa, S., et al. (2002): Visceral larva migrans due to *Ascaris suum* which presented with eosinophilic pneumonia and multiple intra-hepatic lesions with severe eosinophil infiltration—outbreak in a Japanese area other than Kyushu. Intern. Med., 41, 574–579.
 17. Sakurai, M., Hakota, Y., Kodama, S., et al. (2003): Pulmonary eosinophilia caused by visceral larva due to *Ascaris suum* from raw cattle liver. Diagn. Treat., 91, 2297–2299 (in Japanese).
 18. Tokojima, M., Ashitani, J. and Nakazato, M. (2004): A case of eosinophilic pneumonia caused by visceral larva migrans due to *Ascaris suum*. J. Jpn. Assoc. Infect. Dis., 78, 1036–1040 (in Japanese).
 19. Hirakawa, E., Suetsugu, T., Tanoue, A., et al. (2009): Pulmonary eosinophilia caused by visceral larva due to *Ascaris suum*. J. Jpn. Soc. Intern Med., 98, 144–146 (in Japanese).



Case Report

A patient who developed toe necrosis due to poor blood circulation after an interdigital tick bite

Hiromu Yamazaki (MD)^a, Koji Yamaguchi (MD)^{b,*}, Takashi Iwase (MD)^b, Toshiyuki Niki (MD)^b, Kenya Kusunose (MD)^b, Noriko Tomita (MD)^b, Yoshio Taketani (MD)^b, Hirotsugu Yamada (MD)^b, Takeshi Soeki (MD, FJCC)^b, Tetsuzo Wakatsuki (MD)^b, Yutaka Fukunaga (MD)^c, Hideki Nakanishi (MD)^c, Haruhiko Maruyama (MD)^d, Hiroyuki Matsuoka (MD)^e, Masataka Sata (MD, FJCC)^b

^a The Post-Graduate Education Center, Tokushima University Hospital, Tokushima, Japan

^b Department of Cardiovascular Medicine, Institute of Health Biosciences, The University of Tokushima Graduate School, 2-50-1 Kuramoto-cho, Tokushima 770-8503, Japan

^c Department of Plastic Surgery, Institute of Health Biosciences, The University of Tokushima Graduate School, Tokushima, Japan

^d Department of Infectious Diseases, Division of Parasitology, Faculty of Medicine, University of Miyazaki, Miyazaki, Japan

^e Division of Medical Zoology, Department of Infection and Immunity, Jichi Medical University, Shimotsuke, Japan

Received 7 February 2011; received in revised form 12 May 2011; accepted 19 June 2011

KEYWORDS

Tick bite;
Toe necrosis;
Thrombogenic
vasculopathy

Summary A 71-year-old female had worked on a farm in the mountains and noticed itching of the left 3rd toe. She visited a local hospital due to a color change to purple in this area. Attachment of a tick was observed between the left 2nd and 3rd toes, and it was extracted. However, due to persistent pain, she was referred to our department of cardiovascular medicine for close examination and treatment. Lower extremity angiography showed that vascular visualization was poor in the area supplied by the arteries distal to the tick bite site, but the other blood vessels of the toe were clearly visualized. Toe amputation was performed and pathological examination of a surgical specimen revealed that most blood vessels near the necrosis were occluded by thrombi. We speculated that tick bite reactions were associated with thrombogenic vasculopathy. This report shows a patient who developed toe necrosis due to poor blood circulation after an interdigital tick bite.

© 2011 Japanese College of Cardiology. Published by Elsevier Ltd. All rights reserved.

* Corresponding author. Tel.: +81 88 633 7851; fax: +81 88 633 7894.
E-mail address: yamakoji3@clin.med.tokushima-u.ac.jp (K. Yamaguchi).

Introduction

Tick bites are relatively frequently encountered in daily clinical practice. Most patients with tick bites develop dermatitis, but some develop Japanese spotted fever or Lyme disease via the bite. However, there have been no reports of peripheral necrosis due to poor blood circulation following tick bites. We report a patient who developed toe necrosis due to poor blood circulation after an interdigital tick bite.

Case report

A 71-year-old female had previously been healthy, and, although she had annually undergone a human dry dock, no abnormalities had been detected. She had worked on a farm in the mountains on June 7, 2009, noticed itching of the left 3rd toe on June 12, and visited a local hospital due to a color change to purple in this area on June 16. Attachment of a tick was observed between the left 2nd and 3rd toes. The tick was extracted, and minocycline was prescribed. However, due to persistent pain, she was referred to our department of cardiovascular medicine for close examination and treatment on June 23. She was obese, but had no history of smoking, hypertension, diabetes mellitus, or lipid abnormalities. The bilateral dorsalis pedis and posterior tibial arteries were palpable, and the skin temperature of the foot was normal. There was a decrease in

the skin temperature, loss of sensation, and a color change to black distal to the middle phalanx of the left 3rd toe (Fig. 1A). The area between the 2nd and 3rd toes as the site of the tick bite showed erosion, but not redness or swelling (Fig. 1B). No eruptions were observed in the other areas. The thermography (Fig. 1C) showed an increase in the skin temperature in the left compared with the right foot. In the portion peripheral to the proximal interphalangeal joint of the left 3rd toe, the skin temperature showed a decrease in the area corresponding to that showing black necrosis. Lower extremity angiography (Fig. 1D) was performed with selective enhancement of the left external iliac artery. The area extending to the plantar artery and arch was clearly visualized, and no atherosclerotic changes were identified. Vascular visualization was poor in the area supplied by the arteries distal to the tick bite site, but the other blood vessels of the toe were clearly visualized. She was referred to the department of plastic surgery of our hospital on July 3, and toe amputation was considered to be indicated. On July 24, toe amputation was performed (Fig. 2A and B). Pathological examination of a surgical specimen revealed necrosis of the toe tip, and most blood vessels near the necrosis were occluded by thrombi. Between fat tissues, foam cells aggregated, and marked eosinophil infiltration was observed. There were no findings of angiitis (Fig. 2C and D). Her postoperative course was favorable. After walking became possible, she was discharged on August 4.




Figure 1 (A and B) Photo of the left toes (white arrow: tick bite site). (C) Thermography (white arrow: a decrease in the skin temperature was observed in the area corresponding to the site of black necrosis). (D) Lower extremity angiography (white arrow: blood flow interruption was confirmed in the area corresponding to the site of black necrosis).




Figure 2 (A) Surgical specimen. (B) Cross-section of the surgical specimen. (C) Image of the pathological specimen (Cross-section 4, black arrow: blood vessels near the necrosis were occluded by thrombi. In the vascular lumens, granulomatous tissue formed, but reperfusion was observed in some parts. Hematoxylin-eosin staining). (D) Image of the pathological specimen (Cross-section 4, black arrows: microscopic vessels occluded by thrombus, hematoxylin-eosin staining).

Discussion

When this patient visited our department, about 2 weeks had passed since the tick bite. Since she had already been treated by a local doctor, and the tick had been extracted, we could not directly confirm the tick. However, in the southern part of Tokushima Prefecture where she lives, tick bites are often observed, and so the diagnosis of a tick bite made by the previous doctor may be accurate. There has been a report [1] of patients with tissue necrosis in a tick bite area, but no report of patients with necrosis distant from the bite site. Previous studies have shown blood coagulation responses as biological responses to ticks [2] and pathologically confirmed thrombi in about 66% of tick bite areas [3]. Histopathological examination in our patient showed the occlusion of many blood vessels by thrombi in the area near necrosis and the presence of necrosis peripheral to these thrombi. In general, tick bites lead to reactions such as the extravascular leakage of dermal erythrocytes, hardening of collagen fibers, epidermal necrosis, ulceration, dermal neutrophil infiltration, and thrombosis [3]. However, the saliva of ticks contains anti-hemostatic factors and immunosuppressive and anti-inflammatory components, which make tick removal difficult, facilitating long-term blood-feeding [4]. The intractability of areas bitten has been suggested to be associated with the formation of a foreign body granuloma due to remnant tick mouth parts, impairment of the wound healing process by fibrosis-promoting factors contained in the saliva, and ischemia in

the wound area due to various types of local microvasculitis [5]. Since no pathological examination of the tick bite area was performed, detailed findings of this area such as the possible presence of thrombi could not be obtained. However, we speculated that thrombi induced by the tick bite caused peripheral embolism, and impaired peripheral blood circulation resulted in necrosis. We supposed that the involvement of angiitis was probably denied because the value of myeloperoxidase – antineutrophil cytoplasmic antibody was very small. Additionally, we denied the presence of shaggy aorta which was associated with blue toe syndrome by use of contrast-enhanced computed tomography. Coronary angiography was also performed in this patient, but showed no findings of stenosis. Lower extremity angiography also revealed no atherosclerotic findings anywhere except the occluded areas, most strongly suggesting the involvement of thromboembolism. Therefore, tick bites in areas with a few collateral circulation routes are associated with a risk of peripheral necrosis, and require caution. The thermography showed an increase in the foot skin temperature on the affected side. This may be an immune response or biological response such as vascular dilation in the central area due to impaired peripheral blood circulation, but the details were unclear. Dermatitis due to tick bites is often encountered in clinical practice. The most interesting point is the difference between patients with dermatitis not inducing serious conditions and those such as in the present patient who develop marked thrombotic circulatory impairment. Although an intractable case showing

local periarteritis nodosa after a tick bite to the thigh has been reported [6], peripheral necrosis was not induced in this case. We speculate that the degree of a host's immune tissue responses to ticks and the bite site are important factors associated with whether necrosis is induced. Tick bites in the peripheral areas of the four limbs, as in this patient, have the potential to induce necrosis because there are only a few collateral circulation routes, and so require caution. The characteristics of cases that show marked immunohistological responses at the tick bite site are unclear. A previous study showed that inflammatory reactions became marked, and severe inflammatory cell infiltration was histologically confirmed with an increase in the frequency of tick biting [7]. Our patient had no history of treatment for tick bites, but was a farmer, and so it is likely that she had a history of untreated asymptomatic tick bites. Concerning the association with background factors such as the residence, sex, and age, we have encountered no patient with tick bites leading to peripheral necrosis, and, therefore, could not perform statistical analysis. In the future, the accumulation of data on patients with similar necrosis may allow the clarification of more detailed mechanisms of peripheral necrosis.

We encountered a patient who developed toe necrosis due to poor blood circulation after a tick bite. We speculated that tick bite reactions were associated with thrombogenic vasculopathy. Tick bites in peripheral areas of the four extremities with only a few collateral circulation routes are associated with a risk of peripheral necrosis, and careful observation of the course is necessary.

Acknowledgments

We are grateful to Dr Yoshimi Bando in Tokushima University Hospital, Tokushima, Japan for expert assistance in preparing and analyzing histological slides and to Dr Yoshihiro Matsudate and Dr Yoshio Urano in Tokushima Red Cross Hospital, Komatsushima, Japan for introducing the patient.

References

- [1] Pajvani U, Zeikus PS, Basile O, Toback N, Robinson-Bostom L. Thrombogenic vasculopathy with diffuse neutrophilic inflammation: a histologic manifestation of a tick bite. *Cutis* 2006;78:321–4.
- [2] Taylor D. Innate immunity in ticks: a review. *J Acarol Soc Jpn* 2006;15:109–27.
- [3] Murasawa K, Kimura T. Histopathological analysis of 62 cases of tick bite with tick. *Jpn J Dermatol* 2005;115:571–8.
- [4] Bowman AS, Coons LB, Needham GR, Sauer JR. Tick saliva: recent advances and implications for vector competence. *Med Vet Entomol* 1997;11:277–85.
- [5] Isoda K, Murase K, Okumura E, Uchikawa K. Histopathologic study on tick-bite wound. *Clin Parasitol* 1999;10:142–5.
- [6] Isoda K, Amanuma F, Murai T, Okumura E. Two cases of intractable tick-bite wound presenting cutaneous (localized) periarteritis nodosa. *Clin Parasitol* 2002;13:48–50.
- [7] McGinley-Smith DE, Tsao SS. Dermatoses from ticks. *J Am Acad Dermatol* 2003;49:363–92.

寄生虫疾患の各種診断法と漏らさないための 検査システムの提案

丸山 治彦

宮崎大学医学部 感染症学講座 寄生虫学分野

Key Words: 好酸球増多, 便検査, 抗体スクリーニング, 遺伝子スクリーニング

はじめに

宮崎大学医学部寄生虫学教室では、全国の医療機関からの血清診断依頼に対応しており、毎年新規症例の診断に関与している(表 1)。「寄生虫症免疫診断検査申込書」は、(株) エスアールエルを通して、あるいは当研究室のサイトからダウンロードできるようになっているが(<http://www.miyazaki-med.ac.jp/parasitology/detail.htm>)、この申込書には現病歴を簡単に記述するスペースがある。したがってわれわれは、それぞれの症例について、どのような経緯で主治医あるいは医療チームが寄生虫疾患を疑い、宮崎大学へ検査を依頼したのかをある程度知ることができる。その中のほとんどは合理的な検査依頼だが、中にははなはだ理解が難しい例もある。

ここでは、これまでに経験した寄生虫疾患の検査依頼や経過の具体例を通して、わが国における寄生虫疾患診断の問題点を明らかにし、その対策

にどのようなものがあり得るのかを提案する。なお、検査診断業務にともなう得られる臨床データの分析と発表については、宮崎大学医学部医の倫理委員会の承認を得ている。

検査依頼理由

1. 好酸球増多

もっとも多い検査依頼の理由が好酸球増多である。末梢血の好酸球増多はもちろん、胸水や腹水中、心嚢液中や脳脊髄液に多くの好酸球がみとめられた、気管支肺泡洗浄液中に好酸球多数がみとめられた、あるいは生検組織中への好酸球浸潤が著明であった場合などもある。

末梢血好酸球増多以外に異常所見がない例もあるが、好酸球の出現と同時に現れる所見で多いのは肺炎、とくに「抗菌薬を投与したが軽快しない肺炎」である。胸部 X 線所見はさまざまだが、肺野の異常陰影が多発性であり出没を繰り返し

Proposal of systematic screening for the diagnosis of parasitic diseases.

Haruhiko Maruyama

Division of Parasitology, Department of Infectious Diseases, Faculty of Medicine, University of Miyazaki

論文請求先: 丸山治彦 〒889-1692 宮崎市清武町木原 5200 宮崎大学医学部 感染症学講座 寄生虫学分野

表 1 宮崎大学医学部寄生虫学教室における寄生虫症診断実績

寄生虫	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
イヌ回虫・ブタ回虫	68	67	77	100	103	82	101	78	49	48
アニサキス	6	6	6	0	4	4	6	3	2	2
イヌ糸状虫	6	4	4	7	1	5	1	1	0	0
顎口虫	13	10	11	11	0	0	6	7	9	3
鉤虫	6	2	3	0	1	0	1	0	1	1
マンソン孤虫	4	8	6	5	4	3	6	4	5	2
囊虫	2	4	2	4	0	0	0	0	1	0
肺吸虫	37	36	32	45	30	37	46	38	38	45
肝蛭	1	1	8	5	6	2	3	1	1	3
住血吸虫	0	0	1	5	5	6	6	4	4	3
肝吸虫	1	3	1	1	0	0	0	0	0	1
糞線虫	8	21	11	11	2	1	1	2	0	2
回虫	3	3	4	0	1	1	1	2	0	0
広節・日本海裂頭条虫	0	0	2	1	0	2	0	1	0	4
計	155	165	168	195	157	143	178	141	110	114

たり移動する時には、原因が寄生虫である可能性がきわめて高い。申込書によると、たいてい他の原因検査も実行されており、各種細菌陰性、アスペルギルス陰性、細胞診陰性などの記述がみとめられる。マイコプラズマ肺炎と思われる症例で、回復期に好酸球が上昇してきたので検査を依頼する例もある。

呼吸器症状以外には、肝臓の異常陰影をはじめとしてさまざまな症状、所見が付随している。好酸球性髄膜炎では髄膜刺激症状がある。当然のことながら、これらの症例で抗体が陽性なのか陰性なのかは検査してみないと予想はつかない。しかしながら、透析患者にみとめられる好酸球増多症では、抗寄生虫抗体が陽性になることはまれである。

2. 脊髄炎、ぶどう膜炎

特徴的な臨床所見から好酸球増多の有無に関わらず寄生虫感染を疑う場合があり、特に目立つのは脊髄炎とぶどう膜炎である。これはトキソカラなどの回虫類幼虫による幼虫移行症を疑ったもので、脊髄炎では陽性の場合には比較的高い抗

体濃度が得られる。血清と髄液を同時に測定できれば診断の精度を上げることができる。ぶどう膜炎では通常血清抗体濃度は境界値前後であり、診断では眼科医の判断が最も重要である。硝子体液などの局所液が得られれば診断精度は大幅に上昇する。

好酸球性肺炎でも成り立つが、とくに脊髄炎とぶどう膜炎症例では、同じクリニックあるいは医療機関の診療科（それぞれ神経内科と眼科）からの依頼が多い。これは、一度寄生虫症例を経験した医療施設では、同様の症状に対して再び寄生虫感染の可能性を考えるからであろう。逆に言うと、一度も寄生虫疾患に遭遇したことがない、あるいは寄生虫疾患の存在を思いつかない医師しかいない医療施設では、いつまで経っても寄生虫感染を疑うことはないということになる。

3. 慢性の消化器症状

原因不明の下痢で寄生虫検査依頼を受けることがあるが、多くの場合、送付されてくる検体は血清である。好酸球性胃腸炎であれば抗体陽性の結果が得られることもあるが、好酸球増多が見ら

れない場合にはほぼ抗体陰性であり、仮に陽性反応が得られたとしても消化器症状との関連ははっきりしない（抗アニサキス抗体陽性など）。ただし、これらには報告書に便検査を実施するように促すコメントを付けるようにしているので、便検査を実行してもらえれば、検査依頼自体は必ずしも無駄ではない。

4. 炎症所見

頻度は多くはないが、何らかの炎症所見（発熱や CRP 値の上昇など）があるにも関わらず、通常の感染症や自己免疫疾患、悪性疾患が否定的である場合も、診断に苦慮して血清が送付されることがある。検査結果は陰性で寄生虫感染は否定的、との報告を返すことが多いが、寄生虫が鑑別に上がった点で評価すべきであると考えている。

5. その他

皮膚爬行疹や移動性皮下腫瘍も寄生虫感染を疑う所見で、実際にはほとんどの場合感染ありであろうと判断される。これらの症候は比較的わかりやすく見逃すことは少ないと考えられる。また、途上国への渡航歴があったり、患者が欧米以外、とくに東から東南アジアの出身者である場合には、寄生虫疾患が鑑別の対象になりやすい傾向があるように思われる。

寄生虫疾患のサインに気づかなかった症例

以上のように、いくつかの所見を契機にして寄生虫疾患の診断依頼が出されるわけだが、もちろんこの場合医師は寄生虫の存在に考えが及んでいる。問題なのは、寄生虫疾患を示すサインがあるにも関わらず寄生虫感染が全く鑑別に上らず、検査を依頼しない医師である。どのような場面かというと、それは上述した検査依頼理由の裏返しであり、① 好酸球増多があるにも関わらず寄生虫抗体検査をしなかった、② 慢性の下痢がある

にも関わらず便検査をしなかった、というふた通りに大別される。

1. 好酸球増多に注意を払わなかった肺吸虫症例

症例は男性会社員で、ある年の11月初旬から咳と血痰あり、翌年1月に近医を受診したところ胸部 X 線で左胸水と右上肺野に空洞性陰影をみとめた。末梢血好酸球増多（好酸球 18%）があり、ツベルクリン反応も強陽性であった。抗菌剤による治療を受けていたが血痰が持続して胸水も増加したため2月中旬に大規模総合病院を受診した。

総合病院への転院時検査成績では末梢血好酸球数 30%であり、一般細菌検査は常在菌のみ、抗酸菌培養陰性、結核菌に対する PCR は陰性であった。診断がつかないまま退院となったが胸水が増量し、4月初旬に再入院、4月10日の気管支鏡検査で喀痰中に肺吸虫卵が見いだされて診断がついた。

好酸球増多に対して早い段階で寄生虫抗体検査を実施していれば、初診から診断確定まで3ヶ月も要することはなかったであろうと考えられるが、この症例で示唆的と思われるのは、虫卵による診断がついた後に、血清が抗体検査のために送付されてきたことである。形態学的診断に対する評価が低いのか、あるいは抗体による診断が一般的とされるから機械的に検体を送付したのかは不明だが、いずれにしても、検査から得られる情報の診断的価値に対する順位付けに疑問を感じざるを得ない。

同様に当初から好酸球増多がありながら肺結核の治療がおこなわれた症例や、あるいは肺癌を疑って切除術を実施された例がいくつも報告されている^{1),2)}。

2. 慢性の下痢で便寄生虫検査をおこなわなかった糞線虫症例

糞線虫症では時に死亡例が報告されている。2001年に報告された例では、77歳女性が特にきっかけがなく腹部膨満感と全身浮腫および低蛋白血症が出現し、蛋白漏出性胃腸症を疑って治療していたが、大腸内視鏡の際に発見された炎症性変化に対しては抗生剤が投与されていた。この症例では、脂肪便検査の際にラブジチス型幼虫が発見され、上部内視鏡検査で粘膜内に多数の虫体が見とめられた。イベルメクチンを投与したが肺炎を併発して死亡した³⁾。

その他これまでに症例報告されているものでは、免疫抑制剤投与後に下痢が出現したが抗生剤の投与のみで対応し、消化器症状が改善しないために大学病院に転院し、胃粘膜生検で多数の糞線虫成虫を見とめた例⁴⁾、あるいは体重減少、食欲不振、腹痛、下痢などがあつたが諸病院へ入退院を繰り返して診断がつかず、拒食症(疑)とすらされていたが、翌年になって下痢が増強、嘔吐や浮腫も加わって入院し、内視鏡検査にて十二指腸粘膜に虫体を多数認めて診断に至った例などが報告されている⁵⁾。

いずれの例においても、もう少し早く寄生虫を念頭に置いた便検査が実施されていれば、異なった経過をとったであろうことは容易に推測される。

寄生虫疾患の見落としを減らすシステム

寄生虫疾患は、ほとんどの種について検査法が確立しており、どの施設がどの寄生虫検査を実施可能なのかも検索可能である(表2)。したがって、最も重要な点は寄生虫感染症が鑑別診断に含まれることである。しかしながら、卒前卒後教育における寄生虫疾患の占める割合は小さく、日常的に頻繁に遭遇する疾患でもない。したがって、寄生虫疾患の見落としを防ぐためには、皆が知識が

表2 主な寄生虫検査対応施設

原虫性疾患

クリプトスポリジウム、ランブル鞭毛虫、赤痢アメーバ：金沢大医学部寄生虫
赤痢アメーバ：国立感染研寄生動物部、慶応大医学部熱帯医学
トキソプラズマ：千葉大医学部寄生虫、金沢大医学部寄生虫
リーシュマニア：琉球大医学部皮膚科、金沢大医学部寄生虫
マラリア原虫：国際医療センター研究所熱帯医学マラリア研究部など

蠕虫性疾患

抗体スクリーニング：宮崎大医学部寄生虫
切片・虫体からの遺伝子検査：
国立感染研寄生動物部
形態学的鑑別：大分大学医学部寄生虫、目黒寄生虫館
エキノコックス、囊虫：旭川医大寄生虫
リンパ系フィラリア：愛知医大寄生虫
イヌ回虫、旋尾線虫、住血吸虫：
東京医科歯科大寄生虫

ないのが当たり前という点から出発しないといけないだろう。

それではどうすればよいのか。一つのヒントは「寄生虫抗体スクリーニング検査」にあるように思われる。この方式では、寄生虫のことはほとんど知らなくても、「好酸球増多」というキーワードに反応して検体を送付すれば、寄生虫感染の関与についてある程度の情報が得られる。「抗生剤の効かない肺炎」「原因不明の炎症」についても同様である。診療科によっては、これに「脊髄炎」「ぶどう膜炎」「皮膚爬行疹」のような症状も含まれるだろうが、基本的に細かい知識は要求されない。

寄生虫疾患の見落としを防ぐためには、血清診断以外にターゲットを絞らないスクリーニング的な検査法を設定すべきではないだろうか。そこで最もカバーすべき分野は、腸管内寄生虫症であろう。腸管内寄生虫に対しては抗体があまり産生されないため、抗体スクリーニングの検査項目には入っていない。しかしながら上述した糞線虫症のように重大な結果を招くこともあり得る。

幸いなことに、仮に慢性の下痢が続いているとして、その原因が寄生虫であれば、寄生虫由来の細胞成分は必ず便中に排出される。虫卵、幼虫、シスト、栄養体などである。また、寄生虫抗原も排出されている。したがって、「寄生虫抗体スクリーニング検査」のように、「寄生虫抗原または遺伝子スクリーニング検査」を構築できれば、「原因不明の下痢」というキーワードに反応して提出された検体に対して、寄生虫感染の有無について

ある程度の情報を得ることができるようになるだろう（図1）。

おそらく実現可能性という点から考えると、PCR法をベースにした「寄生虫遺伝子スクリーニング検査」の方が現実的であろう。抗原検出のためにはそれぞれの抗原に対して特異抗体を作製する必要があるが、PCR法ならプライマーの塩基配列を注意深く選べば、すぐにでも実行可能である。便検査において、虫卵、幼虫、シストなどを従来法によって検出しようとする、あらかじめ標的に合った検査法をオーダーする必要がある。しかしながら遺伝子スクリーニングであれば、発注側にはとくに要求される知識はない。

今後への提案

これまで、寄生虫学会ではさまざまな診断法が




図1 寄生虫疾患診断のための各種検査法