

Table 2. Univariate and Multivariate Analyses of Host and Viral Factors Associated with Low Hemoglobin Levels (< 11.0 g/dL) at Week 4 of Triple Therapy

Parameter	Univariate Analysis		Multivariate Analysis	
	OR (95% CI)	P	OR (95% CI)	P
Sex (female)	14.3 (4.1–50.0)	< 0.001	29.41 (3.8–250.0)	0.001
Age (> 50 years)	4.3 (1.0–17.5)	0.030	7.3 (1.1–47.6)	0.039
Body weight (< 60 kg)	11.5 (3.4–38.2)	< 0.001		
Body mass index (< 23)	8.4 (2.6–27.1)	< 0.001	17.2 (2.6–112.0)	0.003
Hemoglobin (< 15g/dL)	14.2 (3.5–57.4)	< 0.001		
<i>ITPA</i> gene (CC genotype)		0.062	36.8 (2.5–550.2)	0.009

Abbreviations: OR, odds ratio; CI, confidence level.

them, only age and body weight remained significant in the multivariate analysis.

Discussion

Anemia is a substantial risk in the standard of care therapy with PEG-IFN and RBV.^{4–6} Triphosphorylated RBV accumulates in erythrocytes of patients who receive RBV, increasingly with RBV dose and duration, and causes oxidative damage to erythrocyte membranes toward extravascular hemolysis by the reticuloendothelial system.^{19,20} Inosine triphosphate accumulates also in erythrocytes of individuals who have mutations in the *ITPA* gene, and results in benign red-cell enzymopathy.⁸ The expression of *ITPA* is genetically controlled and reduced in individuals who have point mutations in the *ITPA* gene.^{8–11} As another achievement of GWAS in hepatology,²¹ in the wake of polymorphisms of the *IL28B* gene that influence the response to PEG-IFN and RBV,^{22–24} polymorphisms in the *ITPA* gene has been reported to influence anemia caused by RBV.⁷ How inosine triphosphate protects erythrocytes from hemolysis caused by RBV needs to be sorted out by *in vivo* and *in vitro* experiments. Inosine triphosphate may prohibit the accumulation of RBV in erythrocytes, or rather, it might act directly toward prohibition of hemolysis.

In the present study, 61 patients infected with HCV-1 received triple therapy with PEG-IFN, RBV, and telaprevir in the first 12 weeks followed by PEG-IFN and RBV in the second 12 weeks. Then the RBV dose and hemoglobin were compared between patients with CC and CA/AA genotypes in the *ITPA* gene. Two polymorphisms in the *ITPA* gene, in close linkage disequilibrium with an r^2 value of 0.65,⁷ have been recognized in Caucasians (rs1127354 and rs7270107); the respective CA/AA and AC/CC genotypes decrease the activity of inosine triphosphatase and protect against anemia induced by RBV.^{7,12} Because the Japanese are monoallelic at rs7270107 and possess the AA

genotype exclusively,^{11,25} only polymorphisms at rs1127354 were examined.

Of the 61 patients, 49 possessed the RBV-sensitive CC genotype and the remaining 12 had RBV-resistant CA/AA genotypes. Hemoglobin levels decreased both in patients with CC and CA/AA genotypes. They lowered ≈ 4 g/dL during weeks 8–12 on the triple therapy with telaprevir, and increased thereafter (Fig. 2). Between the two groups of patients, differences in hemoglobin decrease were greatest at week 4 (1.3 g/dL), as in the standard treatment with PEG-IFN and RBV.^{7,12,13}

When anemia and other side effects occurred, doses of RBV, PEG-IFN, and telaprevir were modified. Of the 61 patients studied, 27 (44%) were women and most of them were in old age. Beyond 50 years of age, women are less responsive than men to the standard treatment with PEG-IFN and RBV, probably because estrogens with an antifibrotic potential decrease after menopause.²⁶ Stringent precautions had to be taken, therefore, by reducing the RBV dose in the patients in whom hemoglobin levels decreased <12 g/dL, rather than the conventional threshold of <10 g/dL.

Reductions of RBV dose due to anemia in patients who receive PEG-IFN and RBV are influenced by *ITPA* polymorphisms.¹² Also, in patients who had received the triple therapy the RBV dose had to be reduced more in

Table 3. Univariate and Multivariate Analyses of Host and Viral Factors Associated with Very Low Hemoglobin Levels (<8.5 g/dL) During Triple Therapy

Parameter	Univariate Analysis		Multivariate Analysis	
	OR (95% CI)	P	OR (95% CI)	P
Sex (female)	6.1 (1.5–25.1)	0.007		
Age (>60 years)	6.8 (1.8–26.0)	0.004	10.1 (1.9–53.9)	0.007
Body weight (<60 kg)	23.8 (2.9–200.0)	<0.001	33.3 (3.4–333.3)	0.003
Body mass index (<23)	14.1 (1.7–125.0)	0.001		
Hemoglobin (<14 g/dL)	4.3 (1.2–15.6)	0.023		

Abbreviations: OR, odds ratio; CI, confidence level.

patients with CC than CA/AA genotypes during the first 12 weeks while they received telaprevir ($52 \pm 14\%$ vs. $65 \pm 21\%$ of the target dose, $P = 0.039$). During the second 12 weeks off telaprevir, the RBV dose was somewhat greater in patients with CC than CA/AA genotypes ($47 \pm 24\%$ vs. $43 \pm 20\%$, $P = 0.649$). Thus, the total RBV dose during 24 weeks of therapy was comparable between patients with CC and CA/AA genotypes ($51 \pm 15\%$ and $57 \pm 18\%$, $P = 0.724$). Likewise, the total dose of PEG-IFN ($87 \pm 23\%$ vs. $86 \pm 20\%$ of the target, $P = 0.806$), as well as that of telaprevir ($87 \pm 27\%$ vs. $71 \pm 36\%$ of the target, $P = 0.098$), was no different between patients with CC and CA/AA genotypes. SVR was achieved comparably frequently in them (71% vs. 67% , $P = 0.736$).

Decreases in hemoglobin levels during the first 12 week were similar between the current triple therapy cohort and previous patients receiving PEG-IFN and RBV.^{12,13} The conservative hemoglobin levels chosen for RBV dose reduction may be a possible confounding factor on the impact of *ITPA* variants in anemia, which would have been greater should the RBV dose not be reduced in patients with RBV-sensitive CC genotypes.

ITPA polymorphisms at rs1127354 were associated with RBV-induced anemia in Japanese patients, without involvement of those at rs7270107 reported in Caucasian and African-American patients.¹³ Thus, *ITPA* polymorphisms at rs1127354 would play a major role in protecting patients from RBV-induced anemia. CC/CA genotypes at rs1127354 occurs in 6% of the Caucasian population, much less often in the Oriental population, at 16%.^{25,27} Although AC/CC genotypes at rs7270107 occurs in 13% of Caucasians, they do not exist in Orientals.^{11,25} Obviously, different polymorphisms need to be examined in patients of distinct ethnicities when the influence on RBV-induced anemia is to be evaluated.

In confirmation of our previous report,²⁸ the triple therapy achieved SVR more frequently in patients with CC than CT/TT genotypes of *IL28* at rs12979860 (96% vs. 41% , $P < 0.001$). About two-thirds of studied patients accomplished SVR with the triple treatment, although one-fourth of them were nonresponders to previous IFN-based treatments; they are known to respond poorly to repeated treatments. This would lend further support to the efficacy of triple therapy being higher than treatment with pegylated IFN and RBV.

There are strong points in this study. First, *ITPA* polymorphisms influence RBV-induced anemia in the triple therapy. Second, polymorphisms at rs1127350, without involvement of those at rs7270107, protect against RBV-induced anemia. Third, the triple therapy can be applied with high efficacy by careful monitoring of hemoglobin

and prompt modification of RBV dose. There are weak points in this study as well. First, it was a retrospective cohort study conducted in a small size of patients, especially those with CA/AA genotypes at rs1127350, and included null-responders to previous IFN-based therapies; the real impact of *ITPA* polymorphisms on RBV-induced anemia may have been obscured. Second, the study was conducted in Japanese patients, and the results may or may not be extended to patients of different ethnicities with distinct genetic backgrounds. Hopefully, the results presented herein will promote future studies in which the influence of the *ITPA* polymorphism on RBV-induced anemia will be pursued in larger scale and on patients of various ethnicities around the world.

References

- Shepard CW, Finelli L, Alter MJ. Global epidemiology of hepatitis C virus infection. *Lancet Infect Dis* 2005;5:558-567.
- Hoofnagle JH. Course and outcome of hepatitis C. *HEPATOLOGY* 2002;36:S21-29.
- Seeff LB. Natural history of chronic hepatitis C. *HEPATOLOGY* 2002;36:S35-46.
- Fried MW, Shiffman ML, Reddy KR, Smith C, Marinos G, Goncalves FL Jr, et al. Peginterferon alfa-2a plus ribavirin for chronic hepatitis C virus infection. *N Engl J Med* 2002;347:975-982.
- Manns MP, McHutchison JG, Gordon SC, Rustgi VK, Shiffman M, Reindollar R, et al. Peginterferon alfa-2b plus ribavirin compared with interferon alfa-2b plus ribavirin for initial treatment of chronic hepatitis C: a randomised trial. *Lancet* 2001;358:958-965.
- McHutchison JG, Lawitz EJ, Shiffman ML, Muir AJ, Galler GW, McCone J, et al. Peginterferon alfa-2b or alfa-2a with ribavirin for treatment of hepatitis C infection. *N Engl J Med* 2009;361:580-593.
- Fellay J, Thompson AJ, Ge D, Gumbs CE, Urban TJ, Shianna KV, et al. *ITPA* gene variants protect against anaemia in patients treated for chronic hepatitis C. *Nature* 2010;464:405-408.
- Arenas M, Duley J, Sumi S, Sanderson J, Marinaki A. The *ITPA* c.94C>A and g.IVS2+21A>C sequence variants contribute to missplicing of the *ITPA* gene. *Biochim Biophys Acta* 2007;1772:96-102.
- Cao H, Hegele RA. DNA polymorphisms in *ITPA* including basis of inosine triphosphatase deficiency. *J Hum Genet* 2002;47:620-622.
- Stepchenkova EI, Tarakhovskaya ER, Spitler K, Frahm C, Menezes MR, Simone PD, et al. Functional study of the P32T *ITPA* variant associated with drug sensitivity in humans. *J Mol Biol* 2009;392:602-613.
- Sumi S, Marinaki AM, Arenas M, Fairbanks L, Shobowale-Bakre M, Rees DC, et al. Genetic basis of inosine triphosphate pyrophosphohydrolase deficiency. *Hum Genet* 2002;111:360-367.
- Thompson AJ, Fellay J, Patel K, Tillmann HL, Naggie S, Ge D, et al. Variants in the *ITPA* gene protect against ribavirin-induced hemolytic anemia and decrease the need for ribavirin dose reduction. *Gastroenterology* 2010;139:1181-1189.
- Ochi H, Mackawa T, Abe H, Hayashida Y, Nakano R, Kubo M, et al. *ITPA* polymorphism affects ribavirin-induced anemia and outcome of therapy — a genome-wide study of Japanese HCV patients. *Gastroenterology* 2010;139:1190-1197.
- Hezode C, Forestier N, Dusheiko G, Ferenci P, Pol S, Goester T, et al. Telaprevir and peginterferon with or without ribavirin for chronic HCV infection. *N Engl J Med* 2009;360:1839-1850.
- McHutchison JG, Everson GT, Gordon SC, Jacobson IM, Sulkowski M, Kauffman R, et al. Telaprevir with peginterferon and ribavirin for chronic HCV genotype 1 infection. *N Engl J Med* 2009;360:1827-1838.

16. Mederacke I, Wedemeyer H, Manns MP. Boceprevir, an NS3 serine protease inhibitor of hepatitis C virus, for the treatment of HCV infection. *Curr Opin Investig Drugs* 2009;10:181-189.
17. Ohnishi Y, Tanaka T, Ozaki K, Yamada R, Suzuki H, Nakamura Y. A high-throughput SNP typing system for genome-wide association studies. *J Hum Genet* 2001;46:471-477.
18. Suzuki A, Yamada R, Chang X, Tokuhira S, Sawada T, Suzuki M, et al. Functional haplotypes of PADI4, encoding citrullinating enzyme peptidylarginine deiminase 4, are associated with rheumatoid arthritis. *Nat Genet* 2003;34:395-402.
19. De Franceschi L, Fattovich G, Turrini F, Ayi K, Brugnara C, Manzato F, et al. Hemolytic anemia induced by ribavirin therapy in patients with chronic hepatitis C virus infection: role of membrane oxidative damage. *HEPATOLOGY* 2000;31:997-1004.
20. Russmann S, Grattagliano I, Portincasa P, Palmieri VO, Palasciano G. Ribavirin-induced anemia: mechanisms, risk factors and related targets for future research. *Curr Med Chem* 2006;13:3351-3357.
21. Karlsen TH, Melum E, Franke A. The utility of genome-wide association studies in hepatology. *HEPATOLOGY* 2010;51:1833-1842.
22. Ge D, Fellay J, Thompson AJ, Simon JS, Shianna KV, Urban TJ, et al. Genetic variation in IL28B predicts hepatitis C treatment-induced viral clearance. *Nature* 2009;461:399-401.
23. Suppiah V, Moldovan M, Ahlenstiel G, Berg T, Weltman M, Abate ML, et al. IL28B is associated with response to chronic hepatitis C interferon-alpha and ribavirin therapy. *Nat Genet* 2009;41:1100-1104.
24. Tanaka Y, Nishida N, Sugiyama M, Kurosaki M, Matsuura K, Sakamoto N, et al. Genome-wide association of IL28B with response to pegylated interferon-alpha and ribavirin therapy for chronic hepatitis C. *Nat Genet* 2009;41:1105-1109.
25. Maeda T, Sumi S, Ueta A, Ohkubo Y, Ito T, Marinaki AM, et al. Genetic basis of inosine triphosphate pyrophosphohydrolase deficiency in the Japanese population. *Mol Genet Metab* 2005;85:271-279.
26. Sezaki H, Suzuki F, Kawamura Y, Yatsuji H, Hosaka T, Akuta N, et al. Poor response to pegylated interferon and ribavirin in older women infected with hepatitis C virus of genotype 1b in high viral loads. *Dig Dis Sci* 2009;54:1317-1324.
27. Atanasova S, Shipkova M, Svinarov D, Mladenova A, Genova M, Wieland E, et al. Analysis of ITPA phenotype-genotype correlation in the Bulgarian population revealed a novel gene variant in exon 6. *Ther Drug Monit* 2007;29:6-10.
28. Akuta N, Suzuki F, Hirakawa M, Kawamura Y, Yatsuji H, Sezaki H, et al. Amino acid substitution in hepatitis C virus core region and genetic variation near the interleukin 28B gene predict viral response to telaprevir with peginterferon and ribavirin. *HEPATOLOGY* 2010;52:421-429.

Amino Acid Substitution in Hepatitis C Virus Core Region and Genetic Variation Near the Interleukin 28B Gene Predict Viral Response to Telaprevir with Peginterferon and Ribavirin

Norio Akuta,¹ Fumitaka Suzuki,¹ Miharuru Hirakawa,¹ Yusuke Kawamura,¹ Hiromi Yatsuji,¹ Hitomi Sezaki,¹ Yoshiyuki Suzuki,¹ Tetsuya Hosaka,¹ Masahiro Kobayashi,¹ Mariko Kobayashi,² Satoshi Saitoh,¹ Yasuji Arase,¹ Kenji Ikeda,¹ Kazuaki Chayama,³ Yusuke Nakamura,⁴ and Hiromitsu Kumada¹

Genetic variation near the IL28B gene and substitution of amino acid (aa) 70 and 91 in the core region of hepatitis C virus (HCV) genotype 1b can predict the response to pegylated interferon (PEG-IFN)/ribavirin combination therapy, but its impact on triple therapy of telaprevir/PEG-IFN/ribavirin is not clear. The aims of this study were to investigate the predictive factors of sustained virological response to a 12-week or 24-week regimen of triple therapy in 72 of 81 Japanese adults infected with HCV genotype 1. Overall, sustained virological response and end-of-treatment response were achieved by 61% and 89%, respectively. Especially, the sustained virological response was achieved by 45% and 67% in the 12- and 24-week regimens, respectively. Multivariate analysis identified rs8099917 near the IL28B gene (genotype TT) and substitution at aa 70 (Arg70) as significant determinants of sustained virological response. Prediction of response to therapy based on a combination of these factors had high sensitivity, specificity, and positive and negative predictive values. The efficacy of triple therapy was high in the patients with genotype TT, who accomplished sustained virological response (84%), irrespective of substitution of core aa 70. In the patients having genotype non-TT, those of Arg70 gained high sustained virological response (50%), and sustained virological response (12%) was the worst in patients who possessed both genotype non-TT and Gln70(His70). **Conclusion:** This study identified genetic variation near the IL28B gene and aa substitution of the core region as predictors of sustained virological response to a triple therapy of telaprevir/PEG-IFN/ribavirin in Japanese patients infected with HCV genotype 1b. (HEPATOLOGY 2010;52:421-429)

Abbreviations: aa, amino acid; ALT, alanine aminotransferase; AST, aspartate aminotransferase; γ GTP, gamma-glutamyl transpeptidase; HBsAg, hepatitis B surface antigen; HCC, hepatocellular carcinoma; HCV, hepatitis C virus; IFN, interferon; NPV, negative predictive value; PEG-IFN, pegylated interferon; PPV, positive predictive value

From the ¹Department of Hepatology, Toranomon Hospital, Tokyo, Japan; ²Liver Research Laboratory, Toranomon Hospital, Tokyo, Japan; ³Department of Medical and Molecular Science, Division of Frontier Medical Science, Programs for Biomedical Research, Graduate School of Biomedical Science, Hiroshima University, Hiroshima, Japan; ⁴Laboratory of Molecular Medicine, Human Genome Center, Institute of Medical Science, University of Tokyo, Tokyo, Japan.

Received February 4, 2010; accepted March 17, 2010.

Address reprint requests to: Norio Akuta, M.D., Department of Hepatology, Toranomon Hospital, 2-2-2 Toranomon, Minato-ku, Tokyo, 105-0001, Japan. E-mail: akuta-gi@umin.ac.jp; fax: +81-000-000000.

Copyright © 2010 by the American Association for the Study of Liver Diseases.

Published online in Wiley InterScience (www.interscience.wiley.com).

DOI 10.1002/hep.23690

Potential conflict of interest: Nothing to report.

Hepatitis C virus (HCV) usually causes chronic infection that can result in chronic hepatitis, liver cirrhosis, and hepatocellular carcinoma (HCC).^{1,2} At present, treatments based on interferon (IFN), in combination with ribavirin, are the mainstay for combating HCV infection. In Japan, HCV genotype 1b (HCV-1b) in high viral loads (>100 KIU/mL) accounts for more than 70% of HCV infections, making it difficult to treat patients with chronic hepatitis C.³ Such background calls for efficient treatments of Japanese patients with chronic HCV infection.

Even with pegylated IFN (PEG-IFN) combined with ribavirin, a sustained virological response lasting over 24 weeks after the withdrawal of treatment is achieved in at most 50% of the patients infected with HCV-1b and high viral loads.^{4,5} Recently, a new strategy was introduced in the treatment of chronic HCV infection by

means of inhibiting protease in the NS3/NS4 of the HCV polyprotein. Of these, telaprevir (VX-950) was selected as a candidate agent for treatment of chronic HCV infection.⁶ Later, it was found that telaprevir, when combined with PEG-IFN and ribavirin, gains a robust antiviral activity.^{7,8} Specifically, HCV RNA is suppressed below the limits of detection in the blood in almost all patients infected with HCV-1 during triple therapy of telaprevir with PEG-IFN and ribavirin.⁹ However, treatment-resistant patients who do not achieve sustained virological response by the triple therapy have been reported.⁹⁻¹¹ The underlying mechanism of the response to the treatment is still not clear.

Amino acid (aa) substitutions at position 70 and/or 91 in the HCV core region of patients infected with HCV-1b and high viral loads are pretreatment predictors of poor virological response to PEG-IFN plus ribavirin combination therapy,¹²⁻¹⁴ and also affect clinical outcome, including hepatocarcinogenesis.^{15,16} Furthermore, a recent report showed that aa substitutions in the core region can also be used before therapy to predict very early dynamics (within 48 hours) after the start of triple therapy of telaprevir with PEG-IFN and ribavirin.¹⁷ However, it is not clear at this stage whether aa substitutions in the core region can be used before therapy to predict sustained virological response to triple therapy.

Recent reports showed that genetic variations near the IL28B gene (rs8099917, rs12979860) on chromosome 19 is a host-related factor, which encodes IFN- λ -3, are pretreatment predictors of virological response to 48-week PEG-IFN plus ribavirin combination therapy in individuals infected with HCV-1,¹⁸⁻²¹ and also affect clinical outcome, including spontaneous clearance of HCV.²² However, it is not clear at this stage whether genetic variation near the IL28B gene can be used before therapy to predict sustained virological response to triple therapy.

The present study included 81 patients with HCV-1b and high viral loads who received the triple therapy of telaprevir with PEG-IFN plus ribavirin. The aims of the study were to identify the pretreatment factors that could predict sustained virological response, including viral- (aa substitutions in the HCV core and NS5A regions) and host-related factors (genetic variation near the IL28B gene).

Patients and Methods

Study Population. Between May 2008 and September 2009, 81 patients infected with HCV were

recruited for this study at the Department of Hepatology in Toranomon Hospital in Metropolitan Tokyo. The study protocol was in compliance with the Good Clinical Practice Guidelines and the 1975 Declaration of Helsinki and was approved by the Institutional Review Board. Each patient gave informed consent before participating in this trial. Patients were divided into two groups: 20 (25%) patients were allocated to a 12-week regimen of triple therapy (telaprevir [MP-424], PEG-IFN, and ribavirin) (the T12PR12 group), and 61 patients (75%) were assigned to a 24-week regimen of the same triple therapy for 12 weeks followed by dual therapy of PEG-IFN and ribavirin for 12 weeks (the T12PR24 group).

All of 81 patients met the following inclusion and exclusion criteria: (1) diagnosis of chronic hepatitis C. (2) HCV-1 confirmed by sequence analysis. (3) HCV RNA levels of ≥ 5.0 log IU/mL determined by the COBAS TaqMan HCV test (Roche Diagnostics, Tokyo, Japan). (4) Japanese (Mongoloid) ethnicity. (5) Age at study entry of 20-65 years. (6) Body weight ≥ 35 kg and ≤ 120 kg at the time of registration. (7) Lack of decompensated liver cirrhosis. (8) Negativity for hepatitis B surface antigen (HBsAg) in serum. (9) Negative history of HCC. (10) No previous treatment for malignancy. (11) Negative history of autoimmune hepatitis, alcohol liver disease, hemochromatosis, and chronic liver disease other than chronic hepatitis C. (12) Negative history of depression, schizophrenia or suicide attempts, hemoglobinopathies, angina pectoris, cardiac insufficiency, myocardial infarction or severe arrhythmia, uncontrollable hypertension, chronic renal dysfunction or creatinine clearance of ≤ 50 mL/minute at baseline, diabetes requiring treatment or fasting glucose level of ≥ 110 mg/dL, autoimmune disease, cerebrovascular disorders, thyroidal dysfunction uncontrollable by medical treatment, chronic pulmonary disease, allergy to medication or anaphylaxis at baseline. (13) Hemoglobin level of ≥ 12 g/dL, neutrophil count $\geq 1500/\text{mm}^3$, and platelet count of $\geq 100,000/\text{mm}^3$ at baseline. Pregnant or breast-feeding women or those willing to become pregnant during the study and men with a pregnant partner were excluded from the study. Furthermore, 72 of 81 patients were followed for at least 24 weeks after the completion of triple therapy. The treatment efficacy was evaluated by HCV-RNA negative at the end of treatment (end-of-treatment response) and 24 weeks after the completion of therapy (sustained virological response), based on the COBAS TaqMan HCV test (Roche Diagnostics).

Telaprevir (MP-424; Mitsubishi Tanabe Pharma, Osaka, Japan) was administered at 750 mg or 500 mg

Table 1. Profile and Laboratory Data at Commencement of Telaprevir, Peginterferon and Ribavirin Triple Therapy in Japanese Patients Infected with HCV Genotype 1

Demographic data	
Number of patients	81
Sex (M/F)	44 / 37
Age (years)*	55 (23-65)
History of blood transfusion	24 (29.6%)
Family history of liver disease	13 (16.0%)
Body mass index (kg/m ²)*	22.5 (13.2-32.4)
Laboratory data*	
HCV genotype (1a/ 1b)	1/80
Level of viremia (log IU/mL)	6.7 (5.1-7.6)
Serum aspartate aminotransferase (IU/L)	34 (15-137)
Serum alanine aminotransferase (IU/L)	42 (12-175)
Serum albumin (g/dL)	3.9 (3.2-4.6)
Gamma-glutamyl transpeptidase (IU/L)	36 (9-229)
Leukocyte count (/mm ³)	4,800 (2,800-8,100)
Hemoglobin (g/dL)	14.3 (11.7-16.8)
Platelet count ($\times 10^4$ /mm ³)	17.1 (9.1-33.8)
Alpha-fetoprotein (μ g/L)	4 (2-39)
Total cholesterol (mg/dL)	180 (110-276)
Fasting plasma glucose (mg/dL)	92 (64-125)
Treatment	
PEG-IFN α -2b dose (μ g/kg)*	1.5 (1.3-2.0)
Ribavirin dose (mg/kg)*	11.7 (7.2-18.4)
Telaprevir dose (1,500 / 2,250 mg/day)	10/71
Treatment regimen (T12PR12 group / T12PR24 group)	20/61
Amino acid substitutions in the HCV genotype 1b	
Core aa 70 (arginine / glutamine [histidine] / ND)	47/33/1
Core aa 91 (leucine / methionine / ND)	43/37/1
ISDR of NS5A (wild-type / non wild-type / ND)	76/4/1
Genetic variation near IL28B gene	
rs8099917 genotype (TT / TG / GG / ND)	42/30/2/7
rs 12979860 genotype (CC / CT / TT / ND)	42/32/2/5
Past history of IFN therapy	
Treatment-naive / Relapsers to previous treatment / nonresponders to previous treatment	27/33/21

Data are number and percentages of patients, except those denoted by asterisk (*), which represent the median (range) values. ND, not determined.

three times a day at an 8-hour (q8) interval after the meal. PEG-IFN α -2b (PEG-Intron; Schering Plough, Kenilworth, NJ) was injected subcutaneously at a median dose 1.5 μ g/kg (range: 1.3-2.0 μ g/kg) once a week. Ribavirin (Rebetol; Schering Plough) was administered at 200-600 mg twice a day after breakfast and dinner (daily dose: 600-1000 mg).

PEG-IFN and ribavirin were discontinued or their doses reduced, as required, upon reduction of hemoglobin level, leukocyte count, neutrophil or platelet count, or the development of adverse events. Thus, the dose of PEG-IFN was reduced by 50% when the leukocyte count decreased below 1500/mm³, neutrophil count below 750/mm³, or platelet count below 80,000/mm³; PEG-IFN was discontinued when these counts decreased below 1000/mm³, 500/mm³ or 50,000/mm³, respectively. When hemoglobin decreased to <10 g/dL, the daily dose of ribavirin was reduced from 600 to 400 mg, from 800 to 600 mg

and 1000 mg to 600 mg, depending on the initial dose. Ribavirin was withdrawn when hemoglobin decreased to <8.5 g/dL. However, the dose of telaprevir (MP-424) remained the same, and its administration was stopped when the discontinuation was appropriate for the development of adverse events. In those patients who discontinued telaprevir, treatment with PEG-IFN α -2b and ribavirin was also terminated.

Table 1 summarizes the profiles and laboratory data of the 81 patients at the commencement of treatment. They included 44 males and 37 females, ages 23 to 65 years (median, 55 years).

Measurement of HCV RNA. The antiviral effects of the triple therapy on HCV were assessed by measuring plasma HCV RNA levels. In this study, HCV RNA levels during treatment were evaluated at least once every month before, during, and after therapy. HCV RNA concentrations were determined using the COBAS TaqMan HCV test (Roche Diagnostics). The linear dynamic range of the assay was 1.2-7.8 log IU/mL, and the undetectable samples were defined as negative.

Detection of Amino Acid Substitutions in Core and NS5A Regions of HCV-1b. In the present study, aa substitutions of the core region and NS5A-ISDR (IFN-sensitivity determining region) of HCV-1b were analyzed by direct sequencing. HCV RNA was extracted from serum samples at the start of treatment and reverse transcribed with random primer and MMLV reverse transcriptase (Takara Syuzo, Tokyo). Nucleic acids were amplified by polymerase chain reaction (PCR) using the following primers: (1) Nucleotide sequences of the core region: The first-round PCR was performed with CE1 (sense, 5'-GTC TGC GGA ACC GGT GAG TA-3', nucleotides: 134-153) and CE2 (antisense, 5'-GAC GTG GCG TCG TAT TGT CG-3', nucleotides: 1096-1115) primers, and the second-round PCR with CC9 (sense, 5'-ACT GCT AGC CGA GTA GTG TT-3', nucleotides: 234-253) and CE6 (antisense, 5'-GGA GCA GTC GTT CGT GAC AT-3', nucleotides: 934-953) primers. (2) Nucleotide sequences of NS5A-ISDR: The first-round PCR was performed with ISDR1 (sense, 5'-ATG CCC ATG CCA GGT TCC AG-3', nucleotides: 6662-6681) and ISDR2 (antisense, 5'-AGC TCC GCC AAG GCA GAA GA-3', nucleotides: 7350-7369) primers, and the second-round PCR with ISDR3 (sense, 5'-ACC GCA TGT GGC AGT GCT CA-3', nucleotides: 6824-6843) and ISDR4 (antisense, 5'-GTA ATC CGG GCG TGC CCA TA-3', nucleotides: 7189-7208) primers. ([1,2]; nested PCR.) All samples were initially denatured at 95°C for 2 minutes. The 35 cycles of

amplification were set as follows: denaturation for 30 seconds at 95°C, annealing of primers for 30 seconds at 55°C, and extension for 1 minute at 72°C with an additional 7 minutes for extension. Then 1 μ L of the first PCR product was transferred to the second PCR reaction. Other conditions for the second PCR were the same as the first PCR, except that the second PCR primers were used instead of the first PCR primers. The amplified PCR products were purified by the QIA quick PCR purification kit (Qiagen, Tokyo) after agarose gel electrophoresis and then used for direct sequencing. Dideoxynucleotide termination sequencing was performed with the Big Dye Deoxy Terminator Cycle Sequencing kit (PerkinElmer, Tokyo).

With the use of HCV-J (Access. No. D90208) as a reference,²³ the sequence of 1-191 aa in the core protein of HCV-1b was determined and then compared with the consensus sequence constructed on 81 clinical samples to detect substitutions at aa 70 of arginine (Arg70) or glutamine/histidine (Gln70/His70) and aa 91 of leucine (Leu91) or methionine (Met91).¹² The sequence of 2209-2248 aa in the NS5A of HCV-1b (ISDR) reported by Enomoto et al.²⁴ was determined and the numbers of aa substitutions in ISDR were defined as wildtype (0, 1) or nonwildtype (≥ 2).

Genetic Variation Near the IL28B Gene. Samples for genome-wide association survey were genotyped using the Illumina HumanHap610-Quad Genotyping BeadChip. Genotyping data were subjected to quality control before the data analysis. Genotyping for replication and fine mapping was performed by use of the Invader assay, TaqMan assay, or direct sequencing as described.^{25,26}

In this study, genetic variations near the IL28B gene (rs8099917, rs12979860), reported as the pretreatment predictors of treatment efficacy and clinical outcome,¹⁸⁻²² were investigated.

Statistical Analysis. Nonparametric tests (chi-squared test and Fisher's exact probability test) were used to compare the characteristics of the groups. Univariate and multivariate logistic regression analyses were used to determine those factors that significantly contributed to sustained virological response. The odds ratios (OR) and 95% confidence intervals (95% CI) were also calculated. All *P* values less than 0.05 by the two-tailed test were considered significant. Variables that achieved statistical significance (*P* < 0.05) on univariate analysis were entered into multiple logistic regression analysis to identify significant independent predictive factors. Each variable was transformed into categorical data consisting of two simple ordinal numbers for univariate and multivariate analyses. The

potential pretreatment factors associated with sustained virological response included the following variables: sex, age, history of blood transfusion, family history of liver disease, body mass index, aspartate aminotransferase (AST), alanine aminotransferase (ALT), albumin, gamma-glutamyl transpeptidase (γ GTP), leukocyte count, hemoglobin, platelet count, HCV RNA level, alfa-fetoprotein, total cholesterol, fasting blood sugar, PEG-IFN dose/body weight, ribavirin dose/body weight, telaprevir dose/day, treatment regimen of triple therapy, past history of IFN therapy, genetic variation near the IL28B gene, and aa substitution in the core region, and NS5A-ISDR. Statistical analyses were performed using SPSS (Chicago, IL). Sensitivity, specificity, positive predictive value (PPV), and negative predictive value (NPV) were also calculated to determine the reliability of predictors of the response to therapy.

Results

Virological Response to Therapy. Sustained virological response was achieved by 44 of 72 (61.1%) patients. In all, 64 of 72 (88.9%) patients were considered end-of-treatment response. According to treatment regimen, sustained virological response were achieved by 45.0% (9 of 20 patients) and 67.3% (35 of 52 patients), in the T12PR12 group and the T12PR24 group, respectively. Of eight patients who could not achieve end-of-treatment response, six (75.0%) patients resulted in reevaluation of viral loads regardless of HCV-RNA temporary negative, and the other two patients (25.0%) did not achieve HCV-RNA negative during treatment.

Especially in the T12PR24 group, according to the past history of treatment, sustained virological response were achieved by 76.4% (13 of 17 patients), 86.4% (19 of 22 patients), and 23.1% (3 of 13 patients), in treatment-naive, relapsers to previous treatment, and nonresponders to previous treatment, respectively.

Sustained Virological Response According to Amino Acid Substitutions in Core and NS5A Regions. According to the substitution of core aa 70, a significantly higher proportion of patients with Arg70 substitutions (74.4%) showed sustained virological response than that of patients who showed Gln70(His70) (41.4%) (Fig. 1, *P* = 0.007). In contrast, according to the substitution of core aa 91, the sustained virological response rate was not significantly different between Leu91 (65.0%) and Met91 (56.3%) (Fig. 1). Likewise, according to the numbers of aa substitutions in ISDR, the sustained virological response rate was not significantly different between wildtype




Fig. 1. According to the substitution of core aa 70, a significantly higher proportion of patients with Arg70 substitutions showed sustained virological response than that of patients who showed Gln70(His70) ($P = 0.007$). In contrast, according to the substitution of core aa 91, the sustained virological response rate was not significantly different between Leu91 and Met91. Likewise, according to the numbers of aa substitutions in ISDR, the sustained virological response rate was not significantly different between wildtype and nonwildtype.

(56.3%) and nonwildtype (66.7%) (Fig. 1). Thus, sustained virological response was influenced by the substitution of core aa 70.

Sustained Virological Response According to Genetic Variation Near the IL28B Gene. According to the genetic variation in rs8099917, sustained virological response was achieved by 83.8% (31 of 37 patients), 29.6% (8 of 27 patients), and 0% (0 of 2 patients) in patients with genotype TT, TG, and GG, respectively. Thus, a significantly higher proportion of patients with genotype TT (83.8%) showed sustained virological response than that of patients who showed genotype non-TT (27.6%) (Fig. 2, $P < 0.001$) (Table 2).

According to the genetic variation in rs12979860, sustained virological response was achieved by 83.8% (31 of 37 patients), 34.5% (10 of 29 patients), and 0% (0 of 2 patients), in patients with genotype CC, CT, and TT, respectively. Thus, a significantly higher proportion of patients with genotype CC (83.8%) showed sustained virological response than that of patients who showed genotype non-CC (32.3%) (Fig. 2, $P < 0.001$) (Table 2).

Predictive Factors Associated with Sustained Virological Response. Univariate analysis identified three parameters that correlated with sustained virological response significantly: substitution of aa 70 (Arg70; OR 4.12, $P = 0.007$), genetic variation in rs8099917 (genotype TT; OR 13.6, $P < 0.001$), and rs12979860 (genotype CC; OR 10.8, $P < 0.001$). Two factors were identified by multivariate analysis as independent

parameters that significantly influenced sustained virological response (rs8099917 genotype TT; OR 10.6, $P < 0.001$; and Arg70; OR 3.69, $P = 0.040$) (Table 3).

Assessment of Amino Acid Substitutions in Core Region and Genetic Variation Near the IL28B Gene as Predictors of Sustained Virological Response. The ability to predict sustained virological response by substitution of core aa 70 and rs8099917 genotype near the IL28B gene was evaluated. The sustained virological response rates of patients with a combination of Arg70 or rs8099917 genotype TT were defined as PPV (prediction of sustained virological response). The nonsustained virological response rates of patients with a combination of Gln70(His70) or rs8099917 genotype non-TT were defined as NPV (prediction of nonsustained virological response).

In patients with rs8099917 genotype TT, the sensitivity, specificity, PPV, and NPV for sustained virological response were 79.5, 77.8, 83.8, and 72.4%, respectively. Thus, genotype TT has high sensitivity, specificity, and PPV for prediction of sustained virological response. In patients with Arg70 the sensitivity, specificity, PPV, and NPV were 76.9, 63.0, 75.0, and 65.4%, respectively. Thus, Arg70 has high sensitivity and PPV in predicting sustained virological response. Furthermore, when both predictors were used the sensitivity, specificity, PPV, and NPV were 61.5, 85.2, 85.7, and 60.5%, respectively. When one or more of the two predictors were used the sensitivity, specificity, PPV, and NPV were 94.9, 55.6, 75.5, and 88.2%, respectively. These results indicate that the use of the combination of the above two predictors has high sensitivity, specificity, PPV, and NPV for prediction of sustained virological response (Table 4).




Fig. 2. According to the genetic variation in rs8099917 or rs12979860 near the IL28B gene, a significantly higher proportion of patients with genotype TT or CC showed sustained virological response than that of patients who showed genotype non-TT or non-CC, respectively ($P < 0.001$ or $P < 0.001$, respectively).

Table 2. According to Genetic Variation Near the IL28B Gene, Background at Commencement of Triple Therapy and Treatment Efficacy

	rs8099917 genotype			rs12979860 genotype		
	TT (n = 42)	TG+GG (n = 32)	TT vs. TG+GG P	CC (n = 42)	CT+TT (n = 34)	CC vs. CT+TT P
Demographic data						
Sex (M/F)	22 / 20	18 / 14	NS	22 / 20	19 / 15	NS
Age (years)*	54 (23-65)	56 (36-65)		54 (23-65)	55 (36-65)	NS
History of blood transfusion	15 (35.7%)	9 (28.3%)	NS	15 (35.7%)	9 (26.5%)	NS
Family history of liver disease	6 (14.3%)	6 (18.8%)	NS	6 (14.3%)	6 (17.6%)	NS
Body mass index (kg/m ²)*	22.1 (13.2-32.4)	22.4 (18.7-26.5)	NS	22.1 (13.2-32.4)	22.3 (18.7-26.5)	NS
Laboratory data*						
HCV genotype (1a / 1b)	0 / 42	1 / 31	NS	0 / 42	1 / 33	NS
Level of viremia (log IU/mL)	6.9 (5.4-7.5)	6.6 (5.1-7.4)	NS	6.9 (5.4-7.5)	6.5 (5.1-7.4)	NS
Serum aspartate aminotransferase (IU/L)	38 (15-118)	31 (20-137)	0.036	38 (15-118)	31 (20-137)	0.031
Serum alanine aminotransferase (IU/L)	50 (12-175)	36 (17-136)	0.029	50 (12-175)	35 (17-136)	0.014
Serum albumin (g/dL)	3.9 (3.3-4.6)	3.9 (3.2-4.6)	NS	3.9 (3.3-4.6)	3.9 (3.2-4.6)	NS
Gamma-glutamyl transpeptidase (IU/L)	29 (9-194)	53 (9-154)	0.008	29 (9-194)	53 (9-229)	0.004
Leukocyte count (/mm ³)	4,800 (2,800-8,100)	4,800 (3,000-7,800)	NS	4,800 (2,800-8,100)	4,800 (3,000-7,800)	NS
Hemoglobin (g/dL)	14.3 (12.3-16.5)	14.3 (11.7-16.8)	NS	14.3 (12.3-16.5)	14.3 (11.7-16.8)	NS
Platelet count ($\times 10^4$ /mm ³)	16.8 (9.9-33.8)	17.1 (9.1-24.8)	NS	16.8 (9.9-33.8)	17.8 (9.1-28.8)	NS
Alpha-fetoprotein (μ g/L)	4 (2-39)	5 (2-38)	NS	4 (2-39)	5 (2-38)	NS
Total cholesterol (mg/dL)	184 (112-276)	178 (110-263)	NS	184 (112-276)	178 (110-263)	NS
Fasting plasma glucose (mg/dL)	97 (80-125)	90 (66-111)	0.038	97 (80-125)	91 (66-111)	0.030
Treatment regimen						
T12PR12 group / T12PR24 group	12 / 30	7 / 25	NS	12 / 30	7 / 27	NS
Amino acid substitutions in the HCV genotype 1b						
Core aa 70 (arginine / glutamine [histidine])	30 / 12	13 / 18	0.016	30 / 12	13 / 20	0.009
Core aa 91 (leucine / methionine)	25 / 17	13 / 18	NS	25 / 17	14 / 19	NS
ISDR of NS5A (wild-type / non wild-type)	39 / 3	30 / 1	NS	39 / 3	32 / 1	NS
Past history of IFN therapy						
Treatment-naïve / Relapsers to previous treatment / Nonresponders to previous treatment	16 / 24 / 2	7 / 6 / 19	<0.001	16 / 24 / 2	8 / 7 / 19	<0.001
Treatment efficacy**						
End-of-treatment response (%)	35 (94.6%)	23 (79.3%)	NS	35 (94.6%)	25 (80.6%)	NS
Sustained virological response (%)	31 (83.8%)	8 (27.6%)	<0.001	31 (83.8%)	10 (32.3%)	<0.001

Data are number and percentages of patients, except those denoted by asterisk (*), which represent the median (range) values.

**Treatment efficacy according to rs8099917 genotype was evaluated in 66 patients, and that according to rs12979860 genotype was evaluated in 68 patients.

Predicting Sustained Virological Response by Amino Acid Substitutions in Core Region in Combination with Genetic Variation Near the IL28B Gene. Sustained virological response by core aa 70 in combination with rs8099917 genotype is shown in Fig. 3. In patients with rs8099917 genotype TT, sustained virological response was not different between Arg70 (85.7%) and Gln70(His70) (77.8%). In contrast, in patients with rs8099917 genotype TG and GG, a significantly higher proportion of patients with Arg70 (50.0%) showed sustained virological response than that of patients with Gln70(His70) (11.8%) ($P = 0.038$).

Based on a strong power of substitution of core aa 70 and rs8099917 genotype in predicting sustained virological response (Table 3), how they increase the predictive value when they were combined was evaluated. The results are schematically depicted in Fig. 3.

Together they demonstrate three points: (1) the efficacy of triple therapy was high in patients with genotype TT who accomplished sustained virological response at 83.8%, irrespective of substitution of core aa 70; (2) in patients having genotype TG and GG, those of Arg70 gained high sustained virological response (50.0%); and (3) sustained virological response (11.8%) was the worst in patients who possessed both of genotype TG and GG, and Gln70(His70).

Discussion

Two previous studies (PROVE1 in the US, and PROVE2 in Europe) showed that the T12PR12 and T12PR24 group of telaprevir, PEG-IFN, and ribavirin could achieve sustained virological response rates of 35%-60% and 61%-69%, respectively.^{10,11} In the

Table 3. Multivariate Analysis of Factors Associated with Sustained Virological Response of Telaprevir, Peginterferon and Ribavirin Triple Therapy in Japanese Patients Infected with HCV Genotype 1

Factor	Category	Odds Ratio (95% CI)	P
rs8099917 genotype	1: TG+GG	1	<0.001
	2: TT	10.6 (3.07-36.5)	
Substitution of aa 70	1: Gln70 (His70)	1	0.040
	2: Arg70	3.69 (1.06-12.8)	

Only variables that achieved statistical significance ($P < 0.05$) on multivariate logistic regression analysis are shown. 95% CI: 95% confidence interval.

present Japanese study, the sustained virological response rates were 45% and 67% in the T12PR12 and T12PR24 group, respectively, as in the two previous studies. There were differences at three points between the present study and two previous studies: (1) PEG-IFN in two previous studies was used at a fixed dose of PEG-IFN α -2a, but that of the present study was a body weight-adjusted dose of PEG-IFN α -2b; (2) The body mass index of our patients (median; 23 kg/m²) was much lower than that of the participants of the previous study by McHutchison et al.¹⁰ (median; >25 kg/m²); and (3) The present study was performed based on Japanese patients infected with HCV-1b, except for only one patient with HCV-1a. Especially in PROVE-1, the viral breakthrough rate was higher in HCV-1a subjects compared to HCV-1b, and one of the reasons might be due to the low genetic barrier to the emergence of the R155K variant in HCV-1a.^{10,27} Further studies of a larger number of patients matched for background, including genotype, race, body mass index, treatment regimen, and past history of IFN therapy are required to investigate the rate of the sustained virological response by triple therapy.

IL28A, IL28B, and IL29 (IFN- λ -2, IFN- λ -3, and IFN- λ -1, respectively) are novel IFNs identified recently.^{28,29} They are similar to type I IFNs in terms

of biological activities and mechanism of action, in contrast to their differences in structure and genetics.³⁰ The antiviral effects of IFN- λ against hepatitis B virus and HCV have been reported.³¹ Furthermore, α and λ IFNs act synergistically against HCV.³²⁻³⁴ Recent reports showed that genetic variation near the IL28B gene (rs8099917, rs12979860) are pretreatment predictors of virological response to 48-week PEG-IFN plus ribavirin combination therapy in individuals infected with HCV-1,¹⁸⁻²¹ and also affect clinical outcome, including spontaneous clearance of HCV.²² At the 2009 meeting of the American Association for the Study of Liver Diseases, Thompson et al.³⁵ reported that genetic variation near the IL28B gene also affected the viral suppression in the first 2 to 4 weeks of PEG-IFN plus ribavirin, and this phenomenon probably explains much of the difference in treatment response rate. The present study is the first to report that genetic variation near the IL28B gene significantly also affect sustained virological response by triple therapy. These results should be interpreted with caution because races other than Japanese populations were not included. Any generalization of the results should await confirmation by studies of patients of other races to explore the relationship between genetic variation near the IL28B gene and the response to triple therapy.

The present study indicated that the use of the combination of aa substitution of the core region and genetic variation near the IL28B gene had high sensitivity, specificity, PPV, and NPV for prediction of sustained virological response. The efficacy of triple therapy was high in the patients with TT, irrespective of substitution of core aa 70. In the patients having non-TT, those of Arg70 gained high sustained virological response, and sustained virological response was the worst in patients who possessed both non-TT, and Gln70(His70). Along with a high sustained virological response, combined PEG-IFN and ribavirin are accompanied by severe side effects and entail high

Table 4. Sensitivity, Specificity, Positive Predictive Value (PPV), and Negative Predictive Value (NPV) for Sustained Virological Response, According to Substitution of Core aa 70 and Genetic Variation Near IL28B Gene

	% (Number)			
	Sensitivity	Specificity	PPV*	NPV**
(A) rs8099917 genotype TT	79.5 (31/39)	77.8 (21/27)	83.8 (31/37)	72.4 (21/29)
(B) Substitution at aa 70 of arginine (Arg70)	76.9 (30/39)	63.0 (17/27)	75.0 (30/40)	65.4 (17/26)
(A) and (B)	61.5 (24/39)	85.2 (23/27)	85.7 (24/28)	60.5 (23/38)
(A) and/or (B)	94.9 (37/39)	55.6 (15/27)	75.5 (37/49)	88.2 (15/17)

*PPV; Sustained virological response rates for patients with a combination of Arg70 or rs8099917 genotype TT (prediction of sustained virological response).

**NPV; nonsustained virological response rates for patients with a combination of Gln70(His70) or rs8099917 genotype non-TT (prediction of nonsustained virological response).




Fig. 3. Predicting sustained virological response by aa substitution in core region in combination with genetic variation near the IL28B gene. Efficacy of triple therapy was high in the patients with genotype TT who accomplished sustained virological response at 83.8%, irrespective of substitution of core aa 70. In the patients having genotype TG and GG, those of Arg70 gained a high sustained virological response (50.0%), and sustained virological response (11.8%) were the worst in patients who possessed both genotypes TG and GG, and Gln70(His70).

costs. Hence, the patients who do not achieve sustained virological response need to be identified as early as possible, in order to free them of unnecessary side effects and high costs. The present study is the first to report that the combination of aa substitution of the core region and genetic variation near the IL28B gene are very useful as pretreatment predictors of sustained virological response by triple therapy, and further studies based on a larger number of patients are necessary to investigate the present results.

Other limitations of the present study were that aa substitutions in areas other than the core region and NS5A-ISDR of the HCV genome, such as the interferon/ribavirin resistance determining region (IRRDR),³⁶ were not examined. Furthermore, HCV mutants with aa conversions for resistance to telaprevir during triple therapy, such as the 156S mutation,³⁷ were also not investigated. In this regard, telaprevir-resistant HCV mutants were reported to be susceptible to IFN in both *in vivo* and *in vitro* studies.^{38,39} Thus, viral factors before and during triple therapy should be investigated in future studies and identification of these factors should facilitate the development of more effective therapeutic regimens.

In conclusion, triple therapy with telaprevir, PEG-IFN, and ribavirin in Japanese patients infected with HCV-1 and high viral load achieved high sustained virological response rates. Furthermore, the aa substitution pattern of the core region and genetic variation near the IL28B gene seem to affect treatment efficacy. Further large-scale prospective studies are necessary to investigate whether the present results relate to the efficacy of triple therapy and further understanding of the complex interaction between virus- and host-related

factors should facilitate the development of more effective therapeutic regimens.

Acknowledgment: This study was supported in part by a Grant-in-Aid from the Ministry of Health, Labor and Welfare, Japan.

References

- Niederau C, Lange S, Heintges T, Erhardt A, Buschkamp M, Hürter D, et al. Progress of chronic hepatitis C: results of a large, prospective cohort study. *HEPATOLOGY* 1998;28:1687-1695.
- Kenny-Walsh E. Clinical outcomes after hepatitis C infection from contaminated anti-D immune globulin. *Irish Hepatology Research Group. N Engl J Med* 1999;340:1228-1233.
- Tsubota A, Arase Y, Someya T, Suzuki Y, Suzuki F, Saitoh S, et al. Early viral kinetics and treatment outcome in combination of high-dose interferon induction vs. pegylated interferon plus ribavirin for naive patients infected with hepatitis C virus of genotype 1b and high viral load. *J Med Virol* 2005;75:27-34.
- Manns MP, McHutchison JG, Gordon SC, Rustgi VK, Shiffman M, Reindollar R, et al. Peginterferon alfa-2b plus ribavirin compared with interferon alfa-2b plus ribavirin for initial treatment of chronic hepatitis C: a randomized trial. *Lancet* 2001;358:958-965.
- Fried MW, Shiffman ML, Reddy R, Smith C, Marinos G, Gonçales FL, et al. Peginterferon alfa-2a plus ribavirin for chronic hepatitis C virus infection. *N Engl J Med* 2002;347:975-982.
- Lin C, Kwong AD, Perni RB. Discovery and development of VX-950, a novel, covalent, and reversible inhibitor of hepatitis C virus NS3/4A serine protease. *Infect Disord Drug Targets* 2006;6:3-16.
- Modi AA, Hoofnagle JH. New therapies for hepatitis C. *HEPATOLOGY* 2007;46:615-617.
- Zeuzem S. Telaprevir, peginterferon alfa-2a, and ribavirin for 28 days in chronic hepatitis C patients. *J Hepatol* 2008;49:157-159.
- Lawitz E, Rodriguez-Torres M, Muir AJ, Kieffer TL, McNair L, Khunvichai A, et al. Antiviral effects and safety of telaprevir, peginterferon alfa-2a, and ribavirin for 28 days in hepatitis C patients. *J Hepatol* 2008;49:163-169.
- McHutchison JG, Everson GT, Gordon SC, Jacobson IM, Sulkowski M, Kauffman R, et al.; PROVE 1 Study Team. Telaprevir with peginterferon and ribavirin for chronic HCV genotype 1 infection. *N Engl J Med* 2009;360:1827-1838.
- Hézode C, Forestier N, Dusheiko G, Ferenci P, Pol S, Goeser T, et al.; PROVE 2 Study Team. Telaprevir and peginterferon with or without ribavirin for chronic HCV infection. *N Engl J Med* 2009;360:1839-1850.
- Akuta N, Suzuki F, Sezaki H, Suzuki Y, Hosaka T, Someya T, et al. Association of amino acid substitution pattern in core protein of hepatitis C virus genotype 1b high viral load and non-virological response to interferon-ribavirin combination therapy. *Intervirology* 2005;48:372-380.
- Akuta N, Suzuki F, Kawamura Y, Yatsuji H, Sezaki H, Suzuki Y, et al. Predictive factors of early and sustained responses to peginterferon plus ribavirin combination therapy in Japanese patients infected with hepatitis C virus genotype 1b: amino acid substitutions in the core region and low-density lipoprotein cholesterol levels. *J Hepatol* 2007;46:403-410.
- Donlin MJ, Cannon NA, Yao E, Li J, Wahed A, Taylor MW, et al. Pretreatment sequence diversity differences in the full-length hepatitis C virus open reading frame correlate with early response to therapy. *J Virol* 2007;81:8211-8224.
- Akuta N, Suzuki F, Kawamura Y, Yatsuji H, Sezaki H, Suzuki Y, et al. Amino acid substitutions in the hepatitis C virus core region are the important predictor of hepatocarcinogenesis. *HEPATOLOGY* 2007;46:1357-1364.
- Fishman SL, Factor SH, Balestrieri C, Fan X, Dibisceglie AM, Desai SM, et al. Mutations in the hepatitis C virus core gene are associated

- with advanced liver disease and hepatocellular carcinoma. *Clin Cancer Res* 2009;15:3205-3213.
17. Akuta N, Suzuki F, Hirakawa M, Kawamura Y, Yatsuji H, Sezaki H, et al. Amino acid substitutions in the hepatitis C virus core region of genotype 1b affect very early viral dynamics during treatment with telaprevir, peginterferon and ribavirin. *J Med Virol* 2010;82:575-582.
 18. Ge D, Fellay J, Thompson AJ, Simon JS, Shianna KV, Urban TJ, et al. Genetic variation in IL28B predicts hepatitis C treatment-induced viral clearance. *Nature* 2009;461:399-401.
 19. Tanaka Y, Nishida N, Sugiyama M, Kurosaki M, Matsuura K, Sakamoto N, et al. Genome-wide association of IL28B with response to pegylated interferon-alpha and ribavirin therapy for chronic hepatitis C. *Nat Genet* 2009;41:1105-1109.
 20. Suppiah V, Moldovan M, Ahlensiel G, Berg T, Weltman M, Abate ML, et al. IL28B is associated with response to chronic hepatitis C interferon-alpha and ribavirin therapy. *Nat Genet* 2009;41:1100-1104.
 21. Rauch A, Kuralik Z, Descombes P, Cai T, di Iulio J, Mueller T, et al.: Swiss Hepatitis C and HIV Cohort Studies. Genetic variation in IL28B is associated with chronic hepatitis C and treatment failure — a genome-wide association study. *Gastroenterology* 2010;138:1338-1345.
 22. Thomas DL, Thio CL, Martin MP, Qi Y, Ge D, O'Huigin C, et al. Genetic variation in IL28B and spontaneous clearance of hepatitis C virus. *Nature* 2009;461:798-801.
 23. Kato N, Hijikata M, Ootsuyama Y, Nakagawa M, Ohkoshi S, Sugimura T, et al. Molecular cloning of the human hepatitis C virus genome from Japanese patients with non-A, non-B hepatitis. *Proc Natl Acad Sci U S A* 1990;87:9524-9528.
 24. Enomoto N, Sakuma I, Asahina Y, Kurosaki M, Murakami T, Yamamoto C, et al. Mutations in the nonstructural protein 5A gene and response to interferon in patients with chronic hepatitis C virus 1b infection. *N Engl J Med* 1996;334:77-81.
 25. Ohnishi Y, Tanaka T, Ozaki K, Yamada R, Suzuki H, Nakamura Y. A high-throughput SNP typing system for genome-wide association studies. *J Hum Genet* 2001;46:471-477.
 26. Suzuki A, Yamada R, Chang X, Tokuhira S, Sawada T, Suzuki M, et al. Functional haplotypes of PADI4, encoding citrullinating enzyme peptidylarginine deiminase 4, are associated with rheumatoid arthritis. *Nat Genet* 2003;34:395-402.
 27. Kieffer TL, Sarrazin C, Miller JS, Welker MW, Forestier N, Reesink HW, et al. Telaprevir and pegylated interferon-alpha-2a inhibit wild-type and resistant genotype 1 hepatitis C virus replication in patients. *HEPATOLOGY* 2007;46:631-639.
 28. Sheppard P, Kindsvoegel W, Xu W, Henderson K, Schlutsmeyer S, Whimmore TE, et al. IL-28, IL-29 and their class II cytokine receptor IL-28R. *Nat Immunol* 2003;4:63-68.
 29. Kotenko SV, Gallagher G, Baurin VV, Lewis-Antes A, Shen M, Shah NK, et al. IFN-lambdas mediate antiviral protection through a distinct class II cytokine receptor complex. *Nat Immunol* 2003;4:69-77.
 30. Maher SG, Sheikh F, Scarzello AJ, Romero-Weaver AL, Baker DP, Donnelly RP, et al. IFNalpha and IFNlambda differ in their antiproliferative effects and duration of JAK/STAT signaling activity. *Cancer Biol Ther* 2008;7:1109-1115.
 31. Robek MD, Boyd BS, Chisari FV. Lambda interferon inhibits hepatitis B and C virus replication. *J Virol* 2005;79:3851-3854.
 32. Zhu H, Butera M, Nelson DR, Liu C. Novel type I interferon IL-28A suppresses hepatitis C viral RNA replication. *Virol J* 2005;2:80.
 33. Marcello T, Grakoui A, Barba-Spaeth G, Machlin ES, Kotenko SV, MacDonald MR, et al. Interferons alpha and lambda inhibit hepatitis C virus replication with distinct signal transduction and gene regulation kinetics. *Gastroenterology* 2006;131:1887-1898.
 34. Pagliaccetti NE, Eduardo R, Kleinstein SH, Mu XJ, Bandi P, Robek MD. Interleukin-29 functions cooperatively with interferon to induce antiviral gene expression and inhibit hepatitis C virus replication. *J Biol Chem* 2008;283:30079-30089.
 35. Thompson AJ, Muir A, Sulkowski MS, Afdhal NH, Jacobson IM, Esteban R, et al. Genome wide analysis of patients from the IDEAL study identifies a polymorphism upstream of the IL28B (=IFNL3) gene that is strongly associated with SVR in patients with HCV-1 [Abstract]. *HEPATOLOGY* 2009;50:91A.
 36. El-Shamy A, Nagano-Fujii M, Sasase N, Imoto S, Kim SR, Hotta H. Sequence variation in hepatitis C virus nonstructural protein 5A predicts clinical outcome of pegylated interferon/ribavirin combination therapy. *HEPATOLOGY* 2008;48:38-47.
 37. Lin C, Gates CA, Rao BG, Brennan DL, Fulghum JR, Luong YP, et al. In vitro studies of cross-resistance mutations against two hepatitis C virus serine protease inhibitors, VX-950 and BILN 2061. *J Biol Chem* 2005;280:36784-36791.
 38. Forestier N, Reesink HW, Weegink CJ, McNair L, Kieffer TL, Chu HM, et al. Antiviral activity of telaprevir (VX-950) and peginterferon alpha-2a in patients with hepatitis C. *HEPATOLOGY* 2007;46:640-648.
 39. Zhou Y, Mith U, Hanzelka BJ, Bartels DJ, Wei Y, Rao BG, et al. Phenotypic and structural analyses of hepatitis C virus NS3 protease Arg155 variants: sensitivity to telaprevir (VX-950) and interferon alpha. *J Biol Chem* 2007;282:22619-22628.

Amino Acid Substitution in HCV Core Region and Genetic Variation near the *IL28B* Gene Affect Viral Dynamics during Telaprevir, Peginterferon and Ribavirin Treatment

Norio Akuta^a Fumitaka Suzuki^a Miharuru Hirakawa^a Yusuke Kawamura^a
Hiromi Yatsuji^a Hitomi Sezaki^a Yoshiyuki Suzuki^a Tetsuya Hosaka^a
Masahiro Kobayashi^a Mariko Kobayashi^b Satoshi Saitoh^a Yasuji Arase^a
Kenji Ikeda^a Kazuaki Chayama^d Yusuke Nakamura^c Hiromitsu Kumada^a

^aDepartment of Hepatology, and ^bLiver Research Laboratory, Toranomon Hospital, ^cLaboratory of Molecular Medicine, Human Genome Center, Institute of Medical Science, University of Tokyo, Tokyo, and ^dDepartment of Medical and Molecular Science, Division of Frontier Medical Science, Programs for Biomedical Research, Graduate School of Biomedical Science, Hiroshima University, Hiroshima, Japan

Key Words

Hepatitis C virus · Core region · *IL28B* · Telaprevir · Peginterferon · Ribavirin · Viral dynamics

Abstract

Objectives: Genetic variation near the *IL28B* gene and substitution of aa 70 and 91 in the core region of HCV-1b are useful as predictors of treatment efficacy to telaprevir/pegylated interferon (PEG-IFN)/ribavirin, but its impact on viral dynamics is not clear. **Methods:** This study investigated predictive factors of viral dynamics during 12- or 24-week regimen of triple therapy in 80 Japanese adults infected with HCV-1b. **Results:** After 24 h of commencement of treatment, the proportion of patients with Arg70 and Leu91 substitutions in the core region who showed ≥ 3.0 log drop in HCV RNA level was significantly higher than that of patients with Gln70 (His70) and/or Met91. At 8 and 12 weeks, HCV RNA loss rate of patients with rs8099917 genotype TT near *IL28B* gene was significantly higher than that of patients with non-TT.

Multivariate analysis identified substitution of aa 70 and 91 as a predictor of ≥ 3.0 log fall in HCV RNA level at 24 h (Arg70 and Leu91) and SVR (Arg70), and rs8099917 (TT) as a predictor of HCV RNA loss at 12 weeks and SVR. **Conclusions:** This study identified genetic variation near *IL28B* gene and a substitution of the core region as predictors of viral dynamics during triple therapy.

Copyright © 2011 S. Karger AG, Basel

Introduction

Hepatitis C virus (HCV) usually causes chronic infection that can result in chronic hepatitis, liver cirrhosis, and hepatocellular carcinoma (HCC) [1, 2]. At present, treatments based on interferon (IFN), in combination with ribavirin, are mainstay for combating HCV infection. In Japan, HCV genotype 1b (HCV-1b) in high viral loads (>100 kIU/ml) accounts for more than 70% of HCV infections, making it difficult to treat patients with chronic hepatitis

KARGER

Fax +41 61 306 12 34
E-Mail karger@karger.ch
www.karger.com

© 2011 S. Karger AG, Basel
0300-5526/11/0000-0000\$38.00/0

Accessible online at:
www.karger.com/int

Norio Akuta, MD
Department of Hepatology, Toranomon Hospital
2-2-2 Toranomon, Minato-ku
Tokyo 105-0001 (Japan)
Tel. +81 44 877 5111, E-Mail akuta-gi@umin.ac.jp

C [3]. Such a background calls for efficient treatments of Japanese patients with chronic HCV infection.

Even with pegylated IFN (PEG-IFN) combined with ribavirin, a sustained virological response lasting over 24 weeks after the withdrawal of treatment is achieved in at most 50% of the patients infected with HCV-1b and high viral loads [4, 5]. Recently, a new strategy was introduced in the treatment of chronic HCV infection by means of inhibiting protease in the NS3/NS4 of the HCV polyprotein. Of these, telaprevir (VX-950) was selected as a candidate agent for treatment of chronic HCV infection [6]. Later, it was found that telaprevir, when combined with PEG-IFN and ribavirin, gains a robust antiviral activity [7, 8]. Two previous studies (PROVE1 and PROVE2) showed that the 12- and 24-week regimen of telaprevir/PEG-IFN/ribavirin could achieve sustained virological response rates of 35–60 and 61–69% in patients infected with HCV-1, respectively [9, 10]. Furthermore, a recent study (PROVE3) also showed that the 24- and 48-week regimen of triple therapy could achieve sustained virological response rates of 51 and 53% in HCV-1 infected patients in whom initial PEG-IFN/ribavirin treatment failed, respectively [11].

Amino acid (aa) substitutions at positions 70 and/or 91 in the HCV core region of patients infected with HCV-1b and high viral loads are pretreatment predictors of poor virological response to PEG-IFN plus ribavirin combination therapy [12–14], and also affect clinical outcome, including hepatocarcinogenesis [15, 16]. Furthermore, genetic variations near the *IL28B* gene (rs8099917, rs12979860) on chromosome 19 as host-related factor, which encodes IFN- λ -3, are pretreatment predictors of virological response to 48-week PEG-IFN plus ribavirin combination therapy in individuals infected with HCV-1 [17–20], and also affect clinical outcome, including spontaneous clearance of HCV [21]. A recent report identified genetic variation near *IL28B* gene and aa substitution of the core region as predictors of sustained virological response to triple therapy of telaprevir/PEG-IFN/ribavirin in Japanese patients infected with HCV-1b [22]. However, it is not clear at this stage whether genetic variation near the *IL28B* gene and aa substitution of the core region can be used before therapy to predict viral dynamics during triple therapy.

The present study included 80 patients with HCV-1b and high viral loads, who received the triple therapy of telaprevir with PEG-IFN plus ribavirin. The aims of the study were to identify the pretreatment factors that could predict viral dynamics during treatment, including viral- (aa substitutions in the HCV core and NS5A regions) and host-related factors (genetic variation near *IL28B* gene).

Patients and Methods

Study Population

Between May 2008 and September 2009, 81 patients infected with HCV were recruited to this study at the Department of Hepatology in Toranomon Hospital in metropolitan Tokyo. The study protocol was in compliance with the Good Clinical Practice Guidelines and the 1975 Declaration of Helsinki, and was approved by the institutional review board. Each patient gave an informed consent before participating in this trial. Patients were divided into two groups: 20 (25%) patients were allocated to a 12-week regimen of triple therapy [telaprevir (MP-424), PEG-IFN and ribavirin] (the T12PR12 group), and 61 patients (75%) were assigned to a 24-week regimen of the same triple therapy for 12 weeks followed by dual therapy of PEG-IFN and ribavirin for 12 weeks (the T12PR24 group).

Eighty of the 81 patients met the following inclusion and exclusion criteria: (1) Diagnosis of chronic hepatitis C. (2) HCV-1b confirmed by sequence analysis. (3) HCV RNA levels of ≥ 5.0 log IU/ml determined by the COBAS TaqMan HCV test (Roche Diagnostics, Tokyo, Japan). (4) Japanese (Mongoloid) ethnicity. (5) Age at study entry of 20–65 years. (6) Body weight ≥ 35 kg and ≤ 120 kg at the time of registration. (7) Lack of decompensated liver cirrhosis. (8) Negativity for hepatitis B surface antigen (HBsAg) in serum. (9) Negative history of HCC. (10) No previous treatment for malignancy. (11) Negative history of autoimmune hepatitis, alcohol liver disease, hemochromatosis, and chronic liver disease other than chronic hepatitis C. (12) Negative history of depression, schizophrenia or suicide attempts, hemoglobinopathies, angina pectoris, cardiac insufficiency, myocardial infarction or severe arrhythmia, uncontrollable hypertension, chronic renal dysfunction or creatinine clearance of ≤ 50 ml/min at baseline, diabetes requiring treatment or fasting glucose level of ≥ 110 mg/dl, autoimmune disease, cerebrovascular disorders, thyroidal dysfunction uncontrollable by medical treatment, chronic pulmonary disease, allergy to medication or anaphylaxis at baseline. (13) Hemoglobin level of ≥ 12 g/dl, neutrophil count $\geq 1,500/\text{mm}^3$, and platelet count of $\geq 100,000/\text{mm}^3$ at baseline. Pregnant or breast-feeding women or those willing to become pregnant during the study and men with a pregnant partner were excluded from the study. In this study, all of the 80 patients were evaluated for the pretreatment predictors for viral dynamics during triple therapy, and 77 of the 80 patients were followed up for at least 24 weeks after the completion of treatment. The treatment efficacy was evaluated by 24 weeks after the completion of therapy (sustained virological response), based on the COBAS TaqMan HCV test (Roche Diagnostics).

Telaprevir (MP-424; Mitsubishi Tanabe Pharma, Osaka, Japan) was administered at 750 or 500 mg three times a day at an 8-hour (q8) interval after the meal. PEG-IFN α -2b (PEG-Intron; Schering Plough, Kenilworth, N.J., USA) was injected subcutaneously at a median dose of 1.5 $\mu\text{g}/\text{kg}$ (range 1.3–2.0 $\mu\text{g}/\text{kg}$) once a week. Ribavirin (Rebetol; Schering Plough) was administered at 200–600 mg twice a day after breakfast and dinner (daily dose 600–1,000 mg).

PEG-IFN and ribavirin were discontinued or their doses reduced, as required, upon reduction of hemoglobin level, leukocyte count, neutrophil count or platelet count, or the development of adverse events. Thus, the dose of PEG-IFN was reduced by 50% when the leukocyte count decreased below $1,500/\text{mm}^3$, neutro-

Table 1. Profile and laboratory data at commencement of telaprevir, peginterferon and ribavirin triple therapy in Japanese patients infected with HCV-1b

<i>Demographic data</i>	
Number of patients	80
Sex, M/F	43/37
Age, years*	55 (23–65)
History of blood transfusion	24 (20.0%)
Family history of liver disease	13 (16.3%)
Body mass index*	22.5 (13.2–32.4)
<i>Laboratory data*</i>	
Level of viremia, log IU/ml	6.8 (5.1–7.6)
Serum aspartate aminotransferase, IU/l	34 (15–118)
Serum alanine aminotransferase, IU/l	42 (12–175)
Serum albumin, g/dl	3.9 (3.3–4.6)
Gamma-glutamyl transpeptidase, IU/l	36 (9–229)
Leukocyte count, per mm ³	4,800 (2,800–8,100)
Hemoglobin, g/dl	14.3 (11.7–16.8)
Platelet count, × 10 ⁴ /mm ³	17.3 (9.5–33.8)
α-Fetoprotein, μg/l	4 (2–39)
Total cholesterol, mg/dl	180 (112–276)
Fasting plasma glucose, mg/dl	92 (64–125)
<i>Treatment</i>	
PEG-IFNα-2b dose, μg/kg*	1.5 (1.3–2.0)
Ribavirin dose, mg/kg*	11.5 (7.2–18.4)
Telaprevir dose, 1,500/2,250 mg/day	10/70
Treatment regimen (T12PR12 group/T12PR24 group)	20/60
<i>Amino acid substitutions in the HCV-1b</i>	
Core aa 70, arginine/glutamine (histidine)	47/33
Core aa 91, leucine/methionine	43/37
ISDR of NS5A, wild-type/non-wild-type	76/4
<i>Genetic variation near IL28B gene</i>	
rs8099917 genotype, TT/TG/GG/ND	46/30/2/2
rs12979860 genotype, CC/CT/TT/ND	43/31/2/4
<i>Past history of IFN therapy</i>	
Treatment naive	27
Relapsers to previous treatment	33
Nonresponders to previous treatment	20
Data are numbers and percentages of patients, except those denoted by *, which represent the median (range) values. ND = Not determined.	

phil count below 750/mm³ or platelet count below 80,000/mm³; PEG-IFN was discontinued when these counts decreased below 1,000/mm³, 500/mm³ or 50,000/mm³, respectively. When hemoglobin decreased to <10 g/dl, the daily dose of ribavirin was reduced from 600 to 400, 800 to 600 and 1,000 to 600 mg, depending on the initial dose. Ribavirin was withdrawn when hemoglobin decreased to <8.5 g/dl. However, the dose of telaprevir (MP-424) remained the same, and its administration was stopped when the

discontinuation was appropriate for the development of adverse events. In those patients who discontinued telaprevir, treatment with PEG-IFNα-2b and ribavirin was also terminated.

Table 1 summarizes the profiles and laboratory data of the 80 patients at the commencement of treatment. They included 43 males and 37 females, aged 23–65 years (median 55 years).

Measurement of HCV RNA

The antiviral effects of the triple therapy on HCV were assessed by measuring plasma HCV RNA levels. In this study, HCV RNA levels during treatment were evaluated at least once every month before, during, and after therapy. Furthermore, to investigate the pretreatment predictors for viral dynamics, HCV RNA levels during treatment were evaluated at 7 time points; 24 h, 1, 2, 4, 6, 8 and 12 weeks after the commencement of treatment. HCV RNA levels during treatment were evaluated in 80 (100%), 80 (100%), 80 (100%), 79 (98.8%), 75 (93.8%), 74 (92.5%), and 69 (86.3%) of the 80 patients, at the above time intervals, respectively. HCV RNA concentrations were determined using the COBAS TaqMan HCV test (Roche Diagnostics). The linear dynamic range of the assay was 1.2–7.8 log IU/ml, and the undetectable samples were defined as loss of HCV RNA. Especially, falls in HCV RNA levels at 24 h relative to baseline were investigated as very early dynamics.

Detection of Amino Acid Substitutions in Core and NS5A Regions of HCV-1b

With the use of HCV-J (accession No. D90208) as a reference [23], the sequence of 1–191 aa in the core protein of HCV-1b was determined and then compared with the consensus sequence constructed on 80 clinical samples to detect substitutions at aa 70 of arginine (Arg70) or glutamine/histidine (Gln70/His70) and aa 91 of leucine (Leu91) or methionine (Met91) [12]. The sequence of 2209–2248 aa in the NS5A of HCV-1b (IFN sensitivity-determining region; ISDR) reported by Enomoto et al. [24] was determined, and the numbers of aa substitutions in ISDR were defined as wild-type (0, 1) or non-wild-type (≥2). In the present study, aa substitutions of the core region and NS5A-ISDR of HCV-1b were analyzed by direct sequencing [22].

Genetic Variation near IL28B Gene

Samples for genomewide association survey were genotyped using the Illumina HumanHap610-Quad Genotyping BeadChip. Genotyping data were subjected to quality control before the data analysis. Genotyping for replication and fine mapping was performed by use of the Invader assay, TaqMan assay, or direct sequencing as described previously [25, 26].

In this study, genetic variations near *IL28B* gene (rs8099917, rs12979860), reported as the pretreatment predictors of treatment efficacy and clinical outcome [17–22], were investigated.

Statistical Analysis

Nonparametric tests (χ^2 test and Fisher's exact probability test) were used to compare the characteristics of the groups. Univariate and multivariate logistic regression analyses were used to determine those factors that significantly contributed to viral dynamics and sustained virological response. The ORs and 95%CI were also calculated. All p values less than 0.05 by the two-tailed test were considered significant. Variables that achieved statistical significance ($p < 0.05$) on univariate analysis were entered into




Fig. 1. **a** Very early dynamics according to amino acid substitutions in core region. After 24 h of commencement of the triple therapy, patients with Arg70 and Leu91 (median 2.9 log IU/ml; range 1.8–4.0 log IU/ml) significantly showed the steeper decline of HCV RNA level than those with Gln70 (His70) and/or Met91 (median 2.7 log IU/ml; range 1.6–3.5 log IU/ml). **b** Very early dynamics according to genetic variation near the *IL28B* gene. After 24 h of commencement of the triple therapy, the decline of HCV RNA level of patients with rs8099917 genotype TT (median 2.8 log IU/ml; range 1.6–4.0 log IU/ml) was not significantly different from that of patients with genotype TG and GG (median 2.7 log IU/ml; range 2.2–3.5 log IU/ml).

multiple logistic regression analysis to identify significant independent predictive factors. Each variable was transformed into categorical data consisting of two simple ordinal numbers for univariate and multivariate analyses. The potential pretreatment factors associated with treatment efficacy included the following variables: sex, age, history of blood transfusion, familial history of liver disease, body mass index, aspartate aminotransferase (AST), alanine aminotransferase (ALT), albumin, gamma-glutamyl transpeptidase (γ GTP), leukocyte count, hemoglobin, platelet count, HCV RNA level, α -fetoprotein, total cholesterol, fasting blood sugar, PEG-IFN dose/body weight, ribavirin dose/body

weight, telaprevir dose/day, treatment regimen of triple therapy, past history of IFN therapy, genetic variation near the *IL28B* gene, and amino acid substitution in the core region, and NS5A-ISDR. Statistical analyses were performed using the SPSS software (SPSS Inc., Chicago, Ill. USA).

Results

Virological Response to Therapy and Loss of HCV RNA during Treatment

Sustained virological response was achieved by 63.6% (49 of 77 patients). The disappearance rate of HCV RNA during treatment was 0% (0 of 80), 1.3% (1 of 80), 33.8% (27 of 80), 81.0% (64 of 79), 90.7% (68 of 75), 94.6% (70 of 74), and 89.9% (62 of 69) at 24 hours, 1, 2, 4, 6, 8, and 12 weeks, respectively.

*Very Early Dynamics according to Amino Acid Substitutions in Core Region and Genetic Variation near the *IL28B* Gene*

After 24 h of commencement of the triple therapy, the proportion of patients with Arg70 and Leu91 substitutions who showed ≥ 3.0 log drop in HCV RNA level (45.2%; 14 of 31 patients) was significantly higher than that of patients with Gln70 (His70) and/or Met91 (14.3%; 7 of 49) ($p = 0.004$). Thus, patients with Arg70 and Leu91 (median 2.9 log IU/ml; range 1.8–4.0 log IU/ml) significantly showed the steeper decline of HCV RNA level than those with Gln70 (His70) and/or Met91 (median 2.7 log IU/ml; range 1.6–3.5 log IU/ml) (fig. 1a).

After 24 h of commencement of treatment, the proportion of patients with rs8099917 genotype TT who showed ≥ 3.0 log drop in HCV RNA level (30.4%; 14 of 46 patients) was not significantly different from that of patients with genotype TG and GG (21.9%; 7 of 32). Thus, the decline of HCV RNA level of patients with genotype TT (median 2.8 log IU/ml; range 1.6–4.0 log IU/ml) was not significantly different from that of patients with genotype TG and GG (median 2.7 log IU/ml; range 2.2–3.5 log IU/ml) (fig. 1b).

Hence, the fall in HCV RNA level at 24 h was influenced by aa substitution patterns in the core region, but was independent of genetic variation near *IL28B* gene.

*Rates of Loss of HCV RNA according to Amino Acid Substitutions in Core Region and Genetic Variation near the *IL28B* Gene*

According to the substitution of core aa 70 and 91, the rate of HCV RNA loss of patients with Arg70 and Leu91 was not significantly different from that of patients with

Gln70 (His70) and/or Met91 at each time point (1, 2, 4, 6, 8 and 12 weeks).

According to genetic variation near the *IL28B* gene, the rate of HCV RNA loss at 1, 2, 4 and 6 weeks was not significantly different between rs8099917 genotype TT and non-TT (TG and GG). However, at 8 and 12 weeks, the rate of HCV RNA loss of patients with genotype TT was significantly higher than that of patients with genotype non-TT (fig. 2).

Predictive Factors Associated with ≥ 3.0 log Fall in HCV RNA Level at 24 Hours

Univariate analysis identified two parameters that correlated with ≥ 3.0 log fall in HCV RNA level at 24 h significantly: substitution of aa 70 and 91 (Arg70 and Leu91; OR 4.94, $p = 0.003$) and body mass index (≥ 25.0 ; OR 3.92, $p = 0.022$). Two factors were identified by multivariate analysis as independent parameters that either significantly ($p < 0.05$) or marginally ($p < 0.10$) influenced ≥ 3.0 log fall in HCV RNA level at 24 h [Arg70 and Leu91 (OR 3.99, $p = 0.015$) and body mass index ≥ 25.0 (OR 3.24, $p = 0.061$)] (table 2).

Predictive Factors Associated with Loss of HCV RNA at 2, 4 and 12 Weeks

Univariate analysis identified two parameters that correlated with loss of HCV RNA at 2 weeks significantly: platelet count ($\geq 15.0 \times 10^4/\text{mm}^3$; OR 6.99, $p = 0.014$) and level of viremia (< 7.0 log IU/ml; OR 3.13, $p = 0.045$). One factor was identified by multivariate analysis as independent parameter that either significantly or marginally influenced loss of HCV RNA at 2 weeks (platelet count $\geq 15.0 \times 10^4/\text{mm}^3$; OR 6.99, $p = 0.014$) (table 2).

Univariate analysis identified two parameters that correlated with loss of HCV RNA at 4 weeks significantly: history of blood transfusion (absence; OR 5.71, $p = 0.006$) and body mass index (≥ 20.0 ; OR 4.29, $p = 0.019$). Two factors were identified by multivariate analysis as independent parameters that either significantly or marginally influenced loss of HCV RNA at 4 weeks (history of blood transfusion: absence; OR 4.29, $p = 0.026$, and body mass index ≥ 20.0 ; OR 3.47, $p = 0.069$) (table 2).

Univariate analysis identified two parameters that correlated with loss of HCV RNA at 12 weeks significantly: sex (male; OR 9.52, $p = 0.043$) and genetic variation in rs8099917 (genotype TT; OR 9.00, $p = 0.048$). Two factors were identified by multivariate analysis as independent parameters that either significantly or marginally influenced loss of HCV RNA at 12 weeks (male sex; OR 11.0, $p = 0.036$, and rs8099917 genotype TT; OR 10.3, $p = 0.042$) (table 2).




Fig. 2. Rates of loss of HCV RNA according to genetic variation near the *IL28B* gene. According to genetic variation near the *IL28B* gene, the rate of HCV RNA loss at 1, 2, 4 and 6 weeks was not significantly different between rs8099917 genotype TT and non-TT (TG and GG). However, at 8 and 12 weeks, the rate of HCV RNA loss of patients with genotype TT was significantly higher than that of patients with genotype non-TT.

Predictive Factors Associated with Sustained Virological Response

Univariate analysis identified three parameters that correlated with sustained virological response significantly: substitution of aa 70 (Arg70; OR 3.51, $p = 0.011$), and genetic variation in rs8099917 (genotype TT; OR 11.1, $p < 0.001$) and rs12979860 (genotype CC; OR 10.2, $p < 0.001$). Two factors were identified by multivariate analysis as independent parameters that either significantly or marginally influenced sustained virological response (rs8099917 genotype TT; OR 9.94, $p < 0.001$, and Arg70; OR 3.15, $p = 0.055$) (table 2).

Comparison of Factors Associated with Each Treatment Efficacy Identified by Multivariate Analysis

Table 3 shows independent parameters that either significantly or marginally influenced multivariate logistic regression for each evaluation of treatment efficacy. Multivariate analysis identified substitution of aa 70 and 91 as a predictor of ≥ 3.0 log fall in HCV RNA level at 24 h (Arg70 and Leu91) and sustained virological response (Arg70), and rs8099917 (TT) as a predictor of HCV RNA loss at 12 weeks and sustained virological response. Thus, genetic variation near *IL28B* gene and aa substitution of the core region affect viral dynamics of different phases during triple therapy.

Table 2. Factors associated with treatment efficacy of telaprevir, peginterferon and ribavirin triple therapy in Japanese patients infected with HCV-1b, identified by univariate and multivariate analysis

Factor	Category	Univariate logistic regression		Multivariate logistic regression		
		OR (95% CI)	p	OR (95% CI)	p	
A	≥3.0 log fall in HCV RNA at 24 h					
	Substitution of aa 70 and 91	1: Gln70 (His70) and/or Met91 2: Arg70 and Leu91	1 4.94 (1.70–14.4)	0.003	1 3.99 (1.31–12.2)	0.015
	Body mass index	1: <25.0 2: ≥25.0	1 3.92 (1.22–12.6)	0.022	1 3.24 (0.95–11.1)	0.061
	B	HCV RNA loss at 2 weeks				
	Platelet count, ×10 ⁴ /mm ³	1: <15.0 2: ≥15.0	1 6.99 (1.49–32.8)	0.014	1 6.99 (1.49–32.8)	0.014
	Level of viremia, log IU/ml	1: ≥7.0 2: <7.0	1 3.13 (1.02–9.52)	0.045	– –	– –
C	HCV RNA loss at 4 weeks					
	History of blood transfusion	1: presence 2: absence	1 5.71 (1.66–19.6)	0.006	1 4.29 (1.86–15.6)	0.026
	Body mass index	1: <20.0 2: ≥20.0	1 4.29 (1.26–14.5)	0.019	1 3.47 (0.91–13.3)	0.069
	D	HCV RNA loss at 12 weeks				
Sex		1: female 2: male	1 9.52 (1.08–83.3)	0.043	1 11.0 (1.16–100)	0.036
	rs8099917 genotype	1: TG+GG 2: TT	1 9.00 (1.02–79.5)	0.048	1 10.3 (1.08–98.0)	0.042
	E	Sustained virological response				
rs8099917 genotype		1: TG+GG 2: TT	1 11.1 (3.68–33.5)	<0.001	1 9.94 (3.05–32.4)	<0.001
	Substitution of aa 70	1: Gln70 (His70) 2: Arg70	1 3.51 (1.33–9.26)	0.011	1 3.15 (0.97–10.2)	0.055
	rs12979860 genotype	1: CT+TT 2: CC	1 10.2 (3.33–3.13)	<0.001	– –	– –

Variables that achieved statistical significance ($p < 0.05$) on univariate analysis were entered into multiple logistic regression analysis to identify significant independent predictive factors.

The other significant predictors of HCV RNA loss were platelet count ($\geq 15.0 \times 10^4/\text{mm}^3$) at 2 weeks, history of blood transfusion (absence) at 4 weeks, and sex (male) at 12 weeks.

Discussion

Thompson et al. [27] reported that genetic variation near *IL28B* gene was also associated with increased on-treatment and sustained virological response and effectively predicted treatment outcome in treatment-naïve HCV-1 patients treated with PEG-IFN plus ribavirin. However, HCV RNA loss at 4 weeks (rapid virological

response) was a strong predictor of sustained virological response regardless of genetic variation near the *IL28B* gene. This phenomenon probably explains why it might be important to identify the pretreatment factors that could predict viral dynamics during treatment. The present study is the first to identify the pretreatment factors that could predict viral dynamics during triple therapy in patients infected with HCV-1. These results should be interpreted with caution since races other than Japanese and the patients infected with HCV-1a were not included. Any generalization of the results should await confirmation by studies including patients of other races and with HCV-1a to explore whether genetic variation near *IL28B* gene and aa substitution

Table 3. Comparison of factors associated with treatment efficacy of telaprevir, peginterferon and ribavirin triple therapy in Japanese patients infected with HCV-1b identified by multivariate analysis

Factor	≥3.0 log fall in HCV RNA (at 24 h)	HCV RNA loss (at 2 weeks)	HCV RNA loss (at 4 weeks)	HCV RNA loss (at 12 weeks)	Sustained virological response
Core aa 70 and 91	Arg70 and Leu91 p = 0.015 3.99 (1.31–12.2)*				Arg70 p = 0.055 3.15 (0.97–10.2)*
<i>IL28B</i> rs8099917				genotype TT p = 0.042 10.3 (1.08–98.0)*	genotype TT p < 0.001 9.94 (3.05–32.4)*
Others	body mass index p = 0.061 3.24 (0.95–11.1)*	platelet count p = 0.014 6.99 (1.49–32.8)*	body mass index p = 0.069 3.47 (0.91–13.3)* history of blood transfusion p = 0.026 4.29 (1.86–15.6)*	sex p = 0.036 11.0 (1.16–100)*	

Only variables that achieved statistical significance ($p < 0.05$) or marginal significance ($p < 0.10$) on multivariate logistic regression are shown. * OR (95% CI).

of core region also affect viral dynamics during triple therapy.

Two studies showed that aa substitution of the core region and genetic variation near *IL28B* gene affected viral dynamics during treatment, and sustained virological response to 48-week PEG-IFN plus ribavirin therapy in patients infected with HCV-1 [27, 28]. Furthermore, a recent report also showed that aa substitutions of core region might be used to predict very early dynamics (within 48 h) after the start of triple therapy of telaprevir with PEG-IFN and ribavirin [29]. In the present study, multivariate analysis identified substitution of aa 70 and 91 as a predictor of ≥ 3.0 log fall in HCV RNA level at 24 hours (i.e. viral dynamics of very early phase) and sustained virological response, and rs8099917 as a predictor of HCV RNA loss at 12 weeks (i.e. viral dynamics of later phase) and sustained virological response. This study is the first to report that genetic variation near *IL28B* gene and aa substitution of the core region affect viral dynamics of different phases during triple therapy, and probably explains why the combination of these independent factors is very useful as pretreatment predictors of sustained virological response by triple therapy [22]. The underlying mechanisms of the different viral dynamics to treatment are still unclear, and further studies based on a larger number of patients are necessary to investigate the present results.

Previous data indicated that absence of advanced liver fibrosis and male gender were positive predictors of virological response to 48-week PEG-IFN plus ribavirin therapy [13, 28]. The present study also showed that higher levels of platelet count at 2 weeks, as a surrogate marker of milder liver fibrosis, and male gender at 12 weeks were significant positive predictors of HCV RNA loss during triple therapy. The other positive predictors were absence of history of blood transfusion at 4 weeks and higher levels of body mass index at 24 h and 4 weeks, but the underlying mechanisms are still unclear. Thus, this report identified the pretreatment factors that could predict viral dynamics during triple therapy, but this study, based on a small number of patients, might provide misleading results (e.g. possible type error). Further studies of a larger number of patients are required to explore predictors, including viral- and host-related factors.

The limitations of the present study were that aa substitutions in areas other than the core region and NS5A-ISDR of the HCV genome, such as the interferon/ribavirin resistance determining region (IRRD) [30], were not examined. Furthermore, HCV mutants with aa conversions for resistance to telaprevir during triple therapy, such as the 156S mutation [31], were also not investigated. In this regard, telaprevir-resistant HCV mutants were reported to be susceptible to IFN in both in vivo and in vitro studies [32, 33]. Thus, viral factors before and during triple therapy should be investigated in

future studies, and identification of these factors should facilitate the development of more effective therapeutic regimens.

In conclusion, this study identified genetic variation near *IL28B* gene and aa substitution of the core region as predictors of viral dynamics during triple therapy of telaprevir/PEG-IFN/ribavirin in Japanese patients infected with HCV-1b. Further large-scale prospective studies are necessary to investigate whether the present results relate to the efficacy of the triple therapy, and further under-

standing of the complex interaction between virus- and host-related factors should facilitate the development of more effective therapeutic regimens.

Acknowledgment

This study was supported in part by a Grant-in-Aid from the Ministry of Health, Labor and Welfare, Japan.

References

- Niederer C, Lange S, Heintges T, Erhardt A, Buschkamp M, Hürter D, Nawrocki M, Kruska L, Hensel F, Petry W, Häussinger D: Progress of chronic hepatitis C: results of a large, prospective cohort study. *Hepatology* 1998;28:1687-1695.
- Kenny-Walsh E: Clinical outcomes after hepatitis C infection from contaminated anti-D immune globulin: Irish Hepatology Research Group. *N Engl J Med* 1999;340:1228-1233.
- Tsubota A, Arase Y, Someya T, Suzuki Y, Suzuki F, Saitoh S, Ikeda K, Akuta N, Hosaka T, Kobayashi M, Kumada H: Early viral kinetics and treatment outcome in combination of high-dose interferon induction vs. pegylated interferon plus ribavirin for naive patients infected with hepatitis C virus of genotype 1b and high viral load. *J Med Virol* 2005;75:27-34.
- Manns MP, McHutchison JG, Gordon SC, Rustgi VK, Shiffman M, Reindollar R, Goodman ZD, Koury K, Ling MH, Albrecht JK: Peginterferon alfa-2b plus ribavirin compared with interferon alfa-2b plus ribavirin for initial treatment of chronic hepatitis C: a randomized trial. *Lancet* 2001;358:958-965.
- Fried MW, Shiffman ML, Reddy R, Smith C, Marinos G, Gonçales FL, Häussinger D, Diago M, Carosi G, Dhumeaux D, Craxi A, Lin A, Hoffman J, Yu J: Peginterferon alfa-2a plus ribavirin for chronic hepatitis C virus infection. *N Engl J Med* 2002;347:975-982.
- Lin C, Kwong AD, Perni RB: Discovery and development of VX-950, a novel, covalent, and reversible inhibitor of hepatitis C virus NS5A serine protease. *Infect Disord Drug Targets* 2006;6:3-16.
- Modi AA, Hoofnagle JH: New therapies for hepatitis C. *Hepatology* 2007;46:615-617.
- Zeuzem S: Telaprevir, peginterferon alfa-2a, and ribavirin for 28 days in chronic hepatitis C patients. *J Hepatol* 2008;49:157-159.
- McHutchison JG, Everson GT, Gordon SC, Jacobson IM, Sulkowski M, Kauffman R, McNair L, Alam J, Muir AJ, PROVE1 Study Team: Telaprevir with peginterferon and ribavirin for chronic HCV genotype 1 infection. *N Engl J Med* 2009;360:1827-1838.
- Hézode C, Forestier N, Dusheiko G, Ferenci P, Pol S, Gooser T, Bronowicki JP, Bourlière M, Gharakhanian S, Bengtsson L, McNair L, George S, Kieffer T, Kwong A, Kauffman RS, Alam J, Pawlotsky JM, Zeuzem S, PROVE2 Study Team: Telaprevir and peginterferon with or without ribavirin for chronic HCV infection. *N Engl J Med* 2009;360:1839-1850.
- McHutchison JG, Manns MP, Muir AJ, Terrault NA, Jacobson IM, Afdhal NH, Heathcote EJ, Zeuzem S, Reesink HW, Garg J, Bsharat M, George S, Kauffman RS, Adda N, Di Bisceglie AM, PROVE3 Study Team: Telaprevir for previously treated chronic HCV infection. *N Engl J Med* 2010;362:1292-1303.
- Akuta N, Suzuki F, Sezaki H, Suzuki Y, Hosaka T, Someya T, Kobayashi M, Saitoh S, Watahiki S, Sato J, Matsuda M, Kobayashi M, Arase Y, Ikeda K, Kumada H: Association of amino acid substitution pattern in core protein of hepatitis C virus genotype 1b high viral load and non-virological response to interferon-ribavirin combination therapy. *Intervirology* 2005;48:372-380.
- Akuta N, Suzuki F, Kawamura Y, Yatsuji H, Sezaki H, Suzuki Y, Hosaka T, Kobayashi M, Kobayashi M, Arase Y, Ikeda K, Kumada H: Predictive factors of early and sustained responses to peginterferon plus ribavirin combination therapy in Japanese patients infected with hepatitis C virus genotype 1b: amino acid substitutions in the core region and low-density lipoprotein cholesterol levels. *J Hepatol* 2007;46:403-410.
- Donlin MJ, Cannon NA, Yao E, Li J, Wahed A, Taylor MW, Belle SH, Di Bisceglie AM, Aurora R, Tavis JE: Pretreatment sequence diversity differences in the full-length hepatitis C virus open reading frame correlate with early response to therapy. *J Virol* 2007;81:8211-8224.
- Akuta N, Suzuki F, Kawamura Y, Yatsuji H, Sezaki H, Suzuki Y, Hosaka T, Kobayashi M, Kobayashi M, Arase Y, Ikeda K, Kumada H: Amino acid substitutions in the hepatitis C virus core region are the important predictor of hepatocarcinogenesis. *Hepatology* 2007;46:1357-1364.
- Fishman SL, Factor SH, Balestrieri C, Fan X, Dibisceglie AM, Desai SM, Benson G, Branch AD: Mutations in the hepatitis C virus core gene are associated with advanced liver disease and hepatocellular carcinoma. *Clin Cancer Res* 2009;15:3205-3213.
- Ge D, Fellay J, Thompson AJ, Simon JS, Shianna KV, Urban TJ, Heinzen EL, Qiu P, Bertelsen AH, Muir AJ, Sulkowski M, McHutchison JG, Goldstein DB: Genetic variation in *IL28B* predicts hepatitis C treatment-induced viral clearance. *Nature* 2009;461:399-401.
- Tanaka Y, Nishida N, Sugiyama M, et al: Genome-wide association of *IL28B* with response to pegylated interferon-alpha and ribavirin therapy for chronic hepatitis C. *Nat Genet* 2009;41:1105-1109.
- Suppiah V, Moldovan M, Ahlenstiel G, Berg T, Weltman M, Abate ML, Bassendine M, Spengler U, Dore GJ, Powell E, Riordan S, Sheridan D, Smedile A, Fragomeli V, Müller T, Bahlo M, Stewart GJ, Booth DR, George J: *IL28B* is associated with response to chronic hepatitis C interferon-alpha and ribavirin therapy. *Nat Genet* 2009;41:1100-1104.
- Rauch A, Kutalik Z, Descombes P, et al: Genetic variation in *IL28B* is associated with chronic hepatitis C and treatment failure: a genome-wide association study. *Gastroenterology* 2010;138:1338-1345.
- Thomas DL, Thio CL, Martin MP, Qi Y, Ge D, O'Huigin C, Kidd J, Kidd K, Khakoo SI, Alexander G, Goedert JJ, Kirk GD, Donfield SM, Rosen HR, Tobler LH, Busch MP, McHutchison JG, Goldstein DB, Carrington M: Genetic variation in *IL28B* and spontaneous clearance of hepatitis C virus. *Nature* 2009;461:798-801.